DOE/ER/13592-6

**PHOTOEXCITED CHARGE PAIR
ESCAPE AND RECOMBINATION**

DOE/ER/13592--6

DE92 004030

**Progress Report
for Period November 15, 1990 - November 14, 1991**

**Charles L. Braun
Department of Chemistry
Dartmouth College
Hanover, NH 03755**

November 15, 1991

Prepared for

**THE U. S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-86ER13592**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

Handwritten signature

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ABSTRACT

Progress in four areas is summarized in the following:

I GEMINATE CHARGE PAIR RECOMBINATION IN HEXANE

II FAST CURRENT MEASUREMENTS RESULTING FROM EXCITATION OF CHARGE TRANSFER(CT) STATES

III MEASUREMENT OF THE DIPOLE MOMENT OF EXCITED STATES BY DC CONDUCTIVITY.

IV CHARGE SEPARATION AT MACROSCOPIC INTERFACES BETWEEN ELECTRON DONOR AND ACCEPTOR SOLIDS

In a final section, personnel who have contributed to the project during the past budget period are described.

I GEMINATE CHARGE PAIR RECOMBINATION IN HEXANE

A manuscript describing our most recent work was recently published in The Journal of Physical Chemistry. The abstract of the manuscript follows:

Durene (1,2,4,5-tetramethylbenzene) dissolved in n-hexane was photoionized by 35 psec pulses at 266 nm. Transient absorption at 1064 nm arising chiefly from geminate electrons was detected and used to monitor the recombination of the electron-cation pairs produced by two-photon ionization. An excellent fit to the recombination kinetics was obtained by assuming that the distribution of initial electron-cation separations was of the form $r^2 \text{EXP} = r^2 / (2L^3) \exp(-r/L)$ with a mean radius $3L = 5.7$ nm.

The paper reports the most precise decay kinetics yet obtained for a collection of geminate charge pairs. The work appears to confirm results from our decade-old steady state experiments that suggested that a spherical exponential distribution of electron thermalization lengths is more appropriate than is the assumption that the distribution is gaussian, a hypothesis popular since the earliest suggestions that geminate charge pairs were important to radiation chemistry.

Subsequent to the submission of that manuscript, I visited Tom Scott's lab at NYU to carry out some of the work sketched in last year's progress report. We concentrated on the temperature dependence of the decay kinetics and learned something that is at least mildly surprising. We found that while the decay of a population of geminate pairs slows as the temperature is lowered, the decay kinetics are not controlled solely by the decreasing mobility of electron and cation. Instead, over the temperature range from 230 to 190 K in liquid hexane, the average thermalization length appears to decrease by about 20 percent. Most previous workers have assumed that thermalization lengths would decrease only inversely proportional to the medium density, a predicted change of only 3 percent. It will be important to confirm these results during the next budget period. An attempt to measure the decay kinetics at a second probe wavelength, 1.91 rather than 1.06 micrometer,

did not succeed because of lack of time and inability to achieve good overlap between pump and probe beams. That important experiment also deserves another go.

II FAST CURRENT MEASUREMENTS RESULTING FROM EXCITATION OF CHARGE TRANSFER(CT) STATES

This project continues to present difficulties. This past summer we achieved a breakthrough in purification of the 1:1 phenanthrene/pyromelliticdianhydride CT crystal. We found that we could co-zone-refine the 1:1 material provided that the two components had each been rigorously purified by chemical and physical means including their separate zone refining. The most stubborn impurity is anthracene which must be removed chemically from phenanthrene. That procedure had succeeded in producing phenanthrene with less than 1 ppm anthracene, and at that level, the co-zone-refining succeeded in removing most of the remaining anthracene. Preliminary results indicate that CT state dissociation is strongly suppressed in this clean material. If that result is supported by further work, it will mean that all previous work on photoconductivity arising from CT state dissociation has depended on impurities in the crystals or polymers.

The basic question that this project has attempted to answer is, "How can CT states dissociate given the large Coulomb barrier that holds donor cation and acceptor anion together?" The above result, if confirmed, means that the answer may be, "They can't!" While further work is clearly needed, this project has been temporarily put on the back burner, in order to follow up on the remarkable discovery that follows.

III MEASUREMENT OF THE DIPOLE MOMENT OF EXCITED STATES BY DC CONDUCTIVITY.

We succeeded in demonstrating that pulsed laser excitation of solution-phase molecules present between the electrodes of a conductivity cell gives rise to a fast dc photocurrent that is a direct measure of the change in dipole moment of the molecule upon excitation. A paper describing our initial results has been published and its abstract follows:

We report photocurrent transients arising from the pulsed laser excitation of the dipolar first excited singlet state S_1 of *trans* 4-dimethyl-amino-4'-nitrostilbene (DMANS) in toluene solution. The currents arise from rotational reorientation of DMANS dipoles with respect to the axis of an applied electric field. The method appears to offer a simple and general approach to the measurement of the change in dipole moment upon electronic excitation of a molecule.

It appears that the dipole measurement will allow direct quantitation of the extent of charge transfer in exciplex states, a question of long-standing interest. We are collaborating with Samir Farid and Ian Gould at Eastman Kodak in order to exploit the new method and preliminary results indicate that the fractional charge transfer in exciplex states increases as the electron affinity of the acceptor molecule increases.

We are working to improve the time resolution of the method; a new laser with subnanosecond pulse width will have to be acquired if we are to make substantial progress. We are also attempting to pulse on the dc electric field just prior to the laser pulse in order to allow the use of polar solvents with their high dark conductivity. If that works and if we can achieve high time resolution, we will be able to follow the evolution of dipole moments as CT states are excited to form excited states which then evolve in polar solvents to form solvent separated ion pairs. Thus the new method gives promise of allowing a new and physically revealing view of charge separation and recombination in solution.

IV CHARGE SEPARATION AT MACROSCOPIC INTERFACES BETWEEN ELECTRON DONOR AND ACCEPTOR SOLIDS

We are collaborating with the group of Dr. Ching Tang at Eastman Kodak in order to attempt to understand why charge carriers produced at a donor-acceptor interface separate into free carriers with such high efficiency. Recently, Dr. Tang showed that a photovoltaic cell that he constructed by vacuum evaporation had unusually good engineering efficiency (ca. 2%) as compared with other molecule-based solar cells. In the cell, charge carrier separation is believed to take place at the interface between a thin film donor layer and a thin acceptor layer. Dr. Tang's measurements were largely confined to cw excitation and standard IV curve recording. We are using

pulsed laser excitation and fast current measurements to understand why carrier dissociation at the interface is so favorable.

We have learned that because of the high capacitance of the ultra thin organic cells, all current measurements exhibit an RC "tail" that lasts far longer than the true flow of current within the cell. Analytical solution of the appropriate circuit equations (two resistances and two capacitances) has been achieved and appears to offer a clear approach to retrieval of the microscopic charge carrier motions.

PERSONNEL

Fuxing Hou, an associate professor in the Department of Physics, Peking University, continued on the project until August. Fuxing worked on the fast current measurements needed to gain mechanistic insights into the dissociation pathway of solid-state charge transfer states. Her extensive work on material purification led to the dilemma described under project I above.

Sergei Smirnov is a physicist from the Institute for Kinetics and Combustion in Novosibirsk, Russia. He spent 6 weeks in the group two years ago. He began a new one year appointment in October and is making good progress toward the goal of faster dipole measurements in polar as well as non-polar solvents.

Karyn Grzeskowiak, a third-year graduate student in the group, is heading the collaboration with Farid and Gould to learn about fractional charge transfer in a series of exciplex states.

Henry Spindler '92 worked fulltime on the molecular solar cell project during the past summer and continues his work part-time as he prepares to write a Senior Honors Thesis on his work. Henry decided that our previously crude understanding of the meaning of the transient currents that he recorded must be improved and proceeded to solve the detailed circuit equations. Henry is a candidate for a Rhodes and other scholarships and I feel privileged to have him in the group.

Christopher Morse '93 and **Kimberley Whalen '94** intend to begin undergraduate research projects this coming winter. Chris will do transient dipole measurements on a covalently-linked, donor-acceptor compound that

was sent to us by a Belgian group. Kim will measure exciplex fluorescence lifetimes, an important part of the Grzeskowiak project

END

**DATE
FILMED**

01/24/92