Impact of Observed Hyperfine Splitting on X-Ray Laser Gain

Joseph Nilsen, Jeffrey A. Koch, James H. Scofield, Brian J. MacGowan
Juan C. Moreno, and Luiz B. Da Silva

Lawrence Livermore National Laboratory
Livermore, CA 94550

This paper was prepared for submittal to
the 9th APS Topical Conference on Atomic Processes in Plasmas
San Antonio, Texas
November 15, 1993

November, 1993

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

IMPACT OF OBSERVED HYPERFINE SPLITTING ON X-RAY LASER GAIN

Joseph Nilsen, Jeffrey A. Koch, James H. Scofield, Brian J. MacGowan,

Juan C. Moreno, and Luiz B. Da Silva

Lawrence Livermore National Laboratory, P. O. Box 808, Livermore, California 94550

ABSTRACT

Line broadening mechanisms play an important role in determining the gain of X-ray laser transitions. Typically Doppler broadening is the primary mechanism which determines the linewidth of these transitions. However, we will present cases where the hyperfine effect is the dominant line broadening mechanism. Studying laser lines, which tend to have gain narrowed linewidths, enables us to observe these dramatic hyperfine effects which would be difficult to observe in opacity or Stark broadened lines. In this work we report the observation of hyperfine splitting on an X-ray laser transition and discuss how hyperfine splitting has a major impact on the laser gain. In our experiments we measure the lineshape of the $3p \rightarrow 3s$, $J = 0 \rightarrow 1$ transition in neon-like niobium and zirconium and observe a 28 mÅ splitting between the two largest hyperfine components in the niobium($Z=41$) line at 145.9 Å, in good agreement with theory.^{1,2} In zirconium($Z=40$), no splitting is observed since the hyperfine effect is proportional to the nuclear moment, and zirconium has zero nuclear moment, as is typical for even- Z elements. The hyperfine effect is shown to effect transitions which have a $2p_{1/2}$ vacancy in the closed neon-like core much more than those with a $2p_{3/2}$ vacancy. A comparison of the neon-like niobium laser spectrum with that of zirconium shows a dramatic reduction in the relative intensity of the niobium laser lines with the $2p_{1/2}$ vacancy.³ We also report the unusual behavior noticed recently in low- Z neon-like X-ray lasers in which ions with odd Z lase poorly, if at all, relative to ions with even Z .^{4,5} The hyperfine effect is shown to have a substantial impact on the gain of the low- Z ions with odd Z and helps explain their poor performance.

PREVIOUS EXPERIMENTS

Several different experiments over the last year led us to investigate the possibility of hyperfine effects on the X-ray laser output. First, recent experiments on low- Z neon-like ions have observed lasing on the $3p \rightarrow 3s$, $J = 0 \rightarrow 1$ transition in neon-like titanium($Z=22$), chromium($Z=24$), iron($Z=26$), and nickel($Z=28$) at 326, 285, 255, and 231 Å, respectively. However, when experiments were done with scandium($Z=21$), vanadium($Z=23$), and manganese($Z=25$) there was no obvious sign of lasing while only weak lasing was observed with cobalt($Z=27$) and copper($Z=29$).^{4,6} Second, experiments done on neon-like zirconium($Z=40$) and niobium($Z=41$) observed the pair of $J = 2 \rightarrow 1$ laser lines to be very similar intensity in zirconium while in niobium the shorter wavelength line at 138.6 Å is much stronger than the other $J = 2 \rightarrow 1$ line at 140.4 Å. Also, the $J = 0 \rightarrow 1$ line in niobium is much weaker than it is in zirconium. The above experiments suggested that there was a difference between ions with even Z and odd Z . Since ions with even Z tend to have no nuclear spin while those with odd Z do have nuclear spin and therefore a nuclear moment which leads to hyperfine effects we decided to investigate this possibility in more detail.

Hyperfine splitting affects the gain of the laser line by increasing the linewidth. Since the gain is inversely proportional to linewidth the gain will decrease. If the splitting is large enough, a single line may be split into several weaker lines. For neon-like laser lines the impact of the hyperfine effect is largest for the $J = 0 \rightarrow 1$ line which dominates the spectra of the low- Z neon-like ions. We did a series of experiments to measure the lineshape of the $J = 0 \rightarrow 1$ laser line in neon-like niobium. Niobium was chosen because it has a large nuclear spin, $I = 9/2$, and a large nuclear moment, $\mu = 6.167$, its wavelength is in the range of the available high resolution spectrometer, and it has lased.³

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

gso

EXPERIMENTAL SETUP

Experiments were conducted at Lawrence Livermore National Laboratory (LLNL) on the Nova laser using $\lambda = 0.53 \mu\text{m}$. In a typical experiment the Nova laser illuminates a $125 \mu\text{m}$ thick, 3.0 cm long slab target of niobium. The above length of the target was reduced by a 16% gap in the center which results in an actual length of 2.5 cm . The pump laser beam was a 500 ps square pulse with 2.4 kJ of energy in a $120 \mu\text{m}$ wide (FWHM) by 3.6 cm long line focus, resulting in a peak intensity of 130 TW/cm^2 .

The principal instruments were a time-gated, microchannel plate intensified grazing-incidence grating spectrograph (MCPIGS) and a high-resolution, grazing incidence grating spectrometer; both of these instruments observed the axial output of the X-ray laser. The MCPIGS provided angular resolution over 10 mrad near the X-ray laser axis, used a 1200 line per mm grating, and had spectral coverage of approximately 75 to $340 \text{ \AA}$. The high-resolution, grazing incidence grating spectrometer recorded time-integrated but spatially resolved data using a Princeton Instruments camera with a backside illuminated EEV CCD. This spectrometer was centered on the laser axis with an angular acceptance of 12 mrad and had a measured spectral resolution of 20000 at $146 \text{ \AA}$ and spectral coverage of $2 \text{ \AA}$. The angular or spatial resolution of both instruments was perpendicular to the target surface.

EXPERIMENTAL RESULTS AND ANALYSIS

Figure 1(a) shows the measured intensity (solid line) versus wavelength for the $3p \rightarrow 3s, J = 0 \rightarrow 1$ neon-like niobium laser line at $145.9 \text{ \AA}$. Two components are clearly visible with a separation of $28 \text{ m\AA}$, which is very close to the $32 \text{ m\AA}$ prediction (dotted line) given the $7 \text{ m\AA}$ resolution of the spectrometer. This is the shortest wavelength transition and most highly ionized plasma in which the hyperfine effect has been directly observed with the exception of some very recent accelerator experiments in hydrogen-like bismuth done at GSI Darmstadt.⁷ Figure 1(b) shows the measured intensity versus wavelength for the $3p \rightarrow 3s, J = 0 \rightarrow 1$ neon-like zirconium laser line at $150.4 \text{ \AA}$. The zirconium laser line is now a single narrow line which is slightly gain narrowed and a hundred times brighter than the niobium line. Both experiments used the same length targets and illumination conditions as described previously. A comparison of Figs. 1(a) and 1(b) show how the increased linewidth of the niobium line reduces the gain and dramatically reduces the laser output as compared

Fig. 1. Measure intensity vs wavelength around line center for the $3p \rightarrow 3s, J = 0 \rightarrow 1$ laser line in neon-like niobium (a) at $145.9 \text{ \AA}$ and in neon-like zirconium (b) at $150.4 \text{ \AA}$. Calculated splitting (dotted line) of $32 \text{ m\AA}$ for niobium compares well with $28 \text{ m\AA}$ measured splitting.

with a similar transition in zirconium, an adjacent ion which is not subject to the hyperfine effect.

For vanadium, assuming an ion temperature of 50 eV based on calculations, the hyperfine splitting reduces the gain coefficient of the $J = 0 \rightarrow 1$ laser line at 304 Å by 40% as shown in Fig. 2(a) which presents gain versus wavelength around line center normalized to a gain of unity if there was no hyperfine splitting. The three hyperfine components are shown by dotted lines. Assuming vanadium has the same gain coefficient of 2.6 cm^{-1} as observed for titanium,^{4,5} the hyperfine effect reduces the gain coefficient to 1.6 cm^{-1} for vanadium. For the 3.8 cm long targets tried with vanadium, this would make the vanadium fifty times weaker than the titanium. While this is still within the detectable range of the diagnostics, the hyperfine effect appears to play a major role in the non lasing of vanadium and scandium and the poor lasing of the other odd Z ions. Since the measured titanium gain was time averaged, the peak gain is no doubt larger. Therefore the relative effect of the hyperfine splitting will also be larger.

It is also interesting to calculate the impact of hyperfine splitting on other neon-like ions. The next element above niobium which has a large nuclear moment is indium ($Z=49$). Figure 2(b) shows the gain versus wavelength near line center for the $3p \rightarrow 3s$ neon-like indium lines at 89.0, 89.8, and 116 Å assuming an ion temperature of 1000 eV. The first two lines are the normal pair of $J = 2 \rightarrow 1$ lines, analogous to the selenium 206 and 209 Å lines, while the third is the $J = 0 \rightarrow 1$ line, analogous to the selenium 182 Å line.⁸ One notices a dramatic reduction in gain for the $J = 0 \rightarrow 1$ line. For this line it may be possible to observe all three components. For the pair of $J = 2 \rightarrow 1$ lines, the hyperfine effect is much larger for the 89.8 Å line because this is the transition with the $2p_{1/2}$ vacancy while the 89.0 Å line has the $2p_{3/2}$ vacancy. However, in preliminary experiments we have not been able to achieve lasing on any neon-like indium lines.

CONCLUSIONS

The hyperfine effect is shown to be the dominant line broadening mechanism for the $3p \rightarrow 3s$, $J = 0 \rightarrow 1$ neon-like niobium laser line at 145.9 Å. We measured the lineshape of this transition and observed a 28 mÅ splitting between the two largest hyperfine components, in good agreement with theory. This is the largest hyperfine splittings ever measured on a laser transition.

Fig. 2. Gain vs wavelength near line center for the $3p \rightarrow 3s$, $J = 0 \rightarrow 1$ laser line in neon-like vanadium(a) at 304 Å and for the two $J = 2 \rightarrow 1$ and one $J = 0 \rightarrow 1$ laser lines in neon-like indium(b). The three hyperfine components are given by dotted lines in (a). The gain is normalized to a gain of unity if there were no hyperfine effect.

The hyperfine effect is also shown to effect transitions which have a $2p_{1/2}$ vacancy in the closed neon-like core much more than those with a $2p_{3/2}$ vacancy. A comparison of the neon-like niobium laser spectrum with that of zirconium shows a dramatic reduction in the relative intensity of the niobium laser lines with the $2p_{1/2}$ vacancy. Finally, the hyperfine effect is shown to have a substantial impact on the gain of the low-Z ions with odd Z and helps explain their poor performance.

ACKNOWLEDGEMENTS

The authors would like to thank Jack B. Mitchell, Ron R. Wing, Sharon S. Alvarez, Judy M. Ticehurst, and the Nova facilities crew for providing support for the experiments. We would also like to thank Jay Moskovic and Yair Talmi of Princeton Instruments for the loan of the CCD camera. The support of S. B. Libby, D. A. Nowak and D. L. Matthews is greatly appreciated. Work performed under the auspices of the U. S. Department of Energy by the Lawrence Livermore National Laboratory under contract No. W-7405-ENG-48.

REFERENCES

1. J. Nilsen, J. A. Koch, J. H. Scofield, B. J. MacGowan, J. C. Moreno, and L. B. Da Silva, *Phys. Rev. Lett.* **70**, 3713 (1993).
2. J. H. Scofield and J. Nilsen, "Hyperfine splittings of neonlike lasing lines," University of California report UCRL-JC-115274, Livermore, CA 94550 (1993).
3. J. Nilsen, J. L. Porter, B. J. MacGowan, L. B. Da Silva, and J. C. Moreno, *J. Phys. B* **26**, L243 (1993).
4. T. Boehly, M. Russotto, R. S. Craxton, R. Epstein, B. Yaakobi, L. B. Da Silva, J. Nilsen, E. A. Chandler, D. J. Fields, B. J. MacGowan, D. L. Matthews, J. H. Scofield, and G. Shimkaveg, *Phys. Rev. A* **42**, 6962 (1990).
5. J. Nilsen, B. J. MacGowan, L. B. Da Silva, and J. C. Moreno, "Prepulse technique for producing low-Z Ne-like XUV lasers," *Phys. Rev. A* **48** (in press 1993).
6. J. Nilsen, J. C. Moreno, B. J. MacGowan, and J. A. Koch, *Applied Physics B* **57**, 309 (1993).
7. Thomas U. Kühl, GSI Darmstadt, private communication (August 1993).
8. J. Nilsen and J. H. Scofield, "Wavelengths of neon-like $3p \rightarrow 3s$ X-ray laser transitions," *Physica Scripta* (in press 1994).

END

**DATE
FILMED**

3/18/94