CHARACTERIZATION OF THE MULTIFRAGMENT DECAY CHANNEL IN HIGHLY EXCITED NUCLEI

Progress Report for the Period
March 15, 1992 - September 14, 1993

R.T. de Souza

Indiana University
Bloomington, Indiana, 47405

RECEIVED
MAR 01 1993
OSTI

October, 1992

Prepared for
THE UNITED STATES DEPARTMENT OF ENERGY
UNDER GRANT No. DE-FG02-92ER40714

**CHARACTERIZATION OF THE MULTIFRAGMENT DECAY CHANNEL
IN HIGHLY EXCITED NUCLEI**

**Progress Report for the Period
March 15, 1992 - September 14, 1993**

**R.T. de Souza
Principal Investigator**

**Department of Chemistry
Indiana University
Bloomington, Indiana, 47405**

October, 1992

**Prepared for
THE UNITED STATES DEPARTMENT OF ENERGY
UNDER GRANT DE-FG02-92ER40714**

NOTICE

This report was prepared as an account of work sponsored by the United States government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed or represents that its use would not infringe privately owned rights.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *EP*

Table of Contents

I. ABSTRACT	3
II. INTRODUCTION	4
III. RESEARCH PROGRAM	8
Extraction of the Multifragmentation Time Scale in Intermediate Energy Heavy-Ion Reactions	9
Expansion Effects in Intermediate Energy Heavy-Ion Reactions	21
Development of Low Threshold High Resolution Detectors	33
Development of a Transmission Parallel Plate Avalanche Counter for Ternary Fission Studies	43
Three-body calculations of Neck Emission of Intermediate Mass Fragments During Fission	57
Construction of a Compact Gas Handling system	77
IV. PUBLICATIONS AND ACTIVITIES	80
A. Articles	86
B. Invited Lectures	86
C. Contributed Papers at Professional Meetings	86
D. Professional Activities	88
V. PERSONNEL	89

I. ABSTRACT

Characteristics of multifragment decay in heavy-ion induced reactions at intermediate energies are explored. Evolution of the timescale for emission of intermediate mass fragments (IMF: $3 \leq Z \leq 20$) is investigated. Fragments associated with central collisions in the reactions $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50, 80$, and 110 MeV are emitted on an extremely fast timescale, $\tau \leq 75$ fm/c, comparable to the transit time of the projectile past the target nucleus. With increasing incident energy, mean fragment emission times decrease, consistent with statistical decay from highly excited systems or fast dynamical processes. To examine the importance of expansion effects in multifragmentation, the transverse kinetic energy of fragments was investigated. Evidence for expansion effects in the system $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50, 80$, and 110 MeV was indicated by the charge dependence of the mean isotropic kinetic energy. At the highest incident energy the data suggest the onset of volume emission. Construction and performance of low-threshold high resolution detector telescopes is discussed. Three-body Coulomb trajectory calculations are being used to probe kinematical correlations associated with neck emission of IMFs during fission. Initial design and testing of position sensitive parallel plate avalanche counters for upcoming ternary fission experiments is discussed.

II. INTRODUCTION

During the past year, the research activities of the Indiana University Nuclear Chemistry group supported by DOE Grant No. DE FG02-92ER40714 have experienced substantial growth. Although the research activities of the group remain primarily outside user in nature, a proposed experiment utilizing the Indiana University cyclotron has been approved. Major collaborative experiments were conducted at the National Superconducting Cyclotron Laboratory (Michigan State University) as well as at Laboratoire National Saturne (Saclay, France). An experiment at the Van de Graaf tandem at the University of Rochester constituted a small part of the research program. These collaborative efforts were conducted together with research groups from Washington University (St. Louis, MO), Michigan State University-NSCL (East Lansing, MI), University of Washington (Seattle, WA), University of Rochester (Rochester, NY), Laboratoire National Saturne (Saclay, France), and Ball State University (Muncie, IN). Results of this experimental work are described in Section III of this report.

Three new major experiments were conducted during this period. The first experiment focused on the mechanism of projectile breakup in alpha-cluster nuclei. The reactions studied were : $^{16}\text{O} + ^{197}\text{Au}$ and $^{24}\text{Mg} + ^{197}\text{Au}$ both at $E/A = 60$ MeV. The goal of this experiment is to distinguish between sequential decay and an "instantaneous" breakup mechanism of projectile decay by studying the correlations between alpha particles at forward angles. Analysis is currently underway at Washington University. The second experiment examined the characteristics of multifragmentation in the reaction

$^{84}\text{Kr} + ^{197}\text{Au}$ at $E/A = 35, 55,$ and 70 MeV. This excitation function constitutes the low energy end of the excitation function ($\text{Kr} + ^{197}\text{Au}$ at $E/A = 100, 200,$ and 400 MeV) previously measured at SATURNE. Analysis of some of the data from this experiment is currently underway at Indiana. The third experiment investigated the evolution of the fusion cross-section with increasing incident energy for systems of different fissility. The systems examined were $^{14}\text{N} + \text{Tb}, \text{Sm},$ and Ta at $E/A = 35$ MeV; $^{16}\text{O} + \text{Tb}, \text{Ta},$ and Au at $E/A = 25$ MeV; and $^{14}\text{N} + \text{Au}, \text{Sm},$ and Ta at $E/A = 100$ MeV. Analysis on a subset of this data has just begun at Indiana. In addition, a theoretical collaborative project was initiated with W. Friedman at University of Wisconsin (Madison, WI) to compare experimental signatures with model predictions of multifragment decay.

In addition to conducting new experiments, the main focus in the group involved the analysis of the reaction $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50, 80,$ and 110 MeV. In particular, two aspects of the multifragment decay observed in the system were studied. Two-particle correlation functions of intermediate mass fragments ($\text{IMF}: 3 \leq Z \leq 20$) were used to probe the mean emission time of the IMFs. The experimental data were compared with both two-body Koonin-Pratt calculations, as well as, three-body Coulomb trajectory calculations. Extremely short emission times, $\tau < 75$ fm/c, were deduced. These times are comparable to the transit time of the argon projectile past the gold target nucleus. This data represents one of the first systematic measurements of the evolution of the fragment emission timescale over a broad range of intermediate energies. The second area investigated was the dependence of the moments of the transverse kinetic energy spectra on the charge of the emitted particle. Currently, only statistical models which

reduce the nuclear density substantially following the collision are capable of reproducing the observed fragment multiplicities. Such a reduction in the density might result from an expansion due to the thermal pressure of the system, as well as, a dynamic compression-decompression cycle. Surface emission from an expanding system would result in a boost to the velocity of the emitted particle. This velocity shift would be manifested in the dependence of the moments of the transverse kinetic energy distribution on the mass of the emitted particle. Comparison of experimental data with model calculations indicate strong evidence for expansion effects. The onset of emission from the interior of the nucleus at the highest incident energy is also suggested.

Significant effort was expended towards the design, fabrication, and testing of low-threshold high resolution detector telescopes. These detectors consist of three elements arranged in a collinear geometry. The first element of the telescope is an axial field ionization chamber. The second element is an ion-implanted silicon crystal with a large active area. The large dynamic range of the telescope is provided by the third element, a CsI(Tl) scintillator crystal. A principal feature of these detectors is their compact geometry which allows a high resolution telescope to replace a single element of the MSU Miniball 4π array. The new detectors were utilized in conjunction with the MSU Miniball for three experiments conducted during the spring/summer.

The development of equipment necessary for upcoming approved experiments at both Indiana University Cyclotron Facility and the K1200 at Michigan State University also required significant attention. The principal focus in this area is the design and development of a transmission parallel plate avalanche counter, a gas handling system,

and a PC based CAMAC data acquisition system. All of these activities are described in greater detail in Section III of this report.

During the academic year, the principal investigator devoted 70% of his time to research supported by the grant, while during the summer months of 1992 he devoted 100% of his efforts to the research projects. The postdoctoral fellow (DF) spent all of his time on grant research. The graduate student (SC and BD) are first year graduate students. One of the graduate students (SC) is supported by the NSF through the Indiana University Cyclotron Facility (IUCF). The other graduate student (BD) is currently supported by a teaching assistantship. Consequently, during the academic year each student will spend 50% of his time on grant research. During the summer months, each graduate student will spend 100% of his time on grant research. Two undergraduate students (JM and AS) worked full time on grant research during the summer months. One of the two students (JM) was supported by the NSF through IUCF. A third undergraduate student (AW) participates in a research for credit program towards the completion of an Honors degree in Chemistry. During the academic year, the students work approximately 8-10 hrs./week each and are supported by the DOE grant.

To the best of our knowledge, we have complied with all grant requirements. It is a pleasure to acknowledge the direct support of our research program by the United States Department of Energy. IN addition, we are grateful to the National Science Foundation for having provided indirect support through the use of the facilities of the Indiana University Cyclotron Facility.

III. RESEARCH PROGRAM

Extraction of the Multifragmentation Time Scale in Intermediate Energy Heavy-Ion Reactions

D. Fox and R.T. de Souza

Indiana University Cyclotron Facility
and Department of Chemistry
Indiana University, Bloomington, IN 47405

and

L. Phair, D.R. Bowman[†], N. Carlin^{*}, C.K. Gelbke, W.G. Gong[‡],
Y.D. Kim[§], M.A. Lisa, W.G. Lynch, G.F. Peaslee, M.B. Tsang,
and F. Zhu^{**}

National Superconducting Cyclotron Laboratory
and Department of Physics and Astronomy
Michigan State University, East Lansing, MI 48824

ABSTRACT

The spatial and temporal extent of a system decaying by multifragment emission is deduced. Two-particle intermediate mass fragment correlation functions measured for central $^{36}\text{Ar} + ^{197}\text{Au}$ collisions at $E/A=50\text{-}110$ MeV indicate a rapid decay, < 75 fm/c, of the highly excited system. Furthermore, the behavior of the correlation function at large relative velocities suggests that considerable charge loss occurs prior to fragment emission.

[†] Present address: Chalk River Nuclear Laboratories, Chalk River, Ontario K0J 1J0, Canada.

^{*} Present address: Instituto de Fisica, Universidade de Sao Paulo, C. Postal 20516, CEP 01498, Sao Paulo, Brazil.

[‡] Present address: Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720.

[§] Present address: Indiana University Cyclotron Facility, Indiana University, Bloomington, IN 47405.

^{**} Present address: Brookhaven National Laboratory, Upton NY 11973.

Nuclear systems at high excitation can decay by multiple emission of intermediate mass fragments (IMF: $3 \leq Z \leq 20$) [1-10]. Key to understanding this phenomenon is knowledge of the source size and lifetime. Both dynamical and statistical theories predict breakup of the system on a relatively short timescale $\tau \approx 100$ fm/c [3-5,11]. Both theories, however, require a reduction in the nuclear density to correctly predict fragment emission probabilities since fragment formation and emission is substantially enhanced at low density [1,12-14]. Such a reduction in the density of the system might result either from a rapid dynamical compression-decompression cycle induced in central intermediate energy heavy-ion collisions [1-5] or from the thermal pressure of the system [15-18]. At present, however, little experimental information exists about the evolution of the emission time-scale and density with increasing excitation [19-21].

To characterize the evolution of the IMF emission time scale with increasing incident energy for central collisions, an excitation function has been measured for the system $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50, 80$, and 110 MeV. The experiment was performed at the National Superconducting Cyclotron Laboratory at Michigan State University (MSU-NSCL). Beams of ^{36}Ar , with an intensity of approximately 1×10^8 particles per second, impinged on a ^{197}Au target of areal density 1 mg/cm^2 . Particles emitted in the angular range $9^\circ \leq \theta_{\text{lab}} \leq 160^\circ$ were detected with the MSU Miniball array [22]. Each element of the Miniball consists of a 4 mg/cm^2 plastic scintillator foil backed by a 2 cm thick CsI(Tl) crystal. Particles which punched through the plastic scintillator foils were identified by charge up to $Z=18$. In addition, isotopes of hydrogen and helium were identified. The approximate energy thresholds were $E_{\text{th}}/A \approx 2 \text{ MeV}$ for $Z=3$, $E_{\text{th}}/A \approx 3 \text{ MeV}$ for $Z=10$, and $E_{\text{th}}/A \approx 4 \text{ MeV}$ for $Z=18$. Particles which stopped in the scintillator foil were recorded, but could not be identified by atomic number. Energy calibrations for the forward detectors were obtained by elastic scattering of ^4He , ^6Li , ^{10}B , ^{12}C , ^{16}O , ^{20}Ne and ^{35}Cl beams from a ^{197}Au target. The incident energies ranged from $E/A=4.5$ to 20 MeV . For detectors in rings 1-4, $9^\circ \leq \theta_{\text{lab}} \leq 40^\circ$, these calibrations are estimated to be accurate to within 5%. During the experiment, the trigger condition required that at least two detectors of the Miniball were triggered in order to record an event.

As shown in Fig. 1(a), intermediate energy heavy ion reactions exhibit broad multiplicity distributions. With increasing incident energy the multiplicity distribution extends to larger values of the charged particle multiplicity, N_c , reaching as many as 40 charged particles at $E/A=110$ MeV. A simple geometrical model can be used to relate the charged particle multiplicity, N_c to the impact parameter, b [23] allowing selection of the same impact parameter range at different bombarding energies. The gates on N_c used to select central collisions are depicted in Fig. 1(a). The gates correspond to $b/b_{\max} \leq 0.2$ where the quantity $b_{\max}$ corresponds to the maximum impact parameter which will result in events with two charged particles being detected.

The energy spectra of boron fragments from central $^{36}\text{Ar} + ^{197}\text{Au}$ collisions at $E/A = 110$ MeV emitted in the angular range $\theta_{\text{lab}}=16^\circ\text{-}40^\circ$ are shown in Fig. 1(b). These energy spectra appear to be reasonably described by a single Maxwellian consistent with emission from a single source. The energy spectra of IMFs associated with peripheral collisions, in contrast, are more complex and exhibit a significant non-equilibrium component [24,25]. Thus, restricting the analysis to central collisions might allow comparison of the experimental data to models which stress the equilibrium features of the system formed in central collisions.

The experimental correlation function, $R(v_{\text{red}})$, for two IMFs is defined in terms of the ratio of the coincidence yield, Y_{12} , to the product of the single particle yields, Y_1 and Y_2 .

$$\sum_{p_1, p_2} Y_{12}(p_1, p_2) = C [1 + R(v_{\text{red}})] \sum_{p_1, p_2} Y_1(p_1) Y_2(p_2)$$

where p_1 and p_2 are the laboratory momenta of IMFs 1 and 2, the reduced velocity is $v_{\text{red}} = (p_1/m_1 - p_2/m_2)/(Z_1 + Z_2)^{1/2}$, where m_i is taken as twice the atomic number, Z_i , and C is a normalization constant determined by the requirement that $\langle R(v_{\text{red}}) \rangle = 0$ at large relative momenta where the final state interaction is small. (The use of v_{red} instead of the relative momentum, allows the correlation functions to be summed over different pairs of IMFs [26].)

The correlation functions for two IMFs emitted in central collisions at the three

bombarding energies are shown in Fig. 2(a). The correlation functions are constructed over all IMF pairs with $4 \leq Z_1, Z_2 \leq 9$ emitted into the angular acceptance $16^\circ \leq \theta_{lab} \leq 40^\circ$. Also shown in Fig. 2(a) are data for the same system at 35 MeV/nucleon [26]. At all four beam energies the correlation functions exhibit the same general characteristic of a minimum at small v_{rel} arising from the repulsive Coulomb final-state interaction between the two emitted fragments. With increasing incident energy the width of the minimum at small v_{rel} becomes larger indicating an increase in the magnitude of the final-state interaction between the emitted IMFs. This increased interaction corresponds to a decrease in the spatial-temporal size of the emitting source.

To extract a mean emission time for the IMFs emitted in central collisions, the experimental correlation functions are compared to theoretical calculations using the Koonin-Pratt formalism [25-27]. The curves shown in Fig. 2(a) correspond to calculations with a mean emission time $\tau=5-200$ fm/c using the energy spectra associated with central collisions at $E/A=110$ MeV. Surface emission from a source of radius $R_s=12$ fm and velocity $v_s=0.041c$ was assumed. This velocity corresponds to 50% linear momentum transfer at $E/A = 110$ MeV. Using the energy spectra for central events from the other beam energies results in very similar theoretical correlation functions. Comparison of the experimental correlation functions to the theoretical curves reveals that the mean emission time, according to the Koonin-Pratt formalism, decreases from $\tau \approx 100$ fm/c at $E/A=35$ MeV to $\tau < 50$ fm/c at $E/A=110$ MeV. These emission times are comparable to the transit time of an Ar projectile past a Au nucleus (80 fm/c at 35 MeV, 50 fm/c at 110 MeV).

Strong final state interactions, as depicted by the wide Coulomb minima in the data, represent evidence of either a fast emission time scale or a small source size. Disentangling the spatial and temporal extent of the source requires an assumption of the source radius, R_s . The sensitivity of the Koonin-Pratt calculations to the assumption of R_s is shown in Fig. 2(b). If the source radius is assumed to be 8 fm, the data from central interactions at $E/A=110$ MeV are consistent with $\tau < 75$ fm/c. (For comparison, the radius of a source consisting of two touching carbon nuclei is 5.5 fm.)

Calculations to test the sensitivity of the deduced emission time to the assumption

of the source velocity, v_s were performed. The calculations demonstrated that the deduced mean emission time is fairly insensitive to variations in v_s . In addition, dependence of the extracted emission time scale on the choice of normalization regions was also explored. For all the data and calculations presented, the normalization constant, C , was determined in the region $26 \leq v_{\text{red}} \leq 35$. A maximum variation in τ of ≈ 50 fm/c is observed if different regions of v_{red} , $25 \leq v_{\text{red}} \leq 45$ are used in determining C .

The presence of a large target-like residue may affect the IMF correlation functions [19, 21, 28]. In order to investigate this effect, three-body Coulomb trajectory calculations [25] have been performed. In Fig. 3(a) the results of these calculations for central collisions of $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 110$ MeV are presented. To provide consistency with our previous two-body Koonin-Pratt calculations the initial source was assumed to be a gold nucleus ($Z_s=79$, $A_s=197$) with $R_s=12$ fm, and $v_s=0.04c$. The calculated correlation function exhibits a much steeper rise at large v_{red} than the data. The rise at large v_{red} can be understood since selection of large v_{red} preferentially selects IMF-IMF pairs with a large opening angle. In these cases, the Coulomb interaction of the heavy residue increases the relative momentum of the IMF pair. In contrast, IMF pairs with small relative momenta preferentially correspond to a small opening angle. In these latter cases, the Coulomb interaction of the heavy residue does not affect the relative velocity. Naturally, a reduction of the atomic number of the emitting source reduces this enhancement of the correlation function at large v_{red} . If the source is assumed to be approximately half the charge of a gold nucleus ($Z_s=40$, $A_s=96$, $R_s=9.4$ fm, $v_s=0.04$), then the behavior of the experimental correlation function is reproduced at all values of v_{red} . The choice of the radius for the $Z_s=40$ nucleus assumes that the density is the same as in the case of the gold nucleus. Comparison of the trajectory calculations with the data also indicate a fast mean emission time, $\tau \approx 75$ fm/c. In Fig. 3(b) the trajectory calculations are compared to the correlation function for central collisions at $E/A = 50$ MeV. At this incident energy, the deduced emission time is $\tau \approx 100$ fm/c. The assumption a large source ($Z_s=79$), in this case, also produces too steep of a rise at large v_{red} , a result inconsistent with the experimental data. Comparison of the experimental data from central collisions

at $E/A=80$ MeV with trajectory calculations (not shown) also yield mean IMF emission times of $\tau \approx 100$ fm/c. In all cases, changing R_s , A_s or v_s only results in a small change in τ , $\Delta\tau < 50$ fm/c. Thus, while the behavior of the correlation function at large values of v_{red} exhibits sensitivity to the charge of the emitting system, the shape of the correlation function at small values of v_{red} is dominated by the mean emission time τ .

In summary, two IMF correlation functions have been constructed for central collisions of $^{36}\text{Ar}+^{197}\text{Au}$ at $E/A=50, 80$ and 110 MeV. Comparison with calculations based on both the Koonin-Pratt two-body formalism and three body Coulomb trajectory model indicate a short emission time scale, $\tau \approx 50\text{-}100$ fm/c for central events. Furthermore, with increasing incident energy the spatial-temporal size of the emitting system decreases. The deduced emission times are comparable to the transit time of the Ar projectile past the Au target. The extracted emission times are of comparable magnitude to the expansion and equilibration times predicted by microscopic dynamical theories [1]. Statistical models of sequential decays of hot nuclear systems ($T=13$ MeV) also predict comparable emission times ($\tau \approx 80$ fm/c)[11]. However, in such cases the underlying assumption of complete equilibration between successive emissions may not be valid. Coulomb trajectory calculations indicate that the large v_{red} behavior of the correlation function is sensitive to the charge of the emitting source. Based upon these calculations, it appears that substantial charged particle emission occurs prior to multifragment disintegration. Alternatively, the small charge of the emitting system might also be due to fragment formation and emission from the interior of the excited nuclear system particularly at the highest incident energy.

We would like to acknowledge the valuable assistance of the staff and operating personnel of the K1200 cyclotron at the National Superconducting Cyclotron Laboratory for providing the high quality beams which made this experiment possible. This work was supported by the U.S. Department of Energy under DE FG02-92ER40714 and the National Science Foundation under Grant No. PHY-89-13815 and PHY-90-15957.

References

- [1] W. Bauer, G.F. Bertsch, and S. Das Gupta, Phys. Rev. Lett. **58**, 863 (1987).
- [2] K. Sneppen and L. Vinet, Nucl. Phys. **A480** 342 (1988).
- [3] J. Aichelin, G. Peilert, A. Bohnet, A. Rosenhauer, H. Stocker, and W. Greiner, Phys. Rev. **C37**, 2451 (1988).
- [4] G. Peilert, H. Stocker, W. Greiner, A. Rosenhauer, A. Bohnet, and J. Aichelin, Phys. Rev. **C39**, 1402 (1989).
- [5] D.H. Boal and J.N. Glosli, Phys. Rev. **C37**, 91 (1988).
- [6] L.G. Moretto, Nucl. Phys. **A247**, 211 (1975).
- [7] W.A. Friedman and W.G. Lynch, Phys. Rev. **C28**, 16 (1983).
- [8] W.A. Friedman and W.G. Lynch, Phys. Rev. **C28**, 950 (1983).
- [9] W.A. Friedman, Phys. Rev. **C40**, 2055 (1989).
- [10] W.A. Friedman, Phys. Rev. **C42**, 667 (1990).
- [11] W.A. Friedman, Proceedings of the International Symposium Towards a Unified Picture of Nuclear Dynamics, Nikko, Japan, June 1991, edited by Y. Abe, S.M. Lee, and F. Sakata, AIP Conf. Proc. No. 250, (AIP, New York, 1992) p. 422.
- [12] A. Vicentini, G. Jaucci, and V.R. Pandharipande, Phys. Rev. **C31**, 1783 (1985).
- [13] R.J. Lenk and V.R. Pandharipande, Phys. Rev. **C34**, 177 (1986).
- [14] T.J. Schlagel and V.R. Pandharipande, Phys. Rev. **C36**, 162 (1987).
- [15] G. Bertsch and P.J. Siemens, Phys. Lett. **B126**, 9 (1983).
- [16] H. Sagawa and G.F. Bertsch, Phys. Lett. **B155**, 11 (1985).
- [17] H. Schulz, B. Kampf, H.W. Barz, G. Ropke, and J. Bondorf, Phys. Lett. **B147**, 17 (1984).

- [18] S. Levit and P. Bonche, Nucl. Phys. **A437**, 426 (1984).
- [19] R. Trockel, U. Lynen, J. Pochodzalla, W. Trautmann, N. Brummund, E. Eckert, R. Glasow, K.D. Hildenbrand, K.H. Kampert, W.F. Muller, D. Pelte, H.J. Rabe, H. Sann, R. Santo, H. Stelzer, and R. Wada, Phys. Rev. Lett. **59**, 2844 (1987).
- [20] The mean emission time, τ , and the "half-life" for intermediate mass fragment (IMF) emission, $\tau_{\text{IMF-IMF}}$, defined in Ref. [19], are related as $\tau=1.44 \tau_{\text{IMF-IMF}}$.
- [21] R. Bougault, J. Colin, F. Delaunay, A. Genoux-Lubain, A. Hajfani, C. LeBrun, J.F. Lecolley, M. Louvel, and J.C. Steckmeyer, Phys. Lett. **B232**, 291 (1989).
- [22] R.T. de Souza, N. Carlin, Y.D. Kim, J. Ottarson, L. Phair, D.R. Bowman, C.K. Gelbke, W.G. Gong, W.G. Lynch, R.A. Pelak, T. Peterson, G. Poggi, M.B. Tsang, and H.M. Xu, Nucl. Instr. and Meth. **A295**, 109 (1990).
- [23] C. Cavata, M. Demoulin, J. Gosset, M.C. Lemaire, D.L. Hote, J. Poitou, and O. Valette, Phys. Rev. **C42**, 1760 (1990).
- [24] L. Phair, D.R. Bowman, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa, W.G. Lynch, G.F. Peaslee, R.T. de Souza, M.B. Tsang, and F. Zhu, Nucl. Phys. A (to be published)
- [25] Y.D. Kim, R.T. de Souza, C.K. Gelbke, W.G. Gong, and S. Pratt, Phys. Rev. **C45**, 387 (1992).
- [26] Y.D. Kim, R.T. de Souza, D.R. Bowman, N. Carlin, C.K. Gelbke, W.G. Gong, W.G. Lynch, L. Phair, M.B. Tsang, and F. Zhu, Phys. Rev. **C45**, 338 (1992).
- [27] Y.D. Kim, R.T. de Souza, D.R. Bowman, N. Carlin, C.K. Gelbke, W.G. Gong, W.G. Lynch, L. Phair, M.B. Tsang, F. Zhu, and S. Pratt, Phys. Rev. Lett. **67**, 14 (1991).
- [28] P.A. DeYoung, M.S. Gordon, Xiu qin Lu, R.L. McGrath, John Alexander, D.M. de Castro Rizzo, and L.C. Vaz, Phys. Rev. **C39**, 128 (1989).

Figure Captions

Fig. 1: (a) Charged particle multiplicity, N_c , from $^{36}\text{Ar} + ^{197}\text{Au}$ reactions at $E/A=50, 80$ and 110 MeV. The arrows indicate gates on N_c used to select central events, $b/b_{\text{max}} \leq 0.2$, at each energy based on a simple geometrical model that relates N_c to the impact parameter, b . (b) Energy spectra for boron fragments emitted from central events at $E/A=110$ MeV, and $\theta_{\text{lab}}=19.5^\circ$ (circles), 27° (squares), and 35.5° (triangles).

Fig. 2: (a) Two IMF correlation functions for $4 \leq Z_1, Z_2 \leq 9$ emitted at $\theta_{\text{lab}}=16^\circ\text{-}40^\circ$ for central events from $^{36}\text{Ar} + ^{197}\text{Au}$ at $50, 80$ and 110 MeV/nucleon. Data from Ref. [26] for 35 MeV/nucleon are included for comparison. The results of calculations for the 110 MeV/nucleon data using the Koonin-Pratt formalism are shown as lines, for details see the text. (b) Koonin-Pratt calculations for $R_s=8$ with $v_s=0.041c$ and $\tau=50, 100$, and 200 fm/c.

Fig. 3: Results of three-body Coulomb trajectory calculations for central $^{36}\text{Ar} + ^{197}\text{Au}$ interactions at $E/A=110$ MeV (a) and $E/A=50$ MeV (b). In each panel the data are shown as solid points. The dash-dot curve is the result of three-body Coulomb trajectory calculations in which the source is a Au nucleus with $R_s=12$ fm and $v_s=0.04c$. All other curves represent trajectory calculations for a source with $A_s=96$, $Z_s=40$, $R_s=9.4$, and $v_s=0.04c$.

Figure 1

Figure 2

$^{197}\text{Au}(^{36}\text{Ar}, Z_1 Z_2) \Theta_{\text{lab}}: 16^\circ - 40^\circ, 4 \leq Z_1, Z_2 \leq 9$

Figure 3

Expansion Effects in Intermediate Energy Heavy-Ion Reactions

R.T. de Souza and D. Fox
Indiana University Cyclotron Facility
and Department of Chemistry
Indiana University, Bloomington, IN 47405

W.A. Friedman
Department of Physics, University of Wisconsin, Madison, WI 53706

and

L. Phair, D.R. Bowman[†], C.K. Gelbke, W.G. Gong^{*}, Y.D. Kim[‡]
M.A. Lisa, W.G. Lynch, G.F. Peaslee, M.B. Tsang, and F. Zhu[§]
National Superconducting Cyclotron Laboratory
and Department of Physics and Astronomy
Michigan State University, East Lansing, MI 48824

ABSTRACT

Transverse kinetic energy spectra of fragments in the reaction $^{36}\text{Ar} + ^{197}\text{Au}$ are used as a probe of the multifragmentation mechanism. The mass dependence of the average transverse kinetic energy is examined in the context of statistical decay of an expanding, emitting source model. The role of source expansion during emission is explored. Results suggest the onset of volume emission at the highest incident energy.

[†]Present Address: Chalk River Nuclear Laboratories, Chalk River, Ontario K0J 1J0, Canada.

^{*}Present Address: Lawrence Berkely Laboratory, University of California, Berkeley, CA 94720.

[‡]Present Address: Indiana University Cyclotron Facility, Indiana University, Bloomington, IN 47405

[§]Present Address: Brookhaven National Laboratory, Upton NY 11973.

For highly excited nuclear matter multiple emission of intermediate mass fragments (IMF: $3 \leq Z \leq 20$) is an important deexcitation mode [Bau87a], [deS92a], [Bow92a]. Despite considerable theoretical [Ber83a], [Fri90a], [Aic88a], [Sag85a], [Sch84a] and experimental [deS92a], [Bow92a] investigation of heavy-ion induced multifragmentation the exact nature of the reaction mechanism is unclear.

In the reaction $^{129}\text{Xe} + ^{197}\text{Au}$ at $E/A = 50$ MeV [Bow92a] an average of six IMFs were associated with a central collision of projectile and target nuclei. Dynamical and standard statistical models were unable to predict the large multiplicity of IMFs observed in central collisions [Bow92a].

A model which allows a statistically decaying nucleus to expand to low density under the action of thermal pressure and the restoring nuclear attraction was capable of predicting the observed fragment multiplicities [deS92a], [Bow92a]. The reduction of the nuclear density in the expanding-emitting source (EES) model is a necessary feature in allowing enhanced fragment formation and emission.

To obtain more information about the multifragmentation mechanism, in particular the equilibration and expansion aspects, kinematic features of IMFs in the reaction $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50, 80$, and 110 MeV have been examined. We have focused on the transverse kinetic energy, defined by $E_{\text{trans}} = E \sin^2 \theta$, where E and θ are the laboratory energy and emission angle of the IMF, respectively.

The experiment was performed at the National Superconducting Cyclotron Laboratory at Michigan State University (MSU-NSCL). Beams of ^{36}Ar projectiles impinged on a 1 mg/cm^2 thick target. Charged particles were detected in the angular range $9^\circ \leq \theta \leq 160^\circ$ with the MSU Miniball array [deS90a]. Each element of the Miniball consists of a 4 mg/cm^2 thick plastic scintillator foil backed by a 2 cm thick CsI(Tl) crystal. Particles which punched through the plastic scintillator were identified by atomic number up to $Z=18$. The detector energy thresholds were $E_{\text{th}}/A \approx 3 \text{ MeV}$ for $Z=3$, and $E_{\text{th}}/A \approx 4 \text{ MeV}$ for $Z=18$. Energy calibrations are estimated to be accurate within 10%.

The transverse kinetic energy is especially useful since it suppresses the contribution of peripheral and non-equilibrium processes, such as projectile breakup, in

which the momentum of the fragments is predominately in the beam direction. For central collisions the emitting system acquires a velocity along the beam direction. E_{trans} is also insensitive to this velocity.

We have restricted our attention to collisions at small impact parameters where the probability of forming a single equilibrated source is greatest. These reactions were selected on the basis of the total charged particle multiplicity based on a geometrical relationship between charged particle multiplicity and impact parameter [Cav90a]. All the data presented are associated with central collisions selected by the multiplicity gates indicated in Fig.1a.

Representative transverse kinetic energy spectra of Boron fragments are shown in Fig 1(b) for each of the three incident energies. The high energy portion of the spectra is characterized by a simple exponential. This behavior can be understood in the context of the statistical decay of a thermalized source. Calculations with the EES model indicate that the asymptotic logarithmic slopes of the transverse kinetic energy spectra are nearly equal to the initial (maximum) temperature of the system.

To examine the relationship between the initial temperature of the system and the fragment charge, the logarithmic slopes of the transverse kinetic energy spectra for IMFs emitted from central collisions, have been parameterized with a simple exponential. The values of these slopes (slope parameters) for fragments with $Z=3$ to 9 at each of the three incident energies are shown in Fig 2(a). The parameters are determined to an accuracy of $\approx 10\%$, which represents the uncertainty in the extracted slope parameters due to the selection of the fit region. The constancy of the slope parameters for each bombarding energy suggests that different fragments originate from a common source. For a given fragment, the systematic increase in these parameters with increasing incident energy further suggests that the maximum energy deposited increases accordingly.

To facilitate comparison of the experimental data with models involving the isotropic decay of a source, we introduce the quantity $\langle E_t \rangle$, $\langle E_t \rangle = 3/2 \langle E_{\text{trans}} \rangle$. The quantity $\langle E_t \rangle$ corresponds to the mean kinetic energy of isotropically emitted fragments. The low energy portion of the transverse kinetic energy spectrum dominates the

behavior of the first moment of the distribution, and consequently $\langle E_l \rangle$. This portion of the transverse kinetic energy spectra is influenced by several different factors in addition to the initial excitation of the system.

The dependence of $\langle E_l \rangle$ on the charge of the fragment emitted in a central collisions is shown for each of the three incident energies in Fig. 2(b). $\langle E_l \rangle$ exhibits an essentially linear dependence on the fragment charge. For a given fragment, $\langle E_l \rangle$ also increases with increasing incident energy. This systematic behavior of $\langle E_l \rangle$ strengthens the suggestion that the fragments originate from a common source, and that some degree of equilibrium has been established. To gain insight into the physics underlying this behavior we have examined the charge dependence of $\langle E_l \rangle$ in the context of the EES model.

One can identify contributions to the kinetic energy from three distinct categories: thermal, Coulomb, and collective expansion. A systematic study of EES calculations results in general conclusions for each of the three individual contributions to $\langle E_l \rangle$.

The thermal contribution is provided by the average of $2T$ over the emission process, $\langle 2T \rangle$. The quantity T is the instantaneous temperature at the time of emission. Both particle emission and adiabatic expansion cool the source from its initial temperature. Consequently, $\langle 2T \rangle < 2 T_{\max}$. The average kinetic energy of each fragment is determined by the detailed sequence leading to its production. The calculations also indicate that $\langle T \rangle$ increases only slightly with increasing $T_{\max}$. For example, $\langle T \rangle$ for Boron fragments increases from 6.5 to 7.25 for $T_{\max}$ changing 12 to 18 MeV. The reduced sensitivity of $\langle T \rangle$ to $T_{\max}$ arises from the preferred emission at low densities and lower temperatures in the EES model.

The Coulomb contribution to the average kinetic energy arises from the repulsion between the fragments and the emitting source. If all the emission sequences were identical, the Coulomb energy would be roughly proportional to the Z of the emitted fragment for a large source. However, the dependence is not strictly linear in Z since the decay sequences for the various fragments are different. Calculated Coulomb contributions are considerably less than those associated with the initial source. This result is primarily due to the decrease of the Coulomb barrier with reductions in source

charge and density. Since higher initial excitation causes enhanced particle emission it consequently results in lower average Coulomb repulsion.

Based on the thermal and Coulomb contributions alone a decrease in the average kinetic energy with increasing incident energy would be expected. Standard statistical calculations [Fri83a] for sources with initial $A=175$ and $Z=70$, which do not include expansion, provide average kinetic energies for Fluorine fragments which fall steadily from 60 MeV to 41 MeV for initial temperatures increasing from 12 MeV to 18 MeV. The same calculations also show that the slope of the Z -dependence of the kinetic energy decreases with increasing excitation energy. In marked contrast, the experimental data suggest a nearly constant, Z -dependence, and an increase of the average kinetic energy with increasing incident energy (presumably increasing excitation energy).

The last contribution to the average kinetic energy is associated with the collective expansion of the source. Two factors contribute to expansion of the system: increased thermal pressure and any initial expansion due, for example, to a dynamical compression-decompression cycle [Boa88a]. This expansion provides a boost to the radial component of the thermal velocity, so that the resulting change in the kinetic energy of the fragment depends strongly on its mass. For small excitation, the contribution from the expansion is slight due to low velocities and emission near the turning point of the expansion-contraction oscillation. For larger excitations the emission is more sensitive to the expansion velocity, and the effect of expansion is greatest in reactions where high excitation energy results in an unconstrained expansion. Calculations indicate that for the reaction $^{36}\text{Ar}+^{197}\text{Au}$ at $E/A=110$ MeV such monotonic expansion is most likely to be important. These unconstrained expansions provide the largest average kinetic energy, and the largest Z -dependence.

Comparison of specific EES calculations with the experimental data is shown in Fig. 3. For simplicity, each calculation depicted was performed for a single set, rather than a distribution, of initial conditions. At each incident energy an initial temperature, a few MeV below the maximum (indicated by the asymptotic slope of the spectra), was chosen: 12, 15, and 18 MeV respectively. The source mass and initial expansion energy, E_e , were varied. The cases were limited to those which predict mean multiplicities of

IMFs, and of charged particles, consistent with previous measurements [deS92a]. With this constraint a preference for an effective nuclear compressibility of $K=200$, as against $K=144$ was found. The calculated curves shown are in qualitative agreement with the data. The Z -dependence is quite well matched.

We have also explored the role of volume emission, suggested by sudden freeze-out models [Gro90a]. If fragments are emitted from the interior of the source the contributions to the average kinetic energy from the Coulomb repulsion and collective expansion are substantially reduced from surface emission predictions of the standard EES model. Assuming uniform expansion and density, the mean of squared collective velocity is only 0.6 times its value at the surface, while the Coulomb energy (relative to infinity) is only 0.6 times the Coulomb energy of a particle at the surface. We have calculated average kinetic energy values assuming these kinematic modifications for those fragments emitted when the density of the source has been reduced to less than 0.3 normal nuclear density where volume formation of fragments should be important [Fri89a].

At $E/A = 110$ MeV, the experimental data shown in Fig. 3(c) indicate a weaker Z dependence than the standard EES calculations. The calculations which incorporate volume emission are more consistent with the dependence exhibited by the data, suggesting the onset of volume emission. These findings suggest the intriguing possibility that transverse energy measurements might be used to probe the onset of volume emission at low density.

In conclusion, we have systematically studied two aspects of the transverse kinetic energy spectra of the IMFs produced in central collisions: the asymptotic slopes, and $\langle E_t \rangle$. These features of the transverse kinetic energy distribution are fairly insensitive to preequilibrium emission, as well as, the linear velocity of the emitting source. They also provide a direct link for comparison with models involving equilibrated sources which have isotropic emission of fragments.

The tails of the transverse kinetic energy spectra provide evidence for emission from common equilibrated sources, whose maximum excitation increases with incident energy. Comparison of the experimental data for $\langle E_t \rangle$ with the EES model yielded

insight into the expansion aspect of the emission process. A model incorporating expansion during emission, appears to be required if one is to understand the general features of the data.

While the predictions by EES calculations are generally consistent with the kinetic information presented here, the onset of volume emission is suggested at the highest incident energy.

We would like to acknowledge the staff and operating personnel of the K1200 cyclotron at MSU-NSCL for providing the high quality beams which made this experiment possible. This work was supported by the US Department of Energy under DE-FG02-92ER40714 and the National Science Foundation under Grant Nos. PHY-89-13815, PHY-90-15957, and PHY-90-15255.

References

- [Bau87a] W. Bauer, G.F. Bertsch, and S. Das Gupta, Phys. Rev. Lett. **58**, 863 (1987).
- [deS92a] R.T. de Souza et al., Phys. Lett. **B268**, 6 (1991).
- [Bow92a] D.R. Bowman et al., Phys. Rev. Lett. **67**, 1527 (1992).
- [Ber83a] G. Bertsch and P.J. Siemens, Phys. Lett. **B126**, 9 (1983).
- [Fri90a] W.A. Friedman, Phys. Rev. C **42**, 667 (1990).
- [Aic88a] J. Aichelin et al., Phys. Rev. C **37**, 2451 (1988).
- [Sag85a] H. Sagawa and G.F. Bertsch, Phys. Lett. **B155**, 11 (1985).
- [Sch84a] H. Schulz et al., Phys. Lett. **B147**, 17 (1984).
- [deS90a] R.T. de Souza et al., Nucl. Instr. and Meth. **A295**, 109 (1990).
- [Cav90a] C. Cavata et al., Phys. Rev. C **42**, 1760 (1990).
- [Fri83a] W.A. Friedman and W.G. Lynch, Phys. Rev. C **28**, 16 (1983);
W.A. Friedman and W.G. Lynch, Phys. Rev. C **28**, 950 (1983).
- [Boa88a] D.H. Boal and J.N. Glosli, Phys. Rev. C **37**, 91 (1988).
- [Fri89a] W.A. Friedman in Nuclear Dynamics and Nuclear Disassembly, edited by J.B. Natowitz, World Scientific, Singapore, 1989, pp. 349-364
- [Gro90a] D.H.E. Gross, Rep. Prog. Phys. **53**, 605 (1990).

Figure Captions

Fig. 1: (a) Charged particle multiplicity, N_c , distributions for $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50$, 80, and 110 MeV. Multiplicity cuts corresponding to central collisions are designated with arrows. (b) Transverse kinetic energy distributions, dN/dE_{trans} , of Boron fragments for each of the three incident energies gated on the range of N_c for central collisions shown in panel (a).

Fig. 2: (a) Extracted slopes from the transverse kinetic energy distributions for $3 \leq Z \leq 9$ for central collisions at $E/A = 50, 80$, and 110 MeV. (b) Dependence of $\langle E_t \rangle$ on fragment charge.

Fig. 3: Comparison of measured values of $\langle E_t \rangle$ with the predictions of the EES model. Calculations correspond to a variety of initial source sizes and initial expansion energies as indicated in the figure. All calculations shown are for $K = 200$. Initial temperatures of 12, 15, and 18 MeV were selected for $E/A = 50, 80$ and 110 MeV respectively.

Figure 1

Figure 2

$^{36}\text{Ar} + ^{197}\text{Au}: K=200$

Figure 3

Development of Low Threshold High Resolution Detectors

J. Moore, R.T. de Souza, and D. Fox

Department of Chemistry and
Indiana University Cyclotron Facility

I. Introduction

Investigation of multifragment decay requires exclusive measurements with good phase space coverage. Recently, a new generation of electronic 4π detectors have been developed [1,2]. One of the most recent 4π arrays completed is the MSU Miniball which utilizes a fast plastic-CsI(Tl) phoswich design [3]. Nevertheless, despite the low identification thresholds of the Miniball ≈ 2 MeV/u for $Z=3$, ≈ 3 MeV/u for $Z=10$ and ≈ 4 MeV/u for $Z=18$, for the most violent central collisions a significant fraction of the fragment cross-section falls below these identification thresholds. Moreover, difficulty in calibrating the phoswich detectors, especially for low energy particles suggest the usefulness of supplementing the Miniball with linear, low threshold detector telescopes. During the past year, ten ion chamber-silicon-CsI(Tl) telescopes have been developed. These detectors are based on an earlier design by Kwiatkowski et al. [4]. The new detectors have already been used in two Miniball experiments in which they served as reference detectors for the Miniball detectors located at the same angle. In addition to being used with the Miniball, these detectors are also capable of being used in a standalone mode.

II. Physical Description

A cross-sectional view of an ion chamber telescope is presented in Fig. 1. The detector housing is constructed from stainless steel. The detector is a square based trapezoid, measuring 58 mm on each side at the base, 38 mm on each side at the top, and 154 mm tall. The detector was constructed by welding 4 pieces of 0.04 inch thick stainless steel to stainless steel flanges which form the front and back faces of the

detector. Subsequent to welding, each detector was helium leak checked to ensure its integrity. The front window consists of a 1.5 μm aluminized mylar glued to a window frame 25 mm x 25 mm. Due to its relatively small size, no support wires are used in the window design. The window seals against an 'O'-ring on the front face of the detector. Tests demonstrate that the window design is capable of sustaining a differential pressure of at least 140 torr. Vacuum tight lemo connectors mounted on the back flange of the detector provide access to all input and output signals. Gas feedthroughs for the ionization chamber are also mounted on the back flange.

The first active element of the telescope is an axial field ionization chamber directly following the window. The operating pressure of the ionization chamber 30 torr of CF_4 gas. The ion chamber has a length of 5.5 cm. The ion chamber's axial field is shaped at its edges by 7 square copper rings. These rings are 2 mm thick and are spaced 4 mm apart. The center copper ring serves as the anode of the ion chamber. Due to the weak transverse field in this design, a doubly aluminized polyethylene film with a mass of 50-100 $\mu\text{g}/\text{cm}^2$ is stretched across the anode rings. Both sides of this metallized foil are electrically shorted to the anode ring. This geometry results in efficient collection of charge with only minor energy loss for the particles traversing the foil.

The second element of the detector telescope is a 500 μm ion implanted silicon detector located directly behind the ion chamber. The active area of this detector is 28 mm x 28 mm. The silicon wafer is mounted on a square PC board frame 30 mm x 30 mm, 1.28 mm thick. The silicon detectors were purchased from Micron Semiconductor.

The third element of the detector is a CsI(Tl) scintillator crystal. The crystal is a rectangular block with a face of 25 mm x 25 mm and 30 mm thick. Due to previous experience with the sensitivity of the light output properties to variations in the thallium doping within a single crystal [3], the scintillator crystals were purchased from Hilger Analytical. The silicon detector and the CsI(Tl) crystal are spaced apart by a plexiglas spacer which provides precise positioning relative to the back flange. The CsI(Tl) crystal is wrapped with aluminized mylar on the front face and with teflon tape around the sides to provide a high degree of diffuse reflectivity. The scintillation of the crystal is read out by a 2 cm x 2 cm Hamamatsu photodiode coupled to the back face. The

photodiode signal is processed through a charge sensitive preamplifier located directly behind the photodiode. The preamplifier was fabricated at Indiana University from an existing MSU-NSCL design. To reduce sensitivity of FETs in the preamplifier to electrical discharge in the gas volume, the preamplifier was potted in silicon elastomer (Dow Corning Sylgard 184). The preamplifiers for the ion chamber and the silicon are located external to the can, approximately 25 cm away. The preamplifiers for the ionization chamber and silicon detector are a MSU-NSCL design.

III. Operating characteristics

a) Bench Tests

The detectors were bench tested with an alpha source to ensure complete charge collection in the ionization chamber and to demonstrate position independence of the collected charge.

To determine the intrinsic resolution of the ion chambers, the detectors were placed in a vacuum vessel with a collimated ^{228}Th source in front of it. Preamplifiers for the ion chamber and silicon detector were placed in the chamber connected to the detectors by less than 25 cm of RG-58 cable. The inputs to the preamplifiers were highly sensitive to pickup from the turbomolecular pump used to evacuate the chamber. This source of noise was effectively eliminated by winding the input cables to the preamplifiers around high permeability ferrite cores. The chamber was evacuated to good vacuum ($< 1 \times 10^{-5}$ torr). Preamplified energy signals from the ion chamber and silicon detector were shaped in main amplifiers and processed through standard NIM electronics. The pulse height of the two energy signals was determined in coincidence and readout by a PC based data acquisition system. The resolution of the ion chamber and silicon detector were determined at a variety of pressures (CF_4 gas) as shown in Table 1.

The change in the silicon detector resolution as a function of CF_4 gas pressure is due to energy straggling in the gas which increases with increasing pressure. At 100 torr, $\Delta E_{\text{straggling}} = 109$ keV. The ion chamber signal (based on Ziegler's program STOP) is 2.0 MeV and consequently the resolution is 168 keV (including energy straggling). Thus, the

ion chamber resolution excluding straggling is $(168^2 - 109^2)^{1/2} = 128$ keV (6.4%). The noise in the ion chamber is approximately 100 keV, thus the intrinsic resolution of the ion chamber excluding noise is $(128^2 - 100^2)^{1/2} = 80$ keV (4.0%)

The detector was also scanned with a collimated source to determine any position dependence of the charge collection. The response relative to the center was found to be $\pm 1\%$. Most of the variation was found to occur at the edges as expected.

b) In-beam tests

Elemental identification is provided in these detectors by the ΔE -E technique. This technique relies upon the fact that the rate of ionization (energy loss) is proportional (over a large range) to the atomic number of the incident ion. For the lower energy light ions as well as the heavy ions, the energy loss signal in the ion chamber provide the ΔE information. The residual energy in this case is measured by the silicon detector. If the particle is of sufficient energy to traverse the silicon detector and enter the CsI(Tl) crystal, then the same technique can be utilized between the two latter detectors. In addition, the rate of energy loss has a weak mass dependence allowing identification of mass number for light-ions punching through the silicon detector. The three element telescope can thus be used as a detector with a wide dynamic range to detect particles ranging from high energy light-ions to low velocity heavy fragments.

To determine the resolution of the detectors in separating different elements, the detectors were used in the reaction $^{86}\text{Kr} + ^{197}\text{Au}$ at $E/A = 35, 55, \text{ and } 70$ MeV. The elemental identification achieved by the combination of ion chamber with silicon detector is shown in Figure 2. Elemental bands through $Z=14$ (Si) are clearly distinguishable. The elemental and isotopic identification provided by the combination of silicon detector with CsI(Tl) crystal is shown in Figure 3. Isotopic separation for the isotopes of Carbon is achieved. Unfortunately, the isotopic identification attainable for light-ions via the ΔE -E technique is extremely sensitive to non-uniformities in the thickness of a single silicon wafer. Of an order of 10 detectors from Micron Semiconductor, only 2 detectors had sufficiently small thickness non-uniformities to allow isotopic identification via this technique.

Mass identification can also be achieved via a time-of-flight technique if a sufficiently good start signal is available (accelerator RF, etc.). This technique does not require that the particle "punch-thru" the silicon wafer. Consequently, isotopic separation can be achieved for relatively low energy particles. The present detectors have not been tested in this mode, however similar detectors [4] have achieved a time resolution of 300 ps using fast time pickoff units. This timing resolution together with the intrinsic energy resolution of the detectors results in a mass resolution of 0.3 amu when used in conjunction with an accelerator RF timing signal with ~ 200 ps resolution. Tests to determine the time resolution of the new detectors are underway.

References

- [1] G.D. Westfall, J.E. Yurkon, J. Van der Plicht, Z.M. Koenig, B.V. Jacak, R. Fox, G.M. Crawley, M.R. Maier, B.E. Hasselquist, R.S. Tickle, D. Horn, Nucl. Instr. Meth. **238**, 347 (1985)
- [2] D.G. Sarantites, L.G. Sobotka, T.M. Semkow, V. Abenante, J. Elson, J.T. Hood, Z. Li, N.G. Nicolis, D.W. Stracener, J. Valdes, and D.C. Hensley, Nucl. Instr. Meth. **A264**, 319 (1988)
- [3] R.T. de Souza, N. Carlin, Y.D. Kim, J. Ottarson, L. Phair, D.R. Bowman, C.K. Gelbke, W.G. Gong, W.G. Lynch, R.A. Pelak, T. Peterson, G. Poggi, M.B. Tsang, and H.M. Xu, Nucl. Instr. and Meth. **A295**, 109 (1990)
- [4] K. Kwiatkowski K. Komisarcik, J.L. Wile, S.J. Yennello, D.E. Fields, and V.E. Viola, Nucl. Instr. Meth. **A299**, 166 (1990)

Figure 1 : Cross-sectional view of an ion chamber telescope

Figure 2: Two dimensional spectrum of the energy deposit in the ion chamber vs. the energy deposit in the silicon detector

Figure 3: Two dimensional spectrum of the energy loss in the silicon detector versus the energy deposit in the CsI(Tl) crystal

Table 1

Silicon

Press. (torr)	Peak (ch. #)	FWHM (ch. #)	Resol. (%)
0	506.6	5.50	1.1
60	417.7	8.8	2.1
80	383.8	10.7	2.8
100	348.5	11.5	3.3

Ion Chamber

Press. (torr)	Peak (ch. #)	FWHM (ch. #)	Resol. (%)	V _{anode} (volts)
0				
60	285.7	35.7	12.5	450
80	388.2	38.7	10.0	600
100	496.3	41.7	8.4	570

Development of a Transmission Parallel Plate Avalanche Counter for Ternary Fission Studies.

A. Yonce and R.T. de Souza

Department of Chemistry and
Indiana University Cyclotron Facility
Bloomington, IN 47405

I. Introduction

Heavy nuclei can decay via fission. This decay mode involves the splitting of a nucleus into two new nuclei, referred to as fission fragments. Recent experiments indicate that occasionally a third nucleus is produced during this fission process [1]. The mass of this third nucleus lies between the mass of an alpha particle and the mass of the two major fission fragments. The third nucleus is referred to as an intermediate mass fragment ($3 \leq Z \leq \text{fission fragment}$). The fission fragments and the IMF interact via the Coulomb force. This interaction allows the IMF to act as a test particle for the Coulomb field of the fission fragments. The study of this decay mode can therefore lead to new information on the shape of this nuclear system during fission.

Although fission is a relatively long process on a nuclear time scale, on an atomic time scale it is extremely fast ($\approx 10^{-19}$ s). Therefore, direct measurement of the fission process is impossible. Fission can be induced in a heavy nucleus by bombarding that nucleus with a charged particle. The resulting fission process can be viewed as a deformation of the heavy nucleus. This deformation occurs via a series of oscillations

involving collective motion of the nucleons. The oscillations increase until the neck breaks, producing two smaller nuclei. Occasionally, an IMF is produced from the neck region during this scission process. It is unclear at this point whether or not the IMF is a group of nucleons escaping from the fissioning system (statistical emission) or is torn away when the neck breaks (dynamic ejection). Since nuclei are positively charged, the fission fragments and the IMF are repelled from each other by the mutual Coulomb field. In binary fission, the fission fragments move apart on an axis referred to as the scission axis. In the center of mass, the fission fragments are 180° apart from each other. In a three body (ternary) fission, the third body is involved in the conservation of momentum such that the fission fragments are no longer exactly 180° apart. However, due to the small mass of the IMF relative to the fission fragments, the fragments are still approximately 180° apart. Since the fission fragments are repelled in opposite directions, the components of their Coulomb force on the IMF parallel to the scission axis essentially cancel. Therefore, the IMF is propelled in a direction perpendicular to the scission axis (see Fig. 1).

The scission axis can be calculated if the positions of the two major fission fragments are known. The detection and study of these fragments is accomplished through the use of two position sensitive parallel plate avalanche counters (PPACs) and several ion chamber telescopes (see Fig. 2). The PPACs yield the position and velocity of the two major fission fragments. The ion chamber telescopes determine the atomic number (A), atomic mass (Z), angle (θ) with respect to the scission axis, and energy (E) of the IMF. The design and testing of these two detectors is the focus of this project.

II. Experimental Design

The PPAC consists of an anode foil, a cathode foil, and two planes of wires (Fig. 3). The anode and cathode foils are made up of a 2000 Angstrom thick layer of aluminum (Al) evaporated onto 1.5 μm thick mylar foils. The position-sensing wire planes are located between the anode and the cathode. The anode foil is spaced 3.1 mm from the X-plane. The X and Y wire planes are 1.55 mm apart, and the cathode foil is 3.1 mm from the Y-plane. Two strips of circuit board were used as spacers when mounting the anode and cathode to the main circuit board. Two Lexan rings with O-ring seals were attached to each side of the circuit board with 25 bolts in order to define the gas volume. The use of O-ring seals greatly increases the serviceability of the detector. The wire planes consist of 50 μm diameter Au-W wires. The X-plane contains 50 wires spaced 1.98 mm apart, and the Y-plane contains 30 wires spaced 2.54 mm apart [2]. The X-position sensing wires are perpendicular to the wires of the Y-position sensing plane and vice versa. Each wire is soldered to opposite sides of an etched copper clad epoxy-resin circuit board. Either charge division or the delay line technique can be used to attain position information with the detector. Initial testing on the PPAC was performed using both methods. The discrete resistors used in resistive charge division are inexpensive and rarely in short supply. Delay line readout, however, is inherently a faster method than charge division. Each wire is connected to either one tap of a resistive charge division line or one tap of a lumped delay line. In resistive charge division, equal valued resistors are placed between each of the sensing wires.

During operation, a fission fragment will pass into the gas volume of the detector

and strike a gas molecule. One or two electrons will become separated from the gas molecule, and will accelerate in the applied electric field. These primary electrons will then strike more gas molecules, ionizing them thus generating secondary electrons. This process is repeated resulting in an exponential growth in the number of electrons - an avalanche. As the electrons move past the position-sensing wires, a positive signal is induced on the wires. This signal travels to the resistive chain and splits [3]. A signal in proportion to the amount of resistance between its position and the end of the resistive line (Fig. 4). Comparison of the charge from the two ends of the delay line indicates the position of the fragment. The position X of the traversing particle is given by $X = L(X_R / (X_L + X_R))$ where X_L and X_R refer to the charge corresponding to the left and right ends of the resistive chain, and L is the total length of the coordinate.

Contact between the sensing wires and the resistive charge line is made through traces on the circuit board. Integrated circuit (IC) sockets were soldered to the circuit board and used to attach the resistors. In this design, switching between the two techniques simply requires replacing some IC chips on the detector which are mounted in IC sockets. Ten resistors were soldered to each 14-pin IC chip. There are five chips on the X-axis and three chips on the Y-axis. The IC sockets allow easy replacement of resistors, and the gold contact points help ensure good electrical contact. The traces, however, pose two problems. First, the dimensions of the traces act to make the PPAC sensitive to high frequency noise pickup. Secondly, the O-rings must seal over the traces, and thus the uneven surface generated by the traces represents a potential source of leaks. These problems are addressed in a proposed PPAC design in which the circuit

board is three layered. The traces in the proposed design will be on the middle layer between two pieces of copper clad circuit board. The outer layers of copperclad will provide both a smooth sealing surface as well as electrical shielding from pickup for the traces on the inner layer.

Electrical contact to the anode and cathode foils is provided by two strips of copper clad circuit board, clamped with nylon screws, to the respective electrodes. These copper clad circuit board strips are connected to connectors outside the Lexan rings via traces on the mounting circuit board. At low gas pressures and high voltage, exposed conductors are susceptible to sparking. A major source of sparking is eliminated in this design by removal of over 100 solder points from the gas volume. The placement of the electrical components outside of the active volume of the detector also allows easy access to them. Two pieces of circuit board with cutouts for the entrance of the fission fragments are seated on O-ring containing window frames bolted to the detector. A piece of 1.5 micrometer thick mylar is attached to each piece of circuit board. These windows allow the fission fragments to enter and exit the detector, making possible measurement of the fragments after passage through the detector.

Initial testing of the PPAC was accomplished with a digital multimeter. The multimeter was used to test the voltage drop between different sections of the PPAC and the continuity of the charge division lines. Further testing was carried out in a vacuum chamber with a source. The PPAC was placed inside the chamber horizontally with a ^{252}Cf spontaneous fission source underneath it. The PPAC was AC coupled via Lemo

cables through a flange in the chamber wall to the outside. The chamber and detector were then pumped down to a vacuum pressure of 1×10^{-5} Torr. Next, isobutane gas was pumped into the detector until a pressure of approximately 2-5 Torr was reached. Isobutane gas was used due to the high electron drift velocity in this gas and because a fission fragment only loses approximately 30% of its initial energy in traversing the detector. The appropriate balance between gas pressure in the detector and the applied electric field is very crucial to the correct operation of the detector. If the gas pressure is too high, relative to the voltage applied, the gas cation-electron pairs will recombine before reaching the wire planes. If the gas pressure is too low, relative to the voltage applied, the electrons will not be produced in sufficient numbers to be detected. A high voltage power source was used to apply -650 V of bias to the cathode. The anode plane was AC coupled to a high bandwidth x10 amplifier (Ortec 535) which in turn was coupled to an oscilloscope. Alpha particles and fission fragments from the source could be easily distinguished from the background noise. The alpha particles had amplitudes of 5-10 mV while the amplitudes of the fission fragments were 20-30 mV. The signals had rise times of approximately 5-8 ns (Fig. 5). Positive signals were also seen from the position sensing planes.

Tests are currently underway to investigate the performance of this detector as a function of its operating voltage and pressure. Determination of these characteristic curves is essential for a better understanding of the detectors operation. Design of a more refined PPAC with a three-layer circuit board is just beginning. Using the middle layer for traces will provide shielding from pickup during experimental conditions. Such

a configuration will also provide a substantially smoother surface for the 'O' ring seal which forms the gas-vacuum interface. A new PC based data acquisition CAMAC system which will substantially improve testing of two dimensional position resolution of the detector is presently being implemented.

REFERENCES

- [1] D.E. Fields, Ph.D. Thesis, Indiana University, Bloomington, 1992, (unpublished); preprint INC-40007-88 (1992)
- [2] Butler, M. A. et. al. Development of a Large-Area Position-Sensitive Parallel-Plate Avalanche Detector, DOE Progress Report C00-3496-79
- [3] Eyal Y. and Stelzer H., Two Dimensional Position Sensitive Transmission Parallel Plate Avalanche Counter, Nucl. Instr. Meth. **155**, 157 (1978)
- [4] Knoll G.F., Radiation Detection and Measurement, 2nd ed. John Wiley and Sons, New York. pp.227-228,348. (1989)

Figure Captions

Fig. 1: Schematic of process underlying neck emission phenomena. The essential cancellation of forces parallel to the scission axis on the particle emitted from the neck results in a kinematical focusing orthogonal to the scission axis.

Fig. 2: Proposed experimental setup for investigation of ternary fission.

Fig. 3: Cross-sectional view of the position sensitive parallel plate avalanche counter.

Fig. 4: Schematic illustrating the principle of position determination based on charge division.

Fig. 5: Illustration of the pulse amplitudes and rise times observed from the anode for alpha particles and fission fragments from ^{252}Cf .

Tertiary Fission

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Three-body calculations of Neck Emission of Intermediate Mass Fragments During Fission

S. Chen and R.T. de Souza

Indiana University Cyclotron Facility and

Department of Chemistry

Bloomington, IN 47405

For heavy systems, particularly at high spin, binary fission is an important de-excitation channel. Excited nuclear matter can also decay by statistical emission of neutrons and light charged particles. For sufficiently high excitation, statistical emission of intermediate mass fragments (IMF: $3 \leq Z \leq 20$) is also a significant de-excitation channel. Since fission is an overdamped collective deformation, it is not surprising that the fission timescale at low excitation is long compared to the statistical emission timescale. Studies indicate that fission occurs as essentially the last step of the de-excitation process [1,2]. For heavy systems of sufficient angular momentum and excitation, however, the scission and statistical emission time scales may become comparable.

Emission of α particles from the neck of the scissioning system during spontaneous fission and thermal neutron-induced fission is a reasonably well established phenomenon [3,4,5]. A recent experiment with a light-ion beam has observed ternary fission where the third fragment is an IMF [6]. Characterization of this rare decay mode

is made possible by the unique signature of low energy IMF emission roughly orthogonal to the scission axis. In a recent heavy-ion experiment [7] similar experimental signatures have been attributed to ternary fission. It is presently unclear, however, whether the IMF is emitted statistically by the system close to scission, or arises from a dynamical process such as neck rupture. Investigation of this new decay mode might yield insight into the dynamics of the fission process. Dynamical aspects of the fission process should be manifested in the final state of the emitted IMF due to the sensitivity of the final state to both the initial conditions of the IMF (eg. initial velocity and emission angle) and the Coulomb field of the fissioning system. Study of three-body correlations (IMF-fission fragment-fission fragment) should allow an estimate of the relative time between emission of the IMF and fission.

In the experiment of Ref. 6, $^3\text{He}+^{232}\text{Th}$ at $E_{\text{lab}}=270$ MeV, IMF-fission fragment-fission fragment correlations were studied. The experimental setup used is shown in Fig. 1. Coincident fission fragments were detected in two silicon arrays, each with angular coverage of $\Delta\theta=46.7^\circ$, centered at $\theta_{\text{lab}}=+64^\circ$ and -96° . IMFs were detected in four low-threshold, high resolution ion chamber-Si-CsI telescopes [8]. The energy spectra of ^{10}Be , measured in coincidence with the fission fragments, at $\theta_{\text{lab}}=+100^\circ$, $+160^\circ$, -130° , and -160° are shown in Fig. 2. Comparison of the $\theta_{\text{lab}}=+160^\circ$ spectrum with the other angles reveals that this angle manifests a low energy component not present at the other angles. The detector at $\theta_{\text{lab}}=+160^\circ$ was the IMF detector most sensitive to emission normal to the scission axis defined by the two fission fragment arrays (see Fig. 1). The angular distributions of the low and high energy components of the ^{10}Be spectra relative

to the scission axis are shown in Fig. 3. The high energy component is essentially isotropic while the low energy component is peaked near 97° , with a full-width at half maximum of about 18° . When corrected for kinematic focusing due to center-of-mass motion, this angle corresponds to approximately 90° with respect to the fission separation axis [6]. Similar effects are seen for other IMFs. The low energy component is most pronounced for isotopes which reflect the N/Z of the composite system, Fig. 4. As shown in Fig. 5, relative to the high energy component of the energy spectra, the low energy component increases with increasing fragment charge, Z .

To characterize this phenomenon, it is necessary to perform a kinematically complete determination of the three-body exit channel. In a proposal to the IUCF PAC in June 1992, a measurement of the azimuthal isotropy relative to the scission axis of the fragments emitted from the neck was outlined. One expects azimuthal isotropy relative to the scission axis if the neck fragments are emitted close to the center-of-mass of the scissioning system. This in-plane to out-of-plane ratio should provide an additional test of the proposed neck emission scenario [10]. The folding-angle technique [9] will be used to examine the impact parameter and excitation energy dependence of the new decay mode.

To better understand the kinematic correlations imposed by the neck emission scenario, we have run three-body Coulomb trajectory calculations. Three spatial and three conjugate momenta for each particle predicate that there are a total of 18 degrees of freedom however, conservation of linear and angular momentum reduces the number of degrees of freedom to 12. In the center-of-mass of the fissioning system (no preferred

direction) the number of free variables is further reduced to 10. The behavior of the three charged bodies is approximated by the behavior of three point charges. The error involved in this assumption has been investigated [11] and found to be negligible. The initial calculations are two dimensional in nature--constrained to the fission plane. This simplification reduces the number of degrees of freedom to 7. The coordinate system chosen is depicted in Figure 6. This coordinate system was chosen to facilitate comparison with earlier calculations reported in the literature [12].

The seven initial dynamical variables are: (1) The initial distance D between the two fragments. The line connecting the two fission fragments is defined as the fission axis (x axis). (2) The initial velocity of the heavy fragment V_H is in the direction of the fission axis. (3) The initial velocity of the two fission fragments in the y direction is for the sake of simplicity assumed to be zero. The initial velocity of the light fragment is assumed to be $V_L = V_H R$ where R (the mass ratio) is given by m_H/m_L . Hence the total momentum of the system is not zero but is equal to the initial momentum of the third particle. The error introduced by this assumption is negligible. (4) The initial distance X_0 of the third particle from the origin of the coordinate system along the x axis. (5) The initial distance Y_0 of the third particle from the fission axis. (6) The initial velocity v_{α} of the third particle. (7) The initial angle θ_{α} between the direction of motion of the alpha particle and the direction of the light fragment.

The initial case simulated is neck emission of α particles during spontaneous fission of ^{252}Cf . Since the manifestation of the distorted Coulomb field is largest if the emitted particle has no initial velocity, initial calculations required that the α particle

have zero initial velocity. The geometry of the scissioning system was approximated by two spherical nuclei, disregarding the Coulomb force from the neck. The initial separation of the fragments was chosen to be 16 fm so as to allow 2 fm between adjacent nuclear surfaces. The initial velocity of the two nuclei was assumed to be zero. To test the sensitivity of the final state of the α particle to its initial position, the initial position was chosen randomly in a rectangle 60 fm by 18 fm centered on the spherical nuclei. This random sampling is depicted in Fig. 7.

Comparison of α particle final kinetic energies and angles, based upon the initial position of the alpha particle should allow investigation of the selectivity of neck emission. Depicted in Figs. 8 is the angular distribution of alpha particles. The angular distribution is defined relative to the scission axis. The two prominent features of the distribution are the peak centered at 90° and the rising "wings" at forward (0°) and backward angles (180°). In panel a) the angular distribution is integrated over all possible initial positions sampled in Fig. 7. In panel b), only particles with initial positions $-2 < x_0 < 2$ (neck emission) are shown. Clearly neck emission preferentially selects emission orthogonal to the scission axis. The relevant question, however, is to what degree selection of emission orthogonal to the scission axis uniquely selects the initial position of the alpha particle as originating from the neck. The α particle energy spectra both inclusive and selected on neck emission are shown in Fig. 9. The inclusive energy spectra appears to have a two component nature. Neck emission favors the higher energy emission, although the distribution is broad. This observation can be understood in the context that emission from the periphery of the rectangle shown in Fig. 7 which

is not orthogonal to the scission axis correspond to a larger distance, R , between the emitted alpha and the center of the charge distribution. Consequently, the Coulomb energy is significantly reduced. The initial positions of the particles of low energy $E < 20$ MeV as shown in Fig. 10. As expected, with the exception of particles close to the origin (which experience a negligible force) all of the low energy particles originate from the periphery of the sampled region. Perhaps a more realistic scenario is depicted in Fig. 11 where the correlation between the final kinetic energy of neck particles and their final angular position is explored. The solid line in Fig. 11 corresponds to emission restricted to an initial position within 2 fm of the nuclear surface. The dashed line corresponds to the final angular distribution peaked at approximately 90° relative to the scission axis, whereas the dotted line corresponds to all other final angles. As can be clearly seen, while the orthogonal emission is peaked at somewhat lower kinetic energies, the mean of both distributions is rather similar.

We are currently examining correlations between the energy and angular distribution of the emitted α particles and their initial positions. The effect of initial velocities for all three particle also needs to be explored. The technique we are employing should allow investigation of not only the average behavior of the observed kinematical quantities to the initial position, but also the relationship of the widths of the observed kinematical distributions to variations in the initial conditions.

References

- [1] M. Fatyga et al., Phys. Lett. **B185**, 321 (1987).
- [2] D.J. Hinde et al., Phys. Rev. **C39**, 2268 (1989).
- [3] L.W. Alvarez as reported by G. Farwell, E. Segre, and C. Wiegand, Phys. Rev. **71**, 327 (1947).
- [4] R. Vandenbosch and J.R. Huizenga, Nuclear Fission, Academic Press, New York, 1973.
- [5] C. Wagemans, The Nuclear Fission process, CRC Press Boca Raton, FL, p545ff (1991).
- [6] D.E. Fields et al., Winter Workshop on Nuclear Dynamics, Jackson Hole, Wyoming; D.E. Fields et al., preprint INC-40007-88 (1992).
- [7] J. Boger et al., Phys. Rev. **C41**, R801 (1990).
- [8] K. Kwiatkowski et al. Nucl. Instr. and Meth. **A299**, 166 (1990).
- [9] M. Fatyga et al., Phys. Rev. Lett. **58**, 2557 (1987).
- [10] M. Sowinski et al., Z Phys. **A324**, 87-95 (1986).
- [11] I. Halpern, CERN Report 1963 (unpublished).
- [12] Y. Boneh, Z. Fraenkel, and I. Nebenzahl, Phys. Rev. **156**, 1305 (1967).

Figure Captions

Fig. 1: Experimental setup used in Ref. 6. Fission fragments were detected in two arrays centered at $+64^\circ$ and -96° with angular coverage of $\Delta\theta=46.7^\circ$. IMFs were detected in four ion chamber-Si-CsI telescopes located at $+100^\circ$, $+160^\circ$, -130° , and -160° .

Fig. 2: Kinetic energy spectra of ^{10}Be fragments coincident with angle correlated fission fragments from the reaction $^3\text{He}+^{232}\text{Th}$ at $E_{\text{lab}}=270$ MeV. The spectra are shown for $\theta_{\text{lab}}=+160^\circ$, -160° , $+100^\circ$, and -130° .

Fig. 3: Angular distribution of ^{10}Be fragments relative to the scission axis. The low and high energy components of the ^{10}Be kinetic energy spectrum are shown as the open and closed symbols respectively. In panel (a) the angle of the ^{10}Be is defined relative to the lighter of the fission fragments. In panel (b) the angle is defined by the heavier of the fission fragments.

Fig. 4: Energy spectra of ^7Be , ^9Be , and ^{10}Be detected at $\theta_{\text{lab}}=+160^\circ$ and -160° in coincidence with two fission fragments.

Fig. 5: Kinetic energy spectra at $\theta_{\text{lab}}=+160^\circ$ and -160° for Li, Be, B, and C in coincidence with two fission fragments.

Fig. 6: Coordinate system used for the three body trajectory calculation. The origin for symmetric fission is the midpoint along the scission axis between the two fission fragments. The coordinates X_0, Y_0 represent the initial position of the alpha particle along and orthogonal to the scission axis. Positive x corresponds to the direction of the light fission fragment, while positive y corresponds to the position above the scission axis (as shown).

Fig. 7: Depiction of the initial positions sampled by the alpha particle in the vicinity of the scissioning system. The rectangle is sampled uniformly except for the area occupied by the two spherical nuclei.

Fig. 8: (a) Angular distribution of alpha particles relative to the scission axis integrated over all initial positions. (b) Angular distribution of alpha particles selected on initial positions originating from the region $-2 \text{ fm} < X_0 < 2 \text{ fm}$. This initial position selection dominantly selects particles originating from the neck region.

Fig. 9: (a) Distribution of alpha particle energies, E_α , integrated over all initial positions. (b) Energy distribution of alpha particles selected on particles with initial positions $-1 \text{ fm} < X_0 < 1 \text{ fm}$.

Fig. 10: Initial positions of particles which meet the requirement that their final kinetic energy is less than 20 MeV. (see Fig. 9b)

Fig. 11: Kinetic energy distributions of particles restricted to within 2 fm of the nuclear surface. The dashed and dotted lines correspond to the additional distinction between alpha particles focused orthogonal to the scission axis and alphas at other angles.

Figure 1

Figure 2

Figure 3

Figure 4

■ — — — = + 160°

○ — — — = - 160°

Energy (MeV)

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Construction of a Compact Gas Handling system

A. Sarrazine and R.T. de Souza

Indiana University Cyclotron Facility and

Department of Chemistry

Bloomington, IN 47405

Design and operation of gas detectors requires extended access to a high quality gas handling system. Limited accessibility to existing gas handling systems necessitated the construction of a new system. The features necessary in a high quality gas handling system are control and stability of both gas pressure and flow rate. Control refers to selection of the operating pressure and flow rate, while stability refers to the maintenance of that flow rate over extended periods (1-2 weeks). The operation of avalanche detectors in particular is especially sensitive to this latter feature. At an operating pressure of 2 torr, a variation of 10% in the gas pressure can result in an amplitude variation of 100%.

A schematic of the gas handling system is depicted in Figure 1. The gas handling system is 21 inches in height and 19 inches in width. The system is rack mountable in a standard 19 inch. rack. To provide a gas handling system free of long duration contamination effects, cavitation etc. the entire gas handling system is constructed from stainless steel. All seals internal to the gas handling system are Swagelock. Seals external

to the gas handling system (detector in, detector out, gas bottle, and vacuum pump) are Cajon VCO ('O'- ring). To eliminate operator error arising from the use of multiturn valves, all valves with the exception of the reference valve are two-position valves. The presence of a ballast tank minimizes the effect of pressure fluctuations on the detector.

The detectors both currently in operation as well as under construction suggested an optimal dynamic range in pressure of 100 torr. Moreover, since the gas handling system is intended to be used to supply large area thin window parallel plate avalanche counters (PPAC) which require highly controlled chamber pump-down and let-up cycles, a differential pressure transducer was selected. The pressure transducer selected was a baratron (capacitance manometer) Type 223BD supplied by MKS Laboratories. This transducer has an accuracy (non-linearity, hysteresis, and non-repeatability) of 0.3%. The temperature coefficient is 0.04 of the reading/ $^{\circ}\text{C}$. The baratron provided the most economical transducer with sufficient linearity and stability. Based on previous experience, a solenoid valve capable of a flow rate of 100 cc/min of N_2 gas was chosen. The controller for both the pressure transducer and the solenoid valve is the Type A250 Automatic Pressure/Flow Controller also supplied by MKS Laboratories.

Construction of the gas handling system has recently been completed. Leak checking of all connections with helium gas is currently underway.

GAS IN = gas cylinder
 REF = reference (scattering chamber vacuum)
 SUP = gas supply to the detector
 RET = gas return from the detector
 SV = solenoid valve
 DIF = differential pressure transducer
 FM = flowmeter
 PUMP = roughing pump

Figure 1: Schematic of a compact gas handling system

IV. PUBLICATIONS AND ACTIVITIES

Impact parameter filters for $^{36}\text{Ar} + ^{197}\text{Au}$ collisions
at $E/A=50,80$, and 110 MeV

L. Phair, D.R. Bowman, C.K. Gelbke, W.G. Gong, Y.D. Kim,
M.A. Lisa, W.G. Lynch, G.F. Peaslee, R.T. de Souza,
M.B. Tsang, and F. Zhu,

National Superconducting Cyclotron Laboratory and
Department of Physics and Astronomy
Michigan State University, East Lansing, MI

ABSTRACT

Collisions between ^{36}Ar projectile and ^{197}Au target nuclei at $E/A = 50, 80$, and 110 MeV have been studied with the MSU Miniball, a 4π phoswich array with low detection thresholds. Various impact parameter filters, based upon total charged particle multiplicity, transverse energy, mid-rapidity charge and multiplicity of emitted hydrogen nuclei, are compared and cross-correlated. The relative selectivity of each prescription is evaluated by assessing the suppression of fast particle emission at forward angles for gates placed on low impact parameters.

Fluctuations in Multifragment Decays

L. Phair, M.A. Lisa, D.R. Bowman, C.K. Gelbke,
W.G. Gong, Y.D. Kim, W.G. Lynch, G.F. Pealsee, H. Schulz,
R.T. de Souza, M.B. Tsang, and F. Zhu

National Superconducting Cyclotron Laboratory and
Department of Physics and Astronomy
Michigan State University, East Lansing, MI

ABSTRACT

An analysis of charged particle distributions is performed to search for large fluctuations and signals of intermittency in $^{36}\text{Ar} + ^{197}\text{Au}$ reactions at $E/A = 35\text{-}110$ MeV. When the effects of impact parameter averaging are reduced by appropriate cuts on the total transverse energy, charged particle multiplicity distributions narrower than Poisson distributions are observed. Standard bond-breaking percolation calculations predict overly large fluctuations, while microcanonical and evaporation calculations predict fluctuations smaller than those observed, suggesting that conservation laws prevent intermittency signals of non-trivial origin.

Mechanisms of Intermediate Mass Fragment Formation
from Threshold to $E/A = 100$ MeV

V.E. Viola, J.L. Wile, D.E. Fields, K. Kwiatkowski,
S.J. Yennello, H.M. Xu, M.B. Tsang, R.T. de Souza, E. Renshaw,
J. Pochodzalla, K.B. Morley, W.G. Lynch, W.G. Gong, C.K. Gelbke,
D.J. Fields, and N. Carlin

ABSTRACT

Excitation functions have been measured for the formation of intermediate mass fragments in the reactions of $E/A = 20$ - 100 MeV $^{14}\text{N} + {}^{\text{nat}}\text{Ag}$ and ^{197}Au . The data have been analyzed in terms of a phenomenological moving-source model based on equilibrium, non-equilibrium and projectile-like sources. Systematic features of the data and the moving-source fit parameters are examined.

Intermediate Mass Fragment Emission as a Probe of Nuclear Dynamics

D.R. Bowman, C.M. Mader, G.F. Peaslee, W. Bauer, N. Carlin,
R.T. de Souza, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa,
W.G. Lynch, L. Phair, M.B. Tsang, and C. Williams

National Superconducting Cyclotron Laboratory and
Department of Physics and Astronomy
Michigan State University, East Lansing, MI

N. Colonna, K. Hanold, M.A. McMahan, G.J. Wozniak, and L.G. Moretto
Nuclear Science Division
Lawrence Berkeley Laboratories, Berkeley, CA 94720

and
W.A. Friedman
Department of Physics
University of Wisconsin, Madison, WI

ABSTRACT

Element distributions and fragment multiplicity distributions have been measured for $E/A=50$ MeV $^{129}\text{Xe} + ^{12}\text{C}$, ^{27}Al , ^{51}V , $^{\text{nat}}\text{Cu}$, ^{89}Y , and ^{197}Au reactions using a low threshold 4π detector. Both distributions show a strong correlation with the detected charged particle multiplicity and a large degree of target independence. The measured distributions are compared with hybrid model calculations which incorporate important dynamical aspects in both the preequilibrium and statistical emission phases. Results of these calculations are in reasonable agreement with the data. However, uncertainties with regard to the compressibility of nuclear matter remain due to uncertainties in the coupling of the two models.

Binary Character of Highly Dissipative $^{209}\text{Bi} + ^{136}\text{Xe}$
Collisions at $E_{\text{lab}}/A = 28.2$ MeV

B. Lott, S.P. Baldwin, B.M. Szabo, B.M. Quednau, W.U. Schroder, and J. Toke

Department of Chemistry and
Nuclear Structure Research Laboratory
University of Rochester
Rochester, NY 14627

L.G. Sobotka, J. Bareto, R.J. Charity, L. Gallamore
D.G. Sarantites, and D.W. Stracener

Department of Chemistry
Washington University
St. Louis, MO 63130

R.T. de Souza

National Superconducting Cyclotron Laboratory
Michigan State University
East Lansing, MI

ABSTRACT

Neutrons and light charged particles have been measured over a solid angle close to 4π , in coincidence with fragments detected forward of the grazing angle, in the reaction $^{209}\text{Bi} + ^{136}\text{Xe}$ at $E_{\text{lab}}/A=28.2$ MeV. The α particle emission pattern reveals the binary, dissipative character of these events. This observation rules out definitively any fragmentation or participant-spectator process, which have been advocated to explain the properties of fragments at forward angles. The very high light particle multiplicities associated with fragments of atomic number lower than 30 indicates that these latter fragments, although fast are actually produced in highly dissipative collisions. This feature can be taken as a confirmation that the maximum excitation energy generated in this reaction is significantly lower than the available kinetic energy, as previously established.

A. Articles

"Impact parameter filters for $^{36}\text{Ar} + ^{197}\text{Au}$ collisions at $E/A=50,80$, and 110 MeV ", L. Phair, D.R. Bowman, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa, W.G. Lynch, G.F. Peaslee, R.T. de Souza, M.B. Tsang, and F. Zhu, Nucl. Phys. A (in press)

"Multifragment Emission in $^{36}\text{Ar} + ^{197}\text{Au}$ and $^{129}\text{Xe} + ^{197}\text{Au}$ collisions: Percolation Model", L. Phair, W. Bauer, D.R. Bowman, N. Carlin, R.T. de Souza, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa, W.G. Lynch, G.F. Peaslee, M.B. Tsang, C. Williams, F. Zhu, N. Colonna, K. Hanold, M.A. McMahan, G.J. Wozniak, and L.G. Moretto, Phys. Lett. B285 10 (1992)

"Mechanisms of Intermediate Mass Fragment Formation from Threshold to $E/A = 100\text{ MeV}$ ", V.E. Viola, J.L. Wile, D.E. Fields, K. Kwiatkowski, S.J. Yennello, H.M. Xu, M.B. Tsang, R.T. de Souza, E. Renshaw, J. Pochodzalla, K.B. Morley, W.G. Lynch, W.G. Gong, C.K. Gelbke, D.J. Fields, and N. Carlin, Nucl. Phys. A538 291c (1992)

"Binary Character of Highly Dissipative $^{209}\text{Bi} + ^{136}\text{Xe}$ Collisions at $E_{\text{lab}}/A = 28.2\text{ MeV}$ ", B. Lott, S.P. Baldwin, B.M. Szabo, B.M. Quednau, W.U. Schroder, J. Toke, L.G. Sobotka, J. Bareto, R.J. Charity, L. Gallamore, D.G. Sarantites, D.W. Stracener and R.T. de Souza, Phys. Rev. Lett. 68, 3141 (1992)

B. Invited Lectures

"Multifragment Emission for Highly Excited Systems Formed in the Reaction $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 35, 50, 80$, and 110 MeV .", R.T. de Souza at the Winter Workshop on Nuclear Dynamics, Jackson Hole, WY, January 1992

C. Contributed Papers at Professional Meetings

"Characterizing the Dynamics of Multifragment Decay in the Reaction $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A=50, 80$, and 110 MeV .", D. Fox, R.T. de Souza, L. Phair, D.R. Bowman, N. Carlin, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa, W.G. Lynch, G.F. Peaslee, M.B. Tsang, and F. Zhu, presented by D. Fox at the APS Meeting, Washington, DC, April 1992

"Multifragment Emission for Highly Excited Systems Formed in the Reaction $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 35, 50, 80$, and 110 MeV .", R.T. de Souza, D. Fox, L. Phair, D.R. Bowman, N. Carlin, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa, W.G. Lynch, G.F. Peaslee, M.B. Tsang, H.M. Xu, and F. Zhu, presented by R.T. de Souza at the Winter Workshop on Nuclear Dynamics, Jackson Hole, WY, January 1992

"Evidence for Expansion Effects and the onset of Volume Emission in Nuclear Collisions from Intermediate Mass Fragment Emission in $^{36}\text{Ar} + ^{197}\text{Au}$ at $E/A = 50, 80$, and 110

MeV", R.T. de Souza, D. Fox, W.A. Friedman, L. Phair, D.R. Bowman, N. Carlin, C.K. Gelbke, W.G. Gong, Y.D. Kim, M.A. Lisa, W.G. Lynch, G.F. Peaslee, M.B. Tsang, and F. Zhu, presented by D. Fox, at the Fall Meeting of the Division of Nuclear Physics (APS), Santa Fe, NM, October 1992

"Intermediate Mass Fragment Production in the Reaction $^{84}\text{Kr} + ^{197}\text{Au}$ at $E/A = 35, 55$, and 70 MeV", D. Fox, R.T. de Souza, G.F. Peaslee, D.R. Bowman, M. Chartier, J. Dinius, D. Handzy, W.C. Hsi, C.K. Gelbke, M.A. Lisa, W.G. Lynch, L. Phair, C. Schwarz, M.B. Tsang, C. Williams, R.J. Charity, L.G. Sobotka, and E. van Buren, presented by D. Fox at the Fall Meeting of the Division of Nuclear Physics (APS), Santa Fe, NM, October 1992

D. Professional Activities

R.T. de Souza

Member of the American Physical Society, Division of Nuclear Physics

Member of the American Chemical Society, Division of Nuclear Chemistry and Technology

Member of the American Association for the Advancement of Science

Member of Sigma Xi, Indiana University Chapter

Referee for Journal Articles and Funding Proposals

Member of Indiana University Cyclotron Facility Computers/Electronics Committee

Member of General Chemistry Advisory Committee, Department of Chemistry

Member of Library Committee, Department of Chemistry

V. PERSONNEL

Faculty and Professional Staff

Dr. R.T. de Souza, Assistant Professor of Chemistry

Dr. D. Fox, postdoctoral fellow^{a)}

Graduate Students

Mr. S. Chen, graduate student^{b)}

Mr. B. Davin, graduate student^{c)}

Undergraduate and Summer Students

Mr. J. Moore, undergraduate student^{d)}

Ms. A. Sarrazine, undergraduate student

Ms. A. Wozniowski, undergraduate student^{e)}

- a) Supported by the College of Arts and Sciences, Indiana University
- b) Physics, Supported by the NSF Grant No. PHY-90-15957 (9/1/92 -)
- c) Chemistry, Supported by the Department of Chemistry, Indiana University, Teaching Assistantship (10/1/92 -)
- d) Partial support by NSF Grant No. PHY-90-15957
- e) Chemistry and Physics, Participant in Honors Research program (Credit).

END

DATE
FILMED

4 / 28 / 93