50-91295-2

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2822
UC-420,427

PPPL-2822

THEORY AND APPLICATION OF MAXIMUM MAGNETIC ENERGY
IN TOROIDAL PLASMAS

BY

T.K. CHU

February 1992

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

COPIES OF THIS DOCUMENT IS UNLIMITED

PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

Theory and application of maximum magnetic
energy in toroidal plasmas

PPPL--2822

DE92 008506

by

T. K. Chu

Plasma Physics Laboratory, Princeton University
Princeton, New Jersey 08543

The magnetic energy in an inductively driven steady-state toroidal plasma is a maximum for a given rate of dissipation of energy (Poynting flux). A purely resistive steady state of the piecewise force-free configuration, however, cannot exist, as the periodic removal of the excess poloidal flux and pressure, due to heating, ruptures the static equilibrium of the partitioning rational surfaces intermittently. The rupture necessitates a plasma with a negative q'/q (as in reverse field pinches and spheromaks) to have the same α in all its force-free regions and with a positive q'/q (as in tokamaks) to have centrally peaked α 's.

PACS numbers: 52.55Fa, 41.10-j, 47.25-c, and 52.55Hc

To illustrate the notion of maximum magnetic energy in a driven, current-carrying plasma, consider several resistors (current filaments) carrying parallel current and dissipating energy at a constant rate $V_L I_p$, where V_L is the voltage of the common battery and I_p is the total current. The filaments attract and, when unconstrained, move toward one another, resulting in an ever higher magnetic energy. At the steady state (the state of no motion or, more loosely defined, no time-averaged displacement), the current distribution is a delta-function line current and the stored magnetic energy is the highest. In an inductively driven toroidal plasma, V_L is the loop voltage. If an additional constraint exists so that the distribution of the current density cannot reach the delta-function limit, this constraint sets the limit of (the highest attainable) magnetic energy.

No pressure gradient exists in a force-free region. A tokamak plasma shows a pressure gradient in the entire cross section, indicating that magnetic surfaces marking different force-free regions¹ (with different α 's) exist in some manner. If the neoclassical transport of pressure across these surfaces is slow, the plasma pressure p can be approximated as being discontinuous across the surfaces as well. The present work proposes that a tokamak plasma is piecewise force-free with its

rational surfaces being the surfaces of discontinuity, since field line reconnection^{2,3} can transport any unbalanced pressure and poloidal flux and thus restores equilibrium of the surfaces.

Plasma equilibrium is trivially satisfied in force-free regions. The static equilibrium of a surface, on which a sheet current flows,⁴ is determined by pressure balance. In cylindrical geometry, components of the surface current, determined by the discontinuity of the field across the surface, are

$$I_{s\theta} = \lim_{\Delta r \rightarrow 0} j_{s\theta} \Delta r = -(\partial B_\phi / \partial r) \Delta r = B_{\phi 1} - B_{\phi 2} \quad (1)$$

and $I_{s\phi} = B_{\theta 2} - B_{\theta 1}$. The static equilibrium is achieved when

$(p_1 - p_2) = I_{s\phi} B_{s\theta} - I_{s\theta} B_{s\phi}$. (In a toroidal plasma, add the secondary current so that $\nabla \cdot \mathbf{j}_{s, \text{total}} = 0$.) Use $B_{s\phi} = (B_{\phi 1} + B_{\phi 2})/2$ and $B_{s\theta} = (B_{\theta 1} + B_{\theta 2})/2$ to obtain

$$(p_1 - p_2) = (B_{\phi 2}^2 - B_{\phi 1}^2)/2 + (B_{\theta 2}^2 - B_{\theta 1}^2)/2. \quad (2)$$

Equation (2) can also be obtained from surface integrals (with $\mathbf{B} \cdot \mathbf{n} = 0$ on S) of the equations $\nabla p = \mathbf{j} \times \mathbf{B}$, $\nabla \times \mathbf{B} = \mathbf{j}$ and $\nabla \cdot \mathbf{B} = 0$.

The ratio of supply rates of energy to helicity to a force-free region between two surfaces is¹

$$\alpha(q_s) = \frac{\int_{S'-S''} \mathbf{E} \times \mathbf{B} \cdot \mathbf{n} dS}{2 \int_{S'-S''} \mathbf{E} \times \mathbf{A} \cdot \mathbf{n} dS}, \quad (3)$$

where each integral is the difference of two surface integrals evaluated on S' and S'' , Δr apart. Approximate j as being constant over Δr . Hence, $\alpha(q_s) \approx \alpha(r)$, where

$$\alpha(r) \approx \frac{V_L j(r)}{2 V_L B_\phi} = \frac{j(r)}{2 B_\phi}. \quad (4)$$

The constant- j approximation, however, results in a loss of information on the structure of the field within a force-free region (Ampere's law requires nonuniform field in a force-free region of finite domain). Note that $\alpha_0 \equiv \alpha(0) = 1/q_0 R_0$ (an exact relation), where R_0 is the major radius. When the small variation in B_ϕ [in Eq. (2)] is neglected, $\alpha(r) \propto j(r)$. Thus, the total rate of energy dissipation (for a given rate of helicity dissipation) over the radius a becomes

$$V_L I_p = V_L \int_0^a 2\pi r j(r) dr. \quad (5)$$

The problem is to find the maximum energy, subject to this constraint - equivalently, the constraint of constant I_p .

In each force-free region, B_ϕ and B_θ are related through the force-free condition. (B_r , if finite, is assumed negligible by comparison.) Choose B_θ as the independent parameter. Use the variational method with a Lagrange multiplier, λ' , and obtain

$$\delta W = \delta \int_0^a 2\pi r [B_\theta^2/2 + p/(\gamma - 1) - \lambda' j] dr = 0. \quad (6)$$

Next, relate p to α . With negligible radiative and neoclassical pressure transport, the sustaining influx is helicity (and the accompanying Poynting flux to dissipate this helicity supply) and the final outward flux, due to dissipation, must be a net flow. The variation of p/α is given by $\partial(p/\alpha)/\partial t + \nabla \cdot (\Gamma_{in} + \Gamma_{out}) = 0$. With $\partial(p/\alpha)/\partial t = 0$, the ratio of the fluxes governing p/α is a constant. Integrate the pressure balance equation, $(3/2)\mathbf{v} \cdot \nabla p + (5/2)p \nabla \cdot \mathbf{v} = \mathbf{E} \cdot \mathbf{j}$, where $\mathbf{v}$ is the plasma flow velocity, and obtain

$$p = -(4\alpha/5) \frac{\int_{S'-S''} \mathbf{E} \times \mathbf{A} \cdot \mathbf{n} dS}{\int_{S'-S''} \mathbf{v} \cdot \mathbf{n} dS}. \quad (7)$$

Hence, p is proportional to α . [For large neoclassical transport, as in a highly collisional plasma, or large radiative loss, Eq. (7) becomes a poor approximation.]

A contradiction in the boundary conditions arises. While the force-free configuration requires $\mathbf{B} \cdot \mathbf{n} = 0$, the pressure transport requires $\mathbf{v} \cdot \mathbf{n} \neq 0$. [The net flow is associated with the term $\int_S \mathbf{B}(\mathbf{B} \cdot \mathbf{n}) dS$ when $\mathbf{B} \cdot \mathbf{n} \neq 0$.] Thus, the two processes must occur alternately, as exemplified in the easily observed sawtooth oscillations, and there is no purely resistive steady state. The rupture removes the excess pressure and poloidal flux so that the equilibrium is restored. The validity of Eq. (7) requires the periodicity of the rupture be much shorter than the time needed for neoclassical and radiative transport to flatten the pressure gradient at a surface.

Return to the variational problem. Define Ψ by $B_\theta = d\Psi/dr$ and $\Psi' \equiv d\Psi/dr^2 = (1/q) B_\phi / R_0$. With $p \propto \alpha \propto j$, the variation, together with the boundary condition $\delta\Psi = 0$, requires $j \propto e^{\Psi/\lambda}$, where λ is a constant (Lagrange multiplier). Use this j to solve for Ψ and obtain

$$j/j_0 = p/p_0 = \frac{1}{[1 + (q_a/q_0 - 1)(r^2/a^2)]^2} \quad (8)$$

and $(q_0/q)^2 = j/j_0$. The profile is continuous in rational number space, hence coarse-grained in physical space, as observed.⁵⁻⁷ Equation (8) has been obtained by several authors,⁸⁻¹⁰ but based

on the concept of minimum energy, not that of maximum energy in a driven plasma. Also, for that reason, the condition governing q_0 could not be formulated.

In the limit of $q_0 \rightarrow 0$, j in Eq. (8) approaches a delta-function line current at the magnetic axis. The constraint that limits q_0 to a nonzero value is $\partial p / \partial r = 0$ at $r = 0$. Mathematically, the result can be found from the general force-free configuration that includes the magnetic axis. The toroidal force-free configuration associated with the $q_s = 1$ surface for plasmas having a circular cross section has been obtained by Hastie¹¹ (to the order a^3/R_0^3):

$$q \approx q_0 [1 + (r^2/R_0^2) (29/32 - 1/2 q_0^2)]. \quad (9)$$

[Eq. (9) is applicable to the region $r < r_1$ if the region has a constant α . However, this is not the case. Since $r_1/R_0 \sim 0.1$ in tokamak plasmas, Eq. (9) gives $q_0 \approx .996$, much higher than measured values.] As $r \rightarrow 0$, it gives the desired result. The lowest q_0 below which a monotonically increasing q profile (as in tokamak plasmas) is no longer possible is $q_0 = .743$. The profile in Eq. (8) is thus fully determined.

The following are some consequences of the profile.

1. The profile of dissipation. Since $\alpha(r) \approx j(r)/2B_\phi$, the

rate of dissipation of energy varies as $j(r)$ (while that of helicity as B_ϕ , a constant). The heating rate increases with decreasing r ; the static equilibrium of a surface cannot be maintained.

2. The boundary condition for p . Apply Eq. (2) to the outermost magnetic surface. When p_1 is raised, p_2 also must be raised. Hence, a plasma vacuum ($p_2 = 0$) cannot support an equilibrium.

3. Position of the $q_s = 1$ surface. Set $q = 1$ for $r = r_1$ and $q_0 = .743$ in Eq. (8) and obtain

$$r_1/a = \frac{0.51}{(q_a - .743)^{1/2}}. \quad (10)$$

This cylindrical-geometry result agrees with measured inversion radii of chordal soft X-ray emissions of sawtooth oscillations, r_{inv} , in TFTR¹² (Tokamak Fusion Test Reactor) and TEXT¹³ (Texas Tokamak) plasmas having $q_a \leq 4$, and up to ~60% higher than measured r_{inv} for increasingly higher q_a . (The tokamak at Fontenay-aux-Roses, TFR, shows that r_1 is up to 1.7 times of r_{inv} .¹⁴) It also apparently agrees with the TFR results,¹⁵ described by the empirical formula $r_{inv}/a = 0.5f(q_0, q_a)/q_a^{1/2}$, with f being between .5 and 1.

4. The origin of sawtooth oscillations and concept of an instability. After an internal disruption, j_0 once more

increases in time and $q_0(t) = q_0(0) + (dq_0/dt)t$. The inductance change in the region $r < r_1$ gives $dq_0/dt = -4V_{ind}/\mu_0 I_1 R_0 f_1(q_0)$, where V_{ind} is the inductive part of the loop voltage at $r = 0$, $I_1 = 2\pi r_1^2 B_\phi / \mu_0 R_0$, and $f_1(q_0) = [\ln q_0^2 + 2(1 - q_0) + (1 - q_0)^2 / q_0] / (1 - q_0)^3$. The time required for q_0 to attain the value .743 from $q_0(0)$ is (approximately) the sawtooth period. Sawtooth oscillations are driven by the toroidal electric field which pinches the current channel till the limit $q_0 = .743$. Their period increases with a lowered loop voltage. They are suppressed (disappear) when the voltage becomes zero.

Below the limit $q_0 = .743$, toroidal symmetry of the magnetic axis no longer can be maintained; a helical displacement and plasma motion is generally observed. In the past, a plasma is thought to become unstable when it no longer can maintain a minimum-energy equilibrium. The present theorem inevitably advances the concept that an instability in a plasma, which is driven, is a symmetry-breaking expression in the process of attaining a higher energy as dictated by the driving potential. An ever higher energy can be attained by breaking more symmetry constraints. [A simple one-dimension mechanical analog is a violin string which also undergoes sawtooth oscillations when bowed unidirectionally. In a sawtooth cycle, the string, after the loss of static equilibrium in its slipping from the hair,

continues on to attain higher potential energy (till the ultimate snapback).] Intuitively, one may also expect a correspondence between the structure of the broken symmetry and the throughput from the source and the energy in the plasma.

5. j profile. Figure 1 shows a comparison of measured j profiles in the TEXTOR tokamak⁵ with Eq. (8) (solid curve). The profile shapes agree. $q_0 = .743$ also results in $j_0 / (B_\phi / R_0) = 2.69$; the measured value is $2.62 \pm .20$ (Fig. 6, Ref. 5).

6. p profile. The local scale length of the pressure gradient,¹⁶ $L_p \equiv -p / \nabla p$, can be used to characterize a pressure profile and to bypass its normalization by p_0 . Figure 2 shows the $\bar{a} / L_p$ profile constructed from measured T_e and n_e of the H-mode plasma (in which the effective collision frequency ν_e^* in the edge region is much less than unity¹⁶) produced by electron cyclotron heating in the D-III D tokamak.¹⁷ In the experiment T_i is a fraction of T_e and Z_{eff} (≈ 2.5) is nearly uniform in the cross section, justifying the use of $L_p \approx L_{pe}$. The rise of $\bar{a} / L_p$ at the edge is due to the drop of T_e and n_e beyond the edge. The shape of the two pressure profiles agrees, but in the sawtooth core the measured values are trending towards values that are increasingly lower than those from Eq. (8).

Comparison can also be made in L-mode plasmas in sawtooth-

suppression experiments with lower hybrid waves¹⁸ in the ASDEX tokamak.⁷ Figure 3, curve *a*, shows the a/L_p profile for a plasma in which sawtooth oscillations are suppressed during the wave ($t_{rf}=0.5$ sec, $P_{rf}=1.1$ MW) and only a small residue of the $m=1$ mode is left;¹⁸ and V_L at the plasma surface is finite. This profile agrees with that of Eq. (8), unlabelled curve, for $r/a < 0.6$. The Ohmically heated sawtooth plasma, curve *b*, does not. In both cases, a/L_p rises to ~ 16 at the edge. (The Ohmically heated and the L-mode profiles of the DIII-D experiment¹⁷ are similar to curve *b* of Fig. 3.) Thus, when compared to Eq. (8), p in the edge of an L-mode plasma is lower because of the high collisional and radiative transport; V_p in the core is lower because of the sawtooth oscillations. [Let τ_{q_0} be the time for q_0 to be lowered by $\Delta q_0 = q_0(0) - 0.743$ after an internal disruption, and τ_{p_0} be that for p_0 to increase, due to heating, to the corresponding limit according to Eq. (8). $\tau_{q_0}/\tau_{p_0} \approx (2/3)(\Delta q_0/q_0)/(1-q_0^2) \approx 0.2$. Hence, after an internal disruption, p_0 has not reached its corresponding force-free limit when q_0 has reached its limit.]

7. The bifurcation of a toroidal plasma. Apply Eq. (9) to an RFP plasma and obtain

$$q_a/q_0 \approx 1 - \frac{(\alpha_0 a)^2}{2}. \quad (11)$$

[Alternatively, from Eq. (8) obtain $q_a/q_0 = 1 + a^2 q''/2q_0$ ($q_a < q_0$ for negative q'') and use q'' from Eq. (9) to obtain Eq. (11).] It agrees with measurements,¹⁹ as shown in Fig. 4 [the so-called $F(= q_a/q_0) - \theta(= \alpha_0 a = a/q_0 R_0)$ diagram]. Field reversal occurs at $\alpha_0 a = \sqrt{2}$.

Both tokamak and RFP plasmas have more than one rational surface - both configurations are piecewise force-free. Why centrally peaked α in a tokamak plasma requires monotonically increasing q , while an RFP plasma, with decreasing q , has the same α in all its force-free regions? Because of heating, maintaining a nonuniform- α profile requires, from time to time, a piecewise net removal - "relaxation" - of excess poloidal field and pressure from an inner force-free region to the adjacent outer one so that equilibrium of the separating surface can be restored. At a rational surface q_s , this removal is accomplished by a reconnection of the field lines. Define Ψ_h by $B_h = d\Psi_h/dr$, where the helical field, B_h , at a distance Δr away from the q_s , is given by $B_h = B_\theta - nqB_\phi/m \approx -B_\theta(q'/q)_s \Delta r$. During the reconnection,

$$\partial B_h / \partial t \approx -B_\theta (q'/q)_s V_r. \quad (12)$$

where V_r is the velocity at which the helical flux (and therefore the poloidal flux) is carried by the plasma towards the X-point where it is to be reconnected. V_r is positive (pointing outward) for the inner force-free region and negative for the neighboring outer one. For $q > 0$, any unbalanced interior poloidal flux is removable only for positive q' . Hence, $q_0 = .743$ marks the bifurcation of a toroidal plasma: because of surface rupture, those with monotonically decreasing q (and hence $q_0 < .743$) have the same α in all its force-free regions and with monotonically increasing q (and hence $q_0 > .743$) have centrally peaked α . For q and q' both negative, as in the edge region of an RFP plasma beyond the internal reversal point, α increases towards the interior, as observed.²⁰ The spheromak configuration, with $q > 0$ and $q' < 0$ throughout, has the same constant α in all its force-free regions.

In summary, the magnetic energy of an inductively driven steady-state toroidal plasma is a maximum for a given rate of energy dissipation (a given constant Poynting flux at the boundary). Its configuration is piecewise force-free: it consists of nested, toroidally symmetric and noninfinitesimal regions that are separately force-free. The static equilibrium of the boundary between two adjacent regions is maintained by a pressure balance across the boundary. Because of heating, the

partitioning rational surfaces rupture intermittently. A tokamak plasma has centrally peaked α 's in its piecewise force-free regions. This is because it has a positive q'/q , which allows a piecewise outward-removal of the unbalanced flux and pressure and the restoration of equilibrium of a partitioning surface. In a toroidal plasma having a negative q'/q , such a piecewise outward-removal is not possible and all its force-free regions must have the same time-averaged α .

ACKNOWLEDGEMENT. I benefited from discussions with Wolfgang Stodiek, Allan Reiman, Shoichi Yoshikawa, Martin Kruskal, and Raffi Nazikian. The permission from Dieter Murmann of the ASDEX group at the Max-Planck Institut für Plasmaphysik to use the YAG-laser scattering data for constructing Fig. 3 is gratefully acknowledged. This work is supported by U. S. Department of Energy Contract No. DE-AC02-76-CHO 3073.

References

- ¹ T. K. Chu, "*Theory of minimum dissipation of energy for the steady state*," Princeton Plasma Physics Laboratory Report PPPL-2821, December 1991 (submitted to Phys. Fluids).
- ² H. P. Furth, J. Kileen, and M. N. Rosenbluth, Phys. Fluids, **6**, 459, (1963).
- ³ B. B. Kadomtsev, Fiz. Plazmy **1**, 710 (1975) [Sov. J. Plasma Phys. **1**, 389 (1975)]; also, in *Plasma Physics and Controlled Nuclear Fusion Research 1976*, Berchtesgaden (IAEA, Vienna, 1977), Vol. 1, p. 555.
- ⁴ A. Reiman, Phys. Fluids **23**, 230 (1980); **24**, 956 (1981).
- ⁵ H. Soltiwisch, W. Stodiek, J. Manickam, and A. Schlüter, in *Plasma Physics and Controlled Nuclear Fusion Research 1986*, Kyoto (IAEA, Vienna, 1987) Vol. 1, p 263.
- ⁶ B. Grek, D. Johnson, W. Park, G. Taylor, K. McGuire, D. Mansfield, H. Park, Bull. Am. Phys. Soc. **34**, 1965 (1989).
- ⁷ T. K. Chu, H. Murmann, R. Bartiromo, F. Leuterer, F. Söldner, the ASDEX Team, the Frascati LH Team, and the Princeton LH Team, Bull. Am. Phys. Soc. **34**, 1941 (1989).
- ⁸ D. Biskamp, Comments Plasma Phys. Controlled Fusion, **10**, 165 (1986).
- ⁹ J. Hsu and M. S. Chu, Phys. Fluids **30**, 1221 (1987).
- ¹⁰ B. B. Kadomtsev, Comments Plasma Phys. Controlled Fusion,

11, 153 (1987).

¹¹ J. Hastie, Nuclear Fusion **29**, 96 (1989).

¹² M. Murakami, V. Arunasalam, J. D. Bell, M. G. Bell, M.

Bitter, W. R. Blanchard, F. Boody, N. Bretz, R. Budny, C. E.

Bush, J. D. Callen, J. L. Cecchi, S. Cohen, R. J. Colchin,

S. K. Combs, J. Coonrod, S. L. Davis, D. Dimock, H. F.

Dylla, P. E. Efthimion, L. C. Emerson, A. C. England, H. P.

Eubank, R. Fonck, E. Fredrickson, H. P. Furth, L. R.

Grisham, S. von Goeler, R. J. Goldston, B. Grek, R.

Groebner, R. J. Hawryluk, H. Hendel, K. W. Hill, D. L.

Hillis, W. Heidbrink, R. Hulse, D. Johnson, L. C. Johnson,

R. Kaita, R. Kamperschroer, S. M. Kaye, S. Kilpatrick, H.

Kugel, P. H. LaMarche, R. Little, C. H. Ma, D. Manos, D.

Mansfield, M. McCarthy, R. T. McCann, D. C. McCune, K.

McGuire, D. M. Meade, S. S. Medley, S. L. Milora, D. R.

Mikkelsen, D. Mueller, E. Nieschmidt, D. K. Owens, V. K.

Pare, H. Park, B. Prichard, A. Ramsey, D. A. Ramussen, M. H.

Redi, A. L. Roquemore, P. H. Rutherford, N. R. Sauthoff, J.

Schivell, G. L. Schmidt, S. D. Scott, S. Sesnic, M.

Schimada, J. E. Simpkins, J. Sinnis, F. Stauffer, J.

Strachan, B. Stratton, G. D. Tait, G. Taylor, C. E. Thomas,

H. H. Towner, M. Ulrickson, R. Wieland, J. B. Wilgen, M.

Williams, K-L.Wong, S. Yoshikawa, K. M. Young, M. C.

- Zarnstorff, S. Zweben, *Plasma Physics and Controlled Fusion* **28**, 17 (1986).
- ¹³ J. A. Snipes, "The Dynamics of Sawtooth Phenomena in TEXT," Ph. D. Thesis and Fusion Research Center Report No. 275, The Univ. of Texas, Austin, Texas, p.62 (1985).
- ¹⁴ M. A. Dubois, D. A. Marty, and A. Pochelon, *Nuclear Fusion* **20**, 1355 (1980).
- ¹⁵ Equipe TFR, in *Plasma Physics and Controlled Nuclear Fusion Research 1976*, Berchtesgaden (IAEA, Vienna, 1977), Vol. 1, p. 279.
- ¹⁶ T. K. Chu, "Comments on experimental results of energy confinement of tokamak plasmas," Princeton Plasma Physics Laboratory Report PPPL-2606, April 1989.
- ¹⁷ John Lohr, B. W. Stallard, R. Prater, R. T. Snider, K. H. Burrell, R. J. Groebner, D. N. Hill, Kyoko Matsuda, C. P. Moeller, T. W. Petrie, H. St. John, and T. S. Taylor, *Phys. Rev. Lett.* **60**, 2630 (1988).
- ¹⁸ S. Bernabei, R. Bell, A. Cavallo, T. K. Chu, P. Colestock, J. Felt, W. Hooke, J. Hosea, F. Jobes, E. Mazzucato, E. Meservey, R. Motley, J. A. Murphy, M. Ono, R. Pinsker, J. Stevens, S. von Goeler, and J. R. Wilson, in *Plasma Physics and Controlled Nuclear Fusion Research 1986*, Kyoto (IAEA, Vienna, 1987), Vol. 1, p. 503; T. K. Chu, R. Bell, S. Bernabei, A. Cavallo, S. Guharay, W. Hooke, J. Hosea, F.

Jobes, E. Mazzucato, D. McNeill, E. Meservey, R. Motley, J. Stevens, and S. von Goeler, Nuclear Fusion **26**, 666 (1986); T. K. Chu, Bull. Am. Phys. Soc. **31**, 1471 (1986) and Nuclear Fusion **28**, 1109 (1988).

¹⁹ H. Bodin and A. Newton, Nuclear Fusion **20**, 1255 (1980).

²⁰ Y. Ueda, N. Asakura, S. Matsuzuka, K. Yamagishi, S. Shinohara, Y. Nagayama, H. Toyama, Nuclear Fusion, **27**, 1453 (1987).

Figure Captions

Fig. 1. j/j_0 vs. $(q_a/q_0 - 1)^{1/2} r/a$, Ref. 5: the dashed curve has $q_a = 3.3$, $I_p = 476$ kA, $B_T = 2.51$ T; the dotted curve has $q_a = 3.17$, $I_p = 340$ kA, $B_T = 1.7$ T. $R_0 = 1.75$ m, $a = .46$ m. The solid curve is Eq. (8).

Fig. 2. $\bar{a}/L_p$ vs. $r/\bar{a}$ in an H-mode plasma, Ref. 17 and according to Eq. (8). The measurement used normalized flux coordinate for the D-shaped plasma, somewhat different from the $\bar{a} = \kappa^{1/2} a$ used here in Eq. (8), $\kappa = 1.8$ being the elongation. $R_0 = 1.68$ m, $a = .62$ m, $I_p = .48$ MA, $B_T = 1.07$ T, and $\bar{n}_e \approx 1.2 \times 10^{19} \text{ m}^{-3}$.

Fig. 3. a/L_p vs. r/a in L-mode plasmas, without (curve a) and with (curve b) sawtooth oscillations. The 16-point YAG-laser scattering profile is fitted with a symmetric exponential function with 4 polynomial terms. $R_0 = 1.65$ m, $a = .40$ m, $I_p = .42$ MA, $B_T = 2.8$ T, and $\bar{n}_e \approx 2.8 \times 10^{19} \text{ m}^{-3}$. The unlabelled curve is computed from Eq. (8).

Fig. 4. q_a/q_0 vs. $\alpha_0 a$ ($F-\theta$ diagram) in three RFP plasmas, compiled in Ref. 19. The curve is Eq. (11).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Prof. I. Kawakami, Hiroshima Univ., JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Research Info. Ctr., National Inst. for Fusion Science, JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. S. Tanaka, Kyoto Univ., JAPAN
Prof. M. Goosseens, Astronomisch Instituut, BELGIUM	Library, Kyoto Univ., JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Commission-Europeen, DG. XII-Fusion Prog., BELGIUM	Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	S. Mori, Technical Advisor, JAERI, JAPAN
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL	J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
Documents Office, Atomic Energy of Canada Ltd., CANADA	D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Prof. T.W. Johnston, INRS-Energie, CANADA	Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Dr. C.R. James,, Univ. of Alberta, CANADA	Dr. O. Petrus, AL I CUZA Univ., ROMANIA
Dr. P. Lukáč, Komenského Univerzita, CZECHO-SLOVAKIA	Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
The Librarian, Culham Laboratory, ENGLAND	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS	Dr. L. Stanflo, Univ. of UMEA, SWEDEN
P. Mähönen, Univ. of Helsinki, FINLAND	Library, Royal Inst. of Technology, SWEDEN
Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
Prof. E. Economou, Univ. of Crete, GREECE	Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
Ms. C. Rinzi, Univ. of Ioannina, GREECE	Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
Dr. T. Mui, Academy Bibliographic Ser., HONG KONG	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA	Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Librarian, International Center for Theo Physics, ITALY	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN	Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

END

DATE
FILMED
418192

I