Progress report

DOE/ER/13388--3

DE92 007736

DOE Grant # DE-FG02-85ER13388
 Antenna Organization in Green Photosynthetic Bacteria
 Principal Investigator: Robert E. Blankenship
 Arizona State University

Period Covered: July 1, 1985-June 30, 1987

Introduction

This project is concerned with the structure and function of the unique antenna system found in the green photosynthetic bacteria. The antenna system in these organisms is contained within a vesicle known as a chlorosome, which is attached to the cytoplasmic side of the cell membrane. Additional antenna pigments and reaction centers are contained in integral membrane proteins. Energy absorbed by the bacteriochlorophyll c (BChl c) pigments in the chlorosome is transferred via a "baseplate" array of BChl a antenna pigments into the membrane and to the reaction center. A schematic model of chlorosome structure is shown in Fig. 1 of the main body of the proposal. This project is aimed at increasing our understanding of the organization of the pigments in the chlorosome and how the antenna system functions.

Reporting Period and Personnel

This report covers a period of two years out of a three year grant. In July 1985, just as this grant was beginning, I moved my laboratory to Arizona State University. In January 1986, Dr. Daniel Brune joined my group as a senior postdoctoral associate. In August 1986, George King joined the group as an M.S. candidate, and in December 1986, Dr. Bruce Wittmershaus joined the group as a postdoctoral associate. DOE provided partial support for the two postdocs, with the balance coming from other funded projects that they spend some time working on. Mr. King was supported by DOE one semester during the school year (he was a teaching assistant the other) and this summer has devoted his entire effort to this project.

Spectroscopic Studies of BChl c Oligomers as a Model for Chlorosomes

One of the most interesting aspects of the chlorosome antenna system is that the absorbance properties of the BChl c *in vivo* are almost precisely mimicked by oligomers of BChl c in nonpolar organic solvents (Fig. 2 in Introduction). We have completed a preliminary study of the spectroscopic and photophysical properties of these oligomeric pigments in order to gain insight into the organization of pigments in the chlorosome and have submitted a manuscript for publication (preprint attached as Appendix 1). Recently we have measured circular dichroism (CD) spectra of chlorosomes and oligomers (Fig. PR1). While additional work is needed to reach a final conclusion, our present results indicate that the *in vivo* organization of the BChl c in chlorosomes is very similar to that found in the oligomer that we have been investigating. The existence of direct pigment-pigment interactions as the principal determinants of the spectroscopic properties of the antenna pigments *in vivo* strongly suggests that chlorosomes employ a different organizational principle from that found in all other photosynthetic antennas investigated to date. In these other systems, pigment-protein interactions are of dominant importance, although pigment-pigment interactions may also be involved.

Spectroscopic Measurements on Membranes and Chlorosomes

We have recently completed an initial study of the fluorescence lifetimes of the various emitting pigments in the antenna system of *Chloroflexus aurantiacus*. We have published some of these findings in the Proceedings of the VII International Photosynthesis Congress, and have submitted a manuscript for publication (preprint and reprint attached as Appendices 2 and 3). The most striking finding of this study is the extremely short excited state lifetime of the BChl c pigments, <30 ps under a variety of conditions. Because this lifetime is beyond the resolution of the single photon counting technique, we have begun a study using streak camera detection, which is capable of higher time resolution. In Fig PR2 the dashed line is the excitation function, the solid

line is the data and the theoretical fit with lifetimes of 14 ps and 258 ps. The shorter lifetime is from the BChl c that emits at 750 nm. The longer component is from BChl a that emits at 805 nm. One more series of streak camera experiments is planned in the near future, after which we plan to prepare a manuscript for publication.

Fig. PR1 CD spectra of chlorosomes (solid line) and BChl c oligomers (dashed line).

Fig. PR2 Streak camera measurement of fluorescence from chlorosomes (see text for details).

Biochemical Preparations

An attractive aspect of the chlorosome antenna system is that it has proved quite amenable to biochemical resolution. Procedures have been developed for the isolation of the chlorosome, the chlorosome-depleted membranes, the membrane-bound antenna protein and the reaction center. Lately we have concentrated on further resolution of the chlorosome, which contains 740 nm absorbing BChl c, 795 nm absorbing BChl a, as well as carotenoid. We have developed a method to quantitatively extract the carotenoid without disturbing the other pigments (Fig. 2 in Introduction). This sample will be especially important to investigate using streak camera techniques. The excitation wavelength of the laser system is 532 nm, which is largely absorbed by the carotenoid. Using the carotenoid-depleted samples we can measure fluorescence excitation spectra and lifetimes of samples where BChl c is directly excited, and compare it to normal samples where energy transfer between carotenoid and BChl c is operative. Additional experiments along these lines and also experiments on samples in which the 795 nm absorbing BChl a has been removed, are proposed for the next grant period.

Perhaps the most important development concerning preparative methods concerns our recent finding that the amount of the 795 nm absorbing BChl a and carotenoid found in the chlorosome is strongly dependent on the conditions of isolation. In particular, we have found that the use of strictly anaerobic, reducing conditions yields a sample with much more 795 nm absorbing BChl a, as well as more carotenoid. The fluorescence properties of these samples are also quite different from those of the "standard" preparation, where no special precautions have been taken to exclude oxygen. Absorption and fluorescence spectra of the new as well as the "standard" preparation are shown in Fig. PR3. We were prompted to prepare chlorosomes anaerobically by some apparently anomalous results that we obtained comparing the fluorescence spectra and lifetimes of isolated versus membrane-attached chlorosomes (see Appendix 2). In these studies we found that the isolated chlorosomes had their fluorescence considerably quenched relative to expectations based on the disruption of energy transfer into the membrane pigment proteins. It became apparent that

additional excited state quenching pathways have been opened up in the standard chlorosome preparation, presumably mediated by oxidized or degraded pigments. We now wish to study the lifetimes of chlorosomes prepared by this new technique, and compare the results to membranes prepared in a variety of ways.

An additional important discovery concerning sample integrity is that the energy transfer efficiency from BChl c to the membrane-bound BChl a pigment protein in *Chloroflexus aurantiacus* is exceptionally delicate. Even treatments considered fairly gentle, such as storing whole cells at ice temperatures for a short time can easily lower energy transfer efficiency by 50% or more. Freshly isolated whole cells have essentially 100% efficient energy transfer (see Appendix 2). This result, together with the findings on chlorosomes discussed above, underscore the critical nature of careful sample preparation and characterization. We now think that much of the as yet modest chlorosome literature will have to be reconsidered in light of these findings, and that extreme care in sample preparation is essential to progress in understanding this system.

Fig. PR3 Absorption (A) and fluorescence (B) spectra of chlorosomes prepared aerobically (solid line) and anaerobically (dashed line).

Temperature Dependence of Energy Transfer and Trapping

Chloroflexus is a thermophilic organism, and we normally grow it at 50-55°C in the laboratory. Many of our fluorescence spectra and lifetimes have been measured at room temperature or 77° K. It is therefore important to determine if this temperature difference is important to the overall picture of the antenna system. Fig. PR4 shows emission spectra taken on freshly harvested whole cells at 50°C and 10°C, as well as the temperature dependence of the emission intensities at the three peaks over the range 10 to 55°C. The emission intensity from the 750 nm and 805 nm bands assigned to the BChl c and BChl a chlorosome "baseplate" are essentially independent of temperature over this range, while the intensity of the 885 nm emission from the membrane bound B808/865 antenna species increases approximately threefold in intensity as the temperature decreases, with the midpoint of the change at about 35°C.

We tentatively interpret these results as reflecting a loss of efficiency in trapping by the reaction center as the temperature is lowered, with little effect on the efficiency of energy transfer from chlorosome into the membrane. The transition temperature and shape of the temperature effect are strongly suggestive of a membrane lipid phase transition.

An attractive explanation for these results is that at lower temperatures the membrane loses its fluidity, and diffusional processes (such as quinone movements) that are essential to the cyclic electron transport system are inhibited. Additional experiments testing this idea are planned for the future.

Fig. PR4 Fluorescence emission spectra of *Chloroflexus* whole cells (A) recorded at 10°C (solid line) and 50°C (dashed line). The temperature dependence of the emission is shown in (B).

In vitro Spectroscopy and Photophysics of BChl c

In the course of our studies on the 740 nm absorbing BChl c oligomer, we realized that there is essentially no information in the literature concerning the photophysical properties of BChl c. In particular, no measurements of fluorescence lifetimes or quantum yields have ever been reported for these pigments. We therefore undertook a short project to measure some of these parameters in collaboration with Gilbert Seely, a noted chlorophyll spectroscopist who recently moved to Arizona State University. The results are described in the preprint included as Appendix 4. The fluorescence quantum yields range from 0.1 in methanol to 0.36 in THF, and the lifetimes range from 2.7 ns in methanol to 7.6 ns in DMSO (the lifetime in THF was not measured, but would probably be slightly longer). These experiments, while not of burning current interest, are nevertheless the kind of basic reference data that will be useful in a variety of contexts. In particular, some interesting differences in behavior between BChl c and Chl a (which is chemically akin to BChl c) were observed. These results may give some additional insight into the basic photophysics of Chl a excited states.

Other Oligomers of BChl c

In addition to the monomeric pigment absorbing at 665 nm and the 740 nm absorbing BChl c oligomer, there are at least two other distinct aggregation states of BChl c that can be induced by manipulation of solvent and pigment concentration. One of these species has an absorption maximum at 683 nm and a fluorescence lifetime of 1.1 ns. We think that this species is a dimer of BChl c. Another species, observed in lower polarity solvents, has absorption maximum at 710 nm, and fluorescence maximum at 730 nm, with a lifetime of about 3 ns. Fluorescence excitation and emission spectra of the 710 nm absorbing species are shown in Fig. PR5. The large Stokes shift, wide spectral width and the abnormally large intensity of the 0-1 and 0-2 vibronic bands at 680 nm and 635 nm are clear indications that the equilibrium geometry of the excited state is substantially different from that of the ground state. We are currently uncertain as to the molecular identity of both the 683 nm and 710 nm absorbing species. One possibility is that the 683 nm and 710 nm absorbing species are both dimers, but with different geometries. One could be a head-to-head and the other a (head-to-tail) dimer. It is difficult to obtain these species in pure form, but we have been able to obtain information as to their spectroscopic properties using difference absorption spectra and fluorescence excitation spectra. These species, while apparently not of direct relevance

to the state of BChl c *in vivo* (as is the 740 nm oligomer) are nevertheless of interest in that they provide useful information about the unique behavior of BChl c, which appears to form a richer variety of self-aggregated species than any other chlorophyll-like pigment.

Fig. PR5 Fluorescence excitation (solid line) and emission (dashed line) of the 710 nm absorbing BChl c species.

Fig. PR6 Fluorescence of whole cells of *Chlorobium vibrioforme*.

Other Green Bacteria

So far we have restricted our attention to *Chloroflexus aurantiacus*, and we anticipate that the majority of our work in the near future will also utilize this organism. However, as discussed in the introduction, there are some significant differences in the antenna systems of the Chlorobiaceae compared to those found in *Chloroflexus*. Therefore, we have initiated some experiments utilizing *Chlorobium vibrioforme*, a BChl d containing member of the Chlorobiaceae. Fluorescence emission spectra of whole cells are shown in Fig. PR6. A surprising feature of these spectra is that while the BChl d absorption is, as expected, at 730 nm instead of 740 nm as for the BChl c in *Chloroflexus*, the emission that presumably arises from this pigment is found at 757 nm, compared to 750 nm for *Chloroflexus*. The apparent Stokes shift for BChl d is, therefore, substantially greater than that of BChl c. It is, however, possible that the observed fluorescence is emitted from a minor pigment that is not evident from the absorption spectrum. Additional experiments planned for this and other species of green bacteria are discussed in the Proposed Experiments section.

Fourier Transform Infrared Measurements

The BChl c oligomer proposed as a model for the BChl c in chlorosomes is linked by interactions between the Mg, the C9 carbonyl and the C2 hydroxyethyl (See Fig. 4 in Introduction). This model predicts that the vibrational frequencies of these groups should be affected in predictable ways from their values in monomeric pigments. We have therefore initiated some Fourier Transform Infrared (FTIR) experiments in BChl c oligomers and isolated chlorosomes so as to test this prediction. As discussed in the Introduction and Proposed Experiments section of the proposal, previous information from a Russian group had implicated both these groups in formation of the 740 nm absorbing BChl c oligomer, although no information has yet been as to whether or not similar effects are observed in chlorosomes. We have recently confirmed the results of Bystrova et. al. (50) on BChl c oligomers and extended these measurements to dried films of chlorosomes. Fig. PR7 shows a FTIR difference spectrum in which the IR spectrum of a sample in which the pigment has been converted to monomeric form by pyridine vapor is subtracted from that of an

untreated sample. A peak appears at 1679 cm^{-1} , precisely where the C9 carbonyl absorbs in the monomeric pigment.

These experiments provide the first direct evidence that the molecular environment of BChl c in the chlorosome is similar to that of the BChl c oligomer, and are therefore an important and exciting development. However, the results are still preliminary and we plan additional work utilizing different conditions for reversibly converting the chlorosome pigments to monomeric form and additional experiments on BChl c *in vitro* so as to map out the behavior of these groups under a range of solvents and states of aggregation.

Fig. PR7 Fourier Transform Infrared difference spectra of untreated minus pyridine treated chlorosomes.

Equipment Development

As part of this project, we have constructed a sensitive and versatile spectrophotometer for measurement of fluorescence, absorption, circular and linear dichroism that is optimized for work in the near infrared. This spectral region is largely inaccessible with commercial fluorescence instruments. Our instrument can record fully corrected fluorescence excitation and emission spectra, as well as fluorescence polarization and can also measure absorption spectra of highly scattering samples using an integrating sphere. This instrument is ideal for measurements of fluorescence quantum yields and the efficiency of excitation transfer in a wide range of samples. A block diagram of the spectrophotometer is shown in Fig. PR8. A more complete description may be found in Appendix 2.

We are continuing to upgrade the performance of this instrument, which is unique in the world in some of its capabilities, in particular the ability to measure energy transfer efficiency in highly scattering samples through use of an integrating sphere. Recently, we have added a photoelastic modulator to the instrument so that circular and linear dichroism spectra can be measured (Fig. PR1).

Fig. PR8 Block diagram of the spectrophotometer constructed in our laboratory.

Publications and Presentations

July 1, 1985 to June 30, 1987

DOE Grant # DE-FG02-85ER13388

Antenna Organization in Green Photosynthetic Bacteria

Principal Investigator: Robert E. Blankenship

Arizona State University

Papers Published

Membrane Topology and Photochemistry of the Green Photosynthetic Bacterium *Chloroflexus aurantiacus*. R. E. Blankenship and R. C. Fuller, Chapter in "Photosynthesis III: Photosynthetic Membranes and Light Harvesting Systems" (L. A. Staehelin and C. J. Arntzen, Eds.) Encyclopedia of Plant Physiology, New Series, Vol. 19, Springer Verlag, Heidelberg, 390-399 (1986).

Light Absorption and Fluorescence of BChl c in Chlorosomes from *Chloroflexus aurantiacus* and an *in vitro* Model. D. C. Brune and R. E. Blankenship, "Progress in Photosynthesis Research", J. Biggins, Ed., M. Nijhoff, Pub., Dordrecht, IV, 745-748 (1987).

Manuscripts Submitted for Publication

Antenna Organization in the Green Photosynthetic Bacterium *Chloroflexus aurantiacus*. I. Aggregated Bacteriochlorophyll c as a Model for the 740 nm-absorbing Bacteriochlorophyll c in Chlorosomes. D. C. Brune, T. Nozawa and R. E. Blankenship. Submitted to Biochemistry.

Antenna Organization in Green Photosynthetic Bacteria. II. Energy Transfer in Detached and Membrane-Bound Chlorosomes. D. C. Brune, G. H. King, A. I. Infosino, T. Steiner, M. L. W. Thewalt and R. E. Blankenship. Submitted to Biochemistry.

Fluorescence Quantum Yields and Lifetimes for Bacteriochlorophyll c. D. C. Brune, R. E. Blankenship and G. R. Seely. Submitted to Photochem. Photobiol.

Manuscripts in Preparation

Streak Camera Measurement of BChl c Fluorescence Lifetime in Chlorosomes of the Green Photosynthetic Bacterium *Chloroflexus aurantiacus*. B. P. Wittmershaus, S. Lin, R. S. Knox, D. C. Brune and R. E. Blankenship, to be submitted to FEBS Lett.

Spectroscopic Parameters of Aggregated Forms of Bacteriochlorophyll c. G. H. King, D. C. Brune and R. E. Blankenship, to be submitted to Photochem. Photobiol.

Energy Trapping and Electron Transfer in *Chloroflexus aurantiacus*. R. E. Blankenship, D. C. Brune, J. M. Freeman, G. H. King, J. T. Trost and B. P. Wittmershaus, Proceedings of EMBO Workshop on Green Photosynthetic Bacteria, Nyborg, Denmark, August 19-21, 1987.

Pigment Organization in Chlorosomes of Green Bacteria. R. E. Blankenship, D. C. Brune, G. H. King and B. P. Wittmershaus, Proceedings of Conference on Organization and Function of Photosynthetic Antennas, Freising, Germany, October 12-16, 1987.

Conference Presentations

Light Absorption and Energy Transfer by the BChl c Antenna of *Chloroflexus aurantiacus*. R. Blankenship, A. I. Infosino, D. C. Brune, T. Steiner and M. L. W. Thewalt, Fourteenth Annual American Society for Photobiology Meeting, June 22-26, 1986, Los Angeles, CA. Photochem. Photobiol. 43, 765, (1986).

Light Absorption and Fluorescence of BChl c in Chlorosomes from *Chloroflexus aurantiacus* and an *in vitro* model, D. C. Brune and R. E. Blankenship, VII Inter. Cong. on Photosynthesis, Providence, RI, Aug. 10-15, 1986.

Upcoming Invited Conference Presentations

Pigment Organization in Chlorosomes of Green Bacteria, R. E. Blankenship, Gordon Research Conference on Biochemical Aspects of Photosynthetic Membranes, Colby Sawyer College, New London, N. H., July 27-31, 1987.

Energy Trapping and Electron Transfer in *Chloroflexus aurantiacus*, R. E. Blankenship, D. C. Brune, J. M. Freeman, G. H. King, J. T. Trost and B. P. Wittmershaus, EMBO Workshop on Green Photosynthetic Bacteria, Nyborg, Denmark, August 19-21, 1987.

Pigment Organization in Chlorosomes of Green Bacteria, R. E. Blankenship, D. C. Brune, G. H. King and B. P. Wittmershaus, Conference on Organization and Function of Photosynthetic Antennas, Freising, Germany, October 12-16, 1987.

From: Jnet%"PARSON@UWALOCKE" 10-JUL-1987 17:56
To: BLANKENSHIP
Subj: Collaboration

Received: From UWALOCKE(PARSON) by ASUCP1 with RSCS id 6005
for BLANKENSHIP@ASUCP1; Fri, 10 Jul 87 17:56 MST
Date: Fri, 10 Jul 87 17:53 PDT
From: <PARSON@UWALOCKE>
Subject: Collaboration
To: BLANKENSHIP@ASUCPS

Dear Bob,

This is to confirm that I should be pleased to collaborate with you on calculations of the spectroscopic properties of your models for the chlorosome. The plan would be to start with molecular orbitals for BChl-c, ideally including the effects of electrostatic interactions at the OH-, keto, and Mg atoms. We then would incorporate these orbitals into a treatment similar to the one that we have used for the Rps. viridis reaction center. The program includes intermolecular charge transfer transitions in addition to the local Qy, Qx, Bx and By transitions. Your models are particularly interesting because they provide an opportunity to explore the application of our approach to a system that is structurally very different from the reaction center.

Best wishes, Bill Parson

The University of
Puget Sound

February 6, 1986

Dr. Robert Blankenship
Department of Chemistry
Arizona State University
Tempe, AZ 85287

Dear Bob:

I am writing to confirm our proposed plans for collaboration on the isolation and characterization of the photochemical reaction center from the recently described bacterium, Heliothrix oregonensis. I am looking forward to this joint venture and will be happy to make field trips as needed to collect large masses of material to supply both of our various experimental needs.

I will be prepared to begin the project in 1986 and expect we will continue the collaboration as long as needed to adequately analyze this photochemical system.

Sincerely,

Beverly K. Pierson
Professor and Chairman
Biology Department

BKP:ha

Arizona State University

Department of Botany/Microbiology
Tempe, Arizona 85287
602/965-3414

July 14, 1987

Dr. Robert Blankenship
Department of Chemistry
Arizona State University
Tempe, Arizona 85287

Dear Bob,

This letter is to formally acknowledge our collaboration on the energy transfer aspect of your on-going research with green photosynthetic bacteria.

My contribution would include the processing of cells in preparation for freeze fracturing, the actual freeze fracture procedure and electron microscopy of the replicas.

As you know my primary research interest for the past few years has centered on various aspects of photosynthesis in prokaryotic and eukaryotic organisms. I look forward to a productive cooperative effort with you.

Sincerely yours,

J. R. Swafford
Assistant Professor of Microbiology

Appendices

Antenna Organization in Green Photosynthetic Bacteria

Principal Investigator: Robert E. Blankenship

Arizona State University

Appendix 1: Antenna Organization in Green Photosynthetic Bacteria I. Aggregated Bacteriochlorophyll c as a Model for the 740 nm Absorbing Bacteriochlorophyll c in *Chloroflexus aurantiacus* Chlorosomes. D. C. Brune, T. Nozawa and R. E. Blankenship.

Appendix 2: Antenna Organization in Green Photosynthetic Bacteria II. Excitation Transfer in Detached and Membrane-Bound Chlorosomes from *Chloroflexus aurantiacus*. D. C. Brune, G. H. King, A. Infosino, T. Steiner, M. L. W. Thewalt and R. E. Blankenship.

Appendix 3: Light Absorption and Fluorescence of BChl c in Chlorosomes from *Chloroflexus aurantiacus* and an *in vitro* model. D. C. Brune and R. E. Blankenship.

Appendix 4: Fluorescence Quantum Yields and Lifetimes for Bacteriochlorophyll c. D. C. Brune, R. E. Blankenship and G. R. Seely.

Appendix 5: Membrane Topology and Photochemistry of the Green Photosynthetic Bacterium *Chloroflexus aurantiacus*. R. E. Blankenship and R. C. Fuller.

Reprints & Preprints removed.

END

**DATE
FILMED
4/27/92**