Progress Report for Second Grant Year

Inelastic Electron Scattering from Surfaces

DOE/ER/45083--T1

DE92 008129

Abstract

This report contains highlights of accomplishments of the past year, for the University of California, Irvine and the University of Wisconsin, Milwaukee collaboration on surface excitations, and their interactions with low energy electrons. In addition, we present a summary of future research to be carried out in the coming grant year.

Prof. S. Y. Tong
University of Wisconsin, Milwaukee

Prof. D. L. Mills
University of California, Irvine

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. Highlights of the Last Grant Period

The California-Wisconsin joint research program continues to produce exciting and leading-edge results. In the past grant year, our new results are published in *Phys. Rev. Lett.*, *Science*, *Surface Science Lett.*, etc. Mills and Tong finished an invited review article on high-resolution electron-energy-ion spectroscopy for the *Springer Series in Surface Sciences*. The editor of the prestigious Landoldt-Boernstein Volume has requested that Mills and Tong write separate chapters for its *Volume on the Physics of Solid Surfaces*.

The thrust of our research is to make quantitative analysis and predictions of the structural and dynamical properties of a clean surface or a surface with adsorbed atoms or molecules on it. Our research program is unique in the U. S. in that the theory we developed has a high degree of precision so that direct and quantitative contact can be made with the experimental measurements on the final (scattering) state of the outgoing inelastically scattered electron. While our present efforts are focused on electron energy losses through coupling to vibrational and magnetic degrees of freedom, it is the case that our theoretical program is directly relevant to measurements in a host of other important areas, such as synchrotron radiation based photoemission. In view of increased commitment to fund new synchrotron-radiation based photoemission experiments, it is anticipated that the new data will need quantitative theoretical interpretation. Our theoretical effort can provide such highly precise interpretation of new data because we directly calculate the outgoing photoexcited electron flux, which includes the excitation matrix element effect as well as final state scattering effects. The formalism we have developed and implemented, which now includes a fully relativistic description of the electron-substrate interaction, can be applied to diverse aspects of the electron spectroscopy of materials, including those with magnetic order.

Our program is a coordinated joint effort between researchers at the University of California, Irvine and the University of Wisconsin-Milwaukee. The two principal investigators have collaborated actively during the past ten years, and the joint research program, carried out in parallel with innovative experiments in the U. S. and Europe, is a leading effort in this important area. Our joint program has been responsible for a number of major advances made in recent years. The close collaboration of the two programs can be seen by tracing the methods used in the past few years. The highly precise inelastic electron scattering method was developed at Wisconsin based on the generalized Green's function approach (Tong *et al.* *Phys. Rev. B* **16**, 1459 (1977)). This approach is very versatile and is applicable to complex surface unit cells of clean and adsorbate covered systems. In the last grant period, Mills and coworkers expanded the theory to include spin-dependent diffraction. Now that the California program is working out well, Tong and coworkers will develop a spin-dependent photoemission method based partly on the California approach. This rapid expertise transfer cannot take place without the direct and strong link between the two groups.

II. Work Finished at the University of California, Irvine

a. Spin Dependent Electron Scattering from Ferromagnetic Surfaces; The ΔT Expansion

During the first grant year, we developed and implemented the first spin dependent electron scattering calculations from magnetically ordered substrates, based on *ab initio* crystal potentials (provided kindly by Prof. A. J. Freeman and his colleagues) that have been subject to no adjustment whatsoever. These calculations provided a remarkably quantitative account of spin polarized electron diffraction data (SPLEED) reported by Waller and Gradmann, over a wide range of energies. This analysis was described in last year's progress report, and in our view constitutes a major advance. We are now able, for the first time, to carry through quantitative calculations of spin dependent electron/surface interactions, in a parameter free manner (see item (b) below). Earlier work by R. Feder employed potentials with adjustable parameters, and as a consequence were often confined to selected energies. Analyses by H. L. Davis are impressive in the account they provide of spin orbit scattering from non-magnetic substrates.

The calculations, however, are quite costly in computer time, if one wishes to explore the scattering produced by various models of the magnetic moment distribution near the surface.

During the past grant year, we have developed a new perturbation theoretic approach that appears very accurate, and enables us to explore a wide range of parameter space quickly and accurately.

The approach is to carry out full calculations for the case where the magnetic moments assume their bulk magnitude and orientation everywhere, including the surface. Then if Δt_ℓ is the change in single site t matrix produced by changing moments in layer ℓ by $\Delta\mu_\ell$,

$$f = f_\infty + \sum_\ell \left(\frac{\partial f}{\partial t_\ell} \right) \Delta t_\ell$$

where f_∞ is the scattering amplitude for the "bulk moment everywhere" model.

The above scheme has been demonstrated to reproduce the results of full calculations very accurately. Furthermore, it is far faster. We can now scan a wide range of models of surface magnetic structure quickly and efficiently, as well as wide ranges of energy and angle. We have obtained superb and complete accounts of the extensive data set generated by Waller and Gradmann by this means.

Our earlier study of SPLEED data by Waller and Gradmann examined only four beam energies, while the full data set contains information on dozens of energies. We found earlier, unfortunately, that only one of the four energies chosen by us contains intensities sensitive to the magnetic moment magnitudes near the surface. We were interested in locating other "windows of sensitivity", and our new ability to perform spin polarized scattering calculations quickly has allowed us to explore sensitivity of the entire data set to moment distribution.

Our results are described in a forthcoming paper.⁽¹⁾

1) Alim Ormeci, Burl M. Hall, and D. L. Mills, "Sensitivity of SPLEED Exchange Asymmetries to Surface Magnetic Moment; The Case of Fe (110)", Phys. Rev. B (in press).

b. Theory of Spin Asymmetries in the Transmissivity of Electrons through Ultra Thin Films; The Question of the Spin Dependence of the Electron Mean Free Path in Ferromagnets

A major goal of our present program is to develop a fully quantitative theory of spin dependent electron scattering from magnetic surfaces, and surfaces of ultra thin magnetic films. Spin dependent losses and elastic scatterings are to be studied.

If our program is to be successful, and if we are to have the capability of analyzing data within models that contain few adjustable parameters, substantial improvements in the state of the art are required. Earlier workers employ crystal potentials whose spin dependent pieces are adjusted, with parameters that depend on beam energy. Our work, as described in subsection (a), utilizing excellent potentials supplied by the Freeman group, has demonstrated one may make contact with data in a fully quantitative manner with parameter free potentials, over a wide range of beam energy and angle.

There is another issue. Earlier workers have argued that a critical ingredient that must be present in the theory of spin dependent electron-surface scattering from ferromagnetic surfaces is a spin dependent mean free path. This is another source of uncertainty in the inputs required before theory can confront data.

An experimental group at Irvine headed by Prof. Herbert Hopster has claimed to measure the variation with energy, and the spin asymmetry of the electron mean free path in ferromagnetic Fe. This is done by depositing an ultra thin film of Fe (four atomic layers) on Cu(100), and photoemitting electrons from the Cu 3d bands. These must then propagate through the Fe film before reaching the detector.

We have analyzed this experiment within the framework of our spin dependent multiple scattering theory, and we find, contrary to the interpretation offered by Hopster *et al.*, it is not necessary to invoke any spin dependence in the mean free path to interpret the data. A fully quantitative account may be obtained *by using standard spin independent electron attenuation lengths known from earlier LEED analyses*. The effect observed by Hopster *et al.* instead has its origin in the spin asymmetry of the elastic forward scattering amplitude, which describes electron propagation through the ultra thin film.⁽¹⁾

These results imply that one may account for spin dependent electron-surface interactions in a fully quantitative manner, through use of (i) unadjusted potentials, and (ii) complex inner potentials (spin independent) taken from the LEED literature. This means we are now in a position to carry out quantitative studies of electron probes of magnetic surfaces and ultra thin films, within a framework where the only adjustable parameters relate to the nature of magnetism at the surface.

1) M. P. Gokhale and D. L. Mills, Phys. Rev. Letters **66**, 2251 (1991).

c. Theory of Surface Plasmon Excitation on Simple Metals, in Electron Energy Loss

In the past two years, the electron energy loss method has proved of central importance in the study of the dynamical response of electrons in the outermost atomic layers of metals. The experiments probe certain collective modes of electrons localized at surfaces, known as surface plasmons.

Recent data from Rocca and his colleagues in Genoa calls into question the standard theory of dipole excitation of plasmons. At issue is a rather complex dependence of the excitation cross section on wave vector transfer.

During the last grant year, we have completed the first quantitative study of the energy and angle variation of the surface plasmon excitation cross section, to obtain a fully quantitative account of the data. A key to our analysis is full inclusion of an interference effect in the scattering *amplitude* associated with surface plasmon excitation. As the electron approaches the crystal surface, it may excite the plasmon before striking the crystal surface, thus reflecting off at the scattered energy. Conversely, it may reflect from the surface at the incident energy, then excite the mode as it recedes from the surface, on the outgoing leg of its trajectory. The resulting excitation cross section depends on the relative *phase* as well as amplitude of the two electron surface reflection coefficients just described.

Our results are described in a paper that appeared recently.⁽¹⁾

1) Burl M. Hall and D. L. Mills, Phys. Rev. B **44**, 1202 (1991).

III. Joint Paper Between UC Irvine and UW Milwaukee

a. The Study of Surface Phonons by Electron Energy Loss Spectroscopy: Theoretical and Experimental Considerations

Mills, Tong and Black have reviewed the state-of-the-art applications of the HR-EELS technique.⁽¹⁾ The development of this important area is closely linked to the theoretical predictions made in our earlier publications based on research supported by this DOE joint program. High-resolution EELS is a powerful and versatile technique. It has already provided modern finger-printing services to organic and molecular chemists, as well as to researchers in the field of heterogeneous catalysis.

1) D. L. Mills, S. Y. Tong and J. E. Black, *Springer Series in Surface Sciences*, eds. W. Kress *et al.*, vol. **21**, 193 (1991).

IV. Work Finished at UW Milwaukee

a. Auger Electron Angular Distributions from Surfaces: Forward Focusing or Silhouettes?

We have studied the behavior of Auger and photoexcited electrons⁽¹⁾ as they are forward-scattered by atoms in the surface region. The scattering produces a forward-focusing effect, which acts as a search-light seeking out crystallographic directions inside a solid. This technique provides a rare worm's eye-view of buried atoms in the interface region. In this paper, we pointed out the difference between forward-focusing (the correct interpretation of data) and forward-shadowing, which was suggested by another research group.

1) X. D. Wang, Z. L. Han, B. P. Tonner, Y. Chen, S. Y. Tong in *Science*, vol. 248, p. 1129 (1990).

b. Surface Phonons and Structure of Epitaxial Nickel Layers on Cu(001)

New vibrational modes in epitaxially grown thin films on a Cu(001) substrate are predicted and their existence are confirmed by experimental measurements.⁽¹⁾ The new measurements and accompanying theoretical interpretation also elucidate the novel properties of these localized modes.

1) Y. Chen, S. Y. Tong, J. S. Kim, M. H. Mohamed and L. L. Kesmodel, *Phys. Rev. B* **43**, 6788 (1991).

c. Surface Acoustic Waves and Phonons

We have written a chapter⁽¹⁾ for the Landolt-Boernstein Volume on *The Physics of Solid Surfaces*. The volume is an important reference book in which the best measurements in the subject area are included and critically reviewed. Also, we provide a succinct description of the progress of the field to date.

1) A. A. Maradudin, R. F. Wallis, S. Y. Tong in *Physics of Solid Surfaces*, ed. G. Chiarotti, Landoldt-Boernstein, Berlin (1992).

d. High-Resolution Electron Energy-Loss Spectroscopy Analysis of Ag(001): Discovery of a New Surface Longitudinal Mode Using First-Principles Phonon Calculations

We introduce the first analysis of high-resolution electron energy loss spectroscopy (HREELS) of vibrational modes of Ag(001) using normal mode frequencies and displacements obtained from first-principles microscopic calculations.⁽¹⁾ The first-principles phonon calculations include the full microscopic electron screening at the surface and contain no adjustable parameters. The combined method of using eigensolutions from first-principles calculations and fully multiple scattering inelastic cross section calculations represents the most sophisticated

theoretical treatment of this kind to date. The use of accurate theoretical treatment and experimental data, taken at a range of incident energies, is responsible for the discovery of a new localized longitudinal surface phonon at $\bar{M}$. This new mode has a large cross section at specific incident energies, and this energy range is determined.

1) Y. Chen, S. Y. Tong, M. Rocca, P. Moretto, U. Valbusa, K. P. Bohnen and K. M. Ho, *Surface Sci. Lett.* **250**, L389 (1991).

e. Forward-Focusing Photoelectron Diffraction—A New Structural Tool for Surfaces and Interfaces

We present the use of forward-focusing in *electrons* and forward-shadowing in *positrons* in Cu and Co films.⁽¹⁾ This technique allows layer-by-layer growth to be monitored. The presence of forward-focusing peaks in a system is evidence that an emitting atom is buried one, two or more atomic layers below the surface of a film. The technique thus offers a view of the surface structure from the vantage point of a buried atom.

1) S. Y. Tong, Y. Chen, Hua Li and C. M. Wei, in *Surface Physics*, Gordon and Breach, London (1991).

f. Characterization of Surface Phonons on Cu(001) and Ag(001): First-Principles Phonon Calculations With Experimental and Theoretical Studies of High-Resolution Electron-Energy Loss Spectra

Parameter-free first-principles phonon calculation is combined with multiple scattering electron-energy loss spectra calculation to characterize surface phonons of Cu(001) and Ag(001).⁽¹⁾ Again, the more accurate theory leads to the discovery of two new surface modes on these surfaces, and the inelastic electron scattering cross sections of these modes are investigated. With the discovery of the new modes, and knowledge about their inelastic cross section behavior, data of He-atom scattering and inelastic electron scattering are very satisfactorily explained. A comprehensive picture is established for the surface phonons on Cu(001) and Ag(001).

1) Y. Chen, S. Y. Tong, J. S. Kim, L. L. Kesmodel, T. Rodach, K. P. Bohnen and K. M. Ho, *Phys. Rev. B* **44**, Nov. 15 (1991).

g. Bonding and Structure on Semiconductor Surfaces

We have reviewed the nature of surface bonds on semiconductor surfaces.⁽¹⁾ Bulk-terminated semiconductor surfaces are invariably unstable because of the large number of unpaired dangling bonds which result from the creation of a sharp, idealized vacuum-solid interface (i.e., an ideal surface). To reduce the number of such unpaired bonds, surface atoms often rearrange in position to form new bonds having a higher electron density per bond. In many instances, elastic stress is introduced in the surface region. The balance between electronic and stress energies in the near surface region gives rise to the array of superlattices found

on reconstructed semiconductor surfaces of different materials: e.g., (2×1) , (2×2) , $(\sqrt{3} \times \sqrt{3})R30^\circ$, (7×7) , $c(2 \times 8)$, etc.

1) S. Y. Tong, H. Huang and C. M. Wei, in *Chemistry and Physics of Solid Surfaces*, vol. 8, eds. R. Vanselow *et al.*, Springer-Verlag, Berlin, P. 395 (1990).

V. Projects for the Coming Year

A. At the University of California, Irvine

a. Spin Dependent Low Energy Electron Diffraction from Antiferromagnetic NiO; Intensity of the Half Order Beams

As remarked in our report of progress during the past grant year, we now have in hand fast, efficient and quantitative means of calculating the intensity of the elastic scattering of spin polarized electron beams from surfaces of magnetic materials. While our primary efforts during the coming year will be directed toward initiating new studies of inelastic electron scattering from magnetic degrees of freedom near the surface (see below), we also plan further studies of elastic scattering.

Over twenty years ago, Palmberg reported studies of electron diffraction from the surface of the crystal NiO, which is antiferromagnetic with Neel temperature not far above room temperature. In these experiments, carried out with unpolarized beams, Palmberg reported the observation of half order beams, which appeared below the Neel temperature. Naive considerations suggest the temperature variation of these half order beam intensities should scale as the square of the antiferromagnetic order parameter. The measurements stimulated early theoretical studies of magnetic order near surfaces, and indeed the data reported by Palmberg is consistent with the temperature variation expected for the order parameter near the surface; its surface behavior is very different than that realized in the bulk.

Over the years, these experiments have provoked considerable discussion. Some have raised questions about whether spin exchange scattering is strong enough to account for the intensity of the half order beams. We are now in a position to address this question, in a reliable and fully quantitative manner. We will begin by calculating, with realistic Ni potentials, the intensity of the half order beams, under the assumption that the onset of antiferromagnetism results only in spin ordering, and no shifts in nuclear positions. If, on the basis of this analysis, we are unable to account for the intensity of the half order beams, then we will explore the influence of shifts in nuclear positions induced by the antiferromagnetic order. We hope to decide if the phenomenon reported by Palmberg is intrinsic, or if in fact adsorbates were present and may have influenced the results.

b. Theory of Inelastic Scattering of Electrons from Surface Spin Waves on Ferromagnetic Surfaces

One major purpose of the present program is the development of quantitative theories of electron energy loss by magnetic excitations at crystal surfaces and in ultrathin magnetic films. This work is an extension to spin polarized electron spectroscopy of our successful

theories of off specular energy loss by surface phonons. The projects described in the progress report for the first grant year, and the new study of SPLEED from NiO, are laying the foundations required for our approach to this topic.

We are now ready to address the question of electron energy loss by magnetic excitations. This question will be pursued during the next grant year. Our first task will be to calculate the absolute cross section for inelastic scattering of electrons from surface spin waves, and compare this with our earlier calculations of the absolute magnitude of the off specular phonon cross sections. We shall do this first for models of Ni(100).

One ingredient we require is the spin analogue of the electron-phonon matrix element, i.e. the change in crystal potential produced by tilting of a spin away from its equilibrium position. At the time of this writing, we have completed the formulation of a theory of this basic coupling constant. We are in the stage of assembling the multiple scattering description of the incoming and outgoing electron.

We also require, in effect, the eigenvector associated with the spin wave excitation. We shall generate this from an appropriate localized spin model. The materials we wish to explore in fact are itinerant electron magnets. In parallel with our analysis of the multiple scattering formulation of spin excitation by an incoming electron, we shall develop a theory of excitations near the surface of a model itinerant electron ferromagnet. Our plans in this regard are described in the next subsection.

c. Theory of Spin Excitations Near the Surface of an Itinerant Heisenberg Ferromagnet; Stoner Excitations and Energy Loss

As mentioned in the preceding subsection, our initial studies of spin dependent losses will concentrate on surface spin waves on the surface of a localized spin (Heisenberg) ferromagnet.

However, the materials we wish to study are in fact itinerant electron magnets. In addition to spin waves, these materials contain a broad spectrum of Stoner excitations (single electron-hole excitations, accompanied by a spin flip). The Stoner excitations have in fact been observed in spin polarized electron energy loss spectroscopy, though so far no evidence for spin wave excitation has been seen in the spectra. Thus, it is quite critical to extend the calculation described in the previous subsection, to incorporate *both* the Stoner excitations and the surface spin waves in our discussion of spin fluctuations near the surface.

This requires an itinerant model of the magnetic moment bearing electrons, treated in a framework complete enough so that the collective modes (spin waves) make their appearance. We shall proceed by setting up a Hubbard model, initially for a single band of electrons, and an FCC crystal. Such a model, in the bulk, admits a ferromagnetic ground state in the appropriate parameter regime, in the bulk. Furthermore, within the framework of the random phase approximation (RPA), in the bulk one obtains a description of spin waves and of Stoner excitations. We shall apply this model, with a self consistently determined ground state, to an FCC slab with (100) surfaces.

At the time of this writing, we have generated a self consistent description of the ground state of a finite slab, to obtain useful and interesting results. For instance, we see clear evidence of enhanced surface moments, quite comparable in magnitude with those generated by full *ab initio* studies of real surfaces. In the parameter regime where the bulk is param-

agnetic, but not far from the threshold for the onset of ferromagnetism, we find the ground state of the surface to be ferromagnetic in character, a result similar to those reported by full *ab initio* calculations.

Thus, we are convinced this model, while simple, contains the correct essential physics. It will serve, for our initial studies of electron energy loss from magnetic excitations, the role of our simple force constant models of surface phonon spectra, in the initial stages of our studies of off specular electron losses from surface phonons. We did very well with such simple models, which in many of our calculations are now replaced with input information from *ab initio* theoretical studies of surface phonons.

In the next year, we will develop the dynamical theory of surface spin fluctuations in the itinerant model described above, and integrate this with the energy loss studies described in the previous subsection.

B. At the University of Wisconsin Milwaukee

In the next 12 month grant period, we shall focus on two main topics of considerable importance to the understanding of surface dynamics and the interaction of surfaces with electrons and photons. The first area is to continue the highly successful effort to incorporate first-principle phonon calculations in the multiple scattering evaluation of electron-energy loss cross sections. We shall extend the calculation and analysis to the (110) and (111) surfaces of metals and epitaxially grown ultrathin films. The first principles, phonon eigenvectors, as well as the layer-dependent potentials will be used as inputs in new multiple scattering inelastic electron cross section calculations. The resulting theoretical findings will be compared with data to study surface and resonance modes of these systems. Such studies will establish the reliability and predictive power of quantitative theory; i.e. that geometric structure and dynamical quantities predicted by theory should independently satisfy different sets of data, without adjustable parameters. The analysis on the (110) and (111) surfaces will complete our understanding of electron-phonon interactions in fcc cubic metallic systems.

The second focus area represents a major breakout into a new area of research. We shall develop a new method for spin-dependent photoelectron diffraction and use this new theory to interpret experiments on magnetic thin films. The method of spin-dependent photoelectron diffraction will be used to both study final state diffraction and **imaging**. We currently have a concentrated effort in atomic imaging using diffraction data from emitted electrons. We shall extend this theory to **spin-dependent imaging**. We anticipate new data coming out from beam lines dedicated for diffraction and microscopy. Our efforts in this area will position us well to provide much needed quantitative interpretation of new data when they become available. As we have done in the past for HREELS, we expect that our theoretical predictions will stimulate and affect the direction and progress of this new and exciting area as well as providing a much needed theoretical basis for the experimental efforts.

VI. Papers and Articles Published During The Second Grant Year

1. Alim Ormeci, Burl M. Hall and D. L. Mills, "Sensitivity of SPLEED Exchange Asymmetries to Surface Magnetic Moment; The Case of Fe(110)", Phys. Rev. B (in press).
2. M. P. Gokhale and D. L. Mills, Phys. Rev. Letters **66**, 2251 (1991).
3. Burl M. Hall and D. L. Mills, Phys. Rev. B **44**, 1202 (1991).
4. D. L. Mills, S. Y. Tong and J. E. Black, *Springer Series in Surface Sciences*, eds. W. Kress *et al.*, Vol. **21**, 193 (1991).
5. X. D. Wang, Z. L. Han, B. P. Tonner, Y. Chen, and S. Y. Tong, *Science*, **248**, 1129 (1990).
6. Y. Chen, S. Y. Tong, J. S. Kim, M. H. Mohamed, and L. L. Kesmodel, Phys. Rev. B **43**, 6788 (1991).
7. A. A. Maradudin, R. F. Wallis and S. Y. Tong, in *Physics of Solid Surfaces*, edited by G. Chiarotti, Landoldt-Boernstein, Berlin (1992).
8. Y. Chen, S. Y. Tong, M. Rocca, P. Moretto, U. Valbusa, K. P. Bohnen and K. M. Ho, Surface Sci. Lett. **250**, L389 (1991).
9. S. Y. Tong, Y. Chen, Hua Li and C. M. Wei, in *Surface Physics*, Gordon and Breach, London (1991).
10. Y. Chen, S. Y. Tong, J. S. Kim, L. L. Kesmodel, T. Rodach, K. P. Bohnen and K. M. Ho, Phys. Rev. B **44**, Nov. 15 (1991).

END

**DATE
FILMED**

3 / 24 / 92