Theory of the Dissociation Dynamics of Small Molecules
on Metal Surfaces: Finite Temperature Studies

Progress Report

for Period July 1, 1990 - June 30, 1993

Bret E. Jackson

Department of Chemistry
University of Massachusetts
Amherst, MA 01003

November, 1992

Prepared for

THE U. S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-87ER13744

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability of responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

Abstract:

The goal of this study is to gain a better understanding of metal-catalyzed reactions by examining in detail the dynamics of molecule-metal interactions. Much effort has been focused on treating the molecule quantum mechanically when necessary, and including the effects of finite surface temperature. Recently developed time dependent quantum techniques have been used to compute the dissociative sticking probability of H_2 , HD, and D_2 on Cu and Ni surfaces. All molecular degrees of freedom can now be included either quantum mechanically or classically. The dependence upon translational and internal molecular energy, the angle and site of surface impact, and the details of the molecule-metal interaction potential have been examined. Similar techniques have been used to study the Eley-Rideal mechanism for the recombinative desorption of adsorbed H and D atoms with gas phase H and D atoms. Several useful methods for coupling gas particles to the thermal vibrations of the solid have been developed and used in studies of energy transfer and sticking. The trapping of H_2 and other diatomics in weakly bound molecular precursors to dissociative adsorption is also of interest.

TECHNICAL PROGRESS REPORT

The overall objective of this project has been to develop realistic theoretical models for some of the molecule-metal interactions important in catalysis. In particular, the dissociative adsorption of diatomics on metals, the Eley-Rideal mechanism for recombinative desorption, and the sticking of molecules on surfaces have been examined in detail. These are important steps in a great many catalytic processes, and have been well studied experimentally. Our goal has been to treat the important molecular degrees of freedom, as well as the thermal vibrations of the solid, in a quantum fashion. Efficient time dependent quantum scattering techniques have been developed to do this. A wave function is used to describe the interacting molecule. It is evolved in time via the Schrödinger Equation, scattered from or reacted with the surface, and used to compute all properties of interest. Techniques have been developed for extending these time dependent approaches to finite temperature. Much progress was made during the two and a half years covered by this report, in the following three areas:

1. The primary goal of this project has been to properly model the dissociative adsorption of diatomic molecules on metal surfaces. Six degrees of freedom (DOF) are necessary to fully describe the dissociating molecule. Previous to this project period all quantum studies had been limited to two DOF, making any comparison with experiment tenuous at best. Our intent was to include and examine the effects of all variables in the problem. Several studies were made based upon a separation of these six DOF into two groups. The most important molecular variables are r , the vibration/dissociation coordinate, Z , the center of mass translation perpendicular to the surface, and θ , the polar orientation angle of the molecule relative to the surface normal. Barrier crossing, tunneling, and the splitting of the wave function occur along r and Z , and it is important to treat them quantum mechanically. θ is also important since the barriers to dissociative adsorption are strongly θ dependent, with the $\theta=90^\circ$ orientation (parallel to the surface) usually being favored. In terms of energy flow within the molecule during the collision, the variables r , Z , and θ are highly coupled. We thus represent r , Z , and θ quantum mechanically using a wave

function $\psi(r,z,\theta;t)$. Surface chemistry is very site specific. That is, the barriers to dissociation are strongly dependent upon the position of the molecule over the surface plane (X,Y), and its azimuthal orientation in this plane ϕ . However, these variables are not so strongly coupled to the quantum ones during the collision. Ignoring quantum effects, we treat X, Y, and ϕ classically. The Hamiltonian describing the evolution of $\psi(r,z,\theta;t)$ is obtained by replacing the operators corresponding to X, Y, and ϕ in the full Hamiltonian with the classical variables and their conjugate momenta. Equations of motion for the classical variables depend upon forces which are averaged over the wave function $\psi(r,z,\theta;t)$. The quantum ψ and the classical variables are then propagated simultaneously.

In our first study we examined the dissociation of H₂, HD and D₂ on a Ni(100) surface¹. The effects of rotational orientation were studied by evolving $\psi(r,z,\theta;t)$ for fixed values of X, Y, and ϕ , and comparing the results with the commonly used 2-DOF model, in which only r and Z are included. Our 3-DOF dissociation probability was much lower than for two DOF because of the inclusion of molecular orientations unfavorable for dissociation. Thus, more kinetic energy was required to dissociate, and the apparent activation energy shifted by over 0.1 eV. This underscores the danger of comparing 2-DOF models directly with experiment. The dissociative sticking probability was computed as a function of kinetic energy for several impact sites and azimuthal orientations ϕ . At lower energies the reaction became more site specific. Isotope effects seen experimentally were observed. The effects of internal molecular excitation were also examined. Vibrational excitation of the molecule increased the dissociation probability with nearly perfect efficiency, which may help explain recent studies of H₂ on Cu. Initial rotational excitation decreased the dissociation probability, perhaps due to the fact that a rotating molecule is less able to follow the minimum energy pathway.

The probabilities for rotational excitation of the unreacted molecules were also computed. A noticeable increase in rotational excitation was seen to accompany an increase in dissociation. At energies where dissociation is likely the wave function probes the barrier region of the potential, which is very anisotropic in θ , particularly when compared with non-reactive potentials. This gives rise to anomalous amounts of rotational excitation in the scattered flux, which may serve as a signature of a reactive surface.

These results are in agreement with a study by Holloway and Jackson², based on a simple analytical model.

In our most recent study³ we propagated both the quantum and classical variables, running many trajectories in order to average over all possible initial values for X , Y , and ϕ . This was the first calculation in which all molecular degrees of freedom were included, and the necessary variables were treated quantum mechanically. Two cases were considered: a highly reactive Ni surface where direct dissociation dominated at relatively low beam energies, and a weakly reactive Cu surface where high kinetic energies were required. Dissociation probabilities and scattered ro-vibrational state distributions were computed as a function of beam kinetic energy for H_2 and D_2 . The importance of including all DOF was clearly demonstrated. For example, the potential gradients parallel to the surface tended to focus the incoming molecules into the hollow sites. The lower DOF models only consider a single atop or bridge site, which may not even be accessed by the molecule if X and Y were allowed to move. Similarly, reorientation in ϕ during the reaction had a strong influence on the pathways available for dissociation. Much of the behavior of a given gas-metal system is determined by the way the reaction probability varies between the different impact sites and molecular configurations available, and cannot be described by a one barrier model. Fully classical studies were also made of these systems for comparison. The classical studies did best at high beam energies (Cu), as long as the dissociation probability was sufficiently large. Various sudden approximations which reduce the dimensionality (and difficulty) of the problem were examined. A freezing of the azimuthal orientation angle of the molecule was shown to lead to significant errors in many cases. A diffractive sudden approximation which conserves the center of molecular mass motion parallel to the surface was less severe.

Based upon these results, an number of studies are in progress. Our fully six-DOF model is being used to examine the controversial system of H_2 and D_2 on Cu⁴. This system involves low reaction probabilities, high barriers, and questions as to the role of initial vibrational excitation, all requiring a quantum treatment. This is the first such study to include all important variables. It is also finally possible for us to examine systems such as N_2 on W(100) and O_2 on Pt(111) in a complete and proper manner. Improved models are being studied. A Lanczos method for writing and

propagating the wave function has been developed⁵. Ways of treating r , Z , θ , and ϕ quantum mechanically are under consideration. This allows for a proper quantization of the rotational states as well as an improved description of azimuthal reorientation during dissociation.

2. The second focus of our work has been to include the effects of temperature in molecule-metal interactions, allowing for energy transfer between the gas particle and the vibrations of the solid. Much of our earlier work involved the development of mean field approaches which treated the scattering molecule, as well as the lattice vibrations quantum mechanically. The molecular wave function evolved on a gas-surface interaction potential which was both time and temperature dependent. This aspect of the project has now been phased out, and is currently supported elsewhere. However, during the first year of the project, a review of our earlier work appeared⁶, and two studies of sticking were submitted and published^{7,8}.

The sticking of a gas particle to a surface, in which the kinetic energy of the particle is dissipated into the vibrations of the solid, is an important first step in most catalytic reactions. A general study of gas surface energy transfer was made for He, Ne and Ar on Cu⁷. Sticking probabilities were computed as a function of energy, mass, temperature, and the shape of the gas-surface interaction potential. The sticking probability was found to decay rapidly with decreasing mass and increasing temperature and energy, as seen experimentally. Analytical expressions were derived for the sticking probability at very low temperatures and energies, where quantum effects often dominate. Trapping was found to increase strongly with the attractive well depth. A comparison was also made with the forced oscillator model, which was shown to have several deficiencies.

A detailed study was also made of how sticking and energy transfer depend upon the component of the molecular kinetic energy parallel to the surface⁸. In many experiments sticking and other properties depend only upon the normal component of the energy, while in others, there is parallel energy dependence (non-normal scaling). We found that dissipation of the parallel energy into the solid was slow. However, the corrugation could mix the normal and parallel components of the energy during a collision, leading to non-normal effects. These effects are largest for deep attractive wells,

large corrugation, low energies, high temperatures, and large angles of incidence.

With regard to this aspect of the project, we are still interested in two phenomena involving lattice vibrations. One is to understand the temperature dependence of dissociative adsorption, and it is hoped that our mean field techniques will be useful in this regard. Another is the sticking of diatomics in the weakly bound precursor states which often occur in the pathways to dissociative adsorption. We are currently examining ways to include rotational motion in our sticking models.

3. Our newest endeavor involves the study of Eley-Rideal processes, in which a gas phase molecule reacts with an adsorbed species, with the product molecule then desorbing. In collaboration with Mats Persson, we have used our time dependent quantum techniques to examine these events, which are important for a number of catalytic processes as well as the removal of halogens from semiconductor surfaces. In two papers we studied the dynamics of a gas phase H (or D) atom recombining with an adsorbed H (or D). These initial studies included only two degrees of freedom, with the two atoms in a collinear arrangement perpendicular to the surface^{9,10}. Reaction probabilities and product vibrational state distributions were computed as a function of translational energy for all isotopic combinations. Large amounts of vibrational excitation were found for this very exothermic reaction, in agreement with experiment.

A calculation including all degrees of freedom is now in progress¹¹. A flat surface approximation is made in which three of the variables are conserved. Thus, only three degrees of freedom need to be evolved quantum mechanically, and we are using techniques similar to those developed for dissociative adsorption to treat them. This allows us to get full reactive cross-sections. The code has been fully developed and a number of applications are underway. A parallel project, a collaboration with Bruce Kay which has just started, involves the removal of adsorbed halogen atoms via recombination with gas phase atomic hydrogen. Preliminary two degree of freedom studies suggest that our improved approach, or a model involving partially accommodated but "hot" H atoms may be required to explain the data.

In summary, the past two and a half years have been very productive. Our progress has been similar to that outlined in the original proposal, except that not as many applications of our six-DOF code for dissociative adsorption have been made as we would have liked. This was due primarily to the difficulty of developing a proper approach, which we have now completed. On the other hand, our Eley-Rideal work (not in the original proposal), which evolved as a natural extension of our dissociation studies, has turned out to be extremely interesting and important. Excellent collaborations with Mats Persson (Chalmers) and Bruce Kay (Battelle) have developed. A dissociative adsorption project with Ulrik Nielsen (Aarhus) is also just beginning. During the current project period this work was presented at workshops in Copenhagen and Japan. Invited talks were given at a Gordon Conference, three National ACS meetings, and an AVS meeting. Several posters and departmental seminars have been given. I have been spending about 30% of my time on this project during the semester, and about 50% during the summer. A graduate student, A. J. Cruz, was working 100% on this project. She currently has a position at the University of Humacao, in Puerto Rico, and defended her thesis in December of 1992. My postdoctoral associate S. Kumar has been working full time on this project.

The following papers, referenced above, appeared in print or were submitted during the period of this report, or are currently in progress. I fully expect to submit references 4, 5, and 11, as well as other possible papers, by the end of the project period, 6/30/93.

1. A. J. Cruz and B. Jackson, J. Chem. Phys. 94, 5715 (1991).
2. S. Holloway and B. Jackson, Chem. Phys. Lett. 172, 40 (1990).
3. A. Grüneich, A. J. Cruz and B. Jackson, J. Chem. Phys. (submitted).
4. S. Kumar and B. Jackson, (work in progress).
5. S. Kumar and B. Jackson, (work completed; to be submitted).
6. B. Jackson, Comp. Phys. Comm. 63, 154 (1991).
7. B. Jackson, J. Chem. Phys. 94, 5126 (1991).
8. B. Jackson, J. Chem. Phys. 94, 787 (1991).
9. B. Jackson and M. Persson, J. Chem. Phys. 96, 2378 (1992).
10. B. Jackson and M. Persson, Surf. Sci. 269/270, 195 (1992).
11. M. Persson and B. Jackson, (work near completion).

**DATE
FILMED**

2 / 24 / 93