CHARGE SEPARATION IN PHOTOREDOX REACTIONS

Progress Report

October, 1991 - October, 1992

Larry Kevan

Department of Chemistry

University of Houston

Houston, Texas 77204-5641

October, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY

UNDER GRANT NUMBER DE-FG05-88ER13848

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Progress Report - October 1992

DOE Grant DE-FG05-88ER13848

"Charge Separation in Photoredox Reactions"

Larry Kevan

Department of Chemistry, University of Houston, Houston, Texas 77204-5641

1. Summary of Progress

Development of the control of the location of electron donors and acceptors relative to micelle and vesicle interfaces has advanced by the utilization of electron spin resonance (ESR), deuterium electron spin echo modulation (ESEM) and matrix proton electron nuclear double resonance (ENDOR) spectroscopies.

It has been emphasized that rapidly frozen surfactant solutions are necessary to exploit the measurement of distance discrimination by deuterium ESEM and matrix proton ENDOR spectroscopy. Although a variety of previous experiments have shown that micellar and vesicular structures are retained in rapidly frozen solutions, a recent experimental study has shown that x-doxylstearic acid spin probe differential positions are similar in liquid and frozen dioctadecyldimethylammonium chloride vesicle solutions. ESR of a series of x-doxylstearic acids ($x = 5, 7, 10, 12,$ and 16) solubilized in cationic dioctadecyldimethylammonium chloride vesicles were recorded over 298 - 328 K and analyzed by employing spectral line simulation. The spectra from the vesicle phase are well reproduced by using the microscopic order-macroscopic disorder model of Freed and co-workers with Brownian rotational diffusion. The partial averaging of the magnetic interactions by local anisotropic motions in the vesicles is quantified mainly by an order parameter (S) and the rotational diffusion rate perpendicular to the alkyl chain ($R_{\perp}$). The simulations show that

the doxyl position becomes progressively less ordered (S decreases) as x increases up to $x = 12$, but becomes more ordered at $x = 16$ (S increases). This suggests a U-shaped bent conformation of the doxylstearic acid alkyl chain. $R_{\perp}$ shows an inverse correlation; that is, $R_{\perp}$ increases, plateaus, and decreases with increasing x and supports the S correlation. The U-shaped conformation of the spin probe alkyl chain in liquid vesicle solutions is the same as deduced previously in *frozen* vesicle solutions measured by electron spin echo modulation spectroscopy. There the deuterium modulation depths follow the same trend as the order parameter with the x -doxyl position and have a similar interpretation as to the spin probe conformation.

The efficiency of the photoionization process of tetramethylbenzidine (TMB) solubilized in mixed micelles of anionic-nonionic and cationic-nonionic surfactants as a function of the composition of the mixed micelle has been evaluated. It was possible to change continuously the micelle surface charge from a positive net charge to a neutral to a negative net charge without changes in the size of the micelle or phase separation, in contrast to mixed micelles formed by ionic surfactants. It is possible to separate the contributions to the photoionization yield due to the interface charge and due to the TMB⁺-water interactions. The analysis of the deuterium modulation depth arising from the interaction of the photocation produced in the photoionization process with deuteriums of the nonionic surfactant, selectively deuterated along the alkyl chain or in the polar headgroup, allowed us to monitor the location of the TMB molecule in the mixed micelle as well as changes in the structure of the mixed micelle as a function of the mixed micelle composition.

An ESR, ESE, and ENDOR investigation was performed on a series of alkylmethylviologens which were solubilized in dihexadecyl phosphate (DHP) vesicles and rapidly frozen at 77 K. Photoreduction yields were measured by

ESR, while ESE and matrix ENDOR were used to determine the location of the viologen radical in the lipid bilayer with respect to the water interface and the hydrocarbon interior, respectively. Both the photoreduction yield and the matrix proton ENDOR linewidth increase with increasing alkyl chain length, whereas the normalized deuterium modulation depth decreases. The trend in the ESEM data can be explained by the alkyl chain on the viologen causing a partial insertion of the molecule into the vesicle bilayer with the degree of insertion being greater for longer chain molecules. The ENDOR results corroborate the ESE data. The decrease in deuteron modulation depth correlates with an increase in the proton matrix ENDOR line width; that is, the proton density in the vicinity of the viologen increases as the dictation moves into the bilayer of the vesicle. Both the ESE data and the ENDOR results imply that as the length of the alkyl chain on the viologen increases, the molecule becomes more lipophilic and the viologen moiety is embedded deeper into the vesicle. Addition of cholesterol to the vesicle results in an increase in the photoreduction yield of the radical. ESEM patterns show that adding cholesterol causes increased water penetration into the interface. The ENDOR results indicate that the effective proton density in the interface region is reduced as cholesterol intercalates into the bilayer. A probable interpretation of the increase in photoreduction yield is that added cholesterol reduces the effective interface charge density which lowers the barrier for electron transfer between the chloride counterion and the alkylviologen moiety.

The photoionization yields of N-alkylphenothiazines solubilized in either cationic or anionic micelles at 77 K were studied as a function of both alkyl and surfactant chain length. The results obtained from the analysis of the ESR spectra and the ESEM patterns of photogenerated cations of phenothiazine derivatives solubilized in sodium decyl, dodecyl, and tetradecyl sulfate and in

decyl-, dodecyl-, and tetradecyltrimethylammonium bromide micellar solutions show that the photoyield of the cation generally correlates with the deuterium modulation depth. The photolysis yield and the normalized deuterium modulation depth increase as the alkyl chain on the phenothiazine is lengthened and decrease as the micelle surfactant chain length is increased. These observations support the conclusion that the photoyield is related to the strength of photoproduct cation-water interactions and that the location of the phenothiazine moiety is near the micellar interface. Phenothiazines solubilized in anionic micellar systems show a larger deuterium modulation depth compared with those solubilized in analogous cationic micelles. The photoyields are slightly larger in cationic versus anionic micelles. This increase is attributed to a lower energy barrier for a photoejected electron to transverse a cationic micelle interface. A comparison of the deuterium modulation depths with those from a phenothiazine radical sulfonated on the ring shows that the unsubstituted N-alkylphenothiazine is solubilized in a less polar environment, further from the micellar interface. Micelles formed from shorter chain surfactants have more water penetration at the interface compared to micelles formed from longer chain surfactants. This is supported by an increase in the normalized deuterium modulation depth with a decrease in the micelle surfactant chain length.

The photoionization of alkylphenothiazines in cationic dioctadecylmethylammonium chloride (DODAC), neutral dipalmitoylphosphatidylcholine (DPPC), and anionic dihexadecylphosphate (DHP) vesicles of different surface charge, after addition of cholesterol, was studied by ESR and ESEM methods. Neutral alkylphenothiazines (PC_n , where $n = 1, 3, 6, 9, 12$, and 16) were synthesized and used to study the effects of alkyl chain length. It was found that the photoionization efficiency decreases as a function of alkyl chain length of the

alkylphenothiazine until PC₁₆ at which point the photoionization efficiency increases for all three vesicle/alkylphenothiazine systems studied. The magnitude of the photoyields as measured by ESR decreases in the order DODAC > DPPC > DHP. The addition of cholesterol to these three vesicle/alkylphenothiazine systems reduces the photoyields in all cases. The ESEM data on the relative distances from the vesicle interface are consistent with these findings. The results depend on the alkyl chain length of the alkylphenothiazine, the vesicle surface charge, and the ordering effect of cholesterol.

All current funds are anticipated to be spent or encumbered by January 31, 1993.

2. Publications from October, 1991 to October, 1992

1. Piero Baglioni, Elisabeth Rivara-Minten, Chris Stenland and Larry Kevan, Photoionization of N,N,N',N'-Tetramethylbenzidine in Mixed Micelle of Ionic and Nonionic Surfactants: Electron Spin Echo Modulation and Electron Spin Resonance Studies, *J. Phys. Chem.*, 95, 10169 -10172 (1991).
2. Hugh J.D. McManus and Larry Kevan, Electron Spin Resonance, Electron Spin Echo, and Electron Nuclear Double Resonance Investigations of the Photoreduction Yield of a Series of Alkylmethylviologens in Dihexadecylphosphate Vesicles: The Effect of Added Cholesterol, *J. Phys. Chem.*, 95, 10172 - 10178 (1991).
3. Piero Baglioni, Luigi Dei and Larry Kevan, Application of Electron Spin Resonance and Electron Spin Echo Modulation to the Study of Polymeric Latex Interfaces, *Trends Polym. Sci.*, 1, 73 - 82 (1991).
4. Hugh J.D. McManus, Young Soo Kang and Larry Kevan, Photoreduction of Methylviologen in Organized Molecular Assemblies: Role of the Surfactant Headgroup, *J. Phys. Chem.*, 96, 2274 - 2277 (1992).
5. Masato Sakaguchi, Piero Baglioni and Larry Kevan, Photoreduction of Alkylmethylviologens with α -Tocopherol in Dioctadecyldimethylammonium Chloride Vesicles, *J. Phys. Chem.*, 96, 2772 - 2776 (1992).

6. L. Kevan, Applications of Electron Spin Echo Spectroscopy to Location Control of Alkylphenothiazine Derivatives in Photoinduced Charge Separation Across Vesicles, Micelles and Reverse Micelle Interfaces, *Radiat. Phys. Chem.*, 39, 333 - 344 (1992).
7. L. Kevan, Charge Effects on Photoionization Near Vesicle Interfaces, In *Radiation Research: A Twentieth-Century Perspective*, Vol. II; W. C. Dewey, M. Edington, R. J. M. Fry, E. J. Hall and C. F. Whitmore, Eds.; Academic Press: San Diego, 1992; pp. 115 - 120.
8. Hugh J. D. McManus, Young Soo Kang and Larry Kevan, Electron Spin Resonance, Electron Spin Echo, and Electron Nuclear Double Resonance Studies of the Photoreduction Yield of a Series of Alkylmethylviologens in Sodium Dodecylsulfate and Dodecyltrimethylammonium Chloride Micelles: Effect of the Alkyl Chain Length of the Viologen, *J. Phys. Chem.*, 96, 5622 - 5628 (1992).
9. Peter Bratt, Hugh J. D. McManus and Larry Kevan, Electron Spin Echo Modulation Studies of Doxylstearic Acid Spin Probes in Frozen Dihexadecylphosphate and Dioctadecyldimethylammonium Chloride Vesicles: Interaction of the Spin Probe with Deuterated Water and Effects of Cholesterol Addition, *J. Phys. Chem.*, 96, 5093 - 5096 (1992).
10. Peter J. Bratt, Young Soo Kang and Larry Kevan, Photoionization of Neutral Alkylphenothiazines in Positive, Neutral and Negatively Charged Vesicles: Effects of the Addition of Cholesterol and Alkyl Chain Length, *J. Phys. Chem.*, 96, 5629 - 5633 (1992).
11. Peter J. Bratt and Larry Kevan, Electron Spin Resonance Lineshape Analysis of x-Doxylstearic Acid Spin Probes in Dioctadecyldimethylammonium Chloride Vesicles, *J. Phys. Chem.*, 96, 6849-6852 (1992).
12. Young Soo Kang, Hugh J.D. McManus and Larry Kevan, Comparative Electron Nuclear Magnetic Resonance, Electron Spin Echo Modulation and Electron Nuclear Double Resonance Studies of the Photoionization of N-Alkylphenothiazines in Variously Charged Micelles and Vesicles, *J. Phys. Chem.*, 96, 7373-7478 (1992).
13. Piero Baglioni, Luigi Dei, Larry Kevan and Elisabeth Rivara-Minten, Electron Spin Echo Modulation and Electron Spin Resonance Studies of the Interface of Mixed Micelles Formed of Anionic-Cationic and Ionic-Nonionic Surfactants, In *Mixed Surfactant Systems*, Edited by P. M. Holland and D. N. Rubingh, ACS Symposium Series No. 501; American Chemical Society: Washington, D.C., 1992; Chapter 10 (pp 180 - 193).

**DATE
FILMED**

2 / 2 / 93