DOE/ER/13851 - T2

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DOE/ER/13851--T2

DE93 006902

DEPARTMENT OF ENERGY

Progress Report

Isotope Tracer Studies of Diffusion in Silicates and of
Geological Transport Processes using Actinide Elements

G. J. Wasserburg, Principal Investigator
John D. MacArthur Professor of Geology and Geophysics
California Institute of Technology
Mail Code 170-25
Pasadena, CA 91125

DOE Award #: DE-FG03-88ER13851

Duration of Project period: 2-1-93 to 1-31-94

MASTER

ep

PROGRESS REPORT

High Abundance Sensitivity Mass Spectrometer for U-Th Studies

We have documented in our last Progress Report the abundance sensitivity and ion transmission characteristics of the Finnigan/MAT 262B-RP mass spectrometer (see Table 1). This performance has also been presented at the annual meetings of the American Society for Mass Spectrometry and the American Geophysical Union [Papanastassiou, 1991; Papanastassiou *et al.*, 1991]. We consider that abundance sensitivity at the level of 1×10^{-8} or better for one mass unit difference in the U and Th mass region is acceptable performance for measurement of U and Th in young volcanics. The abundance sensitivity of this system for a two mass unit difference, as is applicable for ^{230}Th measurements, is found to be substantially better (see Table 1). This performance has been obtained by the introduction by Finnigan/MAT, through a collaboration with us, of a retarding potential lens for the central ion beam. This lens permits deceleration of the ion beam, so that ions that have suffered a collision and lost some energy are removed from the beam. These ions of lower energy are those that would have formed the "low-mass" tail of the ion beam. This design provides good performance for a versatile and compact instrument, which is also capable of high precision measurements (at the 10-20 ppm level) by simultaneous ion beam collection in the first stage.

The originally planned design included the use of a mass selective device (a dynamic, RF electric quadrupole) after the retarding potential (RP) lens. However, this part of the design was not implemented by Finnigan/MAT, due to very low (< 1%) transmission of their first trial system, which used an RF-quadrupole not specifically designed to match the ion optics. An ion-optical study of the Finnigan/MAT system carried out at Caltech, indicated that the ion beam was not optimized in angular dispersion for introduction into the RF-quadrupole. We have completed a new ion optical and mechanical design for the second stage of the mass spectrometer. This design includes a static electric quadrupole doublet for beam shaping prior to the deceleration (RP) lens, in contrast to the single static electric quadrupole used by Finnigan/MAT. The deceleration lens has also been redesigned to provide for gradual deceleration

and increased transmission. This system has been installed and checked out. The data on Icelandic basalts, presented later in this report, have been obtained using this redesigned system. We are currently in the process of checking out the RF quadrupole purchased to match the new design parameters. Following this initial testing, the RF quadrupole will be installed and checked on the Finnigan/MAT mass spectrometer. It is our expectation that the increased mass selectivity provided by the RF quadrupole will make it possible to allow a wider energy range of ions through the decelerating lens, thus increasing the overall transmission through the second stage of the mass spectrometer.

Table 1. Abundance sensitivity and transmission for Finnigan/MAT 262B-RP

	Unmodified	Modified ^b
Abundance Sensitivity (U-Th region) ^a		
1 amu	1×10^{-8}	5×10^{-9}
1.5 amu	4×10^{-9}	5×10^{-10}
Transmission (2nd stage)	50%	50%

^aFor low mass side. ^bNote: New ion-optical design for the purpose of adding dynamic (RF) electric quadrupole to improve abundance sensitivity. The transmission is expected to improve when the RF electric quadrupole is installed.

We also document the performance of the first stage of this instrument using simultaneous ion beam measurements. We have prepared and analyzed gravimetric standards of Sr with small enrichments in ⁸⁷Sr, ranging from 15 ppm to 140 ppm. The results are shown in Figure 1a,b, for individual analyses and for averages of several analyses per standard. The results fall within the ± 15 ppm dotted lines. These data demonstrate that ⁸⁷Sr/⁸⁶Sr can be measured with a precision of ± 15 ppm. This precision will be applied to studies of water-rock interactions and of element migration during diagenesis and metamorphism.

²³⁸U-²³⁰Th Disequilibrium in Recent Lavas from Iceland

Iceland is the type locality for a ridge-centered hotspot. Studies of the petrology and

Figure 1a. Enrichments (ζ) in $^{87}\text{Sr}/^{86}\text{Sr}$ determined by individual mass spectrometer analyses (ordinate), using simultaneous ion beam collection with the first stage of the Finnigan/MAT mass spectrometer, versus the gravimetrically calculated enrichments in this ratio (abscissa). The units are parts per million deviations for the $^{87}\text{Sr}/^{86}\text{Sr}$ ratios. The dotted lines indicate a ± 15 ppm range; the solid lines indicate a ± 30 ppm range. The means of individual measurements fall within the ± 15 ppm range, with the uncertainty limits extending to the ± 30 ppm range.

Figure 1b. Same graph for the grand mean averages for each standard, from the individual mass spectrometric analyses shown in Fig. 1a. The results fall well within the ± 15 ppm range and show a resolution of ± 15 to ± 20 ppm in $^{87}\text{Sr}/^{86}\text{Sr}$.

geochemistry of recent volcanic rocks from Iceland have demonstrated the existence of significant variations in chemical composition and in the isotopic ratios of young rocks at various volcanic centers. Several models, falling mainly in one of two types, have been proposed to explain these variations: (1) mantle heterogeneities and (2) crustal contamination. While studies of Sr, Nd and Pb isotopes emphasize mantle heterogeneities, O isotope data suggest crustal contamination of mantle-derived melts. Recent studies of ^{238}U - ^{230}Th systematics in young volcanic rocks from Iceland favor the crustal contamination model, but do not exclude the existence of mantle heterogeneities.

In this study, we applied the high abundance sensitivity TIMS technique to obtain high precision $^{230}\text{Th}/^{232}\text{Th}$ and $^{234}\text{U}/^{238}\text{U}$ ratios on some basalts from Iceland. Two olivine tholeiites (EY3 and EY4A) from the Eyjafjöll volcano system were analyzed. K-Ar dating on EY3 indicates an age of about 10^5 years, while EY4A is thought to be of Holocene age ($\sim 10^4$ y). These two samples were selected for this U-Th study because they both contain abundant phenocrysts of plagioclase, olivine and pyroxene, which may be characterized by different Th/U ratios.

A 1.36 g whole rock sample of EY4A-1, was spiked with ^{229}Th and ^{233}U and ^{236}U and dissolved with HF, HClO_4 , HCl and HNO_3 in a thermally pressurized Teflon bomb. Residues from the dissolution step were treated with HClO_4 and aqua regia. After three dissolution cycles this sample was dissolved completely. A small aliquot (0.2%) was spiked with higher levels of ^{229}Th , and ^{233}U and ^{236}U for determination of the concentrations of ^{232}Th , ^{238}U and the $^{238}\text{U}/^{235}\text{U}$ ratio. Another small aliquot ($\sim 5\%$) was saved. The remaining aliquot ($\sim 95\%$) was used for measurement of $^{230}\text{Th}/^{232}\text{Th}$ and $^{234}\text{U}/^{238}\text{U}$ ratios.

The U and Th isotopic ratios were determined using the modified Finnigan/MAT 262 B-RP mass spectrometer described in the previous section. Briefly, (1) the transmission for the first stage is 35%; the transmission of the second stage is 50%, for an overall transmission of $\sim 20\%$; (2) the abundance sensitivity at 1 amu below ^{238}U or ^{232}U is $\sim 5 \times 10^{-9}$ and at 1.5 amu below ^{238}U or ^{232}Th the abundance

sensitivity is 5×10^{-10} ; (3) for samples of 10^{-6} g of U and of Th, the ionization efficiency for U is $\sim 10^{-3}$, and for Th $\sim 10^{-4}$. These performance characteristics are well suited for the high precision measurements of U and Th isotopic ratios. For example, the mass spectrum of Th from an Icelandic basalt shown in Figure 2, indicates null backgrounds at both sides of ^{230}Th , eliminating the need for correction for a low energy tail from the neighboring, large ^{232}Th beam.

For EY4A-1, the ^{238}U and ^{232}Th concentrations are 6.14×10^{-9} mol/g and 1.94×10^{-8} mol/g, respectively, yielding a $^{232}\text{Th}/^{238}\text{U}$ atomic ratio of 3.16 or a $[^{238}\text{U}/^{232}\text{Th}]$ activity ratio of 0.992 (square brackets denote activity ratios). The data are shown in Table 2. The $^{230}\text{Th}/^{232}\text{Th}$ ratio for EY4A-1 is 6.04×10^{-6} (atomic) or 1.122 (activity). These results indicate a 13% enrichment of ^{230}Th relative to ^{238}U in EY4A-1 ($[^{230}\text{Th}/^{238}\text{U}] = 1.131 \pm 0.013$). A second whole rock sample, EY4A-2, was analyzed as above and yields the same results as EY4A-1. The third whole rock sample, EY4A-3, yields the same $^{230}\text{Th}/^{232}\text{Th}$ ratio as EY4A-1 and EY4A-2, but a 1% higher ^{238}U concentration and a correspondingly lower $[^{230}\text{Th}/^{238}\text{U}]$ of 1.102 ± 0.008 .

We applied the same analytical procedures to the analyses of three whole rock samples of EY3, ranging from 1.5 to 1.8 g. The results indicate a factor of 0.35 lower U and Th concentrations in EY3 than in EY4A. Replicate analyses of EY3 also show about a 4% variation in U concentrations, but only a 1% variation in Th. The $^{230}\text{Th}/^{232}\text{Th}$ and $^{230}\text{Th}/^{238}\text{U}$ ratios in replicate analyses of EY3 vary by 2% and 5%, respectively; this variation is much larger than the analytical errors of about 1%. Currently, we do not know if this difference is due to sample heterogeneity at the 1 to 2 g scale. The $[^{230}\text{Th}/^{238}\text{U}]$ ratios in EY3 also indicate from 9 to 14% enrichments of ^{230}Th relative to the equilibrium value.

We have analyzed separates of plagioclase, magnetite and a mixture of olivine and pyroxene from EY4A to determine the U, Th concentrations and $^{232}\text{Th}/^{238}\text{U}$ ratios, in an attempt to find out if an internal isochron can be established for this sample. The first plagioclase separate (Plag1, 1.01 g) consisted of almost pure plagioclase. The second plagioclase separate (Plag2, 1.69 g) contained some

Figure 2. A mass spectrum of thorium between mass 228 and 232. A ^{229}Th spike was added. The small peak at mass 231 may be from Pa which was not completely separated from this sample. Note that on both sides of ^{230}Th , the background is essentially zero, which eliminates the need for any correction from the neighboring intense ^{232}Th peak.

inclusions or impurities of magnetite, olivine, glass, and rare Fe-sulfide. The U and Th concentrations in Plag1 are 1.65×10^{-10} mol/g and 4.96×10^{-10} mol/g, respectively; these values are ~ 40 times lower than the corresponding values in the whole rock samples. The $^{232}\text{Th}/^{238}\text{U}$ atomic ratio in Plag1 (3.01) is only $\sim 5\%$ lower than in the whole rock. The U and Th concentrations in the second plagioclase separate are $\sim 30\%$ lower than in Plag1, while the $^{232}\text{Th}/^{238}\text{U}$ ratio the same as in the whole rock. The ^{238}U and ^{232}Th concentrations in a magnetite separate (0.1 g) are 2.44×10^{-9} mol/g and 7.67×10^{-9} mol/g, respectively; these values are $\sim 40\%$ lower than in the whole rock. However, the $^{232}\text{Th}/^{238}\text{U}$ ratio (3.17) in magnetite is the same as for the whole rock. The ^{238}U and ^{232}Th concentrations in a mixture of olivine and pyroxene are 2.1×10^{-10} mol/g and 6.58×10^{-10} mol/g, respectively, or ~ 30 times lower than in the whole rock. The $^{232}\text{Th}/^{238}\text{U}$ ratio (3.15) in this mixture is also the same as in the whole rock.

These results indicate that for EY4A the $^{232}\text{Th}/^{238}\text{U}$ ratios in all major phenocrysts, i. e. plagioclase, magnetite, olivine and pyroxene are the same as in the whole rock, but the concentrations are lower from 40% to factors of 40. This suggests that U and Th may be concentrated in a carrier phase (glass?) and its presence in the whole rock and mineral separates dominates the distributions of U and Th in these phases. Given the constant $^{232}\text{Th}/^{238}\text{U}$ ratios in all separated mineral phases of EY4A, it will be difficult to establish an internal isochron for this sample.

Discussion

The present-day [$^{230}\text{Th}/^{238}\text{U}$] ratios in both EY4A and EY3 indicate from 9 to 15% enrichments of ^{230}Th relative to secular equilibrium values. The results are plotted in a $^{230}\text{Th}/^{232}\text{Th}$ versus $^{238}\text{U}/^{232}\text{Th}$ diagram in Figure 3. Data based on α -counting techniques from Sigmarsson *et al.* (1991, 1992) and other workers on volcanic rocks from Iceland are also shown for comparison. The precision of $^{230}\text{Th}/^{232}\text{Th}$ of our new results is a factor of ~ 4 better than that obtained using α -counting. The [$^{238}\text{U}/^{232}\text{Th}$] ratios (0.95 to 1.0) of the Eyjaföll basalts fall within the range of other Icelandic volcanic rocks. The present values of Eyjaföll basalts plot between the field of "primary mantle melts" and the

Table 2. U-Th Results on Volcanic Rocks from Iceland

Sample ^a	Weight g	²³⁸ U nmole/g	²³² Th nmole/g	²³⁰ Th pmole/g	²³⁰ Th/ ²³² Th atomic x10 ⁶	²³⁰ Th/ ²³² Th activity	²³² Th/ ²³⁸ U atomic	²³⁸ U/ ²³² Th activity
EY4A								
-1	1.36	6.139±0.009	19.40±0.06	0.1171±0.0013	6.038	1.122±0.013	3.159	0.9924±0.0032
-2	1.34	6.154±0.016	19.59±0.05	0.1169±0.0008	5.965	1.109±0.008	3.184	0.9848±0.0037
-3	1.36	6.219	19.46	0.1156±0.0009	5.940	1.104±0.008	3.129	1.0021
-Plag1	1.01	0.165	0.496				3.005	1.0434
-Plag2	1.69	0.107	0.341				3.194	0.9817
-Mag	0.10	2.44	7.67				3.167	0.9900±0.0069
-Ol+Px	0.11	0.210	0.658				3.153	0.9945±0.0042
EY3								
-1	1.50	2.197±0.007	7.169±0.011	0.04148±0.0007	5.785	1.075±0.018	3.263	0.9610±0.0032
-2	1.55	2.176±0.007	7.183±0.019	0.04213±0.0003	5.866	1.090±0.009	3.301	0.9499±0.0038
-3	1.80	2.261	7.241	0.0416±0.0003	5.745	1.068±0.006	3.202	0.9792

^aFor mineral separates small amounts were analyzed to determine the level of U, Th concentrations and the range in U/Th.

zone of Icelandic crust in ^{238}U - ^{230}Th radioactive equilibrium. However, it is more appropriate to plot the initial values for this comparison. The initial $^{230}\text{Th}/^{232}\text{Th}_i$ ratios for the Eyjaföll samples can be calculated using the following equation:

$$[^{230}\text{Th}/^{232}\text{Th}]_i = [^{230}\text{Th}/^{232}\text{Th}]_t \exp(-\lambda_{230}t) + ^{238}\text{U}/^{232}\text{Th}_i (1 - \exp(-\lambda_{230}t))$$

The initial values for EY4A, after adjusting for *in situ* decay for 10^4 years, increase by only about 1% but, for EY3, the initial values adjusted for *in situ* decay for 10^5 years increase from 12 to 19%. The initial values for EY4A (1.11 to 1.14) are similar to those from recent basalts from the lateral non-rifting "South Iceland volcanic zone" of Sigmarsson *et al.* (1991, 1992), but the initial values (1.2-1.3) for EY3 are close to the field of some active rift-zone tholeiites (Hemond *et al.*, 1988). Obviously, the true initial values of these samples depend on the exact chronology of these volcanic rocks.

Our high precision Th-U results clearly indicate high enrichments of ^{230}Th relative to the equilibrium value in basalts from the Eyjaföll volcano. The Th/U ratio in different phases of EY4A appears to be very uniform and requires an explanation. We will carry out further analyses of U-Th isotopes as well as Sr, Nd and Pb isotopes on these and other Icelandic volcanic rocks from this volcanic system in the future. This will help define better the $^{230}\text{Th}/^{232}\text{Th}$ vs. $^{87}\text{Sr}/^{86}\text{Sr}$ mantle array [Condomines *et al.*, 1988] and the extent to which the samples reflect mantle source heterogeneities or crustal contamination.

Figure 3. The $(^{230}\text{Th}/^{232}\text{Th})$ - $(^{238}\text{U}/^{232}\text{Th})$ activity isochron diagram from Sigmarsson, Condomines and Fourcade (1992). The typical 2σ error shown corresponds to the α -counting data. The uncertainties for the TIMS data presented here (crosses) are smaller than the symbols used. The broken lines represent the field of rift-zone tholeiites [data from Condomines *et al.*, 1981; Hémond *et al.*, 1988]. C.C. denotes crustal contamination and points to the domain of Icelandic crust in ^{238}U - ^{230}Th radioactive equilibrium. The field of mantle melts expected below Iceland is also shown.

Water-Rock Interaction from U-Th Studies

The problem of U and Th transport in groundwaters and the reactions of these elements with aquifer rocks is of interest in terms of retention/storage and diagenesis. The broad utility of ^{238}U - ^{230}Th dating is fundamentally limited by diagenetic changes. The time range for which ^{230}Th dating on carbonates can be used is, in principle, from 0 ± 5 years to $200,000 \pm 5,000$ years, using TIMS techniques [Edwards *et al.*, 1987]. There is, however, abundant evidence that shifts in $^{234}\text{U}/^{238}\text{U}$ occur in fossil corals [cf Edwards *et al.*, 1987; Chen *et al.*, 1991; Bard *et al.*, 1990] that must be due to interactions with fluids [cf Banner *et al.*, 1990, 1991]. A possible source of excess ^{234}U is obviously from α -recoil, as has been abundantly demonstrated in continental terranes. However, it is not evident that this process operates in carbonate-dominated groundwater systems. We have previously carried out a short study on waters from Barbados and found that no evidence of excess ^{234}U above that in modern sea water. In all fresh waters the $^{234}\text{U}/^{238}\text{U}$ was not distinguishable from the secular equilibrium value. In order to establish whether α -recoil was not operable, we carried out an intensive study on an isolated carbonate island (San Salvador, Bahamas). This locality had been investigated geologically [Curran and White, 1984, 1985] and the ^{230}Th ages were the subject of an extensive study [Chen *et al.*, 1991]. Samples of waters were collected from fresh water wells, brackish creeks, blue holes and hypersaline lakes and the concentrations of U and the $^{234}\text{U}/^{238}\text{U}$ ratios were determined. In addition Ca, Mg, and salinity were measured. The data are shown in Table 3 and Figure 4. The hypersaline lakes are obviously the product of evaporation of seawater (plus spray) and rain water. The ground waters are fresh (0-2 ‰ salinity) and enriched in Ca (76-122 ppm) and in U due to dissolution of carbonate. The $\delta^{234}\text{U}$ in the well-waters ranges from 4.5 to 35 ‰. All other samples yielded $\delta^{234}\text{U}$ less than or equal to that of seawater. In a Ca/U versus $\delta^{234}\text{U}$ diagram (Fig. 3), data from seawater, blue hole and well waters approximately fit a straight line, indicating a two-component mixing between seawater and another end member. The Ca/U ratio of this end member as determined from the intercept between this line and the abscissa at

$\delta^{234}\text{U} = 0$ is $\sim 1.5 \times 10^5$. This corresponds to an old carbonate ($> 2 \times 10^5$ y) at nearly radioactive secular equilibrium. Carbonates of the Pleistocene reef corals which have a present $\delta^{234}\text{U}$ value of 100-110 ‰ can not be the end member. None of these waters can provide a source for the excess ^{234}U found in the fossil corals.

Thorium has been measured for one sample; it shows a low $^{232}\text{Th}/^{238}\text{U} = 1.9 \times 10^{-5}$ and $^{230}\text{Th}/^{238}\text{U} \leq 2 \times 10^{-8}$. This suggests that the waters are depleted in ^{232}Th and highly depleted in ^{230}Th relative to the fossil corals. These results show that diagenesis involving solutions such as these modern groundwaters, in a carbonate terrace, cannot explain the elevated $\delta^{234}\text{U}(\text{T})$ found in some of the Pleistocene corals at San Salvador. The low $^{232}\text{Th}/^{238}\text{U}$ and very low $^{230}\text{Th}/^{238}\text{U}$ values show that Th is much less mobile than U and that $^{230}\text{Th}/^{238}\text{U}$ enrichment in fossil coral skeletons could be caused by Th being very preferentially left behind when U goes into solution.

Table 3. U isotopes and Ca, U concentrations in Bahamian waters

Sample	Salinity	pH	^{238}U pmole/g	$\delta^{234}\text{U}$ (‰)	U ppb	Ca ppm	U/Ca atomic x 10^6
Mrs. Ward's Well	2	7.6	17.29	4.47	4.145	122.3	5.665
Jake Jones's Store Well	0	8	5.193	29.47	1.245	76	2.739
Municipal Building Faucet	0	7.6	9.079	34.94	2.177	111.25	3.271
Little Lake	62	8.4	6.98	81.1	1.674	649.6	0.4306
Storr Lake	79	8.4	16.844	84.05	4.039	856.4	0.7883
Long Lake	77	8.3	7.384	108.3	1.77	846.7	0.3496
Watling's Blue Hole	28	8.1	14.738	83.7	3.534	354	1.669
Pigeon Creek-1	38	8.2	13.3	149.2	3.188	450.6	1.183
Pigeon Creek-2	40	8.3	12.164	141	3.156	461.9	1.142
Seawater	37.5	8.35	14.16	148.5	3.394	422	1.345

SAN SALVADOR, BAHAMAS

Figure 4. Correlation of Ca/U relative to $\delta^{234}\text{U}$. A straight line between wellwaters and seawater would imply a two-component mixing. The fresh ground waters, blue hole and seawater are compatible with a two component mixing. The hypersaline lakes cannot simply be mixtures of evaporated seawater and rain water but require a depletion of U relative to Ca.

Resonance Ionization Mass Spectrometry of Os and Ti

The development of resonance ionization mass spectrometry (RIMS) has provided a new analytical technique that offers extremely high sensitivity, at the part per trillion level, for a wide range of elements. In combination with laser ablation or ion sputtering sources RIMS can provide *in situ* chemical and isotopic analyses free of the isobaric interferences that pose a fundamental limitation to the application of conventional secondary ion mass spectrometry (SIMS). The high sensitivity of RIMS arises from the efficiency and selectivity of the resonant photoionization process. The capability to ionize only the species of interest allows the use of relatively simple, high transmission, low mass resolving power mass spectrometers. Coupled with the near unit efficiency of the resonant ionization process, this capability gives RIMS the potential for high precision analyses on very small samples ($< 10^8$ atoms). While RIMS has been successfully applied to the analysis of nearly 70 elements [Hurst *et al.*, 1975; Parks *et al.*, 1988; Pellin *et al.*, 1987], the use of RIMS for high precision and accuracy isotopic analyses has been inhibited by evidence of large shifts in isotope ratios measured with RIMS. Observed shifts in isotope ratios range from $\sim 3\%$ to nearly 100%, typically with an enhancement of the odd-mass isotopes [Walker and Fassett, 1986; Fairbank *et al.*, 1989; Spiegel *et al.*, 1992]. Although endemic to the resonant ionization process, in most cases the isotope shifts cannot be understood in terms of either laser bandwidth effects or selection-rule differences in ionization probability. These large shifts in measured isotope ratios hinder the recognition of intrinsic isotopic variations and have, in the past, required frequent analyses of standards and severely limited the accuracy of RIMS isotope data. The ubiquitous presence of isotope selective effects, induced by laser excitation and ionization, and the pressing need to develop a clear theoretical understanding of these effects led us to undertake an investigation of the relationship between laser parameters such as intensity, bandwidth and polarization state, and the nature of isotope shifts in RIMS. Our experiments have focused on RIMS studies of Os and Ti; the results are summarized below and are discussed in detail in the appended papers.

In order to explore the physical processes leading to isotopic selectivity in Os, three different types of RIMS experiments were performed: Os isotope ratios were measured as a function of (1) laser wavelength and bandwidth, (2) laser intensity and (3) the polarization state and focussing of the laser radiation. Our data reveal large and, in many cases, systematic variations in isotope ratios with deviations from normal values of up to 20%. Ratios of even-mass Os isotopes are strongly dependent on laser wavelength and intensity even when the laser bandwidth is up to a factor of eight larger than the optical isotope shift. This effect is reduced for transitions with a large oscillator strength. The magnitude of isotope selectivity for both even-even and even-odd mass Os isotope ratios increases with decreasing laser power. Ratios of odd to even mass isotopes exhibit a consistent depletion of odd mass isotopes and depend strongly on the polarization state of the laser radiation; polarization scrambling reduces even-odd isotope selectivity. The results of these experiments are discussed fully in the paper by Wunderlich *et al.* (1992), recently published in *Int. Jour. Mass Spectrom. and Ion Processes*.

The pronounced wavelength dependence of even-mass Os isotope ratios led us to develop a wavelength tuning procedure to permit the reproducible measurement of both even-even and even-odd Os isotope ratios with a precision and accuracy of better than 0.5%. To determine whether or not this technique can be generally applied to eliminate or significantly reduce laser-induced isotope selectivity for other systems, we initiated a RIMS study of Ti. The large shifts ($\sim 40\%$) in even-to-odd mass Ti isotope ratios reported in a previous RIMS experiment [Spiegel *et al.*, 1992] serve as a convenient benchmark against which the effectiveness of our laser tuning procedures can be evaluated.

The wavelength dependent isotope shifts in Ti are significantly different in sign and magnitude from the effects previously observed for Os. For both Os and Ti the intensity of the photoion signal was measured for each isotope as a function of the laser wavelength for several resonance transitions. In Os, the absorption maxima of the heavier isotopes are shifted to progressively longer wavelengths reflecting the (normal) increase of the average squared nuclear charge radius with addition of neutrons. For Ti,

in contrast, the wavelength shift is reversed. The ionization cross-section of the heavier Ti isotopes increases with progressively shorter wavelengths, reflecting a decrease in the nuclear charge radius as the closed neutron shell with magic neutron number 28 is approached. The wavelength dependence of even-even Ti isotope ratios is more pronounced than that observed for Os because (i) the change in the average nuclear charge radius with neutron number is significantly larger for the Ti isotopes and (ii) the electrons removed in the Os resonance transitions (6s electrons) possess a much lower electron density at the nucleus compared to the electrons removed in the Ti experiments (4s electrons).

During the measurements of the wavelength dependence of ionization probabilities for the Ti isotopes (optical isotope shifts), we identified several resonance transitions for which the $^{50}\text{Ti}/^{46}\text{Ti}$ and $^{48}\text{Ti}/^{46}\text{Ti}$ ratios remained constant over a small wavelength interval ($\Delta \sim 0.1 \text{ cm}^{-1}$). In this region both ratios differ from the respective standard ratios only by mass-dependent fractionation. Both ratios yield equal fractionation of $\sim 25 \text{ ‰/amu}$ favoring the lighter Ti isotopes, indicating that laser-induced isotope selectivity is absent. For these same transitions measurements of $^{47}\text{Ti}/^{46}\text{Ti}$ and $^{49}\text{Ti}/^{46}\text{Ti}$ ratios indicated the absence of odd-even isotope shifts larger than 2% for saturating laser intensity in the ionization step. In contrast, we demonstrated large enrichments of the odd-mass isotopes, $\delta^{47}\text{Ti} = 60 \text{ ‰}$ and $\delta^{49}\text{Ti} = 144 \text{ ‰}$, for measurements under nonsaturating conditions. Even-odd Ti isotope ratios measured with RIMS are also sensitive to the degree of polarization of the laser radiation; odd-mass isotopes become increasingly enriched with increasing polarization.

The absence of laser-induced isotope selectivity for the $^{47}\text{Ti}/^{46}\text{Ti}$ and $^{49}\text{Ti}/^{46}\text{Ti}$ ratios under saturating ionization conditions found in our experiments can be compared with the large enhancements of ^{47}Ti and ^{49}Ti reported by Spiegel *et al.* (1992). Using a three photon, two color ionization scheme with an autoionizing state, Spiegel *et al.* found over abundances of the odd-mass isotopes of up to 450‰. Our experiments suggest that such large isotope shifts cannot be attributed to optical isotope shifts or bandwidth effects and point to the need to determine carefully the influence of autoionizing resonances

on even-odd isotope ratios.

In summary, we believe our experiments on Ti and Os have provided critical new information on the nature of laser-induced isotope shifts in RIMS. The results from both our Ti and Os RIMS experiments demonstrate that by carefully choosing resonance transitions and laser operating parameters, both even-even and even-odd isotope ratios free from laser-induced isotope selectivity can be measured using RIMS. These two studies are the first to demonstrate this important result. Although additional work is needed to improve the precision of the Ti isotope ratio measurements, we are confident the techniques developed in these studies will allow the use of RIMS for the determination of unbiased isotope ratios with a precision and accuracy of better than 0.5%.

In addition to the aforementioned paper in *Int. Jour. Mass. Spectrom. Ion Proc.*, four new manuscripts describing our RIMS studies are either under review, in press or have been published. Copies of all five papers are appended.

Self-Diffusion of Mg

We have developed a new technique which allows measurement of self-diffusion rates under conditions of chemical equilibrium across the diffusion interface. The technique takes into account complementary diffusion in both phases and permits simultaneous determination of the activation energies and pre-exponential factors in both phases with very high precision. The solution to the diffusion transport equation in one phase can be used immediately to construct the solution in the coexisting phase. Diffusion data in materials where measurement is inhibited by slow diffusion rates may thus be obtained by the indirect measurement of diffusion parameters in a coexisting material where diffusion is rapid. This technique represents an important advance in studies of diffusion transport and can be applied to many systems of interest in geochemistry and material science.

We applied this technique to determine the rates of Mg self-diffusion in spinel and co-existing silicate melt. The diffusion couple consisted of a gem quality spinel crystal with a normal Mg isotopic

composition and a spinel-saturated, ^{25}Mg -enriched CMAS melt. The bulk composition of the melt matched that of a group of plagioclase-rich, refractory inclusions in the Allende meteorite; small adjustments to the melt composition ensured that the melt and spinel remained in chemical equilibrium at each run temperature. The diffusion couple was annealed in air at temperatures between 1260° and 1553°C for periods of 0.5 to 20h and the resulting gradients in Mg isotopic composition were measured normal to the interface with the PANURGE ion microprobe.

The diffusion coefficients for melt and spinel for each run were calculated from the measured Mg isotope profiles using a model of diffusion in an infinite composite medium. The model incorporates the complementary diffusion of ^{24}Mg , ^{25}Mg , and ^{26}Mg in both phases with the constraint that the Mg concentration on each side of the interface remain invariant. The data exhibit a good linear relationship on an Arrhenius plot, $\ln D$ vs T^{-1} , suggesting that transport of Mg is governed by a single mechanism over the temperature range studied. The activation energy and pre-exponential factor for spinel are 384 ± 7 kJ and 74.6 ± 1.1 cm^2/sec , respectively, while the corresponding values in the melt are 343 ± 25 kJ and 7791.9 ± 1.3 cm^2/sec . The similarity in the activation energies suggests that the potential barriers for diffusion are the same in spinel and melt, while the larger pre-exponential factor reflects the much greater intrinsic jump frequency or jump amplitude in the melt. At a temperature of 1400°C our results show that Mg diffusion in spinel is ~ 15 times slower than reported by Lindner and Åkerström (1958).

The results of our Mg self-diffusion experiments provide constraints on the thermal history of plagioclase-olivine inclusions in meteorites. To preserve Mg isotope heterogeneity in spinel, the inclusions must have experienced maximum melting temperatures of $\sim 1500^\circ\text{C}$ and have initially cooled faster than 50 to 250°C/hr . This cooling rate is comparable to or up to 10 times faster than rates inferred from experimental and textural studies of natural and synthetic assemblages. Coupled with data from our earlier petrographic and isotopic studies of refractory inclusions [Meeker *et al.*, 1983; Huneke *et al.*, 1983; Sheng *et al.*, 1991], the new data on Mg diffusion rates suggest that most refractory

inclusions experienced flash heating in local, hot regions of the solar nebula. More generally, we believe this new approach has widespread application to precise studies of multicomponent diffusion transport and will provide insight into elemental transport mechanisms in melts and crystals. The results of this study were published in *Geochimica et Cosmochimica Acta* [Sheng *et al.*, 1992; copy appended] and represent part of the Ph.D. research of Y. J. Sheng [Sheng, 1992].

References

- J. L. Banner, G. J. Wasserburg, J. H. Chen, and C. H. Moore (1990). *Earth Planet. Sci. Lett.* 101, 296-312.
- J. L. Banner, G. J. Wasserburg, J. H. Chen and J. D. Humphrey (1991). *Earth Planet. Sci. Lett.* 107, 129-137.
- E. Bard, B. Hamelin, R. G. Fairbanks, and A. Zindler (1990). *Nature* 245, 405-410.
- J. H. Chen, H. A. Curran, B. White, and G. J. Wasserburg (1991). *Geol. Soc. Am. Bull.* 103, 82-97.
- J. H. Chen, D. A. Papanastassiou and G. J. Wasserburg (1992). IGC - Kyoto, Japan.
- M. Condomines, C. Hémond, C. J. Allègre (1988). *Earth Planet. Sci. Lett.* 90, 243-262.
- M. Condomines, Ph. Morand, and C. J. Allègre (1981) *Earth Planet. Sci. Lett.* 55, 393-406.
- H. A. Curran and B. White (1984). in Teeter, J. W. ed., *Proceedings of the second symposium on the geology of the Bahamas: San Salvador, Bahamas, CCFL Bahamian Field Station*, p. 71-96.
- H. A. Curran and B. White (1985), in Curran, H. A. ed., *Pleistocene and Holocene carbonate environments on San Salvador Island, Bahamas: Guidebook for Geological Society of America, Orlando, Florida, Annual Meeting Field Trip 2: San Salvador, Bahamas, CCFL Bahamian Field Station*, p. 95-120.
- R. L. Edwards, J. H. Chen, and G. J. Wasserburg (1987). *Earth Planet. Sci. Lett.* 81, 175-192
- W. M. Fairbank, Jr., M. T. Spaar, J. E. Parks and J. M. R. Hutchinson (1980). *Phys. Rev.* A40, 2195-2198.
- C. Hémond, M. Condomines, S. Fourcade, C. J. Allègre, N. Oskarsson, and M. Javoy (1988). *Earth Planet. Sci. Lett.* 87, 273-285.
- J. C. Huneke, J. T. Armstrong and G. J. Wasserburg (1983). *Geochim. Cosmochim. Acta* 47, 1635-1650.

- G. S. Hurst, N. G. Payne, M. H. Nayfeh, J. P. Judish and E. B. Wagner (1975). *Phys. Rev. Lett.* 35, 82-85.
- R. Lindner and Å. Åkerström (1958). *Z. Phys. Chem.* 18, 303-307.
- G. P. Meeker, G. J. Wasserburg and J. T. Armstrong (1983) *Geochim. Cosmochim. Acta* 47, 707-721.
- D. A. Papanastassiou (1991). Invited Talk, ASMS Conference, Nashville, TN.
- D. A. Papanastassiou, J. H. Chen and G. J. Wasserburg (1991). *EOS, Trans. Amer. Geophys. Union* 72, 291.
- J. E. Parks, N. T. Spaar, P. W. Beekman and L. J. Moore (1988). *Resonance Ionization Spectroscopy 1988, Inst. of Phys. Conf. Series* 94, 197-200.
- M. J. Pellin, C. E. Young, W. F. Calaway, J. W. Burnett, B. Jorgensen, E. L. Schweitzer and D. M. Gruen (1987). *Nucl. Instr. Meth.* B18, 446-451.
- Y. J. Sheng (1991) Ph.D. Thesis, California Institute of Technology, 271 pp.
- Y. J. Sheng, I. D. Hutcheon, G. J. Wasserburg (1991). *Geochim. Cosmochim. Acta* 55, 581-599.
- Y. J. Sheng, G. J. Wasserburg and I. D. Hutcheon (1992) *Geochim. Cosmochim. Acta* 56, 2535-2546.
- O. Sigmarsson, C. Hémond, M. Condomines, S. Fourcade, N. Oskarsson (1991). *Geology* 19, 621-624.
- O. Sigmarsson, M. Condomines, S. Fourcade (1992). *Earth Planet. Sci. Lett.* 110, 149-162.
- D. R. Spiegel, W. F. Calaway, A. M. Davis, J. W. Burnett, M. J. Pellin, S. R. Coon, C. E. Young, R. N. Clayton, D. M. Gruen (1992). *Anal. Chem.* 64, 469-475.
- R. J. Walker and J. D. Fassett (1986). *Anal. Chem.* 58, 2923-2927.
- R. K. Wunderlich, I. D. Hutcheon, G. J. Wasserburg and G. A. Blake (1992) *Int. Jour. Mass Spectrom. Ion Proc.* 115, 123-155.

END

**DATE
FILMED**

2 1261 93