12
2-4-92 JSD

PNL-7894
UC-721

Investigation of the Potential of Silica-Bonded Macrocyclic Ligands for Separation of Metal Ions From Nuclear Waste

D. M. Camaioni
N. G. Colton
R. L. Bruening

January 1992

Prepared for the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute

PNL-7894

REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes **any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.** Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401. FTS 626-8401.

Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

INVESTIGATION OF THE POTENTIAL
OF SILICA-BONDED MACROCYCLIC
LIGANDS FOR SEPARATION OF METAL
IONS FROM NUCLEAR WASTE

D. M. Camaioni
N. G. Colton
R. L. Bruening^(a)

MASTER

January 1992

Prepared for
the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Richland, Washington 99352

(a) IBC, Advanced Technologies, Provo, Utah 84603

28

EXECUTIVE SUMMARY

This report describes the testing of some novel separations materials known as SuperLig™ materials for their ability to separate efficiently and selectively certain metal ions from a synthetic, nonradioactive nuclear waste solution. The materials, developed and patented by IBC Advanced Technologies, are highly selective macrocyclic ligands that have been covalently bonded to silica gel. The SuperLig™ materials that were tested are: 1) SuperLig™ 601 for barium (Ba^{2+}) and strontium (Sr^{2+}) separation, 2) SuperLig™ 602 for cesium (Cs^+) and rubidium (Rb^+) separation, 3) SuperLig™ 27 for palladium (Pd^{2+}) separation, and 4) SuperLig™ II for silver (Ag^+) and ruthenium (Ru^{3+}) separation. Our observations show that the technology for separating metal ions using silica-bonded macrocycles is essentially sound and workable to varying degrees of success that mainly depend on the affinity of the macrocycle for the metal ion of interest. It is expected that ligands will be discovered or synthesized that are amenable to separating metal ions of interest using this technology. Certainly more development, testing, and evaluation is warranted.

CONTENTS

EXECUTIVE SUMMARY	iii
INTRODUCTION	1
EXPERIMENTAL PROCEDURES	3
General	3
Separation of Ba^{2+} and Sr^{2+}	3
Separation of Cs^+ and Rb^+	3
Separation of Pd^{2+}	3
Separation of Ru^{3+} and Ag^+	4
RESULTS	5
Separation of Ba^{2+} and Sr^{2+}	5
Separation of Cs^+ and Rb^+	6
Separation of Pd^{2+} in Experiment 1a	7
Separation of Pd^{2+} in Experiment 4	7
Separation of Ru^{3+} in Experiment 1b	8
Separation of Ag^+ in Experiment 1b	8
CONCLUSIONS	11

FIGURES

1 Comparison of IBC and PNL Tests of Superlig™ 601 for Separation of Ba and Sr	13
2 Comparison of IBC and PNL Tests of Superlig™ 602 for Separation of Cs and Rb	14
3 Elution of Ag on Superlig™ 27 During IBC and PNL Tests	15

TABLES

1 Composition of the Test Solution	16
2 Separation of Barium from Test Solution Using SuperLig™ 601	17
3 Separation of Strontium from Test Solution Using SuperLig™ 601	18
4 Separation of Cesium from Test Solution Using SuperLig™ 602	19
5 Separation of Rubidium from Test Solution Using SuperLig™ 602	20
6 Separation of Pd ²⁺ from 450 ml of Test Solution Using a 10-g Column of SuperLig™ 27	21
7 Separation of Pd ²⁺ from 2000 ml of Test Solution Using a 10-g Column of SuperLig™ 27	22
8 Separation of Ru ³⁺ from 450 ml of Test Solution Using a 10-g Column of SuperLig™ II	23
9 Separation of Ag ⁺ From 450 ml of Test Solution Using a 10-g Column of SuperLig™ II	24
10 Ag ⁺ Found in Elutions From SuperLig™ 27 in Experiment 1a	25
11 Visual Characterization of Samples	26

INTRODUCTION

This report derives from a project to investigate the applicability of sequestering agents with high selectivity to mixed waste treatment. Such mixed wastes, which are largely solid and aqueous solutions of ionic compounds in which acutely hazardous and/or radioactive compounds/ions are present in low concentrations relative to nonradioactive ions, are a major focus of DOE's waste cleanup efforts. The conditions present in these wastes require a high degree of selectivity in the sequestering agent. Thus, separation technologies based on molecular recognition of ions are attracting attention.

Described herein are experiments that were performed with novel separations materials that have recently been developed and patented by IBC Advanced Technologies, Inc. The materials, known as SuperLig™ materials, consist of highly selective macrocyclic ligands covalently bound to the surface of silica gel particles. The purpose of the experiments was to evaluate the SuperLig™ materials for their ability to efficiently and selectively separate certain metal ions from a synthetic, nonradioactive nuclear waste solution. The materials tested were 1) SuperLig™ 601 for barium (Ba^{2+}) and strontium (Sr^{2+}) separation, 2) SuperLig™ 602 for cesium (Cs^+) and rubidium (Rb^+) separation, 3) SuperLig™ 27 for palladium (Pd^{2+}) separation, and 4) SuperLig™ II for silver (Ag^+) and ruthenium (Ru^{3+}) separation. The materials and separation procedures were developed by IBC and have been described in progress reports^(a) to Pacific Northwest Laboratory^(b) (PNL). These materials were obtained from IBC. Results of work at IBC as well as experiments performed by PNL staff in collaboration with Dr. Ronald Bruening of IBC are reported herein.

Results for separation of specific metal ions are discussed following a general description of experimental procedures.

(a) Bruening, R. L. 1990. "Exploratory Research in Application of Inorganically-Bonded Macrocyclics to Separation of Se, Zr, Pd, and Sn from Simulated Nuclear Waste." IBC Advanced Technologies, Inc., Final Report to PNL, June 14.

Bruening, R. L. 1990. "Exploratory Research in Application of Inorganically-Bonded Macrocyclics to Separation of Pd, Ru, and Rh from Simulated Single-Shell-Tank Waste." IBC Advanced Technologies, Inc., Final Report to PNL, September 27.

(b) Operated for the U.S. Department of Energy by Battelle Memorial Institute under Contract DE-AC06-76RLO 1830.

EXPERIMENTAL PROCEDURES

General

SuperLig™ materials (packed in glass columns) and the test solution were provided by IBC. The test solution was passed through the columns. Then, the columns were washed with water and/or diluted nitric acid before the sequestered elements were eluted. Flow rates of solutions into the columns were not controlled in these experiments, although IBC's June 1990 Final Report indicated flow rates were held at approximately 30 ml/min. Feed, effluent, wash and elutions were analyzed for elements of interest by atomic absorption (AA), atomic emission (AE) or inductively-coupled plasma (ICP) spectroscopy. Composition of the test solution is provided in Table 1. The specific washes and elutions for each column as well as analytical results are detailed below and tabulated in Tables 2 through 10. Listing of experimental samples as well as visual characterization of samples, feed, and column materials are contained in Table 11.

Separation of Ba^{2+} and Sr^{2+}

A 450-ml volume (approximately 30 bed volumes) of the synthetic test solution was fed into a column containing 10 g of SuperLig™ 601. The column was washed with 1 M HNO_3 (50 ml in one experiment and 100 ml in another).^(a) 0.1 M $\text{Na}_3\text{citrate}$ (100 ml) was added to elute Sr^{2+} followed by 0.03 M Na_4EDTA (100 ml in the one experiment and 75 ml in the other; both solutions were pH~10.5) to elute Ba^{2+} . Concentrations of Ba^{2+} and Sr^{2+} in the feed, effluent, wash, and elutions were determined by AA.

Separation of Cs^+ and Rb^+

30 ml of test solution that had been passed through SuperLig™ 601 were fed into a column containing 20 g of SuperLig™ 602. The column was washed with 45 ml of water, and 75 to 105 ml of 0.01 M HNO_3 was added to elute both Cs^+ and Rb^+ . Concentration of Cs^+ in feed, effluent, wash, and elutions was determined by AA and Rb^+ concentration was determined by AE.

Separation of Pd^{2+}

In the first run, Experiment 1a, 450 ml of test solution were fed into a column containing 10 g SuperLig™ 27. This volume contained ~1 mmol Pd^{2+} , representing ~33% of the 3 mmol ligand load

(a) In the experiments conducted at IBC, the column was washed with 16 M HNO_3 in between loading and unloading to remove Zr and other cations from the blank gel sites.

capacity. The column was washed with 1 M HNO_3 , followed by addition of a solution containing 1 M thiourea/1 M acetic acid to elute Pd^{2+} held in the column.^(a) In the second run, Experiment 4, 2000 ml of synthetic test solution were fed into the SuperLig™ 27 column. The feed volume contained >3 mmol Pd^{2+} needed to load the column to 100% load capacity. The column was washed with 5 M HNO_3 followed by water. A solution of 1 M thiourea/1 M acetic acid was added to elute Pd^{2+} .^(a) Concentrations of Pd^{2+} in thiourea/acetic acid elutions were determined by ICP.

Separation of Ru^{3+} and Ag^+

In this experiment, referred to as Experiment 1b, 450 mL of test solution that had been passed through SuperLig™ 27 were fed into a column containing 10 g of SuperLig™ II. This volume contained ~2.1 mmol Ru^{3+} and 0.1 mmol Ag^+ , representing ~73% of the 3-mmol ligand load capacity. The column was washed with 1 M HNO_3 followed by addition of a hot (60°C) solution containing 1 M thiourea/1 M acetic acid/0.02 M TiCl_3 to elute Ru^{3+} and Ag^+ .^(b) Concentration of Ag^+ in feed, effluent, wash, and elutions was determined by AA. Concentration of Ru^{3+} in the elutions was determined by ICP.

-
- (a) The IBC experiment washed the column with 16 M HNO_3 to remove Zr from the blank gel sites followed by an acetic acid wash to displace the nitric acid and prevent oxidation of thiourea. Pd was eluted with 1 M thiourea/1 M acetic acid.
- (b) In the IBC experiments, 9 M HBr was used to elute Ag.

RESULTS

Separation of Ba^{2+} and Sr^{2+}

Analytical results for SuperLig™ 601 used to separate Ba^{2+} and Sr^{2+} from synthetic waste solution (Experiments 2 and 5) as well as results contained in IBC's June 1990 Final Report are summarized in Tables 2 and 3 and depicted graphically in Figure 1. Data from the June 1990 Final Report indicate that Sr^{2+} and Ba^{2+} were efficiently removed (100%) from the test solution and selectively eluted with $\text{Na}_3\text{citrate}$ and Na_4EDTA , respectively; approximately 100% of the Sr^{2+} and 10% of the Ba^{2+} was eluted by the $\text{Na}_3\text{citrate}$ elution. The remainder of the Ba^{2+} was eluted by the Na_4EDTA . The report states that a 10-g column was used; in fact, however, a 20-g column was used.^(a) Data from Experiments 2 and 5 obtained from a 10-g column indicate that 90 to 98% of the Ba^{2+} and 73 to 86% of the Sr^{2+} were separated from the test solution. Virtually quantitative removal of Ba^{2+} and Sr^{2+} was observed until breakthrough. Depending on the experiment, 26 to 42% of the Sr^{2+} eluted with the wash, and the remainder with $\text{Na}_3\text{citrate}$. Approximately 35% of the Ba^{2+} was eluted with the wash and $\text{Na}_3\text{citrate}$, and between 50 and 60% with Na_4EDTA . Despite breakthrough, much of the Ba^{2+} eluted by Na_4EDTA was substantially free of Sr^{2+} and the other ions initially present in the feed solution. Only Zr was present in any significant quantity in the elutions which followed the wash step. Zr was not found in the IBC experiments because the columns were washed with 16 M nitric acid to remove Zr from the plain silica gel sites. Apparently, 5 M nitric acid, which was used in our experiments to wash the column, is not a strong enough acid to completely displace Zr from silica gel.

The feed volume (450 ml) contained ~1.1 mmol Ba^{2+} and 0.8 mmol Sr^{2+} , representing ~63% of the 3-mmol ligand load capacity.^(b) The binding constants of Superlig™ 601 for Ba^{2+} and Sr^{2+} are sufficiently large (10^4 and 10^3 , respectively)^(c) that greater than 90% of the load capacity should be usable if the column is well equilibrated with feed.^(d) That breakthrough occurred indicates that flow rates may have been too fast to permit equilibration with the column material or that channeling of the feed solution through portions of the column resulted in less than optimum contact between column material and test solution. Observations supporting the latter conclusion are 1) the column was dry packed and 2) gray,

(a) R. L. Bruening, IBC, Advanced Technologies, Inc. private communications (letters dated 4/15 and 6/25/91).

(b) IBC June 1990 Final Report.

(c) Equilibrium data supplied by IBC.

(d) The equilibrium load capacity, f , is calculated from metal ion concentration, $[M]$, using the equilibrium expression, $K = f/(1-f)[M]$.

vertical streaks developed in the lower half of the column, indicative of channel formation. In retrospect, smaller loadings of test solution, e.g., 250 to 300 ml (see Figure 1), would have given better separations with the 10-g quantity of column material.

Separation of Cs⁺ and Rb⁺

Analytical results for SuperLig™ 602 used to separate Cs⁺ and Rb⁺ from synthetic waste solution (Experiments 3 and 6) as well as corresponding results contained in IBC's June 1990 Final Report are summarized in Tables 4 and 5 and depicted in Figure 2. Data from the June 1990 Final Report show that Cs⁺ and Rb⁺ were quantitatively separated from the test solution and readily eluted with dilute HNO₃. The report states that a 10-g column was used in the experiments, however the experiments were in fact performed with a 20-g column.^(a) The report also states that 1) Rb⁺ and Cs⁺ could only be separated from 30 ml aliquots of the test solution by the column because of weak binding constants between the elements and ligand, and 2) the separation was almost chromatographic, also because the binding constants were weak. The 30-ml volume of test solution contained ~0.12 mmol Cs⁺ and 0.03 mmol Rb⁺, ~2.5% of the 6-mmol ligand load capacity.^(b) Given the concentration of Cs⁺ in the test solution and the binding constant for the ligand ($K = 50$),^(c) the maximum load capacity of the column should be 17%^(d) if complete equilibration were maintained during loading of the column.

Results from Experiment 3 differ from the IBC test run in that the column showed diminished load capacity as evidenced by breakthrough of Cs⁺ in the wash. This first run, however, may have conditioned the column, as the column exhibited improved performance on the second trial (Experiment 6), and the elution profile resembled the elution profile obtained by IBC. No breakthrough occurred during the 30-ml loading or the first 30-ml wash, which suggests that the 20-g column might handle a larger load volume. Good separation of Cs⁺ and Rb⁺ from the other elements in the test solution was observed; however, Cs⁺ and Rb⁺ could not be separated from each other.

Material balances on the two elements were essentially quantitative; however, the volume of elution containing Cs⁺ and Rb⁺ was approximately three times that of the volume of the original test solution feed to the column. Unlike the experiments with Sr²⁺ and Ba²⁺, a step-gradient elution using sequestering agents to facilitate release of Cs⁺ and Rb⁺ from the column could not be improvised because few sequestering agents bind Cs⁺ or Rb⁺ better than the silica-bound macrocycle on the column.

(a) R. L. Bruening, IBC, Advanced Technologies, Inc., private communications (letters dated 4/15 and 6/25/91).

(b) IBC June 1990 Final Report

(c) Equilibrium data supplied by IBC.

(d) This calculation neglects competition with Rb for the bound ligands. In practice, a smaller load capacity would be observed.

Alternatively, IBC suggests that it can take advantage of the greater affinity that the macrocycle has for Ba^{2+} ions. For example, Ba^{+2} solution, instead of 0.01 M nitric acid, should elute and concentrate Cs^+ and Rb^+ .

Separation of Pd^{2+} in Experiment 1a

Results for the separation of Pd^{2+} from a 450-ml synthetic test solution using SuperLig™ 27, as well as corresponding results contained in IBC's June 1990 Final Report, are summarized in Table 6. Data from the June 1990 Final Report indicate Pd^{2+} was efficiently separated (100%) from the test solution using a 10-g column and efficiently eluted (100%) with a 100-ml thiourea/acetic acid solution. Results from PNL experiments are comparable. Based on Pd^{2+} concentrations determined by ICP analysis of the eluted samples, mass balance for Pd^{2+} is 107%. The concentration of Pd^{2+} in the combined eluted samples is 1.06 ppm compared with 0.22 ppm in the feed solution. Thus, the Pd^{2+} was concentrated by a factor greater than four during the separation.

Total amounts of other elements present in the elutions were 8 mg Zr, 9 mg Ag^+ , and 7 mg Ru^{3+} . Zr is present for the same reasons previously discussed in the section on separations of Ba^{2+} and Sr^{2+} . Ru^{3+} is present because it also binds to SuperLig™ 27, and the amount of Pd^{2+} fed to the column was not sufficient to occupy all the binding sights in the column. Ag^+ was not expected to be present based on previous experiments performed by IBC. The Ag^+ may not have been bound to the column material but instead was deposited at the top of the column as a precipitate that was subsequently solubilized and eluted by the thiourea eluent. See the section below on the separation of Ag^+ in Experiment 1b.

Separation of Pd^{2+} in Experiment 4

Analytical results for Experiment 4 as well as corresponding results contained in IBC's June 1990 Final Report are summarized in Table 7. Data from the June 1990 Final Report indicate 3.16 mmol Pd^{2+} were separated from the 5.50 mmol Pd^{2+} fed into the column and subsequently eluted (100%) with 1 M thiourea/1 M acetic acid solution. The data for Experiment 4 are incomplete as the effluent, wash, and first elution samples were not analyzed for Pd^{2+} concentration; however, three elution samples, E7 through E9, for which analytical data are available, total 87 ml and account for 327 mg or 97% of the estimated 336 mg Pd^{2+} load capacity. Thus, based on the available data, Pd was concentrated by a factor of about 17. As observed in Experiment 1a, Zr(IV), Ru^{3+} and Ag^+ were eluted with Pd^{2+} . Some Ru^{3+} was still obtained because, although Ru^{3+} does not bind as well as Pd^{2+} with the SuperLig™ 27 ligand, it has slower kinetics of release, and therefore some of it still remained on the column even though the column was overloaded with Pd^{2+} .

Separation of Ru^{3+} in Experiment 1b

Analytical results for SuperLig™ II used to separate Ru^{3+} from synthetic waste solution are summarized in Table 8. No corresponding results are available from the June 1990 Final Report, although results from the September 1990 Final Report indicate 40 to 50% of Ru^{3+} bound by SuperLig™ II was recovered using a 1 M thiourea/1 M acetic acid solution. The September 1990 Final Report notes that SuperLig™ II bound Ru^{3+} strongly and released Ru^{3+} to other ligands slowly. The report also notes reduction of Ru^{3+} to Ru^{2+} with thiourea yielded a Ru-thiourea complex which eluted off the column; however, any Ru^{2+} which did not quickly form a complex with thiourea rebound to the SuperLig™ material and again became kinetically difficult to elute.

In Experiment 1b, TiCl_3 was added to the thiourea/acetic acid elution in an attempt to speed up reduction of Ru^{3+} to Ru^{2+} and hopefully facilitate formation of the Ru^{2+} thiourea complex. The new elution did not increase Ru recovery, as only 37% of the Ru eluted from the column compared with 40 to 50% eluted using just thiourea/acetic acid. The concentration of Ru in the 100 ml of eluent was 1.7 times greater than the feed solution. The observations here that Ru^{3+} can be partially extracted in a more concentrated form is promising. However, more work is needed to discover ways of obtaining complete recovery of Ru or to find other ligands that release Ru^{3+} more rapidly.

The SuperLig™ II column retained a dark bluish green color, indicating that Ru was still held in the column. Deep blue solutions are formed when chloro complexes of Ru^{3+} are reduced in HCl by Ti^{3+} . Addition of 16 M HNO_3 to the column at a later date yielded a blue-colored elution which turned yellow on standing overnight. Ru^{2+} chloro complexes are air sensitive and will oxidize to a yellow complex.^(a) The column material was a light, orange-brown color after the addition of the concentrated HNO_3 .

Separation of Ag^+ in Experiment 1b

Analytical results for SuperLig™ II used to separate Ag^+ from synthetic solution as well as corresponding results contained in IBC's Final Report are summarized in Table 9. Data from IBC's report indicate that Ag^+ was not bound by the SuperLig™ 27 column (used to remove Pd^{2+} from the feed solution), and that Ag^+ was all removed, within experimental error, from the Pd^{2+} -depleted test solution by the SuperLig™ II column. One hundred percent (100%) of the Ag^+ was recovered with 9 M HBr.

Experiments performed at PNL did not reproduce these experiments. According to IBC results (Table 10 and Figure 3), the effluent from SuperLig™ 27, which served as the feed solution for SuperLig™ II, should have contained the bulk of the Ag^+ concentration, as Ag^+ was supposed to pass

(a) Cotton, F. A., and G. Wilkinson, *Advanced Inorganic Chemistry*, 3rd Ed.; Interscience Publishers: New York; p. 1007.

through the column. Unlike the IBC experiments, the bulk of the Ag^+ , 6.2 mg (48%) of the 13.1 mg Ag^+ thought to be fed into the column, appeared in the elutions rather than the effluent from SuperLig™ 27.

The actual concentration of Ag^+ in the effluent from SuperLig™ 27 was not determined before the solution was fed into the SuperLig™ II column. The concentration of Ag^+ in the solution used to elute both Ru^{3+} and Ag^+ from the SuperLig™ II column accounted for an additional 0.5 mg Ag^+ , which still left approximately half of the Ag^+ thought to be fed into the columns unaccounted for. No Ag^+ turned up in the EDX analysis of SuperLig™ II material nor did any appear in an acid digestion of SuperLig™ 27 material, which suggests that any Ag^+ added to the columns had been removed.

Analysis of the sludge observed at the bottom of the feed solution container showed the sludge to contain approximately 22 mg/g Ag. Some of this sludge was observed to filter out at the tops of the columns as the test solution was added, and at times the glass wool plugs atop the SuperLig™ materials had to be removed and rinsed with distilled water to maintain reasonable flow rates through the columns. These observations are consistent with a conclusion that the amount of Ag^+ originally fed into the SuperLig™ was probably not 13.1 mg.

In an attempt to determine how much Ag^+ might actually have been in the solution that was passed through the SuperLig™ 27 column and subsequently fed into the SuperLig™ II column, the effluent in Experiment 4 was analyzed. The concentration of Ag^+ in the first 500 mL of feed effluent that passed through the column was ~1 mg/L. If the feed concentration of Ag^+ into SuperLig™ II in Experiment 1b is assumed to be 1 mg/L, the mass balance for Ag then becomes ~106%.

CONCLUSIONS

Some of the SuperLig™ experiments performed at PNL reproduced experiments performed at IBC; others did not. Separations of Pd^{2+} with SuperLig™ 27, Ru^{3+} with SuperLig™ II and Cs^+ and Rb^+ with SuperLig™ 602 were consistent with the experimental results obtained by IBC. Ba^{2+} and Sr^{2+} were separated from the test solution using SuperLig™ 601; however, separations were not as clean as reported by IBC. Premature breakthrough of the ions coupled with observations of gray vertical streaks through the lower half of the column after elution strongly suggests that channelling occurred, ostensibly due to inefficient packing of the column. Also, the IBC results were obtained using twice as much SuperLig™ material. Experiments for separation of Ag^+ were compromised because the Ag concentration in the test solution was uncertain as a result of precipitation. Thus, material balances were low for Ag . Ag^+ appeared to be sequestered by both SuperLig™ 2 and 27 columns, although more than likely the precipitate filtered out at the top of the SuperLig™ 27 column and was subsequently solubilized by the thiourea eluent.

Accordingly, the observed differences in PNL and IBC experiments appear to be mainly attributable to experimental procedures, rather than to failure of the SuperLig™ materials themselves. The technology is basically sound and works with varying degrees of success depending mainly on the affinity of the supported ligand for the metal ion of interest. Because the field of ion-recognition research is large and active, it is expected that suitable ligands will be discovered or synthesized to separate metal ions of interest with this technology. Certainly more development, testing, and evaluation is warranted.

Figure 1. Comparison of IBC and PNL Tests of Superlig™ 601 for Separation of Ba (□) and Sr (Δ): A, IBC Test Result (20-g column); B and C, PNL Experiments 2 and 5, Respectively (10-g column, see Appendix, Tables 2 and 3)

Figure 2. Comparison of IBC and PNL Tests of Superlig™ 602 for Separation of Cs (x) and Rb (O): A, IBC Test Result; B and C, PNL Experiments 3 and 6, Respectively (see Appendix, Tables 4 and 5)

Figure 3. Elution of Ag on Superlig™ 27 During IBC and PNL Tests: (A) IBC Final Report, (B) PNL Experiment 4 (see Table 10)

Table 1. Composition of the Test Solution^(a)

<u>Element/Specie</u>	<u>Target μg/ml</u>	<u>ICP μg/ml</u>	<u>AA/AE μg/ml</u>
Ru	450	464	*
Ba	350	303	350
Pd	310	219	*
Ag	20	32	29
Cs	550	550	550
Rb	80	*	84
Sr	150	150	160
Gd	2400	2025	*
Y	80	71	*
Zr	700	723	*
Mo	700	645	*
Rh	100	92	*
Cd	20	16	*
Sn	10	25	*
Sb	3	11	*
Te	100	150	*
La	250	242	*
Ce	500	417	*
Pr	250	219	*
Nd	800	1457	*
Sm	200	81	*
Eu	30	34	*
U	1260	1054	*
PO ₄ ³⁻	400	*	*
Fe	300	40	*
Cr	50	44	*
Ni	20	18	*
Na	High	87115	*

(a) The test solution was designed to represent a nuclear waste stream, e.g., PUREX raffinate, in 5 M nitric acid.

* No analysis performed.

Table 2. Separation of Barium from Test Solution Using SuperLig™ 601

	IBC Report			PNL Experiment 2			PNL Experiment 5		
	$\mu\text{g/mL}$	mL	μg	ID	$\mu\text{g/mL}$	mL	$\mu\text{g/mL}$	mL	μg
Original Solution	350	450	<u>157,500</u>	Feed	350	450	350	450	<u>157,500</u>
Effluent	ND ^a	450		B1	<1	100	4.0	300	1,200
				B2	<1	100			
				B3	<1	100			
				B4	35	150			
16 M HNO ₃ 1 M HNO ₃	ND ^a	50		B5	110	50	140	100	14,000
	ND ^a	50							
0.1 M Na ₃ citrate	<1	25		B6(b)	110	25	100	25	2,500
	17	25	425	B7	440	25	480	50	24,000
	230	25	5,750	B8	770	25	520	25	13,000
	388	25	9,700	B9	770	25			
0.03 M Na ₄ EDTA	2519	25	62,975	B10	840	25	640	25	16,000
	2978	25	74,450	B11	1700	25	960	50	48,000
	133	25	3,325	B12	680	50			
	21	25	525	B13	25	50			
		<u>750</u>	<u>157,500</u>			<u>750</u>	230	<u>775</u>	<u>143,700</u>
			(100%)						(91%)

(a) ND: None Detected.

(b) B6-B10 contained Zr in the following respective quantities ($\mu\text{g/mL}$): 252, 363, 66, 17, 15.

(c) F4 contained 121 $\mu\text{g/mL}$ Zr.

Table 3. Separation of Strontium from Test Solution Using SuperLig™ 601

	IBC Report			PNL Experiment 2			PNL Experiment 5		
	$\mu\text{g/mL}$	mL	μg	ID	$\mu\text{g/mL}$	mL	$\mu\text{g/mL}$	mL	μg
Original Solution	184 ^a	450	<u>82,800</u>	Feed	60	450	160	450	<u>72,000</u>
Effluent	ND ^a	450		B1	<1	100	4.6	300	1,400
				B2	<1	100			
				B3	<1	100			
				B4	69	150			
16 M HNO ₃ 1 M HNO ₃	ND ^a	50		B5	370	50	290	100	29,000
0.1 M Na ₃ Citrate	357	25	8,925	B6	530	25	230	25	5,700
	1734	25	43,350	B7	990	25	310	50	15,000
	1392	25	34,800	B8	190	25	12	25	300
	107	25	2,675	B9	<10	25			
0.03 M Na ₄ EDTA	<1	25	<u>89,750</u>	B10	<5	25	32	25	80
	<1	25		B11	<5	25			
	<1	25		B12	<5	50			
	<1	25		B13	<5	50			
		<u>750</u>				<u>750</u>		<u>775</u>	<u>68,480</u>
			(108%)						(95%)
									<u>71,700</u>
									(100%)

(a) ND: None Detected.

(b) B6-B10 contained Zr in the following respective quantities ($\mu\text{g/mL}$): 252, 363, 66, 17, 15.

(c) F4 contained 121 $\mu\text{g/mL}$ Zr.

Table 4. Separation of Cesium from Test Solution Using SuperLig™ 602

	IBC Report			PNL Experiment 3				PNL Experiment 6			
	$\mu\text{g/mL}$	<u>mL</u>	<u>μg</u>	ID	$\mu\text{g/mL}$	<u>mL</u>	<u>μg</u>	ID	$\mu\text{g/mL}$	<u>mL</u>	<u>μg</u>
Feed Solution (Ba and Sr removed)	592	30	<u>17,760</u>	B1	530	30	<u>15,900</u>	F2	550	30	<u>17,000</u>
Effluent	<5	30		C1	<2	30		G1	2.2	30	66
H ₂ O	<5	15		C2	21	15	310	G2	2.5	30	75
	28	15	420	C3	190	15	2,900				
				C4	220	15	3,300	G3	140	15	2,100
0.01 M HNO ₃	299	30	8,970	C5	170	15	2,500	G4	240	15	3,600
	250	30	7,500	C6	150	30	4,500	G5	220	30	6,600
	20	30	600	C7	54	30	1,600	G6	130	30	3,900
	4	30	120	C8	15	30	450				
		180	<u>17,610</u>			180	<u>15,560</u>			150	<u>16,341</u>
			(99%)				(98%)				(99%)

Table 5. Separation of Rubidium from Test Solution Using SuperLig™ 602

	IBC Report			PNL Experiment 3			PNL Experiment 6				
	µg/mL	mL	µg	ID	µg/mL	mL	µg	ID	µg/mL	mL	µg
Feed Solution (Ba and Sr removed)	85a	30	<u>2,550</u>	B1	69	30	<u>2,070</u>	F2	88	30	<u>2,640</u>
Effluent	<5	30		C1	<1	30		G1	0.3	30	9.0
H ₂ O	<5	15		C2	1.7	15	25	G2	<0.4	30	
	3	15	45	C3	17	15	250				
				C4	26	15	390	G3	11	15	170
0.01M HNO ₃	36	30	1,080	C5	25	15	370	G4	34	15	510
	44	30	1,320	C6	22	30	660	G5	41	30	1,200
	4	30	120	C7	8.2	30	250	G6	24	30	720
	<1	30		C8	2.5	30	75				
		<u>180</u>	<u>2,565</u>			<u>180</u>	<u>2,020</u>			<u>150</u>	<u>2,609</u>
			(101%)				(98%)				(99%)

Table 6. Separation of Pd²⁺ from 450 ml of Test Solution Using a 10-g Column of SuperLig™ 27

	IBC Report			PNL Experiment 1a ^(a)			
	<u>ug/mL</u>	<u>mL</u>	<u>ug</u>	<u>ID</u>	<u>ug/mL</u>	<u>mL</u>	<u>ug</u>
Original Solution	325 ^(b)	450	<u>146,250</u>	Feed	219	450	<u>98,600</u>
Effluent	<1	450		(b)			
16 M HNO ₃ + 0.1 M Acetic Acid	<1	75					
1 M HNO ₃				(b)		100	
1 M Thiourea/0.1 M Acetic Acid	2925 27	50 50	146,250 1,350				
1 M Thiourea, 1 M Acetic Acid				D1 ^(c)	2120	50	106,000
				D2 ^(d)	3	<u>50</u>	<u>150</u>
		<u>625</u>	<u>147,600^(e)</u>			550	<u>106,150</u>
			(101%)				(107%)

(a) IPC analysis provided by IBC.

(b) Effluent and wash from SuperLig™ 27 fed directly into SuperLig™ II; concentration of Pd²⁺ in effluent and wash undetermined.

(c) Other metal ions present (μg/mL): Ag, 67; Zr, 70; Ru, 49.

(d) Other metal ions present (μg/mL): Ag, 19; Zr, 14; Fe, 17.

(e) 5.8 mg Ru also present.

Table 7. Separation of Pd²⁺ from 2000 ml of Test Solution Using a 10-g Column of SuperLig™ 27

	IBC Report			PNL Experiment 4 ^(a)			
	<u>µg/mL</u>	<u>mL</u>	<u>µg</u>	<u>ID</u>	<u>µg/mL</u>	<u>mL</u>	<u>µg</u>
Original Solution	274 ^(b)	2000	<u>580,000</u>	Feed	219	2000	<u>438,000</u>
Effluent	<5	500		E1		500	
	32	500	16,000	E2		500	
	192	500	96,000	E3		500	
	250	300	75,000	E4		500	
	258	100	25,800				
	268	100	26,800				
100 ml 5 M HNO ₃ + 15 ml H ₂ O				E5		115	
150 ml 16 M HNO ₃ + 25 ml 1 M Acetic Acid	43	175	7,525				
1 M Thiourea/1 M Acetic Acid	6200	50	310,000	E6		13	
	515	50	25,750	E7 ^(b)	10,910	28	305,480
				E8	607	35	21,245
				E9	18	24	432
	8	<u>50</u>	<u>400</u>				
		2325	<u>336,150</u>			2215	<u>324,150</u>
			(112%) ^(c)				(97%) ^(c)

(a) ICP analysis provided by IBC.

(b) Other metal ions present (µg/mL): Ag, 180; Zr, 214; Ru, 34.

(c) Percent of column capacity (3 m equiv).

Table 8. Separation of Ru³⁺ from 450 ml of Test Solution Using a 10-g Column of SuperLig™ II

	PNL Experiment 1b ^(a)		
	<u>µg/mL</u>	<u>mL</u>	<u>µg</u>
Original Solution	464	450	<u>208,800</u>
SuperLig™ II			
A1 Effluent		100	
A2 Effluent		100	
A3 Effluent		100	
A4 Effluent		150	
A5 1 M HNO ₃		100	
A6 1 M Thiourea, 1M	717 ^(c)	50	35,850
A7 Acetic Acid, 0.02M TiCl ₃	836	50	41,800
SuperLig™ 27			
D1 1 M Thiourea,	98	50	4,900
D2 1 M Acetic Acid, 0.02 M TiCl ₃	7	50	850
			<u>83,400</u>
			(40%) ^(b)

(a) IPC analyses provided by IBC.

(b) Other metal ions present (µg/mL): Zr, 33; Pd, 34.

(c) Ru visibly remaining in column (deep blue color); 16M HNO₃ removed more of the Ru.

Table 9. Separation of Ag⁺ From 450 ml of Test Solution Using a 10-g Column of SuperLig™ II

	IBC Report			PNL Experiment 1b			
	<u>µg/mL</u>	<u>mL</u>	<u>µg</u>	<u>ID</u>	<u>µg/mL</u>	<u>mL</u>	<u>µg</u>
Feed Solution (Pd ²⁺ removed)	5	450	<u>2,250</u>	(a)	29	450	<u>13,050</u>
Effluent	ND ^(b)	450		A1	0.06	100	6
				A2	≤0.05	100	
				A3	<0.05	100	
				A4	<0.05	150	
1M HNO ₃				A5	<0.05	100	
16M HNO ₃	ND ^(b)	100					
H ₂ O	ND ^(b)	100					
Hot 1 M Thiourea, 1M Acetic Acid, 0.02M TiCl ₃				A6 ^(c)	9.3	50	470
				A7	<0.5	50	
9M HBr	45	<u>50</u>	<u>2,250</u>				
		700	<u>2,250</u>			650	<u>476^(d)</u>
			(100%)				(3.6%) ^(e)

(a) 450 ml effluent from SuperLig™ 27 fed directly into SuperLig™ 2; concentration of Ag in effluent undetermined.

(b) ND: None Detected.

(c) Other metal ions present (µg/mL): Zr, 33; Pd, 34.

(d) Samples D1 and D2 (elutions from SuperLig™ 27) contained 5000 µg and 1100 µg Ag respectively (47% Ag accounted for in these two samples)

(e) In Experiment 4, concentration of Ag after passing through SuperLig™ 27 was ~1 µg/mL. If effluent concentration in Experiment 1b above is 1 µg/mL, then mass balance is ~106%.

Table 10. Ag⁺ found in Elutions From SuperLig™ 27 in Experiment 1a

	IBC Report			PNL Experiment			
	<u>ug/mL</u>	<u>mL</u>	<u>ug</u>	<u>ID</u>	<u>ug/mL</u>	<u>mL</u>	<u>ug</u>
Original Solution	5	2000	<u>10,000</u>	Feed	29	2000	<u>58,000</u>
Effluent	3	500	1,500	E1	0.96	500	480
	3	500	1,500	E2	1.5	500	750
	4	500	2,000	E3	1.1	500	550
	5	300	1,500	E4	1.9	500	950
	7	100	700				
	4	100	400				
100 ml 5 M HNO ₃ + 15 ml H ₂ O				E5	5.6	115	93
150 ml 16 M HNO ₃ + 25 ml Acetic Acid	4	175	700				
1 M Thiourea/1 M Acetic Acid	<1	50		E6	16	13	210
	<1	50		E7	120	28	3,400
				E8	6.5	35	230
				E9	<0.5	24	
	<1	<u>50</u>					
		2325	<u>8,300</u>			2215	<u>7,210</u>
			(83% Feed) (0% Eluent)			(5.8% Feed, Wash) (6.7% Eluent)	

Table 11. Visual Characterization of Samples

Sample ID	Appearance
Experiment 1b	
A-1	Pale gold, no precipitate
A-2	Pale gold, no precipitate
A-3	Pale gold, no precipitate
A-4	Pale gold, no precipitate
A-5	Pale peach, no precipitate
A-6	Green solution, olive crystals on walls, precipitate on bottom, possibly trans-RuCl ₄
A-7	Dark aqua-blue and white precipitate on bottom
Experiment 3	
B-1	Dark brown clear liquid, no precipitate
B-2	Dark brown clear liquid, no precipitate
B-3	Dark brown clear liquid, no precipitate
B-4	Dark brown clear liquid, no precipitate
B-5	(lighter) dark brown clear liquid, no precipitate
B-6	pale peach; very fine, small amount black precipitate on bottom
B-7	Clear, colorless, no precipitate
B-8	Clear, colorless; small amount fine white precipitate on bottom
B-9	Clear, colorless, no precipitate
B-10	Clear, colorless, small amount of fuzzy gray precipitate on bottom
B-11	Clear, colorless, ~1/4" fuzzy gray precipitate on bottom
B-12	Clear, colorless, fine gray precipitate adhered to bottom
B-13	Pale gold; fine, gray flakes on bottom
C-1	Tan, clear liquid, very fine tan precipitate on bottom
C-2	Gold liquid, flaky precipitate on bottom
C-3	Dark gold liquid, no precipitate
C-4	Dark gold liquid, no precipitate
C-5	Tan liquid, no precipitate
C-6	Pale, peach, fluffy white/tan precipitate on bottom
C-7	Clear, colorless, flaky precipitate on bottom
C-8	Clear, colorless, very fine white precipitate on bottom
Experiment 1a	
D-1	Dark brown liquid, ~1/4" gold precipitate on bottom
D-2	Pale gray solution, fine white precipitate on bottom

Table 11. (contd)

Sample ID	Appearance
Experiment 4	
E-1	Dark brown liquid, no precipitate
E-2	Dark brown liquid, no precipitate
E-3	Dark brown liquid, no precipitate
E-4	Dark brown liquid, no precipitate
E-5	(lighter) dark brown liquid, tan precipitate on bottom
E-6	light tan liquid, brown and tan precipitate on bottom
E-7	Orange-brown liquid, crystals (same color) on bottom, yellow precipitate as well (orange crystals possibly palladium diacetate)
E-8	Golden orange liquid, fine white precipitate adhered to bottom
E-9	Yellow liquid, fine white precipitate adhered to bottom
Experiment 6	
F-1	Dark brown solution
F-2	Dark brown solution
F-3	(lighter) dark brown solution
F-4	Pale, peach solution, very fine white precipitate on bottom
F-5	Clear, colorless solution
F-6	Clear, colorless solution
F-7	Clear, colorless solution
F-8	Clear, colorless solution, very fine precipitate on bottom
F-9	Clear, colorless solution
G-1	Clear, colorless solution
G-2	(light) dark brown solution, no precipitate
G-3	Dark brown solution, no precipitate
G-4	Dark brown solution, no precipitate
G-5	Pale brown solution, no precipitate
G-6	Clear, colorless solution
Columns	
SuperLig™ 27	Gold color
SuperLig™ 2	Red brown and aqua blue; light brown after cleaning with 16 M HNO ₃
SuperLig™ 601	Light gray with dark gray vertical streak on side of lower half of column
SuperLig™ 602	Light gray

DISTRIBUTION

No. of
Copies

No. of
Copies

OFFSITE

- 12 DOE/Office of Scientific and
Technical Information

R. L. Bruening
IBC, Advanced Technologies, Inc.
P.O. Box 656
Provo, UT 84603

ONSITE

- DOE Richland Field Office

S. Wolf

5 Westinghouse Hanford Company

K. D. Boomer, H5-49
J. J. Holmes, L5-55
J. A. Hunter, L5-31
R. G. Cowan, L5-31
J. P. Sloughter, T6-07

ONSITE

- 33 Pacific Northwest Laboratory

S. A. Bryan, P7-25
D. M. Camaioni, K2-10 (15)
N. G. Colton, P8-38
J. L. Cox, P8-38
B. P. Hay, K6-81
L. K. Holton, Jr., P7-43
M. A. Lilga, P8-38
G. J. Lumetta, P7-35
J. R. Morrey, P7-14
J. L. Straalsund, K1-79
M. R. Thompson, P7-07
D. W. Wester, P7-25
K. D. Wiemers, P7-14
Publishing Coordination
Technical Report Files (5)

END

DATE
FILMED
314192

I