DOE/ER/13710--T1

DE92 009045

CELLULAR ENERGY METABOLISM

Final Technical Report
for Period May 1, 1987 to April 30, 1991

Michael Glaser
Department of Biochemistry
University of Illinois
Urbana, Illinois 61801

June 1991

Prepared for
The U.S. Department of Energy
DE-FG02-87ER13710

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

APPENDIX: PROGRESS REPORT

Summary: Studies have been carried out on adenylate kinase which is an important enzyme in determining the concentrations of the adenine nucleotides. An efficient method has been developed to clone mutant adenylate kinase genes in E. coli. Site-specific mutagenesis of the wild type gene also has been used to obtain forms of adenylate kinase with altered amino acids. The wild type and mutant forms of adenylate kinase have been overexpressed and large quantities were readily isolated. The kinetic and fluorescence properties of the different forms of adenylate kinase were characterized. This has led to a new model for the location of the AMP and ATP bindings sites on the enzyme and a proposal for the mechanism of substrate inhibition. Crystals of the wild type enzyme were obtained that diffract to at least 2.3 Å resolution. Experiments were also initiated to determine the function of adenylate kinase in vivo. In one set of experiments, E. coli strains with mutations in adenylate kinase showed large changes in cellular nucleotides after reaching the stationary phase in a low phosphate medium. This was caused by selective proteolytic degradation of the mutant adenylate kinase caused by phosphate starvation.

The overall objectives of this research were to determine the structure and dynamic properties of adenylate kinase and to understand its role in determining the concentrations of the adenine nucleotides in vivo. Research has been carried out on all the specific aims of the original research application. One specific aim was to isolate adenylate kinase from wild type and mutant strains of E. coli.

The isolation of adenylate kinase has been made much easier by using the cloned gene to overproduce the enzyme in E. coli. The wild type gene was cloned onto the low copy number plasmid RP4 using Mu-mediated transposition and screening for complementation of a temperature-sensitive adenylate kinase (adk) mutant. Five positive colonies were obtained that contained the hybrid RP4:Mucts adk⁺ plasmid. These strains showed 3- to 5-fold increases in adenylate kinase activity (under standard assay conditions) and the enzyme was no longer temperature-sensitive indicating the presence of the wild type adk gene. The hybrid plasmid from one strain was isolated and used to subclone the adenylate kinase gene.

Complete HindIII digestion of the plasmid gave more than six fragments that were each larger than 10 kb. The digest was self-ligated and used to transform KG10, a recA1, adk-6 temperature-sensitive strain. Two plasmids of interest were recovered that conferred resistance to ampicillin and tetracycline. The smallest plasmid (28 kb) designated pLP315, contained a single HindIII site and did not complement the adk-6 temperature-sensitive gene of KG10. The second plasmid, designated pLP315-C, was made up of pLP315 with a 13 kb HindIII insert (fragment C) that carried the adk⁺ gene. KG10 (adk-6) produced a mutant adenylate kinase with a

lower mobility on SDS-polyacrylamide gels whereas other adk mutants [e.g., KG3 (adk-31) and CV2(adk-2)] produced thermolabile adenylate kinases with the same mobility as the wild type enzyme (Fig. I). The introduction of pLP315-C into KG10 led to approximately a 3-fold overproduction of the wild type adenylate kinase in the mutant background (Fig. I, lanes 3 and 4), indicating the presence of adk⁺ on the plasmid. Subsequent subcloning of the 13 kb HindIII fragment into pUC13 (designated pLP105) and a further trimming to a 1.4 kb ClaI fragment into pBR322 (designated pLP101) led to approximately a 60-fold overproduction of the wild type adenylate kinase (Fig. I, lanes 1 and 2).

Fig. I. Western blot analysis of adenylate kinases. 50 μ g of total protein were analyzed from crude extracts of each strain. Lane 1: LP101 (adk⁺ on pBR322 in KG10); lane 2: LP105 (adk⁺ on pUC13 in KG10); lane 3: LP8 (adk⁺ on pLP315 in KG10); lane 4: LP3 (adk⁺ on RP4 in KG10); lane 5: KL218 (adk⁺); lane 6: CV2 (adk-2); lane 7: KG3 (adk-31); lane 8: KG10 (adk-6). The arrowhead indicates the mutant adenylate kinase from KG10 (in lanes 1-4 and 8). Note the other mutant adenylate kinases (lanes 6 and 7) have the same mobility as the wild type adenylate kinase (lane 5). Also note the adk⁺ gene cloned on the low copy number plasmid (pLP315-C) was expressed in KG10 (lanes 3 and 4) and was greatly overproduced when cloned into high copy number plasmid vectors (lanes 1 and 2).

Some strains with high copy number plasmids containing the adenylate kinase gene grew slowly and appeared sick while other strains grew fairly well and overproduced the enzyme over 60-fold. Starting with a specific activity over 60-fold higher allows adenylate kinase to be purified in essentially one step by affinity chromatography. The enzyme was bound to a Gel Blue A column and was specifically eluted with the inhibitor Ap₅A. The enzyme was greater than 95% pure at this stage and the remaining contaminating proteins were removed by gel filtration on Sephadex G-75. Large quantities of the enzyme can now be routinely purified by this procedure.

Since a number of independently isolated temperature-sensitive adk mutants have been isolated, it is desirable to characterize the properties of these altered enzymes. The first step was to clone the adk genes in order to facilitate isolation of the enzymes and to sequence the genes. Often when a wild type gene is available, cloning of mutant alleles is done by in vivo homologous DNA recombination (5,11). This method, though theoretically simple, has its limitation especially since the frequency of homogenotization can be lower than 0.02% when applied to small segments of DNA cloned into conventional high copy number plasmid vectors (5,2). Screening of thousands of colonies is necessary to recover a few desired mutant phenotypes which is laborious for mutants with no method for positive selection such as temperature-sensitive mutants. Consequently, a simple and efficient method was worked out to substantially increase the frequency of homogenotization and identify the temperature-sensitive homogenotes (7).

In order to clone the adk-6 gene, a plasmid carrying the wild type gene was placed in a host containing the mutant allele, i.e. KG10. Since homologous recombination is recA⁺ dependent, a plasmid carrying the recA⁺ gene (either pLC24-31 or pLC21-33) was first used to transform KG10 followed by introduction of pLP315-C through conjugation. The presence of both plasmids in KG10 was selected by growth on plates containing mitomycin c (for the recA⁺ plasmid) and tetracycline (for pLP315-C) and confirmed by the pattern of restriction enzyme digestion. After restreaking the cells on the plates containing mitomycin C and tetracycline at 30°C to allow homologous genetic recombination to proceed, the cells were resuspended in LB-broth containing tetracycline. Serial dilutions were made and the cells were replica plated on nitrocellulose filter discs. A frequency of 15% homogenotization was obtained. The temperature-sensitive candidates were restreaked on plates with streptomycin and tetracycline to select KG10 containing pLP315-C* (now carrying the adk-6 gene instead of the adk⁺ gene). Restriction analysis showed that the recA⁺ plasmid was lost without the mitomycin c selection which made isolation of pLP315-C* easier.

Four temperature-sensitive isolates of KG10[pLP315-C*] (designated LP11 to LP14) were examined on Western blots. They showed a few-fold overproduction of the mutant adenylate kinase with no trace of the wild type enzyme as expected (Fig. II, lanes 2 to 5). Two randomly picked temperature-resistant colonies that arose during the isolation of KG10[pLP315-C*] were examined as well. One had a few-fold more wild type enzyme than mutant enzyme, as expected for KG10[pLP315-C] that had not undergone recombination and homogenotization, so the wild type gene remained on the plasmid and the mutant gene remained on the bacterial chromosome (Fig. II, lane 8). The other temperature-resistant colony, however, was just the opposite with a few fold more mutant enzyme than wild type enzyme (Fig. II, lane 7). This is consistent with having the wild type gene on the chromosome and the mutant gene on the plasmid which could be an intermediate in the homogenotization process.

The pLP315-C* plasmid carrying the adk-6 gene was isolated from LP12 and found to have the same restriction enzyme pattern as the

original plasmid carrying the wild type gene, indicating homologous DNA recombination. The 13 kb HindIII fragment (C*) was then inserted into the HindIII site of pUC19 which could be used to produce very large amounts of the mutant adenylate kinase. The resulting plasmid was used as a template for nucleotide sequencing.

Fig. II. Western blot analysis of adenylate kinases from strains carrying adk-6 plasmids obtained from replica plating. 50 μ g of total protein from each crude extract was analyzed. Lane 1: KG10 (adk-6); lanes 2 to 5: LP11 to LP14 (KG10[pLP315-C/adk-6]); lane 6: KL218 (adk⁺); lane 7: KG10[pLP315-C/adk-6] with adk-6 on the chromosome replaced by adk⁺ as a result of homogenotization. lane 8: KG10[pLP315-C/adk⁺]. TS stands for the temperature-sensitive mutant enzyme and WT stands for the wild type enzyme.

The sequencing covered the complete coding region along with the 5' and 3' flanking regions of the adk-6 gene. One point mutation was found in the adk-6 gene and this was a A-T to T-A transversion in the middle of the gene to give an a substitution of glutamine-107 for leucine-107 (Fig. III).

The copy number of the plasmid used for gene homogenotization could be a crucial factor in increasing the efficiency (9). The low copy number plasmid vectors commonly available are not ideally suited for use in gene homogenotization. One of the most commonly used low copy number plasmids, pSC101 (3) has a copy number of eight (9). Other plasmids such as pRK290 or pRK404 that are constructed from RK2 (4) also have copy numbers of 8-10/cell as measured by adenylate kinase overproduction. The 28 Kb plasmid, pLP315 derived from RP4 in this work is a low copy number (1 to 3 copies/cell as judged by adenylate kinase activity), broad host range and self-transmissible cloning vector. This plasmid carries both ampicillin and tetracycline resistance, and has single HindIII, EcoRI and BamHI restriction sites which allow cloning of rather big inserts up to 40 Kb. The plasmid pLP315 and its derivatives are small enough so that they can be easily isolated by routine plasmid minipreps (8) and they can be readily transformed into commonly used strains for transformation such as JM83 and

HB101. They also can be subsequently introduced into other desired host strains through efficient conjugation. Thus, pLP315 is a potentially very useful low copy number cloning vector.

Fig. III. Sequencing of the point mutation in the adk-6 gene. Both the adk⁺ (on pLP101) and the adk-6 (on pLP202) genes were sequenced with five oligodeoxynucleotide primers. Three primers were made in order to sequence from the 5' end of the gene (5'-CTCGCCATTAACCG, 5'-AGGCTCAGTTCATC and 5'-TTCCTGTTGGACGG) and two primers to sequence from the 3' end of the gene (5'-CCGGCCTGAGATTGC and 5'-CTTCAGCAACCGGC). Panel A: partial nucleotide sequence of the adk⁺ gene, CTG codes for leucine. Panel B: a point mutation in the adk-6 gene, CAG codes for glutamine.

Another specific aim given in the original application involved using the cloned gene to vary the level of adenylate kinase in the cell in order to determine the effects on the concentrations of adenine nucleotides and on cell metabolism. By inserting the gene into low and high copy number plasmids the amount of adenylate kinase in the cell could be varied. Some experiments have been carried out to determine the level of adenine nucleotides in these cells by labeling the nucleotides with ³²[P]phosphate, extracting the nucleotides and separating them by thin-layer chromatography. The nucleotide spots are then cut out and counted. The experiments showed that increasing the amount of adenylate kinase caused significant changes in the adenine nucleotides and the mass action ratio (see Table I on page 11 of the grant application). There were also differences observed between stationary and logarithmically growing cells.

Further studies on the *in vivo* behavior of adenylate kinase were side-tracked because of two things. The first involved the behavior of strains with mutant adenylate kinases when they reached the stationary phase. Very large changes took place in the cellular adenine nucleotide concentrations, as determined by [³²P]phosphate labeling, compared with the wild type strain (8). This was very exciting. Further investigation lead to the observation that the mutant adenylate kinases, in contrast to the wild type, were being proteolyzed when the cells reached stationary phase (Fig. IV). This was triggered by phosphate starvation since

a low phosphate medium was being used to obtain high efficiency labeling with [32 P]phosphate. Increasing the phosphate concentration in the medium eliminated the proteolyses. The selective proteolysis caused by phosphate starvation represents a novel observation, but it was not exactly what we were looking for in terms of how adenylate kinase alters adenine nucleotides concentrations. Future studies will be carried out at a phosphate concentration where the enzymes are stable.

Fig. IV Western-blot analysis showing the loss of adenylate kinase from adk mutant strains in the stationary phase. Cells were grown at 30°C in 56-LP minimal medium. (A) 50 μ g of total protein from the log and stationary phase crude extracts of CV2(adk-2) (lanes 1 and 2), KG3(adk-31) (lanes 3 and 4), KG10(adk-6) (lanes 5 and 6) and KL218(adk+) (lanes 7 and 8), respectively, were analyzed.

The second thing that side-tracked the in vivo work was the attempts made to produce mutants using site-specific mutagenesis with altered kinetic properties. The initial choice of which mutants to make was based on the X-ray structure of the pig muscle enzyme and the location of the AMP and ATP binding sites assigned by Pai et al. (10). As mentioned previously this assignment turned out to be incorrect and caused more effort to be placed on the last specific aim which involved the structure and dynamic properties of adenylate kinase.

The x-ray crystallography experiments have proceeded quite rapidly (1). Two methods have been used to crystallize the protein. The first method, used to narrow the range of possible crystallization conditions, was vapor diffusion, or more specifically, the hanging drop method. The second method, used to produce large crystals, is free interface diffusion. Here, an interface is created between the protein solution (low salt concentration and low density) and the precipitant solution (high salt concentration and high density). The slow diffusion across the interface initiates crystal growth. Crystals of E. coli adenylate kinase suitable for x-ray analysis were obtained by placing 7.1 μ l of 3.6 M ammonium sulfate, 2 mM MgCl, 0.1 M imidazole pH 7.0 into a depression slide cavity. 5 μ l of 26 mg/ml adenylate kinase, 0.5 M ammonium sulfate, 2 mM Ap_5A , and 0.23 M imidazole pH 7.9 was carefully layered over the high salt drop. The depression was sealed with a silicon-coated coverslip and

petroleum jelly. Crystals appeared within one week and grew to dimensions of 0.7 x 0.12 x 0.06 mm in about two weeks at 1°C. These crystals are typically rectangular bars with well-developed faces.

The unit cell dimensions as determined by diffractometry are: $a = 73.2 \text{ \AA}$, $b = 85.0 \text{ \AA}$, $c = 79.2 \text{ \AA}$. Based on the solvent content of the two known adenylate kinase structures and a calculated V_m of $2.6 \text{ \AA}^3/\text{dalton}$, there are probably eight molecules per unit cell and two molecules per asymmetric unit. Diffraction can be seen on small angle precession photos to at least $2.3 \text{ \AA}$ resolution. Systematic absence along the 100 and 010 axes, and the lack of such absences along the 001 axis, determine the space group to be of $P2_12_12$.

Crystals of diffraction quality also have been obtained in the presence of nucleotide substrates. These are in a different crystal form from the other ones, but they have not been further characterized as yet.

Several mutations have been generated in the adenylate kinase gene and the enzymes have been studied kinetically and by fluorescence spectroscopy. Three mutations are of particular interest and their characterization had led to a new model for the location of the AMP and ATP binding sites and also suggest a new mechanism for AMP substrate inhibition (6). Phe86 and tyr133, which are in two locations on the enzyme with each one having close contact with the adenine moieties of Ap_5A according to previous crystallographic results, have been independently changed to trp and other amino acids. The temperature-sensitive adk-6 mutation resulted from leu107 being changed to gln (7). Leu 107 is part of the central β -sheet and is near the original AMP site by crystallographic analysis while phe86 was near the original ATP site.

The mutagenesis of phe86 and tyr133 was done using oligodeoxynucleotide primers in M13mp18. The complete coding region of the two genes was sequenced to insure that the proper mutations were obtained and no secondary mutations were generated. The genes were put back in pUC18 in order to overproduce and isolate the enzymes. Both the trp86 and the trp133 enzymes were active and could be purified using the procedure worked out for the wild type enzyme.

Kinetic results for the altered adenylate kinases are summarized in Table A. The decrease in V_{max} for the wild type adenylate kinase at higher concentration of AMP was due to the AMP substrate inhibition. The phe86-trp mutation caused a 3- to 6-fold increase in the apparent K_M value for ATP at low (200 μM) and high (10 mM) AMP concentrations. However, the trp86 mutant enzyme did not show substrate inhibition at high concentrations of AMP and showed a dramatic increase (~44-fold) in the K_M for AMP. In contrast, the trp133 mutation resulted in little change in its kinetic properties as compared with the wild type enzyme. The gln107 enzyme showed a 6-fold increase in the K_M for ATP, a 7-fold

decrease in the $V_{\max}$, but with the K_M for AMP and AMP substrate inhibition unperturbed.

Table A. Comparison of the Kinetic Properties of Mutant Adenylate Kinases with the Wild Type Adenylate Kinase

	<u>K_M</u> for ATP (μM)		<u>V_{max}</u> (unit/mg)	
<u>Enzyme</u>	(AMP=0.2 mM)	(AMP=10 mM)	(AMP = 0.2 mM)	(AMP = 10 mM)
wild type	90 ± 14	52 ± 10	900 ± 45	400 ± 18
phe86→trp	287 ± 32	292 ± 50	250 ± 24	1430 ± 69
tyr133→trp	153 ± 19	-	1120 ± 75	-
leu107→gln	542 ± 160	-	125 ± 18	-

<u>Enzyme</u>	<u>K_M for AMP (μM) *</u>	<u>V_{max} (unit/mg) *</u>	<u>K_I for AMP μM *</u>
wild type	48 ± 10	1316 ± 158	1500 ± 160
phe86→trp	2100 ± 220	1461 ± 36	None
tyr133→trp	43 ± 8	1420 ± 150	1900 ± 204
leu107→gln	38 ± 8	185 ± 16	1700 ± 600

*For these measurements the ATP concentration was held constant at 5 mM.

The data indicate that phe86 is involved in the AMP binding site rather than the ATP binding site since the phe86K_M for AMP. The rather small increase in the K_M for ATP suggests that phe86 is not directly involved in the binding of ATP. Instead the change in the K_M for ATP could be the result of an interaction between the two nucleotide binding sites since the binding of one substrate has been shown to affect the binding of the other.

The introduction of a single tryptophan residue in place of phe86 in the binding site created an intrinsic fluorescent probe to monitor the binding of nucleotides. The emission maximum of the fluorescence from trp86 was at 328 nm in comparison to 338 nm for trp133 which is consistent with the location of trp86 in the

hydrophobic cleft of the nucleotide binding pocket. The binding of nucleotide substrates and their analogues all caused significant changes in the fluorescence quantum yield and showed saturation kinetics which allowed their dissociation constants to be readily determined. The phe86→trp mutant adenylate kinase showed much higher affinity for ATP than AMP which is consistent with their K_M 's (6).

The tyr133→trp adenylate kinase had nearly normal kinetic constants and showed no fluorescence changes upon nucleotide binding indicating that this residue was not located in the region of a nucleotide binding site contrary to the conclusions of previous X-ray studies. This and other results suggest the tyr133 likely plays a structural role rather than being part of a nucleotide binding site on the enzyme.

The leu107→gln mutation caused a significant change in the structure of the enzyme as evidenced by the decrease in V_{max} and the temperature-sensitivity. Even though this weakens any argument about the role of this residue in the binding of a specific nucleotide it is pertinent that the mutation caused a 6-fold change in the K_M for ATP without any change in the K_M for AMP. In conjunction with other data, it suggests that leu107 is located nearer the ATP binding site than the AMP binding site. The side-chain of leu107 may exert its influence directly on ATP binding or by changing the packing of this key helix in relation to the central β -sheet. Further studies will be necessary to clarify the precise role of this residue in ATP binding.

Based on the effects of the amino acid substitutions on the kinetic and fluorescence properties of AMP and ATP binding, a proposed model for the location of the nucleotide binding sites is shown in Figure V. The model shows the AMP site near phe86 and the ATP site near leu107. Characterization of the properties of these mutant enzymes has also led to a proposed mechanism for AMP substrate inhibition (6).

FIGURE V. A molecular model for the location of the ATP and AMP binding sites on *E. coli* adenylate kinase. It is proposed that the AMP site is near phe86 (right hand side), and the ATP site is near leu107 (lower left). Tyr133 (upper left) appears not to be directly involved in nucleotide binding. The amino acids where substitutions were examined are shown with yellow dotted surfaces corresponding to the van der Waals radii of each atom, and stick drawings represent the nucleotides.

Bibliography

1. Althoff, S., Zambrowicz, B., Liang, P., Glaser, M., and Phillips, G.N., Jr. (1988) J. Mol. Biol. **199**, 665-66.
2. Chatteraj, D.K., Cordes, K., Berman, M.L., and Das, A. (1984) Gene **27**, 213-222.
3. Cohen, S.N., and Chang, A.C.Y. (1977) J. Bacteriol. **132**, 734-737.
4. Ditta, G., Schmidhauser, T., Yakobson, E., Lu, P., Liang, X-W., Finlay, D.R., Guiney, D., and Helinski, D.R. (1985) Plasmid **13**, 149-153.
5. Horwitz, A.H., Heffernan, L., Cass, L., Miyada, C.G., and Wilcox, G. (1980) Molec. Gen. Genet. **179**, 615-625.
6. Liang, P., Phillips, G. N. Jr., and Glaser, M. (1991) Proteins **9**, 22-36.
7. Liang, P., and Glaser, M. (1989) Gene **80**, 20-28.
8. Maniatis, T., Fritsch, E. E., and Sambrook, J. (1982) Molecular Cloning. A Laboratory Manual, Cold Spring Harbor Press, NY.
9. Meacock, P.A., and Cohen, S.N. (1980) Cell **20**, 529-542.
10. Pai, E.F., Sachsenheimer, W., Schirmer, R.H., and Schulz, G.E. (1977) J. Mol. Biol. **114**, 37-45.
11. Stauffer, G.V., Zurawski, G., and Bennett, G.N. (1979) Plasmid **2**, 498-502.

Publications attributable to DOE funding:

- Althoff, S., Zambrowicz, B., Liang, P., Glaser, M., and Phillips, G. N., Jr. (1988) "Crystallization and Preliminary X-ray Analysis of Escherichia coli Adenylate Kinase," J. Mol. Biol. **199**, 665-666.
- Liang, P., and Glaser, M. (1989) "Efficient Cloning of a Mutant Adenylate-Kinase-Encoding Gene from Escherichia coli," Gene **80**, 21-28.
- Liang, P., Phillips, G. N. Jr., and Glaser, M. (1991) "Assignment of the Nucleotide Binding Sites and the Mechanism of Substrate Inhibition of Escherichia coli Adenylate Kinase," Proteins **9**, 22-36.

END

**DATE
FILMED
5/05/92**