Final Technical Report for February 15, 1990 – February 14, 1992

DOE/ER/40557--2

DE92 008138

1 Introduction

During the first two years of this program, we have successfully investigated two new methods for the production of polarized muonic helium. This work was done at LAMPF in collaboration with a group from Syracuse University. Traditionally, polarized muonic helium has been formed by stopping polarized muons in unpolarized atoms. Unfortunately, because of depolarization in the muon cascade to the ground state, residual polarizations are only $\sim 3\%$. [1, 2, 3] The two methods we have developed both achieve much higher muon polarizations. To accomplish our goals, we first developed an appropriate muon beam for use with our small gas targets. This work is summarized by Holmes *et al* in a NIM article. [4] During the summer of 1990, we stopped unpolarized negative muons in nuclear polarized ^3He . The muons were polarized in the cascade to the ground state through the hyperfine interaction with the nucleus. The resulting muon polarizations were $7.2 \pm 0.8\%$ for a 100% nuclear polarized target. While higher polarizations are clearly desirable for practical purposes, this experiment yielded important insights on spin interactions that occur in muonic atoms. Also, the ability to rapidly reverse the target polarization gave the experiment good sensitivity with minimal systematic effects. This work is summarized in a Physical Review Letter, a preprint of which is attached to this proposal as appendix A. [5] Last summer, we completed an experiment in which we polarized muonic helium by direct spin interactions with an optically pumped Rb vapor. In this technique, the muonic helium atom is polarized through a combination of charge exchange and spin exchange with the polarized valence electron of the Rb vapor. In contrast to the technique described in appendix A, the ^3He nuclei are not polarized. This last technique yielded dramatic polarizations approaching 50%.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

2 Laser polarized muonic ^3He : stopping muons in polarized ^3He

We summarize here the results of the first experiment carried out under this grant. This experiment (LAMPF experiment #1164) is described in more detail in a Physical Review Letter that is attached to this proposal as appendix A.

When an unpolarized negative muon is stopped in a nuclear polarized ^3He target, the muon becomes polarized by the hyperfine interaction in the process of cascading to the $1s$ ground state. As was discussed by Kuno *et al.*, the final muon polarization depends strongly on the n level at which the hyperfine splitting becomes comparable to the level width.[6] Collisional effects can also limit the final polarization. The "repolarization" of negative muons stopped in polarized nuclear targets was first demonstrated by Kadono *et al.* using ^{209}Bi in which the nuclear spins were oriented by placing a ferromagnetic compound BiMn in a magnetic field of 5-8 kG at temperatures of around 60 mK.[7]

The ^3He targets for experiment 1164 were polarized by spin-exchange with optically pumped Rb vapor.[8] The targets were 1" glass spheres containing 7-8 atm. of ^3He gas, 75 Torr N_2 gas, and a few mg of Rb. The glass cell walls were only $\sim 100\mu\text{m}$ thick in order to maximize the number of muons that penetrated into the cell and stopped in the ^3He .

To polarize the targets, the cells were heated to obtain Rb number densities of $\sim 10^{15} \text{ cm}^{-3}$. The Rb vapor was optically pumped with 3-5 Watts of laser light. The polarized Rb valence electron spin-exchanges with the helium nuclei, leading to ^3He nuclear polarizations. In order to achieve high polarization, the ^3He spin relaxation time must be much longer compared to the spin-exchange time. To this end we developed a technique for cleaning the ^3He gas that resulted in 7-8 atmosphere targets cells with ^3He spin relaxation times on the order of 100 hours. These lifetimes were dominated by spin relaxation during ^3He - ^3He collisions, and compare quite favorably with previously reported lifetimes for high pressure cells.[9]

The nuclear ^3He polarization was monitored by the new *frequency shift* method.[10] The polarized ^3He nuclei cause an effective magnetic field that is experienced by the Rb atoms in the cell. A voltage controlled oscillator is locked to the Rb magnetic resonance frequency, and a frequency shift is observed that is proportional to the ^3He polarization. The proportionality constant, which is a fundamental constant for the Rb- ^3He system, has been determined in a separate set of measurements.

The frequency shift technique has several advantages for target polarimetry. Since the quantity measured is a frequency, it can be measured quite accurately. In fact, the absolute accuracy of the measurement is only limited by the knowledge of the aforementioned proportionality constant, which can be improved at any later time. Also, the target cell can be probed optically, which greatly simplifies the required instrumentation in the beam. Both the improvements in the wall relaxation times and this new form of polarimetry should be important in future experiments employing these targets.

The geometry of the experiment was quite simple. A pair of thin (40μ thick) plastic scintillators positioned just in front of the target detected the incoming muons. Two telescopes, located above and below the target, detected the decay electrons from the stopped muons. Each electron telescope contained a pair of scintillators. Beam line extensions were used to reduce the air through which the muons traveled to less than 2 cm. Gated scalars counted the number of electrons for each telescope that were detected within $4.4 \mu\text{sec}$ of a muon's arrival.

The polarization of the captured muons was determined from the up-down asymmetry in the gated decay electrons,

$$A_{\text{exp}} = \frac{N_{\uparrow\uparrow} - N_{\uparrow\downarrow}}{N_{\uparrow\uparrow} + N_{\uparrow\downarrow}}, \quad (1)$$

where $N_{\uparrow\uparrow}$ ($N_{\uparrow\downarrow}$) is the number of electrons that decayed parallel (antiparallel) to the magnetic holding field, which was adiabatically rotated every two minutes. It was also possible to choose arbitrarily the direction of the target spins with respect to the magnetic field. The expected sign of A_{exp} is then negative (positive) if the target spins are parallel (antiparallel) to the magnetic field. The experimental asymmetries, adjusted to correspond to the average target polarization of 36%, are displayed in Fig. (2). The experimental asymmetry of approximately 0.3% was obtained in about 100 hours of beam time with an error of less than 5%. The cleanliness of this data attests to the power of the rapid polarization reversals possible with laser techniques. Our final result is $P_{\mu}^N = 0.072 \pm 0.008$ where P_{μ}^N is the final muon polarization normalized to a 100% polarized ^3He target. Our measurement leads to better understanding of spin interactions muonic atoms, and provides important practical information on the use of lasers for producing polarized muonic ^3He . It should be possible to apply these same techniques to the formation of polarized muonic ^{21}Ne and ^{129}Xe , where even higher muon polarizations would be expected. Highly polarized muonic ^{21}Ne is of interest for possible time-reversal experiments.

Figure 1: Raw asymmetries adjusted to a constant helium polarization $P_{He} = 36\%$, and fraction of decay electrons from helium $f = 39\%$. As indicated by the solid line, the sign of the asymmetry conforms to that expected from the relative orientation of the target spin to the applied magnetic field.

3 Polarizing Muons by Direct Interactions with Polarized Rb

3.1 Theory

We describe here the second of the two techniques we have demonstrated for polarizing muonic helium using lasers. This work was performed during the summer of 1991 and represents the first phase of approved LAMPF experiment #1231. With this technique we polarized muonic helium through direct interactions with optically pumped Rb vapor. In contrast to the experiments from the summer of 1990, the helium nuclei were unpolarized and the optical pumping was done with the target in the beam. The results were fairly spectacular. We observed time averaged polarizations in 3He and 4He of approximately 22% and 37% respectively. The polarization increased with time in a roughly exponential fashion, with time constants of 1.5–3.0 μs . We observed saturation polarizations reaching 39% and 58%. This is an order of magnitude increase over any previously achieved polarizations in muonic helium, and easily in a range that is useful for measurements of g_P .

When a muon stops in the gas, a muonic helium ion $(\mu^- \text{He})^+$ is formed. Before the ion can interact with the optically pumped Rb vapor, it must first be neutralized. Neutral muonic helium reacts chemically like hydrogen, and the spin interaction between Rb and hydrogen are then mimicked in the Rb-muonic helium system. To this end, CH_4 was added to the target cells. Methane, with an ionization potential of 12.6 eV, will neutralize the muonic helium ion in a simple charge exchange reaction of the form[11]:

The $(\mu^- \text{He})^+$ core will act as a kind of pseudo-nucleus with a charge of +1. The neutral muonic helium atom will then become polarized through spin-exchange collisions with the Rb vapor:

where arrows indicate valence electron polarization. This reaction is chemically identical to Rb-hydrogen spin-exchange which has been extensively studied.[12] Following a spin-exchange collision, the electron spin polarization of the neutral muonic helium will be transferred to the muonic core via the hyperfine interaction. The only caveat is that when the μ^- spin couples to the ^3He nuclear spin, spin-one triplet states will be formed 75% of the time, and spin-zero singlet states will be formed 25% of the time. The spin exchange collisions described by (3) will only be effective for polarizing the the spin-one triplet states, so the maximum possible polarization will be 75%.

The neutral muonic ^3He polarization will evolve with time according to the relation

$$P_{He} = \langle P_{Rb} \rangle \frac{3}{4} (1 - e^{-f_N / \tau_{ex}}) \quad (4)$$

where P_{He} is the polarization of the muonic core of the atom, $\langle P_{Rb} \rangle$ is the polarization of the Rb vapor averaged over the volume in which the muons stop, $f_N \approx 3/11$ is the nuclear slowing down factor which results from the large spin-capacity of the nucleus, and $1/\tau_{ex}$ is a spin-exchange rate given by

$$\frac{1}{\tau_{ex}} = \langle \sigma v \rangle [\text{Rb}] . \quad (5)$$

Here $\langle \sigma v \rangle$ is the velocity averaged cross section for spin exchange collisions, and $[\text{Rb}]$ is the number density of Rb atoms. Clearly, P_{He} will only reach high values if the effective spin-exchange rate f_N / τ_{ex} is at least comparable to the muon decay rate of $\approx 4.5 \times 10^5 / \text{sec}$. Otherwise, the muonic helium will not become polarized before it decays.

The muon polarization can be greatly enhanced by removing the CH_4 , in which case the $(\mu^- \text{He})^+$ ion will form a molecular ion with the surrounding He atoms [13]. The $(\mu^- \text{He})\text{He}^+$ molecules are broken up by the polarized Rb vapor in a charge induced disassociative reaction of the form [14]:

where arrows indicate polarized valence electrons. The net result is that the $(\mu^- \text{He})^+$ ion picks up a polarized electron from the Rb atom. The cross-section for this process is on the same order as the spin-exchange process. The neutral muonic helium atom will then undergo spin-exchange collisions with the polarized Rb vapor as described above. The effective time constant with which the muonic atoms become polarized is shortened. As we will discuss below, our best performance resulted from leaving the CH_4 out of the cells, and allowing the $(\mu - \text{He})^+$ ions to become polarized from the Rb.

3.2 Experimental set-up and results

The experimental apparatus is illustrated in Fig. 3.2. As with our earlier work, the target cells were about 2.5–3.0 cm in diameter, and contained about 8 atm. of helium, as well as smaller amounts of N_2 , several milligrams of Rb, and in some cases a few percent of CH_4 . Through a combination of strong focusing and collimation, we achieved a beam spot with an area of $\sim 1 \text{ cm}^2$. The beam tune, which also featured a fairly narrow momentum spread, is described in detail in reference [4]. The cell was illuminated with a Spectra Physics Ti:Sapphire laser pumped by a continuous wave Ar^+ laser, which provided about 5 watts of cw light at the Rb D_1 resonance of 795 nm. The Rb polarization, and consequently the muon polarization, was reversed by rotating the $1/4$ wave plate that was used to circularly polarize the laser light. The incoming muons were tagged by the thin scintillators at the exit to the beam pipe, and their decay electrons were detected by a pair of telescopes. Each telescope comprised two scintillators which were positioned above and below the target.

The muon polarization was determined by the up down asymmetry in the gated decay electrons,

$$A_{\text{exp}} = \frac{N_{\uparrow\uparrow} - N_{\uparrow\downarrow}}{N_{\uparrow\uparrow} + N_{\uparrow\downarrow}} \quad (7)$$

where $N_{\uparrow\uparrow}(N_{\uparrow\downarrow})$ is the number of electrons that decayed parallel(antiparallel) to the Rb polarization direction as determined by the angular position of the $1/4$ wave plate

Figure 2: Shown schematically is the apparatus used to polarize muonic helium through direct interactions with polarized Rb vapor.

which fixed the circular polarization of the laser light. Our experimental up/down asymmetry was diluted by two factors — the analyzing power of the decay is about $1/3$, and only about 28% of the decay electrons detected were from muons stopped in helium. As in our earlier work, the fraction of muons stopped in the gas was determined by analyzing the time spectra of the muon decays.

The asymmetries were large enough that we were able to study the time dependence of the build up of polarization. We considered two variables in the target cells — the presence or lack of CH_4 , and the isotope of He, either ^3He or ^4He . We thus studied four different target cells, representing the four possible combinations. Our results are shown in Fig. 3.2.

There are three trends that we see in our data. Since only 75% of the $(\mu^-^3\text{He})^+$ ions are in triplet states, we expect the cell containing ^3He will saturate at lower polarizations than cells containing ^4He . In fact, assuming that $\langle P_{\text{Rb}} \rangle$ is the same for both cells, we would expect a ratio of saturation polarizations of 0.75. The ratio observed for the cells shown in (a) and (c) is 0.72 ± 0.07 . Moreover, the nuclear slowing down factor f_N increases with increasing nuclear spin so we expect cells containing ^3He to polarize more slowly than cells containing ^4He . This is because the “nucleus”

Figure 3: The time dependence of the asymmetries are shown for each of the four target cells studied. In the top row, we show two cells containing ^4He : (a) contains no CH_4 , and (b) contains 3% CH_4 . In the bottom row, we show two cells containing ^3He : (c) contains no CH_4 , and (d) contains 5% CH_4 . The measured asymmetries have been normalized to the fraction of stops in the He gas to make it possible to compare data from different runs.

of muonic ^3He will have a spin of 1 (for the triplet states) whereas the “nucleus” of muonic ^4He will have a spin of $1/2$. Finally, as discussed above, if the cross section for the breakup of HeH^+ under collisions with Rb is large enough, we expect the polarization to increase more quickly in the cells which do not contain methane.

All of these trends are evident in our data, convincing us that we understand most of the gross aspects of the spin polarization processes that we are observing. Most importantly for future work, our observed polarizations are extremely high. We note also an additional advantage of this technique which is very important for our measurement of g_p . The lifetime of a polarized Rb atom is on the order of a millisecond. Thus, any impurities in the target that might potentially depolarize the Rb or the muonic helium ion would have to interact on a very short timescale. We can therefore relax our requirements that pristine conditions be maintained in the cell. For instance, the cell can contain metal electrodes, so that it can be used as a drift chamber. This is very different from the situation described in section 2 in which long (~ 100 hours) spin relaxation times were crucial to producing high ^3He

nuclear polarizations.

3.3 SLAC Target cell development

We have made progress on development of the polarized ^3He target cell for SLAC experiment E142. The spin relaxation time of a target cell is extremely important to achieving high polarizations. In the development work for LAMPF, all of our cells were spherical, and had volumes of 7-8 cm³. In contrast, the SLAC target cells have volumes more than an order of magnitude larger, and are cylindrical in shape. The first SLAC target cells that were built had lifetimes of only minutes. These results suggested that collisions with the walls were causing strong relaxation. As discussed earlier, a good cell has a spin relaxation time that is dominated by ^3He - ^3He collisions, which should be about 80 hours for a 10 atm. cell. We suspect that the raw tubing has surface irregularities or deposits of paramagnetic impurities that increase wall relaxation. We were less sensitive to these problems when using spherical cells which were blown out from a molten blob of glass. Our hypothesis is that small flecks of steel are embedded on the inner surface of the tubing. During the melting and blowing of a sphere, these flecks are homogeneously distributed throughout the glass, resulting in a nearly pristine, fire polished surface. When raw tubing is used for making a cylindrical cell, however, these flecks of metal can cause serious problems.

Acting on our suspicions, we have explored several methods of preparing the surface of the glass tubing from which we make our cells. In order to distinguish wall relaxation rates from ^3He - ^3He bulk relaxation, we have conducted a series of cell tests at low ^3He pressures. We have had excellent results with several different acid etching techniques. Etching with nitric acid alone yields 30-130 hour wall relaxation times, a dramatic increase from the several minute relaxation times first observed. A combined etch of nitric and hydrochloric acids produced greater than 200 hour wall relaxation times, essentially eliminating wall relaxation as an obstacle to long lifetime cells. We believe that the two different acids are needed to attack all the metals commonly found in the steel alloys that may exist as residues from the manufacturing process. We have now produced half-sized target cells containing 7-8 atms. of ^3He with better than 40 hour relaxation times. The 40 hour relaxation times already meet the E142 proposal specifications. The remaining discrepancy between the 40 hour lifetime and the expected 80 hour lifetime is probably due to impurities in the ^3He gas. The larger cell volume requires much more gas, making cleaning difficult. We are in the process of installing a liquid He trap in our gas handling system to cryogenically remove any

non-helium impurities from the ^3He . We will soon combine the acid etching technique with our improved gas cleaning ability to make the first full sized target cells.

References

- [1] P. A. Souder et al., Phys. Rev. Lett. **34**, 1417 (1975).
- [2] P. A. Souder et al., Phys. Rev. A **22**, 33 (1980).
- [3] H. Orth, Hyperfine Interactions **19**, 829 (1984); K.-P. Arnold et al., Verhandl. DPG **4**, 315 (1982).
- [4] R. Holmes, D.-H. Kim, K.S. Kumar, P.A. Souder, D. Benton, G.D. Cates, N. Newbury, and C. Pillai, Nucl. Inst. and Meth. in Phys. Res. A **303**, 226 (1991).
- [5] N.R. Newbury, A.S. Barton, P. Bogorad, G.D. Cates, M. Gatzke, B. Saam, L. Han, R. Holmes, P.A. Souder, J. Xu, and D. Benton, Phys. Rev. Lett., **67**, 3219 (1991).
- [6] Y. Kuno, K. Nagamine, and T. Yamazaki, Nuc. Phys. A **475**, 615 (1987); J. Congleton, Ph.D. Thesis, TRIUMF, (unpublished, 1992).
- [7] R. Kadono et al., Phys. Rev. Lett. **57**, 1847 (1986).
- [8] M. A. Bouchiat, T. R. Carver, C. M. Varnum, Phys. Rev. Lett. **5**, 373 (1960); W. Happer et al., Phys. Rev. A **29**, 3092 (1984); X. Zeng et al., Phys. Rev. A **31**, 260 (1985).
- [9] B. Larson, O. Häusser, P.P.J. Delheij, D.M. Whittal, and D. Thiessen, Phys. Rev. A **44**, 3108 (1991).
- [10] S. R. Schaefer et al., Phys. Rev. A **39**, 5613 (1989).
- [11] H. von Koch, Arkiv För Fysik **28**, 529 (1965); Y. Chiu et al., J. Chem. Phys. **88**, 6814 (1988); Y. Nakai, T. Shirai, T. Tabata and R. Ito, Atomic and Nuclear Data Tables **37**, 69-101 (1987).
- [12] S.G. Redsun, R.J. Knize, G.D. Cates, and W. Happer, Phys. Rev. A **42**, 1293 (1990).
- [13] G. Reifenröther, E. Klempt and R. Landua, Phys. Lett. B **191**, 15 (1987).

- [14] W.J. van der Zande, W. Koot, D.P. de Bruijn and C. Kubach, Phys. Rev. Lett. 57, 1219 (1986).

(removed & copied separately)

END

END

END

**DATE
FILMED**

3 / 24 / 92