DOE/ER/60340--8

DE92 007907

California Basin Study (CaBS)
DOE West Coast Basin Program

Grant DE-FG05-85ER60340

Progress Report VIII

For work performed during 15 Nov 1989 - 14 Nov 1990

by

Lawrence F. Small
College of Oceanography
Oregon State University
Corvallis, Oregon 97331-5503

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. INTRODUCTION

The overall objective of our research continues to be elucidation of the transport pathways and transformations of organic matter in the California Basins region, with particular reference to the role of macrozooplankton in upper waters. We have concentrated on C and N pathways and fluxes to date, and will continue to investigate these further (seasonal aspects, and the role of zooplankton carnivory in zooplankton-mediated C and N flux, for example).

During this report period we have continued to analyze data and write manuscripts for publication, based on the eight cruises in Santa Monica Basin in which we have participated prior to the present grant year (CaBS-1, -2, -3, -4, -5, -7, -8, -10, in October 1985, February 1986, May 1986, October 1986, April 1987, October 1987, February 1988, and September/October 1988, respectively). CaBS-6 and CaBS-9 were hydrographic cruises only, on a smaller vessel (CaBS-2 was mainly a hydrographic cruise as well, but we were able to obtain some data in this February 1986 period).

During this report period we also have participated in cruise CaBS-11, in January/-February 1990, and as this progress report is being written we are preparing to participate in CaBS-12, the final cruise of this series, in July 1990. CaBS-12 is very important to us from the standpoint of seasonal coverage, as it is the only mid-summer cruise of the series. Samples from CaBS-11 are currently being worked up in the laboratory.

The CaBS group has agreed to develop summary papers for a compendium volume of Progress in Oceanography, and L.F. Small is the guest editor in charge of assembling and initially editing the manuscript for Progress in Oceanography. We hope to have all manuscripts in press in 1991. Our own contribution will be a summary of the role of macrozooplankton in transforming the small particle field, and in contributing to the vertical flux of particulate C and N, in Santa Monica Basin.

II. REPORT OF PROGRESS

A. Fecal Pellet Production Rates

From the eight cruises in which we have fully analyzed data sets, we have developed night and day biomass profiles, gut evacuation rates, gut pigment concentrations, and fecal pellet C/Pigment ratios for two feeding modes (strict herbivory and herbivory/detritivory), all for three size classes of zooplankton in Santa Monica Basin. These herbivory and herbivory/detritivory data have been turned into night and day (and 24-hr) estimates of fecal carbon production rates through the photic zone by the three size groups, and these rates in turn have been compared to primary production and new production in the photic zone, and to trap-

measured total carbon flux out of the photic zone [Table 1, showing results for a spring (May), fall (October) and winter (February) month, as examples]. Analyses for three October cruises and two spring cruises (April and May) have recently been reported (Small et al., 1989 - see Appendix). We have another paper (Small and Ellis, in prep.), in which we examine the potential role of carnivorous feeding to carbon flux in Santa Monica Basin. The effect of carnivory is not shown in Table 1, but our initial analyses indicate that carnivory is very important in the region, and unless this contribution is included, the role of macro-zooplankton in transforming particulate C into sinking flux will be greatly underestimated (see later). On the other hand, all of the herbivory/detritivory contribution appears to be herbivory, within the errors of our analytical and computational methods.

Our analytical methods for assessing fecal carbon production rates are briefly as follows. First we determine the initial gut pigment levels in fresh collections of the three size groups of zooplankton, night and day, and above and below the top of the nitracline. Gut pigment is chlorophyll plus pheophytin, determined by fluorometry and converted to mass concentration units (gut pigment per unit dry weight of the animals). Conversion to gut pigment per unit body C is accomplished by knowing the percentage body C, measured in CHN analyzer. Next we determine the presumed C/Pig ratio in fecal pellets 1) as if the animals were eating non-selectively from the total suspended particle field, and 2) as if they were eating only the phytoplankton fraction from the total suspended particle field. This is done as follows. The POC/Total Pigment ratio in the suspended particle field is computed for the regions above and below the top of the nitracline (within the photic zone), as representative of the food source for herbivorous/detritivorous feeders above and below the top of the nitracline. This ratio is then multiplied by $0.30/0.67 = 0.45$, where 0.30 is the fraction of ingested POC that is egested (i.e., a 70% assimilation efficiency), and 0.67 is the average amount of total ingested particulate pigment not lost during gut passage (from compendium data from many authors). The resulting ratio is that presumed to be the fecal pellet C/Pig ratio under herbivory/detritivory. For an estimate of strictly herbivorous feeding, the POC/Total Pigment ratio of the suspended particle field is replaced by the ratio $(\text{PON} \times 5.7)/\text{Chl}$, where $(\text{PON} \times 5.7)$ is the Redfield-derived estimate of phytoplanktonic carbon (Redfield $\text{POC}/\text{PON} = 5.7$ by weight), and Chl is the chlorophyll concentration (not total pigment). This $(\text{PON} \times 5.7)/\text{Chl}$ ratio is also multiplied by 0.45 (see above), to yield an estimate of the fecal pellet C/Pig ratio under strict herbivory. As indicated earlier for the CaBS region, strict herbivory was predominant (under our definition), because the POC/PON ratios in the suspended particle fields rarely exceeded 5.7; that is, the suspended particle field was judged to be all living phytoplankton on most occasions (surely this was not perfectly true, but true within the limits of the Redfield POC/PON ratio to elucidate phytoplankton from detrital particles).

Following establishment of both gut pigment concentrations in animals and presumed C/Pig ratios in fecal pellets, we ran gut evacuation rate experiments on the three animal size groups, night and day. Briefly, this entailed following the decline of total gut pigment through time, to establish rates of gut clearance. Hourly gut evacuation rates were converted to rates expected through the total night period (and through the total daylight period) of the

Table 1. Primary production, new production, carbon flux, and fecal pellet production above and below the top of the nitracline and through the complete photic zone during three selected months (cruises) in Santa Monica Basin, California. Fecal pellet production assumes herbivorous/-detritivorous feeding only. Percentage comparisons are also given. Thanks to R. Eppley and M. Landry for certain data.

		May 1986			October 1987			February 1988		
		0-16m	16-39m	0-39m	0-50m	50-72m	0-72m	0-39	39-55	0-55
1	Prim. Prod. (mgC m ⁻² d ⁻¹)	634	620	1254	532	102	634			425*
2	New Prod. (mgC m ⁻² d ⁻¹)	39	149	188	21	83	104			85*
3	Trap Flux (mgC m ⁻² d ⁻¹)	-	-	259	-	-	241			387
4	Col.2/Col.1 x 100 (%)	6.2	24.0	15.0	4.0	81.4	16.4			20.0
5	Col.3/Col.1 x 100 (%)	-	-	20.7	-	-	38.0			91.0
6	Col.2/Col.3 x 100 (%)	-	-	72.6	-	-	43.2			22.0
Pellet Production (mgC m ⁻² time period ⁻¹)										
7	200-500 μm, night	8.2	20.6	28.8	20.4	3.9	24.3	12.6	3.1	15.7
8	200-500 μm, day	15.4	12.9	28.3	7.9	1.7	9.6	5.8	1.9	7.7
9	500-1000 μm, night	6.5	8.2	14.7	2.2	0.4	2.6	4.5	0.9	5.4
10	500-1000 μm, day	5.8	6.1	11.9	0.4	0.1	0.5	1.4	0.3	1.7
11	1000-2000 μm, night	2.4	3.9	6.3	0.7	0.2	0.9	2.9	0.5	3.4
12	1000-2000 μm, day	1.2	1.7	2.9	0.1	0.1	0.2	0.7	0.2	0.9
Pellet Production [mgC m ⁻² (24 hr) ⁻¹]										
13	200-500 μm	39.5	53.4	92.9	31.7	6.4	38.1	27.9	6.9	34.8
14	500-2000 μm	15.8	20.0	35.8	3.4	0.8	4.2	9.5	1.9	11.3
15	Col.13/Col.1 x 100 (%)	6.2	8.6	7.4	6.0	6.2	6.0	-	-	8.2
16	Col.14/Col.1 x 100 (%)	2.5	3.2	2.9	0.6	0.8	0.7	-	-	2.7
17	Col.13/Col.2 x 100 (%)	101.1	35.9	49.4	151.1	7.7	36.6	-	-	40.9
18	Col.14/Col.2 x 100 (%)	39.5	13.4	19.0	16.3	1.0	4.1	-	-	13.3
19	Col.13/Col.3 x 100 (%)	-	-	35.9	-	-	15.8	-	-	9.0
20	Col.14/Col.3 x 100 (%)	-	-	13.8	-	-	1.8	-	-	2.9

* = estimated from previous Jan/Feb cruise data with similar hydrographic/nutrient/light patterns.

24-hr day simply by multiplying by the number of nighttime (and daylight) hours present during the month of sampling.

The resultant fecal pellet production rates for each size group, day and night, above and below the top of the nitracline (as in Table 1), were computed from the following formula:

$$F = P \times R \times E$$

where: F = fecal carbon production rate (either herbivorous or herbivorous/detritivorous, as mg fecal C/mg body C/hr; P = gut pigment level, as mg Pig/mg body C; R = fecal C/fecal Pig (either the herbivorous or herbivorous/detritivorous ratio); and E = number of gut pigment evacuations/night or day period, from the gut pigment evacuation rate experiments. To put F on a population basis above and below the top of the nitracline, we multiplied the appropriate F by the integrated concentration of each zooplankton size group above and below the top of the nitracline (concentrations in mg body C/m², from the biomass profiles). Thus, population F was in units of mg fecal C/m²/day or night period. Evaluating animal concentrations for day and night periods, above and below the top of the nitracline, allowed us to take into account the effects of diel vertical migration and any other shifts in population abundance within the photic zone.

Any combination of size-group, day-night, or above-within nitracline pellet production rates can be examined by developing the data as above. For example, to obtain the overall fecal C production rate by all three size classes combined over a 24-hr period through the complete photic zone, one simply sums the fecal C production rates for all three size groups during both day and night periods above and within the nitracline (Table 1). In Table 1 we also examined the combined role of the 500-1000 µm and 1000-2000 µm groups, separate from the role of the smallest size class (200-500 µm), figuring that fecal pellets from the two larger size classes would be those most likely to make it intact into sediment traps just below the photic zone, and therefore those most likely to be visible and "countable" under a microscope. It should be noted that such rates and rate combinations are directly comparable to trap-measured C fluxes out of the photic zone, as well as to primary and new production rates (Table 1).

Finally, it should be noted here that the precision with which F can be calculated depends upon the propagation of errors associated with the calculations of P, R and E. The approximate standard error around any calculated mean F (s_F) can thus be determined as follows, ignoring interaction terms:

$$s_F = \left[F^2 \left(\frac{s_P^2}{P^2} + \frac{s_R^2}{R^2} + \frac{s_E^2}{E^2} \right) \right]^{1/2}$$

where s_P^2 , s_R^2 , and s_E^2 are respectively the mean squares of the means of P, R and E. The propagated error term can be calculated for each F (i.e., for each size group, day and night, above and within the nitracline, on each cruise date). We have done this for our cruise data analyzed to date, and have calculated that the propagated error (s_F) averaged approximately 19% of mean F values (range of 18-24%). The propagated error associated with each population F (i.e., F times appropriate biomass B) can be calculated by including the term s_B^2/B^2 in the above equation for s_F . Our experience to date with replicated net hauls suggests that the mean standard error for biomass concentrations of a given size group taken within minutes of each other is less than 10%.

In addition to transforming the suspended-particle field into fecal matter of different chemical composition, the macrozooplankton also eat microzooplankton, nauplii, and other animal matter. We have begun to directly measure C/Pig ratios in fresh fecal pellets, and have found these ratios to be substantially higher than those ratios computed for herbivory and herbivory/detritivory. Thus, there must be a substantial amount of carnivory, i.e., the macrozooplankton (particularly the largest of our three size groups) must have been obtaining their daily rations by eating a lot of carbon-rich animal food containing little or no chlorophyllous pigment. This meant 1) that zooplankton were potentially contributing even more carbon to total flux than heretofore recognized in Table 1, for example, and 2) that total fecal pellet production could no longer be directly compared to primary and new production (because the source carbon for pellets was now from both animal and plant material, not just plants). We are in the process of determining the effect of carnivory on total zooplankton fecal C production, and early calculations suggest that the effect will be to at least double the potential fecal C contribution to total C flux at the base of the photic zone in Santa Monica Basin. It is readily seen that such information could have large impact on our ability to infer carbon flux from satellite measurements of surface pigments only.

B. Fecal Contributions to Total Particulate Carbon Flux

Using all available data for autumn (September/October) and winter (January/February) in Santa Monica Basin, we have developed mean C/N and C/Pigment ratios for the suspended particle fields, zooplankton bodies (200-2000 μm), zooplankton fecal pellets (from 200-2000 μm animals), and sediment trap material at the base of the photic zones during those two seasons (Fig. 1), to show that macrozooplankton represent an important pathway through which particulate C and N are transformed into fecal particles having different ratios than the suspended particles but similar ratios to sediment trap material. This suggests that zooplankton graze the small particle field and also consume tiny animal matter (high in carbon but devoid of pigment), subsequently use appropriate ratios of C to N for their own body composition (body C/N = 4.0 in Fig. 1), and then void some of the rest as high C/N fecal pellets which become a large part of the trap material. Some C and N are "lost" as dissolved respiratory and excretory products as well.

Work on $\delta^{15}\text{N}$ was begun over a year ago with Dr. Mark Altabet of WHOI, and two autumn CaBS sample suites have been analyzed (Fig. 1, and see Appendix). Work continues

on CaBS-11 samples (Jan/Feb 1990) and will proceed on our CaBS-12 (July 1990) cruise. Note that the mean $\delta^{15}\text{N}$ value in animal bodies is substantially higher than the mean value in the suspended particles, and that the mean fecal pellet $\delta^{15}\text{N}$ value is intermediate. This suggests again that the animals process N from different sources, the two logical sources being the small particle field with low $\delta^{15}\text{N}$ and small animal matter with relatively high $\delta^{15}\text{N}$, to yield fecal matter with intermediate $\delta^{15}\text{N}$ values. Data are few at present, however, and these suppositions need verification in upcoming work. A major significance of $\delta^{15}\text{N}$, from other non-CaBS work we have done, does seem to be that it can help distinguish food sources of N for zooplankton (animal or plant) even when C/N ratios are indistinguishable (in certain open-ocean environments C/N ratios of suspended particles, animals, fecal material, and trap contents aren't all that distinguishable).

III. PUBLICATIONS

Listed below is an updated list of papers, beginning with the most recent, written under the full or partial support of CaBS. Items (2) and (4) are attached as an Appendix.

- (1) Small, L.F. and S.G. Ellis (In prep.). Fecal carbon production by zooplankton: the effects of body size and carnivorous feeding.
- (2) Altabet, M.A. and L.F. Small (1990). Nitrogen isotopic ratios in fecal pellets produced by marine zooplankton. *Geochim. Cosmochim. Acta* 54:155-163.
- (3) Huh, C-A., L.F. Small, S. Niemi, B.P. Finney, B.M. Hickey, N.B. Kachel, D.S. Gorsline and P.M. Williams (1990). Sedimentation dynamics in the Santa Monica-San Pedro Basin off Los Angeles: radiochemical, sediment trap and transmissometer studies. *Cont. Shelf Res.* 10:137-164.
- (4) Small, L.F., M.R. Landry, R.W. Eppley, F. Azam and A.F. Carlucci (1989). Role of plankton in the carbon and nitrogen budgets of Santa Monica Basin, California. *Mar. Ecol. Prog. Ser.* 56:57-74.
- (5) CaBS Group (1989). Elemental cycling and fluxes off the coast of Southern California. *EOS* 70:146-155.
- (6) CaBS Group (1988). The Coastal Transition Zone program. *EOS* 69:698-707.
- (7) Huh, C-A., D.L. Zahnle, L.F. Small and V.E. Noshkin (1987). Budgets and behaviors of uranium and thorium series isotopes in Santa Monica Basin sediments. *Geochim. Cosmochim. Acta* 51:1743-1754.

IV. APPENDIX

1. Role of plankton in the carbon and nitrogen budgets of Santa Monica Basin, California.
2. Nitrogen isotopic ratios in fecal pellets produced by marine zooplankton.

Fig. 1 Composite relationships (means $\pm$ 1 s.d.) in the Santa Monica Basin photic zone in (a.) September/October and (b.) January/February. The large standard deviations associated with C/Pig in suspended particles and in fecal pellets were mainly the result of combining data within and outside the pigment max in the water column (these data can be separated for more detailed analyses). Thanks to M. Altabet, R. Eppley and M. Landry for certain data used herein.

END

DATE
FILMED
3126192

1