Grant DE-FG03-91ER20028
"A Genetic Analysis of Adh1 Regulation"

Progress report (period: June 1991 to February 1992)

Principle Investigator:
Michael Freeling, Professor
Department of Plant Biology
111 Genetics and Plant Biology Building
University of California, Berkeley
Berkeley, CA 94720
(510) 642-0924

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Progress report and non-competitive proposal

The overall goal of our research proposal is to understand the meaning of the various cis-acting sites responsible for Adh1 expression in the entire maize plant.

Studies on the TATA box and analysis of revertants of the Adh1-3F1124 allele.

1. Previously we isolated and characterized a revertant Adh1-3F1124r17 allele, which bears a deletion of 18 nucleotides removing the TATA box and its immediate vicinity. In this revertant the level of Adh1 expression increased from 6 to 60% of wild type in anaerobic roots and scutellum while simultaneously expression in pollen decreased from 100% to 30%. Our most recent findings are, that in pollen Adh1 transcripts accumulate to wild type levels, while in anaerobic roots transcripts accumulate at 60% levels, a number which agrees only in root tissue with the observed enzymatic activity. While in roots multiple transcription initiation sites, clustered around the wild type site, are utilized, in pollen transcription initiates at least 25 nucleotides further upstream. These data suggest that the TATA box region influences post-transcriptional processes in the male gametophyte but not in somatic root tissue. Different organs may provide surrogate TATA function in different ways, leading to organ-specific differences. We published this work recently in the EMBO Journal (Kloeckener-Gruissem et al. 1992 see appendix).

Our observations raise further questions which we will address during the next granting period: (1) What is the precise Adh1 transcription initiation site in the revertant pollen? RNase protection and primer extension experiments will show the answer. (2) Which post-transcriptional step is causing the reduced enzyme activity in pollen? We will establish a polysomal profile from male tassel tissue and pollen and use a wild type and a revertant-specific transcription probe to determine whether translation of the 5'-extended revertant Adh1 mRNA can be initiated or whether translational elongation may be effected. (3) Are the revertant-specific elongated transcripts specifically made in gametophytic tissues? We will investigate the transcription pattern in immature ears. We conducted initial in situ experiments which demonstrate that ADH activity is differentially found in the developing immature ear, prior to fertilization. We are currently analyzing Adh1 transcripts from these tissues.

2. We proposed to discover the meaning of the Adh1 TATA box sequences by site-directed mutagenesis. We completed the construction of all relevant substitution/deletion promoter constructs. Their effect on transcription was assayed transiently after electroporation into two different tissue culture lines: the Black Mexican Sweet (BMS) cell line which was derived from mesocotyl tissue and a more embryo like cell line P3377. Although we have not yet been able to assign a single dramatic effect to a particular nucleotide of the TATA box, the following general observations and conclusions can be stated: (i) The two different cell lines exhibit a differential level of expression when the same mutant construct was tested;

indicating differential expression in vitro, just as we observed in vivo in revertant r17. (ii) The embryonic cells express almost every mutant construct better in comparison to the BMS cells; suggesting that less differentiated tissues can tolerate the TATA box mutations better than a more differentiated cell. (iii) The largest of all tested deletions (the 18 bp deletion which mimics the r17 deletion), results in 20% level of reporter gene expression. This level of expression is not as high as the 60% expression observed in vivo in roots but is closer to the 30% in pollen. Again, we interpret these data just as proof of differential gene expression in different tissues in vivo and in vitro and that each tissue requires specific factors which, in combination, allow proper interpretation of the meaning of a cis-acting site. The electroporation experiments also demonstrate that another transient expression system needs to be employed, one which comes closer to the in vivo situation. For these purposes we proposed to improve the utilization of the particle gun as a mean to deliver DNA into living tissues. We will report progress on the mechanistic aspect of those experiments below; here we will focus on initial experiments of bombardment of seedling roots with wild type and the 18 bp deletion constructs. Up to date we tested only aerobic seedling roots for reporter gene expression and found a drastic decrease in expression when tissues were bombarded with the deletion mutant construct in comparison to the wild type construct.

Besides repeated verification of these initial observations we are planning to test the expression in anaerobic roots, shoots, immature ears and tassels.

3. A third experimental approach to the understanding of the meaning of the TATA box sequences focuses on protein factors that recognize and bind the TATA box. We proposed to isolate the gene(s) encoding the transcription factor TFIID, which was shown in yeast and mammalian cells to bind the TATA box and to play a role in initiation of transcription. Substantial progress was made towards isolation and characterization of the maize TFIID genes. In contrast to animals and yeast, the maize genome carries two unlinked loci as determined by southern blot analysis and cDNA cloning. The cDNA clone for locus one (TFIID-1) was identified based on sequence homology to primers obtained from animal TFIID. A fragment was obtained in a PCR reaction which was used to screen a maize seedling cDNA library. With the identified locus one, a second cDNA clone, representing the expression of the locus two, was isolated. We studied the expression pattern of the two loci with the primary question why plants (note that Arabidopsis also carries two TFIID loci as reported by Chua's laboratory) have two loci. Are these loci differentially expressed with respect to gametophytic vs somatic tissue or is their expression developmentally regulated. We know now, that both loci are actively transcribed and that they are differentially expressed only with respect to somatic tissue. Experimental complications, primarily based on the fact that both genes are very similar in sequence composition and exhibit only one short region of non-homology to use for loci-specific probing, have posed problems in the identification of loci-specific transcripts.

Goal-oriented future experiments will focus on in vitro expression of both maize TFIID loci and the isolation of antibodies. At this point we will combine the TATA box mutant constructions with the TFIID studies and both, the over expressed TFIID proteins and their antibodies will be used in binding assays to analyze the requirements of the sequences within and surrounding the Adh1 TATA box.

Studies on cis-acting sequences required for root-specific Adh1 expression.

We proposed to analyze a revertant Adh1-3F1124-r53, which exhibits a novel tissue-specific effect on Adh1 expression: 50% activity in the scutellum and pollen and no activity in aerobic and anaerobic roots. We postulated that this pattern fits a knock-out of a root-specific box, rather than an anaerobic-specific box which exerts its effects in the roots.

During our attempts to obtain a promoter fragment using the PCR reaction we discovered that homology to one (or both) of the primers was disturbed by the mutation. We performed detailed Southern Blot analyzes which focused on short restriction fragments encompassing the 1 kbp BamHI-HindIII promoter region and concluded that this mutation is not a simple deletion or insertion but rather a combination of both or an inversion. It became clear that precise nucleotide information is needed of the r53 promoter region which extends the initial limits set by the primers as described in our proposal. We are currently involved in the cloning of the 11 kbp genomic BamHI fragment which contains the entire r53 locus and up- and downstream sequences into bacteriophage lambda. Upon sub-cloning of the promoter region we will perform DNA sequencing reactions to identify the lesion in the r53 allele.

The initial phenotypic characterization of the r53 mutant as described above and in the proposal was extended to monitor ADH1 expression in other organs, tissues and cells. We obtained preliminary evidence that shows a novel hypoxic inducibility of ADH1 activity in the normally not inducible aleurone layer of an imbibed kernel. We will spend continuous effort on the analysis of this revertant allele in the whole plant, at all developmental stages and under various environmental conditions. As did revertant r17, this revertant offers new surprises about sequence requirements and their consequences on Adh1 expression.

Refinement of the use of the particle gun

The availability of the particle gun as a mean to deliver DNA constructs, whose transient expression can be measured in living tissues and cells, is essential to our studies. We are continuously working to improve the frequency of successful DNA delivery. We tested some of the obvious technical aspects and decided to conduct our experiments according to the best results which are listed below. One major breakthrough in the ease of measuring bombardment success came with the availability of a plasmid

pC17 which contains the maize color gene C1/ B-Peru driven by a 35S promoter (provided by Michael Fromm). We discovered that expression of this gene within 24 hours after bombardment of seedlings carrying the genotypes of our Adh1-3F, Adh1-3F1124 and Adh1-3F1124r53 alleles results in red colored epidermal and sub-epidermal cells, reflecting frequencies of success. Using this color reaction as a measuring unit we found consistently fewer spots in seedling roots versus seedling shoots. We find best results if every material is assaulted with two shots, we determined the optimal distance between the position of the macroprojectile and the living material and the use of two metal screens of different mesh sizes placed between the projectile and the target material. It should be noted that "best" and "optimal" reflects the momentary conditions and we continue with improvements.

Screening for more, different mutants that affect Adh1 expression differentially.

1. Search for mutations in genes coding for trans-acting proteins.

We chose the transposon Mutator system as the mutagen. Mu-active plants (in which transposon activity was monitored by an unstable color gene "bronze") which carried a normal, active Adh1-1F allele, were grown to maturity. Alive, ADH1 positive pollen grains were collected and treated with allyl alcohol vapors, which kill ADH positive pollen. Ideally, only ADH mutants with no enzyme activity will be able to survive. These pollen grains were crossed to tester ears, carrying an active Adh1 allele whose products migrate at a distinguishable position in an electrical field. The scutellum of the resulting kernels were subjected to ADH analysis; regardless of the enzyme activity pattern, every kernel from this experiment is currently growing and will be self pollinated as well as outcrossed to tester plants. Progeny will be subjected to further tests of ADH enzyme activity as well as DNA analysis. Some numbers may demonstrate the powerful selection of this procedure: We started with approximately 5×10^7 pollen grains and yielded approximately 100 kernels of which 10 lack an active ADH1 enzyme in the scutellum. We anticipate several scenarios of events: (i) The mutation occurred within the Adh1 locus: these mutants will be saved. (ii) The mutation occurred at an unlinked site: these mutants are what we are truly searching for. This unlinked site might code for a trans-acting factor which is either generally or differentially required for Adh1 expression. Analysis of the Adh1 expression pattern and DNA structure will allow us to identify which mutant belongs to which category.

2. Search for additional mutants derived from a partial revertant of Adh1-3F1124.

Revertant r16 was described in our proposal. It carries 30% of wild type activity in the scutellum and roots but shows no alteration in pollen expression levels (=100% of wild type). The transposon Mu3 did not excise

and is still sitting in the Adh1 promoter. In situ ADH activity in aleurone cells indicates that this allele, like the mutant 1124 is also unstable. We will take advantage of this instability and with varying selection conditions we will screen for either up or down mutation. In preparation for this screen, we are currently constructing a maize line that carries a morphological marker linked to the Adh1-3F1124r16 allele. As marker we chose the knotted phenotype, a dominant mutation which is located one map unit apart from Adh1. Homozygous r16 plants were crossed to a Knotted plant which carries an Adh1-1S allele. Heterozygous kernels are now being self pollinated and will be analyzed for recombination events leading to the Adh1-r16 allele linked to Knotted.

Functional analysis of a non-glycolytic anaerobic protein.

We continued work on one of the anaerobically inducible genes, ANP31. Repeated DNA sequence analysis of a full-length cDNA clone led us to rename the gene as ANP27. The clone hybridizes to a message of approximately 1.1 kb that is induced 2-3 fold in hypoxic seedling roots and shoots. We can not detect transcripts in aerobic tissue, nor in cold- or heat-treated seedlings. The cDNA contains an open reading frame of 765 nucleotides, encoding a polypeptide of 255 amino acids with a predicted molecular mass of 27.5 kDal. The protein is prolin (10%) and arginine (10%) rich; yet it does not show obvious sequence homology to known proteins rich in these two amino acids. We are currently constructing the appropriate plasmids to obtain in vitro expressed polypeptides and consequently to induce the production of antibodies. With these molecules in hand we hope to elucidate more of the (hopefully regulatory) functional aspects of this gene.

Publications resulting from work supported by this grant:

Kloeckener-Gruissem B, Vogel JM and M Freeling (1992) The TATA box promoter region of maize Adh1 affects its organ-specific expression. EMBO J. 11, 1, 157-166.

The general scope of the proposal does not change. No budget changes are requested.

END

**DATE
FILMED**

4/10/92