AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

PREPARATION OF UNSYMMETRICALLY LABELED OXYGEN MOLECULES AND THEIR USE TO ELUCIDATE OXYGEN METABOLISM

Evan H. Appelman
Chemistry Division, Argonne National Laboratory

Takashi Ogura and Teizo Kitagawa
Institute for Molecular Science, Okazaki, Japan

Constantinos Varotis, Yong Zhang and Gerald T. Babcock
Department of Chemistry, Michigan State University, East Lansing

A novel technique has been developed to synthesize in large quantities an O₂ molecule that is unsymmetrically labeled with the oxygen isotopes oxygen-16 and oxygen-18. This unusual molecule, ¹⁶O¹⁸O, has been synthesized by Dr. Evan H. Appelman at Argonne National Laboratory in work supported by the DOE Division of Chemical Sciences and has been utilized as a unique spectroscopic probe of the mechanism of metabolism in living organisms. Preparation of this molecule requires prior synthesis of the exotic precursor hypofluorous acid, HOF, and Dr. Appelman has recently developed an improved method for synthesizing this precursor that permits the preparation of essentially unlimited quantities of the labeled oxygen. This, in turn, has made it practical to use this oxygen in Raman-spectroscopic studies of the mechanism by which O₂ is reduced by the enzyme cytochrome oxidase, a key step in the metabolic utilization of oxygen by living organisms. These studies have involved Dr. Appelman in collaboration with two different research groups: that of Professor Teizo Kitagawa at the Institute for Molecular Science, Okazaki, Japan, and that of Professor Gerald T. Babcock at Michigan State University. The results indicate that the interaction of O₂ with the iron of the enzyme may be represented:

The results of the collaboration with the Michigan State group have been accepted for publication in the Proceedings of the National Academy of Sciences.

MASTER

PREPARATION OF UNSYMMETRICALLY LABELED OXYGEN MOLECULES AND THEIR USE TO ELUCIDATE OXYGEN METABOLISM

Evan H. Appelman

Chemistry Division, Argonne National Laboratory

Takashi Ogura and Teizo Kitagawa

Institute for Molecular Science, Okazaki, Japan

Constantinos Varotis, Yong Zhang and Gerald T. Babcock

Department of Chemistry, Michigan State University, East Lansing, MI

Some 20 years ago, the compound hypofluorous acid, HOF, was first synthesized at Argonne National Laboratory by Drs. Martin Studier and Evan Appelman. Appelman subsequently showed this compound to be an unusually powerful reagent for effecting oxidation and oxygen-transfer reactions. He further demonstrated that HOF could be labeled with the oxygen isotope ^{18}O , and in that case it could provide a convenient method for labeling other molecules with oxygen-18, i.e.,

Of especial interest was the fact that this approach made possible the synthesis of "unsymmetrically" labeled hydrogen peroxide and oxygen molecules in a neat state:

(Heretofore these unsymmetrical molecules could only be synthesized as components of statistical mixtures containing a variety of isotopic species.)

It was soon realized that these unsymmetrically labeled oxygen molecules could be very useful as probes in Raman-spectroscopic studies of the bonding of O_2 and H_2O_2 in complexes of biological interest. The first application was in a collaboration with Professors D. F. Shriver and I. M. Klotz and their colleagues at Northwestern University to determine the configuration in which oxygen molecules

were bonded to the iron in hemoglobin. At issue was a long-standing controversy as to whether the oxygen was bonded in an unsymmetrical end-on configuration, as in diagram A below, or in a symmetrical side-on or "bridged" configuration, as in diagram B. If unsymmetrically labeled oxygen were used and the resonance Raman spectrum of the resulting complex examined, configuration A should show two resonance-enhanced Raman bands corresponding to the Fe—O stretch, whereas configuration B should show only one such band. In fact, two bands were observed, indicating that the complex actually had the end-on configuration.

More recently, a collaboration was undertaken with Drs. Thomas Loehr and Joann Sanders-Loehr and their colleagues at the Oregon Graduate Center to study systems that served as models for biochemical oxygen bonding. The bonding of hydrogen peroxide to the $\text{Fe}^{\text{III}}(\text{edta})$ complex was examined, and in this case it was found that a symmetrical side-on or "bridged" structure was formed.

For many years, however, practical applications of unsymmetrically-labeled oxygen and hydrogen peroxide were limited by the fact that HOF was an unstable and potentially dangerous compound that could only be made in very small quantities. A change in this situation was presaged about five years ago, when Professor Shlomo Rozen, an organic chemist at Tel Aviv University in Israel, accidentally observed that passage of elemental fluorine through wet acetonitrile produced a powerfully oxidizing species. He turned to Argonne for help in interpreting this observation, and Appelman was able to demonstrate that the oxidizing species was actually hypofluorous acid that was being stabilized by complexing to the acetonitrile.

Appelman realized that this offered a safe and convenient method for preparing hypofluorous acid in arbitrarily large quantities, and that it could therefore also permit the production of corre-

spondingly large quantities of unsymmetrically labeled molecular oxygen. This, in turn, made possible new applications that had been unfeasible so long as only very small amounts of the unsymmetrically labeled oxygen were available. In particular, biochemists have been keenly interested in understanding the reduction of molecular oxygen by the enzyme cytochrome oxidase, a key step in the course of biological metabolism. A variety of oxygen-containing intermediates have been postulated in the complex mechanism of this enzymatic reduction. Distinction among various possibilities for such intermediates could be greatly facilitated if Raman-spectroscopic experiments could be carried out using unsymmetrically labeled O_2 . However, since this reduction is a rapid dynamic process, it is usually studied in a flow system, and much greater quantities of the labeled O_2 would be required than had been needed for the equilibrium studies of hemoglobin and the iron-EDTA complex. The new method for synthesis of HOF, leading to the ability to make relatively large amounts of $^{16}O^{18}O$, made such an experiment feasible, and collaborations were entered into with Professor Teizo Kitagawa and his group at the Institute for Molecular Science at Okazaki, Japan, and with Professor Gerald Babcock and his group at Michigan State University. The results indicate that a Raman band previously assigned to the iron(III)-peroxyspecies $Fe^{3+}-OOH^-$ actually belonged to a subsequently formed iron(IV) species, $Fe^{4+}=O$, so that the sequence of interaction between O_2 and the iron of the enzyme may be represented:

This reaction scheme is shown in greater detail on the accompanying figure.

The collaborative study with the Michigan State group has been accepted for publication in Proceedings of the National Academy of Sciences.

This work has been supported by the Division of Chemical Sciences, Office of Basic Energy Sciences, U.S. Department of Energy.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Cytochrome Oxidase Catalysis

**DATE
FILMED**

7/6/94

END