DOE Report [for annual report]
University of Wisconsin-Madison
Madison, WI 53706

DE92 012123

DOE/ER/13707--2

Enzymology of Biological Nitrogen Fixation (DE-FG02-87ER13707)
R. H. Burris, Department of Biochemistry

Two genes involved in the regulation of nitrogenase activity, draT and draG, were cloned and found to be contiguous on the Azospirillum brasilense chromosome. The nifH gene, encoding dinitrogenase reductase, is near to draT with an intervening gap of 1.9 kb. The organization of these genes in Azospirillum lipoferum and Rhodospirillum rubrum is similar, but nifH and draT are separated by only 400 bp in the organisms. A. brasilense draTG is very similar to draTG in R. rubrum with 91.8% similarity and 85.3% identity at the amino acid level. Apparently A. brasilense uses the normal ATG initiation codon for draT, and draG. The genes for A. brasilense were able to restore function to appropriate mutants of R. rubrum. The heterologous expression of A. brasilense draTG in R. rubrum was not fully normal, as it responded more slowly to darkness and more quickly to ammonia than wild type cells. Our mutational analysis of the draTG region of A. brasilense confirms the function of these genes in the regulation of nitrogenase activity, but it also revealed minor but demonstrable differences in the control systems of R. rubrum and A. brasilense.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DOE Grant: DE-FG02-87ER13707 - Enzymology of Biological Nitrogen Fixation
R. H. Burris, Department of Biochemistry
University of Wisconsin-Madison
Madison, WI 53706

Progress Report

Two genes involved in the regulation of nitrogenase activity, draT and draG, were cloned and found to be contiguous on the A. brasilense chromosome. The nifH gene, encoding dinitrogenase reductase, is near to draT with a 1.9-kb intervening gap and is divergently transcribed. Similar organization of these genes has been found in A. lipoferum and R. rubrum, but in the latter organism nifH and draT are separated by only 400 bp.

A. brasilense draTC is extremely similar to draTC of R. rubrum at both amino acid and nucleotide levels, but the similarity was lower than that of their nifH genes; they showed 91.8% similarity and 85.3% identity at the amino acid level. Particularly in draT, there seem to be several regions of extreme conservation, probably reflecting the involvement of those regions of the gene product in its catalytic function or its regulation.

R. rubrum draT has the unusual initiation codon TTG, which was verified by determination of the N-terminus of DRAT from that organism. This initiation codon is relatively rare in E. coli, and it usually supports poor translation of the gene, which is consistent with the low amount of DRAT in cells. The start codon of A. brasilense draT is less clear. We propose that it uses a normal ATG as initiation codon, preceded by a Shine-Dalgarno sequence of GAGAG, but there are two other possibilities. A TTG codon is located at the tenth position, but is not preceded by a reasonable Shine-Dalgarno sequence. More interesting, a comparison with the R. rubrum sequence suggests that the 17th codon of the A. brasilense sequence might be the start site. CTG, like TTG, is typically a leucine codon, but it has not previously been identified as a start codon. Since A. brasilense DRAT has not been purified and its N-terminus is unknown, we cannot reach a firm conclusion about the start codon.

draG appears to use ATG as an initiation codon, and it can encode a protein of 32.4 kDa, consistent with the characterization of the purified protein. The sequence also revealed the first 42 nucleotides of an ORF immediately downstream of draG that was very similar to a similarly positioned ORF of R. rubrum. In R. rubrum, the product of this ORF appears to have a secondary role in the regulation of the DRAG/T system. The sequence comparison of the draTC to GenBank showed similarity only for R. rubrum; it was not similar to the ADP-ribosyltransferase of bacterial toxins.

The functionality of the identified genes was shown both by mutational analysis in A. brasilense and by their ability to complement appropriate mutants of R. rubrum. The dra mutants behaved rather similarly to those described in R. rubrum. In particular, the ability of a draG mutant to display active nitrogenase upon derepression, and to respond irreversibly to ammonium, strongly suggests that DRAT activity itself is subject to post-translational regulation as seen in R. rubrum.

The expression of the A. brasilense dra genes in R. rubrum revealed several other interesting features of the system. The ability of DRAT/G from A. brasilense to respond to cycles of darkness/light indicates sufficient similarity to the R. rubrum products that they still interact with the light/dark signal transduction pathway. Although there have been several studies of signal transduction to this system, including the effects of energy charge, pyridine nucleotide pools and amino acid pools, neither the small molecules nor protein factors involved are known.

Residual nitrogenase activity was observed in wild type and dra mutants of R. rubrum after a shift to ammonia or darkness, but no residual activity was found in A. brasilense after ammonia addition. R. rubrum transconjugants containing A. brasilense draTC genes also displayed significant residual activity. This suggests that the residual activity probably can be attributed to other factors in R. rubrum, rather than a specific difference in the DRAG/T system. Consistent with this, no residual nitrogenase activity has been reported in ammonia-treated K. pneumoniae when draTC genes from either R. rubrum or A. lipoferum were expressed in that organism. Higher residual nitrogenase activities (20-30%) in R. rubrum transconjugants exposed to darkness or ammonium chloride may reflect greater DRAG accumulation (multicopy of draG gene) or inappropriate regulation of the heterologous DRAG.

The heterologous expression of A. brasilense draTC in R. rubrum was not completely "normal," since it showed a slower response to darkness and a faster response to ammonia than did wild-type R. rubrum. A. brasilense itself shows a relatively fast response to ammonia, so that then DRAG/T regulatory systems in the two organisms are not identical.

Whereas no hybridization of a nifJ probe has been reported in the region between draT and nifH, a putative NtrA-dependent promoter has been described upstream of nifH which would be transcribed toward dra. However, A. brasilense strains UB5 and UB6, containing insertion mutations in the region between draT and nifH, showed no detectable effect on nitrogenase or its regulation by DRAG/T.

Our mutational analysis of the draTC region of A. brasilense confirms the function of these genes in the regulation of nitrogenase activity. It also demonstrates that the draTC genes are not essential for growth, nor are there duplicated genes or redundant functions in this organism. While the general behavior of the DRAG/T system in A. brasilense is similar to that seen in R. rubrum, there are a number of interesting differences in the timing and extent of the regulatory response that make the comparative analysis of the two systems important.

The project will continue to support the investigation of the control system for nitrogenase in Azospirillum spp. as described in accompanying reprints and the paper that will be submitted soon by Zhang, Burris and Roberts.

Progress has been slow in our efforts to determine the kinetics of N_2 reduction and protein reduction by nitrogenase, but we hope that continuing difficulties with the rapid reaction equipment can be resolved so that the quantitative time-course of reduction of N_2 and H^+ can be established more accurately.

We have confirmed that Acetobacter diazotrophicus grows on N_2 at a pH below 3, whereas most nitrogenases are inoperative below pH 5. The organism utilizes a variety of substrates, but it metabolizes glucose and acetate particularly rapidly. Again we have been unable with varied methodologies to demonstrate any exchange reaction accompanying the fixation of N_2 by Azotobacter vinelandii.

A 943 page book on "Biological Nitrogen Fixation" has been assembled and edited by G. Stacey, R. H. Burris and H. J. Evans. It appeared in November 1991, but the publishers have dated it 1992.

Publications:

R. H. Burris. Acetylene Reduction to Measure Biological Nitrogen Fixation. Current Contents [Citation Classics] 33, #38, 18 (Life Sciences) and 21, #38, 18 (Agriculture, Biology & Environmental Sciences) (1990).

G. P. Roberts, P. W. Ludden, R. H. Burris, W. P. Fitzmaurice, H.-A. Fu, G. Nielsen, J.-H. Liang, L. Lehman, D. Woehle, D. Lies, H. Wirt, S. Montgomery, R. Davis and Y. Bao. The Genetics and Biochemistry of the Reversible ADP-Ribosylation Systems of Rhodospirillum rubrum and Azospirillum lipoferum. In, "Nitrogen Fixation: Achievements and Objectives," pp. 475-481. Gresshoff, Roth, Stacey and Newton, eds. Chapman and Hall, New York, 1990.

R. H. Burris. Congress Summary: the Molecular Perspective. In, "Nitrogen Fixation: Achievements and Objectives," pp. 845-850. Gresshoff, Roth, Stacey and Newton, eds. Chapman and Hall, New York, 1990.

H. I. Miller, R. H. Burris, A. K. Vidaver and N. A. Wivel. Risk-Based Oversight of Experiments in the Environment. Science 250, 490-491 (1991).

R. H. Burris. Nitrogenases. J. Biol. Chem. 266, 9339-9342 (1991).

R. H. Burris, A. Hartmann, Y. Zhang and H. Fu. Control of Nitrogenase in Azospirillum sp. Plant and Soil 137, 127-134 (1991); In, "Nitrogen Fixation," pp. 89-96. M. Polsinelli, R. Materassi and M. Vincenzini, eds., Kluwer Academic Publishers, Dordrecht, Netherlands, 1991.

J. Liang, G. M. Nielsen, D. P. Lies, R. H. Burris, G. P. Roberts and P. W. Ludden. Mutations in the draT and draG Genes of Rhodospirillum rubrum Result in Loss of Regulation of Nitrogenase by Reversible ADP-Ribosylation. J. Bacteriol. 173, 6903-6909 (1991).

G. Stacey, R. H. Burris and H. J. Evans, eds. Biological Nitrogen Fixation, 943 pp. Chapman & Hall, New York, 1992.

H. J. Evans and R. H. Burris. Highlights in Biological Nitrogen Fixation during the Last 50 Years. In, "Biological Nitrogen Fixation," pp. 1-42. G. Stacey, R. H. Burris and H. J. Evans, eds. Chapman & Hall, New York.

Reprints removed

END

**DATE
FILMED**

6 / 11 / 92