APR 27 1992

LITERATURE AND DATA REVIEW FOR THE
SURFACE-WATER PATHWAY: COLUMBIA RIVER
AND ADJACENT COASTAL AREAS

Hanford Environmental Dose
Reconstruction Project

W. H. Walters
R. L. Dirkes
B. A. Napier

April 1992

Prepared for
review and evaluation
by the Technical Steering Panel

Pacific Northwest Laboratory
Richland, Washington 99352

DISCLAIMER

This report was prepared under the direction of the HANFORD ENVIRONMENTAL DOSE RECONSTRUCTION PROJECT Technical Steering Panel by Battelle Memorial Institute's Pacific Northwest Laboratories operating the Pacific Northwest Laboratory for the U.S. Department of Energy (DOE). While funding for the work was provided by DOE, the work is not under DOE direction or control. The views and opinions of the authors expressed in this document do not necessarily reflect those of the United States Government or any agency thereof. Reference herein to any specific commercial product, process or service by trade name, trademark, manufacturer or otherwise does not necessarily constitute or imply its endorsement, recommendation or favoring by the U.S. Government or any agency thereof, nor by Battelle Memorial Institute.

Printed in the United States of America

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401. FTS 626-8401.

Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

LITERATURE AND DATA REVIEW FOR THE
SURFACE-WATER PATHWAY: COLUMBIA RIVER
AND ADJACENT COASTAL AREAS

April 1992

This document has been reviewed and approved
by the Technical Steering Panel.

J. E. Till, Chair
Technical Steering Panel

Date

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

for

PREFACE

In 1987, the U.S. Department of Energy (DOE) directed the Pacific Northwest Laboratory (PNL), which is operated by Battelle Memorial Institute, to conduct the Hanford Environmental Dose Reconstruction (HEDR) Project. The DOE directive to commence project work followed a recommendation by the Hanford Health Effects Review (HHER) Panel in 1986. The HHER Panel was formed to consider the potential health implications of historic Hanford Site releases of radioactive materials. The HEDR Project objective is to estimate radiation doses to individuals and population groups from exposure to radioactive emissions that began in 1944 from the Hanford Site.

An 18-member Technical Steering Panel (TSP) was selected by the Vice Presidents for Research at the major universities of Washington and Oregon to direct the project work. The TSP consists of experts in the various technical fields relevant to HEDR project work and representatives from the states of Washington, Oregon, and Idaho; Native American tribes; and the public.

A December 1990 Memorandum of Understanding between the Secretaries of the DOE and the U.S. Department of Health and Human Services (DHHS) transferred responsibility for managing the DOE's dose reconstruction and exposure assessment studies to the DHHS. This transfer resulted in the current process to establish a contract between Battelle and the Centers for Disease Control (CDC), which is an agency of the DHHS, to continue performance of the project. The TSP will continue as technical director for the work conducted by Battelle.

The scope of this report is to 1) describe the segment of the Columbia River system and the adjacent coastal areas that were involved in the transport and distribution of radionuclides released at Hanford; 2) review and summarize the availability of river and coastal area monitoring data (Hanford and offsite sources) providing radionuclide concentrations for water, sediment, and biota; 3) review and summarize the reports and studies pertaining to the release and transport of radionuclides in the river; and 4) calculate preliminary dose estimates for selected locations and times to include freshwater and marine food sources.

Because of the extent and complexity of the database and related information, an analysis of the reviewed material is beyond the scope of this report. While it is recognized that the data quality will vary among the many sources and sampling techniques, the data analysis and data quality evaluation will be carried out in later studies during FY 1992.

ABSTRACT

As part of the Hanford Environmental Dose Reconstruction Project, Pacific Northwest Laboratory reviewed literature and data on radionuclide concentrations and distribution in the water, sediment, and biota of the Columbia River and adjacent coastal areas. Over 600 documents were reviewed including Hanford reports, reports by offsite agencies, journal articles, and graduate theses. Certain radionuclide concentration data were used in preliminary estimates of individual dose for the 1964-1966 time period. This report summarizes the literature and database review and the results of the preliminary dose estimates.

The initial sampling of river water began in 1945 and of riverbed sediments in 1948. The sampling was confined to the Hanford to Pasco (Washington) reach of the Columbia River until 1950 when water samples were collected at Bonneville Dam. The sampling of river biota in the Hanford area began in 1950. Environmental monitoring by Hanford contractors increased significantly from 1951 through 1957 both in the Hanford area and downstream. From 1958 to the present, the monitoring program at Hanford was streamlined and now involves mainly the Hanford to McNary Dam reach of the river.

Comprehensive monitoring and studies by offsite agencies began in 1960 and continued into the early 1980s. Much of the work was conducted below The Dalles Dam out to the coastal areas of Washington and Oregon. The work of these agencies focused mainly on the uptake of radionuclides by freshwater and marine biota and the distribution of the reactor effluent along the coastal areas. Other work investigated the uptake of radionuclides by river sediments. Only gross beta measurements are available from 1945 through 1957 from both Hanford and offsite sources. After 1957, gamma spectrum measurements were able to determine activity levels of specific radionuclides.

For this report, preliminary dose estimates were calculated for several locations from Richland, Washington to Willapa Bay, Washington. Maximally exposed individuals were those who consumed significant quantities of fish or seafood. The important radionuclides were zinc-65, arsenic-76, and phosphorus-32. The drinking water pathway was the most important for typical

individuals, who were assumed to have eaten much less fish or seafood. The important radionuclides were arsenic-76, neptunium-239, and zinc-65.

EXECUTIVE SUMMARY

This literature review investigated the availability of radiological data and information for the Columbia River and adjacent coastal areas (see Plates 1 and 2 in pockets at the back of this report for locations mentioned in this summary and in the report). A major objective was to locate data on the concentrations of radionuclides in water, sediment, and biota for the surface-water transport pathway. Because the database will be used to reconstruct past radioactivity levels for the surface water system, the size, geographic extent, and continuity (location and time) of the historical database are of prime importance. The historical nature of the HEDR Project rules out supplementing the existing database through new data collection; therefore, the use of mathematical methods will be required to fill any serious gaps in the database. Mathematical approaches that could be used to fill the gaps will be investigated during the follow-on development work on the conceptual model.

Studies of the reactor effluent by Hanford and offsite contractors provided a significant amount of information on the directional movement of the reactor effluent in the river and adjacent coastal areas. The results of these studies identified some of the reasons for variability in radionuclide concentrations in the water, sediment, and biota. They also provided information on the uptake and release and distribution of radionuclides by river and coastal sediments.

During the literature review, data were selected from several locations along the river for preliminary dose estimates. The locations extended from Richland, Washington, downstream to Astoria, Oregon, and along the Washington coast to Willapa Bay. The dose estimates considered treated drinking water, freshwater fish, and various seafoods.

MONITORING LOCATIONS AND TIME PERIODS

From 1945 to 1950, monitoring of radionuclide concentrations in water and sediment (sediment sampling began in 1948) was conducted primarily in the vicinity of the reactors and downstream to the 300 Area, which is just

upstream of Richland. The only sampling farther downstream was at Richland and Pasco. The Yakima River was sampled as an indicator of background levels of radioactivity in water samples. There were no sampling locations downstream of Pasco until 1950, when water sampling began at Bonneville Dam. Sediment sampling was conducted only in the Hanford reach (from upstream of the reactors to the Umatilla-McNary Dam location). Most of the sediment samples were collected near the reactors, but others were collected at Richland, Kennewick, and the Pasco bridge. No monitoring or investigative studies were conducted by offsite contractors during this period. Thus, there are no radiological data available downstream of Pasco until 1950, and then only for the Bonneville Dam location. Also, only gross beta readings were taken since the samplings predated gamma spectroscopy equipment.

Beginning in 1951, and lasting until 1958 when gamma spectroscopy equipment became available, many more water sampling locations were added downstream of Pasco. Locations were added at McNary Dam in 1951. By 1957, twelve locations, between McNary Dam and Vancouver, Washington, had been added. Most of the water sampling was conducted during 1953 and 1957, with only intermittent sampling for the other years. From 1951 to 1957, sediment sampling was extended to McNary Dam and to Paterson, Washington, about 18 miles downstream of McNary. Sediment was not sampled at any other downstream locations.

Routine sampling of fish commenced in 1950, but only in the Hanford reach of the river. Various studies of biota continued and the number of samples increased during the 1950s, with most, but not all, of the effort remaining in the Hanford reach. Hanford contractors began sampling shellfish near the mouth of the Columbia River in 1953. From 1950 to 1977, fish and shellfish were routinely sampled and analyzed by Hanford contractors, with much of the effort concentrated on freshwater whitefish in the Hanford reach and Willapa Bay oysters from the Washington coast. Both whitefish from the river and oysters from Willapa Bay were popular sources of food for humans.

It was also during the early 1950s that the U.S. Public Health Service (USPHS) began its water quality and biota studies. This effort, which lasted about 3 years, was the first radiological monitoring program by an offsite agency for the Columbia River. The USPHS water quality and biota studies

began in 1951 and continued into 1953. The results provide the first independent database for comparison with the Site monitoring by Hanford contractors. The sampling work was limited to the length of river from just upstream of the reactors to Paterson, Washington, downstream of McNary Dam. Only brief, exploratory surveys were conducted at Bonneville Dam, at Portland, Oregon, and at the mouth of the Columbia River.

Although the sampling locations and frequency increased significantly from 1945 through 1957, the results were reported as gross beta concentrations. While gross beta concentrations are not suitable for direct use in dose calculations, they can be used in preliminary dose estimates as described in Section 10 of this report.

After 1957, concentrations of specific radionuclides became available for water, sediment, and biota samples collected by Hanford contractors. Beginning about the same time, Hanford contractors began streamlining the monitoring program. Sediment, which was not sampled in 1958, was sampled once more in 1959, but sampling was discontinued after that year because information on radionuclide concentrations in sediment was not directly usable in dose estimates. Hanford contractors continued to collect water samples but the number of locations decreased, significantly, to several locations of specific interest. Most of these were in the Hanford to McNary Dam reach of the river. After 1958, water sampling locations downstream of McNary were limited to five sites: 1) Paterson, Washington (1959 only); 2) The Dalles, Oregon (1963 and 1964); 3) Hood River, Oregon (1959 only); 4) Bonneville Dam (1964 through 1975); and 5) Vancouver, Washington (1959 through 1963).

Beginning about 1960, offsite agencies commenced a series of studies and monitoring programs which involved the states of Washington and Oregon, their state universities, and the U.S. Geological Survey (USGS). Much of the work was focused on the Columbia River estuary and adjacent coastal areas and was oriented toward transport processes. Most of the data are not useful for dose calculations but do provide very good information regarding the disposition of the reactor effluent. The data that are the most useful are those radionuclide concentrations measured in fish and shellfish; these data were found in various reports, journal articles, and graduate theses. Some of these data

were used to calculate the preliminary dose estimates presented in Section 10. Other useful data are radionuclide concentrations in water and biota from the Oregon State Department of Health. From 1961 through 1977, Department staff routinely sampled Columbia River water from The Dalles downstream to the mouth of the river, as well as estuarine and coastal fish and shellfish. Some sampling continued into 1983 but was very limited. Although the sampling was mostly quarterly for each year, the results provide at least a partial data-base for evaluation of activity levels in water and biota at downstream locations. Another use would be in testing the credibility of computed water concentrations for downstream locations and time periods.

REACTOR EFFLUENT RELEASES AND TRANSPORT

A review of the Hanford contractor reports indicates a significant amount of variability in effluent releases from the reactor outfalls in the river. As a group, the reports discussing the variation in effluent activity levels identify six primary factors for the variations: 1) the increase in the number of reactors on-line and the operating power levels, 2) the activation of natural elements in raw river water, 3) chemical additives, 4) the rupture of fuel elements, 5) the purging of reactors to remove radioactive film buildup, and 6) the length of time effluent stayed in the retention basins (amount of decay).

As the number of reactors increased from three in 1945 to eight by 1955, the quantity of radionuclides released to the river increased. Beginning in 1957, the operating power levels were also gradually increased with correspondingly higher radionuclide releases to the river. The natural increase or decrease in the elements in raw river water is a seasonal event, with higher concentrations of these elements occurring during the spring flood season. The concentrations were also source (drainage-basin) dependent. Chemical additives were continually used during reactor operations. Fuel element ruptures and reactor purging were discrete events lasting hours and occurring on an intermittent basis. Basin retention times were not constant and were gradually shortened over the years as the need warranted.

Once the reactor effluent was discharged from the outfalls, it traveled downstream in a relatively narrow plume several hundred feet wide. Although the plume gradually mixed laterally, it was still distinguishable at Pasco. Beyond Pasco the flow patterns in the McNary reservoir and Snake River inflow preclude any definitive measurements. The plume tended to hug the reactor shoreline (west shoreline) downstream past Richland to the Yakima River confluence. Beyond that point, the plume was found along the Pasco shore, possibly as a result of being forced across the channel by the Yakima River inflow. Water concentration measurements along the plume centerline were upwards of five to seven times as high, in the vicinity of the reactors, compared with those in the ambient river water. Near Pasco, the concentrations were much closer but differences were still distinguishable.

Another factor that possibly contributed to the higher water concentrations along the plant shore was the shoreline springs that released retention basin leakage through river bank soils. Basin leakage releases through the shoreline were occurring at B, C, D, and F reactors during the 1950s.

The rate of transport of the effluent downstream to the Columbia River mouth is of key importance with respect to radionuclide decay and activity levels. Hanford contractors began in 1955 to develop a relationship between river discharge and downstream travel time of effluents. By the early 1960s, a general knowledge of travel times from the reactor area to Astoria, Oregon, was available. Travel times from the reactor area to eight locations from Pasco, Washington, to Astoria, Oregon, for four ranges of discharge were estimated using field data. Since McNary Dam was in place in 1955 and The Dalles Dam was filling, the times are not valid for the years prior to 1955. Travel times for those years would be shorter and would need to be determined mathematically.

SEDIMENT AND RADIONUCLIDE TRANSPORT STUDIES

By the 1950s, the role of river sediment in the uptake and storage of radionuclides and the effect of sediments on concentrations in the water were recognized. Hanford contractors and the USGS conducted a number of studies into sediment storage effects. It was also determined that the sediments

could release radionuclides back into the river water, causing an increase in concentration exceeding that expected from effluent releases. This process provided another factor for variation in the water concentrations.

During the early 1960s, in cooperation with Hanford contractors, the USGS began studying the uptake of radionuclides by river sediments. These cooperative studies continued through most of the 1960s, with final reports being published in the early 1970s. The reports produced data on sediment concentrations and a considerable amount of information on uptake and release of radionuclides by sediment. Water concentration data are also included in the reports. The USGS studies were strongly oriented toward river processes and did not provide the lengthy and consistent sampling needed for dose calculations. Nonetheless, the results provide some data to assist in determining activity levels at locations downstream of McNary Dam during the sampling period.

Comprehensive programs to monitor the radioactivity in shoreline sediments began in 1959. Before that time, only exploratory measurements had been made in the vicinity of the reactors. Shoreline monitoring continued into the late 1970s, but was not conducted on a continuing yearly basis. These shoreline surveys included the surfaces of channel islands and the measurement of exposure rates for boaters on the river. The highest overall rates were found along the reactor areas, as would be expected, with the maximum measured on the island offshore of the 100-D area. Below the reactor areas, the exposure rates decreased but not uniformly. The survey results reported by Sula (1980) found no definitive downstream decrease in shoreline exposure rates from below the reactor areas to the confluence of the Snake River.

PRELIMINARY DOSE ESTIMATES

Using radionuclide concentration data from the various monitoring programs and studies summarized in this report, preliminary dose estimates for the years 1964 through 1966 were calculated for five locations along the river: 1) Richland, Washington, 2) McNary Dam area, 3) Bonneville Dam area, 4) Astoria, Oregon, and 5) Willapa Bay, Washington. The estimates considered a maximally exposed individual and a typical individual.

Maximally exposed individuals were assumed to have eaten significant quantities of fish or seafood from the Columbia River. Included in the doses were contributions from drinking water, swimming, boating, and exposure to shoreline contamination. For the maximum individual, the most important pathway was consumption of nonmigratory fish. The important radionuclides were zinc-65, arsenic-76, and phosphorus-32, with zinc-65 the most important at downstream locations because of its relatively longer half-life. The doses decreased downstream except for slight increases at Bonneville with the addition of salmon to the diet and at Astoria with the addition of shellfish to the diet.

Drinking water was the most important exposure pathway for typical individuals since they were assumed to eat very little fish. The important radionuclides in the Hanford reach (from the reactor area to Umatilla and McNary Dam) were arsenic-76, neptunium-239, and zinc-65. At locations farther downstream, zinc-65 and phosphorus-32 were the most important. Examples of annual doses for 1964 are shown in Table ES.1.

TABLE ES.1. Preliminary Estimates of Annual Doses from Columbia River Pathways for 1964

<u>Location</u>	<u>Maximally Exposed Individual (rem)^(a)</u>	<u>Typical Individual (rem)</u>
Richland	1.6×10^{-1}	7.8×10^{-3}
McNary	1.7×10^{-2}	1.0×10^{-3}
Bonneville	2.1×10^{-2}	3.8×10^{-4}
Astoria	4.0×10^{-2}	2.0×10^{-3}
Willapa Bay	1.6×10^{-2}	1.6×10^{-3}

(a) rem = roentgen equivalent man

FOLLOW-ON WORK

The data and information summarized in this report will be analyzed during follow-on work conducted in FY 1992 to determine the extent to which the data can be used to support dose estimates. The amount, continuity, and variability of river data, together with the changes in the river system (e.g., dams) have a direct bearing on the data's usefulness and computational techniques proposed for reconstructing concentrations in water, sediment, and, possibly, biota. The data analysis will be used to produce a conceptual model of the Columbia River that will describe the processes affecting radionuclide transport and distribution and the relationship between the key variables. From the conceptual model work, several alternative approaches will be proposed for providing the necessary database to support dose estimates.

ACKNOWLEDGMENTS

This report was prepared for the Technical Steering Panel, which serves as the technical director of the Hanford Environmental Dose Reconstruction Project, as a summary of data and information on the Columbia River dose pathway. In accomplishing this task, the authors received assistance and reviews from many people.

The authors wish to express appreciation to S. B. Yabusaki and W. A. Perkins for their valuable and timely assistance in the collection and review of the offsite documents and to C. E. Cushing for preparation of the information on Columbia River ecology.

Thanks are extended to L. K. Grove for the initial editing of the draft report in preparation for the review process and to J. W. Thielman, K. R. Hanson, and J. E. Danko for the final editing and assembly of the report. A special thanks to B. G. Miller for his expertise in computer graphics.

ACRONYMS AND ABBREVIATIONS

CDC	Centers for Disease Control
CRAG	Columbia River Advisory Group
CREDDP	Columbia River Estuary Data Development Program
DHHS	U.S. Department of Health and Human Services
DOE	U.S. Department of Energy
HEDOP	Hanford Environmental Dose Overview Panel
HEDR	Hanford Environmental Dose Reconstruction
HHER	Hanford Health Effects Review (Panel)
PNL	Pacific Northwest Laboratory
PNRBC	Pacific Northwest River Basin Commission
SESP	Surface Environmental Surveillance Project
TSP	Technical Steering Panel
USAEC	U.S. Atomic Energy Commission
USGS	U.S. Geological Survey
USPHS	U.S. Public Health Service

UNITS

cfs	cubic feet per second
Ci	curies
dpm/g	disintegrations per minute per gram
dpm/L	disintegrations per minute per liter
m ³	cubic meter
μCi/g	microcuries per gram
μCi/L	microcuries per liter
μg/m ³	micrograms per cubic meter
μR/hr	microroentgens per hour
mL	milliliter
mrep/hr	millireps per hour
mR/hr	milliroentgens per hour
nCi/m ³	nanocuries per cubic meter
nCi/g	nanocuries per gram
ppm	parts per million
pCi/g	picocuries per gram
pCi/L	picocuries per liter
pCi/kg	picocuries per kilogram
RM	river mile
rem	roentgen equivalent man

RADIONUCLIDES

¹²⁴Sb antimony-124
⁷⁶As arsenic-76
¹⁴⁰Ba barium-140
¹⁴⁴Ce cerium-144
¹³⁷Cs cesium-137
⁵¹Cr chromium-51
⁵⁸Co cobalt-58
⁶⁰Co cobalt-60
⁶⁴Cu copper-64
¹⁵²Eu europium-152
¹⁵⁴Eu europium-154
⁷²Ga gallium-72
¹²⁹I iodine-129
¹³¹I iodine-131
⁵⁵Fe iron-55
⁵⁹Fe iron-59
¹⁴⁰La lanthanum-140
⁵⁴Mn manganese-54
⁵⁶Mn manganese-56

²³⁹Np neptunium-239
⁹⁵Nb niobium-95
³²P phosphorus-32
²³⁹Pu plutonium-239
²⁴⁰Pu plutonium-240
⁴⁰K potassium-40
¹⁰³Ru ruthenium-103
¹⁰⁶Ru ruthenium-106
¹⁵³Sm samarium-153
⁴⁶Sc scandium-46
²⁴Na sodium-24
⁸⁹Sr strontium-89
⁹⁰Sr strontium-90
⁹⁹Tc technetium-99
¹³²Te tellurium-132
³H tritium
⁶⁵Zn zinc-65
⁹⁵Zr zirconium-95

CONTENTS

PREFACE	iii
ABSTRACT	v
EXECUTIVE SUMMARY	vii
ACKNOWLEDGMENTS	xv
ACRONYMS AND ABBREVIATIONS	xvii
1.0 INTRODUCTION	1.1
2.0 TECHNICAL APPROACH	2.1
3.0 SURFACE-WATER PATHWAY	3.1
3.1 PATHWAY FROM HANFORD TO McNARY DAM	3.3
3.2 McNARY DAM TO BONNEVILLE DAM	3.4
3.3 BONNEVILLE DAM TO COLUMBIA RIVER MOUTH	3.5
3.4 COASTAL AREAS	3.6
4.0 COLUMBIA RIVER ECOLOGY	4.1
4.1 PHYTOPLANKTON	4.2
4.2 PERIPHYTON	4.5
4.3 MACROPHYTES	4.7
4.4 ZOOPLANKTON	4.7
4.5 BENTHIC ORGANISMS	4.8
4.6 FISH	4.8
4.6.1 Anadromous Species	4.9
4.6.2 Resident Species	4.12
5.0 REACTOR OPERATIONS AND EFFLUENT WATER COMPOSITION	5.1
5.1 TIMETABLE OF REACTOR OPERATIONS	5.1
5.2 ELEMENTS IN COLUMBIA RIVER WATER	5.4

5.3	CHEMICAL ADDITIVES	5.5
5.4	PURGES	5.6
5.5	ACCIDENTAL RELEASES	5.7
5.5.1	Shoreline Releases	5.8
5.5.2	Fuel Element Ruptures	5.8
5.5.3	300 Area Waste Pond Dike	5.10
5.6	BASIN RETENTION TIME	5.11
5.7	EFFLUENT DISCHARGE TO RIVER	5.11
5.8	ACTIVITY LEVELS AND VARIABILITY OF RELEASES	5.12
6.0	COLUMBIA RIVER MONITORING AT HANFORD	6.1
6.1	1944 THROUGH 1957	6.1
6.2	1958 THROUGH 1971	6.5
6.3	1972 THROUGH 1990	6.9
7.0	SUMMARY OF RESULTS OF COLUMBIA RIVER MONITORING BY HANFORD CONTRACTORS	7.1
7.1	WATER MONITORING DATA	7.2
7.2	SEDIMENT MONITORING DATA	7.3
7.3	BIOTA MONITORING DATA	7.8
8.0	SPECIAL STUDIES AND SURVEYS AT HANFORD	8.1
8.1	EFFLUENT PLUME DISPERSION	8.1
8.2	SHORELINE RADIATION SURVEYS	8.3
8.2.1	Survey of 1959	8.3
8.2.2	Survey of 1961-1962	8.4
8.2.3	Shoreline Surveys Reported in 1966	8.6
8.2.4	Post-Reactor Shutdown Surveys	8.8
8.3	DOWNRIVER TRAVEL TIMES	8.10

9.0	MONITORING AND STUDIES BY OFFSITE AGENCIES	9.1
9.1	STATE OF WASHINGTON	9.1
9.2	UNIVERSITY OF WASHINGTON	9.2
9.3	STATE OF OREGON	9.4
9.4	OREGON STATE UNIVERSITY	9.6
9.4.1	Columbia River and Estuary	9.10
9.4.2	Coastal and Ocean	9.10
9.5	U.S. GEOLOGICAL SURVEY	9.15
9.5.1	Initial Studies at Hanford	9.15
9.5.2	U.S. Geological Survey Cooperative Studies	9.16
9.6	U.S. PUBLIC HEALTH SERVICE	9.22
10.0	SCREENING CALCULATIONS OF RADIATION DOSE AT VARIOUS DOWNSTREAM LOCATIONS FOR SEVERAL YEARS	10.1
10.1	RADIONUCLIDE CONCENTRATIONS IN RIVER WATER	10.2
10.2	RADIONUCLIDE CONCENTRATIONS IN AQUATIC BIOTA	10.4
10.3	EXPOSURE SCENARIOS AND PATHWAYS USED IN SCREENING CALCULATIONS	10.10
10.4	EXPOSURE PATHWAY MODELS	10.13
10.5	DOSE RESULTS	10.13
10.6	SCALING MID-1960s DOSES TO OTHER TIMES	10.19
11.0	REFERENCES	11.1
	APPENDIX A - COLUMBIA RIVER HYDROLOGIC DATA SOURCES	A.1
	APPENDIX B - TABLE OF RADIONUCLIDES	B.1
	APPENDIX C - BIBLIOGRAPHY	C.1

FIGURES

(Plates are in pockets in the back of the report)

Plate 1	Locations of Water and Sediment Sampling by Hanford Contractors	
Plate 2	The Hanford Reach - Showing Surveys of Shoreline Radioactivity Exposure Rates and the Effluent Plume Centerline	
3.1	Columbia River Profile Showing Conditions in 1944 and Dam and Reservoir Construction in Later Years	3.2
4.1	Simplified Food Web of Columbia River	4.2
4.2	Radionuclide Concentration in Columbia River Net Plankton for Chromium-51, Zinc-65, Iron-59, Phosphorus-32 and Manganese-54 . .	4.4
4.3	Concentrations of Zinc-65 in Columbia River Phytoplankton Following Closure of Hanford Reactors	4.5
4.4	Radionuclide Transport by Phytoplankton	4.6
4.5	Expected Occurrence of Salmonids in Hanford Reach of Columbia River	4.10
4.6	Hanford Reach Salmon Redd Counts	4.11
5.1	100-D Area Water System	5.2
5.2	Operating Periods for the Once-Through-Cooled Hanford Production Reactors	5.3
5.3	Monthly Average Radioactivity Release Rate in Ci/day for Reactor Effluent at the Point of Release to the Columbia River	5.13
7.1	Total Beta Concentrations in Columbia River Water at Pasco, Washington 1945-1971	7.3
7.2	Annual Average Radionuclide Concentrations in Columbia River Water at the Richland Pumphouse and Pasco Pump Station	7.7
7.3	Annual Average Radionuclide Concentrations in Columbia River Water at McNary Dam	7.8
7.4	Annual Average Radionuclide Concentrations in Columbia River Water at Bonneville Dam	7.9
7.5	Annual Average Radionuclide Concentrations in Columbia River Water at Vancouver, Washington	7.10

7.6	Monthly Mean Phosphorus-32 Concentrations in Columbia River Whitefish, 1964-1966	7.11
7.7	Monthly Mean Zinc-65 Concentrations in Columbia River Whitefish, 1964-1966	7.12
7.8	Zinc-65 Concentrations in Willapa Bay Oysters, 1959-1977	7.14
9.1	Columbia River and Coastal Sampling Locations for 1961-1967 Studies by the State of Oregon	9.5
9.2	Columbia River and Coastal Sampling Locations for 1967-1977 Studies by the State of Oregon	9.9
9.3	Winter Plume Pattern of February 1966 Based on Chromium-51 Activity	9.12
9.4	Summer Plume Pattern of June 1965 Based on Chromium-51 Activity	9.13
10.1	Locations for Which Dose Estimates Were Made	10.1

TABLES

ES.1	Preliminary Estimates of Annual Doses from Columbia River Pathways for 1964	xi
4.1	Selected Fish Species Categorized by Interest Groups	4.9
5.1	Fuel and Target Element Failures Removed from the Reactors	5.9
6.1	Summary of Routine Locations for Sampling Columbia River Water, 1945-1957	6.3
6.2	Summary of Routine Locations for Sampling Columbia River Sediment, 1948-1957	6.4
6.3	Summary of Routine Locations for Sampling Columbia River Water, 1958-1971	6.7
6.4	Analyses for River Water Samples, 1958-1971	6.8
6.5	Summary of Routine Locations for Sampling Columbia River Water, 1972-1990	6.10
7.1	Annual Average Radionuclide Concentrations in Columbia River Water at the Richland Pumpouse	7.4
7.2	Annual Average Radionuclide Concentrations in Columbia River Water at the Pasco Pump Station	7.5
7.3	Annual Average Radionuclide Concentrations in Columbia River Water at McNary Dam	7.5
7.4	Annual Average Radionuclide Concentrations in Columbia River Water at Bonneville Dam	7.6
7.5	Annual Average Radionuclide Concentrations in Columbia River Water at Vancouver, Washington	7.6
7.6	Radionuclide Concentrations in Surface Sediments in McNary Reservoir in 1971 and 1976	7.11
7.7	Annual Average Radionuclide Concentrations in Willapa Bay Oysters, 1959-1977	7.13
8.1	Travel Times of Peak Concentrations in the Columbia River 100-D Reactor to Various Downstream Points, Measured During July 1964	8.11
9.1	Range of Yearly Concentrations of Zinc-65 and Chromium-51 at Columbia River Locations for 1962 Through 1967	9.7

9.2	Range of Yearly Concentrations of Zinc-65 at Oregon Coast Locations for 1962 Through 1967	9.8
9.3	Maximum and Average Gross Beta Activities at River Mile 362 . .	9.23
10.1	Annual Mean Radionuclide Concentrations at Various Locations in Columbia River Water	10.3
10.2	Radionuclide Concentrations in Selected Monitored Biota in the Columbia River Estuary at Astoria	10.5
10.3	Reported Average Radionuclide Concentrations in Selected Pacific Coastal and Ocean Biota	10.6
10.4	Freshwater Fish Bioaccumulation Factors Used in This Analysis .	10.7
10.5	Exposure Pathway Parameters Assumed for Maximally Exposed Individuals	10.11
10.6	Exposure Pathway Parameters Assumed for Typical Individuals . .	10.12
10.7	Summary of Annual Doses from Columbia River Pathways for the Maximally Exposed Individual	10.14
10.8	Detail of Annual Doses Reported in Table 10.7 for Astoria . . .	10.15
10.9	Detail of Annual Doses Reported in Table 10.7 for Bonneville . .	10.15
10.10	Summary of Annual Doses from Columbia River Pathways for the Typical Individual	10.16
10.11	Annual Average Total Beta Concentrations, Columbia River at Pasco, Washington	10.20

1.0 INTRODUCTION

The Surface-Water Transport Task is part of the Hanford Environmental Dose Reconstruction (HEDR) Project, which was established to estimate the radiation doses people may have received from operations that began at the Hanford Site in 1944. The technical work is being conducted by Battelle staff at the Pacific Northwest Laboratory (PNL) under the direction of an independent Technical Steering Panel (TSP).

This task addresses the radioactivity in the Columbia River, which received cooling-water effluent from the eight Hanford once-through-cooled reactors^(a) and was the major pathway for waterborne radionuclides. The pathway began at the Hanford Site and continued downstream past the mouth of the Columbia River to the adjacent coastal and ocean areas. The overall objective of the task is to provide radionuclide concentrations at locations along the pathway for water, sediment, and biota for use in final dose estimates. Preliminary dose estimates for several years are provided in Section 10.

The purpose of this report is to summarize the significant results of a literature and database review conducted during FY 1991. The review included available reports and data pertaining to the river from both Hanford contractors and offsite sources. The period of interest was from 1944 to 1990, with emphasis on the period from 1944 through 1972, when some or all of the eight once-through-cooled reactors were operating. Although a detailed summary of all the literature and data gathered is beyond the scope of this report, all of the documents reviewed are listed either in the references or the appendixes. The data that were determined to be potentially useful or descriptive for the river pathway are presented as maximum values or ranges of values in tables and graphics.

Following the Executive Summary, this report contains 11 sections, three appendixes, and two plates. Section 2 presents the technical approach of the study. In Section 3, the surface-water pathway from the Hanford reactors to

(a) In once-through-cooling, water drawn from the river passes through the reactor core and is returned to the river.

the coastal areas is described. Section 4 discusses the ecology of the Columbia River, with an emphasis on the fish pathway. Factors affecting the composition and variability of reactor effluent water are summarized in Section 5. Section 6 describes the development of Columbia River monitoring programs at Hanford from 1945 to the present, and Section 7 summarizes the monitoring results. Section 8 discusses the results of special studies conducted by Hanford contractors that directly pertained to the transport processes of the river. Section 9 summarizes monitoring and studies by off-site agencies. Section 10 presents the preliminary results of dose calculations for selected time periods and data sets from Hanford and offsite sources. Section 11 provides a list of references. Appendix A provides a brief summary of the Columbia River hydrologic data that are available for use in evaluating radiological data and transport calculations. Appendix B is a table of radionuclides discussed in the report and their half-lives. Appendix C is a bibliography of additional references examined during the preparation of this report but not cited in the text. Two plates are provided in pockets in the back of the report to help the reader follow the discussions that refer to locations and activities along the Columbia River from the Hanford Site to the river mouth.

2.0 TECHNICAL APPROACH

The literature review covered two general categories of source documents: 1) those prepared by Hanford Site contractors and 2) those prepared by offsite agencies. The documents originating at the Hanford Site were identified through the HEDR Information Resources Task (Task 11), as described by Shipler (1992).

The documents prepared by offsite agencies were obtained by the principal investigators directly from the originating agencies. Most of these offsite sources were known, although the extent of their document holdings was not. The agencies that have radiological data and information for the Columbia River pathway are the state health organizations of Washington and Oregon and the state universities that offer related programs of study in marine science. The primary federal agencies are the U.S. Geological Survey (USGS) and the U.S. Public Health Service (USPHS). Nonradiological river data were obtained from both the USGS and the U.S. Army Corps of Engineers.

Each document was reviewed for possible use in the HEDR Project. All of the Hanford-originated environmental monitoring reports (mostly data) are part of a permanent database regardless of the amount of data and information included. Titles appear in the list of references (Section 11) if they are cited in the body of this report; otherwise they appear in Appendix C. Topical reports on special studies of the Columbia River system originating from Hanford are similarly listed. Any special studies that related to the transport and deposition of radionuclides in the Columbia River are summarized in this report, together with information regarding reactor operations and effluent releases. Much of this special study information was scattered throughout numerous reports and has been assembled in various sections in the report.

Visits were made to offsite agency libraries (state universities and science centers) and catalog searches were conducted onsite. All documents found in this way that contained radiological data and information on the Columbia River and adjacent coastal areas were reviewed for possible use in the project. The list of references in each document was also reviewed for other source documents. Other agencies were contacted directly for assistance

in locating copies of their river-related publications. The data and information from offsite agencies are organized by agency in this report.

The data and information from all sources were reviewed for radionuclide activity level measurements in Columbia River water. Activity levels in river and coastal sediment and biota sampled within or affected by the reactor effluent pathway were also reviewed. Because an extensive presentation of monitoring data is beyond the scope of this report, only ranges of activity levels, maximum values, or averages are included in the summaries. These values are included to provide some idea of the levels present during the years of Hanford operations. River and coastal locations and the time periods of the monitoring and special studies are discussed. No calculations or data analyses were conducted, except for calculation of the preliminary dose estimates presented in Section 10. Because the documents spanned a period from 1945 to 1990, the reported units of activity levels changed. No conversion of units to a common system was attempted for this report; therefore, all units given are the same as in the original documents.

3.0 SURFACE-WATER PATHWAY

From September 1944 to January 1971, the Columbia River served as a source of process water and a medium for disposal of cooling water from a once-through process for as many as eight production reactors at the Hanford Site. A ninth reactor (100-N) had a closed-loop cooling system and used the river as a source of secondary cooling water. The effluent water from N Reactor was released to a holding crib and trench system that allowed the water to percolate into the underlying geologic media where most of the radionuclides were retained by the soil. From the time the first reactor (100-B) went on-line in September 1944 until 1971, cooling water was discharged into the river in a continuous release, as long as at least one of the reactors was operating. Several days after the startup of 100-B, the first radioactive effluent reached the mouth of the Columbia River and began contributing to a plume that extends into the Pacific Ocean and along the coastal areas. In Plate 1 (in a pocket in the back of this report), the Columbia River pathway is shown as the river exists today with all the dams in place. Figure 3.1 shows the river profile as it was in 1944 and the sequence of dam construction from 1953 to 1967.

Once the effluent was discharged into the river, the concentration and distribution of radioactivity in the river water became largely dependent on river discharge and the uptake of radionuclides by sediment and algae. The primary hydrologic factors were the seasonal variations in water volume and flow velocities. The water volume directly affects the concentration by dilution. The flow velocity controls the rate of transport through the system and, therefore, influences the degree of radionuclide decay that has occurred by the time any one point downstream is reached. The sediment tends to accumulate significant quantities of certain radionuclides in the channel bed. Through this process, the sediment removes some of the radionuclides from the water and acts as a "sink" for a portion of the radioactivity. This sediment sink also provides a medium for certain biota to take up radionuclides, thus introducing them into the food chain.

FIGURE 3.1. Columbia River Profile Showing Conditions in 1944 and Dam and Reservoir Construction in Later Years

For purposes of discussion, the Columbia River pathway is separated into three river reaches and the coastal area impacted by the reactor effluent. The reach from Hanford to McNary Dam includes the reactors and is where Hanford contractors conducted most of their monitoring and studies. The reach from McNary Dam to Bonneville Dam included only one reservoir in 1944, but by 1967 was totally controlled by dams and reservoirs. The reach from Bonneville Dam to the mouth includes the river's estuary and is where tidal processes dominate the transport of radionuclides.

There are numerous references available from which descriptive information on the Columbia River can be obtained. The primary references used in this section are the reports and maps listed in Appendix A, especially the documents published by the Columbia River Estuary Data Development Program (CREDDP 1984a, 1984b). The text by Pruter and Alverson (1972) was another primary source. The river mile values, as called out in the text, were obtained from an index prepared by the Pacific Northwest River Basins Commission (PNRBC 1972).

3.1 PATHWAY FROM HANFORD TO McNARY DAM

The effluent pathway began at the reactor (100-B) farthest upstream, at river mile 384 (RM 384) above the mouth of the Columbia River. As other reactors came on-line over the years following 1944, the area in which effluents were released extended from RM 384 downstream to RM 369. The releases to the river from the retention basins came from outfall lines (pipes) near the river bottom and took the form of a narrow plume that gradually spread and dispersed downstream. Because of the shoreline location and proximity of the reactor outfalls to each other, these plumes tended to coalesce and hug the Richland side of the river. The various channel islands, the roughness of the channel bed (i.e., the presence of boulders), the location of pools and riffles, and the curvature of the river's natural flow all affected the rate at which the plume spread and mixed with the river water. Under some flow conditions, the plume was not entirely mixed over the full river width until it approached Pasco in the vicinity of RM 330. The phenomenon of downstream plume mixing, as it occurs in the reach from Hanford to McNary, will be discussed in more detail in Section 8. The river system from the Hanford reactors to McNary Dam is shown in Plates 1 and 2.

Three tributaries join the Columbia River between Hanford and McNary Dam: the Yakima River at RM 335, the Snake River at RM 324, and the Walla Walla River at RM 315. All three, but especially the Snake River, dilute the effluent and contribute a significant volume of sediment to the Columbia River.

The mean annual discharge of the Columbia River at Hanford is 121,512 cubic feet per second (cfs). The mean annual discharges for the tributaries are as follows: the Yakima River at Kiona, Washington (3661 cfs); the Snake River at Ice Harbor Dam (53,948 cfs); the Walla Walla River at Touchet, Washington (593 cfs) (Williams and Pearson 1985).

When the first reactors came on-line during the 1940s, McNary Dam (RM 292) did not exist. Construction of the dam began during the early 1950s and culminated with the raising of the pool upstream of the dam in December 1953. This created Lake Wallula, a body of water with a maximum depth of about 100 feet and a length of about 62 miles. Before the construction of

McNary Dam at RM 292, the flow, with its effluent and sediment loads, passed through Wallula Gap and past Umatilla, Oregon (near the present damsite) under free-flow conditions. Since construction of the dam, the flow velocity within the influence of Lake Wallula is considerably reduced, and much of the sediment load is trapped behind the dam. However, as is true of the other Columbia River dams, some of the trapped sediment is resuspended and transported downstream by seasonal high discharges.

3.2 McNARY DAM TO BONNEVILLE DAM

The reach of river from McNary Dam to Bonneville Dam is shown in Plate 1. From 1944 to 1956, there were no dams on the river between the McNary-Umatilla, Oregon, site (RM 292) and Bonneville Dam (RM 146.1) as shown in Figure 3.1. The flow was unrestricted until the upper limit of the Bonneville reservoir was reached. This length of the Columbia River included many ancestral Native American fishing grounds, such as the Celilo Falls. With the construction of The Dalles Dam (RM 191.5) and raising of the reservoir pool in 1956, the Celilo Falls fishing ground was inundated (Pruter 1972). By 1967 the John Day Dam (RM 215.6) was also in operation, and the river in this reach consisted of three reservoir pools with no open channel flow. After 1969, Native American fishing grounds were largely confined to tributary streams and the reservoirs of Bonneville Dam and The Dalles Dam (Pruter 1972). With the construction of The Dalles and John Day dams, the river water and effluent flow velocity were reduced and much of the sediment inflow was trapped in the reservoirs. Another characteristic of this length of river is that the river flows through the Columbia River Gorge with no appreciable flood plain both before and after dam construction.

Numerous small rivers and creeks discharge into the river between McNary and Bonneville Dams. The three largest rivers, in downstream order, are John Day River (RM 218), the Deschutes River (RM 204.1), and the Klickitat River (RM 180.4). With the combined inflows, some dilution effects occurred, but nothing comparable to that associated with the Snake River upstream of McNary Dam. The mean annual discharges of the tributaries are as follows: the Klickitat River near Pitt, Washington (1607 cfs) (Williams and Pearson 1985);

the John Day River at McDonald's Ferry, Oregon (2103 cfs); the Deschutes River near Biggs, Oregon (5869 cfs) (Moffatt, Wellman, and Gordon 1990).

3.3 BONNEVILLE DAM TO COLUMBIA RIVER MOUTH

Below Bonneville Dam, the river enters the estuary of the Columbia River where the width, depth, and flow characteristics (tides, multiple channels) of the Columbia River change considerably (Plate 1). At about RM 52 the river is less than 1 mile wide, but it increases to nearly 9 miles at about RM 20. There are several bays and headlands within the estuary, which consists of multiple channels separated by numerous islands, bars, and shoals. Deep areas (e.g., Gray's Point), where water depths approach 100 feet (Neal 1972), can be found in the estuary apart from the main channel.

The three largest tributaries contributing inflow below Bonneville Dam are the Willamette River (RM 101.5), the Lewis River (RM 87), and the Cowlitz River (RM 68). These tributaries all enter at a considerable distance upstream of the estuary. The mean annual discharges of the tributaries are as follows: the Willamette River near Portland, Oregon (33,310 cfs) (Moffatt, Wellman, and Gordon 1990); the Cowlitz River at Castle Rock, Washington (9330 cfs); the Lewis River at Ariel, Washington (4899 cfs).

At some point in the vicinity of RM 23 to 25, the maximum intrusion of ocean water occurs and marks the upstream extent of the Columbia River estuary (Neal 1972). Some disagreement exists concerning the definition of an estuary. Early definitions considered the entire tidal portion of a river to be the estuary; more recent practice is to consider only the segment of river subject to salinity intrusion as the estuary boundary (Neal 1972), although tidal effects can extend further upstream into the freshwater areas. For the Columbia River, tidal fluctuations have been observed as far upstream as Bonneville Dam during low flow conditions. Under any flow conditions, tides are strong enough to reverse the flow as far as 53 miles upstream.

Because of the tidal conditions and flat channel gradient, Columbia River sediments deposit in the estuary. The upstream reaches of the Columbia River and its tributary system are the major source of sediment for the estuary; the size range of sediment is limited to the finer fractions, as is

typical for estuaries. These finer fractions consist of fine sand, silt, and clay. These are also the sediment sizes that most readily sorb with radionuclides. Deposits of coarse sand and gravel are rare in the estuary and are found mostly in areas of extreme scour where transport velocities are high. Much of the silt and clay fractions remain in suspension or are resuspended often enough to be eventually flushed from the estuary out to the continental shelf. Only about 20% of the silts and clays transported to the estuary tend to remain there, primarily in peripheral bays and inactive (sluggish) channels of the middle to upper estuary (CREDDP 1984a, 1984b). The remaining limited range of sediments (fine to medium sand) is transported either along the estuary bed or by intermittent suspension. Because velocities in the estuary are reduced, fine to medium sand tends to persist as the dominant sediment size.

3.4 COASTAL AREAS

The initial mixing of Columbia River water (and associated reactor effluent) with ocean water occurs within the estuary as a result of tidal currents, wind-generated wave action, variation in water density, and the turbulent flow regime of the river. As the water and effluent move away from the mouth of the river, wind-wave action and open-sea processes become the dominant mixing processes. The volume of water leaving the estuary is largely determined by the seasonal drainage basin runoff to the Columbia River. As the water-sediment-effluent mixture is transported away from the river mouth, it gradually increases in both salinity and volume as it mixes with ambient ocean water both horizontally and vertically. The outflow tends to maintain the form of a plume with distinct boundaries that are sharper near the mouth and become less distinct with increasing distance along the plume. The plume extends over a latitude range of about 1000 kilometers and a maximum seaward distance of about 600 kilometers (Barnes, Duxbury, and Morse 1972). The plume is mostly an ocean surface process that transports dissolved radionuclides, but which would have the capacity to transport the finer sediment sizes, such as clays and silts, with sorbed radionuclides originating from the river.

The ocean plume has two dominant seasonal patterns, largely controlled by wind direction: during winter the plume lies north of the river mouth and

inshore as a result of southerly winds, and during the summer the plume is directed south and offshore by northerly winds. The northerly plume movement extends beyond the Strait of Juan de Fuca, and the southerly movement extends to the border between Oregon and California (Barnes, Duxbury, and Morse 1972).

The coarser sediments that contain radionuclides and that pass the mouth of the Columbia River are mostly sands transported by bottom currents along the continental shelf. The directional pattern of these currents was determined during the 1960s by studying the movement of seabed drifters (perforated plastic discs) placed near the mouth of the river (Morse, Gross, and Barnes 1968; Barnes, Duxbury, and Morse 1972). Once the drifters were away from the direct influence of the river outflow, the predominant direction of movement was northward and toward the Washington coastal areas (e.g., Willapa Bay and Gray's Harbor). This evidently occurred regardless of the season. The northward pattern of movement extended to the Strait of Juan de Fuca and beyond to Vancouver Island, British Columbia. The study determined that the northward movement of drifters is affected in the area of the Strait with some movement of the drifters into Puget Sound. According to Barnes, Duxbury, and Morse (1972), the Strait and submarine valley that extends seaward across the shelf apparently act as a partial barrier to northward movement. The authors presumed the Strait would have a similar effect on any sediment transported to this location from the Columbia River.

4.0 COLUMBIA RIVER ECOLOGY

The Columbia River supports a large and diverse community of plankton, periphyton, macrophytes, benthic invertebrates, and fish. It is the fifth largest river in North America and has a total length of about 2000 kilometers (~1240 miles) from its origin in British Columbia to its mouth at the Pacific Ocean. Eleven hydroelectric dams were constructed on the Columbia River between 1933 and 1968. As a result, except for the Hanford reach, the ecosystem has been changed from free-flowing throughout its length to one that is now a series of large flow-through lakes. This change from a lotic to lentic habitat severely altered the aquatic habitat and resulted in significant changes in the biotic communities. Organisms originally adapted to a flowing-water regime either had to adapt to a still-water environment or perish. New still-water forms invaded these newly created habitats. As mentioned previously, a major Native American fishery located at Celilo Falls on the Columbia River above The Dalles was eradicated when The Dalles Dam was closed.

Major tributaries to the upper Columbia River include the Spokane, Okanagon, and Wenatchee Rivers. No tributaries enter the Columbia River during its passage through the Hanford Site. Several major tributaries enter the Columbia River, especially in the lower reaches. These include the Yakima and Snake Rivers, which enter the McNary pool; the Umatilla and John Day Rivers, which enter the John Day pool; and the Deschutes River, which enters the Bonneville pool. The Willamette River enters the Columbia River below Bonneville Dam.

The Columbia River is a very complex ecosystem because of its size, the number of manmade alterations, the diversity of the biota, and the size and diversity of its drainage basin. Streams in general, especially smaller ones, usually depend on organic matter from outside sources (terrestrial plant debris) to provide energy for the ecosystem. Large rivers, particularly the Columbia River with its series of large reservoirs, contain significant populations of primary energy producers (algae, plants) that contribute to the basic energy requirements of the biota. Phytoplankton (free-floating algae) and periphyton (sessile algae) are abundant in the Columbia River and provide

food for herbivores such as immature insects and certain fishes, which in turn are consumed by carnivorous species. Figure 4.1 is a simplified diagram of the food-web relationships in selected Columbia River biota and represents probable major energy and contaminant pathways. This is largely a consumptive-based food web based on known feeding habits.

4.1 PHYTOPLANKTON

Phytoplankton species identified from the Hanford reach include diatoms, golden or yellow-brown algae, green algae, blue-green algae, red algae,

FIGURE 4.1. Simplified Food Web of Columbia River (adapted from Cushing 1991)

and dinoflagellates. Diatoms are the dominant algae in the Columbia River phytoplankton, usually representing more than 90% of the populations. The main genera include *Asterionella*, *Cyclotella*, *Fragilaria*, *Melosira*, *Stephanodiscus*, and *Synedra* (Neitzel, Page, and Hanf 1981). Plankton populations in the Hanford reach are typical of those forms found in lakes and ponds. They are influenced by communities that develop in the reservoirs of upstream dams, particularly Priest Rapids Reservoir, and by manipulation of water levels below through dam operation in downstream reservoirs. A number of algae found as free-floating species in the Hanford reach of the Columbia River are actually derived from the periphyton; they are detached and suspended by the current and frequent fluctuations of the water level. Phytoplankton and zooplankton populations at Hanford are largely transient, flowing from one reservoir to another. There is generally insufficient time for characteristic endemic groups of phytoplankton and zooplankton to develop in the Hanford reach.

The peak concentration of phytoplankton is observed in April and May, with a secondary peak in late summer/early autumn (Cushing 1967a). The spring pulse in phytoplankton density is probably related to increasing light and water temperature rather than to availability of nutrients because sufficient phosphate and nitrate nutrient concentrations are always present. Minimum numbers of phytoplankton are present in December and January. Green algae (Chlorophyta) and blue-green algae (Cyanophyta) occur in the phytoplankton community during warmer months, but in substantially fewer numbers than the diatoms.

Phytoplankton respond rapidly to changes in ambient concentrations of radionuclides by absorption and, more importantly, by adsorption because of their large surface-to-volume ratios. Phytoplankton are a primary food source for filter-feeding organisms such as caddisfly larvae and clams. Watson et al. (1969) documented the rapid loss and uptake of radionuclides during a brief closure of the Hanford reactors that resulted in a rapid lowering and subsequent increase of radionuclides in the water (Figure 4.2). In addition, Cushing et al. (1981) describe the relatively rapid loss of radionuclides

Radionuclide Concentration in Columbia River Net Plankton

R9204041

FIGURE 4.2. Radionuclide Concentration in Columbia River Net Plankton for Chromium-51 (^{51}Cr), Zinc-65 (^{65}Zn), Iron-59 (^{59}Fe), Phosphorus-32 (^{32}P) and Manganese-54 (^{54}Mn) (Watson et al. 1969)

from phytoplankton following permanent closure of the plutonium-producing reactors at Hanford. This decline was found at Hanford, McNary Reservoir, and Bonneville Reservoir, with downstream decreases of lessening magnitude (Figure 4.3). Cushing (1967a) presented data showing the magnitude of radio-nuclide transport of phosphorus-32 and zinc-65 by phytoplankton (in terms of nanocuries per cubic meter [nCi/m^3] of water) at different times of the year. Transport was greatest in spring and late summer, coincident with increases in phytoplankton populations (Figure 4.4).

FIGURE 4.3. Concentrations of Zinc-65 in Columbia River Phytoplankton Following Closure of Hanford Reactors. The value 1067 picocuries per gram (pCi/g) dry weight was found at McNary in June 1971 (Cushing et al. 1981).

a) Concentration of Phosphorus-32 and Zinc-65 by Net Plankton

b) Concentration of Net Plankton and Associated Phosphorus-32 and Zinc-65

FIGURE 4.4. Radionuclide Transport by Phytoplankton (Cushing 1967a)

4.2 PERIPHYTON

Communities of periphytic species ("benthic microflora") develop on suitable solid substrata wherever there is sufficient light for photosynthesis. Peaks of production occur in spring and late summer (Cushing 1967b). Dominant genera are the diatoms *Acanthes*, *Asterionella*, *Cocconeis*, *Fragilaria*, *Gomphonema*, *Melosira*, *Nitzschia*, *Stephanodiscus*, and *Synedra* (Page and Neitzel 1978; Page, Neitzel, and Hanf 1979; Beak Consultants Inc. 1980; Neitzel, Page and Hanf 1981). This community is a significant food item for some fish species, especially suckers and carp (see below).

Periphyton, like phytoplankton, have a large surface-to-volume ratio and thus accumulate high levels of radionuclides (Cushing 1967b). They also react similarly in terms of rapid uptake and decreased rates in relation to ambient water concentrations. At low water levels, periphyton could contribute to the external dose received by people frequenting the shoreline.

4.3 MACROPHYTES

Macrophytes are sparse in the Columbia River because of the strong currents, rocky bottom, and frequently fluctuating water levels. Rushes (*Juncus* spp.) and sedges (*Carex* spp.) occur along the shorelines of the slack-water areas. Macrophytes are also present along gently sloping shorelines that are subject to flooding during the spring freshet and daily fluctuating river levels. Commonly found plants include *Lemna*, *Potamogeton*, *Elodea*, and *Myriophyllum*. Where they exist, macrophytes have considerable ecological value. They provide food and shelter for juvenile fish and spawning areas for some species of warm-water game fish. However, should some of the exotic macrophytes (Eurasian millfoil) increase to nuisance levels, they may encourage increased sedimentation of fine particulate matter. These changes could have a significant impact on the trophic relationships of the Columbia River.

4.4 ZOOPLANKTON

Zooplankton populations in the Hanford reach of the Columbia River are generally sparse. In the open water regions, crustacean zooplankters are dominant. Dominant genera are *Bosmina*, *Diaptomus*, and *Cyclops*. Densities are lowest in winter and highest in summer. Summer peaks are dominated by *Bosmina* and range up to 4500 organisms/cubic meter (m^3). Winter densities are generally less than 50 organisms/ m^3 . *Diaptomus* and *Cyclops* dominate in winter and spring, respectively (Neitzel, Page, and Hanf 1983). Zooplankton are important food items for juvenile salmonids when they are living in the interstitial water among the rocks.

4.5 BENTHIC ORGANISMS

Benthic organisms are found either attached to or closely associated with the substratum. All major freshwater benthic taxa are represented in the Columbia River. Insect larvae such as caddisflies (Trichoptera), midges (Chironomidae), and black flies (Simuliidae) are dominant. Dominant caddisfly species are *Hydropsyche cockerelli*, *Cheumatopsyche campyla*, and *C. enonis*. Other benthic organisms include molluscs, sponges, and crayfish. Peak larval insect densities are found in late fall and winter, and the major emergence is in spring and summer.^(a) Stomach contents of fish collected in the Hanford reach revealed that benthic invertebrates are important food items for nearly all juvenile and adult fish. There is a close relationship between food organisms in the stomach contents of fish and those in the benthic and invertebrate drift communities. Thus, this community is directly linked in the food pathway leading to man. Certain molluscs and crayfish are potential food items for people and thus also are involved directly in food pathways leading to man.

4.6 FISH

Gray and Dauble (1977) list 43 species of fish in the Hanford reach of the Columbia River. The brown bullhead (*Ictalurus nebulosus*) has been collected since 1977, bringing the total number of fish species identified in the Hanford reach to 44. Table 4.1 presents a selected list of species important as sport and commercial species and those important to Native Americans.

There are two groups of fish present in the Columbia River based on their life histories: 1) anadromous species, and 2) resident species. The predilection of these two groups in terms of their potential for accumulating radionuclides differs considerably. The anadromous species hatch in freshwater, grow and migrate to the ocean, and eventually return to freshwater to

(a) E. G. Wolf. 1976. "Characterization of the benthos community." Section 4 in Final Report on Aquatic Ecological Studies Conducted at the Hanford Generating Project 1973-1974. WPPSS Columbia River Ecological Studies, Vol. 1. Battelle, Pacific Northwest Laboratories, Richland, Washington.

TABLE 4.1. Selected Fish Species Categorized by Interest Groups

<u>Sport Fishing</u>	<u>Commercial Fishing</u>	<u>Native American Fishing</u>
Steelhead trout	Steelhead trout	Steelhead trout
Chinook salmon	Chinook salmon	Chinook salmon
Mountain whitefish	Coho salmon	Coho salmon
Walleye	Sockeye salmon	Sockeye salmon
Smallmouth bass	White sturgeon	White sturgeon

spawn. They are carnivorous, and actively feed as juveniles in the river and as they mature in the ocean, but do not feed during the spawning migration. Although they are most susceptible to accumulating radionuclides from the Columbia River food web during their brief residence as juveniles, there is evidence of some species accumulating radioactivity prior to their return to the Columbia River estuary (Kujala 1966). Muscle samples from spawned-out chinook salmon collected in the Hanford reach in 1988 indicated no measurable influence from radionuclides released to the Columbia River (Jaquish and Bryce 1989). On the other hand, resident fish species can spend essentially their entire life span feeding on contaminated food and being exposed to external contaminants, thus offering the potential for much higher levels of contamination.

4.6.1 Anadromous Species

Anadromous species that use the Columbia River as a migration route include the chinook salmon, sockeye salmon, coho salmon, and steelhead trout; these are also the species with the greatest economic importance (Table 4.1). Both the fall chinook and the steelhead trout spawn in the Hanford reach (from the reactor area to Umatilla and McNary Dam), and these populations may potentially accumulate radionuclides during their freshwater rearing period. The various life stages of salmon and steelhead trout and the time when each is present in the Hanford reach is shown in Figure 4.5.

The relative contribution of upper river bright stocks to fall chinook salmon runs in the Columbia River increased from about 24% of the total in the early 1980s to 50% to 60% of the total by 1988 (Dauble and Watson 1990).

FIGURE 4.5. Expected Occurrence of Salmonids in Hanford Reach of Columbia River

The progressive damming of the Columbia River, beginning with the construction of Bonneville Dam in the lower reaches, has resulted in a concurrent reduction in the runs of anadromous salmonids. This reduction is attributed to a combination of factors, including blockage by the dams during adult migration, destruction of spawning habitat, and loss of downstream migrants as they passed through the turbines or were increasingly preyed upon by the large population of predator species that thrive in the lentic habitats created by the dams. The destruction of other mainstream Columbia River spawning grounds by dams has increased the relative importance of the Hanford reach for spawning (Watson 1970, 1973). Dauble and Watson (1990) present data illustrating the fluctuation in the numbers of salmon redds (nests for salmon eggs in gravel) observed in the Hanford reach from 1944 through 1988; these varied from nearly zero in 1956 to about 8000 in 1989 (Figure 4.6).

FIGURE 4.6. Hanford Reach Salmon Redd (Nest) Counts (Dauble and Watson 1990)

The upper estimates of steelhead spawning population in the Hanford reach based on dam counts for the years 1962 to 1971 were about 10,000 fish. The estimated annual sport catch for the period 1963 to 1968 in the reach of the river from Ringold to the mouth of the Snake River was approximately 2700 fish (Watson 1973).

The American shad, another anadromous species, may also spawn in the Hanford reach. The upstream range of the shad has been increasing since 1956 when fewer than ten adult shad ascended McNary Dam. Since then, the number ascending Priest Rapids Dam, immediately upstream from Hanford, has risen to many thousands each year and the young-of-the-year have been collected in the Hanford reach. Unlike salmonides, the shad is not dependent on specific current and bottom conditions for spawning and has apparently found favorable conditions for reproduction throughout much of the Columbia and Snake Rivers.

4.6.2 Resident Species

Resident fish species are ecologically important to the structure and function of the Columbia River ecosystem. The resident fish population at Hanford is diverse, yet characteristic of many northwest rivers. Resident fish of importance to sport fishermen and Native Americans include the mountain whitefish, white sturgeon, smallmouth bass, crappie, channel catfish, walleye, and yellow perch. Large populations of rough fish including carp, shiners, suckers, and squawfish are also present. A large database exists on radionuclide concentrations in several resident fish species, dating from the 1940s to the present time (Davis, Watson, and Palmiter 1956; Watson et al. 1970).

The range of the mountain whitefish includes southwestern Canada and the northwestern United States. Whitefish are a popular sport fish and may compete with other desirable species, including trout and juvenile salmon, for food and space. Whitefish are present year-round in the Hanford reach and feed primarily near the bottom on aquatic insect larvae, including caddisflies (Trichoptera), midges and black flies (Diptera), mayflies (Ephemeroptera), and stoneflies (Plecoptera). Additionally, they sometimes prey on fish eggs (including their own), fish larvae, and small fish and may take eggs of salmon

or trout. Whitefish containing measurable body burdens of Hanford-related radionuclides have been collected in the lower reaches of the Yakima River below the Hanford Site and near Priest Rapids Dam above the Site. This provides evidence that the resident population in the Hanford reach migrates to these locations (Watson and Davis 1957).

White sturgeon are a long-lived species that reside year-round in the Hanford reach. They are primarily bottom feeders and can take up radionuclides both by ingesting contaminated sediments and via the aquatic food chain. Concentrations of various radionuclides were determined for white sturgeon tissues during radiological studies conducted in 1966 and 1967 (Watson et al. 1970). However, no attempt was made to determine relationships between fish age and radionuclide concentration in carcass or muscle tissue.

Because sturgeon cannot move upstream through fish ladders, their movements since 1960 have been restricted to the area bounded by Priest Rapids, McNary, and Ice Harbor dams. However, seasonal movement throughout the McNary Pool is known to occur (Haynes 1978), and migration range could have been greater for sturgeon that existed before construction of McNary and Priest Rapids dams. Thus, even for sturgeon captured near former production reactors, exposure may not be directly related to radionuclide concentrations present at that location.

Large-scale suckers and carp are numerous in the Hanford reach and have feeding habits conducive to accumulating contaminants. They, like the sturgeon, ingest large quantities of fine detritus and periphyton, both of which have large surface-to-volume ratios and thus adsorb high levels of contaminants. However, there is no evidence that these species contain elevated tissue burdens of radionuclides (Jaquish and Bryce 1989).

There is extensive literature concerning radionuclide burdens of species consumed by man. They include data on zinc-65 levels in man from consuming whitefish (Foster and Honstead 1967) and zinc-65 concentrations in oysters from Willapa Bay, Washington (Seymour 1966). This literature is discussed in Section 7.3. Some of this information is used in Section 10.

5.0 REACTOR OPERATIONS AND EFFLUENT WATER COMPOSITION

The radionuclide composition and activity level of the cooling water discharged to the river varied considerably as the result of several factors including the number of reactors and their power levels, seasonal changes in the parent elements in raw river water (i.e., the elements activated as they passed through the reactor core), the chemicals used in water treatment, the corrosion rates of reactor piping and fuel element cladding, the occasional purging of radioactive film from reactor components, and the length of time the effluent was retained in the basins before discharge. Another factor was radionuclide releases from periodic fuel element ruptures (slug ruptures). The wide variations in these factors, together with the hydrologic variables of the Columbia River and dam construction, produced a complex variation in the combination of river water and reactor effluent during the years of reactor operation. The concentrations and distributions of radionuclides in the river were never constant, in either time or location, throughout the operational years.

The withdrawal and processing of raw river water involved filtration and chemical additives to provide the desired quality for reactor cooling water. Emergency operating procedures were also part of the water system design. Cooling water released from the reactors passed through a retention basin and spillway system to the outfall lines where the water was discharged to the river. An example design of a reactor water system is shown in Figure 5.1 (Ballinger and Hall 1991).

5.1 TIMETABLE OF REACTOR OPERATIONS

Reactor operating periods are shown in Figure 5.2. The 100-B reactor (RM 384) was the first on-line in September 1944, followed by 100-D (RM 377.6) in December of the same year. The 100-F reactor (RM 369) came on-line in February 1945. These three reactors contributed all of the radioactivity discharged to the river until about November 1949, when 100-H (RM 372.5) came on-line. In October 1950, 100-DR (RM 377.6) came on-line, followed by 100-C (RM 383.6) in November 1952. The last of the once-through-cooled reactors,

S9203065.3

FIGURE 5.1. 100-D Area Water System (Ballinger and Hall 1991)

FIGURE 5.2. Operating Periods for the Once-Through-Cooled Hanford Production Reactors (based on Foster 1972)

100-KW (RM 381.8) and 100-KE (RM 381.4), came on-line in January and April 1955, respectively. Because design power levels for older reactors were substantially increased and design power levels were high for C and K reactors, peak plutonium production and discharge of maximum amounts of radioactive effluent did not begin until about 1958.

In 1963, President Johnson ordered a large-scale reduction in plutonium production, and the Hanford reactors were scheduled for retirement. Between 1964 and 1970, all eight once-through-cooled reactors were taken off-line permanently (Figure 5.2). The first reactor to be shut down was 100-DR on December 30, 1964. Its closure was followed in mid-1965 by the closures of 100-H and 100-F. The remaining reactors were all down temporarily during July and August 1966 because of a labor strike on July 8. In June 1967, 100-D was taken off-line, followed by 100-B in 1968, 100-C in April 1969, and 100-KW in February 1970. The last of the eight once-through-cooled reactors, 100-KE, was shut down permanently in January 1971 (Ballinger and Hall 1991).

5.2 ELEMENTS IN COLUMBIA RIVER WATER

During the early 1950s, scientists at the Hanford Site recognized that the isotopic composition of effluent water was not constant, but displayed a regular seasonal fluctuation and irregular daily fluctuations as well as the expected differences between reactors (Honstead 1954). During the early 1960s, studies at the Hanford Site investigated the presence of natural elements in the river water to determine the relative concentrations and their sources. These elements were of interest because of their activation during the passage of cooling water through the reactor pile. The elements of interest were sodium, copper, phosphorus, sulfur, manganese, arsenic, uranium, lanthanum, iron, cobalt, zinc, and scandium.

The reports of this work concluded that concentrations of elements depended on the relative volumes of water supplied by two separate drainage basins (Silker 1963). One basin in Canada includes the headwaters of the Columbia River and the Kootenai River basin, and the other is the combined drainage of the Pend Oreille and Spokane River watersheds in the United States. Both sources drain about 30,000 square miles and supply approximately equal amounts of water to the Columbia River until mid-May, when the Columbia and Kootenai Rivers crest to contribute about 80% of the Columbia River flow at Grand Coulee Dam.

The following conclusions were reached regarding the relationship between the two drainage basin runoffs and the elemental concentrations in the raw river water: zinc and cobalt appeared to originate primarily in the Canadian area of the basin, and the changes in concentration closely parallel those in the upper Columbia River discharge hydrograph. Concentrations of sodium, uranium, sulfur, and phosphorus are lower in the Canadian area, causing a relative decrease in concentrations of these elements during the peak flow period (Silker 1963).

Concentrations of manganese, copper, arsenic, and uranium were from two to five times higher in Spokane River water than in the upstream segment of the Columbia River (Silker 1963). The increased concentration of these elements tends to correspond to peak flows from the Spokane River in late April.

Lanthanum exhibited a similar late-April trend. Iron concentrations were variable and scandium was consistently low.

5.3 CHEMICAL ADDITIVES

Certain radionuclides in the reactor effluent originated from neutron activation of chemicals added to the process water. As the water was withdrawn from the river, water quality variables (e.g., pH, turbidity, temperature, chemistry, bacteriology) were carefully monitored to determine the amounts of chemical compounds to be added. Because the natural pH of the river water is in the range of 8.0 to 8.6, sulfuric acid (H_2SO_4) was added in large amounts to bring the water to a pH of 7.0 that was desired for cooling water. From 3 to 10 parts per million (ppm) of alum [$\text{Al}_2(\text{SO}_4)_3 \cdot 18\text{H}_2\text{O}$] were added as a coagulant as soon as water was withdrawn from the river to chemically reduce certain elements that were present in the water as solids and colloids, such as arsenic, scandium, and uranium. Approximately 2 ppm sodium dichromate ($\text{Na}_2\text{Cr}_2\text{O}_7$) were added shortly before the water entered the reactor to inhibit corrosion of the aluminum tubing and fuel element jackets. An organic filter aid, Separan, was added occasionally in very small amounts just prior to filtration (Hall and Jerman 1960; Perkins 1961).

Two reports offered some estimates as to the percentage of radionuclides the various additives contributed to the effluent. Hall and Jerman (1960) concluded that many of the radionuclides found in reactor effluent water had their primary source in the river except for chromium-51, which was supplied by the sodium dichromate. This chemical also contributed much of the elemental sodium. Hall and Jerman further concluded that as much as 25% of the phosphorus-32 may have resulted from activation of the sulfuric acid.

Perkins (1961) concluded that nearly 100% of the chromium-51 and 25% of the sodium-24 were neutron-activation products of sodium dichromate. At least 40% of the phosphorus-32 was produced from the sulfur in the sulfuric acid and the alum. According to Perkins, impurities in the alum also produced 60% of the gallium-72, 50% of the samarium-153, and 25% of the iron-59. The contribution to all other radionuclides from neutron activation of chemical additives was less than 10%.

5.4 PURGES

To dislodge and remove a radioactive film that built up on the surface of fuel elements and process tubing, purges were run on the average of about two to three times per month. These purges were necessary because this film would occasionally become thick enough to partially restrict the flow of water through the tubing (Hall and Jerman 1960). During the purging process, a slurry of diatomaceous earth and process water was run through the tubes in the belief that the abrasive action of the slurry would dislodge the film.

Reports on the purging process written during the 1950s identified several radionuclides that were elevated by the diatomaceous earth slurry process. A report by Healy (1952) specifically mentions copper-64, iron-59, and phosphorus-32 as being increased by the purging process. This finding was made by comparing the normal activity levels in cooling water entering the retention basin with those levels following purging. A later report by Koop (1957a) identified phosphorus-32 and zinc-65 concentrations being increased in the river because of purging. According to Koop, the concern regarding these two radionuclides was their uptake in Columbia River whitefish flesh, especially during the warm-water season of the river. This season extended from about the first of July to the first of November.

Based on tests conducted in 1955, a report by Koop (1956) recommended control measures for purging operations. These included a time limitation of one hour per purge, a maximum of 25 ppm of diatomaceous earth (100 ppm had been previously used), no more than one purge per 48-hour period (all reactors combined), and purges during reactor operations only on those days when the river temperature was less than 15°C. Purging while a reactor was operating, as opposed to being shut down, produced several times the activity level (net difference) in the effluent, based on analysis of samples from the retention basin (Healy 1952).

Another purging process involved the use of a chemical cleaner, Turco 4306-B, to remove radioactive scale from the surfaces of rear face reactor piping. The purpose of using this cleaner was to lower rates of radiation exposure for workers near the rear face of a reactor. Turco 4306-B, which is a chelating and reducing agent and a strong acid, was tested in September

1957; the results were reported by Koop (1957b). The cleaner was tested during reactor outage time using a concentration of 6 ounces of Turco per gallon of cooling water, heated to about 85°C, and pumped through reactor tubing and down the rear face of the reactor to the storage basin. The flow rate used was about 5% of normal reactor operation flow. Disposal of the cleaning solution was handled differently than the other purging process. The solution was transferred from the storage basin to the empty west side retention basin, which was about half full when cleaning was completed. The coolant flow was then switched to the east basin and reactor operation resumed. Three days later, the full effluent flow was switched back to the west basin, which filled in about an hour, at which time the contents began flowing into the river. The normal reactor flow rate was maintained in the west basin for 4 hours.

Koop (1957b) stated that the concentrations of radionuclides from Turco purging were not significantly different from the normal reactor effluent samples, although two radionuclides, iron-59 and zinc-65, were found in higher concentrations in basin inlet samples, which contained cleaning solution, than in operating effluent. In the same report, Koop later stated that the radionuclides of concern, with respect to potential pollution, were iron-59, zinc-65, and neptunium-239. The recommended disposal was to release the spent cleaning solution to the river at as slow a rate as was practical.

Another study, reported by Perkins (1959), presented results of testing the efficiency of the Turco cleaning solution on process tubing and associated fuel element jackets. The report presents a detailed discussion of Turco purging effects within the reactor system, but does not provide much information on releases to the river.

5.5 ACCIDENTAL RELEASES

Accidental releases of radionuclides to the river occurred from retention basin leakage (shoreline releases); fuel element ruptures; and in one instance, the failure of a 300 Area waste pond dike. The shoreline releases occurred as river bank springs that were fed by retention basin leakage into the underlying soils. Fuel element ruptures were intermittent events that

began in 1948 and increased in frequency as more reactors were added and power levels increased. The failure of the 300 Area waste pond dike was a single event occurring in October 1948.

5.5.1 Shoreline Releases

By the late 1950s, the condition of the retention basins for B, C, D, and F reactors had raised concerns regarding the leakage of the basins contributing to shoreline releases to the river (Koop, McCormack, and Hall 1958). The concerns were the contamination of drinking water for workers at downstream reactors (more specifically, at reactors downstream of B, C, and D reactors--no reactor was downstream of F reactor) and the radionuclide uptake by aquatic life in the river. The bottom organisms along the shoreline area were food for whitefish and waterfowl, both of which were consumed by humans.

The report by Koop, McCormack, and Hall (1958) mentioned that the deterioration of the basins involved both surface and bottom leakage but that the release rates were not known. Although the shoreline releases were considered to be significant, some decontamination was assumed from filtration and ion exchange within the soil column. The authors also assumed that the soil between the basins and the river had limited capacity and that the activity level of the water emerging from the shoreline would eventually approach that of the basin effluent. According to the authors, this trend had been confirmed by the increase in the amount of radionuclides in ground water flowing into the 181-B forebay and approximately a twofold increase since 1955 in the radionuclide concentrations from river bank hot springs at the 100-F Area.

5.5.2 Fuel Element Ruptures

During reactor operations, the cladding (jacket) of individual fuel elements can fail and release fission products and uranium to the cooling water. McCormack and Schwendiman (1959) discuss two principal rupture types: 1) a severe rupture, usually a side rupture or fragmented element, and 2) a split or end-cap failure.

The first two fuel element ruptures occurred in 1948 followed by three more in 1950 (DeNeal 1965). Beginning in 1951, the number of ruptures per

year increased significantly as shown in Table 5.1. Jerman, Koop, and Owen (1965) stated that severe ruptures resulted in a weight loss of about 150 grams per fuel element, while the other types of ruptures resulted in a weight loss of about 9 grams per fuel element. The authors did not state whether the weights included other materials beside uranium. They estimated that the average weight loss was 12 kilograms per year for the years from 1955 through 1958.

Two fission products released during ruptures were strontium-89,90 and iodine-131 (McCormack and Schwendiman 1959; Jerman, Koop, and Owen 1965).

TABLE 5.1. Fuel and Target Element Failures Removed from the Reactors (DeNeal 1965)

<u>Calendar Year</u>	<u>Total Failures All Types</u>
1948	2
1949	0
1950	3
1951	115 ^(a)
1952	142
1953	93
1954	211
1955	242
1956	191
1957	201
1958	174
1959	71
1960	130
1961	86
1962	95
1963	68
1964	97
1965 ^(b)	49
Total	1970

(a) Includes 13 lithium target elements.

(b) Through May 1965. Includes 17 thoria target elements.

McCormack and Schwendiman (1959) stated that ruptures contributed about 20% of the strontium-89,90 in the Columbia River at Pasco and about 4% of the gross fission product activity. Jerman, Koop, and Owen (1965) considered iodine-131 to be the most significant radionuclide. They estimated that 2400 grams of irradiated uranium containing 920 curies of iodine-131 were lost from the 97 ruptures in 1964. For comparison, they stated that about 500 kilograms of natural uranium in the cooling water passed through the eight reactors, creating about 800 curies of iodine-131.

As of 1965, the most severe rupture occurred on May 12, 1963 (Hall 1963; Jerman, Koop, and Owen 1965). The rupture occurred in one of the process tubes of the 100-KE reactor and involved a zircalloy-clad experimental fuel element. About 450 grams of uranium were lost, indicating a release of about 170 curies (Jerman, Koop, and Owen 1965).

5.5.3 300 Area Waste Pond Dike

A description of this pond, its associated problems, and the dike failure that occurred on October 25, 1948, is presented in a report by Singlevich and Paas (1949). The following description of the pond and details of the dike failure are summarized from that report.

The pond covered an area of 490,000 square feet and was about 5 feet deep. The walls were constructed of crushed rock and earth; the bottom was earthen, through which waste solution continuously infiltrated. Estimates indicated that about 2 to 3 pounds of uranium were discharged daily into the pond from the 300 Area. However, this estimate neglected sources of uranium from other buildings. At the time of the report, exact quantities of uranium discharged into the pond were not known.

Following the failure of a weak point in the northwest corner of the dike, approximately 14.5 million gallons of waste solution were discharged into the river between 2:30 PM and 4:00 PM on October 25. The waste traveled about 1000 to 1500 feet over the ground to the river. Rough calculations indicated that about 12 to 60 pounds of uranium were included in the waste solution. River water samples were taken from the 300 Area down to Portland, Oregon, at strategic locations (500 samples). The maximum single result was

2280 disintegrations per minute per liter (dpm/L) of alpha activity about 30 yards downstream of the 300 Area. Trace quantities of uranium were found about 1 mile above Richland and normal alpha activity was detected below Richland and at Portland.

5.6 BASIN RETENTION TIME

The purpose of the retention basins was to hold reactor effluent until many of the shorter-lived radionuclides had decayed substantially before discharging effluent to the river. Retention time depended on the flow rate through the reactors and varied from one reactor to the next. In 1945, the optimum time was reported as 6 hours (Parker 1945).

By 1960, the retention time varied from 30 minutes to 3 hours, which reduced the activity levels in the effluent released to the river by a factor of two to three, but was not long enough to reduce the activity of those radionuclides of major interest by a significant amount (Hall and Jerman 1960).

A study of basin retention times was conducted in 1953 to provide data for calculating decay correction factors to determine more accurate values for beta particle activity density in the water leaving the basins. The results were reported by Soldat and Quimby (1953). A summary of basin retention times based on flow rate data for 100-B, 100-D, 100-DR, 100-H, and 100-F is given by Pilcher and Norton (1953).

5.7 EFFLUENT DISCHARGE TO RIVER

The effluent was discharged into the river from each reactor by gravity flow through a pipe 42 to 60 inches in diameter that was placed along the river bottom and ran from an open discharge structure (spillway) at the bank. The pipe or outfall line usually extended to the channel center and was buried for most of its length a few feet below the riverbed (Honstead, Healy, and Paas 1951). Thus, the outfall lines discharged the effluent at a point 300 to 700 feet from the reactor shore where the velocity of flow was faster, to produce as much near-field diffusion as possible.

There were two exceptions to this typical effluent discharge design. At the 100-D reactor, the river channel is divided by a mid-channel island, with the higher velocities occurring on the opposite side of the island. The outfall line for the combined 100-D and 100-DR effluents was extended along the near-channel bottom, over the island, and into the far channel to take advantage of the faster flow velocities. The length of outfall line crossing the island was perforated with 1/2-inch holes every 20 feet to prevent air pockets in the line. This design allowed continuous venting of effluent water onto the island surface. The other exception was at 100-F, where the original line failed and was replaced in 1946 by a new line that extended only 300 feet from the retention basin (Honstead, Healy, and Paas 1951). As a result, effluent was discharged about 200 to 250 feet from the shore into slower flow velocities. The resulting effluent plume hugged the reactor shoreline and produced elevated activity levels for some distance downstream.

Problems arose during seasons of high river flow because the hydraulic head differential between the discharge structure and the water surface elevation decreased. This decrease would lower the discharge capacity of the line and cause the effluent basin to overflow into a bypass spillway that released into the river at the shoreline. Diversion of basin inflow to the bypass spillway was also common during power peaking and concurrent high flow rates through the reactors (Honstead 1954; Hall and Jerman 1960).

5.8 ACTIVITY LEVELS AND VARIABILITY OF RELEASES

According to a report by Parker (1954), the contamination of the Columbia River by reactor effluent had been studied "with increasing intensity since the start of the Hanford operations." Parker stated that, at the time of the report, about 8000 Ci/day of radioactive material were released to the river. The radiochemical composition was variable and influenced by seasonal turbidity changes and modifications of water treatment. A brief discussion in Parker's report seems to indicate that calculations of river concentrations had been attempted, but that the results were not as reliable as desired because 100% material balance was impossible. No specific details were provided on the calculation procedure.

A report by Hall and Jerman (1960) includes some tabulated data of the activity levels in effluent releases to the river. For the years 1957 through 1959, the tabulated data for each of the eight reactors includes the monthly average cooling water discharge, monthly average total beta releases (Ci/day), and monthly average releases (Ci/day) for phosphorus-32, arsenic-76, zinc-65, chromium-51, and neptunium-239. A monthly average release rate for total beta is shown in Figure 5.3. The cause of the unusually high monthly release rates

FIGURE 5.3. Monthly Average Radioactivity Release Rate in Ci/day for Reactor Effluent at the Point of Release to the Columbia River (Hall and Jerman 1960)

that occurred in 1957 (Figure 5.3) is not known at this time; however, there are several possibilities. One of the possibilities was the initial use of Separan, which improved the efficiency of filter-bed backflushing, the inflow of elemental manganese from the Spokane River watershed, or the disturbance of upstream riverbed sediments at the Priest Rapids Dam construction site. The most likely cause is thought to be the activation of elemental manganese from the Spokane River system, producing manganese-56. According to Hall, there was a significantly higher level of elemental manganese in the Columbia River at that time.^(a)

A report by Bogan (1956) summarizes the results of several earlier studies and provides some insight into the difficulty in correlating reactor power levels (total beta activity produced), river flow, and measured downstream beta activity. Bogan states that the results were both inconsistent and inconclusive. In some instances, higher values were found in the river than would be expected based on input calculations and, in another instance, just the opposite was found.

(a) Personal communication to W. H. Walters from R. B. Hall, March 1992.

6.0 COLUMBIA RIVER MONITORING AT HANFORD

The potential for the contamination of the Columbia River as a result of Hanford operations was recognized even before the construction of the first plutonium production reactors. Initial monitoring began at the Hanford Site in 1945 and was conducted by site contractors. Since 1945, the emphasis and detail of the monitoring programs have changed, as has the reporting frequency. Also, beginning about 1950, offsite (e.g., federal and state) agencies conducted monitoring programs and various studies at river locations downstream of Hanford, including the coastal areas. This section presents an overview of the river monitoring and topical studies conducted by Hanford contractors. Monitoring and topical studies conducted by offsite agencies are discussed in Section 9.

The history of Columbia River monitoring is divided into three periods: 1944 through 1957, 1958 through 1971, and 1972 through 1990. These particular periods correspond reasonably well with 1) significant developments in analytical techniques that greatly enhanced the type of information generated by the monitoring programs, 2) programmatic modifications that resulted in changes in the rationale for and purposes of the programs, and 3) operational changes that resulted in significant reductions in the contaminant loading to the river as a result of Hanford operations.

6.1 1944 THROUGH 1957

The first studies of the effects of radioactivity on the Columbia River ecosystem were proposed during the summer of 1943, before the beginning of construction of the original reactors along the river. The emphasis of the early studies was on the accumulation of radioactivity by aquatic organisms, primarily fish. Routine Columbia River monitoring was initiated during 1945, shortly after the startup of the original three plutonium production reactors along the river at Hanford.

The primary objective of the original surveillance efforts was only the detection of reactor-created radioisotopes, not their quantification. Releases to the river were controlled by self-imposed dose limits because

there were no standards or concentration guides for radionuclide discharges to the river (Parker 1952). Dose rate measurements were made at several locations, including the point of release, sites just downstream of the reactor outfalls, and points as far downriver as Pasco. Shortly after the startup of the reactors, the need not only to detect but to measure the quantity of radionuclides in the river was recognized and routine sampling was started.

Locations for routine Columbia River water sampling from 1944 through 1957 are shown in Table 6.1. Initially, the program included sampling locations at the 100-B, 100-D, and 100-F Areas, the Hanford townsite, the 300 Area, and Richland. This scheme was quickly extended as far downstream as Pasco. During the late 1940s through 1957, the Columbia River monitoring program was expanded significantly, providing a large amount of data (primarily gross beta activities) from several locations. Plate 1 illustrates the various water sampling locations used at various times during the years 1944 through 1957. Grab samples were collected at various times from upstream of the Hanford Site to the Portland, Oregon/Vancouver, Washington, area.

In addition to the routine Columbia River water monitoring programs, comprehensive surveys were conducted to determine the plume dispersion of reactor effluents during the 1950s and early 1960s. Special studies, including extensive sampling, were conducted to determine the horizontal and vertical mixing characteristics of the Columbia River along the Hanford reach to better understand the fate of reactor-generated contaminants. These studies are discussed in Section 8 of this report.

Analytical capabilities were limited during the early years. Laboratory techniques were newly developed and improvements were constantly being made throughout the late 1940s and 1950s. Until the advent of gamma energy spectroscopy in 1957-1958, total alpha and total beta measurements were essentially the only data available. Five-hundred-milliliter (mL) samples were evaporated, leaving a sample residue that was then counted for total beta activity with a thin-window counter. Alpha activity was determined using a standard alpha counter. Detection levels were reported as being approximately 5×10^{-5} microcuries per liter ($\mu\text{Ci/L}$) beta and 2 dpm/L alpha activity.

TABLE 6.1. Summary of Routine Locations for Sampling Columbia River Water, 1945-1957

Location ^(a)	1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957
Columbia River													
Wills' Ranch	-(b)	-	-	-	X	X	X	X	X	X	X	X	X
Above 100-B	X	X	X	X	X	X	-	-	-	-	-	-	-
181-B	X	X	X	X	X	X	X	X	X	X	X	X	X
181-C	-	-	-	-	-	-	-	-	X	X	X	X	X
Allard Station	-	-	-	-	-	X	X	X	X	X	X	X	-
181 KW	-	-	-	-	-	-	-	-	-	-	X	X	X
181 KE	-	-	-	-	-	-	-	-	-	-	X	X	X
181-D	X	X	X	X	X	X	X	X	X	X	X	X	X
181-H	-	-	-	-	X	X	X	X	X	X	X	X	X
Below 100-H	-	-	-	-	-	X	X	X	X	X	X	X	X
181-F	X	X	X	X	X	X	X	X	X	X	X	X	X
Below 100-F	-	-	-	-	-	X	X	X	X	X	X	X	X
Forster's Ranch	-	-	-	-	X	X	X	-	-	-	-	-	-
Hanford S. Bank	X	X	X	X	X	X	X	X	X	X	X	X	X
Hanford Middle	-	X	X	X	X	X	X	X	X	X	-	-	-
Hanford N. Bank	X	X	X	X	X	X	X	X	X	X	-	-	-
300 Area	X	X	X	X	X	X	X	X	X	X	X	X	X
Byers Landing	-	-	-	-	-	-	-	-	X	X	X	X	X
Richland (Dock)	X	X	X	X	X	X	X	X	X	X	X	X	X
Kennewick Highlands	-	-	-	-	-	X	X	X	X	X	X	X	X
Pasco Pump Plant	-	-	-	-	-	-	-	-	-	X	X	X	X
Pasco Bridge													
Pasco	X	X	X	X	X	X	X	X	X	X	X	X	X
Kennewick	-	-	-	-	X	X	X	X	X	X	X	X	-
Sacajawea Park	-	-	-	-	-	-	X	X	X	X	X	X	X
McNary Dam - Below	-	-	-	-	-	-	X	X	X	X	X	X	X
McNary Pool	-	-	-	-	-	-	-	-	-	X	X	-	-
Paterson, WA	-	-	-	-	-	-	X	X	X	X	X	X	X
Arlington, OR	-	-	-	-	-	-	-	-	X	-	X	-	X
Mary Hill Ferry, OR	-	-	-	-	-	-	-	-	X	-	-	-	X
Celilo Falls, OR	-	-	-	-	-	-	-	-	X	-	-	-	X
The Dalles, OR	-	-	-	-	-	-	-	-	X	-	X	-	X
Hood River, OR	-	-	-	-	-	-	-	-	X	-	-	-	X
Cascade Locks, OR	-	-	-	-	-	-	-	-	X	-	-	-	-
Troutdale, OR	-	-	-	-	-	-	-	-	X	-	-	-	X
Stevenson, WA	-	-	-	-	-	-	-	X	X	-	-	-	X
Bonneville Dam, OR/WA	-	-	-	-	-	X	-	X	X	-	-	-	X
Portland, OR	-	-	-	-	-	-	-	X	X	-	X	-	-
Vancouver, WA	-	-	-	-	-	-	-	-	-	-	X	-	X
Yakima River													
Prosser	X	X	X	X	X	-	-	-	-	X	X	X	X
Horn	-	-	-	X	X	-	-	-	-	X	X	X	X
Mouth	-	-	-	X	X	-	X	X	X	X	X	X	X
Snake River													
Mouth	-	-	-	-	-	-	X	X	X	X	X	X	X

(a) For an idea of where water was sampled, see Plate 1 in a pocket in the back of this report.
 (b) - indicates no samples collected.

Monitoring of Columbia River sediment, or mud as it was referred to in the early days, was initiated in 1948. Sediment samples were routinely collected at various locations through 1957, as shown in Table 6.2. Sediment samples were collected from two points at each sampling location. The first

TABLE 6.2. Summary of Routine Locations for Sampling Columbia River Sediment, 1948-1957

Location ^(a)	Year Sampled									
	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957
Columbia River										
Wills' Ranch	x	x	x	x	x	x	x	x	x	x
Allard Station	x	x	x	x	x	x	x	x	-	-
100-H Area	x	x	x	x	x	x	x	x	x	x
Below 100-F	x	x	x	x	x	x	x	x	x	x
Hanford Ferry	x	x	x	x	x	x	x	x	x	x
300 Area	x	x	x	x	x	x	x	x	x	x
Byers Landing	-(b)	-	-	-	x	x	x	x	x	x
Richland (Dock)	x	x	x	x	x	x	x	x	x	x
Kennewick Highlands	-	-	x	x	x	x	x	x	x	x
Clover Island	-	-	-	-	x	-	-	-	-	-
Pasco Bridge										
Pasco	x	x	x	x	x	x	-	-	-	-
Kennewick	-	x	x	x	x	x	x	x	x	x
Sacajawea Park	-	-	-	x	x	x	x	x	x	x
McNary Dam	-	-	-	x	x	x	x	-	-	-
McNary Cold Springs										
South	-	-	-	-	-	x	-	-	-	-
Middle	-	-	-	-	-	x	-	-	-	-
North	-	-	-	-	-	x	-	-	-	-
McNary Pool	-	-	-	-	-	-	x	-	-	-
McNary Below Dam	-	-	-	-	-	-	x	x	x	-
Paterson	-	-	-	x	x	x	x	x	x	x
Yakima River										
Prosser	-	-	-	-	-	-	x	x	x	x
Horn	-	-	-	-	-	-	x	x	x	x
Snake River										
Mouth	-	-	-	x	x	x	x	x	x	x

(a) For an idea of where sediment was sampled, see Plate 1.

(b) - indicates no samples collected.

was on shore, just above the water level, and the second was from the river bottom, approximately 5 feet into the river. Sediment sampling locations during the period 1944 through 1957 are presented in Plate 1 (in a pocket at the back of this report).

During this period, direct surveys of the sediment provided raw data indicative of contamination levels present. In addition, gross beta results were typically reported for the sediment samples collected and analyzed by a laboratory.

As discussed earlier, before and during the early days of operations at Hanford, there was a great deal of interest and concern about the impact of discharging radioactive material into the Columbia River. Numerous special

studies, both laboratory and field-oriented, were undertaken during the early days of Hanford operations. Columbia River fish, primarily whitefish, have been routinely sampled at Hanford since 1950. Whitefish were chosen for routine sampling because they were available year-round and had some of the highest concentration factors.^(a) Other species that have been monitored include trout, salmon, sucker, carp, bass, crappie, perch, squawfish, chisel-mouth, chub, catfish, and sturgeon (e.g., Davis, Watson, and Palmiter 1956). Sampling locations were generally defined by area rather than specific site because of the mobility of the fish. These areas typically included sites upstream of operating facilities, sites directly downstream of operating reactors, and sites downstream of the Hanford Site at areas popular with the local fisherman.

Routine monitoring results from various shellfish, primarily oysters, obtained from the marine environment near the mouth of the Columbia River are documented from 1953 through 1978. Willapa Bay, the major oyster-rearing area nearest the mouth of the Columbia River, was chosen as the primary sample collection site for oysters. As in the case of water samples, the analysis of fish and shellfish samples from 1944 through 1957 consisted of gross beta measurements.

6.2 1958 THROUGH 1971

By 1957, the Columbia River monitoring activities had developed into an extensive program of sampling river water and sediments at a number of locations downstream of Priest Rapids Dam (Wilson 1962). In addition, samples of various aquatic biota in the river were routinely taken, as were shellfish from coastal regions near the mouth of the river. During the late 1950s, the purpose of the Columbia River monitoring program changed from detection of Hanford-derived contaminants to quantification and dose evaluation. As a result, the monitoring program was modified to generate the optimum amount and type of river data for evaluating the potential radiation doses received by the public living near and using the Columbia River.

(a) J. P. Corley, personal communication to W. H. Walters, March 1992.

Routine river water sampling during 1958 through 1971 provided an indication of the burden of radioactive materials added to the river as a result of Hanford operations. Data obtained through the program were used in estimating doses that might have been received through the surface-water pathway. In addition, the sampling program provided insight on seasonal changes and long-term trends in the concentrations of radionuclides resulting, in part, from changes in operating procedures and practices. Routine monitoring also provided a mechanism for detecting and evaluating the effects of abnormal releases, such as slug ruptures and system purges.

The Columbia River monitoring program established in the early 1960s is the predecessor of today's Surface Environmental Surveillance Project (SESP) surface-water monitoring task. Sampling sites (see Plate 1) were selected to provide specific data. Upstream locations were established to determine background concentrations so that the contribution of Hanford effluents could be distinguished from that of fallout from weapons tests. The first downstream points of water withdrawal for use as a public drinking water supply were established as routine sample locations as well. Points farther downstream, such as Bonneville Dam and Vancouver, served as indicators of the amounts of radioactivity being discharged via the river into the Pacific Ocean or taken up within the river environment. Table 6.3 lists the Columbia River water sampling locations that were routinely monitored as part of the monitoring program during the years 1958 through 1971.

Sampling methods and equipment changed somewhat over the years. Grab samples were standard during the early years. Composite sampling was initiated later to provide better estimates of the average radionuclide concentrations in the river at a given location over time. In addition, composite sampling provided some assurance that short-term elevated releases were not missed because of the sampling protocol and frequency. Continuous sampling systems (filter/resin) were put into service in later years to increase the sample size sufficiently to detect radionuclides at very low concentrations. With the introduction of gamma energy spectroscopy in 1957, it became practical to obtain isotopic analyses of river water samples on a routine basis.

TABLE 6.3. Summary of Routine Locations for Sampling Columbia River Water, 1958-1971

Location ^(a)	Year Sampled													
	1958 ^(b)	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969 ^(c)	1970	1971
Priest Rapids Dam	-(d)	-	-	-	-	-	-	-	X	X	X	X	-	-
Wills' Ranch	-	X	-	-	-	-	-	-	-	-	-	-	-	-
Vernita Bridge	-	-	-	-	-	-	-	X	X	-	-	-	X	X
Vernita Ferry	-	-	-	-	X	X	X	X	-	-	-	-	-	-
Hanford S. Bank	-	X	X	X	X	X	-	-	-	-	-	-	-	-
Ringold	-	-	-	-	-	-	-	X	X	X	X	-	-	-
300 Area	-	-	-	-	X	X	-	-	-	-	-	-	-	-
Richland Pumphouse	-	-	-	-	-	X	X	X	X	X	X	X	X	X
Richland (Dock)	-	X	-	-	X	-	-	-	-	-	-	-	-	-
Pasco Pump Plant	-	X	X	X	X	X	X	X	-	-	-	-	-	-
Pasco Bridge	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pasco	-	X	-	-	-	-	-	-	-	-	-	-	-	-
Sacajawea Park	-	X	-	-	-	-	-	-	-	-	-	-	-	-
McNary Dam - Below	-	X	-	-	-	-	X	X	X	X	X	X	X	-
Paterson, WA	-	X	-	-	-	-	-	-	-	-	-	-	-	-
The Dalles, OR	-	-	-	-	-	X	X	-	-	-	-	-	-	-
Hood River, OR	-	X	-	-	-	-	-	-	-	-	-	-	-	-
Bonneville Dam, OR/WA	-	-	-	-	-	-	X	X	X	X	X	X	X	X
Vancouver, WA	-	X	X	X	X	X	-	-	-	-	-	-	-	-

(a) For an idea of where water was sampled, see Plate 1.

(b) Specific locations not provided. Assume similar to 1957 because concentrations reported for river "stretches" from the 100 Areas to Portland.

(c) Nonradiological parameters initiated (Vernita, 100-D, 100-F, 300 Area, Richland Pumphouse).

(d) - indicates no samples collected.

Following this improvement in radiation detection capabilities, measurement of specific radionuclides became the norm. Table 6.4 summarizes the radionuclide analyses during 1958 through 1971.

For the 1958 through 1971 time period, samples of Columbia River sediments were collected only in 1959. Routine sampling of the shoreline and surface river bottom sediments was discontinued in 1960, primarily because the data were not useful in direct determination of doses, as sediment was not in the food chain.^(a) Subsequent annual reports likewise were void of sediment sampling data. The sediment sampling locations for 1959, which was the last year of routine monitoring of activity levels in sediment include the following:

(a) Personal communication to W. H. Walters from J. K. Soldat and R. F. Foster, March 1992.

- Wills' Ranch
- Hanford Ferry
- the 300 Area
- Byers Landing
- Richland (Dock)
- the Pasco and Kennewick sides of the Pasco Bridge
- Sacajawea Park
- below McNary Dam
- Paterson.

Special studies were conducted throughout this period, providing some indication of the levels of Hanford-derived radionuclides associated with Columbia River sediments. These studies are summarized in Section 9.5. Core samples were collected from behind McNary Dam on various occasions between 1958 and 1971.

TABLE 6.4. Analyses for River Water Samples, 1958-1971

Year	Analyses								
	Total Alpha	Total Beta	Gamma ^(a) Scan	³ H ^(b)	³² P	⁹⁰ Sr	¹³¹ I	²³⁹ Pu	U
1958	X	X	X	-(c)	X	X	-	-	X
1959	X	X	X	-	X	X	X	-	X
1960	X	X	X	-	X	X	X	-	X
1961	X	X	X	-	X	X	X	-	X
1962	X	X	X	-	X	X	X	-	X
1963	X	X	X	-	X	X	X	-	X
1964	X	X	X	-	X	X	X	-	X
1965	X	X	X	X	X	X	X	-	X
1966	X	X	X	X	X	X	X	-	X
1967	X	X	X	X	X	X	X	-	X
1968	X	X	X	X	X	X	X	-	X
1969	X	X	X	X	X	X	X	-	X
1970	X	X	X	X	X	X	X	X	X
1971	X	X	X	X	X	X	X	X	X

(a) Including primarily sodium-24, barium-140, scandium-46, chromium-51, manganese-56, copper-64, zinc-65, zirconium/niobium-95, iodine-131, cesium-137, and neptunium-239.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: tritium, phosphorus-32, strontium-90, iodine-131, plutonium-239, and uranium.

(c) - indicates not analyzed.

As in the case of water samples, the advent of gamma energy spectroscopy provided the ability to generate radionuclide-specific data for the sediment samples collected. Core sampling investigations also included isotopic analysis.

The collection of fish from the Columbia River and shellfish from coastal areas near the mouth of the river continued to be an integral part of the Columbia River monitoring program from 1958 through 1971. As had been the case during the earlier years, whitefish remained the primary species sampled. Sampling locations were again identified as areas along the river. Samples were typically collected from areas upstream of operating facilities, immediately downstream of the operating reactors, and downstream of Hanford at areas known to be frequented by the local fishermen.

Shellfish, primarily oysters, continued to be routinely collected throughout the years 1958 through 1971. Oysters reared in Willapa Bay were obtained from a commercial market in Portland, Oregon. Sample analysis included gamma scans, which provided radionuclide-specific data, with zinc-65 being of most interest.

6.3 1972 THROUGH 1990

The Columbia River monitoring program, as it evolved during the late 1950s and early 1960s, has continued from 1972 to the present. As sampling and analytical techniques were improved, the program was modified, enhancing the quality and usefulness of the data obtained.

Samples of river water continued to be collected routinely from several locations from 1972 to the present. The primary emphasis of the Columbia River monitoring program since its restructuring in the early 1960s has been the evaluation of the potential dose to those persons using and/or consuming the river water. Following the shutdown of the once-through-cooled production reactors, the concentrations of radionuclides in the river water decreased significantly (Robertson et al. 1973). Levels of radionuclides fell to near detection limits relatively quickly at several locations. As a result, the more distant sample locations were gradually eliminated from the routine river

water sampling network. Routine water sampling locations during the years 1972 through the present are listed in Table 6.5.

As was the case during the earlier periods, grab, composite, and continuous sampling systems were used for the collection of river water samples. Like sample collection methods, analytical procedures have undergone numerous improvements to provide adequate sensitivities. Standard analytical procedures were not sensitive enough to detect the low concentrations of radionuclides present in the river water following the shutdown of the original reactors. Radiological analyses of water samples included gross alpha, gross beta, gamma scan, tritium, strontium-90, technetium-99, iodine-129, plutonium-239,240, and isotopic uranium.

With the shutdown of the original production reactors, direct discharges of large quantities of radionuclides into the river were eliminated. However, radionuclides were known to be present in the ground water beneath the Hanford

TABLE 6.5. Summary of Routine Locations for Sampling Columbia River Water, 1972-1990

Location ^(a)	Year Sampled																		
	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Richland Pumphouse	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Vernita	x	x	x	x	x	x	x	(b)	-	-	-	-	-	-	-	-	-	-	-
Bonneville Dam- River	x	x	x	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Below 181-N	x	x	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100-B Area-River	-	x	x	x	x	x	x	x	x	-	x	x	x	-	-	-	-	-	-
Hanford Powerline Crossing	-	-	x	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
300 Area	-	-	-	-	-	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Priest Rapids- River	-	-	-	-	-	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Hanford Townsite- River	-	-	-	-	-	-	-	-	-	x	x	x	-	-	-	-	-	-	-

(a) For an idea of where sediment was sampled, see Plate 1.

(b) - indicates no samples collected.

Site and, in some cases, to be approaching and entering the river along the Hanford shoreline (Myers, Fix, and Raymond 1977).

The ground-water discharges or river bank springs have been sampled periodically over the years. Documentation of these monitoring activities, however, is not abundant and is typically contained in project and personal files. Springs near the 300 Area retention basin and sewage leaching trenches were routinely sampled and analyzed for various biological, chemical, and radiological parameters. Springs along the 100-N Area resulting from disposal of liquid waste have been and are today monitored routinely (Rokkan 1988).

Springs located in the contaminated ground-water plumes from the 100-N Area, the 200 Area (emerging at the Hanford townsite), and the 300 Area have been sampled routinely since 1984. Special studies conducted in 1982-1983 and 1988 included sampling several contaminated springs along the Hanford reach (McCormack and Carlile 1984; Dirkes 1990). Analyses of spring samples were similar to those performed on the river water samples, and sampling results were consistent with those from nearby ground-water monitoring wells.

The routine collection of Columbia River sediment samples, discontinued in 1960, was reestablished in 1988. Initially, samples were collected from behind Priest Rapids Dam upstream of Hanford and behind McNary Dam downstream of Hanford. Subsequently, sediment sampling locations have been established at the White Bluffs slough, the 100-F Area slough, the Hanford townsite slough, and Richland. The routine sampling reestablished in 1988 consisted of collecting surface sediments using a clamshell-type sampling dredge designed to collect approximately the top 6 inches of sediment material. Sample analyses consisted of gamma scans, strontium-90, isotopic plutonium, and isotopic uranium.

Even before 1988, special studies were conducted at McNary Dam to investigate the content of radionuclides in the sediments and to observe how soon after the closure of the once-through-cooled reactors clean sediments were deposited on top of the radiologically contaminated sediment material (Robertson et al. 1973; Robertson and Fix 1977; Beaseley et al. 1981; Beaseley

and Jennings 1984). In the special studies, core samples were collected from the pool sediments behind McNary Dam (Robertson and Fix 1977).

The collection of various fish species continued into the period from 1972 through 1990. As in the past, whitefish was the primary species sought. The number and diversity of fish samples routinely collected and the associated radionuclide concentrations declined steadily over the years, following the decline of radioactivity in the river as a result of the shutdown of the reactors (Cushing et al. 1981). The primary constituents of concern in the river environment included strontium-90, tritium, and the gamma-emitters.

The collection of samples of oysters from Willapa Bay continued into the years following closure of the once-through-cooled reactors. The routine collection and analysis of oysters from Willapa Bay was discontinued in 1978 (Houston and Blumer 1978). Zinc-65 was still the radionuclide of concern, although its levels diminished following the reactor shutdown, as the remaining zinc-65 decayed.

From 1944 through 1957, reports that presented the results of the environmental monitoring programs (including Columbia River monitoring) were issued at various intervals and under different titles by authors from Hanford Site contractors. Over the years, the frequency of the monitoring reports generally decreased. During the very early days of Hanford operations (January 1945 through August 1946), reports were issued weekly. Semimonthly reports were issued from September 1946 to May 1947. From June 1947 through the end of 1957, monthly environmental monitoring reports were issued; quarterly reports were also prepared from January 1947 through the end of 1957. An annual report was also prepared summarizing all the monitoring results for 1957.

Beginning in 1958, annual environmental monitoring reports were issued, providing improved documentation of the findings of the river monitoring program. Routine environmental monitoring data are available either within these reports or in a data appendix. In addition to the routine environmental monitoring reports, topical reports were issued presenting results of special studies of the river. Such studies were conducted to fill a particular need.

The special studies are discussed in Sections 7 and 9.5. These studies primarily address investigations of river processes, as well as transport and distribution of radioactivity in the Hanford reach and, to a limited extent, downstream of the river mouth. This reporting system remains in place today.

7.0 SUMMARY OF RESULTS OF COLUMBIA RIVER MONITORING BY HANFORD CONTRACTORS

The amount of radioactivity entering the river varied throughout the period from 1944 through 1971 in response to the startup and shutdown of various reactors, the operation of individual reactors at different power levels, reactor closures for maintenance and refueling, water treatment modifications, the frequency and severity of fuel-cladding failures, and other operational features, as discussed in Section 5 of this report. As the number and power levels of the reactors increased, the amount of radioactivity discharged to the river increased. Consequently, radionuclide releases to the Columbia River were highest during the late 1950s through the early 1960s.

An extensive amount of Columbia River monitoring data has been generated through the monitoring programs and activities described in the previous section. Sample results generated as part of the routine surveillance programs have been reported periodically over the years in weekly, semimonthly, monthly, quarterly, and annual reports. To document the results of special investigations and sampling activities, topical reports generally were issued. In some cases, summary reports have contained data previously reported in periodic status reports.

An eight-volume set, A Compilation of Basic Data Relating to the Columbia River, was prepared by several staff members of the Hanford Atomic Products Operation under the direction of R. F. Foster in 1962 (Foster et al. 1962). Much of the raw data generated during the early years of Hanford had been classified, but was made publicly available in 1962 as Volume 8 (under separate title) of this multivolume compilation (Soldat 1962b).

In 1990 an entire book, Aquatic Bioenvironmental Studies: The Hanford Experience 1944-84, was dedicated to the summary of various studies conducted on the Columbia River and related to operations at Hanford (Becker 1990). This publication serves as an excellent reference on the history of Hanford, as well as providing extensive summaries of the data available through years of monitoring and study of the Columbia River ecosystem. While the book focuses primarily on the monitoring of biota and related laboratory studies, much information is also provided on the Hanford-derived contaminants and

their behavior and fate in the river water and sediment. It should be considered a primary source of summary information regarding the surface-water (Columbia River) pathway and may be useful in evaluating the adequacy of monitoring data for the purpose of reconstructing potential doses during the HEDR Project.

The following sections provide brief summaries of the Columbia River monitoring data generated by Hanford contractor monitoring programs (as opposed to those of offsite agencies) during the years of Hanford operations.

7.1 WATER MONITORING DATA

During the early years of Hanford operations, until the advent of gamma energy spectroscopy in 1957, gross radioactivity in Columbia River water was measured only as total beta concentrations. Although total beta measurements in themselves are not directly useful in the determination of the potential doses, they are indicative of the contaminant loading to the river and serve to identify those years in which concentrations of radioactivity present in the river water as a result of Hanford operations were greatest. Figure 7.1 provides a graphic presentation of the total beta concentrations observed in Columbia River water at Pasco for the years 1947 through 1963. Consistent with the effluent discharges to the river, total beta levels in river water at Pasco were highest during the late 1950s and early 1960s.

The development of gamma energy spectroscopy in 1957 permitted the measurement of specific radionuclides in Columbia River water. Annual average radionuclide concentrations as reported (prior to 1971) in historical annual reports and as contained in the Hanford SESP database (since 1971) are provided in Tables 7.1 through 7.5 for the Richland Pumphouse, Pasco Pump Station, McNary Dam, Bonneville Dam, and Vancouver water sampling locations for any of the years between 1957 and 1990 when data were collected at each site. Figures 7.2 through 7.5 graphically present the data included in Tables 7.1 through 7.5. Apparent in the data is the decrease in radionuclide concentrations with increasing distance from the reactor discharge points and with time following the shutdown of the first of the reactors in December

S9204054.1a

FIGURE 7.1. Total Beta Concentrations in Columbia River Water at Pasco, Washington 1945-1971 (see Table 10.11)

1964. Radionuclide concentrations in river water rapidly declined to levels at or below detection limits following the closure of the last of the original reactors. As this occurred, the remote river water sampling effort gradually ceased, in accord with the decrease in contaminant levels to those which were no longer detectable or of significance with respect to dose evaluations.

7.2 SEDIMENT MONITORING DATA

Although they are not a direct contributor to dose, radionuclides in sediments may be resuspended during high river flows and reintroduced into the aquatic foodchain and may subsequently contribute to the dose potentially received by persons using the river. As in the case of river water, the analyses of sediment samples collected from 1944 through 1957 were limited to total beta measurements. However, direct surveys of the sediment material that provided raw count rate data were also performed. Routine collection of

TABLE 7.1. Annual Average Radionuclide Concentrations in Columbia River Water at the Richland Pumphouse

Year (a)	Radionuclide Concentration, pCi/L											
	³ H (b)	²⁴ Na	³² P	⁴⁶ Sc	⁵¹ Cr	⁵⁶ Mn	⁶⁵ Zn	⁷⁶ As	⁹⁰ Sr	¹³¹ I	¹³² Te	¹³⁷ Cs
1963	..(c)	3400	260	-	8800	-	380	1200	1	8	-	-
1964	-	3500	300	-	12000	-	450	1200	1	19	-	-
1965	-	3100	140	-	7000	-	180	1000	1	10	-	-
1966	-	2600	140	30	3600	290	200	420	1	18	-	-
1967	1500	2600	190	60	3200	520	220	400	1	8	-	-
1968	1700	2200	92	100	1500	250	86	320	0.6	7.4	-	-
1969	1900	1600	73	72	720	1000	72	310	0.5	4	-	-
1970	1100	900	28	43	300	100	34	130	<MDC (d)	<MDC	-	-
1971	779	43.9	22.1	-	127	-	18.1	11.2	0.844	1.64	-	10.4
1972	56.8	-	0.305	-	115	-	5.53	-	0.363	1.16	-	7.37
1973	623	-	-	-	42	-	7.17	-	0.334	1.01	-	2.22
1974	508	-	-	-	9.96	-	3.87	-	0.277	2.1	-	1.63
1975	373	-	-	-	10.3	-	3.6	-	-	0.647	-	2.79
1976	261	-	-	-	17.4	-	1.33	-	0.239	-	-	0.661
1977	585	-	-	-	27.2	-	2.75	-	0.321	-	-	0.783
1978	429	-	-	-	21.7	-	1.96	-	0.474	-	-	2.25
1979	355	-	-	-	16.1	-	-0.514	-	0.336	-	-	2.1
1980	265	-	-	-	1.72	-	1.39	-	0.195	-	-	0.413
1981	199	-	-	-	1.49	-	0.0305	-	0.23	-	-	0.176
1982	216	-	-	-	17.3	-	0.459	-	0.171	-1.34	-	0.211
1983	135	-	-	-	-	-	-0.746	-	0.29	-	-	0.153
1984	169	-	-	-	-	-	-0.639	-	0.169	-	-	0.0935
1985	152	-	-	-	-	-	0.0645	-	0.158	-	-	-0.105
1986	149	-	-	-	-	-	0.0295	-	0.16	-	-	0.347
1987	128	-	-	-	-	-	-0.261	-	0.131	-	-	-0.138
1988	135	-	-	-	-	-	-0.724	-	0.119	-	-	0.121
1989	128	-	-	-	-	-	0.204	-	0.0745	-	-	0.0594

(a) Data not collected from before 1963 and in 1990.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: tritium, sodium-24, phosphorus-32, scandium-46, chromium-51, manganese-56, zinc-65, arsenic-76, strontium-90, iodine-131, tellurium-132, and cesium-137.

(c) - means not determined.

(d) <MDC = below minimum detection concentration.

sediment samples was discontinued in 1959, shortly after the capability to measure specific radionuclides became available.

Special studies of radionuclide concentrations in river sediments, primarily behind McNary Dam, were conducted during and after the shutdown of the original reactors (Fisher 1971; Fix 1975; Nelson et al. 1966; Robertson and

TABLE 7.2. Annual Average Radionuclide Concentrations in Columbia River Water at the Pasco Pump Station

Year (a)	Radionuclide Concentration, pCi/L											
	³ H (b)	²⁴ Na	³² P	⁴⁶ Sc	⁵¹ Cr	⁵⁶ Mn	⁶⁵ Zn	⁷⁶ As	⁹⁰ Sr	¹³¹ I	¹³² Te	¹³⁷ Cs
1959	-(c)	1139	155.7	-	4209	-	206.2	1458.3	0.638	-	-	-
1960	-	1500	200	35	5000	-	300	1500	0.6	12	-	-
1961	-	1800	260	33	5700	-	340	1200	0.4	10	-	-
1962	-	1600	180	30	4300	87	220	470	0.7	6	-	-
1963	-	1600	190	-	6700	-	220	750	1	8	-	-
1964	-	1500	200	-	6800	-	240	670	1	12	-	-
1965	-	-	87	-	4100	-	180	-	-	7	-	-

(a) Data not collected before 1957 and after 1965.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: tritium, sodium-24, phosphorus-32, scandium-46, chromium-51, manganese-56, zinc-65, arsenic-76, strontium-90, iodine-131, tellurium-132, and cesium-137.

(c) - means not determined.

TABLE 7.3. Annual Average Radionuclide Concentrations in Columbia River Water at McNary Dam

Year (a)	Radionuclide Concentration, pCi/L											
	³ H(b)	²⁴ Na	³² P	⁴⁶ Sc	⁵¹ Cr	⁵⁶ Mn	⁶⁵ Zn	⁷⁶ As	⁹⁰ Sr	¹³¹ I	¹³² Te	¹³⁷ Cs
1964	-(c)	-	70.3	-	3478	-	76.5	-	-	6.7	-	-
1965	-	-	47.5	-	2264	-	67.6	-	-	3.9	-	-
1966	-	-	76.2	-	1849	-	61.0	-	-	7.1	-	-
1967	-	-	49.5	-	1748	-	82.2	-	-	4.7	-	-
1968	-	-	39.2	-	769	-	51.1	-	-	3.8	-	-
1969	-	-	33.0	-	333	-	44.2	-	-	2.6	-	-

(a) Data not collected from before 1964 and after 1969.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: tritium, sodium-24, phosphorus-32, scandium-46, chromium-51, manganese-56, zinc-65, arsenic-76, strontium-90, iodine-131, tellurium-132, and cesium-137.

(c) - means not determined.

Fix 1977). Five years after the shutdown of the final once-through-cooled reactor, the short- and intermediate-lived radionuclides had decayed away. However, measurable concentrations of a few long-lived radionuclides remained in the deep sediments behind McNary Dam. Sediments free of Hanford-originated radionuclides covered the contaminated radionuclides behind McNary Dam at a rate of approximately 15 to 30 inches per year between 1971 and 1976 (Robertson and Fix 1977).

TABLE 7.4. Annual Average Radionuclide Concentrations in Columbia River Water at Bonneville Dam

Year (a)	Radionuclide Concentration, pCi/L											
	³ H (b)	²⁴ Na	³² P	⁴⁶ Sc	⁵¹ Cr	⁵⁶ Mn	⁶⁵ Zn	⁷⁶ As	⁹⁰ Sr	¹³¹ I	¹³² Te	¹³⁷ Cs
1964	-(c)	-	28	-	2400	-	63	-	-	5	-	-
1965	-	-	23	-	1700	-	70	-	-	3	-	-
1966	-	-	23	-	1300	-	43	-	-	3	-	-
1967	-	-	25	-	1400	-	62	-	-	3	-	-
1968	-	-	15	20	530	-	30	-	-	3.2	-	-
1969	-	-	14	-	240	-	25	-	-	-	-	-
1970	-	-	5	-	100	-	10	-	-	-	-	-
1971	-	-	-	2.6	-	-	5.9	-	-	-	-	-
1972	-	-	-	0.15	-	-	1.1	-	-	-	-	-
1973	-	-	-	-	-	-	8	-	-	-	-	-
1974	-	-	-	<MDC (d)	<MDC	-	<MDC	-	-	-	-	<MDC
1975	-	-	-	<MDC	<MDC	-	<MDC	-	-	-	-	<MDC

(a) Data not collected from before 1964 or after 1976.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: tritium, sodium-24, phosphorus-32, scandium-46, chromium-51, manganese-56, zinc-65, arsenic-76, strontium-90, iodine-131, tellurium-132, and cesium-137.

(c) - means not determined.

(d) <MDC = below minimum detection concentration.

TABLE 7.5. Annual Average Radionuclide Concentrations in Columbia River Water at Vancouver, Washington

Year (a)	Radionuclide Concentration, pCi/L											
	³ H (b)	²⁴ Na	³² P	⁴⁶ Sc	⁵¹ Cr	⁵⁶ Mn	⁶⁵ Zn	⁷⁶ As	⁹⁰ Sr	¹³¹ I	¹³² Te	¹³⁷ Cs
1959	-(c)	-	31	-	2100	-	37	-	-	-	-	-
1960	-	-	41	18	2100	-	75	-	0.4	4	-	-
1961	-	-	68	24	2100	-	90	-	0.4	-	-	-
1962	-	-	38	20	1800	-	64	-	0.4	3	-	-
1963	-	-	30	-	2600	-	60	-	1	4	-	-
1964	-	-	50.3	-	3317	-	55.8	-	-	4.2	-	-

(a) Data not collected from before 1959 and after 1964.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: tritium, sodium-24, phosphorus-32, scandium-46, chromium-51, manganese-56, zinc-65, arsenic-76, strontium-90, iodine-131, tellurium-132, and cesium-137.

(c) - means not determined.

FIGURE 7.2. Annual Average Radionuclide Concentrations in Columbia River Water at the Richland Pumphouse and Pasco Pump Station (for sodium-24, phosphorus-32, chromium-51, zinc-65, and arsenic-76)

SB202009.2

FIGURE 7.3. Annual Average Radionuclide Concentrations in Columbia River Water at McNary Dam (for phosphorus-32, chromium-51, and zinc-65)

Sediment samples were also collected from The Dalles Dam and Bonneville Dam. Radionuclide concentrations in these sediments were lower than those observed in the McNary reservoir sediments. Radionuclide concentrations in sediment samples collected during 1976 were reported in disintegrations per minute per gram (dpm/g). Table 7.6 provides a comparison of radionuclide concentrations in surface sediment behind McNary Dam during 1971 and 1976.

7.3 BIOTA MONITORING DATA

Concern for the aquatic biota and the effects of the reactor effluent on the biota were expressed very early in the development of the Hanford Site,

S9202009.3

FIGURE 7.4. Annual Average Radionuclide Concentrations in Columbia River Water at Bonneville Dam (for phosphorus-32, chromium-51, and zinc-65)

even before construction was begun. Numerous laboratory and field investigations have been conducted over the years to investigate the effects of reactor effluent on biota. Summaries of these studies have been documented, including discussion of the study results (Becker 1990). Consumption of fish contaminated with Hanford effluent radionuclides was recognized as a primary pathway through which the public could be exposed to radionuclides originating at Hanford.

Phosphorus-32 and zinc-65 were the two radionuclides of concern in Columbia River fish because these isotopes accumulate significantly in the edible muscle tissue. It was estimated that these two radionuclides

S8202009.4

FIGURE 7.5. Annual Average Radionuclide Concentrations in Columbia River Water at Vancouver, Washington (for phosphorus-32, chromium-51, and zinc-65)

contributed more than 90% of the calculated dose resulting from the consumption of fish. Figures 7.6 and 7.7 indicate the monthly mean phosphorus-32 and zinc-65 concentrations in muscle tissue of whitefish collected at Ringold during the years 1964 through 1966. The 90% confidence interval is also provided in the figures. Data on whitefish are presented because their radionuclide concentrations indicate the highest levels observed in Columbia River sport fish. Concentrations in suckers were typically the highest for any large Columbia River fish (Davis et al. 1958). However, whitefish were thought to be more important from the standpoint of potential dose because

TABLE 7.6. Radionuclide Concentrations in Surface Sediments in McNary Reservoir in 1971 and 1976 (from Robertson and Fix 1977)

Element	Typical Concentrations in dpm/g Dry Sediment	
	April 1971	August 1976
Iron-55	1100	30 (est.)
Zinc-65	240	0.14
Scandium-46	120	0.34
Cobalt-60	60	2.7
Europium 152,154	51	2.2
Manganese-54	25	0.32
Cesium-137	9	2.7
Plutonium 239,240	0.06	0.03

S8912060.2

FIGURE 7.6. Monthly Mean Phosphorus-32 Concentrations in Columbia River Whitefish, 1964-1966

FIGURE 7.7. Monthly Mean Zinc-65 Concentrations in Columbia River Whitefish, 1964-1966

sports fishermen harvest large numbers of these fish, as opposed to suckers, which are rarely eaten (Foster and Junkins 1960).

Various marine organisms that indicate the presence of Hanford-originated radionuclides have been collected routinely from areas near the mouth of the Columbia River. Oysters generally contained higher concentrations of zinc-65 than other marine organisms (Foster and Wilson 1964). Table 7.7 presents the radionuclide concentrations observed in Willapa Bay oysters for the years 1959 through 1977. The collection of oysters from Willapa Bay was discontinued during 1978 because Hanford-originated radionuclides had declined to levels that were generally less than detectable. Figure 7.8 graphically displays the zinc-65 concentrations reported in the annual environmental reports. Evident from the figure is the relatively rapid decline in radionuclide concentrations following the shutdown of the once-through-cooled reactors.

TABLE 7.7. Annual Average Radionuclide Concentrations in Willapa Bay Oysters, 1959-1977

Year (a)	Radionuclide Concentration, pCi/L									
	³² P(b)	⁴⁰ K	⁴⁶ Sc	⁵¹ Cr	⁵⁸ Co	⁶⁰ Co	⁶⁵ Zn	⁹⁰ Sr	¹³¹ I	¹³⁷ Cs
1959	-(c)	-	-	-	-	-	51	-	-	-
1960	0.57	2.4	0.12	0.54	-	-	55	0.0032	<MDC(d)	<MDC
1961	1.7	1.4	0.11	0.59	-	-	67	-	<MDC	<MDC
1962	2.9	2.1	-	-	1.0	<MDC	91	-	-	0.34
1963	3.9	6.5	-	-	0.82	<MDC	80	-	-	0.31
1964	4.7	2.0	-	-	-	-	54	-	-	<MDC
1965	3.7	2.0	-	-	-	-	39	-	-	0.13
1966	3.0	2.1	-	-	-	-	28	-	-	0.10
1967	3.4	2.1	-	-	-	-	32	-	-	0.10
1968	1.9	2.2	-	-	-	-	25	-	-	0.10
1969	3.3	2.0	-	-	-	-	19	-	-	0.16
1970	0.61	1.9	-	-	-	-	13	-	-	0.04
1971	1.3	1.8	-	-	-	-	4.6	-	-	0.04
1972	-	1.7	-	-	-	-	1.7	-	-	0.025
1973	-	1.7	-	-	-	-	0.57	-	-	0.02
1974	-	1.6	-	<MDC	-	<MDC	0.14	-	-	<MDC
1975	-	1.7	-	-	-	-	<MDC	-	-	<MDC
1976	-	1.4	-	-	-	-	<0.08	-	-	<0.04
1977	-	1.4	-	-	-	-	<0.08	-	-	<0.04

(a) Data not collected before 1959 and after 1977.

(b) From left to right, abbreviations in column headings stand for the following radionuclides: phosphorus-32, potassium-40, scandium-46, chromium-51, cobalt-58, cobalt-60, zinc-65, strontium-90, iodine-131, and cesium-137.

(c) - means not determined.

(d) <MDC = below minimum detection concentration.

FIGURE 7.8. Zinc-65 Concentrations in Willapa Bay Oysters, 1959-1977

8.0 SPECIAL STUDIES AND SURVEYS AT HANFORD

This section discusses effluent plume dispersion and shoreline radiation surveys in the Hanford reach of the Columbia River. Plate 2 (in a pocket in the back of this report) will help the reader track the effluent plume and locate the areas surveyed at various times.

8.1 EFFLUENT PLUME DISPERSION

Monitoring of radioactivity in the Columbia River began in 1945 with routine sampling of river water for total gross beta activity. The sampling was conducted at areas near the operating reactors (initially 100-B, 100-D, and 100-F), the Hanford Ferry Landing, the 300 Area, Richland, and Pasco. The horn of the Yakima River was sampled for background levels. Sampling stations were added as more reactors came on-line.

Early in the sampling program scientists realized that radioactivity was not uniformly distributed across the channel and downstream. An early report by Turner (1947) stated that "there is not much mixing of this effluent by the time it reaches [the] Hanford [townsite]." Observations from 1947 indicate that the radioactive "channel" within the river appeared to hug the south bank (reactor shore).

By 1950 the pattern of effluent dispersion in the Hanford reach was well established. The centerline of the effluent plume from the 100-B reactor to Pasco is shown in Plate 2. Immediately below 100-B, the discharge point farthest upstream, the beta activity was confined to a narrow plume that gradually widened to about 400 feet in the 100-D Area (Paas and Singlevich 1950). At the 100-D Area, the high-velocity path of flow was directed toward the opposite shore to the north of 100-D Island. However, the maximum measured beta activity remained along the reactor shore. Although the activity became more diffuse toward the north shore, there were no indications of cross-channel mixing (Paas 1951a). The plume continued to hug the reactor shore between 100-H and 100-F, especially when high water elevations caused effluent to be discharged over a spillway directly into the river (Paas and Singlevich 1951a). Immediately downstream of the 100-F Area, the maximum zone

of radioactivity tended to move directly toward the reactor side of the river (Paas and Singleovich 1950), even though the high-velocity flow was directed toward the opposite shore around an island at RM 366-367. It was assumed that the island forced the plume toward the reactor shore (Paas 1951b), but the maximum activity in this area may have actually been a consequence of the shortness of the outfall line at 100-F.

The maximum beta activity generally occurred near the Hanford Ferry Landing, possibly because all reactors were contributing at this point. By the time it reached this point, the plume was about 5 miles long and 500 feet wide. Downstream of the Hanford Ferry Landing the mixing across the river was better, although the plume could still be discerned along the shore at Richland (Honstead, Healy, and Paas 1951).

The plume distribution in the river was confirmed by ferro-floc dispersion studies conducted in September 1950 and April 1951. Deep red ferro-floc sludge from the sedimentation basins was released to the river and used as a tracer. Aerial photography indicated that the plume formed a narrow band that persisted for 6 to 10 miles downstream from the release point (Rostenbach 1956).

In 1949 and again in 1951, special studies were conducted to determine the effects of the Yakima River confluence on the plume pathway. Results indicated that the Yakima River water diluted the radioactivity by as much as a factor of two along the west shore and forced the higher radioactivity levels toward the east (Pasco) shore. These effects persisted for at least 7 miles downstream (Paas and Singleovich 1951b).

Radioactivity distribution studies in the McNary reservoir in 1954 indicated little cross-sectional variation. The maximum beta activity was located in the middle of the river, while minimum concentrations fluctuated between the Washington and Oregon shorelines (Paas 1954).

Only one significant plume dispersion study was carried out below the reactor areas. Tracer dye was released into the river from the 300 Area in August 1961, October 1961, and January 1962. Aerial photography and water samples were used to define the plume distribution (Backman 1962, 1963). The

objectives were to determine whether complete mixing occurred by the time the dye reached Pasco and to identify the dispersion configuration for low-flow conditions.

All three tracer tests produced similar results. For each test, the dye tended to remain near the 300 Area shore, and the plume passed directly over the location of the Richland pumping station intake. At this point the plume was almost 3 miles long. Downstream of the proposed pumping station, the plume moved east of the fourth island but left pools of dye along the west shore. More mixing was observed off the downstream tip of the island than at any other location. Lateral dispersion was almost complete by the time the plume reached Pasco. Vertical sampling indicated there were no variations in dye concentration except within about 300 feet of the release point, suggesting that vertical dispersion was complete within a much shorter mixing length.

All raw data collected during downstream dispersion studies undertaken before 1960 were compiled in a section of a rather large monitoring report (Soldat 1962a). The data include river velocity profiles, ferro-floc distributions, radioactivity density profiles, and the results of early dye tests.

8.2 SHORELINE RADIATION SURVEYS

The earliest information on a shoreline radiation survey was reported by Paas (1953). A survey of 100-D Island, using portable instruments, showed readings ranging from 500 counts/minute to 5000 counts/minute around the island perimeter; readings of 35 millireps per hour (mrep/hr) (1 rep = 0.93 rad) were measured at locations adjacent to the point where the 100-D outfall discharged effluent. These readings were not related to any unit area. Beta activity in island mud was 0.2 $\mu\text{Ci/g}$, but no alpha particle emissions were detected in the mud (Paas 1953). The locations of the shoreline radiation surveys are shown in Plate 2.

8.2.1 Survey of 1959

A shoreline survey was conducted during March and April of 1959 to correlate a new sensitive scintillator for the measurement of gamma dose rates

from reactor effluent in the Columbia River. The river discharge was approximately in the range of 100,000 to 125,000 cfs. The survey results were reported by McConiga and Rising (1959).

The survey combined scintillator readings with pulse pencil and portable instrument readings taken along the river from 1.5 miles upstream of the reactors to about 57 miles downstream. Gamma pulse pencils, in sets of five, were placed 3 feet above the ground on stakes on both shorelines at each survey river mile beginning at 1.5 miles above the reactors to 20 miles downstream of that point. From the 20-mile point, pencils were placed at each mile on alternate shores. At least one group of pencils was placed on each island, but the exact locations were not described.

Scintillator measurements in the boat over effluent bubbles varied from 0.6 to 1.8 milliroentgens per hour (mR/hr) to a maximum of 2.5 mR/hr with the scintillator held outside the boat. At 6 miles below the reactors, dose rates decreased to about 0.06 mR/hr through the boat and 0.1 mR/hr outside the boat. Shoreline readings from the pencils indicated a dose rate of 0.1 mR/hr to Richland and on some islands. A reading of 15 mR/hr was measured on 100-D Island. The Pasco pumping plant and Sacajawea Park recorded 0.022 mR/hr and 0.018 mR/hr, respectively. Background was about 0.018 mR/hr according to the authors.

8.2.2 Survey of 1961-1962

During 1961 and 1962, three surveys were conducted to determine average exposure rates on beaches and islands for 33 locations between Ringold and Richland (McConnon 1962). (Sampling locations are shown on Plate 2.) The locations were chosen for their attractiveness to swimmers, sunbathers, and boaters. The surveys were conducted in July 1961, early October 1961, and late March 1962. The river discharge at the time of the July survey was approximately 175,000 cfs following a June peak of approximately 500,000 cfs. By October, the discharge was reduced to 60,000 cfs, and by the end of March, the discharge was about 50,000 cfs. Measurements were taken in approximately the same areas during the first two surveys. For the third survey, measurements were repeated from Ringold to the 300 Area, but emphasis was placed on areas opposite Richland.

Four types of radiation detection instruments were used: 1) a portable Geiger-Müller (GM) detector for extent and intensity of contamination, 2) a 40-liter ionization chamber for extremely low exposure rates, 3) a 5-inch plastic scintillation detector for exposure rates in boats, and 4) small "pencil" ionization chambers for measuring integrated exposure rates over a 1-week period per location. The small pencil instruments were used both on the shoreline and in the water. At shoreline locations, they were placed at ground level and 3 feet above ground to measure exposure from either lying or standing on the beach. Pencils were also submerged in 4 to 8 feet of water about 10 to 15 feet offshore to estimate exposure rates from swimming.

The July results showed beach activity levels from about 200 counts/minute to a maximum of 800 counts/minute (portable GM detectors). The highest readings were on the upper half of Island Z; however, there was not a consistent decrease downstream near Richland. The contamination was not associated with scattered particles but seemed to be spread uniformly over the surveyed surfaces. The pencil results indicated ground-level rates ranging from 0.06 to 0.19 mR/hr with levels at the 3-foot height being about half the ground values. Pencil instruments submerged offshore indicated a range from 0.05 to 0.09 mR/hr. The 40-liter chamber showed exposure rates slightly higher than the pencil instruments and a lack of any distinct change downstream near Richland (McConnon 1962).

During the October survey, measurements were taken on beaches that had been submerged during the July survey; the rates were significantly higher. The GM detector recorded measurements as high as 5000 counts/minute. The 40-liter chamber measured a maximum value of 1.1 mR/hr. The pencil range at ground level was from 0.06 to 0.21 mR/hr, with the exposure rate in water about twice that in July. There was no significant decrease downstream near Richland (McConnon 1962).

The March survey focused more on the shorelines from the 300 Area to Richland. The measurements taken with the GM detectors and the 40-liter chamber indicated no substantial increase in exposure over the October survey. Sand samples from five locations were analyzed for the contributing

radionuclides. The major contributors were zinc-65 and chromium-51 in approximately equal amounts, with neptunium-239 and lanthanum-140 contributing to a lesser degree.

8.2.3 Shoreline Surveys Reported in 1966

Two shoreline surveys were reported in 1966. (Sampling locations are shown on Plate 2.) The first survey included specific locations from Vernita Ferry Landing (upstream of the reactors) to Sacajawea Park at the Snake River confluence. The second survey was of the reactor areas and extended from the 100-B reactor (RM 384) to White Bluffs (RM 370).

Vernita Ferry Landing to Sacajawea Park Survey

This survey, reported by Grande (1966), is dated January 31, 1966, but may have been conducted sometime in 1965. No specific survey date is mentioned in the report. However, most of the report is concerned with a comparison between counting instruments.

For each site along the shoreline, gamma activity measurements were taken at the beach surface and at elevations of 3 feet and 5 feet above the surface. At each site, three locations were chosen: 1) near the maximum flood level, 2) the maximum weekly flow level, and 3) the water line existing at the time of measurement.

Based on the initial GM detector survey of the general areas in question, the highest readings were measured at the river side of the Hanford Slough and the north shore of the old Hanford Ferry Landing (Plate 2). Readings taken from mossy rocks at these locations ranged from 1000 to 3000 counts/minute with a maximum of 15,000 counts/minute. The readings remained in this range from the reactor locations to the 300 Area but decreased downstream from that point to 250 counts/minute at Richland and 150 counts/minute at the Finley lagoon (opposite the Snake River confluence).

Counting rate normally decreased rapidly with increasing distance and height from the shoreline, except for one location. At a point in the Hanford Slough, the activity level increased from 150 counts/minute at the surface to 400 counts/minute at heights of 3 to 5 feet. This was because of improved geometry for the gamma radiation originating at the shoreline and river.

Vegetation that received irrigation water from the river at least once per year showed a significant concentration of radionuclides (200 to 1000 counts/minute). Readings at the water's edge at Ringold ranged from 1000 to 3500 counts/minute.

The major contributor to exposure rates was zinc-65, with small additional contributions from sodium-24 and chromium-51. Accumulations of radionuclides were found mainly in algae, among the rocks, and on native shoreline vegetation (Grande 1966).

Reactor Areas Survey

This second survey, reported by Lodge (1966), was conducted in March and April 1966. This extensive GM detector survey extended from the 100-B reactor (RM 384) to White Bluffs (about RM 370). The reactor farthest downstream, 100-F, is at RM 369 (Plate 2). Measurements were made at the water line, the daily water elevation, and the annual high-water mark of 44 locations on the reactor shore, 25 far-shore locations, and 12 island locations. For son purposes, an aerial survey, using a sodium iodide scintillation cal, was conducted concurrently with the GM detector survey.

The objectives of the survey were to determine shoreline radiological conditions, check instrumentation, and determine major contributing radionuclides (Lodge 1966). Radioactivity levels were highest at the water line on the islands opposite and below the 100-D Area. Maximum shoreline readings were observed along the reactor side of the river just below the 100-K Area and again below the 100-N Area. On the far shore, background levels were measured as far downstream as the 100-D Area. Between the 100-D Area and White Bluffs, the readings increased, with the highest at the bluffs. The maximum spot activity level was 78,000 counts/minute at the downstream end of 100-D Island, and a maximum shoreline level of 12,000 counts/minute was found on the next island downstream (Island E). The locations of these maxima are most likely the result of leaks from the combined 100-D and 100-DR perforated outfall line extending across 100-D Island. For the reactor shoreline, the water-line maximum levels ranged from 100 to 4000 counts/minute; the daily wet-line maxima ranged from 100 to 1000 counts/minute; the annual high-water

maxima ranged from 100 to 350 counts/minute. The major contributing radionuclides were zinc-65; chromium-51; manganese-54,56; and scandium-46.

8.2.4 Post-Reactor Shutdown Surveys

Two shoreline radiation surveys were conducted after the shutdown of the eight once-through-cooled reactors. The purpose of these surveys was to assess the association of radionuclides with sediments in the river system. The first of these two surveys was reported in 1975 and included a detailed aerial radiation survey flown March 26 and April 28, 1974. The survey also included analysis of sediment samples from the shoreline, island, and slough areas shown in the aerial surveys to have the highest activity levels and analysis of sediment cores from Priest Rapids (background) and McNary dams (Fix 1975). The aerial survey was conducted along the Hanford reach, from the vicinity of the Vernita Bridge to several miles below the Snake River confluence. An additional aerial survey was included for several miles downstream of McNary Dam.

The highest activity levels were found along the slough north of the old Hanford townsite and in the slough between 100-D and 100-H reactors. Radiation levels in the slough north of the old townsite included 0.022 mR/hr (maximum survey reading) from cobalt-60 and 0.001 mR/hr from cesium-137. Radiation levels in the slough between 100-D and 100-H were 0.014 mR/hr from cobalt-60 and 0.003 mR/hr from cesium-137. Similar activity levels were measured along the 100-F slough and on the islands upstream of the 300 Area downriver to those adjacent to Columbia Park. The radiation levels were highly variable and indicate no decrease in activity downstream.

A 22-inch-long core taken somewhere along the reactor shore at Hanford indicated that the primary radionuclides contributing to exposure rates were cobalt-60, cesium-137, and europium-152. Also present were manganese-54, cobalt-58, zinc-65, ruthenium-106, cesium-134, cerium-144, and europium-154. Analysis of cores from McNary reservoir showed that the maximum activity was from europium-152, cobalt-60, and zinc-65, with contributions from scandium-46, manganese-54, and cesium-137.

The second post-shutdown survey was conducted during the spring and summer months of 1979 and reported by Sula (1980). The survey area included shorelines and islands between the uppermost point of reactor discharge (100-B) and the confluence of the Snake River, almost a 60-mile length of river. From these measurements, three basic types of contamination were identified: 1) a uniform layer of contamination over the entire area, 2) areas of higher contamination referred to as "contamination deposits," and 3) discrete particles containing cobalt-60.

Uniformly distributed contamination consisted of a constant level of radiation, slightly higher than background, that extended from 100-B to the Snake River. The average exposure rate was 11 ± 3 microroentgens per hour ($\mu\text{R/hr}$); the background radiation level was 7 ± 1 $\mu\text{R/hr}$. This slightly higher exposure rate was attributed to cobalt-60 and europium-152 everywhere, with cesium-137 contributing at the 100-N Area and on the Hanford townsite peninsula. There was no downstream decrease in activity levels.

Contamination deposits, exhibiting exposure rates that were significantly higher than the uniform contamination rate, were located throughout the reach. Ninety-two areas exhibited exposure rates exceeding 25 $\mu\text{R/hr}$. These areas were attributed to the presence of contaminated sediments that had been concentrated by river processes. The areas ranged from a few square meters to several thousand square meters, usually in dense vegetation. The highest contamination deposits were at the White Bluffs slough area (40 $\mu\text{R/hr}$), the Hanford townsite peninsula (45 $\mu\text{R/hr}$), and the island at RM 344 near the 300 Area (38 $\mu\text{R/hr}$). The remaining contamination deposits were in the 25- to 30- $\mu\text{R/hr}$ range and appeared to be evenly distributed over the survey area. Samples of soil and vegetation indicated that the radionuclides in the deposits consisted of a mixture of cobalt-60, cesium-137, and europium-152 in approximately equal proportions.

Discrete particles of contamination containing cobalt-60 were found along the river, with the highest concentrations on the group of islands between 100-D and Locke Island. Locations with the greatest number of particles were the island at RM 375 (below 100-D Island); the 100-F Area floodplain, and the islands at RM 367, RM 353, and RM 350. The particles tended to

decrease in number downstream and were found both in flat, rocky, unvegetated areas and above the daily high-water level. The particles were metallic flakes, possibly fragments of stellite valve and pump components used in production reactors. Fourteen particles were recovered and found to contain from 1.7 to 24 μCi of cobalt-60.

8.3 DOWNRIVER TRAVEL TIMES

Beginning in 1955, Hanford scientists began studies to determine downstream travel times of reactor effluent released to the Columbia River. The primary reason for the studies, according to the initial report on the studies (Soldat 1956), was the problem of evaluating the hazards resulting from the discharge of radioactive effluent and the subsequent consumption of river water in cities downstream of Hanford. Knowledge of travel time was required to establish decay correction factors for river water samples collected for monitoring and to determine what sampling times at downstream locations would represent specific Hanford operating conditions.

The initial approach to determining these travel times was to use a system of river surface floats with the objective of measuring the minimum travel time to various points downstream for a range of river discharges. The tests were confined to the reach from the reactors to Pasco. Two types of floats, rod floats and cork floats, were used. In either case, most of the float was submerged just below the surface to reduce the effect of wind. Beginning in April 1955 and ending in August 1955, intermittent tests were conducted for river discharges ranging from 86,000 to 360,000 cfs. The study results provided a minimum travel time to the Pasco-Kennewick area of 22.4 hours at 90,000 cfs and 11.2 hours at 360,000 cfs. Because of the backwater effects of the McNary reservoir, these travel times were much slower than those possible under the free-flow conditions that existed until dam construction.

During the next 5 to 6 years, the travel times were slightly refined to provide more detail and improved graphical presentation of the information (Soldat 1962a). Comparisons were made between data collected by the U.S. Army Corps of Engineers and by Hanford contractors; data were also compared with results from a mathematical equation used to calculate flow times.

During 1964, studies were conducted to determine downriver flow times from the reactors to any point downstream as far as Astoria, Oregon (Nelson, Perkins, and Haushild 1965, 1966). These studies used two tracer methods to determine the travel times. One method used the decay of sodium-24 (15.0 hours) as an index of travel time to various points downstream. The other method, which was used simultaneously, measured the time required for a "peak" of iodine-131 activity to reach the locations. Measurements were made during January and July 1964. The results were tabulated (Table 8.1) from the reactors to Vancouver, Washington, for low, intermediate, and high discharges to the Columbia River and from the reactors to Astoria, Oregon, for very high discharges to the river.

TABLE 8.1. Travel Times of Peak Concentrations in the Columbia River 100-D Reactor (RM 377.6) to Various Downstream Points, Measured During July 1964

Downstream Points	River Mile	Low Discharge		Intermediate Discharge		High Discharge		Very High Discharge	
		Time, days	Discharge, cfs x 10 ³	Time, days	Discharge, cfs x 10 ³	Time, days	Discharge, cfs x 10 ³	Time, days	Discharge, cfs x 10 ³
Pasco	330	1.0 ^(a)	65	0.67	173	0.48 ^(a)	309	0.43 ^(a)	432
Finley	324	1.6	66	---	---	---	---	---	---
Umatilla	290	6.0	89	2.6	222 ^(b)	1.7	560 ^(b)	1.3	624 ^(b)
Biggs	208	7.9	95	---	---	---	---	---	---
The Dalles	191	9.5	102	4.6	209	2.5	577	2.3	627
Hood River	170	10.7	105	---	---	---	---	---	---
Vancouver	107	14.6	108	7.2	192 ^(c)	3.7	585 ^(c)	3.6	627 ^(c)
Astoria	14	---	---	---	---	---	---	5.7	657

(a) Estimated from Soldat (1962a).

(b) Includes Snake and Walla Walla Rivers.

(c) Upstream of Willamette River.

9.0 MONITORING AND STUDIES BY OFFSITE AGENCIES

Reports regarding Columbia River and coastal area contamination by Hanford reactor releases have been produced by state and federal agencies with specific interests in the river. These agencies included the states of Washington and Oregon and their state universities with marine science programs and laboratories (University of Washington and Oregon State University). Two federal agencies provided radiological data and information for the Columbia River system: the U.S. Geological Survey (USGS) and the U.S. Public Health Service (USPHS) under the U.S. Department of Health, Education, and Welfare. The USGS conducted a lengthy and comprehensive study of sediment and radionuclide transport in cooperation with Hanford Site contractors. The USPHS conducted a water-quality sampling program to obtain a database for evaluating the effects of McNary Dam on the Columbia River. Results of this program also include data on radioactive effluent releases from Hanford. Although the USPHS program was called a water-quality study, a significant number of biota samples were collected and analyzed for activity levels.

9.1 STATE OF WASHINGTON

For 1961 to 1962 and 1969 through 1976, samples of biota, water, and sediment were analyzed for concentrations of certain radionuclides. Biota considered were clams, oysters, and crabs. Samples were collected at several locations along the Columbia River and the Washington coast.

A radiological surveillance was conducted from September 1961 to April 1962 by the Washington Department of Health (1962) for statewide river systems. Only gross gamma and gross beta measurements were made, and no specific radionuclides were identified.

Reports by Mooney (1970, 1972, 1974, 1975, 1976) present the results of radiation surveillances. These reports present results of a continuing program documenting the significant reduction of radioactivity levels in the Columbia River during the systematic shutdown of the original Hanford production reactors. The radionuclides considered were zinc-65, chromium-51, phosphorus-32, and scandium-46. Most of the concentrations were relatively

low (i.e., below a certain detection level or not analyzed). This was true for both the water column and the biota (e.g., oysters, salmon, halibut).

From 1977 through 1987, water column and sediment samples were analyzed for tritium, strontium-90, and some gamma-emitters. The samples were collected from two locations: the Hanford Site and the Columbia River in the vicinity of the Trojan Nuclear Plant. For the gamma-emitters, concentrations were generally either below specific detection levels or were not analyzed. The primary radionuclides in the river were tritium and strontium-90. The report by Mooney (1977) shows a range of less than 200 to 430 picocuries per liter (pCi/L) of tritium for the Columbia River at Richland for the months of April, May, and June 1977. A strontium-90 composite for the same period at Richland was 0.23 pCi/L. Tritium activity for the Columbia River at Longview, Washington, varied from less than 200 to 410 pCi/L. Similar results were documented for the years up through 1987. The results were listed in tables and presented graphically. However, the description of methods used is incomplete.

9.2 UNIVERSITY OF WASHINGTON

Reports from the University of Washington were published from 1960 through 1970. Radionuclide concentrations were determined for the water column, sediment, and several species of biota. Zinc-65 was the primary isotope considered. Others were chromium-51, phosphorus-32, scandium-46, potassium-40, and several others to a very limited extent. The sample locations were the coastal areas of Washington and Oregon, including Puget Sound. Specific locations were North Head (just north of the Columbia River mouth), Willapa Bay, and Hood Canal, all in Washington.

No regular monthly or yearly monitoring programs were carried out by the University of Washington. Although all the studies were conducted by the Laboratory of Radiation Biology and some reports support others, each has its own small and restricted data set. Radionuclide concentrations in the water column and sediment were determined primarily to support analysis of various species of biota (e.g., plankton, shellfish).

Studies of radioactivity in plankton were conducted from 1961 through 1963 and reported by Seymour,^(a) Seymour and Lewis (1964), and Lewis and Seymour (1965). The sampling locations were along the Washington and Oregon coastlines and Puget Sound. The radionuclides of primary interest were zinc-65 and chromium-51. Seymour^(a) reported the range of zinc-65 concentration in plankton sampled in 1961 to be from 6 to 980 pCi/g dry weight. Seymour and Lewis (1964) reported maximum concentrations of zinc-65 in plankton as ranging from 110 to 1300 pCi/g dry weight.

Naidu (1963) reported on zinc-65 concentrations in Willapa Bay oysters based on samples gathered during February, April, May, and June 1963. Concentrations of zinc-65 ranged from 450 to 618 pCi/g dry weight. The corresponding range for Willapa Bay plankton ranged from 33.8 to 448 pCi/g dry weight. Concentrations in the mud ranged from 2.69 to 7.49 pCi/g dry weight. The data analysis and discussion present a reasonably complete assessment of zinc-65 in Willapa Bay for the short period of sampling.

Isakson (1969) studied phosphorus-32 activity in biota at North Head on the Washington coast near the mouth of the Columbia River. The sampling period was from October 1965 to September 1966. Most of the data are presented graphically for tissue from clams, mussels, barnacles, anemones, and algae. The data analysis and discussion present a detailed assessment of phosphorus-32 activity for this location.

The most significant publication from the University of Washington is a text edited by Pruter and Alverson (1972), The Columbia River Estuary and Adjacent Pacific Ocean Waters. The text contains 34 papers on the biological and chemical characteristics of the estuary and coastal areas, including several papers on sediment; most of the information concerns biological studies. Nine of the papers (Chapters 25-33) deal with radionuclides in the estuary and coastal system. The primary radionuclides investigated are zinc-65, chromium-51, scandium-46, and phosphorus-32. Only six of the papers contain data of potential use for estimating dose. Most of the concentrations

(a) Progress report prepared by A. H. Seymour, 1961, Laboratory of Radiation Biology, University of Washington, Seattle, Washington.

are associated with biota from the estuary bottom and coastal shelf, although some water column and sediment concentrations are presented. The reported studies cover the period from 1960 to 1969. The most important contributions are the analysis and discussions of the Columbia River estuarine and coastal area processes and how they distribute radioactivity.

9.3 STATE OF OREGON

The Health Division of the Oregon State Department of Human Resources initiated a study in June 1961 to identify and monitor the activity levels of neutron activation and fission product radionuclides in the lower Columbia River and Oregon coastal areas. The primary concern was the continual release to the Columbia River of radionuclides in the cooling water effluent of the Hanford reactors. The objective of the study was to determine the extent of the distribution of the radionuclides and their uptake by biological organisms and sediments. The work was supported by a contract with the Division of Radiological Health of the USPHS, with supporting funds from the Oregon State Board of Health. When the contract expired on July 31, 1967, the Oregon State Board of Health continued to maintain the program to serve three purposes: 1) to establish a continuous background of radiological data to be used in assessing radioactivity from future nuclear power installations on the lower Columbia River, 2) to monitor the changes in levels of radioactivity resulting from reactor shutdowns at Hanford, and 3) to provide a basis for the evaluation of radionuclide intake of individuals consuming fish, shellfish, and other Columbia River foods containing radioactivity.

The state's original upstream sampling location was the John Day Dam site (RM 215.6), but its use was discontinued in November 1963 because of filling operations and construction extending downstream from the damsite (Figure 9.1). After 1963, The Dalles Dam (RM 191.5) was the upstream sampling location. The other Columbia River locations, in downstream order, were Rooster Rock State Park (RM 128.4), Goble (RM 74.0), Beaver Army Terminal (RM 53.6), and Astoria (RM 13.7), all in Oregon.

Oregon coastal locations (in miles below the mouth of the Columbia River) were Seaside (28 miles), Cannon Beach (35 miles), Nehalem River jetty

S9112088.6

FIGURE 9.1. Columbia River and Coastal Sampling Locations for 1961-1967 Studies by the State of Oregon (Toombs and Cutler 1968)

(50 miles), Tillamook Bay (62 miles), Agate Beach-Yaquina Bay (140 miles), and Coos Bay (260 miles). Both the river and coastal locations, with the type of sampling conducted at each, are mapped in Figure 9.1.

The results of the study are documented in reports by Toombs and Cutler (1968) and Toombs and Paris (1978). The 1968 report, covering the period from 1961 to 1967, states that ten radionuclides were detected using gamma spectrometry and radiochemical procedures. Zinc-65, chromium-51, and phosphorus-32 were of special concern because of their abundance and biological significance. The highest levels were found in sediments and algae. The most widely distributed radionuclide, zinc-65, was found in significant concentrations, especially in edible shellfish. Activity levels of zinc-65 and

chromium-51 in water and sediment from the Columbia River for 1962 through 1967 are shown in Table 9.1. Activity levels of zinc-65 in water, sediment, and shellfish from Oregon coast locations for 1962 through 1967 are shown in Table 9.2.

From 1967 to 1977, the studies continued much as they had been conducted from 1961 to 1967, except special studies were conducted in the vicinity of Tillamook Bay in 1970 and 1971 because of the potential for exposure of coastal residents through seafood. The locations and type of sampling are shown in Figure 9.2.

Radiological data were also reported in a set of data tables in a surveillance report for selected rivers in Oregon (Oregon Department of Human Resources 1985). The surveillance covered from 1961 to 1983 for the Columbia, Snake, Willamette, and Klamath Rivers, and the coastal river system.

For the Columbia River, six monitoring stations were used (McNary Dam, The Dalles Dam, Rooster Rock, Cascade Locks, Beaver Army Terminal, and Astoria). The stations were not always monitored over the entire period; the sampling frequency was usually quarterly for each year but sometimes less frequent. Water, sediment, and algae were sampled. The surveillance measured gross alpha, gross beta, tritium, phosphorus-32, zinc-65, and chromium-51. However, not all samples were analyzed for each radionuclide, as shown by numerous NA (not analyzed) notations in the data tables. This pattern may indicate a selective sample analysis at each location, as the notations are not consistent for any radionuclide.

9.4 OREGON STATE UNIVERSITY

Oregon State University did not conduct any lengthy continuous monitoring programs, and therefore, no large or comprehensive databases were developed. Potentially useful data are found in small quantities, usually concerned with one type of fish or shellfish or with a particular location or phenomenon. Most of the reports and field programs represent limited or individual studies; many supported student dissertation work. In addition to dissertations, these studies also resulted in a number of journal articles and symposium proceedings. The basic purposes of the studies varied. There was a

TABLE 9.1. Range of Yearly Concentrations of Zinc-65 and Chromium-51 at Columbia River Locations for 1962 Through 1967. Activity levels are in pCi/L for water and in pCi/g (wet or drained weight) for sediment and algae. Number of samples is shown in parentheses.

Location	Range of Yearly Concentrations					
	Zinc-65			Chromium-51		
	Water	Sediment	Algae	Water	Sediment	Algae
John Day Dam						
1962-63	<35-81 (5)	1.2- 82.1 (9)	-(a)	2550-5280 (5)	-	-
1963-65	<35 (2)	2.5- 6.7 (2)	-	5790-10303 (2)	-	-
The Dalles Dam						
1963-65	-	0.7 (1)	-	-	-	-
1965-67	<35-144 (8)	0.9-116.5 (8)	-	837-3700 (8)	13.4-149.9 (8)	-
Rooster Rock						
1962-63	<35-74 (4)	0.2- 1.4 (4)	-	1670-3240 (4)	-	-
1963-65	<35-135 (22)	0.7- 36.0 (22)	-	725-5760 (22)	-	-
1965-67	<21-407 (24)	0.5- 31.3 (24)	-	<100-2520 (24)	4.4- 50.8 (24)	-
Goble						
1962-63	<35 (5)	5.2-230.0 (4)	-	930-1780 (5)	-	-
1963-65	<35-51 (23)	1.2- 99.6 (23)	18.7-225.0 (16)	350-5400 (23)	-	-
1965-67	<21-78 (24)	0.9-127.1 (24)	18.7-340.7 (14)	<100-1830 (24)	4.6-195.1 (24)	298.1-349.7 (14)
Beaver Army Terminal						
1962-63	<35-131 (7)	2.2-128.0 (6)	-	1300-2620 (5)	-	-
1963-65	<35-81 (23)	3.4-128.0 (21)	36.1-137.9 (16)	<100-4670 (23)	-	-
1965-67	<21-100 (24)	0.4-164.8 (24)	9.4- 65.0 (6)	<80-2184 (24)	4.6- 90.1 (24)	247.5-398.9 (6)
Astoria						
1962-63	<35 (7)	62.9 (1)	9.1-208.0 (10)	600-2250 (7)	-	-
1963-65	<35-37 (23)	1.5- 52.8 (19)	7.3- 82.7 (14)	<100-2720 (23)	-	-
1965-57	<21-60 (24)	1.9- 85.6 (23)	8.3- 48.9 (7)	<100-1390 (24)	10.1-552.9 (23)	239.0-304.2 (7)

(a) - Indicates no sampling.

TABLE 9.2. Range of Yearly Concentrations of Zinc-65 at Oregon Coast Locations for 1962 Through 1967. Activity levels are in pCi/L for water and in pCi/g (wet or drained weight) for sediment and shellfish. Number of samples is shown in parentheses.

Location	Range of Yearly Concentrations						
	Water	Sediment	Clams (Razor)	Clams (Soft Shell)	Crabs (Dungeness)	Mussels	Dysters
Seaside Beach							
1962-63	- (a)	-	5.4-36.2 (3)	-	-	-	-
1963-65			12.6-23.8 (3)	-	-	-	-
1965-67			7.7- 9.4 (3)	-	-	-	-
Cannon Beach							
1962-63	-	-	-	-	-	18.3-32.5 (3)	-
1963-65	-	-	-	-	-	20.0-28.8 (2)	-
1965-67	-	-	-	-	-	12.9-21.8 (2)	-
Nehalem River Jetty							
1962-63	-	-	-	-	-	10.1-32.0 (4)	-
1963-65	-	-	-	-	-	17.1-20.2 (3)	-
1965-67	-	-	-	-	-	12.8-21.4 (2)	-
Tillamook Bay							
1962-63	<35 (7)	0.1-0.9 (17)	-	0.4-5.3 (16)	10.0-27.4 (12)	-	7.1-37.0 (13)
1963-65(b)	<35 (22)	0.1-0.5 (19)	-	0.9-5.3 (21)	9.4-16.1 (21)	-	9.6-23.4 (23)
1965-67	<21-<35 (24)	<0.1-0.6 (22)	-	0.4-4.0 (18)	4.7-9.3 (20)	-	5.0-21.6 (23)
Yaquina Bay							
1962-63	-	-	-	-	13.9 (1)	4.9 (1)	3.2 (1)
1963-65	-	-	-	-	-	5.0-5.1 (2)	2.3-3.3 (2)
1965-67	-	-	-	-	2.2 (1)	2.1-3.9 (2)	1.1-2.0 (2)

(a) - Indicates no sampling.

(b) For 1963-65, gross beta activity levels in water ranged from 53 to 820 pCi/L.

S9112088.5

FIGURE 9.2. Columbia River and Coastal Sampling Locations for 1967-1977 Studies by the State of Oregon (Toombs and Paris 1978)

considerable amount of interest in the dispersion of the Columbia River plume along the coastal areas and seaward. Coupled with the plume dispersion was the affinity of certain commercial fishes for radionuclides. In at least one case, a secondary objective was to use the uptake of radionuclides to differentiate salmon of Asiatic stock from those of North American stock. Determining the decrease of radioactivity in fish and benthic marine life with distance from the river mouth was another objective.

For this review and discussion, the reports were grouped into two general locations: the Columbia River and estuary, and the coastline with adjacent ocean areas. Within these two basic groups, the reports and articles are separated into specific subjects for review purposes.

9.4.1 Columbia River and Estuary

There were several studies of Columbia River water, sediment, and the association of sediment with radionuclides, excluding the biological regime of the river. The studies extended as far upstream as The Dalles Dam and downstream to the river mouth, with most of the work being conducted in the estuary. The major radionuclides investigated were chromium-51 and zinc-65 over the period from 1963 to 1968. The study objectives were, collectively, to test methods of radionuclide measurement in water and sediment, to determine the chemistry of chromium-51 in river water (including its relationship with salinity), to identify the association of zinc-65 with sediment, and to determine the forms (dissolved or particulate) of chromium-51 and zinc-65 in transport. Most of the data are displayed in graphic form, with very few actual data points listed. The importance of these reports is in the study of specific processes and testing equipment. Reports and dissertations listing data or information of potential use for estimating dose are those by Cutshall and Osterberg (1964); Forster and Guthrie (1968); Hanson (1967); Jennings (1966); and Larsen, Renfro, and Forster (1968).

The studies investigating the uptake of radionuclides by biota were primarily concerned with using the presence of radionuclides to attain a better understanding of the physical, chemical, and biological phenomena taking place (radionuclide transfer through the food web), plus the effects of the Hanford production reactors being shutdown. The radionuclides in question were zinc-65, chromium-51, and, to a lesser extent, scandium-46 and manganese-54. The period of sampling extends intermittently from 1963 to 1971. The data in the reports are mostly presented in graphical form. The tabulated data that may be useful are sparse. The biota investigated included algae, rooted vegetation, various freshwater fish, crab, and starry flounder. The reports providing potentially useful data are by Johnson, Cutshall, and Osterberg (1966) and Renfro (1966).

9.4.2 Coastal and Ocean

Studies in the coastal zone and adjacent ocean areas investigated the directional movement and areal extent of the Columbia River plume and its effect on the food web. The radionuclides of interest were mostly zinc-65 and

chromium-51, with some interest in manganese-54 and certain radionuclides derived from fallout (from sources other than Hanford). The period extends from 1961 to 1970. The majority of the field studies focused on radionuclide concentrations in biota, with limited emphasis on water column concentrations and sediment. The decrease in activity levels with distance from the river mouth and coastal areas was of interest. Many of the species of biota sampled were noncommercial (e.g., starfish, sea cucumber, sea urchin, and others). The samples were gathered from various distances seaward along the coastal shelf and used as "tracers" for the various radionuclides.

Columbia River Plume

The dispersion of the Columbia River plume was studied using chromium-51 and salinity as tracers (Frederick 1967a, 1967b). Water samples were collected at the surface along Washington and Oregon coastal areas and analyzed for salinity and concentrations of chromium-51. The salinity and chromium-51 concentrations were then used to define the direction of dispersion of the winter and summer plumes, as shown in Figures 9.3 and 9.4, respectively. Within the boundary of the sampling pattern, the winter plume tended to hug the Washington coast and the summer plume dispersed along the Oregon coast in a slightly seaward direction.

A subsequent study by Pearcy and Forster (1968) analyzed the concentration of zinc-65 in biota (pelagic animals) relative to the plume boundaries. Samples of biota and water were collected near the surface to about a 5-meter (16.4-foot) depth. The results indicated that concentrations in the biota roughly corresponded to those in the plume water; that is, concentrations in both were higher near the origin of the plume at the mouth of the river.

Three earlier studies used zinc-65 in various species of biota to evaluate the decrease in radioactivity with distance from the mouth of the Columbia River (Osterberg 1962; Pearcy and Osterberg 1963; Mellinger 1966). Osterberg's (1962) results indicated that zinc-65 is a reasonably good tracer for the Columbia River plume, and a significant amount of information regarding radionuclide content of marine biota off the Oregon coast is provided in the report. The other two reports evaluate much smaller data sets.

S9112088.7

FIGURE 9.3. Winter Plume Pattern of February 1966 Based on Chromium-51 Activity (Frederick 1967b)

S9112088.8

FIGURE 9.4. Summer Plume Pattern of June 1965 Based on Chromium-51 Activity (Frederick 1967a)

Radionuclide Concentrations in Tuna and Salmon

A study of radioactivity in tuna livers was reported by Percy (1966) and Percy and Osterberg (1967). The radionuclides identified were zinc-65 and manganese-54. The period was from 1962 to 1966, with sampling locations extending from offshore northern Oregon southward to Baja California. Data tabulated by Percy and Osterberg (1967) indicate that, for the northern Oregon area, the ranges of zinc-65 concentrations in pCi/g of liver ash were 283 to 1050 pCi/g in 1963, 224 to 733 pCi/g in 1965, and 159 to 1067 pCi/g in 1966. For the southern Oregon area, the zinc-65 concentrations were 39.8 to 133 pCi/g in 1963 and 320 to 729 pCi/g in 1964. The one 1964 sample from the Baja California coast yielded a concentration of 46.4 pCi/g. The concentrations of manganese-54 were much less, ranging from 3.6 to 88.6 pCi/g in all areas considered.

Radioactivity in Pacific salmon was studied during 1964 and 1965. The initial findings were reported by Kujala (1966), followed by brief reports by Kujala and Forster (1968) and Forster and Loeffel (1968). The study by Kujala (1966) analyzed salmon for zinc-65 and manganese-54 at ten locations from Bristol Bay, Alaska, to Eureka, California. The range of concentrations for zinc-65 (in pCi/g dry weight) in chinook salmon was from 1.77 (Bristol Bay, Alaska) to 81.87 (Eureka, California). For sockeye salmon, the range was 0.86 (Bristol Bay, Alaska) to 5.52 (Barkley Sound, Canada). Sockeye salmon were not caught south of Canada. For coho salmon, the range was 3.61 (Cook Inlet, Alaska) to 59.28 (Depoe Bay, Oregon). All values were reported as averages for each location. The concentrations of manganese-54 were much less, with a maximum average for all salmon of 8.8 pCi/g dry weight.

The concentrations of zinc-65 in a "tagged" salmon study by Forster and Loeffel (1968) were given for two locations on Vancouver Island, Canada, and three locations off the coast of Alaska. Zinc-65 concentrations for the Vancouver Island salmon ranged from 339 to 518 pCi/g of ash; for the Alaska locations, from 7.3 to 450 pCi/g.

9.5 U.S. GEOLOGICAL SURVEY

During the early 1960s, the USGS commenced a series of studies to investigate the role of river sediment in the uptake and transport of radionuclides. The work was conducted in collaboration with the U.S. Atomic Energy Commission (USAEC) and Hanford Site contractors. The length of river involved extended from the vicinity of Pasco to Longview, Washington, just upstream of the Columbia River estuary. The overall objective of the studies was to determine quantitatively, to the extent possible, the transport and disposition of Hanford radionuclides in and along the Columbia River. Specific studies addressed the uptake and release of radionuclides by river sediment, the transport rates of certain radionuclides, and the inventory of radionuclides in riverbed sediments. This effort followed initial studies conducted at Hanford during the 1950s. The field work for the cooperative studies began in 1962 and continued intermittently to 1966. Preliminary reports were prepared during the 1960s by the USGS and General Electric Company until 1964 and by Pacific Northwest Laboratory after 1964. The final results were not published until the early 1970s.

9.5.1 Initial Studies at Hanford

During 1956 and 1957, an attempt was made to assess the magnitude of radionuclide uptake by sediment between the reactors and McNary Dam. The results of this effort were reported by Nielsen and Perkins (1957). Quantities of certain radionuclides in reactor effluent were measured at the reactors and downstream in an attempt to obtain an estimate of sediment uptake along the Hanford reach. Each reactor basin was sampled for a period of 2 to 3 days in November 1956 and March 1957. The river at Pasco was sampled daily starting 1 day later to allow for travel time. In the first 50 miles below the reactor outfalls, about half of the sodium-24 and copper-64; two-fifths of the arsenic-76; one-third of the phosphorus-32, zinc-65, and neptunium-239; and one-fifth of the chromium-51 were lost. The authors assumed that the net reduction in activity levels, after allowances for travel time (decay), resulted from uptake by sediment.

Nielsen and Perkins also conducted a depletion study for the reach between Pasco and Vancouver. Concentrations of phosphorus-32, chromium-51,

zinc-65, and neptunium-239 were measured in samples from twelve stations during 3 days in January 1957. Depletion of phosphorus-32 and chromium-51 averaged 15%, while depletion for zinc-65 was 65%.

The general conclusion drawn from these studies was that accumulation in sediment may account for a large part of the depletion of radionuclides from the water column, although Nielsen and Perkins' results were only representative of the conditions during sampling. Also, the conclusion that zinc-65 and chromium-51 were the major radionuclides in river sediments was based on one sample taken from behind McNary Dam. That sample contained 357×10^{-6} microcuries per gram ($\mu\text{Ci/g}$) of zinc-65, 87×10^{-6} $\mu\text{Ci/g}$ of chromium-51, and 5×10^{-6} $\mu\text{Ci/g}$ of cobalt-60.

9.5.2 U.S. Geological Survey Cooperative Studies

The results of these studies were first published in several Hanford documents and USGS open-file reports. These were mostly progress reports that included discussions of results to date and supporting data. The final results were published in a series of USGS professional papers that included all of the data from field sampling tabulated in appendixes.

Interim Hanford and USGS Reports

The first progress report presented preliminary results on the inventory of radionuclides in the river and the processes of sorption and release of radionuclides by sediment (Nelson, Perkins, and Nielsen 1964). The radionuclides of importance were zinc-65, chromium-51, scandium-46, cobalt-60, and manganese-54. Although the study team realized that core sampling was the most direct way to obtain an inventory of streambed radionuclides, they also knew that hundreds of samples would be required for a comprehensive survey. Therefore, an alternative approach was agreed upon that involved sampling water and sediment (suspended and bed material) at specific locations.

Beginning in July 1962 and continuing until September 1963, water, suspended sediment, and surficial streambed samples were collected several times per week at the highway bridges at Pasco, Hood River, and Vancouver. All water samples for a 2-week sampling period were composited and filtered, and the water and colloids were analyzed for cross-sectional and vertical

variations in radionuclide concentration. Transport rates calculated for chromium-51 and zinc-65 at Pasco and Vancouver indicated that up to 30% of the chromium-51 was lost to decay and very little tended to adsorb to sediment. However, much of the zinc-65 was taken up by sediment, and its resuspension during high river discharges (spring runoff) yielded higher than usual zinc-65 concentrations at Vancouver in May.

Analysis of the radionuclides associated with sediment particle sizes indicated that chromium-51 and scandium-46 were present in the finer sizes, while cobalt-60 and zinc-65 were more prevalent in the coarse fractions because of the presence of organic material. Studies were then conducted to determine the ionic form of certain radionuclides in reactor effluent and river water. Zinc-65, scandium-46, and manganese-54 were all determined to be predominantly cationic, and all associated freely with sediment downstream of McNary Dam. Chromium-51 was principally anionic in behavior, although a later article by Nelson et al. (1966) concluded that both hexavalent (anionic) and trivalent (cationic) forms of chromium occurred in the river and that chromium-51 in the cationic trivalent form was easily adsorbed by sediment.

The progress of radionuclide transport studies was reported by Nelson (1965). This report discusses the riverbed characteristics where sediment deposition would be expected to accumulate, such as behind dams, in slack-water areas along the river, and others. Nelson calculated the zinc-65 inventory from the reactor area to the Snake River to be about 1500 curies. This calculation was based on an estimate of zinc-65 concentration in coarse sand and gravel of 70 curies per square mile (Ci/mi^2) and an assumption that the extent of all types of deposition areas was reasonably estimated.

Nelson also investigated the uptake of radionuclides by biota. Algae scraped from rocks downstream of Pasco showed substantial concentrations of zinc-65 (21,800 dpm/g) and chromium-51 (78,000 dpm/g). The role of biota was not included in any of Nelson's calculations, based on the assumption that the biota did not amount to much volumetrically.

The results of the first year's sampling were presented in an open-file report of the USGS (Haushild et al. 1966). The report contains a compilation and analysis of all data collected during the 15-month sampling period. Data

are presented for radionuclide, sediment, and water measurements at Pasco, Hood River, and Vancouver, and at stations near the mouths of the Snake and Willamette Rivers. The radionuclides analyzed were sodium-24, scandium-46, chromium-51, manganese-54, cobalt-60, copper-64, zinc-65, zirconium/niohium-95, ruthenium/rhodium-103, lanthanum-140, cerium-141, and neptunium-239.

The report also discussed a special study, conducted during 2 days in March 1964, which indicated that radionuclide concentrations measured during the morning sampling period were 15% lower than the average concentration for 1 day because of significant diurnal fluctuations (hydropeaking) in discharge from Priest Rapids Dam. Differences in concentration values resulted from increases and decreases in travel time, which affected the decay of radionuclides between the reactors and Pasco. Variation in hydropeaking discharge would also modify the degree of dilution. These effects were particularly significant at low flow.

Estimates of the total quantity of radionuclides transported past each station indicated that approximately 277,000 curies were transported past Vancouver from January 1963 through September 1963. Chromium-51 was by far the largest volume (95%), with zinc-65 at 3%, and other radionuclides at 2%. Discharge of radionuclides in solution was highest during late winter and early spring, except for chromium-51. Maximum levels for chromium-51 were recorded for July through September. According to Haushild et al. (1966), these maxima either were anomalous or indicated a change in the practices controlling the addition of sodium dichromate to the reactor cooling water. The 277,000 curies transported past Vancouver represented about 77% of the total transported past Pasco. Decay was assumed to have reduced the Vancouver activity levels by about 19%; the remainder was assumed to have been retained in the riverbed. The authors recommended that future work investigate the variation in storage under changing hydrologic and hydraulic river conditions.

The second year of field work was carried out from January 1964 to January 1965. The first reported results came out in June 1965 (Perkins, Nelson, and Haushild 1965). The radionuclides considered in the study were scandium-46, chromium-51, manganese-54, cobalt-58, cobalt-60, zinc-65,

zirconium/niobium-95, ruthenium-106, antimony-124, and barium-140. Water samples were collected weekly at the highway bridge locations at Pasco and Vancouver, but less frequently at Hood River. From analysis of these samples, the sorption and transport of radionuclides with suspended sediment, the uptake of radionuclides by sediment (water column depletion), and the riverbed inventory (Pasco to Vancouver) were investigated. The radionuclides that tended to stay in solution were chromium-51, ruthenium-106, antimony-124, and barium-140. Those associated with sediments were scandium-46, manganese-54, cobalt-58, cobalt-60, zinc-65, and zirconium/niobium-95. Perkins, Nelson, and Haushild (1965) concluded that about 75% of the depletion occurred upstream of Hood River, probably behind McNary Dam.

Based on the data from the January 1964 to January 1965 sampling year, another report was prepared by Nelson et al. (1966). They used the January 1964 to January 1965 data, together with data from supplementary sampling, to determine depletion, deposition, scouring, and inventories of radionuclides and to estimate where and by what materials the radionuclides were held. Data from the 1962-1963 sampling year were not included. The radionuclides considered were scandium-46, chromium-51, manganese-54, cobalt-58, iron-59, cobalt-60, zinc-65, zirconium/niobium-95, ruthenium/rhodium-106, antimony-124, cesium-137, and barium/lanthanum-140. The reach of river considered was from Pasco to Vancouver.

Percent depletion was computed for each month and a radionuclide inventory estimated based on the assumption that radionuclide deposition for prior years was similar to that observed for 1964. The inventory calculated for the reach between Pasco and Vancouver was 11,000 to 38,000 curies. Of this total, 90% was zinc-65 and chromium-51.

Based on radionuclide concentrations in effluent and river water, the depletion, inventory, and amount of zinc-65 scoured (resuspended) in the reach from the reactors to McNary Dam were calculated. An average water column depletion of about 65% indicated a deposition of 5500 curies of zinc-65 in the reach. It was estimated that 30% of this was scoured and transported

downstream during the spring discharge of 1964. These results were based on weekly samples taken from May to October 1964, but were assumed to be representative for the entire year.

A special study that investigated the scouring of sediment from behind McNary Dam during high river discharges was begun in May 1965. Weekly sampling at the highway bridge at Umatilla (just downstream of the dam) was added to the sampling program. To quantify the amount of scour, ratios of zinc-65 (associated primarily with sediment) to chromium-51 (primarily in solution) in reactor effluent were compared with ratios in river water. The results indicated that the ratio of zinc-65 to chromium-51 increased by a factor of five during the high discharge period.

In 1969, Nelson and Haushild reported on an attempt to estimate the radionuclide inventory in bed sediments from the reach between the reactors and McNary Dam (Nelson and Haushild 1969). Two estimation methods were used. The first method was to determine the radionuclide concentrations in different bed sediments and estimate the extent of these sediment types within the reach. The second method was to use radionuclide data for Pasco and Umatilla to calculate the amount of radionuclides deposited behind McNary Dam. Both methods were based on the assumption of quasi-equilibrium between the river and the radionuclide input from the reactors, such that the number of curies of each radionuclide added to the streambed in a year balanced those lost by radioactive decay. This seems to be an assumption of constant input of specific radionuclides from the source, an assumption that is invalid. The results indicate that about 16,230 to 17,300 curies were contained in the bed sediments between the reactors and McNary Dam. From the bed sampling method, it was estimated that 1430 curies were stored between the reactors and Pasco, and an additional 14,800 curies were between Pasco and the dam. The radionuclide discharge data, which were valid only for that length of reach between Pasco and the dam, yielded an estimate of 17,300 curies.

USGS Professional Papers

The final reported results were published in 1973 and 1975 (Hubbell and Glenn 1973; Glenn 1973; Haushild et al. 1973; Haushild, Dempster, and Stevens 1975). According to these authors, the major radionuclides were zinc-65,

chromium-51, manganese-54, scandium-46, and cobalt-60. A number of other radionuclides were discussed, but the reports indicate that these other radionuclides contributed only a very small percentage of the radioactivity in the river.

The report by Glenn (1973) presents the results of a detailed study of riverbed sediments and the five major radionuclides sampled at nine locations from Pasco to the Columbia River estuary. The sampling period was April 21 to May 12, 1966. Radionuclide levels in sediment were related to cation-exchange capacity, and some rudimentary models were proposed. A considerable amount of information on sediment particle size and mineralogy is presented. Although the reported results do not include data useful for dose calculations, the report does present a discussion of the river sediment processes that is very useful for conceptualizing the water-sediment-contaminant complex.

Haushild et al. (1973) reported on the study of radionuclide transport from Pasco to Vancouver. The sampling period was from January 1964 to September 1966. The three sampling locations were Pasco, Umatilla, and Vancouver. Samples were obtained about three to four times per month and analyzed for both solute and particulate concentrations of the radionuclides of interest. The results are listed in an extensive set of tables in the report appendix. These water column concentrations were used to estimate radionuclide transport rates for eight radionuclides originating in the Hanford reactors. For 1964-1966, the combined discharges averaged 9190 curies per week at Pasco and 6630 curies per week at Vancouver. The approximate order of their abundance and the average percentage of the combined discharge were chromium-51 (96.4%), zinc-65 (2.5%), scandium-46 (0.5%), iron-59 (0.2%), antimony-124 (>0.1%), manganese-54 (>0.1%), cobalt-58 (<0.1%), and cobalt-60 (<0.1%).

A study of the radionuclide content of bed sediments (including core samples) from the Columbia River estuary was reported by Hubbell and Glenn (1973). Many measurements of gross gamma radiation were taken in situ at river cross sections spaced several miles apart. The cross sections began upstream at RM 64 and ended near the mouth at RM 2. These measurements were used in inferring concentrations of individual radionuclides based on a few

core samples that were counted in a laboratory. The results were used to calculate a rough estimate of the radionuclide inventory for the entire length of river sampled. The estimated total number of curies was 8700, with zinc-65 contributing 2100 curies (24%), chromium-51 contributing 5300 curies (61%), and the remainder divided about equally among scandium-46, manganese-54, cobalt-60, ruthenium-106, and zirconium/niobium-95. The results of the gamma measurements and core sample analysis are tabulated in the report appendix.

The final project report (Haushild, Dempster, and Stevens 1975) presents the results of bed sediment analysis and inventory calculations for the Columbia River from the Hanford reactors to Longview, Washington, below Bonneville Dam. A radionuclide inventory was computed for individual reaches and for the entire length under study. For 1965, the total radionuclide inventory was 37,000 curies, of which 60% was chromium-51 and 34% was zinc-65. The report's appendix lists radionuclide concentrations and particle-size distributions of surficial-sediment samples and gross gamma count rates of in situ sediment from the McNary reservoir to Longview, Washington. Also included are tables of core sample analysis results for reservoirs behind McNary, The Dalles, and Bonneville dams.

9.6 U.S. PUBLIC HEALTH SERVICE

Planning for this sampling program began in 1950 with meetings between USPHS staff, the USAEC, General Electric Company (Hanford), and the Columbia River Advisory Group (CRAG). The purpose of the program was to develop a water-quality database for determining what effects dams would have on the Columbia River. (Several dams were either proposed or under construction at that time.) Also, the database was to be used to evaluate the effect of radioactive effluent releases from Hanford on the normal stream purification factors. The study began in 1951 and continued into 1953. The results were published in a report by Robeck, Henderson, and Palange (1954).

The principal study area was from Priest Rapids at RM 400, just upstream of the Hanford reactors, to Paterson, Washington, at RM 278 and below McNary Dam. The program was initiated on July 23, 1951, when a few samples were collected for chemical analysis. Sampling for the complete chemical, biologi-

cal, bacteriological, and radiological program began on September 26, 1951, and continued into March 1953. Brief surveys were also conducted at other areas both upstream and downstream of Hanford. The downstream areas were the Bonneville reservoir; the area immediately upstream of Portland, Oregon; and the mouth of the Columbia River.

Ranges for sampling cross sections of the Columbia River were established in each study area and identified by river miles above the mouth of the river. Biological samples were generally collected in shallow water near the shorelines. Water samples were taken at three to ten points across the cross section; this was eventually standardized to five points at most locations. Water samples were collected four times a week. The ranges above and within the Hanford Site were sampled twice a month; those below the site were sampled weekly. Biological samples were collected mostly at semimonthly or monthly intervals. All radiological results were presented as gross beta activity densities. Maximum and average gross beta activities for various types of samples at RM 362, several miles below the reactors, are shown in Table 9.3. Other fish caught and analyzed for gross beta activity included salmon, carp, chub, squawfish, bass, sunfish, crappie, and sculpin.

TABLE 9.3. Maximum and Average Gross Beta Activities at River Mile 362. Activity levels are in pCi/L for water and in pCi/g for biota.

<u>Sample Type</u>	<u>Maximum</u>	<u>Average</u>
Water	19	6
Plankton	80,000	20,000
Filamentous algae	13,000	6,000
Caddisfly larvae	10,000	7,000
Juvenile fish (shiners)	9,000	1,300
Adult fish (suckers)		
Bone	5,000	1,200
Muscle	1,100	300

Activity levels were also determined for a limited number of bottom mud samples collected at RM 191 (The Dalles, Oregon), RM 167 (Hood River, Oregon), and RM 150 (Bonneville reservoir). Gross beta activity varied from 13 to $210 \times 10^{-7} \mu\text{Ci/g}$. Alpha activity varied from 1.8 to $5.9 \times 10^{-7} \mu\text{Ci/g}$.

The well-written USPHS report presents a considerable amount of field and laboratory data. All sampling and laboratory procedures are described and sample calculations are included. The data tables are easy to read and are listed in the appendixes. Many of the results are presented graphically. A graphical comparison between Hanford Site data and USPHS results (at RM 362) for gross beta activity levels in river water and juvenile fish (shiners) indicates that USPHS results were approximately 1.5 to 2 times higher than Hanford Site data. The numerical values of the Hanford data are not listed in the report, but both sets of results are described as being representative of a cross section location near the west shore (i.e., on the reactor side of the river). The data listed in this report will provide an independent database for comparison with Hanford monitoring results for the period 1951-1953. The year 1953 is probably the best for comparison purposes because of an extensive sampling program conducted downstream of McNary Dam.

10.0 SCREENING CALCULATIONS OF RADIATION DOSE AT VARIOUS DOWNSTREAM LOCATIONS FOR SEVERAL YEARS

To provide a scoping estimate of the magnitude of possible radiation doses to individuals along the Columbia River that resulted from aquatic releases of Hanford-originated radionuclides, a set of screening dose estimates for maximally and typically exposed individuals was prepared. The basis of these estimates is the environmental monitoring data described in Sections 7 and 9 of this report. Estimates are provided here, by major exposure pathway and radionuclide, for five representative locations along the Columbia River for the years 1964 through 1966. The five locations are illustrated in Figure 10.1.

The choice of the 1964 through 1966 time period for these screening dose estimates was based on several factors. First, before the 1960s, it was not easy to discriminate among the various radionuclides represented by gross beta measurements, so it was important to select a period in the 1960s. Second,

FIGURE 10.1. Locations for Which Dose Estimates Were Made

data from various sources were available for the specific years selected. Third, the selected time period is of interest because it includes a year, 1964, when all reactors were operating and years when some reactors were shut down (see Figure 5.2).

10.1 RADIONUCLIDE CONCENTRATIONS IN RIVER WATER

The literature and databases summarized in Sections 7 and 9 of this report provided information on the radionuclide content of the river water. Monitoring was routinely performed at a few locations along the river; these monitoring points provide the most consistent and coherent set of available data. The Hanford monitoring database was queried for all samples, and the data for the Richland Pump house, McNary Dam, and Bonneville Dam were summarized into annual means. Data were selected for the years 1964, 1965, and 1966 to match other available sources. Data for Astoria, Oregon, for the same period are available from the Oregon State Board of Health (Toombs and Cutler 1968). Data for all radionuclides reported for the selected years are presented in Table 10.1. For Richland and the McNary reservoir, the data are shown with annual mean values derived from the monthly numbers used in HEDR Phase I calculations, as reported by Richmond and Walters (1991). In some instances, the Phase I data appear to be more complete than the monitoring data, because these values reflect Phase I modeling calculations based on effluent monitoring data and are not environmental monitoring data per se.

The radionuclide concentration values in Table 10.1 show predictable behavior; the concentration uniformly decreases with increasing distance downstream. This is most noticeable in the short-lived radionuclides, such as sodium-24, neptunium-239, or iodine-131. It is less apparent in longer-lived radionuclides, such as chromium-51 or zinc-65, for which most of the decrease can be attributed to dilution by inflowing tributaries or to uptake by and deposition in bed sediments.

Gaps in the data are noticeable; not all radionuclides are reported for all years or locations. Extrapolation was used temporally, but not spatially, to calculate dose estimates. Thus, for any one location, if a concentration of a radionuclide was reported for 1 or 2 years but not at another time,

TABLE 10.1. Annual Mean Radionuclide Concentrations at Various Locations in Columbia River Water

Location/ Radionuclide	Radionuclide Concentrations (pCi/L)		
	1964	1965	1966
<u>Richland^(a)</u>			
Sodium-24	3500/3600	3100/3100	2600/2600
Phosphorus-32	300/170	140/120	140/140
Scandium-46	-/- ^(b)	-/-	30/-
Chromium-51	12000/8900	7000/5200	2600/3500
Manganese-56	-/2800	-/-	290/-
Copper-64	-/5100	-/770	-/1400
Zinc-65	450/250	180/240	200/210
Arsenic-76	1200/1200	1000/1100	420/740
Strontium-90	1/-	1/-	1/-
Iodine-131	19/-	10/-	18/-
Neptunium-239	-/2100	-/1200	-/880
<u>McNary^(a)</u>			
Sodium-24	-/150	-/-	-/-
Phosphorus-32	70/80	50/60	80/80
Chromium-51	3500/4100	2300/2500	1850/1850
Copper-64	-/20	-/-	-/-
Zinc-65	77/77	68/75	61/62
Iodine-131	6.7/-	3.9/-	7.1/-
Neptunium-239	-/470	-/300	-/190
<u>Bonneville^(c)</u>			
Phosphorus-32	28	23	23
Chromium-51	2400	1700	1300
Zinc-65	63	70	43
Iodine-131	5	3	3
<u>Astoria^(d)</u>			
Phosphorus-32	-	18	11
Chromium-51	1500	930	1600
Zinc-65	<35	<35	<35
Zirconium/Niobium-95	<13	<13	<13
Ruthenium-103,106	<10	<10	<10

(a) Information on the left from Hanford Monitoring Database; on the right from Richmond and Walters (1991).

(b) Dashes (-) indicate no data available.

(c) Source: Hanford Monitoring Database.

(d) Source: Toombs and Cutler (1968).

the highest value reported was used as an estimate of the missing value. This conservative assumption was made to ascertain whether radionuclides that were not monitored could have made significant contributions to dose in the other years. Additionally, the larger of the two values was used in all calculations as an additional conservative measure.

Because the HEDR Phase I report (Richmond and Walters 1991) was one of the sources of information for the analysis, the input concentrations in this screening study are compatible with the Phase I inputs (well within a factor of two). The inputs for the Richland location are also generally compatible (within factors of two to four) with the concentrations used in the HEDR "dominant radionuclides" study (Napier 1991).

10.2 RADIONUCLIDE CONCENTRATIONS IN AQUATIC BIOTA

Large amounts of data are available for fish in the Hanford reach of the Columbia River and immediately downstream. Many of these data were used in the Phase I dose calculations. However, monitoring data for biota are much more limited for areas below McNary Dam. Available data for biota in the Columbia River estuary near Astoria, Oregon, are presented in Table 10.2, for the same period as the water data in Table 10.1. The data are much more limited than the data from the Hanford reach, and only a few radionuclides and a few species are represented. Measurements of Pacific coastal biota relevant to the selected locations are summarized in Table 10.3. Because the North Head area is at the mouth of the Columbia River across from Astoria, data from North Head were used in the calculation to represent the Astoria area.

Essentially no data are available on radionuclide concentrations on salmon returning from the Pacific Ocean to the Columbia River to spawn (Section 9.4.2). This is partly because runs of salmon in the Columbia were greatly depleted by the mid-1960s (Becker 1985) and partly because the concentrations of Hanford-originated radionuclides in the salmon tended to be lower than in the resident fish (Jaquish and Bryce 1989). However, the limited data shown in Table 10.3 indicate that Hanford-related radionuclides in Pacific salmon increased near the mouth of the Columbia. The concentrations of zinc-65 and manganese-54 in salmon caught off the Oregon coast near

TABLE 10.2. Radionuclide Concentrations in Selected Monitored Biota in the Columbia River Estuary at Astoria

<u>Biota/Radionuclide</u>	<u>Radionuclide Concentrations (pCi/kg)</u>		
	<u>1964</u>	<u>1965</u>	<u>1966</u>
<u>Flounder</u> ^(a)			
Zinc-65	19,000	17,000	14,000
<u>Sculpin (incl. bone)</u> ^(a)			
Zinc-65	10,000	9,400	7,400
<u>Crab Muscle</u> ^(b)			
Zinc-65	37,000	37,000	37,000
Manganese-54	20	20	20
Chromium-51	300	300	300
<u>Freshwater Clams</u> ^(c)			
Zinc-65		250	250
Manganese-54		17	17
Cobalt-60		1	1

(a) Annual averages reported by Renfro, Forster, and Osterberg (1972).

(b) Tennant and Forster (1969).

(c) Johnson, Cutshall, and Osterberg (1966).

Astoria and Depoe Bay approached those calculated for fish resident in the river at the Astoria location. The entry in Table 10.3 for herring indicates that concentrations in the salmon were essentially the same as in their primary food supply. Because many of the radionuclides of interest are obtained mainly from food-web uptake (Poston and Klopfer 1986) and because the salmon do not eat significantly once they enter the river (Poston and Klopfer 1986), their river-entry concentration has been assumed in this analysis to remain essentially constant as the fish migrate upstream past Astoria to Bonneville.

TABLE 10.3. Reported Average Radionuclide Concentrations in Selected Pacific Coastal and Ocean Biota

Biota-Location/Radionuclide	Radionuclide Concentrations (pCi/kg)					
	1963	1964	1965	1966	1967	1968
Mussels-North Head ^(a)						
Zinc-65	120,000	-	-	-	38,000	24,000
Mussels-/Westport ^(a)						
Zinc-65	52,000	-	-	10,000 ^(b)	13,000 ^(c)	10,000
Dover Sole-Ocean ^(d)						
Chromium-51	-	100	-	-	-	-
Zinc-65	-	200	-	-	-	-
Oysters-Willapa Bay ^(e)						
Zinc-65	100,000 ^(f)	62,000	52,000	-	-	-
Chinook Salmon-Astoria ^(g)						
Manganese-54	-	8	-	-	-	-
Zinc-65	-	10,000	-	-	-	-
Coho Salmon-Depoe Bay ^(g)						
Manganese-54	-	170	-	-	-	-
Zinc-65	-	12,000	-	-	-	-
Herring-Depoe Bay ^(g)						
Manganese-54	-	44	-	-	-	-
Zinc-65	-	10,000	-	-	-	-

- (a) Seymour (1970).
 (b) Mellinger (1966).
 (c) Larsen (1970).
 (d) Jonsson and Seymour (1965).
 (e) Seymour (1966).
 (f) Naidu (1963).
 (g) Kujala (1966).

Although it is often an oversimplification, the concentration of radionuclides in fish can be related to the concentrations of those radionuclides in water by means of a bioaccumulation factor (pCi/kg of fish per pCi/L of water). For this screening analysis, the bioaccumulation factors developed for Phase I were used where available. Because the HEDR Phase I model used actual distributions based on the numerous available measurements of zinc-65 and phosphorus-32 in fish (PNL 1991), bioaccumulation factors for these and for the radionuclides not considered in Phase I were taken from analyses specific to Hanford and the Columbia River (ERDA 1975). The values used are

shown in Table 10.4. The values used represent an average bioaccumulation across numerous species commonly caught and eaten. Each of these values is discussed below.

The bioaccumulation factor for arsenic-76 is based on measurements from the Hanford reach of the Columbia River in 1961. This factor is based on measurements from 28 fish caught between April and December. The calculated bioaccumulation ranged from 30 to 910, with a mean of about 200. For reference, the current default used in Hanford Site-related dose calculations is 300. The radiation dosimeter software system, GENII, was used for these calculations (Napier et al. 1988).

The bioaccumulation factor for cobalt-60 is taken from ERDA (1975), based on Hanford conditions. Recommendations for cobalt-60 range from 27 to 320, based on water conditions. The value currently recommended by Poston and Klopfer (1986) is 330 for mesotrophic systems (such as the Columbia River). The current GENII default is also 330.

TABLE 10.4. Freshwater Fish Bioaccumulation Factors
Used in this Analysis

<u>Element</u>	<u>Bioaccumulation Factor</u>
Arsenic	200.0 ^(a)
Cobalt	330.0 ^(b)
Chromium	5.0 ^(a)
Copper	10.0 ^(a)
Iodine	15.0 ^(b)
Manganese	70.0 ^(a)
Sodium	1.0 ^(a)
Neptunium	25.0 ^(a)
Phosphorus	170.0 ^(b)
Ruthenium	10.0 ^(b)
Scandium	2.0 ^(b)
Strontium	30.0 ^(b)
Zinc	64.0 ^(b)
Zirconium	330.0 ^(b)

(a) HEDR Phase I input (PNL 1991).

(b) ERDA (1975).

The factor for chromium-51 is based on Hanford data from 1961. The data indicate that most fish in the Columbia River at that time were below the detection limit for chromium-51. These data provide a range of 0.8 to 5. The value of 5 was conservatively chosen at the top of the range. The current GENII default is 20.

Copper-64 has a half-life of only 12.8 hours. As a toxic metal, it would be expected to accumulate in the liver, not flesh. Hanford data indicate a value of <10 for copper-64. The value for stable copper is much higher; the GENII default is 2500.

Literature recommendations for iodine-131 bioaccumulation range from 15 (Thompson et al. 1972) to 40 (Vanderploeg et al. 1975). A value of 15 was derived for Hanford (ERDA 1975). The current GENII default is 50, derived from Poston and Klopfer's (1986) recommendation for iodine-129.

Manganese-56 has only a 2.6-hour half-life. Published values for stable or long-lived manganese reach as high as 1000. Vanderploeg et al. (1975) developed a relationship, derived from filtered Columbia River data by Silker (1964). Poston^(a) suggests a value of no more than 70 for unfiltered water. The current GENII default is based on Poston and Klopfer's (1986) generic recommendation of 400.

Sodium-24 has a short half-life compared with its biological turnover time. This suggests a value of 1.0, based on rapid isotopic distribution within fish tissue fluids. Longer-lived sodium is homostatically regulated and independent of water sodium concentrations. A generic value of 70 is used as a default in GENII.

The bioaccumulation factor for neptunium-239 is based on Hanford measurements. Data from 1961 range from a minimum of about 4 to a maximum of about 40. A seasonally weighted average of 25 was used. For longer-lived neptunium, Poston and Klopfer (1986) recommend values of 50 for piscivorous species, 250 for planktivorous species, and 2500 for bottom feeders. The current GENII default is 500, based on this recommendation.

(a) Personal communication to B. A. Napier from T. M. Poston, May 1990.

Actual fish monitoring data were used in Phase I calculations for phosphorus-32. The reported values of bioaccumulation range from 20 to 100,000. Many studies have reported on the Columbia River (e.g., Foster, Soldat, and Essig 1966). Soldat derived a recommended value of 170 for the Columbia River (ERDA 1975). The current GENII default for unspecified rivers is 1500 (Poston and Klopfer 1986).

Few data exist to defend a bioaccumulation value for ruthenium-106. The reported values range from 0.1 to 170. Thompson et al. (1972) recommend 10, which was used by ERDA (1975) for the Columbia River. The current GENII default is 100.

Most sources indicate a bioaccumulation factor for stable scandium of between 20 and 104. The scandium-46 value used by ERDA (1975) was 2. Poston and Klopfer (1986) indicate that this may be a reasonable value for the Columbia River. The current GENII default is 100. A sensitivity analysis for scandium-46 indicates that even using a value of 100, the dose from scandium-46 varies by only a factor of four; most exposure to scandium-46 comes from pathways other than fish. This 50-fold increase of the scandium-46 bioaccumulation factor would result in less than a 1% change in the total dose to the maximally exposed individual.

Bioaccumulation of strontium-90 is directly proportional to the amount of stable calcium in water. The bioaccumulation factor can range over three to four orders of magnitude. The value recommended by Poston and Klopfer (1986) for generic fresh waters is 50; this is the GENII default. The value of 30 used in the analysis is from ERDA (1975); this is not significantly different.

Bioaccumulation for zinc-65 can range from 100 to 2500. Values reported for the Columbia River range from 4 to 40, with single values of 132 and 155 (Poston and Klopfer 1986). The current GENII default is 500. The value of 64 used in ERDA (1975) approximates the results of using 1964-1966 data in Phase I. Therefore, a value of 64 was used in this analysis.

Zirconium-95 bioaccumulation ranges from 40 to 460 in fresh water (Poston and Klopfer 1986). Poston and Klopfer recommend a value of 200. The default in GENII is 200. The ERDA (1975) value of 330 for the Columbia River is near the upper end of the range.

10.3 EXPOSURE SCENARIOS AND PATHWAYS USED IN SCREENING CALCULATIONS

Humans living along the river may have been exposed to radionuclides carried in the water, deposited in the sediments, accumulated in fish and other aquatic foods, and irrigated onto soils and crops. For any one of these pathways, a wide range of exposures may have occurred to different individuals, depending on their habits and activities. For the purpose of screening the possible magnitude of exposures to individuals in the population, two basic "types" of individuals were postulated. The first is a "maximally exposed" individual, who would have relatively large exposures from each of the pathways; the second is a more "typical" individual, whose exposures would be more representative of the average population. The parameters used in the calculation for the maximally exposed individual are shown in Table 10.5, and those for the typical individual are shown in Table 10.6.

The external exposure pathways are those for which the individual would merely have been in a location where radioactive contamination was present to receive a dose of external radiation. These include boating in contaminated water, swimming in contaminated water, standing along a shoreline where radionuclides were in contact with the sediments, or standing in fields or lawns irrigated with the contaminated water. The major determinant of the dose received was the concentration in the water or soils and the amount of time spent there. Irrigation is assumed only for areas upstream of Bonneville Dam in these calculations.

For the drinking water pathway, it was assumed that the individual consumed water taken from the Columbia River. This pathway is applicable only along the river, because salt water precludes drinking from the ocean bays. In these analyses, all water has been assumed to be treated prior to

TABLE 10.5. Exposure Pathway Parameters Assumed for Maximally Exposed Individuals

<u>Pathway</u>	<u>Assumed Value</u>	<u>Notes</u>
<u>External Exposure</u>		
Boating	500 h/yr	
Swimming	100 h/yr	
Shoreline	500 h/yr	
Irrigated Soil	4000 h/yr	Richland/McNary/Bonneville only
<u>Drinking Water Consumption</u>		
Drinking Water	730 L/yr	River locations only; HEDOP ^(a) default
Treatment	Yes	
<u>Aquatic Food Consumption</u>		
Resident Fish	40 kg/yr	Considered at all locations
Anadromous Fish	100 kg/yr	Astoria/Bonneville only
Crabs	20 kg/yr	Astoria only
Mussels	20 kg/yr	Astoria only
Clams	20 kg/yr	Astoria only
Oysters	20 kg/yr	Willapa Bay only
Holdup times	none	Consumed same day as caught
<u>Irrigation</u>		
Irrigation Rate	40 in./yr	Richland/McNary only
Foliar Interception Fraction	0.25	Sprinkler irrigation
Vegetable Diet		HEDOP defaults
Milk Diet		HEDOP defaults
<u>Inhalation</u>		
Resuspension Mass Loading	100 $\mu\text{g}/\text{m}^3$ ^(b)	
Exposure	8760 h/yr	

(a) Hanford Environmental Dose Overview Panel (McCormack, Ramsdell, and Napier 1984).

(b) Micrograms per cubic meter.

consumption. The drinking water cleanup factors known to be applicable to the Richland and Pasco alum flocculation cleanup processes have been used for all locations. Only the amount of water consumed has been allowed to vary.

TABLE 10.6. Exposure Pathway Parameters Assumed for Typical Individuals

Pathway	Assumed Value	Notes
<u>External Exposure</u>		
Boating	5 h/yr	HEDOP ^(a) default
Swimming	10 h/yr	HEDOP default
Shoreline	17 h/yr	HEDOP default
Irrigated Soil	100 h/yr	4 h/weekend in summer on lawns; Richland/McNary
<u>Drinking Water Consumption</u>		
Consumption	440 L/yr	River locations only; HEDOP default
Treatment	Yes	
<u>Aquatic Food Consumption</u>		
Resident Fish	1 kg/yr	
Anadromous Fish	5 kg/yr	Astoria/Bonneville only
Crabs	2 kg/yr	Astoria only
Mussels	2 kg/yr	Astoria only
Clams	2 kg/yr	Astoria only
Oysters	2 kg/yr	Willapa Bay only
Holdup times	none	Consumed same day as caught
<u>Irrigation</u>		Not used (except lawns)
<u>Inhalation</u>		
Resuspension Mass Loading	10 $\mu\text{g}/\text{m}^3$	
Exposure	8760 h/yr	

(a) Hanford Environment Dose Overview Panel (McCormack, Ramsdell, and Napier 1984).

The assumed rates of nonmigratory fish, salmon, and shellfish consumption are given in Tables 10.5 and 10.6. These values are somewhat arbitrary, but are believed to be reasonable estimates for maximal and typical consumers, respectively. Although some groups may consume more than the quantities shown, their doses may be derived from direct multiples of those presented. An example of this is presented in Section 10.5.

Irrigation with water taken from the Columbia River below Hanford would not have been a pathway of exposure for many people. Most irrigation water in

the area was and is taken either from the Columbia River upstream at Grand Coulee Dam or from tributary streams. The largest area irrigated with downstream water during the period of reactor operations was the Riverview area near Pasco, Washington. A few thousand people were directly affected in the Riverview area. The assumed irrigation rate of 40 inches per year is typical for Columbia Basin farming practices (McCormack, Ramsdell, and Napier 1984). Consumption rates used in the calculations are those of the Hanford Environmental Dose Overview Panel (McCormack, Ramsdell, and Napier 1984).

Inhalation was a secondary pathway, but one that may have resulted from dust blowing off irrigated land. People are assumed to have been exposed to this source all year.

The pathways and parameters selected for the maximally exposed individual for this screening analysis are consistent with the HEDR "dominant radionuclides" study (Napier 1991). This screening analysis extends the dominant radionuclides report by the addition of the typical individual.

10.4 EXPOSURE PATHWAY MODELS

The calculations were performed using the publicly available GENII computer software system (Napier et al. 1988). The GENII system is composed of seven linked computer programs and their associated data libraries. The computer programs are of three types: user interfaces, internal and external dose factor generators, and environmental dosimetry programs. All steps of code development have been documented.

10.5 DOSE RESULTS

The doses estimated for the maximally exposed individual at five locations for the period 1964 through 1966 are summarized in Table 10.7. The table presents the total estimated annual dose in rem, the radionuclide or radionuclides that contributed the largest percentage of the total dose, and the exposure pathway that contributed the largest percentage of the total dose. As indicated in the table, the most important exposure pathway was consumption of nonmigratory (resident) fish. The radionuclides contributing most to dose varied slightly with distance downstream. At Richland, the most important

TABLE 10.7. Summary of Annual Doses from Columbia River Pathways for the Maximally Exposed Individual

Location/Detail	Dose (rem)		
	1964	1965	1966
Richland	1.6×10^{-1}	1.0×10^{-1}	8.2×10^{-2}
Dominant Radionuclide/%	Arsenic-76/34%	Arsenic-76/49%	Arsenic-76/40%
	Zinc-65/25%	Zinc-65/21%	Zinc-65/22%
Dominant Pathway	Fish/86%	Fish/89%	Fish/87%
McNary	1.7×10^{-2}	1.4×10^{-2}	1.4×10^{-2}
Dominant Radionuclide/%	Zinc-65/39%	Zinc-65/46%	Phosphorus-32/48%
	Phosphorus-32/39%	Phosphorus-32/36%	Zinc-65/39%
Dominant Pathway/%	Fish/64%	Fish/64%	Fish/65%
Bonneville (with salmon)	2.1×10^{-2}	2.1×10^{-2}	2.0×10^{-2}
Dominant Radionuclide/%	Zinc-65/90%	Zinc-65/90%	Zinc-65/90%
Dominant Pathway/%	Fish/95%	Fish/95%	Fish/95%
Astoria (with salmon)	4.0×10^{-2}	4.0×10^{-2}	4.0×10^{-2}
Dominant Radionuclide/%	Zinc-65/93%	Zinc-65/93%	Zinc-65/93%
Dominant Pathway/%	Fish + Seafood/99%	Fish + Seafood/99%	Fish + Seafood/99%
Willapa Bay	1.6×10^{-2}	1.3×10^{-2}	1.3×10^{-2}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%
Dominant Pathway/%	Oysters/100%	Oysters/100%	Oysters/100%

radionuclides were arsenic-76 and zinc-65, with a significant contribution also from phosphorus-32. At locations farther downstream, the relative importance of zinc-65 increased and that of the other radionuclides decreased. This change is a direct result of the radioactive decay of the shorter-lived materials. The calculated doses also decreased as distance downstream increased, reflecting the decay and dilution of the radionuclides listed in Table 10.1. However, the dose increases slightly at Bonneville with the addition of the salmon pathway and at Astoria with the addition of shellfish. Details relating to these additional pathways are shown in Tables 10.8 and 10.9.

The total doses reported include contributions from the external pathways of swimming, boating, and exposure to contaminated soils and sediments, as well as internal doses from drinking water, eating fish, and eating irrigated foods. The notation "fish" in Table 10.7 represents nonmigratory freshwater fish. The notation "seafood" includes crabs, mussels, and freshwater

TABLE 10.8. Detail of Annual Doses Reported in Table 10.7 for Astoria

	Dose (rem)		
	1964	1965	1966
Standard Pathways	4.4×10^{-3}	4.3×10^{-3}	4.0×10^{-3}
Dominant Radionuclide/%	Zinc-65/41%	Zinc-65/42%	Zinc-65/45%
Dominant Pathway/%	Fish/90%	Fish/90%	Fish/90%
Crab/Mussel/Clams	2.0×10^{-2}	2.0×10^{-2}	2.0×10^{-2}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%
Salmon	1.6×10^{-2}	1.6×10^{-2}	1.6×10^{-2}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%

TABLE 10.9. Detail of Annual Doses Reported in Table 10.7 for Bonneville

	Dose (rem)		
	1964	1965	1966
Standard Pathways	5.1×10^{-3}	5.0×10^{-3}	3.7×10^{-3}
Dominant Radionuclide/%	Zinc-65/55%	Zinc-65/64%	Zinc-65/51%
Dominant Pathway/%	Fish/93%	Fish/93%	Fish/94%
Salmon	1.6×10^{-2}	1.6×10^{-2}	1.6×10^{-2}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%

clams, as well as salmon. In Tables 10.8 and 10.9, the notation "standard pathways" includes the external pathways plus ingestion of resident fish.

The doses reported in Table 10.7 reflect an assumed consumption rate of 40 kg/yr of resident fish. However, concerns have been expressed that a few individuals may have subsisted on a diet consisting almost entirely of fish. For the extreme case of an individual eating 1 kg/d of fish, a total effective dose of about 1.3 rem/yr may be derived for the year 1964 at Richland. This is probably an upper bound, because a significant fraction of this dose is caused by short-lived radionuclides (e.g., arsenic-76); the decay that occurs between catching and eating the fish has not been accounted for in this calculation.

The doses estimated for typical individuals at the five locations for the period 1964 through 1966 are summarized in Table 10.10. Because the typical individual would have eaten very little fish, the doses were dominated by the drinking water pathway. From 70% to 80% of the dose estimated for the

TABLE 10.10. Summary of Annual Doses from Columbia River Pathways for the Typical Individual

Location/Detail	Dose (rem)		
	1964	1965	1966
Richland	7.8×10^{-3}	5.1×10^{-3}	4.2×10^{-3}
Dominant Radionuclide/%	Arsenic-76/26%	Arsenic-76/35%	Arsenic-76/29%
	Neptunium-239/22%	Neptunium-239/20%	Sodium-24/18%
	Zinc-65/19%		Neptunium-239/17%
Dominant Pathway/%	D. Water/72%	D. Water/68%	D. Water/70%
McNary	1.0×10^{-3}	7.7×10^{-4}	7.4×10^{-4}
Dominant Radionuclide/%	Neptunium-239/40%	Neptunium-239/32%	Phosphorus-32/31%
	Zinc-65/27%	Zinc-65/32%	Zinc-65/28%
Dominant Pathway/%	D. Water/87%	D. Water/83%	D. Water/80%
Bonneville (without salmon)	3.8×10^{-4}	3.5×10^{-4}	2.6×10^{-4}
Dominant Radionuclide/%	Zinc-65/53%	Zinc-65/66%	Zinc-65/54%
Dominant Pathway/%	D. Water/70%	D. Water/70%	D. Water/70%
Astoria (without salmon)	2.8×10^{-4}	2.7×10^{-4}	2.6×10^{-4}
Dominant Radionuclide/%	Zinc-65/46%	Zinc-65/48%	Zinc-65/50%
	Phosphorus-32/19%	Phosphorus-32/20%	D. Water/71%
Dominant Pathway/%	D. Water/70%	D. Water/70%	
Astoria shellfish	2.0×10^{-3}	2.0×10^{-3}	2.0×10^{-3}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%
Ocean salmon	7.9×10^{-4}	7.9×10^{-4}	7.9×10^{-4}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%
Willapa Bay	1.6×10^{-3}	1.3×10^{-3}	1.3×10^{-3}
Dominant Radionuclide/%	Zinc-65/99%	Zinc-65/99%	Zinc-65/99%
Dominant Pathway/%	Oysters/100%	Oysters/100%	Oysters/100%

typical individual is a result of drinking water derived from the Columbia River. The cities of Richland, Pasco, and Kennewick were and are the main users of Columbia River water for public drinking supplies. A few small towns downstream of these cities also use the Columbia River as a source of water. Because the drinking water pathway dominated, the spectrum of important radionuclides was slightly different from that for the maximally exposed individual: arsenic-76, neptunium-239, and zinc-65 were important in the regions

nearest Hanford. As the short-lived arsenic-76 and neptunium-239 decayed, the zinc-65 and phosphorus-32 remained at the downstream locations.

The screening doses presented in Tables 10.7 and 10.10 are generally compatible with the HEDR Phase I results for Richland and McNary (PNL 1991), as well as with results of the HEDR "dominant radionuclide" study for Richland (Napier 1991). All results for the typical individual are within a factor of two of the median doses reported in the Phase I Columbia River Summary Report (PNL 1991), although they tend to be somewhat lower. All results for the maximally exposed individual are within a factor of two of the 95th percentile values reported in the Phase I report, although the results tend to be somewhat higher. Another similarity with the earlier calculations is that the organ receiving the largest dose would have been the gastrointestinal tract, indicating the importance of the short-lived radionuclides that are not readily absorbed by the body. These similarities are expected because the water concentrations and bioaccumulation factors used are very close to the Phase I estimates.

The calculated external pathway doses are always quite small. The largest doses were for the maximally exposed individual in Richland. The sum from boating, swimming, shoreline exposure, and exposure to irrigated fields would always have been less than 1 mrem to the typical individual; external doses would have been less than 10% of the total for the maximally exposed individual.

The dose from ingestion of irrigated crops would also have been small. The largest calculated dose to a maximally exposed individual, who consumed nearly 500 kilograms of vegetables and milk, was less than 10% that of eating 40 kilograms of resident fish. An individual who was typical in all respects except for irrigating a garden and eating predominantly from it would have received, at most, double the dose from the drinking water pathway.

Dose from inhalation of resuspended soil contaminated by irrigation water was negligible: less than one-one millionth of the dose from eating fish.

Although calculations for drinking water give a dose that is only 5% to 10% as large as that calculated for the maximally exposed individual from eating fish, drinking water would have been the major pathway for the majority of the population, represented by the typical individual. The drinking water doses estimated here for "typical" individuals are all within a factor of two of those reported in the 1964, 1965, and 1966 Hanford Site annual reports (Foster and Wilson 1964; Foster, Soldat, and Essig 1966; Honstead, Essig, and Soldat 1967).

The dose from eating salmon migrating up the river would be about the same as the dose from eating an equivalent amount of resident fish at Astoria. Because the salmon did not eat, and therefore presumably did not greatly increase in radionuclide concentration as they migrated upstream, they would probably have contributed a pound-for-pound dose that is less than that from resident fish at locations closer to Hanford. This assumption will have to be investigated further before any definitive statements can be made about the dose resulting from salmon consumption above Bonneville Dam.

The doses reported in Tables 10.8 and 10.9 for consumption of salmon are based on an assumed consumption rate of 100 kg/yr. For the possible case of a subsistence fisherman consuming up to 1 kg/d, the dose would be about 0.7 rem/yr from zinc-65. It is possible that other radionuclides could contribute incrementally to this upper-bound estimate for locations in the river nearer to Hanford.

As described in Section 9.4, the plume from the Columbia River out into the Pacific Ocean tended to travel in a northerly direction along the Washington coast in the winter and in a southerly direction and slightly seaward along the Oregon coast in the summer (Kujala 1966). For this reason, the concentrations of radionuclides in shellfish on the Washington coast (e.g., Willapa Bay) tended to be higher than those along the Oregon coast. The major radionuclide in Willapa Bay oysters was zinc-65. The dose resulting from consumption of the oysters would have been directly proportional to the quantity of oysters consumed. For a nominal amount of 20 kg/yr, the dose would have been around 15 mrem. A dose of the same general magnitude was attributable to consumption of 20 kg/yr each of mussels, crabs, and clams from the estuary of

the Columbia River. Additional information concerning the diets of groups who consume significant quantities of these foods is needed before definitive statements can be made about the doses they received. The results indicate that consumption of estuary shellfish may have been more important than consumption of resident fish, depending on the relative rates of intake.

10.6 SCALING MID-1960s DOSES TO OTHER TIMES

Radionuclide monitoring data are most comprehensive for the decade of the 1960s. Before that time, there was no technology for easily discriminating among the various radionuclides represented by the gross beta measurements. It is therefore much more difficult to piece together a complete picture of contaminants in the river for periods earlier than about 1960. However, a rough estimate can be made on the basis of the gross beta measurements. Annual total beta measurements derived from the literature are presented in Table 10.11. The initial construction of the reactors in the late 1940s and early 1950s, the increase in individual reactor power levels in the late 1950s, and the gradual shutdown of the reactors in the late 1960s are all reflected in the Columbia River total beta measurements shown in Table 10.11. If the individual doses are assumed to scale in rough proportion to the beta measurements, this table provides evidence that the largest doses probably occurred in the late 1950s through the mid-1960s.

As an example of scaling of the doses based only on the gross beta measurements, an attempt is made to predict the dose to a maximally exposed individual in 1966 in Richland on the basis of the estimated 1964 dose reported in Table 10.7 and the measurements in Table 10.11. The 1964 dose estimate is 0.16 rem. The 1964 and 1966 annual average gross beta measurements are 11,000 and 5700 pCi/L, respectively. The estimate for 1966 can be calculated as

$$(0.16 \text{ rem}) \frac{5700 \text{ pCi/L}}{11,000 \text{ pCi/L}} = 0.083 \text{ rem}$$

The calculated estimate from Table 10.7 is 0.082 rem, indicating reasonable accuracy by this technique. Because reactor operating parameters and water

TABLE 10.11. Annual Average Total Beta Concentrations, Columbia River at Pasco, Washington

<u>Year</u>	<u>Concentration pCi/L</u>	<u>Reference</u>
1945	470	Clukey (1957)
1946	210	Clukey (1957)
1947	180	Clukey (1957)
1948	110	Clukey (1957)
1949	220	Clukey (1957)
1950	520	Clukey (1957)
1951	990	Clukey (1957)
1952	1,410	Clukey (1957)
1953	2,120	Clukey (1957)
1954	1,750	Clukey (1957)
1955	2,160	Clukey (1957)
1956	3,000	Clukey (1957)
1957	5,080	Clukey (1957)
1958	9,150	Junkins and McConiga (1959)
1959	9,400	Foster and Junkins (1960)
1960	11,000	Nelson (1961)
1961	11,000	Nelson (1962)
1962	7,500	Wilson (1964)
1963	11,000	Wilson (1964)
1964	11,000 ^(a)	Essig (1970)
1965	8,250 ^(a)	Essig (1970)
1966	5,700 ^(a)	Essig (1970)
1967	6,200 ^(a)	Essig (1970)
1968	4,100 ^(a)	Essig (1970)
1969	3,400 ^(b)	Ballinger and Hall (1991)
1970	1,600 ^(b)	Ballinger and Hall (1991)
1971	100 ^(b)	Ballinger and Hall (1991)

(a) Based on number of operating reactors and ratios to phosphorus-32 and zinc-65.

(b) Based on number of reactor operating months.

treatment changed over time, it is probable that estimates for earlier years will not be as good as this example. Also, because the mix of radionuclides changes with radiological decay downstream of Richland, this technique will not work as well for the downstream locations. However, this example provides some evidence for the conclusion that scaling on gross beta concentrations provides an indication of the relative magnitude of the possible doses.

11.0 REFERENCES

Backman, G. E. 1962. Dispersion of 300 Area Liquid Effluent in the Columbia River. HW-73672, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Backman, G. E. 1963. Determination of River Dispersion Factors. HW-SA-3056, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Ballinger, M. Y., and R. B. Hall. 1991. A History of Major Hanford Facilities and Processes Involving Radioactive Material. PNL-6964 HEDR, Pacific Northwest Laboratory, Richland, Washington.

Barnes, C. A., A. C. Duxbury, and B. A. Morse. 1972. "Circulation and Selected Properties of the Columbia River Effluent at Sea." In The Columbia River Estuary and Adjacent Ocean Waters, eds. A. T. Pruter and D. L. Alverson, pp. 41-80. University of Washington Press, Seattle, Washington.

Beak Consultants Inc. 1980. "Aquatic Ecological Studies Near WNP-1, 2 and 4. August 1978-March 1980." WPPSS Columbia River Ecology Studies, Vol. 7. Prepared for Washington Public Power Supply System by Beak Consultants Inc., Portland, Oregon.

Beaseley, T. M., L. A. Ball, J. E. Andrews III, and J. E. Halverson. 1981. "Hanford-Derived Plutonium in Columbia River Sediments." Science 214:913-915.

Beaseley, T. M., and C. D. Jennings. 1984. "Inventories of $^{239,240}\text{Pu}$, ^{241}Am , ^{137}Cs , and ^{60}Co in Columbia River Sediments from Hanford to the Columbia River Estuary." Environmental Science and Technology 18:207-212.

Becker, C. D. 1985. Anadromous Salmonids of the Hanford Reach, Columbia River: 1984 Status. PNL-5371, Pacific Northwest Laboratory, Richland, Washington.

Becker, C. D. 1990. Aquatic Bioenvironmental Studies: The Hanford Experience 1944-84. Elsevier Science Publishing Company, New York.

Bogan, R. H. 1956. Review and Analysis of Hydrological and Radiochemical Columbia River Surveys. HW-45124, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Clukey, H. V. 1957. Hanford Atomic Project and Columbia River Pollution, HW-54243 Rev., General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Columbia River Estuary Data Development Program (CREDDP). 1984a. The Dynamics of the Columbia River Estuarine Ecosystem, Volume 1. Columbia River Estuary Study Taskforce (CREST), Astoria, Oregon.

Columbia River Estuary Data Development Program (CREDDP). 1984b. Sedimentary Processes and Environments in the Columbia River Estuary. Columbia River Estuary Study Taskforce (CREST), Astoria, Oregon.

Cushing, C. E., Jr. 1967a. "Concentration and transport of ^{32}P and ^{65}Zn by Columbia River plankton." Limnol. Oceanogr. 12(2):330-332.

Cushing, C. E., Jr. 1967b. "Periphyton productivity and radionuclide accumulation in the Columbia River, Washington, U.S.A." Hydrobiologia 29(1-2):125-139.

Cushing, C. E., D. G. Watson, A. J. Scott, and J. M. Gurtisen. 1981. "Decrease of Radionuclides in Columbia River Biota Following Closure of Hanford Reactors." Health Physics 41(1):59-67.

Cushing, C. E., ed. 1991. Hanford Site National Environmental Policy Act (NEPA) Characterization. PNL-6415 Rev. 4, Pacific Northwest Laboratory, Richland, Washington.

Cutshall, N., and C. Osterberg. 1964. "Columbia River Filter Samples." In Radioanalysis of Oceanic Organisms in the Pacific Ocean off Oregon, Progress Report, 1 March 1963 through 29 February 1964, ed. C. L. Osterberg, pp. 29-31. Oregon State University, Corvallis, Oregon.

Dauble, D. D., and D. G. Watson. 1990. Spawning and abundance of fall chinook salmon (*Oncorhynchus tshawytscha*) in the Hanford Reach of the Columbia River, 1948-1988. PNL-7298, Pacific Northwest Laboratory, Richland, Washington.

Davis, J. J., D. G. Watson, and C. C. Palmiter. 1956. Radiological Studies of the Columbia River Through December, 1955. HW-36074, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Davis, J. J., R. W. Perkins, R. F. Palmer, W. C. Hanson, and J. F. Cline. 1958. "Radioactive Materials in Aquatic and Terrestrial Organisms Exposed to Reactor Effluent Water." In Proceedings of the 2nd International Conference on the Peaceful Uses of Atomic Energy. Vol. 18, pp. 423-428.

DeNeal, D. L. 1965. Historical Events - Reactors and Fuels Fabrication. RL-REA-2247, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Dirkes, R. L. 1990. 1988 Hanford Riverbank Springs Characterization Report. PNL-7500, Pacific Northwest Laboratory, Richland, Washington.

Energy Research and Development Administration (ERDA). 1975. Final Environmental Impact Statement on Waste Management Operations, Hanford Reservation. ERDA-1538, Vol. 1-2, U.S. Energy Research and Development Administration, Washington, D.C.

Essig, T. H. 1970. Age-Dependent Radiation Dose Estimates for Residents of the Hanford Environs, 1945 - 1968. Master's thesis, Washington State University, Pullman, Washington.

Fisher, W. L. 1971. Radionuclide Concentrations in a Recent McNary Dam Sediment Core. BNWL-C-74, Pacific Northwest Laboratory, Richland, Washington.

Fix, J. J. 1975. "Association of Long-Lived Radioactivity with Sediment Along the Columbia River Shoreline, Islands, Bottom, and Slough Areas." CONF-7509067-20, from 3rd ERDA Environmental Protection Conference, Chicago.

Forster, W. O., and T. Guthrie. 1968. "Specific Activity of Zinc-65 in Natural Water." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 through 30 June 1968, ed. J. E. McCauley, pp. 72-81. RLO-1750-28, Oregon State University, Corvallis, Oregon.

Forster, W. O., and R. Loeffel. 1968. "Activity of Tagged Salmon in the North Pacific Ocean." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 through 30 June 1968, ed. J. E. McCauley, pp. 102-103. RLO-1750-28, Oregon State University, Corvallis, Oregon.

Foster, R. F. 1972. "History of Hanford and Its Contribution of Radionuclides to the Columbia River." In The Columbia River Estuary and Adjacent Ocean Waters, A. T. Pruter and D. L. Alverson, eds. University of Washington Press, Seattle, Washington.

Foster, R. F., et al. 1962. A Compilation of Basic Data Relating to the Columbia River. HW-69368, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and J. F. Honstead. 1967. Accumulation of Zinc-65 from Prolonged Consumption of Columbia River Fish. Health Phys. 13(1):39-43.

Foster, R. F., and R. L. Junkins. 1960. Off-Project Exposure from Hanford Reactor Effluent. HW-63654 Rev., General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and R. H. Wilson. 1964. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1963. HW-80991, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., J. K. Soldat, and T. H. Essig. 1966. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1965. BNWL-0316, Pacific Northwest Laboratory, Richland, Washington.

Frederick, L. C. 1967a. Dispersion of the Columbia River Plume Based on Radioactivity Measurements. Ph.D. dissertation, Oregon State University, Corvallis, Oregon.

Frederick, L. C. 1967b. "Tracking the Winter Plume by Large Volume Chemistry." In Ecological Studies in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1966 through 30 June 1967, ed. J. E. McCauley, pp. 34-38. RLO-1750-22, Oregon State University, Corvallis, Oregon.

Glenn, J. L. 1973. Relations Among Radionuclide Content and Physical, Chemical, and Mineral Characteristics of Columbia River Sediments. USGS Professional Paper 433-M, U.S. Department of the Interior, Washington, D.C.

Grande, L. A. 1966. Gamma Spectrum, Count Rate, and Dose Rate Measurements on the Columbia Riverbank from Vernita to Sacajawea. BNWL-CC-468, Pacific Northwest Laboratory, Richland, Washington.

Gray, R. H., and D. D. Dauble. 1977. "Checklist and relative abundance of fish species from the Hanford Reach of the Columbia River." Northwest Sci. 51(3):208-215.

Hall, R. B. 1963. Environmental Effects of a Fuel Element Failure. HW-79073, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Hall, R. B., and P. C. Jerman. 1960. Reactor Effluent Water Disposal. HW-63653, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Hanson, P. J. 1967. Vertical Distribution in the Columbia River Estuary. Master's thesis, Oregon State University, Corvallis, Oregon.

Haushild, W. L., G. R. Dempster, and H. H. Stevens. 1975. Distribution of Radionuclides in the Columbia River Streambed, Hanford Reservation to Longview, Washington. USGS Professional Paper 433-0, U.S. Department of the Interior, Washington, D.C.

Haushild, W. L., R. W. Perkins, H. H. Stevens, G. R. Dempster, and J. L. Glenn. 1966. Progress Report: Radionuclide Transport in the Pasco to Vancouver, Washington Reach of the Columbia River, July 1962 to September 1963. USGS-PP-433-N, USGS Open-File Report, U.S. Geological Survey, Portland, Oregon.

Haushild, W. L., H. H. Stevens, J. L. Nelson, and G. R. Dempster. 1973. Radionuclides in Transport in the Columbia River from Pasco to Vancouver, Washington. USGS Professional Paper 433-N, U.S. Department of the Interior, Washington, D.C.

Haynes, J. M. 1978. Movement and habitat studies of chinook salmon and white sturgeon. PNL-2471, Pacific Northwest Laboratory, Richland, Washington.

Healy, J. W. 1952. Effect of Purging During Pile Operation on the Effluent Water. HW-24578, General Electric Company, Hanford Works, Richland, Washington.

Honstead, J. F. 1954. Columbia River Survey 1951, 1952, 1953. HW-32506, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Honstead, J. F., T. H. Essig, and J. K. Soldat. 1967. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1966. BNWL-0439, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F., J. W. Healy, and H. J. Paas. 1951. Columbia River Survey Preliminary Report. HW-22851, General Electric Company, Hanford Works, Richland, Washington.

Houston, J. R., and P. J. Blumer. 1978. Environmental Surveillance at Hanford for Calendar Year 1977. PNL-2614, Pacific Northwest Laboratory, Richland, Washington.

Hubbell, D. W., and J. L. Glenn. 1973. Distribution of Radionuclides in Bottom Sediments of the Columbia River Estuary. USGS Professional Paper 433-L, U.S. Department of the Interior, Washington, D.C.

Isakson, J. S. 1969. Total Phosphorus and Phosphorus-32 in Seawater, Invertebrates, and Algae from North Head, Washington, 1965-66. Master's thesis, University of Washington, Seattle, Washington.

Jaquish, R. E., and R. W. Bryce, eds. 1989. Hanford Site Environmental Report for Calendar Year 1988. PNL-6825, Pacific Northwest Laboratory, Richland, Washington.

Jennings, C. D. 1966. Radioactivity of Sediments in the Columbia River Estuary. Master's thesis, Oregon State University, Corvallis, Oregon.

Jerman, P. C., W. N. Koop, and F. E. Owen. 1965. Release of Radioactivity to the Columbia River from Irradiated Fuel Element Ruptures. RL-REA-2160, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Johnson, V., N. Cutshall, and C. Osterberg. 1966. "Distribution of Hanford Induced Gamma Emitters in a Fresh Water Mollusc: *Anodonta wahlamatisensis*." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 through 30 June 1966, ed. J. E. McCauley, pp. 105-109. RLO-1750-8, Oregon State University, Corvallis, Oregon.

Jonsson, G., and A. H. Seymour. 1965. Gamma-Emitting Radionuclides in Dover Sole Caught off the Mouth of the Columbia River. Laboratory of Radiation Ecology, University of Washington, Seattle, Washington.

Junkins, R. L., and W. M. McConiga. 1959. Hanford Mid-Year Report to Columbia River Advisory Group - Through June 1959. HW-61029 Rev., General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Koop, W. N. 1956. Influences of Purges During Reactor Operation on the Concentration of Radioisotopes in the Columbia River. HW-45125, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Koop, W. N. 1957a. Effect of Reactor Purges When River Temperature Exceeds 15°C. HW-50601, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Koop, W. N. 1957b. Disposal of Decontaminating Agents for Reactor Rear Face Piping. HW-53372, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Koop, W. N., J. D. McCormack, and R. B. Hall. 1958. Effects of Direct Disposal of Reactor Effluent Water to the River. HW-55249, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kujala, N. F. 1966. Artificial Radionuclides in Pacific Salmon. Master's thesis, Oregon State University, Corvallis, Oregon.

Kujala, N. F., and W. O. Forster. 1968. "Radioisotope Measurements in Pacific Salmon." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 through 30 June 1968, ed. J. E. McCauley, pp. 104-105. RLO-1750-28, Oregon State University, Corvallis, Oregon.

Larsen, I. L. 1970. "Determination of Zn-65 Specific Activity in Various Tissues of the California Sea Mussel, *Mytilus californianus*, from Stations along the Pacific Coast." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1969 through 30 June 1970, ed. J. E. McCauley, pp. 53-60. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Larsen, I. L., W. C. Renfro, and W. O. Forster. 1968. "In-Situ Probe Detection of Radioactivity in the Columbia River Estuary." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 through 30 June 1968, ed. J. E. McCauley, pp. 39-40. RLO-1750-28, Oregon State University, Corvallis, Oregon.

Lewis, G. B., and A. H. Seymour. 1965. "Distribution of Zinc-65 in Plankton from Offshore Waters of Washington and Oregon, 1961-1963." Ocean Science and Engineering 2:956-967.

Lodge, J. D. 1966. Radiation Survey of the Columbia River Shoreline Near the Hanford Reactor Areas. BNWL-CC-835, Pacific Northwest Laboratory, Richland, Washington.

McConiga, M. W., and F. L. Rising. 1959. Scintillator Correlation and Columbia River Radiation Survey, March-April 1959. HW-60880, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McConnon, D. 1962. Dose-Rate Measurements of Beaches and Islands on the Columbia River Between Ringold and Richland. HW-72229 REV, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McCormack, W. D., and J.M.V. Carlile. 1984. Investigation of Ground-Water Seepage from the Hanford Shoreline of the Columbia River. PNL-5289, Pacific Northwest Laboratory, Richland, Washington.

McCormack, W. D., J. V. Ramsdell, and B. A. Napier. 1984. Hanford Dose Overview Program: Standardized Methods and Data for Hanford Environmental Dose Calculations. PNL-3777 Rev. 1, Pacific Northwest Laboratory, Richland, Washington.

McCormack, J. D., and L. C. Schwendiman. 1959. Significance of Rupture Debris in the Columbia River. HW-61325, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Mellinger, P. J. 1966. "The Concentration of Zinc-65 in *Mytilus californianus* as a Function of Distance from the Columbia River." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 through 30 June 1966, ed. J. E. McCauley, pp. 113-118. RLO-1750-8, Oregon State University, Corvallis, Oregon.

Moffatt, R. L., R. E. Wellman, and J. M. Gordon. 1990. Statistical Summaries of Streamflow Data in Oregon: Volume 1--Monthly and Annual Streamflow, and Flow-Duration Values. U.S. Geological Survey Open-File Report 90-118. Prepared in Cooperation with the Oregon Water Resources Department, Portland, Oregon.

Mooney, R. R. 1970. Environmental Radiation Surveillance in Washington State, Ninth Annual Report, July 1969 - June 1970. State of Washington, Department of Social and Health Services, Olympia, Washington.

Mooney, R. R. 1972. Environmental Radiation Surveillance in Washington State, Eleventh Annual Report, July 1971 - June 1972. State of Washington, Department of Social and Health Services, Olympia, Washington.

Mooney, R. R. 1974. Environmental Radiation Surveillance in Washington State, Thirteenth Annual Report, July 1973 - June 1974. State of Washington, Department of Social and Health Services, Olympia, Washington.

Mooney, R. R. 1975. Environmental Radiation Surveillance in Washington State, Fourteenth Annual Report, July 1974 - June 1975. State of Washington, Department of Social and Health Services, Olympia, Washington.

Mooney, R. R. 1976. Environmental Radiation Surveillance in Washington State, Fifteenth Annual Report, July 1975 - June 1976. State of Washington, Department of Social and Health Services, Olympia, Washington.

Mooney, R. R. 1977. Environmental Radiation Surveillance in Washington State, Sixteenth Annual Report, July 1976 - June, 1977. State of Washington, Department of Social and Health Services, Olympia, Washington.

Morse, B. A., M. G. Gross, and C. A. Barnes. 1968. "Movement of Seabed Drifters Near the Columbia River." Journal of Waterways & Harbors Division, ASCE Proceedings 94(WW1):93-103.

Myers, D. A., J. J. Fix, and J. R. Raymond. 1977. Environmental Monitoring Report on the Status of Ground Water Beneath the Hanford Site January-December 1976. BNWL-2199, Pacific Northwest Laboratory, Richland, Washington.

Naidu, J. R. 1963. Stable and Radioactive Zinc in Willapa Bay. Master's thesis, University of Washington, Seattle, Washington.

Napier, B. A. 1991. Selection of Dominant Radionuclides for Phase I of the HEDR Project. PNL-7231 HEDR, Pacific Northwest Laboratory, Richland, Washington.

Napier, B. A., R. A. Peloquin, J. V. Ramsdell, and D. L. Streng. 1988. GENII - The Hanford Environmental Radiation Dosimetry Software System. PNL-6584, Vol. 1-3, Pacific Northwest Laboratory, Richland, Washington.

Neal, V. T. 1972. "Physical Aspects of the Columbia River and Its Estuary." In The Columbia River Estuary and Adjacent Ocean Waters, eds. A. T. Pruter and D. L. Alverson, pp. 19-40. University of Washington Press, Seattle, Washington.

Neitzel, D. A., T. L. Page, and R. W. Hanf, Jr. 1981. "Mid-Columbia River microflora." J. Freshwat. Ecol. 1(5):495-505.

Neitzel, D. A., T. L. Page, and R. W. Hanf, Jr. 1983. "Mid-Columbia River zooplankton." Northwest Sci. 57(2):112-118.

Nelson, I. C., ed. 1961. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1960. HW-68435, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nelson, I. C., ed. 1962. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1961. HW-71999, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nelson, J. L. 1965. "Distribution of Sediments and Associated Radionuclides in the Columbia River Below Hanford." In Hanford Radiological Sciences Research and Development Annual Report for 1964, pp. 3.80-3.88. BNWL-36, Pacific Northwest Laboratory, Richland, Washington.

Nelson, J. L., and W. L. Haushild. 1969. Accumulation of Radionuclides in Bed Sediments of the Columbia River Between Hanford Reactors and McNary Dam. BNWL-SA-2571, Pacific Northwest Laboratory, Richland, Washington.

Nelson, J. L., R. W. Perkins, and W. L. Haushild. 1965. Determination of Columbia River Flow Times Using Radioactive Tracers Introduced by the Hanford Reactors. BNSA-176, Pacific Northwest Laboratory, Richland, Washington.

Nelson, J. L., R. W. Perkins, and W. L. Haushild. 1966. "Determination of Columbia River Flow Times Downstream from Pasco, Washington, Using Radioactive Tracers Introduced by the Hanford Reactors." Water Resources Research 2(1):31-39.

Nelson, J. L., R. W. Perkins, and J. M. Nielsen. 1964. Progress in Studies of Radionuclides in Columbia River Sediments. HW-83614, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nelson, J. L., R. W. Perkins, J. M. Nielsen, and W. L. Haushild. 1966. "Reactions of Radionuclides from the Hanford Reactors with Columbia River Sediments." In Disposal of Radioactive Wastes Into Seas, Oceans, and Surface Waters, pp. 139-161. International Atomic Energy Agency, Vienna, Austria.

Nielsen, J. M., and R. W. Perkins. 1957. The Depletion of Radioisotopes from the Columbia River by Natural Processes. HW-52908, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Oregon Department of Human Resources. 1985. Environmental Radiological Surveillance Report for Surface Waters in the State of Oregon, 1961 - 1983: Data Tables. Oregon Department of Human Resources, Salem, Oregon.

Osterberg, C. L. 1962. Radioactivity in Oceanic Organisms. Ph.D. dissertation, Oregon State University, Corvallis, Oregon.

Paas, H. J. 1951a. Radioactive Contamination in the Environs of the Hanford Works for the Period April, May, June, 1951. HW-22313, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. 1951b. Radioactive Contamination in the Environs of the Hanford Works for the Period October, November, December, 1950. HW-21566, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. 1953. Radioactive Contamination in the Hanford Environs for the Period January, February, March, 1953. HW-28009, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J. 1954. Radioactive Contamination in the Hanford Environs for the Period January, February, March, 1954. HW-31818, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J., and W. Singlevich. 1950. Radioactive Contamination in the Environs of the Hanford Works for the Period January, February, March, 1950. HW-18446, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J., and W. Singlevich. 1951a. Radioactive Contamination in the Environs of the Hanford Works for the Period July, August, September, 1950. HW-20700, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J., and W. Singlevich. 1951b. Radioactive Contamination in the Environs of the Hanford Works for the Period January, February, March, 1951. HW-21214, General Electric Company, Hanford Works, Richland, Washington.

Pacific Northwest Laboratory (PNL). 1991. Columbia River Pathway Report. PNL-7411 HEDR Rev. 1, Pacific Northwest Laboratory, Richland, Washington.

Pacific Northwest River Basin Commission (PNRBC). 1972. River Mile Index Main Stem Columbia River. Hydrology and Hydraulics Committee, PNRBC.

Page, T. L. and D. A. Neitzel. 1978. "Columbia River benthic macrofauna and microflora near WNP 1, 2, and 4: January through December 1977." In Aquatic Ecological Studies Near WNP through December 1977. WPPSS Columbia River Ecology Studies, Vol. 5, Section 4. Battelle, Pacific Northwest Laboratories, Richland, Washington.

Page, T. L., D. A. Neitzel, and R. W. Hanf. 1979. "Columbia River benthic macrofauna and microflora near WNP 1, 2, and 4: January through August 1978." In Aquatic Ecological Studies Near WNP 1, 2, and 4, January through August 1978. WPPSS Columbia River Ecology Studies, Vol. 6, Section 4. Battelle, Pacific Northwest Laboratories, Richland, Washington.

Parker, H. M. 1945. Status of Problem of Measurement of the Activity of Waste Water Returned to the Columbia River. HW-7-2346, E. I. duPont de Nemours and Company, Hanford Engineer Works, Richland, Washington.

Parker, H. M. 1952. Permissible Limits - Release of Reactor Effluent to the Columbia River. HW-24356, General Electric Company, Hanford Works, Richland, Washington.

Parker, H. M. 1954. Columbia River Situation - A Semi-Technical Review. HW-32809, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Pearcy, W. G. 1966. "Radioecology of Albacore (*Thunnus alalunga*) from the Northeastern Pacific." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 through 30 June 1966, ed. J. E. McCauley, pp. 148-152. RLO-1750-8, Oregon State University, Corvallis, Oregon.

Pearcy, W. G., and W. O. Forster. 1968. "Zinc-65 Specific Activities: Pelagic Animals and Surface Water in the Vicinity of the Columbia River Plume." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 through 30 June 1968, ed. J. E. McCauley, pp. 154-159. RLO-1750-28, Oregon State University, Corvallis, Oregon.

Pearcy, W. G., and C. L. Osterberg. 1963. "Notes on the Vertical Distribution of Zinc-65 and Zirconium-95 from Oceanic Animals." In Species Composition and Distribution of Marine Nekton in the Pacific Ocean off Oregon, Final Progress Report, 1 September 1961 through 1 September 1963, pp. 81-89. TID-19608, Oregon State University, Corvallis, Oregon.

Pearcy, W. G., and C. L. Osterberg. 1967. "Radioecology of Albacore from the Northeastern Pacific." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1966 through 30 June 1967, ed. J. E. McCauley, pp. 90-100. RLO-1750-22, Oregon State University, Corvallis, Oregon.

Perkins, R. W. 1959. The Origin of Reactor Effluent Water Radioisotopes. I. The Effect of Turco Purging on the Radioisotope Concentration in Reactor Effluent Water and Radioisotope Deposition on Metal Surfaces. HW-60890, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Perkins, R. W. 1961. Source of Reactor Effluent Water Radioisotopes. HW-69969, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Perkins, R. W., J. L. Nelson, and W. L. Haushild. 1965. Behavior and Transport of Radionuclides in the Columbia River Between Hanford and the Ocean. BNSA-16, Pacific Northwest Laboratory, Richland, Washington.

Pilcher, G. E., and H. T. Norton. 1953. The Calculation of Beta Particle Emitter Concentration in Hanford Reactor Effluent Water. HW-27584, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Poston, T. M., and D. C. Klopfer. 1966. A Literature Review of the Concentration Ratios of Selected Radionuclides in Freshwater and Marine Fish. PNL-5484, Pacific Northwest Laboratory, Richland, Washington.

Pruter, A. T. 1972. "Review of Commercial Fisheries in the Columbia River and in Contiguous Ocean Waters." In The Columbia River Estuary and Adjacent Pacific Ocean Waters, eds. A. T. Pruter and D. L. Alverson, pp. 81-120. University of Washington Press, Seattle, Washington.

Pruter, A. T., and D. L. Alverson, eds. 1972. The Columbia River Estuary and Adjacent Pacific Ocean Waters. University of Washington Press, Seattle, Washington.

Renfro, W. C. 1966. "Transfer of Radionuclides Through Estuarine Food Webs." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 through 30 June 1966, ed. J. E. McCauley, pp. 37-39. RLO-1750-8, Oregon State University, Corvallis, Oregon.

Renfro, W. C., W. O. Forster, and C. L. Osterberg. 1972. "Seasonal and Areal Distributions of Radionuclides in the Biota of the Columbia River Estuary." In The Columbia River Estuary and Adjacent Ocean Waters, Bioenvironmental Studies, eds. A. Pruter and D. Alverson, pp. 777-798. University of Washington Press, Seattle, Washington.

Richmond, M. C., and W. H. Walters. 1991. Estimates of Columbia River Radionuclide Concentrations: Data for Phase I Dose Calculations. PNL-7248 HEDR, Pacific Northwest Laboratory, Richland, Washington.

Robeck, G. G., C. Henderson, and R. C. Palange. 1954. Water Quality Studies on the Columbia River. U.S. Public Health Service, U.S. Department of Health, Education, and Welfare, Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio.

Robertson, D. E., and J. J. Fix. 1977. Association of Hanford Origin Radionuclides with Columbia River Sediment. BNWL-2305, Pacific Northwest Laboratory, Richland, Washington.

Robertson, D. E., W. B. Silker, J. C. Langford, M. R. Petersen, and R. W. Perkins. 1973. Transport and Depletion of Radionuclides in the Columbia River. BNWL-SA-4203, Pacific Northwest Laboratory, Richland, Washington.

Rokkan, D. J. 1988. Westinghouse Hanford Company 100 Areas Environmental Releases for 1987. WHC-EP-0165, Westinghouse Hanford Company, Richland, Washington.

Rostenbach, R. E. 1956. Columbia River Survey: 1954 - 1955. HW-43529, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Seymour, A. H. 1966. "Accumulation and Loss of Zinc-65 by Oysters in a Natural Environment." In Proceedings of the Symposium on the Disposal of Radioactive Wastes into Seas, Oceans, and Surface Waters, pp. 605-619. International Atomic Energy Agency, Vienna, Austria.

Seymour, A. H. 1970. Columbia River Studies, Annual Progress Report, 1969-1970. Laboratory of Radiation Ecology, University of Washington, Seattle, Washington.

Seymour, A. H., and G. B. Lewis. 1964. Radionuclides of Columbia River Origin in Marine Organisms, Sediments, and Water Collected from the Coastal and Offshore Waters of Washington and Oregon 1961-1963. UWFL-86, University of Washington, Seattle, Washington.

Shipler, D. B. 1992. FY 1992 Task Plans for the Hanford Environmental Dose Reconstruction Project. PNL-7757 HEDR, Pacific Northwest Laboratory, Richland, Washington.

Silker, W. B. 1963. Variations in Elemental Concentrations in the Columbia River. HW-SA-3147, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Silker, W. B. 1964. "Variations in Elemental Concentrations in the Columbia River." Limnol. Oceanog. 9:540-545.

Singlevich, W., and H. J. Paas. 1949. Radioactive Contamination in the Environs of the Hanford Works for the Period October, November, December 1948. HW-13743, General Electric Company, Hanford Works, Richland, Washington.

Soldat, J. K. 1956. Columbia River Travel Time Measurements By Float Methods. HW-41275, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Soldat, J. K. 1962a. Columbia River Flow Time Calculations. HW-58312 REV, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Soldat, J. K. 1962b. A Compilation of Basic Data Relating to the Columbia River. Section 8 - Dispersion of Reactor Effluent in the Columbia River. HW-69369, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Soldat, J. K., and G. R. Quimby. 1953. Effective Retention Time of the Hanford 100 Reactor Effluent Retention Basins. HW-28830, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Sula, M. J. 1980. Radiological Survey of Exposed Shorelines and Islands of the Columbia River Between Vernita and the Snake River Confluence. PNL-3127, Pacific Northwest Laboratory, Richland, Washington.

Tennant, D. A., and W. O. Forster. 1969. "Seasonal Variation and Distribution of Zn-65, Mn-54 and Cr-51 in Tissues of the Crab Cancer magister Dana." Health Physics 18:(6)649-657.

Thompson, S. E., C. A. Burton, D. J. Quinn, and Y. C. Ng. 1972. Concentrations Factors of Chemical Elements in Edible Aquatic Organisms. UCRL-50564 Rev. 1, Lawrence Livermore Laboratory, Livermore, California.

Toombs, G. L., and P. B. Cutler. 1968. Comprehensive Final Report for Lower Columbia River Environmental Radiological Survey in Oregon, June 5, 1961 - July 31, 1967. Oregon State Board of Health, Salem, Oregon.

Toombs, G. L., and R. D. Paris. 1978. Comprehensive Report for Lower Columbia River Environmental Radiological Survey in Oregon, 1967 - 1977. Oregon State Health Division, Portland, Oregon.

Turner, L. D. 1947. The Trend of Contamination Observed in the Air, the Columbia River, Vegetation, and Waste at the Hanford Engineer Works for the Period January 1, 1947 to March 25, 1947. HW-3-5511, General Electric Company, Hanford Works, Richland, Washington.

U.S. Geological Survey. 1975. Water Resources Data for Washington, 1975. U.S. Geological Survey Water-Data Report WA 75-1. Prepared in Cooperation with the State of Washington and with other agencies, Tacoma, Washington.

Vanderploeg, H. A., D. C. Parzyck, H. H. Wilcox, J. R. Kerchner, and S. V. Kaye. 1975. Bioaccumulation Factors for Radionuclides in Freshwater Biota. ORNL-5002, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Washington Department of Health. 1962. Radiological Survey of Surface Water Environment in the State of Washington. Washington Department of Health, Olympia, Washington.

Watson, D. G. 1970. Fall chinook salmon spawning in the Columbia River near Hanford 1947-1969. BNWL-1515. Pacific Northwest Laboratory, Richland, Washington.

Watson, D. G. 1973. "Fall chinook salmon population census." In BNWL-1750-Part 2, p. 6.16. Pacific Northwest Laboratory, Richland, Washington.

Watson, D. G., C. E. Cushing, C. C. Coutant, and W. L. Templeton. 1969. "Effect of Hanford Reactor Shutdown on Columbia River Biota." In Proceedings of the Second National Symposium on Radioecology, D. J. Nelson and F. C. Evans, eds., pp. 291-299. CONF-670503, Ann Arbor, Michigan.

Watson, D. G., C. E. Cushing, C. C. Coutant, and W. L. Templeton. 1970. Radioecological studies on the Columbia River. BNWL-1377, Pacific Northwest Laboratory, Richland, Washington.

Watson, D. G., and J. J. Davis. 1957. Concentration of radioisotopes in Columbia River whitefish in the vicinity of the Hanford Atomic Products Operation. HW-48523, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Williams, J. R., and H. E. Pearson. 1985. Streamflow Statistics and Drainage-Basin Characteristics for the Southwestern and Eastern Regions, Washington, Volume II. Eastern Washington. U.S. Geological Survey Open-File Report 84-145-B. Prepared in Cooperation with the State of Washington Department of Ecology, Tacoma, Washington.

Wilson, R. H. 1962. The Columbia River Environmental Monitoring Program. HW-75076, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Wilson, R. H., ed. 1964. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1963. HW-80991, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

APPENDIX A

COLUMBIA RIVER HYDROLOGIC DATA SOURCES

APPENDIX A

COLUMBIA RIVER HYDROLOGIC DATA SOURCES

This appendix summarizes the major sources of hydrologic, hydraulic, and other related data that describe the Columbia River flow regime and provide data for transport calculations. Some sources contain river data specific to the Hanford Site, while others include water quality data (e.g., sediment, water temperature) or fish and wildlife information. Because several categories of data and information are often included in the separate references, each reference is listed individually with a brief summary of the kinds of pertinent data and information it contains.

1. Streamflow Statistics and Drainage Basin Characteristics for the Southwestern and Eastern Regions, Washington, Vol. II: Eastern Washington. USGS Open-File Report 84-145B.

Monthly discharge for all years of record for the U.S. Geological Survey gauging stations on the Columbia River and its tributaries.

2. Woods, V. W. 1954. A Summary of Columbia River Hydrographic Information Pertinent to Hanford Works, 1894-1954. HW-30347, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Daily discharge (in cubic feet per second) at the Trinidad, Washington, gauge (RM 441), about 50 miles upstream of Hanford, from January 2, 1943, to December 31, 1953. Various staff gauges at the Hanford Site were calibrated relative to the Trinidad discharge readings, assuming negligible drainage basin inflow between Trinidad and the Hanford gauges. The Hanford gauges used mean sea level as datum. Evidently, temporary gauges were set at four locations until others could be permanently established at the 100-B, 100-D, and 100-F Areas. The gauge locations (daily readings) and periods of record are as follows:

<u>Location</u>	<u>Period of Record</u>
Coyote Rapids	July 7 - December 31, 1943
Wahluke	July 9 - December 26, 1943
White Bluffs	July 5 - December 30, 1943
Hanford	July 8 - December 30, 1943
181-B Pumphouse	July 15, 1944 - December 31, 1953
181-D Pumphouse	March 1, 1944 - December 8, 1944 ^(a)
	March 2, 1944 - December 31, 1953
181-F Pumphouse	July 1, 1944 - November 4, 1944 ^(a)
	April 24, 1945 - December 31, 1954

(a) Intermittent readings taken during construction.

3. General Electric Company. 1962. A Compilation of Basic Data Relating to the Columbia River. HW-69368, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Compilation of all river data, both published and unpublished, including such categories as river flows, radioactivity, sediment, effluent studies, water quality, and possible effects of Hanford effluent on river biota. Examples of water quality variables are temperature, pH, and turbidity. Most of the data were collected near the reactor areas. Other locations were Trinidad, Priest Rapids, Richland, and the McNary reservoir.

4. Honstead, J. F., J. W. Healy, and H. J. Paas. 1951. Columbia River Survey Preliminary Report. HW-22851, General Electric Company, Hanford Works, Richland, Washington.
5. Honstead, J. F. 1954. Columbia River Survey 1951, 1952, 1953. HW-32506, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Water samples, depth soundings, and velocity measurements were taken concurrently at about nineteen locations between 100-B and Pasco, Washington, at three different flow conditions (rising hydrograph, peak flow, and stable low flow). A fourth survey was conducted below a single reactor during a time when it was the only effluent source. The data are presented as river cross sections with velocity contours (isovels). Analysis of the water samples provided data for plotting the profile of the effluent plume.

6. U.S. Army Corps of Engineers. 1949. Report on Flood of May-June 1948. Columbia River and Tributaries (Upstream from Snake River). Seattle District, Seattle, Washington.

Flood hydrographs of the upper Columbia River and tributaries, increments of the lower Columbia River flood peak at The Dalles, and a water-surface profile for July 21, 1949.

7. Robeck, G. G., C. Henderson, and R. C. Palange. 1954. Water Quality Studies on the Columbia River. U.S. Department of Health, Education, and Welfare, Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio.

- a. Discharge and river stage curves for Columbia River near Hanford and near Umatilla for July 1951 - June 1953.

- b. Flow-time curves for the Columbia River from RM 368.5 (reactor area) to RM 292 (McNary Dam site) for discharges ranging from 46,000 to 500,000 cubic feet per second. Curves were developed for conditions before and after impoundment by McNary Dam.

- c. Routine physical and chemical measurements of water quality, including turbidity, temperature, and pH. Readings were taken approximately once weekly from January 1951 to February 1953 for about 12 cross sections between Priest Rapids and the McNary Dam site. Measurements were also taken on the Yakima River at Kiona, Enterprise, and the Richland Highway bridge, on the Snake River at Page and near the mouth, and on the Walla Walla River at the US Highway 410 bridge and the US Highway 395 bridge. The turbidity and temperature data are plotted.

8. Columbia River Estuary Data Development Program.

The Columbia River Estuary Data Development Program (CREDDP) was a 6-year program of study authorized by the U.S. Congress in October 1978 and completed in 1984. The objectives were to increase understanding of the ecology of the Columbia River estuary and to provide information useful in making decisions about land and water use. The research was divided into thirteen work units. Three of the units described and mapped the productivity and biomass patterns of the estuary's primary producers and their levels.

Seven units dealt with the higher trophic levels in the estuarine food web. These included zooplankton and larval fish, benthic infauna, epibenthic organisms, fish, avifauna, wildlife, and marine mammals. The goals of these units were to describe and map the abundance patterns of the invertebrate and vertebrate species and their relationships to physical factors.

The other three work units are, perhaps, the most relevant to the HEDR Project. These units dealt with sedimentation and shoaling, currents, and circulation. The goals were to characterize and map the bottom sediment distribution, to characterize sediment transport, to determine the causes of bathymetric change, and to determine and model circulation patterns, vertical mixing, and salinity patterns. Included with these reports is a detailed set of planform and bathymetric maps and portfolios.

9. River Maps and Cross Sections

Various sets of river channel maps and cross sections are available from the U.S. Army Corps of Engineers North Pacific Division Office in Portland, Oregon, and from district offices in Walla Walla, Washington; Seattle, Washington; and Portland, Oregon.

- a. 1894 - river surveys and depth soundings.
- b. 1955 - detailed sounding measurements in maps PD-7-24/0 to PD-7-24/22.
- c. 1963 - Columbia River longitudinal channel profile.
- d. 1986 - 145 cross sections from Priest Rapids to the Yakima River.

Navigation charts of the Columbia River from Priest Rapids Dam to the mouth are available from the U.S. Coast and Geodetic Survey.

10. Sedimentologic Data

The primary sources for these data are the U.S. Geological Survey and the U.S. Army Corps of Engineers division and district offices.

a. Daily record of water discharge, sediment concentration, and total sediment discharge for the following locations and dates:

<u>Location</u>	<u>Date</u>
Columbia River at Pasco	August 1962 - September 1966
Snake River at Pasco	August 1962 - September 1964
Columbia River at Umatilla	August 1965 - September 1966

b. Mean weekly suspended sediment concentrations and mean weekly sand and total sediment discharges at Pasco, Umatilla, and Vancouver for 1964 through 1966.

c. Mean particle-size distribution of surficial sediment samples from the riverbed (mid-channel and shoreward) at Pasco, Hood River, and Vancouver taken intermittently during 1962 through 1965. Each sample is dated as to the day of its collection.

APPENDIX B

TABLE OF RADIONUCLIDES

Table of Radionuclides

Radionuclide	Half-Life	Radiation ^(a)
Antimony-124	60 days	β^- , γ
Arsenic-76	26.5 hours	β^- , γ
Barium-140	12.8 days	β^- , e^- , γ
Cerium-144	284 days	β^- , e^- , γ
Cesium-137	30.0 years	β^- , e^- , γ
Chromium-51	27.8 days	e^- , γ
Cobalt-58	71.3 days	β^+ , γ
Cobalt-60	5.26 years	β^- , γ
Copper-64	12.8 hours	β^- , e^- , β^+ , γ
Europium-152	12 years	β^- , β^+ , e^- , γ
Europium-154	16 years	β^- , e^- , γ
Gallium-72	14.12 hours	β^- , γ
Iodine-129	1.7×10^7 years	β^- , e^- , γ
Iodine-131	8.05 days	β^- , e^- , γ
Iron-55	2.6 years	γ
Iron-59	45.6 days	β^- , γ
Lanthanum-140	40.22 hours	β^- , γ
Manganese-54	303 days	e^- , γ
Manganese-56	2.58 hours	β^- , γ
Neptunium-239	2.35 days	β^- , γ , e^-
Niobium-95	35 days	β^- , γ
Phosphorus-32	14.28 days	β^-
Plutonium-239	24,390 years	α , e^- , γ
Plutonium-240	6,580 years	α , e^- , γ
Potassium-40	1.42×10^9 years	β^- , β^+ , γ
Ruthenium-103	39.6 days	β^- , γ
Ruthenium-106	368 days	β^- , γ
Samarium-153	46.8 hours	β^- , e^- , γ
Scandium-46	83.9 days	β^- , γ

Table of Radionuclides

Radionuclide	Half-Life	Radiation ^(a)
Sodium-24	15.0 hours	β^- , γ
Strontium-89	57.7 days	β^- , γ
Strontium-90	28.1 years	β^- (D.R.)
Technetium-99	6.04 hours	γ , e^-
Tellurium-132	78 hours	β^- , e^- , γ
Tritium	12.26 years	β^-
Zinc-65	245 days	β^+ , e^- , γ
Zirconium-95	65 days	β^- , γ (D.R.)

(a) Conversion electrons (e^-) are listed if they are prominent in the electron spectrum. Decay products may give rise to daughter radiation. This is indicated, where prominent, by the notation (D.R.).

Sources: The Table of Isotopes, by C. M. Lederer, J. M. Hollander, and I. Perlman (6th ed.; New York: John Wiley & Sons, Inc., 1967).

Villforth, J. C. and G. R. Shultz. 1970 rev. Radio-logical Health Handbook. Bureau of Radiological Health and The Training Institute Environmental Control Administration, U.S. Department of Health, Education and Welfare, Rockville, Maryland.

APPENDIX C

BIBLIOGRAPHY

APPENDIX C

BIBLIOGRAPHY

All of the documents listed here were examined in the literature review, although they have not been cited in the body of the report.

Adley, F. E., and W. K. Crane. 1950. Meeting of the Columbia River Advisory Group November 21-23, 1949. HW-15861, General Electric Company, Hanford Works, Richland, Washington.

Adley, F. E., and W. K. Crane. 1950. Meeting of the Columbia River Advisory Group, March 6-7, 1950. HW-17595, General Electric Company, Hanford Works, Richland, Washington.

Andersen, B. V. 1959. Hanford Environmental Monitoring Annual Report for 1958. HW-61676, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V. Undated. Regional Monitoring - Data Summary All Results for Year 1957. HW-89066, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., and J. K. Soldat. 1955. Radioactive Contamination in the Hanford Environs for the Period July, August, September, 1955. HW-39429, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., and J. K. Soldat. 1956. Radioactive Contamination in the Hanford Environs for the Period October, November, December, 1955. HW-40871, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., and J. K. Soldat. 1956. Radioactive Contamination in the Hanford Environs for the Period January, February, March, 1956. HW-43012, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., and J. K. Soldat. 1956. Radioactive Contamination in the Hanford Environs for the Period April, May, June, 1956. HW-44215, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., and J. K. Soldat. 1957. Regional Monitoring Program. HW-52038, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., M. W. McConiga, and J. K. Soldat. 1956. Radioactive Contamination in the Hanford Environs for the Period July, August, September, 1956. HW-46726, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Andersen, B. V., M. W. McConiga, and J. K. Soldat. 1957. Radioactive Contamination in the Hanford Environs for the Period October, November, December, 1956. HW-48374, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Atwood, J. M., R. B. Hall, and S. L. Nelson. 1955. Recirculation Loop Slug Rupture Incident at 100-H. HW-40115, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Bainard, W. D. 1955. Chemical Contamination of Drinking Water by Sodium Dichromate at Low River Flows. HW-38469-RD, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Barnes, C. A., and M. G. Gross. 1966. Distribution at Sea of Columbia River Water and Its Load of Radionuclides. RLO-1725-41, University of Washington, Department of Oceanography, Seattle, Washington.

Beasley, T. M., and R. L. Holton. 1979. Cycling of Transuranic Radionuclides in the Columbia River, its Estuary, and the North Pacific Ocean. RLO-2227-T30-2, Oregon State University, Corvallis, Oregon.

Beasley, T. M., C. D. Jennings, and D. A. McCullough. 1986. "Sediment Accumulation Rates in the Lower Columbia River." In Journal of Environmental Radioactivity (3)2:103-123.

Becker, C. D. 1973. Aquatic Bioenvironmental Studies in the Columbia River at Hanford 1945-1971. BNWL-1734, Pacific Northwest Laboratory, Richland, Washington.

Beetle, T. M. 1972. "Statistical Study of Radiation Dose for Fishermen." BNWL-SA-4449, Pacific Northwest Laboratory, Richland, Washington.

Bosley, C. E., and G. F. Gately. 1981. Polychlorinated biphenyls and chlorinated pesticides in Columbia River white sturgeon (*Acipenser Transmontanus*). U.S. Fish and Wildlife Service, Nordland, Washington.

Bovingdon, J. R. 1966. Radiation Survey of Islands and Beaches on the Columbia River between Hanford and the 300 Area. BNWL-CC-1386, Pacific Northwest Laboratory, Richland, Washington.

Bramson, P. E. 1972. Environmental Status of the Hanford Reservation for 1970. BNWL-C-96, Pacific Northwest Laboratory, Richland, Washington.

Bramson, P. E., and J. P. Corley. 1972. Environmental Surveillance at Hanford for Calendar Year 1971. BNWL-1683, Pacific Northwest Laboratory, Richland, Washington.

Bramson, P. E., and J. P. Corley. 1972. Environmental Status of the Hanford Reservation for 1971. BNWL-B-228, Pacific Northwest Laboratory, Richland, Washington.

Bramson, P. E., and J. P. Corley. 1973. Environmental Surveillance at Hanford for Calendar Year 1972. BNWL-1727, Pacific Northwest Laboratory, Richland, Washington.

Bramson, P. E., J. P. Corley, and W. L. Nees. 1973. Environmental Status of the Hanford Reservation for Calendar Year 1972. BNWL-B-278, Pacific Northwest Laboratory, Richland, Washington.

Brauer, F. P., and H. G. Rieck. 1973. "I-129, Co-60 and Ru-106 Measurements on Water Samples from the Hanford Project Environs." BNWL-SA-4478, Pacific Northwest Laboratory, Richland, Washington.

Bugbee, S. J. 1943. Proposed Locations of Monitoring Stations. DUH-11853, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Bulow, W. F. 1945. Vegetation (Sage) Sampling and Analysis 7-1-45 Through 12-21-45: Water Samples and Analysis 4-23-45 Through 12-24-45, Columbia River, Hanford Site and Tri-Cities. HW-06660-T, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Carey, A. G., Jr. 1969. "Zinc-65 in Echinoderms and Sediments in the Marine Environment Off Oregon." In Symposium on Radioecology, Proceedings of the Second National Symposium, eds. D. J. Nelson and F. C. Evans, pp. 380-388. U.S. AEC CONF-670503, U.S. Atomic Energy Commission, Washington, D.C.

Carey, A. G., Jr., and N. Cutshall. 1972. "Zinc-65 Specific Activities from Oregon and Washington (U.S.A.) Continental Shelf Sediments and Benthic Invertebrate Fauna." Presented at Symposium on Interaction of Radioactive Contaminants with Constituents of the Marine Environment, July 10-14, 1972. RLO-2227-T12-26, IAEA/SM-158/17, Seattle, Washington.

Carey, A. G., Jr., and J. E. McCauley. 1966. "Radioecology and Ecology of Benthic Animals." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 June 1965 Through 30 July 1966, ed. J. E. McCauley. pp. 119-125. U.S. AEC RLO-1750-8, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Carey, A. G., Jr., W. G. Pearcy, and C. L. Osterberg. 1966. "Artificial Radionuclides in Marine Organisms in the Northeast Pacific Ocean Off Oregon." Disposal of Radioactive Wastes into Seas, Oceans, and Surface Waters, pp. 303-319. International Atomic Energy Agency, Vienna, Austria.

Carter, L. A., J. P. Corley, et al. 1961. Compilation of Basic Data Relating to the Columbia River Sections I-VI. HW-69368, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Clukey, H. V. 1956. Radiation Aspects for River Navigation Through Hanford Project. HW-47152, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Clukey, H. V. 1957. Hanford Atomic Project and Columbia River Pollution. HW-54243-REV, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Clukey, H. V., and M. W. McConiga. 1958. Calculation of Drinking-Water Radiation Exposures from Columbia River Water Sources. HW-57411, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Cohn, W. E. 1944. Tolerable Concentrations of Fission Products in Air and Water. DUH-04517, E. I. duPont de Nemours & Company, Hanford Engineer Works Richland, Washington.

Coleman, R. M. 1944. Measurement of Low Intensity Gamma Radiation. DUH-3070, Clinton Laboratories, Knoxville, Tennessee.

Conklin, A. W. 1987. Summary of Preliminary Review of Hanford Historical Documents - 1943 to 1957. EPS-87-201, State of Washington, Department of Social and Health Services, Olympia, Washington.

Coopey, R. W. 1948. Accumulation of Radioactivity as Shown by a Limnological Study of the Columbia River in the Vicinity of The Hanford Works. HW-11662, General Electric Company, Hanford Works, Richland, Washington.

Coopey, R. W. 1953. Abundance of the Principal Crustacea of the Columbia River and the Radioactivity They Contain. HW-25191, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Coopey, R. W. 1954. "The Abundance of the Principal Crustacea of the Columbia River and the Radioactivity They Contain." In Biology Research - Annual Report 1953, pp. 14-15. HW-30437-3, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Corley, J. P. 1968. "Hanford Experience with Composite Environmental Dose Estimates." BNWL-SA-1452, Pacific Northwest Laboratory, Richland, Washington.

Corley, J. P. 1969. Environmental Surveillance in the Vicinity of Hanford for November, 1968. BNWL-0778-11, Pacific Northwest Laboratory, Richland, Washington.

Corley, J. P. 1973. Environmental Surveillance at Hanford for Calendar Year 1970 Data. BNWL-1669-ADD, Pacific Northwest Laboratory, Richland, Washington.

Corley, J. P., C. B. Wilson, and T. H. Essig. 1970. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1969. BNWL-1505, Pacific Northwest Laboratory, Richland, Washington.

Corley, J. P., and C. B. Wooldridge. 1968. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1967. BNWL-0983, Pacific Northwest Laboratory, Richland, Washington.

Cory, R. L., W. O. Forster, W. C. Renfro, and J. W. Nauman. 1970. "Radioactivity of Three Fresh-Water Molluscs (*Anodonta wahlamatisensis*, *Anodonta oregonensis*, *Carbicula fluminea*) of the Tidal Columbia River and Their Associated Sediments." In Hydrobiology, Bioresources of Shallow Water Environments, eds. W. G. Weist, Jr. and P. E. Greeson, pp. 162-173. The American Water Resources Association, Miami Beach, Florida.

Cushing, C. E. 1965. Periphyton Productivity and Radionuclide Accumulation in the Columbia River, Washington, USA. BNWL-SA-0360, Pacific Northwest Laboratory, Richland, Washington.

Cushing, C. E., and D. G. Watson. 1966. Accumulation and Transport of Radionuclides by Columbia River Biota. BNWL-SA-0623, Pacific Northwest Laboratory, Richland, Washington.

Cushing, C. E., and D. G. Watson. 1973. Decline of Radioactivity in Biota Following Shutdown of Hanford Reactors. BNWL-SA-4718, Pacific Northwest Laboratory, Richland, Washington.

Cushing, C. E., D. G. Watson, J. M. Gurtisen, and A. J. Scott. 1980. Decrease of Radionuclides in Columbia River Biota Following Closure of the Hanford Reactors. PNL-SA-08407, Pacific Northwest Laboratory, Richland, Washington.

Cutshall, N. 1966. "Benthos-Nekton-Radiochemistry Cruise." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 Through 30 June 1966, ed. J. E. McCauley, pp. 110-111. U.S. AEC RLO-1750-8, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Cutshall, N. 1974. "Turnover of Zinc-65 in Oysters." Health Physics 26:327-331.

Cutshall, N. H. 1967. Chromium-51 in the Columbia River and Adjacent Pacific Ocean. Master's thesis, Oregon State University, Corvallis, Oregon.

Cutshall, N., and L. Larsen. 1964. "River Water Analyses." In Radioanalysis of Oceanic Organisms in the Pacific Ocean Off Oregon, Progress Report, 1 March 1963 Through 29 February 1964, ed. C. L. Osterberg, pp. 7-11. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Cutshall, N. H., and W. C. Renfro. 1970. "Zn-65 in Oregon-Washington Continental Shelf Sediments." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1969 Through 30 June 1970, ed. J. E. McCauley, pp. 29-36. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Davis, J. J., and C. L. Cooper. 1951. Effect of Hanford Pile Effluent upon Aquatic Invertebrates in the Columbia River. HW-20055, General Electric Company, Hanford Works, Richland, Washington.

Davis, J. J., R. W. Perkins, R. F. Palmer, W. C. Hanson, and J. F. Cline. 1958. "Radioactive Materials in Aquatic and Terrestrial Organisms Exposed to Reactor Effluent Water," pp. 423-428. In Proceedings of the Second United Nations International Conference on the Peaceful Uses of Atomic Energy, Vol. 18. United Nations, Geneva.

Davis, J. J., D. G. Watson, and C. C. Palmiter. 1956. Radiobiological Studies of the Columbia River Through December, 1955. HW-36074, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Davis, J. J., D. G. Watson, C. C. Palmiter, and R. W. Coopey. 1954. "1953 Radiological Survey of the Columbia River," pp. 8-13 in Biology Research - Annual Report 1953. HW-30437, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Dickinson, P. 1946. Site Survey - Water, Vegetation, and Special Sample Data, 1/20/46 Through 8/29/46. HEW-00616-T, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Dirkes, R. L. 1990. Hanford Riverbank Springs Characterization Report - 1988. PNL-7500, Pacific Northwest Laboratory, Richland, Washington.

E. I. duPont de Nemours & Company. 1943. Weekly Progress Report - Technical Division for Period Ending July 31, 1943. DUH-10771, Hanford Engineers Works, Richland, Washington.

Ecker, R. M., W. H. Walters, and F. L. Thompson. 1983. N Reactor Thermal Plume Characterization Study during Dual-Purpose Mode of Operation. UNI-2620, UNC Nuclear Industries, Richland, Washington.

Eisenacher, P. L. 1948. Beta Water Monitor. HW-09584, General Electric Company, Hanford Works, Richland Washington.

Environmental Monitoring at Hanford for 1986. 1987. PNL-6120, Pacific Northwest Laboratory, Richland, Washington.

Erickson, J. L., and R. R. Mooney. 1987. State of Washington Environmental Radiation Program, Twenty-Fifth Annual Report, January, 1986-December, 1986. State of Washington, Department of Social and Health Services, Division of Health, Office of Radiation Protection, Olympia, Washington.

Erickson, J. L., G. L. Ziegler, and R. R. Mooney. 1985. State of Washington Environmental Radiation Program, Twenty-Third Annual Report, January, 1984-December, 1984. State of Washington, Department of Social and Health Services, Division of Health, Office of Radiation Protection, Olympia, Washington.

Essig, T. H., and J. P. Corley. 1968. Comparison of Radionuclide Releases from Hanford with Environmental Dose Estimates, 1963-1967. BNWL-0886, Pacific Northwest Laboratory, Richland, Washington.

Essig, T. H., and J. P. Corley. 1969. Criteria Used to Estimate Radiation Doses Received by Persons Living in the Vicinity of Hanford. BNWL-1019, Pacific Northwest Laboratory, Richland Washington.

Essig, T. H., G. W. R. Endres, J. K. Soldat, and J. F. Honstead. 1972. Concentrations of Zn-65 in Marine Foodstuffs and Pacific Coastal Residents. BNWL-SA-4206, Pacific Northwest Laboratory, Richland, Washington.

Essig, T. H., and R. B. Hall. 1966. Environmental Status of the Hanford Project 1965 Annual Summary. BNWL-CC-0913, Pacific Northwest Laboratory, Richland, Washington.

Essig, T. H., and R. B. Hall. 1967. Environmental Status of the Hanford Reservation for December, 1966. BNWL-CC-0637-12, Pacific Northwest Laboratory, Richland, Washington.

Evans, D. W., and N. H. Cutshall. 1972. "Effects of Ocean Water on the Soluble-Suspended Distribution of Columbia River Radionuclides." Presented at IAEA Symposium on Interaction of Radioactive Contaminants with Constituents of the Marine Environment, July 10-14, 1972. Seattle, Washington.

Fisher, W. L. 1971. Radiological Considerations of Opening the Columbia River for Public Access Upstream from 100-F Area, Part 1. BNWL-B-071-PT1, Pacific Northwest Laboratory, Richland, Washington.

Fisher, W. L., and D. H. Denham. 1969. Radiological Considerations of Opening the Columbia River for Recreational Purposes - Ringold to 100-F Area. BNWL-CC-2363, Pacific Northwest Laboratory, Richland, Washington.

Fisher, W. L., and C. B. Wilson. 1970. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1968. BNWL-1341, Pacific Northwest Laboratory, Richland, Washington.

Fix, J. J. 1975. Environmental Surveillance at Hanford for Calendar Year 1974. BNWL-1910, Pacific Northwest Laboratory, Richland, Washington.

Fix, J. J., and P. J. Blumer. 1975. Environmental Surveillance at Hanford for Calendar Year 1974 Data. BNWL-1910-ADD, Pacific Northwest Laboratory, Richland, Washington.

Fix, J. J., P. J. Blumer, G. R. Hoenes, and P. E. Bramson. 1977. Environmental Surveillance at Hanford for Calendar Year 1976. BNWL-2142, Pacific Northwest Laboratory, Richland, Washington.

Forster, W. O., P. Hanson, and I. L. Larsen. 1968. "Medium Volume Precipitation Techniques Combined with In Situ Probe Detection, 'Mark Twain Cruise'." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967-30 June 1968, ed. J. E. McCauley, pp. 82-87. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Forster, W. O., and R. Loeffel. 1968. "Activity of Tagged Salmon in the North Pacific Ocean." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 June 1967 - 30 July 1968, pp. 102-103. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Foster, R. F. 1945. Fish Life Observed in the Columbia River on September 27, 1945. HW-7-2514, E. I. duPont de Nemours & Company, Hanford Engineers Works, Richland, Washington.

Foster, R. F. 1946. Occasional Heavy Mortalities Among Fish Held in 100-F Area Effluent Water and Some Effects of "Calol" on Steelhead Trout Fingerlings. HW-7-04243, E. I. duPont de Nemours & Company, Hanford Engineers Works, Richland, Washington.

Foster, R. F. 1947. Fish Laboratory Report for First Half of October, 1947. HW-07822, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1947. Fish Laboratory Report for the Last Half of October, 1947. HW-07951, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1947. Fish Laboratory Report for the Last Half of November, 1947. HW-08166, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1947. Fish Laboratory Report for the First Half of December, 1947. HW-08308, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1947. Fish Laboratory Report for the Last Half of December, 1947. HW-08421, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Aquatic Biology Laboratory Report for June, 1948. HW-10642, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the First Half of January, 1948. HW-08591, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the Last Half of January, 1948. HW-08733, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the First Half of February, 1948. HW-08902, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the Last Half of February, 1948. HW-09052, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the First Half of March, 1948. HW-09229, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the Last Half of March, 1948. HW-09395, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the First Half of April, 1948. HW-09530, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the Last Half of April, 1948. HW-09731, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the First Half of May, 1948. HW-09838, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the Last Half of May, 1948. HW-09960, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Fish Laboratory Report for the First Half of June, 1948. HW-10170, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1948. Occurrence of Dead Whitefish in the Columbia River During December, 1948. HW-11861, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1949. Proposed Radiobiological-Ecological Survey of the Columbia River. HW-12574, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1949. Radiological Problems in the Columbia River. HW-23793, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1949. Supplement to Proposed Radiobiological-Ecological Survey of the Columbia River. HW-12575, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1950. Abnormally High Activity in Aquatic Organisms Below the 100-H Area. HW-18438, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1950. Proposed Monitoring of Effluent Water with Chinook Salmon, Fiscal Year 1950. HW-19016, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1951. Rough Draft-Monthly Fish Laboratory and Aquatic Biology Reports Covering Periods December, 1946 Through December, 1950. HW-20277-RD, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1952. "Biological Problems Associated with the Discharge of Pile Effluent into the Columbia River," pp. 11-13 in Biology Research - Annual Report 1951. HW-25021, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F. 1958. Effect on Fish of Increasing the Temperature of the Columbia River. HW-54858, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1959. Behavior of Radionuclides in Food Chains Fresh Water Studies. HW-SA-1725, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Analysis of Radiological Data for the Month of February, 1961. HW-68851, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Analysis of Radiological Data for the Month of April, 1961. HW-69618, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Analysis of Radiological Data for the Month of May, 1961. HW-70103, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Analysis of Radiological Data for the Month of July, 1961. HW-70689, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Analysis of Radiological Data for the Month of August, 1961. HW-71041, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Summary of Radiological Data for the Month of October, 1961. HW-71678, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1961. Summary of Radiological Data for the Month of November, 1961. HW-72012, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Summary of Radiological Data for the Month of December, 1961. HW-72338, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Summary of Radiological Data for the Month of January, 1962. HW-72691-1, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Summary of Radiological Data for the Month of February, 1962. HW-72691-2, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Summary of Radiological Data for the Month of March, 1962. HW-72691-3, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Summary of Radiological Data for the Month of April, 1962. HW-72691-4, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Summary of Radiological Data for the Month of May, 1962. HW-72691-5, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Radiological Status of the Hanford Environs for June, 1962. HW-74307-06, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Radiological Status of the Hanford Environs for July, 1962. HW-74307-07, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Radiological Status of the Hanford Environs for August, 1962. HW-74307-08, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Radiological Status of the Hanford Environs for September, 1962. HW-74307-09, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Radiological Status of the Hanford Environs for October, 1962. HW-74307-10, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1962. Radiological Status of the Hanford Environs for November, 1962. HW-74307-11, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Accumulation of Zn-65 from Prolonged Consumption of Columbia River Fish. HW-SA-3060, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for December, 1962. HW-74307-12, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for January, 1963. HW-76525-01, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for February, 1963. HW-76525-02, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for March, 1963. HW-76525-03, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for April, 1963. HW-76525-04, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for May, 1963. HW-76525-05, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for June, 1963. HW-76525-06, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for July, 1963. HW-76525-07, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for August, 1963. HW-76525-08, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for September, 1963. HW-76525-09, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for October, 1963. HW-76525-10, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1963. Radiological Status of the Hanford Environs for November, 1963. HW-76525-11, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for December, 1963. HW-76525-12, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for January, 1964. HW-80892-01, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for February, 1964. HW-80892-02, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for March, 1964. HW-80892-03, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for April, 1964. HW-80892-04, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for May, 1964. HW-80892-05, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for June, 1964. HW-80892-06, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for July, 1964. HW-80892-07, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for August, 1964. HW-80892-08, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for September, 1964. HW-80892-09, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for October, 1964. HW-80892-10, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1964. Radiological Status of the Hanford Environs for November, 1964. HW-80892-11, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for December, 1964. HW-80892-12, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for January - March, 1965. BNWL-77-03, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for April, 1965. BNWL-77-04, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for May, 1965. BNWL-77-05, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for June, 1965. BNWL-77-06, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for July, 1965. BNWL-77-07, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for August, 1965. BNWL-77-08, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for September, 1965. BNWL-77-09, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for October, 1965. BNWL-77-10, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1965. Radiological Status of the Hanford Environs for November, 1965. BNWL-77-11, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1966. Radiological Status of the Hanford Environs for December, 1965. BNWL-77-12, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1969. Hanford Story and Its Contribution of Radionuclides to the Columbia River Waters. BNWL-SA-2812, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1970. Effects of Hanford Reactors on Columbia River and Adjacent Land Areas. BNWL-SA-3679, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1970. Radiation Exposure to the Local Population near the Hanford Plant. BNWL-SA-3455, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F. 1973. Sources and Inventory of Radioactivity in the Aquatic Environment. BNWL-SA-4614, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F., and R. W. Coopey. 1948. Aquatic Biology Laboratory. HW-23791, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F., R. W. Coopey, and J. J. Davis. 1949. Cursory Survey of the Radioactivity in Biological Materials of the Lower Columbia River. HW-12573, General Electric Company, Hanford Works, Richland Washington.

Foster, R. F., J. P. Corley, et al. 1964. Report to the Working Committee for Columbia River Studies on Progress Since September, 1962 for Projects Carried out by the General Electric Company at Hanford. HW-80649, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., J. J. Davis, and P. A. Olson. 1954. Studies on the Effect of the Hanford Reactors on Aquatic Life in the Columbia River. HW-33366, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., J. F. Honstead, and R. T. Jaske. 1970. Progress Report on Columbia River Studies July-December, 1969. BNWL-CC-2403, Pacific Northwest Laboratory, Richland Washington.

Foster, R. F., and D. McConnon. 1962. Relationships Between the Concentration of Radionuclides in Columbia River Water and Fish. HW-SA-2688, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and I. C. Nelson. 1961. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1960. HW-68435, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and I. C. Nelson. 1961. Evaluation of Radiological Conditions in the Vicinity of Hanford January-March, 1961. HW-70411, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and I. C. Nelson. 1961. Evaluation of Radiological Conditions in the Vicinity of Hanford April-June, 1961. HW-70552, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and I. C. Nelson. 1961. Evaluation of Radiological Conditions in the Vicinity of Hanford July-September, 1961. HW-71203, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and I. C. Nelson. 1962. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1961. HW-71999, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and I. C. Nelson. 1962. Evaluation of Radiological Conditions in the Vicinity of Hanford January-March, 1962. HW-73366, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., P. A. Olson, and R. H. Whittaker. 1952. Effect of Effluent Retention Time on Activity Densities in Aquatic Organisms. HW-24107, General Electric Company, Hanford Works, Richland, Washington.

Foster, R. F., and J. K. Soldat. 1966. Evaluation of the Exposure that Results from the Disposal of Radioactive Wastes into the Columbia River. BNWL-SA-0601, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F., J. K. Soldat, and T. H. Essig. 1966. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1965. BNWL-0316, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F., J. K. Soldat, and T. H. Essig. 1967. Evaluation of Radiological Conditions in the Vicinity of Hanford January-June, 1966. BNWL-0391, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F., and J. W. Vanderbeek. 1962. Evaluation of Radiological Conditions in the Vicinity of Hanford April-June, 1962. HW-74398, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and J. W. Vanderbeek. 1962. Evaluation of Radiological Conditions in the Vicinity of Hanford July-September, 1962. HW-75431, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and R. H. Wilson. 1963. Evaluation of Radiological Conditions in the Vicinity of Hanford January-March, 1963. HW-77533, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and R. H. Wilson. 1963. Evaluation of Radiological Conditions in the Vicinity of Hanford April-June, 1963. HW-78395, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and R. H. Wilson. 1964. Evaluation of Radiological Conditions in the Vicinity of Hanford January-June, 1964. HW-83723, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Foster, R. F., and R. H. Wilson. 1965. Evaluation of Exposure from Multiple Environmental Sources in the Environs of the Hanford Plants. BNSA-053, Pacific Northwest Laboratory, Richland, Washington.

Foster, R. F., and R. H. Wilson. 1965. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1964. BNWL-0090, Pacific Northwest Laboratory, Richland Washington.

Foster, R. F., and R. H. Wilson. 1965. Evaluation of Radiological Conditions in the Vicinity of Hanford January-June, 1965. BNWL-0165, Pacific Northwest Laboratory, Richland Washington.

Fryar, R. M. 1952. Technical Activities Report Water Studies, February, 1952. HW-23646, General Electric Company, Hanford Works, Richland, Washington.

Fuquay, J. J., and D. W. Pearce. 1969. Pacific Northwest Laboratory Report for 1968 to the USAEC Division of Biology and Medicine, Volume 1 Life Sciences; Part 2 Ecological Sciences. BNWL-1050-PT2, Pacific Northwest Laboratory, Richland, Washington.

Gamertsfelder, C. C. 1947. Effects on Surrounding Areas Caused by the Operations of the H.E.W. HW-7-5934, General Electric Company, Hanford Works, Richland, Washington.

Gamertsfelder, C. C., and J. K. Green. 1963. Hanford Radiological Sciences Research and Development Annual Report for 1962. HW-77609, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Gamertsfelder, C. C., and J. K. Green. 1964. Hanford Radiological Sciences Research and Development Annual Report for 1963. HW-81746, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Geier, R. G. 1965. Contamination Control - Columbia River - Monthly Report November, 1965. DUN-0255, Douglas United Nuclear, Inc., Richland, Washington.

General Electric Company. 1952. Aquatic Biology Laboratory (to December 31, 1951). HW-24800-97, General Electric Company, Hanford Works, Richland, Washington.

General Electric Company. 1953. Biology Research - Annual Report 1952. HW-28636, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

General Electric Company. 1954. Biology Research - Annual Report 1953. HW-30437, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

General Electric Company. 1964. Standard Practices Radiochemical Procedures for Off-Site Environmental Samples. HW-81277-RD, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

General Electric Company. Undated. Concentration of Gross Radioactivity in the Columbia River. GEH-18774, General Electric Company, Hanford Works, Richland, Washington.

Hall, R. B. 1957. Radioactivity in 100-Area Water. HW-52410, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Hall, R. B., and R. H. Wilson. 1962. Evaluation of Discharging Radioactive Wastes into Fresh Water Streams. HW-SA-2748, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Hanson, P. 1967. "Dissolved Chrome-51 in the Estuary." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1966 Through 30 June 1967, ed. J. E. McCauley, pp. 20-21. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Hanson, P. J., and W. O. Forster. 1969. "Measurement of Columbia River Flow Time from Hanford Reactors to Astoria, Oregon - Summer, 1966." Water Resources Research 5 (5):1129-1131.

Hanson, W. C., and R. L. Browning. 1953. "Biological Monitoring," pp. 180-185 in Biology Research - Annual Report 1952. HW-23636, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Haushild, W. L., G. R. Dempster, and H. H. Stevens. 1973. Distribution of Radionuclides in the Columbia River Streambed from the Nuclear Reactors, Hanford Reservation, to Longview, Washington. TID-26531, U.S. Department of Interior, Geological Survey, Portland, Oregon.

Healy, J. W. 1946. Accumulation of Radioactive Elements in Fish Immersed in Pile Effluent Water. HEW-3-3442, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Healy, J. W. 1947. Trend of Contamination in the Air, the Columbia River, Vegetation, and Waste at the HEW for the Period 3/25/47 - 6/30/47. HW-07317, General Electric Company, Hanford Works, Richland, Washington.

Healy, J. W. 1952. Effect of Purging during Pile Operation on the Effluent Water. HW-24578, General Electric Company, Hanford Works, Richland Works, Richland, Washington.

Healy, J. W., B. V. Andersen, H. V. Clukey, and J. K. Soldat. 1958. "Radiation Exposure to People in the Environs of a Major Production Atomic Energy Plant," pp. 309-318. In Proceedings of the Second United Nations International Conference on the Peaceful Uses of Atomic Energy, Waste Treatment, and Environmental Aspects of Atomic Energy. United Nations, Geneva.

Healy, J. W., and C. C. Gamertsfelder. 1947. H.I. Environs Reports - September, 1946 Through May, 1947. HW-7-5042, General Electric Company, Hanford Works, Richland, Washington.

Healy, J. W., and R. E. Rostenbach. 1955. Influence of Hanford Reactors on Domestic Use of Columbia River Water. HW-36862, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Herde, K. E. 1947. Radioactivity in Various Species of Fish from the Columbia and Yakima Rivers. HW-3-5501, General Electric Company, Hanford Works, Richland, Washington.

Herde, K. E. 1948. One Year Study of Radioactivity in Columbia River Fish. HW-11344, General Electric Company, Hanford Works, Richland, Washington.

Herde, K. E. 1948. Radioactivity of Pile Effluent and Its Biological Significance. HW-11509, General Electric Company, Hanford Works, Richland, Washington.

Honstead, J. F. 1952. Monitoring Survey - Richland to ARCO Period May 9-11, 1952. HW-24727, General Electric Company, Hanford Works, Richland, Washington.

Honstead, J. F. 1955. Isotopic Dilution of P-32 in the River. HW-36014, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Honstead, J. F. 1957. Dispersion of Dissolved Material in the Columbia River. HW-49008, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Honstead, J. F. 1966. Dietary Sources of Radioactivity for Richland Residents. BNWL-CC-0926, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F. 1969. Columbia River Studies July-December, 1968. BNWL-CC-1992, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F. 1969. Recreational Use of the Columbia River - Evaluation of Environmental Exposure. BNWL-CC-2299, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F. 1970. Quantitative Evaluation of Environmental Factors Affecting Population Exposure near Hanford. BNWL-SA-3203, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F., and D. N. Brady. 1965. Uptake and Retention of P-32 and Zn-65 from the Consumption of Columbia River Fish. BNSA-045, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F., and T. H. Essig. 1967. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1966, Appendices. BNWL-0439-APP, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F., and T. H. Essig. 1968. Evaluation of Radiological Conditions in the Vicinity of Hanford January-June, 1967. BNWL-0665, Pacific Northwest Laboratory, Richland, Washington.

Honstead, J. F., T. H. Essig, and J. K. Soldat. 1967. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1966. BNWL-0439, Pacific Northwest Laboratory, Richland, Washington.

Houston, J. R., and P. J. Blumer. 1979. Environmental Surveillance at Hanford for Calendar Year 1978. PNL-2932, Pacific Northwest Laboratory, Richland, Washington.

Houston, J. R., and P. J. Blumer. 1980. Environmental Surveillance at Hanford for Calendar Year 1979. PNL-3283, Pacific Northwest Laboratory, Richland, Washington.

Hungate, F. P., and E. G. Swezea. 1961. Hanford Biology Research Annual Report for 1960. HW-69500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Jaquish, R. E., and R. W. Bryce. 1989. Hanford Site Environmental Report for Calendar Year 1988. PNL-6825, Pacific Northwest Laboratory, Richland, Washington.

Jaquish, R. E., and R. W. Bryce. 1990. Hanford Site Environmental Report for Calendar Year 1989. PNL-7346, Pacific Northwest Laboratory, Richland, Washington.

Jaquish, R. E., and P. J. Mitchell, eds. 1987. Environmental Monitoring at Hanford for 1987. PNL-6464, Pacific Northwest Laboratory, Richland, Washington.

Jennings, C. D., and C. Osterberg. 1969. "Sediment Radioactivity in the Columbia River Estuary." In Symposium on Radioecology, Proceedings of the Second National Symposium, eds. D. J. Nelson and F. C. Evans. National Technical Information Service, Springfield, Virginia.

Johnson, V. G. 1966. "Retention of Zinc-65 by Columbia River Sediment." Master's thesis, Oregon State University, Corvallis, Oregon.

Johnson, V. G. 1978. "Decline of Radioactivity in the Columbia River and Estuary: Rates and Mechanisms." Ph.D. disseration, Oregon State University, Corvallis, Oregon.

Johnson, V., N. Cutshall, and C. Osterberg. 1964. "Geochemical Studies." In Radioanalysis of Oceanic Organisms in the Pacific Ocean Off Oregon, Progress Report, 1 March 1963 Through 29 February 1964, ed. C. L. Osterberg, pp. 55-66. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Johnson, V., N. Cutshall, and C. Osterberg. 1966. "Retention of ⁶⁵Zn by Columbia River Sediment." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 Through 30 June 1966, ed. J. E. McCauley, pp. 21-28. RLO-1750-8, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Johnson, V., N. Cutshall, and C. Osterberg. 1967. "Retention of ⁶⁵Zn by Columbia River Sediment." Water Resources Research 3(1):99-102.

Jones, W. M., C. D. Jennings, and N. H. Cutshall. 1975. Cycling of Fe-55 and Zn-65 in Columbia River Carp Following Reactor Shutdown. IAEA-SM-197-19. International Atomic Energy Agency, Vienna, Austria.

Jonsson, G., and A. H. Seymour. 1965. Gamma-Emitting Radionuclides in Dover Sole Caught Off the Mouth of the Columbia River. Laboratory of Radiation Biology, University of Washington, Seattle, Washington.

Junkins, R. L. 1960. Analysis of Radiological Data for the Month of October, 1960. HW-67439, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Junkins, R. L., E. C. Watson, I. C. Nelson, G. E. Backman, and R. C. Henle. 1960. Evaluation of Radiological Conditions in the Vicinity of Hanford July Through September 1960. HW-67390, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Junkins, R. L., E. C. Watson, I. C. Nelson, and R. C. Henle. 1959. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1959. HW-64371, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kennedy, V. C. 1965. Mineralogy and Cation-Exchange Capacity of Sediments from Selected Streams. USGS Professional Paper 433-D, U.S. Department of the Interior, Washington, D.C.

Klingeman, P. C., and W. J. Kaufman. 1965. Transport of Radionuclides with Suspended Sediment in Estuarine Systems. Report 65-15, University of California, Sanitary Engineering Laboratory, Berkeley, California.

Kornberg, H. A. 1951. Columbia River Symposium, Second Meeting, 21-22 May, 1951, Richland, Washington. HW-22181, General Electric Company, Hanford Works, Richland, Washington.

Kornberg, H. A. 1952. Biology Research - Annual Report 1951. HW-25021, General Electric Company, Hanford Works, Richland, Washington.

Kornberg, H. A. 1955. Biology Research - Annual Report 1954. HW-35917, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A. 1956. Biology Research - Annual Report 1955. HW-41500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A. 1957. Biology Research - Annual Report 1956. HW-47500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A. 1958. Hanford Biology Research Annual Report for 1957. HW-53500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A. 1964. Progress in Biological Research. HW-83613, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A., and J. J. Davis. 1959. Hanford Biology Research Annual Report for 1958. HW-59500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A., and E. G. Swezea. 1962. Hanford Biology Research Annual Report for 1961. HW-72500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A., and E. G. Swezea. 1963. Hanford Biology Research Annual Report for 1962. HW-76000, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A., and E. G. Swezea. 1964. Hanford Biology Research Annual Report for 1963. HW-80500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kornberg, H. A., and D. E. Warner. 1960. Hanford Biology Research Annual Report for 1959. HW-65500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Kramer, H. A. 1948. Columbia River Flood 1948. HW-31882, General Electric Company, Hanford Works, Richland, Washington.

Kujala, N. F., I. L. Larsen, and C. L. Osterberg. 1969. "Radioisotope Measurements of the Viscera of Pacific Salmon." In Symposium on Radioecology, Proceedings of the Second National Symposium, eds. D. J. Nelson and F. C. Evans, pp. 440-449. National Technical Information Service, Springfield, Virginia.

Kujala, N., and C. L. Osterberg. 1967. "Columbia River Estuary Fishes." In Vol. 1, Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1966 Through 30 June 1967, ed. J. E. McCauley, pp. 10-12. RLO-1750-22, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Lutton, J. M., and W. C. A. Woods. 1953. Columbia River Water as a Pile Coolant. HW-28467, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Matsumoto, W. Y. 1955. Radioactive Contamination in Reactor Effluent Using Unfiltered Columbia River Water as a Coolant. HW-40699, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McCauley, J. E., ed. 1969. Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1968 - 30 June 1969. RLO-1750-54, Oregon State University, Corvallis, Oregon.

McCauley, J. E., and C. L. Osterberg. 1966. Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 - 30 June 1966. RLO-1750-08-V1, Oregon State University, Corvallis, Oregon.

McConiga, M. W. 1960. MPC'S for Ingestion of Rare Earths in Columbia River Water. HW-66026, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McConiga, M. W., J. M. Selby, and J. K. Soldat. 1958. Radioactive Contamination in the Hanford Environs for the Period July, August, September, 1957. HW-54841, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McConiga, M. W., J. M. Selby, and J. K. Soldat. 1958. Radioactive Contamination in the Hanford Environs for the Period October, November, December, 1957. HW-54842, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McConiga, M. W., and J. K. Soldat. 1957. Radioactive Contamination in the Hanford Environs for the Period January, February, March, 1957. HW-51009, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McConiga, M. W., and J. K. Soldat. 1957. Radioactive Contamination in the Hanford Environs for the Period April, May, June, 1957. HW-52803, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

McCormack, W. D., and J.M.V. Carlile. 1984. Investigation of Ground-Water Seepage from the Hanford Shoreline of the Columbia River. PNL-5289, Pacific Northwest Laboratory, Richland, Washington.

McMurray, P. R. 1956. Columbia River Aspects of Increased Production. HW-41049, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Mooney, R. R. 1979. State of Washington Environmental Radiation Program, Eighteenth Annual Report, July, 1978-June, 1979. State of Washington, Department of Social and Health Services, Health Services Division, Olympia, Washington.

Mooney, R. R. 1980. State of Washington Environmental Radiation Program, Nineteenth Annual Report, July, 1979-June, 1980. State of Washington, Department of Social and Health Services, Health Services Division, Olympia, Washington.

Mooney, R. R. 1981. State of Washington Environmental Radiation Program, Twentieth Annual Report, July, 1980-June, 1981. State of Washington, Department of Social and Health Services, Health Services Division, Olympia, Washington.

Nakatani, R. E., and R. F. Foster. 1966. Hanford Temperature Effects on Columbia River Fishes. BNWL-CC-0591, Pacific Northwest Laboratory, Richland, Washington.

Nees, W. L., and J. P. Corley. 1974. Environmental Surveillance at Hanford for Calendar Year 1973. BNWL-1811, Pacific Northwest Laboratory, Richland, Washington.

Nees, W. L., and J. P. Corley. 1975. Environmental Status of the Hanford Reservation for Calendar Year 1973. BNWL-B-336, Pacific Northwest Laboratory, Richland, Washington.

Nelson, I. C. 1963. Ringold - A Hanford Environmental Study. HW-SA-2882, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nelson, I. C., and R. F. Foster. 1964. Ringold (Farms) - A Hanford Environmental Study. HW-78262-REV, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nelson, J. L., and W. L. Haushild. 1968. Accumulation of Radionuclides in Bed Sediments of McNary Reservoir of the Columbia River. BNWL-SA-1730, Pacific Northwest Laboratory, Richland, Washington.

Nelson, J. L., and W. L. Haushild. 1969. Accumulation of Radionuclides in Bed Sediments of the Columbia River Between Hanford Reactors and McNary Dam. BNWL-SA-2571, Pacific Northwest Laboratory, Richland, Washington.

Nelson, J. L., R. W. Perkins, and W. L. Haushild. 1965. Determination of Columbia River Flow Times Using Radioactive Tracers Introduced by the Hanford Reactors. BNSA-176, Pacific Northwest Laboratory, Richland Washington.

Nelson, J. L., R. W. Perkins, and J. M. Nielsen. 1964. Progress in Studies of Radionuclides in Columbia River Sediments. HW-83614, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nelson, J. L., R. W. Perkins, J. M. Nielsen, and W. L. Haushild. 1966. Reactions of Radionuclides from the Hanford Reactors with Columbia River Sediments. BNWL-SA-0317, Pacific Northwest Laboratory, Richland, Washington.

Nielsen, J. M. 1973. Pacific Northwest Laboratory Annual Report for 1972 to the USAEC Division of Biomedical and Environmental Research Volume 2 Physical Sciences, Part 2 Radiological Sciences. BNWL-1751-PT2, Pacific Northwest Laboratory, Richland, Washington.

Nielsen, J. M., and R. W. Perkins. 1957. Depletion of Radioisotopes from the Columbia River by Natural Processes. HW-52908, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Nielsen, J. M., and R. W. Perkins. 1962. Removal of Radioisotopes from the Columbia River by Natural Processes. HW-SA-2411, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Norton, H. T. 1957. Turbulent Diffusion of River Contaminants. HW-49195, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

O'Farrell, T. P., and V. K. Christensen. 1971. Bibliography of Environmental Research: Ecosystems Department 1952-1970. BNWL-1549, Pacific Northwest Laboratory, Richland, Washington.

Olson, P. A. 1947. Fish Laboratory Report for First Half of July, 1947. HW-07132, General Electric Company, Hanford Works, Richland, Washington.

Olson, P. A. 1947. Fish Laboratory Report for Last Half of July, 1947. HW-07267, General Electric Company, Hanford Works, Richland, Washington.

- Olson, P. A. 1947. Fish Laboratory Report for First Half of August, 1947. HW-07325, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A. 1947. Fish Laboratory Report for Last Half of August, 1947. HW-07432, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A. 1947. Fish Laboratory Report for First Half of September, 1947. HW-07547, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A. 1948. Fish and Fish Problems of the Hanford Reservation. HW-11642, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A. 1948. Some Effects of Pile Area Effluent Water on Young Silver Salmon. HW-08944, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A., and R. H. Foster. 1946. Fish Laboratory Report - December, 1946 Through June, 1947. HW-3-5160, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A., and R. F. Foster. 1952. Accumulation of Radioactivity in Columbia River Fish in the Vicinity of the Hanford Works. HW-23093, General Electric Company, Hanford Works, Richland, Washington.
- Olson, P. A., and R. F. Foster. 1958. Effect of Separan on Rainbow Trout. HW-55292, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.
- Onishi, Y. 1977. Mathematical Simulation of Sediment and Radionuclide Transport in the Columbia River. BNWL-2228, Pacific Northwest Laboratory, Richland, Washington.
- Osterberg, C. 1962. "Zn-65 Content of Salps and Euphausiids." Limnology and Oceanography 7(4):478-479.
- Osterberg, C. 1965. "Radioactivity from the Columbia River." Ocean Science and Ocean Engineering 2:968-979.
- Osterberg, C., N. Cutshall, and J. Cronin. 1965. "Chromium-51 as a Radioactive Tracer of Columbia River Water at Sea." Science 150(3703):1585-1587.
- Osterberg, C., D. Jennings, and N. Cutshall. 1964. "In situ Measurement of Gamma Emitters in the Columbia River." In Radioanalysis of Oceanic Organisms in the Pacific Ocean Off Oregon, Progress Report, 1 March 1963 Through 29 February 1964, ed. C. L. Osterberg, pp. 13-29. Department of Oceanography, Oregon State University, Corvallis, Oregon.
- Osterberg, C., L. D. Kulm, and J. V. Byrne. 1963. "Gamma Emitters in Marine Sediments near the Columbia River." Science 139(3558):916-917.

Osterberg, C., and L. Larsen. 1964. "Radionuclides in Migratory Albacore Tuna." In Radioanalysis of Oceanic Organisms in the Pacific Ocean Off Oregon, Progress Report, 1 March 1963 Through 29 February 1964, ed. L. Osterberg, pp. 32-34. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Osterberg, C., J. Pattullo, and W. Pearcy. 1963. "Zinc-65 in Euphausiids as Related to Columbia River Water Off the Oregon Coast." In Species Composition and Distribution of Marine Nekton in the Pacific Ocean off Oregon, Final Progress Report, 1 September 1961 through 1 September 1963, pp. 107-129. Oregon State University, Corvallis, Oregon.

Osterberg, C., W. G. Pearcy, and H. Curl Jr. 1963. "Radioactivity and Its Relationship to Oceanic Food Chains." J. of Marine Research 22(1):2-12.

Osterberg, C., and J. Wiese. 1964. "Trace Element Analysis." In Radioanalysis of Oceanic Organisms in the Pacific Ocean Of Oregon, Progress Report, 1 March 1963 through 29 February 1964, ed. C. L. Osterberg, pp. 37-44. Oregon State University, Corvallis, Oregon.

Paas, H. J. 1951. Radioactive Contamination in the Environs of the Hanford Works for the Period July, August, September, 1951. HW-23133, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. 1952. Radioactive Contamination in the Environs of the Hanford Works for the Period October, November, December, 1951. HW-24203, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. 1952. Radioactive Contamination in the Environs of the Hanford Works for the Period January, February, March, 1952. HW-25866, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. 1952. Radioactive Contamination in the Environs of the Hanford Works for the Period April, May, June, 1952. HW-26493, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. 1953. Radioactive Contamination in the Environs of the Hanford Works for the Period July, August, September, 1952. HW-27510, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J. 1953. Radioactive Contamination in the Environs of the Hanford Works for the Period October, November, December, 1952. HW-27641, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J. 1953. Radioactive Contamination in the Hanford Environs for the Period April, May, June, 1953. HW-29514, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J. 1953. Radioactive Contamination in the Hanford Environs for the Period July, August, September, 1953. HW-30174, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J. 1954. Radioactive Contamination in the Hanford Environs for the Period October, November, December, 1953. HW-30744, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J., and G. E. Pilcher. 1954. Radioactive Contamination in the Hanford Environs for the Period April, May, June, 1954. HW-33896, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Paas, H. J., and W. Singlevich. 1950. Radioactive Contamination in the Environs of the Hanford Works for the Period April, May, June, 1949. HW-17434, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J., and W. Singlevich. 1950. Radioactive Contamination in the Environs of the Hanford Works for the Period July, August, September, 1949. HW-18615, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J., and W. Singlevich. 1950. Radioactive Contamination in the Environs of the Hanford Works for the Period October, November, December, 1949. HW-17003, General Electric Company, Hanford Works, Richland, Washington.

Paas, H. J. and W. Singlevich. 1950. Radioactive Contamination in the Environs of the Hanford Works for the Period April, May, June, 1950. HW-19454, General Electric Company, Hanford Works, Richland, Washington.

Park, K., M. J. George, Y. Miyake, K. Saruhashi, Y. Katsauragi, and T. Kanazawa. 1965. "Strontium-90 and Caesium-137 in Columbia River Plume, July 1964." Nature 208(5015):1084-1085.

Parker, H. M. 1945. Water Sampling for Radioactivity. HW-7-2071, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Parker, H. M. 1946. Tolerance Dose: Interpretation of Permissible Daily Dose. DUH-12744, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Parker, H. M. 1947. Action Taken on the Spot Contamination in the Separations Plant Areas. HW-07920, General Electric Company, Hanford Works, Richland, Washington.

Parker, H. M. 1954. Columbia River Situation. HW-32809, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Patterson, C. M. 1947. H.I. Report for the 100 and 300 Areas for the Period Ending September 15, 1947. HW-07564, General Electric Company, Hanford Works, Richland, Washington.

Pearce, D. W., and J. K. Green. 1965. Hanford Radiological Sciences Research and Development Annual Report for 1964. BNWL-0036, Pacific Northwest Laboratory, Richland, Washington.

Pearcy, W. G. 1966. "Specific Activities of Oceanic Animals." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1965 Through 30 June 1966, ed. J. E. McCauley, pp. 145-147. RLO-1750-8, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Pearcy, W. G. 1967. "Radioecology of Benthic Fishes." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1966 Through 30 June 1967, ed. J. E. McCauley, pp. 102-110. RLO-1750-22, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Pearcy, W. G., and R. R. Clays. 1972. "Zinc-65 and DDT Residues in Albacore Tuna Off Oregon in 1969." California Marine Research Committee CalCOFI Report 16:66-73.

Pearcy, W. G., and R. J. Eagle. 1968. "Radio-Zinc, Total Zinc, and Specific Activities of the Hake, *Merluccius Productus*." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1967 Through 30 June 1968, ed. J. E. McCauley, pp. 161-162. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Pearcy, W. G., and W. O. Forster. 1968. "Zinc-65 Specific Activities: Pelagic Animals and Surface Water in the Vicinity of the Columbia River Plume." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 Through 30 June 1968, ed. J. E. McCauley, pp. 154-159. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Pearcy, W. G., and C. L. Osterberg. 1964. "Vertical Distribution of Radionuclides as Measured in Oceanic Animals." Nature 204(4957):440-441.

Pearcy, W. G., and C. L. Osterberg. 1967. "Depth, Diel, Seasonal, and Geographic Variations in Zinc-65 of Midwater Animals Off Oregon." International Journal of Oceanology and Limnology 1:103-116.

Pearcy, W. G., and C. L. Osterberg. 1968. "Zinc-65 and Manganese-54 in Albacore, *Thunnus alalunga*, from the West Coast of North America." Limnology and Oceanography 13(3):490-498.

Pendleton, R. C. 1957. Contamination of River Organisms Grown in Retained and Nonretained Reactor Effluent. HW-48381, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Perkins, R. W., J. L. Nelson, and W. L. Haushild. 1965. Behavior and Transport of Radionuclides in the Columbia River Between Hanford and the Ocean. BNSA-16, Pacific Northwest Laboratory, Richland, Washington.

Peterson, D. S., and R. R. Mooney. 1983. State of Washington Environmental Radiation Program, Twenty-Second Annual Report, July, 1982-December, 1983. State of Washington, Department of Social and Health Services, Division of Health, Olympia, Washington.

Peterson, D. S., and R. R. Mooney. 1986. State of Washington Environmental Radiation Program, Twenty-Fourth Annual Report, January, 1985-December, 1985. State of Washington, Department of Social and Health Services, Division of Health, Olympia, Washington.

Pilcher, G. E. 1952. Statistical Analysis of Measured River Contamination at Selected Shoreline Locations. HW-23642, General Electric Company, Hanford Works, Richland, Washington.

Pilcher, G. E., J. K. Soldat, and Z. E. Carey. 1955. Radioactive Contamination in the Hanford Environs for the Period July, August, September, 1954. HW-36504, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Price, K. R. 1986. Environmental Monitoring at Hanford for 1985. PNL-5817, Pacific Northwest Laboratory, Richland, Washington.

Price, K. R., J.M.V. Carlile, R. L. Dirkes, R. E. Jaquish, M. S. Trevathan, and R. K. Woodruff. 1985. Environmental Monitoring at Hanford for 1984. PNL-5407, Pacific Northwest Laboratory, Richland, Washington.

Price, K. R., J.M.V. Carlile, R. L. Dirkes, and M. S. Trevathan. 1984. Environmental Surveillance at Hanford for Calendar Year 1983. PNL-5038, Pacific Northwest Laboratory, Richland, Washington.

Proceedings of the Second United Nations International Conference on the Peaceful Uses of Atomic Energy, Waste Treatment, and Environmental Aspects of Atomic Energy. 1958. United Nations, New York.

Reiman, R. T., and T. S. Dahlstrom. 1990. An Aerial Radiological Survey of the Hanford Site and Surrounding Area, Richland, Washington, July-August, 1988. EGG-10617-1062, EG&G, Las Vegas, Nevada.

Renfro, W. C. 1968. "Transfer of Radionuclides through Marine Food Webs." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 June 1967 Through 30 July 1968, ed. J. E. McCauley, pp. 17-22. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Renfro, W. C. 1968. "Columbia River Estuary Fishes." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, July 1 1967-June 30 1968, ed. J. E. McCauley, pp. 27-31. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Renfro, W. C. 1968. "Transfer of Radionuclides Through Marine Food Webs." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967-30 June 1968, ed. J. E. McCauley, pp. 17-22. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Renfro, W. C. 1970. "Ecological Half-Lives of ^{65}Zn in Willapa Bay Biota." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, July 1 1969-June 30 1970, ed. J. E. McCauley, pp. 45-50. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Renfro, W. C., and N. D. Farrow. 1970. "Cycling of Radionuclides in the Columbia River Estuary." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1969-30 June 1970, ed. J. E. McCauley, pp. 37-41. Department of Oceanography, Oregon State University, Corvallis, Oregon.

Renfro, W. C., and C. Osterberg. 1969. "Radiozinc Decline in Starry Flounders After Temporary Shutdown of Hanford Reactors." In Symposium on Radioecology, Proceedings of the Second National Symposium, eds. D. J. Nelson and F. C. Evans, pp. 372-379. National Technical Information Service, Springfield, Virginia.

Renfro, W. C., and M. Thompson. 1967. "Transfer of Radionuclides Through Estuarine Food Webs." In Ecological Studies of Radioactivity in the Columbia River and Adjacent Pacific Ocean, Progress Report, 1 July 1966 Through 30 June 1967, ed. J. E. McCauley, pp. 15-18. RLO-1750-22, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Rieck, H. G., and R. W. Perkins. 1956. Radium Determinations in Soil, Vegetation, and Water by Radon Counting. HW-32210-REV, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Robertson, D. E., and J. J. Fix. 1977. Association of Hanford Origin Radionuclides with Columbia River Sediment. BNWL-2305, Pacific Northwest Laboratory, Richland, Washington.

Robertson, D. E., W. O. Forster, and H. G. Rieck. 1968. "An Environmental and Radioecological Study at a Northeast Pacific Ocean Station 350 Miles Off Newport, Oregon." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 Through 30 June 1968, ed. J. E. McCauley, pp. 51-67. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Rostenbach, R. E. 1954. Radioactivity Levels of the Columbia River below Richland, Washington for the Period July, August, September, 1953. HW-30834, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1954. Radioactivity Levels of the Columbia River below Richland, Washington for the Period October, November, December, 1953. HW-31558, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1954. Radioactivity Levels of the Columbia River below Richland, Washington for the Period January, February, March, 1954. HW-32449, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1954. Radioactivity Levels of the Columbia River Below Richland, Washington for the Period April, May, June, 1954. HW-34135, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1955. Radioactivity Levels of the Columbia River Below Richland, Washington for the Period July, August, and September, 1954. HW-38048, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1955. Radioactivity Levels of the Columbia River Below Richland, Washington for the Period October, November, and December, 1954. HW-38049, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1955. Radioactivity Levels of the Columbia River Below Richland, Washington for the Period January, February, March, 1955. HW-38077, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Rostenbach, R. E. 1955. Radioactivity Levels of the Columbia River Below Richland, Washington, April, May, June, 1955. HW-39632, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Sayre, W. W., H. P. Guy, and A. R. Chamberlain. 1963. Uptake and Transport of Radionuclides by Stream Sediments. USGS Professional Paper 433-A, U.S. Department of the Interior, Washington, D.C.

Schiffman, R. H. 1961. Uptake of Strontium from Diet and Water by Rainbow Trout. HW-72107, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Schoning, R. W., and T. R. Merrill. 1951. Indian Dip-Net Fishery at Celilo Falls on the Columbia River. OFC-C17, Oregon Fish Commission, Portland, Oregon.

Selby, J. M., and J. K. Soldat. 1958. Summary of Environmental Contamination Incidents at Hanford 1952 - 1957. HW-54636, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Seymour, A. H. 1960. A Program to Study the Radionuclides in Water, Bottom and Biological Samples from near the Mouth of the Columbia River. Laboratory of Radiation Biology, University of Washington, Seattle, Washington.

Seymour, A. H. 1966. "Accumulation and Loss of Zinc-65 by Oysters in a Natural Environment." In Proceedings of the Symposium on the Disposal of Radioactive Wastes into Seas, Oceans, and Surface Waters, pp. 605-619. International Atomic Energy Agency, Vienna, Austria.

Seymour, A. H. 1970. Columbia River Studies, Annual Progress Report, 1969-1970. Laboratory of Radiation Ecology, University of Washington, Seattle, Washington.

Seymour, A. H. 1970. Columbia River Studies, Annual Progress Report, 1969-1970. RLO-2225-T1-02, Laboratory of Radiation Ecology, University of Washington, Seattle, Washington.

Seymour, A. H. 1970. Radiological Aspects of Nuclear Power and the Aquatic Environment. RLO-2225-T1-01, Laboratory of Radiation Ecology, University of Washington, Seattle, Washington.

Seymour, A. H., and V. A. Nelson. 1971. Decline of Zinc-65 in Marine Mussels Following the Shutdown of Hanford Reactors. RLO-2225-T1-09, Laboratory of Radiation Ecology, University of Washington, Seattle, Washington.

Seymour, F. P. 1946. Study of Total Amounts of Active Waste Released in all Manners by the H.E.W. Process to Date. HW-7-5463, General Electric Company, Hanford Works, Richland, Washington.

Silker, W. B. 1958. Strontium-90 Concentrations in the Hanford Environs. HW-55117, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Singlevich, W. 1947. Trend of Contamination in the Air, the Columbia River, Rain, Sanitary Water, Vegetation, and Wastes, at the Hanford Works for the Period July, August, September, 1947. HW-08549, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of December, 1947. HW-08486, General Electric Company, Hanford Works, Richland Washington.

Singlevich, W. 1948. Trend of Contamination in the Air, Columbia River, Rain, Sanitary Water, Vegetation and Wastes at the Hanford Works and Vicinity for the Period October, November, December, 1947. HW-09496, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of January, 1948. HW-08783, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of February, 1948. HW-09103, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of March, 1948. HW-09590, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. Radioactive Contamination in the Environs of the Hanford Works and Vicinity for the Period January, February, March, 1948. HW-10242, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of April, 1948. HW-09730, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. Radioactive Contamination in the Columbia River and in the Air and Radiation Levels Measured in the Air at Hanford Works and Vicinity for 1945, 1946, 1947, and Early 1948. HW-09871, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of May, 1948. HW-10049, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. Radioactive Contamination in the Environs of the Hanford Works for the Period April, May, June, 1948. HW-11333, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of July, 1948. HW-10730, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of September, 1948. HW-11264, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1948. H.I. Environs Report for Month of October, 1948. HW-11534, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. Comparison of Background Measurements for Radioactivity in Some Water Sources at the Hanford Works with Sources from the Colorado Raw Materials Office Area. HW-13501, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. Radioactive Contamination in the Environs of the Hanford Works for the Period July, August, September, 1948. HW-12677, General Electric Company, Hanford Works, Richland Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of December, 1948. HW-12131, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of January, 1949. HW-12325, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of February, 1949. HW-12629, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of March, 1949. HW-12948, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of April, 1949. HW-13175, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of May, 1949. HW-13570, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of June, 1949. HW-13804, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of July, 1949. HW-14040, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of August, 1949. HW-14401, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of September, 1949. HW-14598, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of October, 1949. HW-14953, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1949. H.I. Environs Report for Month of November, 1949. HW-15377, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of January, 1950. HW-16015, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of February, 1950. HW-17150, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of March, 1950. HW-17409, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of April, 1950. HW-17657, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of May, 1950. HW-17944, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of June, 1950. HW-18242, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of July, 1950. HW-18447, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of August, 1950. HW-18741, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of September, 1950. HW-19046, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of October, 1950. HW-19332, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1950. H.I. Environs Report for Month of November, 1950. HW-19625, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1951. H.I. Environs Report for Month of December, 1950. HW-19896, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1951. H.I. Environs Report for Month of January, 1951. HW-20286, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W. 1951. Review of Columbia River Studies Program - H. I. Development Division. HW-21074, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W., and H. J. Paas. 1949. Radioactive Contamination in the Environs of the Hanford Works for the Period January, February, March, 1949. HW-14243, General Electric Company, Hanford Works, Richland, Washington.

Singlevich, W., and C. M. Patterson. 1948. H.I. Environs Report for Month of November, 1948. HW-11833, General Electric Company, Hanford Works, Richland, Washington.

Snyder, J. H. 1955. Effect of Low River Flows on the Operation of Hanford Production Works Resulting from Operation of the Proposed Priest Rapids Dam. HW-39449, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Soldat, J. K. 1968. Statistical Study of the Habits of Fishermen Utilizing the Columbia River Below Hanford. BNWL-SA-1451, Pacific Northwest Laboratory, Richland, Washington.

Soldat, J. K., G. E. Pilcher, D. L. Reid, and Z. E. Carey. 1955. Radioactive Contamination in the Hanford Environs for the Period October, November, December, 1954. HW-36505, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Soldat, J. K., and R. C. Thorburn. 1951. Radiological Survey of the Pasco, Washington Water Treatment Plant. HW-22862, General Electric Company, Hanford Works, Richland, Washington.

Speer, D. R., J. J. Fix, P. J. Blumer. 1976. Environmental Surveillance at Hanford for Calendar Year 1975. BNWL-1979, Pacific Northwest Laboratory, Richland, Washington.

Sula, M. J., and P. J. Blumer. 1981. Environmental Surveillance at Hanford for Calendar Year 1980. PNL-3728, Pacific Northwest Laboratory, Richland, Washington.

Sula, J. M., J.M.V. Carlile, K. R. Price, and W. D. McCormack. 1983. Environmental Surveillance at Hanford for Calendar Year 1982. PNL-4657, Pacific Northwest Laboratory, Richland, Washington.

Sula, M. J., W. D. McCormack, R. L. Dirkes, K. R. Price, and P. A. Eddy. 1982. Environmental Surveillance at Hanford for Calendar Year 1981. PNL-4211, Pacific Northwest Laboratory, Richland, Washington.

Tennant, D. A., and W. O. Forster. 1968. "Seasonal Variation of ^{65}Zn , ^{54}Mn , and ^{51}Cr Radioactivity and Specific Activity in Tissues of Cancer Magister Dana." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, July 1, 1967 Through June 30, 1968, ed. J. E. McCauley, pp. 10-16. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Tennant, D. A., and W. O. Forster. 1970. "Seasonal Variation and Distribution of ^{65}Zn , ^{54}Mn , and ^{51}Cr in Tissues of the Crab Cancer Magister Dana." Health Physics 18(6):649-657.

Thomas, C. W., G. E. Pilcher, K. C. Knoll, and B. V. Andersen. 1955. Radioactive Contamination in the Hanford Environs for the Period January, February, March, 1955. HW-36506, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Thomas, C. W., V. K. Schlatter, G. E. Pilcher, B. V. Andersen, J. K. Soldat, and D. L. Reid. 1955. Radioactive Contamination in the Hanford Environs for the Period April, May, June, 1955. HW-38566, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Tipton, W. J. 1975. Aerial Radiological Survey of the U.S. ERDA's Hanford Reservation (Survey Period: 1973-1974). EGG-1183-1661, EG&G, Las Vegas, Nevada.

Turner, L. D. 1947. Trend of Contamination Observed in the Air, Columbia River, and Vegetation, at the Hanford Engineer Works for 1946. HW-3-5402, General Electric Company, Hanford Works, Richland Washington.

Vermeere, W. R. 1968. "Biological Tracer of Radioactive Columbia River Effluent upon Entering the Marine Environment." In Ecological Studies of Radioactivity in the Columbia River Estuary and Adjacent Pacific Ocean, Progress Report, 1 July 1967 Through 30 June 1968, ed. J. E. McCauley, pp. 109-115. RLO-1750-28, Department of Oceanography, Oregon State University, Corvallis, Oregon.

Watson, D. G. 1960. "Chinook Salmon Spawning in the Vicinity of Hanford - 1959," pp. 180-182 in Hanford Biology Research - Annual Report for 1959. HW-65500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, D. G., C. E. Cushing, C. Coutant, and W. L. Templeton. 1970. Cycling of Radionuclides in Columbia River Biota. BNWL-SA-3364, Pacific Northwest Laboratory, Richland, Washington.

Watson, D. G., C. E. Cushing, C. C. Coutant, and W. L. Templeton. 1970. Radioecological Studies on the Columbia River. BNWL-1377-PT1, Pacific Northwest Laboratory, Richland, Washington.

Watson, D. G. 1952. "Observations on Spawning and Migration of Chinook Salmon in The Columbia River in the Vicinity of Hanford Works," pp. 14-18, in Biology Research - Annual Report 1951. HW-25021, General Electric Company, Hanford Works, Richland, Washington.

Watson, D. G. 1970. Fall Chinook Salmon Spawning in the Columbia River near Hanford 1947-1969. BNWL-1515, Pacific Northwest Laboratory, Richland, Washington.

Watson, D. G., and J. J. Davis. 1957. Concentration of Radioisotopes in Columbia River Whitefish in the Vicinity of the Hanford Atomic Products Operation. HW-48523, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, D. G., J. J. Davis, and W. C. Hanson. 1960. "Zinc-65 in Marine Mollusks near the Mouth of the Columbia River," pp. 183-187 in Hanford Biology Research - Annual Report 1959. HW-65500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, D. G., J. J. Davis, and W. C. Hanson. 1960. "Dispersion of Gamma Emitters in Marine Organisms near the Mouth of the Columbia River," pp. 151-155 in Hanford Biology Research - Annual Report 1960. HW-69500, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, D. G., J. J. Davis, and W. C. Hanson. 1962. Interspecies Differences in Accumulation Gamma Emitters by Marine Organisms near the Columbia River Mouth. HW-SA-2857, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, E. C. 1960. Analysis of Radiological Data for the Month of November, 1960. HW-67752, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, E. C. 1961. Analysis of Radiological Data for the Month of December, 1960. HW-68127, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Watson, E. C. 1961. Analysis of Radiological Data for the Month of January, 1961. HW-68471, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

West, J. M. 1946. Activity of Pile Discharge Water. HW-7-4673, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

West, J. M. 1946. Activity of Pile Discharge Water, Water Activity as a Function of Power Level. HW-7-3692, E. I. duPont de Nemours & Company, Hanford Engineer Works, Richland, Washington.

Whelan, G., D. W. Damschen, and R. D. Brockhaus. 1987. Columbia River Statistical Update Model - Version 4.0 (COLSTAT4): Background Documentation and User's Guide. PNL-6041, Pacific Northwest Laboratory, Richland, Washington.

Wiese, G., and C. Osterberg. 1964. "Radiological Study of the Columbia River Estuary." In Radioanalysis of Oceanic Organisms in the Pacific Ocean of Oregon, Progress Report, 1 March 1963 through 29 February 1964, ed. C. L. Osterberg, pp. 78-85. Oregon State University, Corvallis, Oregon.

Wildung, R. E., R. C. Routson, and R. L. Schmidt. 1972. Seasonal Changes in Particle Size Distribution, Composition, and Strontium Exchange Capacity of Particulate Matter Suspended in the Columbia River. BNWL-1638, Pacific Northwest Laboratory, Richland, Washington.

Wilson, C. B., and T. H. Essig. 1970. Environmental Status of the Hanford Reservation for January-June, 1969. BNWL-CC-2478, Pacific Northwest Laboratory, Richland, Washington.

Wilson, C. B., and T. H. Essig. 1970. Environmental Status of the Hanford Reservation for July-December, 1969. BNWL-CC-2583, Pacific Northwest Laboratory, Richland, Washington.

Wilson, R. H. 1962. Columbia River Environmental Monitoring Program. HW-75076, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Wilson, R. H., and R. F. Foster. 1963. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1962. HW-76526, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Wilson, R. H., and R. F. Foster. 1964. Evaluation of Radiological Conditions in the Vicinity of Hanford for 1963. HW-80991, General Electric Company, Hanford Atomic Products Operation, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Status of the Hanford Reservation for November-December, 1967. BNWL-CC-1197-6, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Status of the Hanford Reservation for January-June, 1968. BNWL-CC-1850, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1969. Environmental Status of the Hanford Reservation for July-December, 1968. BNWL-CC-2026, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for January, 1968. BNWL-0778-01, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for February, 1968. BNWL-0778-02, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for March, 1968. BNWL-0778-03, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for April, 1968. BNWL-0778-04, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for May, 1968. BNWL-0778-05, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for June, 1968. BNWL-0778-06, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for July, 1968. BNWL-0778-07, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for August, 1968. BNWL-0778-08, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for September, 1968. BNWL-0778-09, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1968. Environmental Surveillance in the Vicinity of Hanford for October, 1968. BNWL-0778-10, Pacific Northwest Laboratory, Richland, Washington.

Wooldridge, C. B. 1969. Environmental Surveillance in the Vicinity of Hanford for December, 1968. BNWL-0778-12, Pacific Northwest Laboratory, Richland, Washington.

DISTRIBUTION

No. of
Copies

No. of
Copies

OFFSITE

Technical Steering Panel

D. S. Barth
University of Nevada
4505 Maryland Parkway
Las Vegas, NV 89154

W. A. Bishop
2503 Wedgewood Court SE
Olympia, WA 98501

M. L. Blazek
Oregon Department of Energy
625 Marion Street N.E.
Salem, OR 97310

G. G. Caldwell
Director
Tulsa City-County Health Dept
4616 East 15th Street
Tulsa, OK 74112

S. N. Davis
Dept. of Hydrology & Water
Resources
Bldg. 11
University of Arizona
Tucson, AZ 85721

N. J. Germond
224 Iron Mountain Blvd.
Lake Oswego, OR 97034

P. C. Klingeman
Civil Engineering Dept.
Oregon State University
Appejin Hall 202
Corvallis, OR 97331-2302

K. J. Kopecky
Fred Hutchinson Cancer
Research Center
1124 Columbia Street
Seattle, WA 98104

P. D. McGavran
Dept. of Health and Welfare
450 W. State Street, 4th Floor
Boise, ID 83720-5450

R. L. Morrill
Dept. of Geography
DP-10
University of Washington
Seattle, WA 98195

A. H. Murphy
Dept. of Atmospheric Sciences
Oregon State University
Corvallis, OR 97331-2209

D. W. Price
Agricultural Economics
Hulbert Hall Room 211
Washington State University
Pullman, WA 99164-6210

M. A. Robkin
Radiological Sciences SB-75
University of Washington
Seattle, WA 98195

G. S. Roessler
202 Nuclear Sciences Center
University of Florida
Gainesville, FL 32611

B. Shleien
2421 Homestead Drive
Silver Springs, MD 20902

No. of
Copies

A. P. Slickpoo, Sr.
P. O. Box 331
809 Nez Perce Lane
Kamiah, ID 83536

J. E. Till
Rt. 2 Box 122
Neeses, SC 29107

D. E. Walker, Jr.
P.O. Box 4147
Boulder, CO 80306

Other

2 DOE Office of Scientific and
Technical Information
Technical Information Center
P.O. Box 62
Oak Ridge, TN 37830

B. G. Brooks, EH-421
Department of Energy
Room J-112
Germantown, MD 20545

18 K. CharLee
Office of Nuclear Waste Mgmt.
Department of Ecology
99 South Sound Center
Mail Stop PV-11
Olympia, WA 98504

6 W. A. Glass
13874 E. 47th Lane
Yuma, AZ 85365

M. Sage
Centers for Disease Control
1600 Clifton Road
Atlanta, GA 30333

K. F. Sparks
Shea & Gardner
1800 Massachusetts Ave., N.W.
Washington, DC 20036

No. of
Copies

J. Thomas
HEAL
1720 N. Ash
Spokane, WA 99205

ONSITE

2 DOE Richland Field Office

R. F. Brich, TSD A5-55
M. W. Tiernan, TSD A5-55

37 Pacific Northwest Laboratory

C. E. Cushing	P7-59
J. M. Daer (5)	K6-89
D. H. Denham	A3-60
R. L. Dirkes	K6-13
W. T. Farris	K3-54
M. D. Freshley	K6-77
R. O. Gilbert	K7-34
R. H. Gray	K1-30
G. L. Harvey	K1-36
C. M. Heeb	K6-42
T. A. Ikenberry	K3-54
A. H. McMakin	K6-86
B. A. Napier	K3-54
J. V. Ramsdell	K6-03
R. E. Rhoads	K6-64
J. C. Simpson	K7-34
D. L. Stewart	K6-91
W. L. Templeton	K1-30
W. H. Walters (5)	K6-09
G. L. Wilfert	K6-54
Public Reading Room (5)	A1-65
Publishing Coordination	K1-06
Records Center	K3-70
Technical Library (2)	P8-55

Priest
Rapids
Dam

Mile 150

Index Map of Part of the Columbia River Basin

Plate 1

Locations of Water and Sediment Sampling by Hanford Contractors

- △ Water Sample Locations**
- Sediment Sample Locations**

State centerline
during high flow

H Island

Locker Island

White Bluffs
Slough

White Bluffs
Towpound

White Bluffs
Boat Ramp

Grant County
Franklin County

Hanford Slough

Hanford Township

Hanford Site
Boundary

Alouette Island

Plate 2

The Hanford Reach - Showing Surveys of Shoreline Radioactivity Exposure Rates and the Effluent Plume Centerline

Surveys of Shoreline Radioactivity Exposure Rates

1953 Shoreline Survey (Paas 1953). D Island only.

- Ⓐ - ⓓ 1959 Shoreline Survey (McConiga and Rising 1959) - Both shorelines surveyed each mile from point Ⓐ to point ⓓ. Alternate shorelines surveyed each mile from point ⓓ to point ⓔ.

○ 1962 Shoreline Survey (McConnon 1962).

■ 1966 Shoreline Survey (Grande 1966).

- Ⓐ - Ⓑ 1966 Shoreline Survey (Lodge 1966). Reactor shoreline surveyed every quarter mile and opposite shoreline surveyed every half mile from point Ⓐ to point Ⓑ. Both shorelines surveyed every half mile from point Ⓑ to point Ⓒ. Islands D, E, and H, and Locke Island surveyed.

▲ 1974 Shoreline Survey (Fix 1975)

- Ⓐ - Ⓕ 1978-79 Shoreline Survey (Sula 1980) Islands and selected recreational shoreline areas surveyed from point Ⓐ to point Ⓕ.

Effluent Plume Centerline

→ Effluent Plume Centerline based on transect data

→ Assumed effluent plume centerline (no transect data available).

Plate 2

The Hanford Reach - Showing Surveys of Shoreline Radioactivity Exposure Rates and the Effluent Plume Centerline

Surveys of Shoreline Radioactivity Exposure Rates

1953 Shoreline Survey (Paas 1953). IJ Island only.

- (A) - (D) 1959 Shoreline Survey (McConiga and Rising 1959). Both shorelines surveyed each mile from point (A) to point (D). Alternate shorelines surveyed each mile from point (D) to point (E).

○ 1962 Shoreline Survey (McConnon 1962).

■ 1966 Shoreline Survey (Grande 1966).

- (A) - (B) 1966 Shoreline Survey (Lodge 1966). Reactor shoreline surveyed every quarter mile and opposite shoreline surveyed every half mile from point (A) to point (B). Both shorelines surveyed every half mile from point (B) to point (C). Islands D, E, and H, and Locke Island surveyed.

▲ 1974 Shoreline Survey (Fix 1975).

- (A) - (F) 1978-79 Shoreline Survey (Sula 1980). Islands and selected recreational shoreline areas surveyed from point (A) to point (F).

Effluent Plume Centerline

————→ Effluent Plume Centerline based on transect data

————→ Assumed effluent plume centerline (no transect data available).

END

**DATE
FILMED**

6 / 9 / 92