Spectroscopic Probes of Vibrationally Excited Molecules
at
at Chemically Significant Energies

DOE/ER/13965--4

DE92 011026

Progress Report
for Period August 15, 1991 - August 14, 1992

Thomas R. Rizzo

University of Rochester
Rochester, New York 14627

March 1992

Prepared for

The U. S. Department of Energy
under grant No. DE-FG02-88ER13965

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTERDISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *se*

Table of Contents

Abstract	3
Project Overview	4
Double Resonance Spectroscopy of Dissociating Molecules	4
Vibrational Overtone Spectroscopy of Jet-Cooled Molecules.....	7
References	8

Abstract

These experiments apply multiple-laser spectroscopic techniques to investigate the bond energies, potential surface topologies, and dissociation dynamics of highly vibrationally excited molecules. Infrared-optical double resonance pumping of light atom stretch vibrations in H_2O_2 and HN_3 prepares reactant molecules in single rovibrational states above the unimolecular dissociation threshold on the ground potential surface, and laser induced fluorescence detection of the OH or NH fragments monitors the partitioning of energy into individual product quantum states. Product energy partitioning data from H_2O_2 dissociation provide a stringent test of statistical theories as well as potential energy surface calculations. Ongoing work on HN_3 seeks to determine the height of the barrier to dissociation on the singlet potential energy surface.

Our most recently developed spectroscopic scheme allows the measurement of high vibrational overtone spectra of jet-cooled molecules. This approach uses CO_2 laser infrared multiphoton dissociation followed by laser induced fluorescence product detection to measure weak vibrational overtone transitions in low pressure environments. Application of this scheme to record the $\Delta v_{\text{OH}}=4$ and $\Delta v_{\text{OH}}=5$ transitions of CH_3OH cooled in a supersonic free-jet demonstrates both its feasibility and its utility for simplifying high vibrational overtone spectra.

Project Overview

This project involves the development and application of new spectroscopic techniques for investigating small polyatomic molecules with chemically significant amounts of vibrational energy. Traditional spectroscopic techniques provide little information about the high energy regions of molecular potential energy surfaces that control chemical bond breaking. Highly excited species produced by exothermic reactions in combustion processes sample these regions of the potential, and a better understanding of their surface topology and bond dissociation energies will allow more accurate models of their dissociation dynamics.

In the last budget period we have continued (and completed) infrared-optical and optical-infrared double resonance studies of hydrogen peroxide (H_2O_2)¹⁻³ and have begun similar studies of the spectroscopy and unimolecular dissociation dynamics of hydrazoic acid (HN_3).⁴ In addition, we have developed a new experimental scheme which allows the measurement of light atom stretch overtone transitions of molecules cooled in supersonic molecular beams. Our first application of this new approach has been to measure the $\Delta v_{\text{OH}}=4$ and $\Delta v_{\text{OH}}=5$ transitions of jet-cooled methanol.⁵

Double Resonance Spectroscopy of H_2O_2 and HN_3

We have now completed (for the time being) our analysis of the spectroscopy and dissociation dynamics of H_2O_2 . Using infrared-optical double resonance, we first excite H_2O_2 molecules *via* an OH stretch fundamental or combination band with an optical parametric oscillator and then promote the initially excited molecules to a high vibrational overtone level with a pulsed dye laser. This technique prepares reactant molecules in single rovibrational states at energies above their unimolecular dissociation threshold on the ground potential energy surface. We monitor the absorption of the second photon, and hence the

dissociation process, *via* laser induced fluorescence of the OH fragments with a third tuneable laser. Two types of information result from these experiments. In one case we scan the frequency of the vibrational overtone excitation laser to generate a simplified overtone excitation spectrum of the parent reactant molecule. In the other case, we fix the frequency of the infrared and overtone excitation lasers to produce reactants with well specified vibrational and rotational quantum numbers and then scan the frequency of the laser that probes the OH fragments. The product excitation spectrum so generated allows us to determine how the excess energy is distributed among the internal degrees of freedom of the products.

We have in the last year completed our analysis of the rotationally resolved overtone transitions to states above the unimolecular dissociation threshold.^{2, 3} The dependence of the spectra on the rotational quantum numbers J and K help us to determine the nature of the couplings that cause redistribution of the initial excitation energy prior to dissociation. Highly resolved overtone transitions to predissociative levels also indicate that near threshold, the relative timescales of IVR and unimolecular dissociation do not differ as much as statistical theories of unimolecular dissociation assume. We have also completed analysis of the product energy partitioning data from 36 different reactant rovibrational states ranging in energy from 3–2000 cm^{-1} above the unimolecular dissociation threshold.¹ These results indicate that the predictions of Phase Space Theory are in reasonable agreement with our measured product distributions, however clear deviations from statistical behavior exist. This result indicates that for molecules as small as H_2O_2 , the density of states is sufficiently small that statistical theories may not apply as well as to larger systems. The state-specific data that we have obtained provides a stringent test of calculated potential energy surfaces of H_2O_2 as well as dynamical calculations carried out on such surfaces.

We have recently begun to apply our infrared-optical double resonance technique to investigate the unimolecular dissociation dynamics of hydrazoic acid (HN_3). The dissociation dynamics of this molecule

are expected to be fundamentally different from that of H_2O_2 insofar as a curve crossing between the ground state and a low lying dissociative excited state can lead to dissociation via a spin forbidden channel to produce $\text{NH} (^3\Sigma) + \text{N}_2$. At sufficiently high energies on the ground potential surface, there is a competition between dissociation on the spin allowed singlet surface and spin forbidden triplet surface which is sharply dependent upon the reactant energy. Moreover, the topology of the exit channel on the singlet surface is not well characterized.⁶ There appears to be a barrier along the coordinate to produce $\text{NH} (^1\Delta) + \text{N}_2$, however estimates of this barrier range from 450 to 1740 cm^{-1} .⁶

We have begun to perform both infrared-optical and optical-infrared double resonance studies of this molecule to investigate its dissociation dynamics on both the singlet and triplet surface. To date we have obtained simplified vibrational overtone spectra in both the $5\nu_{\text{NH}}$ and $6\nu_{\text{NH}}$ vibrational overtone levels and are in the process of assigning these spectra. We are also attempting to access combination bands containing $6\nu_{\text{NH}}$ and one quantum of low frequency vibration which will put us right at the energy of the barrier on the singlet surface in an effort to help us determine the height of the barrier. Moreover, by observing the rotational state dependence of the overtone spectra near the of the barrier, we can actually estimate the rotational constants of the molecule at the transition state.

We have also done some preliminary work on HOCl . In addition to its important in the chemistry of the atmosphere, this molecule should provide an important meeting place for experiment and theory. By virtue of its small size it should lend itself well to both potential surface calculations and dynamical simulations.

Vibrational Overtone Spectroscopy of Jet-Cooled Molecules

In order to extract dynamical information from high vibrational overtone spectra, one must eliminate

the severe spectral congestion which is inherent to room temperature samples. In our studies of H_2O_2 and HN_3 described above, the double resonance nature of the excitation process completely eliminates spectral congestion and allows the preparation of reactant molecules in single rovibrational states. However, because vibrational overtone excitation promotes such a small number of molecules to the highly excited states, one must detect this absorption with extreme sensitivity. For H_2O_2 and HN_3 , we achieve such sensitivity by detecting the OH or NH product fragments via laser induced fluorescence. This limits us to investigating the spectroscopy of vibrational levels which lie above the unimolecular dissociation threshold and hence we must choose molecules for study which have particularly low dissociation energies. Although photoacoustic spectroscopy has the sensitivity to detect weak vibrational overtone transitions, the high pressures required for this approach are incompatible with spectral simplification techniques such as infrared-optical double resonance or cooling in a supersonic expansion

We have recently developed a technique which allows the detection of vibrational overtone transitions to non-dissociative levels in low pressure environments where spectral simplification methods are applicable. This approach uses infrared multiphoton dissociation (IRMPD) of the vibrationally excited molecules produced after overtone excitation. Spectroscopic detection (of the overtone absorption) is achieved by monitoring dissociation products *via* LIF. This method relies on our ability to selectively dissociate vibrationally excited molecules *via* IRMPD without dissociating the large excess of ground state molecules. We have demonstrated such selectivity in preliminary experiments on CH_3OH cooled in a supersonic free jet. Our initial results show that the $\Delta\nu_{\text{OH}}=4$ and $\Delta\nu_{\text{OH}}=5$ vibrational overtone transitions, which have widths of $100\text{-}150\text{ cm}^{-1}$ when recorded by room temperature photoacoustic spectroscopy, are broadened primarily by thermal inhomogeneous spectral congestion. Individual features in the $\Delta\nu_{\text{OH}}=5$ spectrum show linewidths of $< 1\text{ cm}^{-1}$.

While or initial applications of this approach have been to measure direct overtone spectra in a supersonic free jet, it should also lend itself to detecting infrared-optical double resonance overtone excitation as well as stimulated emission pumping of high vibrational states.

References

1. X. Luo and T. R. Rizzo, J. Chem. Phys. **96**, (in press) (1992).
2. X. Luo and T. R. Rizzo, J. Chem. Phys. **96**, (in press) (1992).
3. P. R. Fleming, X. Luo, and T. R. Rizzo, in *Mode Selective Chemistry*, edited by B. Pullman and J. Jortner (Kluwer, Dordrecht, 1991).
4. T. A. Seckel, X. Luo, and T. R. Rizzo, (work in progress) .
5. R. D. F. Settle and T. R. Rizzo, J. Chem. Phys. (submitted) (1992).
6. M. H. Alexander, H. Werner, T. Hemmer, and P. J. Knowles, J. Chem. Phys. **93**, 3307 (1990).

END

**DATE
FILMED
5/18/92**