DOE/ER/40562--T1

Technical Progress Report

DOE/ER/40562--T1

DE93 009945

on

Department of Energy Division on Nuclear Physics

Grant

DE-FG02-90ER40562

Measurements of Rare Composite Objects and High Sensitivity Searches for
Novel Forms of Matter Produced in High Energy Heavy Ion Collisions

For the Period

April 1992 - March 1993

Yale University
Physics Department
554 J.W. Gibbs Laboratory
260 Whitney Avenue
New Haven, CT 06511

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

1 Overview

This document outlines the development of our group's research in the field of high energy heavy ion collisions.

Our group, a subset of the Yale experimental high energy physics group (called YAUG), initially became interested in high energy heavy ions in 1988. At that time, we joined E814 at Brookhaven's AGS in order to search for a novel form of matter called *strange quark matter*. Since the start of our involvement in E814, our group's experience and interest in heavy ions has grown, and we now are building a new experiment at the AGS, E864, in order to pursue a wide range of physics topics.

Our physics program consists of two complementary parts. First, we are searching for new particles. These include the aforementioned strange quark matter, as well as strange chiral solitons and multistrange hypernuclei. These and other new phenomena may result from the extreme conditions present in relativistic heavy ion collisions, which is a regime that is still relatively new and untested. Second, we plan to perform "bread and butter" measurements of known particles, such as the coalescence production of nuclei and antinuclei. Knowledge of coalescence production is not only crucial to interpreting the limits in strange matter searches, but also is useful in probing the relativistic heavy ion collision process.

While E814's data analysis is still proceeding, no additional experimental runs will take place in the E814 apparatus and our involvement in this experiment is coming to a close.

Our efforts will primarily be the construction, running, and analysis of E864. E864's program was approved by BNL in 1991, and its funding was granted last year by the DOE. E864 is just getting under way, and it will involve the efforts of our group and our collaborators' beyond the last running period which is currently planned to take place in FY1997.

We also plan to continue our heavy ion program at BNL's RHIC accelerator. We envision this to be a "fixed-target" experiment using a gas jet or pellet target that would extend our studies of new particles in the high energy regime offered by this new accelerator.

2 E814

E814 was designed to study both global event characteristics and forward fragments resulting from heavy ion collisions. The detector combined 4π calorimetry around the target with a high resolution spectrometer for particles which pass through a 1° cone about the beam axis. The spectrometer arm was used in the single-particle searches and measurements discussed below.

2.1 Motivation

Chin and Kerman first made the striking suggestion that multiquark ($n_{quarks} \gg 3$) states with comparable numbers of u , d , and s quarks would have metastable lifetimes as long as 10^{-4} sec.¹ Witten, in 1984, renewed interest in this topic when he argued that strange matter systems may actually be more stable than nuclear matter.² About the same time, Farhi and Jaffe performed a detailed study of the phenomenology of strange matter.³ Using the MIT Bag Model, Farhi and Jaffe studied both extended strange matter systems as well as small droplets which they called *strangelets*.

The essential reason for the stability of strange matter states comes from the Pauli Exclusion principle. This principle asserts that an extended three-flavor state may be more stable than a comparably-sized state populated with just two quark flavors. In the approximation that quark matter is a Fermi gas enclosed in a bag, and for simplicity that the quarks are massless, the total energy of the system is proportional to

$$E \propto N_f^{-\frac{1}{4}},$$

where N_f is the number of quark flavors in the system. Increasing N_f from 2 to 3 gives an energy savings of $1 - (\frac{2}{3})^{\frac{1}{4}} \simeq 10\%$, or roughly 100 MeV/baryon. Though a massive strange quark would give less binding, this large energy savings may compensate for the strange quark mass. Thus the exclusion principle, when combined with the system's small coulomb repulsion (as well as other, dynamical reasons), makes strange matter plausible.

¹S.A. Chin and A.K. Kerman, Phys. Rev. Lett. **43**, 1292 (1979).

²E. Witten, Phys. Rev. D**30**, 272 (1984).

³E. Farhi and R.L. Jaffe, Phys. Rev. D**30**, 2379 (1984).

Some results of the study by Farhi and Jaffe are shown in Fig. 1. The minimum E/A (energy per baryon) is plotted versus A in the top curve; the bottom shows S (the number of strange quarks) vs. A . The conclusion of these curves is that strangelets with $A \gtrsim 8$ may be metastable, and the charge-to-mass ratio which yields the most stability is in the range $0.1 \lesssim Z/A \lesssim 0.3$.

As progress was made in E814, we became more interested in studying antiproton production, as well as the production of light nuclei in relativistic heavy ion collisions. Antiproton production will begin to provide a handle on the characteristics of heavy ion collisions. The production of light nuclei will give an important first look at coalescence in heavy ion collisions at the AGS. While coalescence has been demonstrated at BEVALAC energies, it is still unproven in the AGS energy regime. These production topics are not only interesting in their own right: they will provide the necessary tools to interpret null results from rare particle searches.

2.2 Strangelet Searches in E814

Heavy ion collisions at the AGS create regions with a high strangeness content and baryon density, and thus may be the best environment in which to search for strange matter. There are two classes of strangelet production models in heavy ion collisions. The first are quark-gluon plasma (QGP) based models, where the plasma preferentially emits antistrange quarks in the form of mesons, resulting in a strangelet. The second are coalescence models, where hyperons and nucleons are made in the interaction, and then the baryons near each other in $(\vec{x}, \vec{p})$ space bind to form a strangelet. Both models would predict strangelet production peaked at the center of mass, but they give very different results for cross sections. A plasma-based prediction⁴ estimates that the probability of a strangelet being produced in a collision is between 1 and 10^{-6} per quark-gluon plasma produced. Though no formal coalescence calculation on strangelet production has been performed, it is expected that the production will be roughly 10^{-12} per collision for ^{28}Si beams and nuclear targets.

Our strangelet searches in E814 are now concluded. The first run's upper limits were documented in the literature.⁵ Analysis of data from a second

⁴H.C. Liu and G.L. Shaw, Phys. Rev. D30, 1137 (1984).

⁵J. Barrette *et al.*, Phys. Lett. B252, 550 (1990).

Figure 1: Results from the study by Farhi and Jaffe. See text for details.

run, which included an improved trigger, has also been completed.⁶ The upper limits set by E814 are at the level $\sigma_S/\sigma_{int} < 10^{-4} - 10^{-5}$, where σ_S and σ_{int} are the strangelet and total cross sections, respectively. This range of upper limits is relevant to a strangelets of typical parameters, and assumes a reasonable production model. These limits address some parameters in the QGP production model, but are well below the sensitivity needed to address the coalescence models.

2.3 Antimatter and Light Nucleus Production in E814

Our involvement in E814 is now primarily in the analysis of antiproton and light nucleus production data.

We have completed the antiproton analysis in E814. The results have been the subject of a thesis⁷, as well as submitted for publication⁸. The analysis measured the antiproton cross section as a function of multiplicity in Si collisions with Al, Cu, and Pb targets.

Data concerning the coalescence production of light nuclei is also presently being analyzed. Figure 2 shows some preliminary results. As can be seen from the figure, the analysis will measure deuteron, triton, and alpha particle production at $p_t = 0$ for different values of event multiplicity. An estimate of the coalescence radius and also a measure of the freezeout volume (in the thermodynamic model) will be presented.⁹ The results will be sent for publication as well.

3 E864

3.1 Introduction

Many of the goals in E864 overlap with E814's, however E864 is specifically designed for high sensitivity, single-particle searches, and is 5 or 6 orders

⁶F.S. Rotondo, *Limits on Strange Quark Matter Production*, Yale University thesis, 1991.

⁷*Production of Antiprotons in Relativistic Heavy Ion Collisions*, S.V. Greene, Yale University thesis

⁸J. Barrette *et al.*, *Antiproton Production in Relativistic Si-Nucleus Collisions*, submitted to Physical Review Letters.

⁹J.V. Germani, Yale University thesis, in preparation.

Figure 2: Differential cross sections of coalesced nuclei. Preliminary E814 results.

of magnitude more sensitive than E814. This high sensitivity and broad acceptance in E864 allow for the search of many exotic states which include but are not restricted to strange matter. Such states, such as strange chiral solitons¹⁰ and bare color, may be more readily produced in relativistic heavy ion collisions than in simpler $p+p$ collisions. The high sensitivity is designed to address the coalescence level of production of the exotic particle searches.

3.2 The E864 Detector

The detector layout is shown in Fig. 3. M1 and M2 are magnets. H1, H2, and H3 are scintillation counter hodoscopes. S1, S2, and S3 are straw tube stations. The calorimeter is a tower geometry, "spaghetti" design.

The straw tubes will measure the rigidity of a track. Each of the hodoscopes will measure the charge, time-of-flight, and vertical position of a particle. The calorimeter will measure the shower energy and its time-of-flight. Thus, individual measurements will have great redundancy: time-of-flight will be measured 4 times (3 times by the hodoscopes, and once by the calorimeter), charge will be measured 3 times in the hodoscopes, and particles will make spatial hits in each of the straw stations and hodoscope planes. Furthermore, the charged particle masses will be measured twice, once using momentum and time-of-flight, and once using cluster energy and time-of-flight.

The great redundancy designed into E864 provides the necessary background rejection to perform meaningful particle searches.

3.3 Yale's Responsibilities in E864

Our group's responsibilities include:

Overall experiment management: It is natural for the Yale group (in which both cospokesmen reside) to perform experiment management tasks. These include both internal management as well as accepting managerial responsibilities relating outside agencies.

Test beam coordination: The detector groups in E864 are also planning to participate in a test beam at BNL this year. Our group is responsible for setting up the running program for our systems, as well as

¹⁰V.B. Kopeliovich *et al.*, Phys. Lett. B242, 145 (1990).

E864

Detector	Fiducial Region	Number of Elements
S1	103.78h x 25.2v	6 planes, 253 tubes each (4mm tubes)
S2	189.51h x 40.0v	6 planes, 462 tubes each (4mm tubes)
H1	224.07h x 48.0v	205 slats, 1.093 cm wide
H2	314.27h x 64.0v	205 slats, 1.533 cm wide
S3	415.53h x 80.0v	6 planes, 1013 tubes each (4mm tubes)
H3	451.41h x 84.0v	205 slats, 2.202 cm wide
CAL	560.0h x 110.0v	754 modules (10cm x 10 cm x 117cm deep)

Figure 3: The E864 detector layout.

coordinating our effort with BNL and other test-beam users who will share the test beam line.

Beam design: The E864 beam line is presently being designed by the AGS liaison physicist D.Beavis in close interaction with our group. Indeed, we are checking all transport runs. This design will take into account the physical restrictions of our position at the AGS. Thus the beam will be made as short as possible so as not to interfere with access doors at the AGS building, as well as steer clear of regions where extended access is needed. This work is proceeding in close conjunction with BNL liaisons.

Data acquisition (DA) system: Our group is also beginning to design a DA system which can absorb the high data flow produced by E864. Using the DA system of E791 as a model (this system actually is more than what E864 needs), we plan to use the same techniques of parallel readout and event building, as well as concurrently writing to multiple tape drives.

4 Future Plans

4.1 E814

During the next year, our effort in E814 will continue with the analysis of light nucleus production data collected in February 1991. At the completion of this analysis, a paper outlining results will be published and a thesis will be written. It is expected that these tasks will be finished in 1993.

4.2 E864

The design decisions for detectors are now largely completed, and the construction of prototype detectors has begun. A test beam run is planned at BNL in 1993, with the precise time yet to be determined by time constraints at the AGS. A detailed construction plan for the next 3 years is provided in the document *E864: Management Plan*, which will arrive at the DOE shortly.

The E864 beam design will be completed in 1993.

The actual DA system configuration will be determined in 1993, and components will be procured and assembled in early 1994 for a first data run which will occur in the fall of 1994.

4.3 Rare Particle Searches at RHIC

We plan to move the E864 apparatus, with some modification, in a gas jet fixed-target configuration at RHIC. There are some advantages of a RHIC setup over a comparable AGS setup. The "focussing" due to the Lorentz boost of the higher energy RHIC beam would allow a modest-sized experiment to have a rather large acceptance. In particular, the E864 detectors can be spread out longitudinally and form a system with interesting acceptance and resolution for RHIC (fixed-target) energies. Further, the RHIC beams will have a 100% macro duty cycle (the AGS has a 20% duty cycle). The variable-energy RHIC beam could also be used to study energy-dependent production effects. In some models the higher energy available in the RHIC fixed-target mode is predicted to yield enhanced production rates. The RHIC beam would allow for interaction rates up to about 10^8 Hz without adversely affecting other experiments.

We plan to submit a BNL Letter of Intent for this activity in the next year.

END

**DATE
FILMED**

5 / 10 / 93