NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

DOE/ER/45118--7

DE92 010032

CONDENSED MATTER PHYSICS
AT
SURFACES AND INTERFACES OF SOLIDS

Progress Report
For the period February 1, 1991 - January 31, 1992

Eugene Mele

University of Pennsylvania
Philadelphia, Pennsylvania 19104

January 1992

Prepared for

THE U. S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-84ER45118

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Progress Report:
Condensed Matter Physics at Surfaces and Interfaces of Solids

E.J. Mele
Department of Physics
University of Pennsylvania
Philadelphia, PA 19104

Prepared: November 1991

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. Introduction

This research program is focused on structural and elastic properties of crystalline solids and interfaces between solids. We are particularly interested in novel forms of structural ordering and the effects of this ordering on the lattice dynamical properties. We are currently studying structural and vibrational properties of the surfaces of the elemental alkaline earths (particularly Be), and structural phenomena in the doped fullerenes.

II. Summary of Work in Progress:

1. Surface Phonon Spectra for Be

Elemental Be crystallizes in the hcp crystal structure. Among the hcp metals it is unique in that it exhibits the smallest c/a ratio. Recent diffraction studies have revealed that for the clean (0001) surface (normal to the hexagonal planes) there is an exceptionally large interlayer relaxation in which the first interlayer distance expands relative to its bulk value. This is unusual for close packed elemental surfaces which typically exhibit a contraction of the first interlayer spacing. Finally, other low index high symmetry surfaces of Be are characterized by more exotic longer range reconstruction patterns, which have been incompletely explored to date.

We have begun a lattice dynamical study of the bulk and surface vibrational properties of Be. The goal of the work is to interpret some of the

structural anomalies noted above. More generally, we believe that these phenomena may be related to the non free electron character of the conduction states in Be. Be is a semimetal, with a strong pseudogap appearing in the electronic spectrum near the Fermi energy, and this is likely to have significant consequences for the low frequency nonlocal electronic polarizability which is relevant for lattice dynamics.

As a preliminary stage in this study we have completed an investigation of the bulk lattice dynamical properties of Be, and have extended the model to the (0001) surface using a Green's function technique to study the vibrational spectra. The bulk model is a harmonic model including central and noncentral interactions which are directly fit to the bulk phonon spectra of Be (measured by inelastic neutron scattering some time ago). The model indeed reveals the importance of "extended" interatomic interactions. In fact a fit to the bulk spectra requires interactions over four coordination shells parallel to the basal (hexagonal) planes, and third neighbor in the stacking direction. This is a significantly more extended force field than is required for the alkali metals for example, and is even more extended than the interaction model for Mg which is the next heavier alkaline earth in the hcp structure.

We have applied this model to the (0001) surface, assuming an ideal termination of the crystal with no relaxation of the surface interaction parameters. Representative results for the surface and bulk phonon spectra along the Γ -M symmetry line are shown below. These reveal that even within this simplified model, the Rayleigh vibration deviates substantially from the classical prediction (derived from a continuum elastic theory) and is anomalously soft at the zone boundary. Interestingly this softening is not

obtained for the dispersion along the Γ -K symmetry direction. Experiments on a different termination of the crystal $(\bar{2} \ 1 \ 0 \ 0)$ surface have in fact revealed an anomalous Rayleigh wave with this character, but the details of the mechanism leading to this phenomenon both for the (0001) surface and the $(\bar{2} \ 1 \ 0 \ 0)$ surface are still being studied. Continuation of this work is described in Section III of the proposal.

2. Alkali Ordering In Doped Fullerites

Shortly after the discovery of condensed solid phases of the fullerites (C_{60}) it was discovered that these compounds could be chemically doped to conducting, and superconducting states by the incorporation of alkali metals. The situation is loosely analogous to the well studied intercalation chemistry of the graphites. The alkali doped phases of these compounds have been studied most extensively to date. The experiments show, that in addition to interesting transport properties, there are an interesting and rich array of structural phenomena associated with the incorporation of alkali metals into the fullerite host. The experimental situation is evolving rapidly, but to summarize the currently available data, it is known that saturation doping with the heavy alkali metals, leads to a transition of the undoped fcc phase to a saturated bcc phase with stoichiometry M_6C_{60} . At intermediate concentrations there are stable phases at $x = 3$ in the fcc phase (this is the superconducting phase), and a metallic (but nonsuperconducting) compound at $x=4$ in the body centered tetragonal structure. There are indications of a stable structure at a composition $x=1$ which is thought to exhibit the rocksalt structure.

We are developing a theory for structural ordering in the alkali doped fullerites. The theory is similar in some respects to the theory for stage ordering in the alkali intercalated graphites. We are completing two important but preliminary stages of this study (results will be prepared for publication soon).

The first of these applies a simple Ising lattice gas model to describe the alkali ordering in the fullerite host. This is appropriate for the doped phases with $x \leq 3$ in which the C₆₀ occupy lattice sites of the fcc Bravais lattice. The available alkali sites are the high symmetry interstitials, namely the octahedral and tetrahedral interstitial sites in the primitive cell. Of course the interactions among the alkali ions on these sites is primarily repulsive, and is described by a quadratic interactions coupling Ising spin variables on each of the available sites, representing the occupation number. The important physical point is that the Ising model on this interstitial lattice is actually quite nontrivial: as the chemical potential for the alkali is varied there is a competition between the tendency to occupy the octahedral sites (which has the lowest self energy) and the tetrahedral sites for which the repulsive energy is minimized. The model can potentially exhibit an array of modulated equilibrium structures as temperature and chemical potential are varied; we have so far been exploring only the $q=0$ structures, which already exhibit 2 first order transitions, and the possibility of a broken symmetry phase in which the occupation of the tetrahedral sites is unequal. A model phase diagram in the T - x plane is given below which exhibits miscibility gaps for $3 > x > 1$ and $1 > x > 0$. We refining these data, and extending the scheme to study the possibility of equilibrium phases for $q \neq 0$.

Figure 1: Phase diagram for the Ising lattice gas model in the T - x plane. Shaded regions are equilibrium thermodynamic phases at the given values of T and x , with the open regions at low temperature representing miscibility gaps induced by first order phase transitions in the model. At very low temperature there are three distinct phases corresponding to the empty lattice, saturation of the octahedral interstitial site ($x=1$) and saturation of the octahedral and tetrahedral interstitial sites ($x=3$). The range of equilibrium structures represented at high temperatures is limited only by the range of chemical potentials included in the scan.

A more striking aspect of the structural problem in the fullerites is the structural transition in the host induced by alkali doping. As noted above the saturation phase is in the bcc crystal structure, and there are intermediate, stable, non cubic compounds. We are developing a description of this phenomenon within the framework of a Landau theory. Briefly, all of the structures observed to date can be described by subjecting the parent fcc structure to a homogenous (relatively large amplitude) strain. Our observation is that these phenomena might therefore be describable in the framework of a macroscopic nonlinear elastic theory. In fact it is known that the fcc-bcc martensitic transition can be well described with a Landau expansion of the elastic energy to quartic order in the strains. Our program is to study the renormalization of the coefficients in the Landau expansion due to alkali doping, and then to calculate the equilibrium structure at fixed temperature and alkali chemical potential. The results of one such calculation are given below. This preliminary phase diagram indicates a strongly first order transition between a low density fcc structure and a high density bcc structure, with an intermediate cubic phase in the A-15 structure. The sensitivity of the predicted diagram to parameters in the potential, the effect of external symmetry breaking fields (anisotropic stress, etc.) are important aspects of the theory which remain to be developed.

III. Proposed Work:

1. Bulk and Surface Lattice Dynamics of Alkaline Earths Metals

We have written and tested a general computer code which applies a Green's function method to extract the surface lattice dynamical properties for a relatively complex harmonic interaction model on a general lattice. [This work is largely the contribution of R.L. Sameth who is now completing a M.S. with my group.] Our immediate goals are to apply this to the interaction model which we have derived for bulk Be. Of interest are the dispersions of the low frequency Rayleigh branches, and other localized surface modes at high frequency in the surface projected Brillouin zone. We hope to couple this activity with the experimental work of Plummer's group in our Department who is tuning up a high performance high resolution inelastic electron scattering spectrometer to study surface vibrational phenomena. Be (0001) will likely be the first sample studied in this system. It is possible to develop a simple phenomenological model of the low frequency $S(q,\omega)$ in this system by positing a dependence of the "dynamic charges" to interatomic displacement; in the absence of more complete calculations of the electronic response functions, we expect that this will be a valuable tool for interpreting the q dependence of the observed EELS spectra.

A longer range goal of this program is to couple this phenomenology to a more complete theory of the electronic properties of the semimetallic surface. We have had some success carrying out such a theory, within a tight

binding formalism for semiconductor surfaces. Similar models have been developed for lattice dynamics for transition metals and related systems. This type of extension appears to be particularly relevant for Be whose bulk phonon spectrum is already quite complex, and might be more readily analyzed with an explicit calculation of the nonlocal interactions mediated by the conduction electrons in this system. This remains a longer range goal of this research problem.

2. Structural Phenomena in the Doped Fullerites.

We are currently refining the predictions of the Ising lattice gas model for the equilibrium phases of the alkali doped fullerites. This model, though quite simple and compact, is already yielding an interesting phase diagram for the doped system, even restricted to the subspace of $q=0$ (homogeneous) intercalation patterns. Furthermore this model is very likely to exhibit equilibrium superlattice structures with $q \neq 0$ when we generalize this study. Such phases have not yet been observed experimentally, although we note that the available experimental cuts across this phase diagram are far from complete, particularly at low temperature where these modulated structures are most likely to appear.

We are planning a continuation of our studies using the Landau theory to describe the doping induced structural transition in this system. This is an ambitious program, of which several parts are still being developed. We are currently attempting to fit the various coefficients which appear in the Landau expansion of the free energy to the results of molecular dynamics simulations,

which use classical interatomic potentials to study the equilibrium structures of the doped fullerites at finite temperatures. This fitting procedure has so far met with mixed success; although we are hopeful of extracting a tractable model for the Free energy which will allow us to explore the cubic and noncubic doped phases of this system.

IV. Students

There are three students who have participated in various aspects of this program, one of whom is directly supported through this DOE grant. The students are R.L. Sameth who has been working on the surface phonon problem, and who is completing his M.S. here, M. Deshpande (DOE supported), a second year Ph.D. student who is carrying out the Ising lattice gas calculations, and S. Hong (currently supported on a TA in the Department of Physics), also a Ph.D. student who is helping to develop the Landau theory of the doping induced structural transformations in fullerites.

V. Publications and Presentations

1. "Thermal Vibrational Amplitudes on Si (111) and Si (001), Physical Review B 40, xxx (1990) E.J. Mele with O.L. Alerhand and J.D. Joannopoulos
2. "Staging in Doped Fullerites" Workshop on Fullerites, University of Pennsylvania, August 1991., E.J. Mele and S. Hong

Our work on the Ising lattice gas theory for doped fullerites is in preparation for publication.

VI. Current and Pending Support

NSF DMR 90 08256: Structure and Excitations of Reduced Dimensional Systems
\$45,000 12/01/90 to 11/30/91

NSF DMR 88 19885: Elementary Excitations in Condensed Matter Systems
(NSF-MRL) \$40,141 3/1/91 to 2/28/92

DOE DE FR02 84 ER45118: Condensed Matter Physics at Surfaces and Interfaces
of Solids
(this proposal) \$138,000 2/1/91 to 1/31/94

NSF: Physics Of Condensed Phases in New Advanced Materials:
(submitted, November 1991, with A.B. Harris and C.L. Kane)

END

**DATE
FILMED**

4 / 21 / 92