BNL-NCS--17541

DE92 010079

ENDF-201
ENDF/B-VI SUMMARY DOCUMENTATION

Compiled and Edited by
P.F. Rose
Brookhaven National Laboratory

October 1991

MASTER

NATIONAL NUCLEAR DATA CENTER

BROOKHAVEN NATIONAL LABORATORY
ASSOCIATED UNIVERSITIES, INC.
UPTON, LONG ISLAND, NEW YORK 11973

UNDER CONTRACT NO. DE-AC02-76CH00016 WITH THE
UNITED STATES DEPARTMENT OF ENERGY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency, contractor or subcontractor thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor or subcontractor thereof.

Printed in the United States of America
Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

NTIS price codes:
Printed Copy: A21; Microfiche Copy: A01

Table of Contents

Contents of ENDF/B-VI	PAGE
Contributors to Documentation	v
Introduction	xvi
Appendix (Corrections to Release 1)	1
	471

Evaluation Summaries

Isotope	Material Number	
Hydrogen-1	125	7
Hydrogen-2	128	13
Helium-3	225	15
Lithium-6	325	18
Lithium-7	328	33
Beryllium-9	425	41
Boron-10	525	48
Boron-11	528	69
Carbon-Natural	600	78
Nitrogen-14	725	82
Nitrogen-15	728	91
Oxygen-16	825	101
Fluorine-19	925	120
Vanadium-Natural	2300	124
Chromium-Isotopes (50,52,53,54)	2425,2431,2434,2437,2437	131
Manganese-55	2525	144
Iron-Isotopes (54,56,57,58)	2625,2631,2634,2637	149
Iron-56 (High energy, n + p sub-libraries)	2631	168
Cobalt-59	2725	174
Nickel-Isotopes (58,60,61,62,64)	2825,2831,2834,2837,2843	181
Nickel-59	2828	197
Copper-Isotopes (63,65)	2925,2931	198
Yttrium-89	3925	211
Niobium-Natural	4125	216
Palladium-105	4634	222
Palladium-107	4640	225

Evaluation Summaries (continued)

Isotope	Material Number	PAGE
Indium-Natural	4900	228
Indium-115	4931	234
Cesium-134	5528	238
Barium-134	5637	241
Barium-135	5640	245
Barium-136	5643	249
Barium-137	5646	253
Neodymium-147	6040	256
Promethium-147	6149	260
Samarium-147	6234	263
Samarium-151	6246	268
Europium-151	6325	271
Europium-152	6328	278
Europium-153	6331	285
Europium-154	6334	291
Europium-155	6337	298
Holmium-165	6727	302
Erbium-166	6837	309
Erbium-167	6840	313
Rhenium-185	7525	317
Rhenium-187	7531	321
Gold-197	7925	325
Lead-Isotopes (206,207,208)	8231,8234,8237	336
Bismuth-209	8325	351
Uranium-235	9228	356
Uranium-236	9231	376
Uranium-238	9237	386
Neptunium-237	9346	397
Neptunium-239	9352	411
Plutonium-239	9437	415
Plutonium-240	9440	434
Plutonium-241	9443	446
Americium-241	9543	453
Americium-243	9549	457
Berkelium-249	9752	465
Californium-249	9852	468

Contents of ENDF/B-VI

The contents of ENDF/B-VI are described in the following tables. All new or extensively modified evaluations as well as older evaluations converted to ENDF-6 format are included. The list includes the contents of the incident neutron, proton, and deuteron sublibraries, the thermal scattering law sublibrary, and two high energy evaluations for ^{56}Fe . The contents of the photon interaction sublibrary, decay data sublibrary, and the fission product yield sublibrary are not included.

Neutron Sublibrary

MAT No. Material Usage

0125	H-1	Standard, Neutron transport, Gamma production
0128	H-2	Neutron transport, Gamma production
0131	H-3	Neutron transport
0225	He-3	Standard, Neutron transport
0228	He-4	Neutron transport
0325	Li-6	Standard, Neutron transport, Gamma production
0328	Li-7	Neutron transport, Gamma production, Covariances
0425	Be-9	Neutron transport, Gamma production
0525	B-10	Standard, Neutron transport, Gamma production
0528	B-11	Neutron transport, Gamma production
0600	C	Standard, Neutron transport, Gamma production, Covariances
0725	N-14	Neutron transport, Gamma production
0728	N-15	Neutron transport, Gamma production
0825	O-16	Neutron transport, Gamma production
0828	O-17	Neutron transport
0925	F-19	Neutron transport, Gamma production, Covariances
1125	Na-23	Neutron transport, Gamma production, Covariances
1200	Mg	Neutron transport, Gamma production
1225	Mg-24	Activation
1325	Al-27	Neutron transport, Gamma production, Covariances
1400	Si	Neutron transport, Gamma production, Covariances
1525	P-31	Neutron transport, Gamma production
1600	S	Neutron transport, Gamma production
1625	S-32	Neutron transport, Gamma production

MAT No. Material Usage

1700	Cl	Neutron transport, Gamma production
1837	Ar-40	Activation
1900	K	Neutron transport, Gamma production
1931	K-41	Activation
2000	Ca	Neutron transport, Gamma production
2125	Sc-45	Dosimetry
2200	Ti	Neutron transport, Gamma production
2225	Ti-46	Dosimetry
2228	Ti-47	Dosimetry
2231	Ti-48	Dosimetry
2237	Ti-50	Activation
2300	V	Neutron transport, Gamma production, Covariances
2425	Cr-50	Neutron transport, Gamma production, Covariances
2431	Cr-52	Neutron transport, Gamma production, Covariances
2434	Cr-53	Neutron transport, Gamma production, Covariances
2437	Cr-54	Neutron transport, Gamma production, Covariances
2525	Mn-55	Neutron transport, Gamma production, Covariances
2625	Fe-54	Neutron transport, Gamma production, Covariances
2631	Fe-56	Neutron transport, Gamma production, Covariances
2634	Fe-57	Neutron transport, Gamma production, Covariances
2637	Fe-58	Neutron transport, Gamma production, Covariances
2725	Co-59	Neutron transport, Gamma production, Covariances
2825	Ni-58	Neutron transport, Gamma production, Covariances
2828	Ni-59	Activation
2831	Ni-60	Neutron transport, Gamma production, Covariances
2834	Ni-61	Neutron transport, Gamma production, Covariances
2837	Ni-62	Neutron transport, Gamma production, Covariances
2843	Ni-64	Neutron transport, Gamma production, Covariances
2925	Cu-63	Neutron transport, Gamma production, Covariances
2928	Cu-65	Neutron transport, Gamma production, Covariances
3100	Ga	Neutron transport, Gamma production
3231	Ge-72	Fission product
3234	Ge-73	Fission product
3237	Ge-74	Fission product
3243	Ge-76	Fission product

MAT No. Material Usage

3325	As-75	Fission product
3425	Se-74	Fission product
3431	Se-76	Fission product
3434	Se-77	Fission product
3437	Se-78	Fission product
3443	Se-80	Fission product
3449	Se-82	Fission product
3525	Br-79	Fission product
3531	Br-81	Fission product
3625	Kr-78	Neutron transport, Fission product
3631	Kr-80	Neutron transport, Fission product
3637	Kr-82	Neutron transport, Fission product
3640	Kr-83	Neutron transport, Fission product
3643	Kr-84	Neutron transport, Fission product
3646	Kr-85	Fission product
3649	Kr-86	Neutron transport, Fission product
3725	Rb-85	Neutron transport, Fission product
3728	Rb-86	Fission product
3731	Rb-87	Neutron transport, Fission product
3825	Sr-84	Fission product
3831	Sr-86	Fission product
3834	Sr-87	Fission product
3837	Sr-88	Fission product
3840	Sr-89	Fission product
3843	Sr-90	Fission product
3925	Y-89	Neutron transport, Gamma production, Covariances
3928	Y-90	Fission product
3931	Y-91	Fission product
4000	Zr	Neutron transport
4025	Zr-90	Neutron transport
4028	Zr-91	Neutron transport
4031	Zr-92	Neutron transport, Activation
4034	Zr-93	Fission product
4037	Zr-94	Neutron transport
4040	Zr-95	Fission product

MAT No. Material Usage

4043	Zr-96	Neutron transport
4125	Nb-93	Neutron transport, Gamma production, Covariances
4128	Nb-94	Fission product
4131	Nb-95	Fission product
4200	Mo	Neutron transport, Gamma production
4225	Mo-92	Activation
4231	Mo-94	Fission product
4234	Mo-95	Fission product
4237	Mo-96	Fission product
4240	Mo-97	Fission product
4243	Mo-98	Activation
4246	Mo-99	Fission product
4249	Mo-100	Activation
4325	Tc-99	Fission product
4425	Ru-96	Fission product
4431	Ru-98	Fission product
4434	Ru-99	Fission product
4437	Ru-100	Fission product
4440	Ru-101	Fission product
4443	Ru-102	Fission product
4446	Ru-103	Fission product
4449	Ru-104	Fission product
4452	Ru-105	Fission product
4455	Ru-106	Fission product
4525	Rh-103	Fission product
4531	Rh-105	Fission product
4625	Pd-102	Fission product
4631	Pd-104	Fission product
4634	Pd-105	Fission product
4637	Pd-106	Fission product
4640	Pd-107	Fission product
4643	Pd-108	Fission product
4649	Pd-110	Fission product
4725	Ag-107	Neutron transport
4731	Ag-109	Neutron transport, Activation

MAT No. Material Usage

4737	Ag-111	Fission product
4800	Cd	Neutron transport
4825	Cd-106	Fission product
4831	Cd-108	Fission product
4837	Cd-110	Fission product
4840	Cd-111	Fission product
4843	Cd-112	Fission product
4846	Cd-113	Neutron transport
4849	Cd-114	Fission product
4853	Cd-115m	Fission product
4855	Cd-116	Fission product
4900	In	Neutron transport, Gamma production, Covariances
4925	In-113	Fission product
4931	in-115	Activation, Dosimetry
5025	Sn-112	Fission product
5031	Sn-114	Fission product
5034	Sn-115	Fission product
5037	Sn-116	Fission product
5040	Sn-117	Fission product
5043	Sn-118	Fission product
5046	Sn-119	Fission product
5049	Sn-120	Activation
5055	Sn-122	Activation
5058	Sn-123	Fission product
5061	Sn-124	Activation
5064	Sn-125	Fission product
5067	Sn-126	Fission product
5125	Sb-121	Fission product
5131	Sb-123	Fission product
5134	Sb-124	Fission product
5137	Sb-125	Fission product
5140	Sb-126	Fission product
5225	Te-120	Fission product
5231	Te-122	Fission product
5234	Te-123	Fission product

MAT No.	Material	Usage
5237	Te-124	Fission product
5240	Te-125	Fission product
5243	Te-126	Fission product
5247	Te-127m	Fission product
5249	Te-128	Fission product
5253	Te-129m	Fission product
5255	Te-130	Fission product
5261	Te-132	Fission product
5325	I-127	Activation
5331	I-129	Fission product
5334	I-130	Fission product
5337	I-131	Fission product
5349	I-135	Fission product
5425	Xe-124	Neutron transport
5431	Xe-126	Neutron transport
5437	Xe-128	Neutron transport
5440	Xe-129	Neutron transport
5443	Xe-130	Neutron transport
5446	Xe-131	Neutron transport
5449	Xe-132	Neutron transport
5452	Xe-133	Fission product
5455	Xe-134	Neutron transport
5458	Xe-135	Fission product
5461	Xe-136	Neutron transport
5525	Cs-133	Neutron transport
5528	Cs-134	Fission product
5531	Cs-135	Fission product
5534	Cs-136	Fission product
5537	Cs-137	Fission product
5637	Ba-134	Fission product
5640	Ba-135	Fission product
5643	Ba-136	Fission product
5646	Ba-137	Fission product
5649	Ba-138	Neutron transport, Gamma production
5655	Ba-140	Fission product

MAT No. Material Usage

5728	La-139	Fission product, Activation
5731	La-140	Fission product
5837	Ce-140	Fission product
5840	Ce-141	Fission product
5843	Ce-142	Fission product
5846	Ce-143	Fission product
5849	Ce-144	Fission product
5925	Pr-141	Neutron transport
5928	Pr-142	Fission product
5931	Pr-143	Fission product
6025	Nd-142	Fission product
6028	Nd-143	Neutron transport
6031	Nd-144	Fission product
6034	Nd-145	Neutron transport
6037	Nd-146	Neutron transport
6040	Nd-147	Fission product
6043	Nd-148	Neutron transport
6049	Nd-150	Neutron transport
6149	Pm-147	Neutron transport, Fission product
6152	Pm-148	Fission product
6153	Pm-148m	Fission product
6155	Pm-149	Fission product
6161	Pm-151	Fission product
6225	Sm-144	Fission product
6234	Sm-147	Neutron transport, Fission product
6237	Sm-148	Fission product
6240	Sm-149	Neutron transport
6243	Sm-150	Fission product
6246	Sm-151	Neutron transport, Fission product
6249	Sm-152	Neutron transport
6252	Sm-153	Fission product
6255	Sm-154	Fission product
6325	Eu-151	Neutron transport, Gamma production
6328	Eu-152	Neutron transport, Fission product
6331	Eu-153	Neutron transport, Gamma production

MAT No. Material Usage

6334	Eu-154	Neutron transport, Fission product
6337	Eu-155	Neutron transport, Fission product
6340	Eu-156	Fission product
6343	Eu-157	Fission product
6425	Gd-152	Neutron transport
6431	Gd-154	Neutron transport
6434	Gd-155	Neutron transport
6437	Gd-156	Neutron transport
6440	Gd-157	Neutron transport
6443	Gd-158	Neutron transport
6449	Gd-160	Neutron transport
6525	Tb-159	Fission product
6528	Tb-160	Fission product
6637	Dy-160	Fission product
6640	Dy-161	Fission product
6643	Dy-162	Fission product
6646	Dy-163	Fission product
6649	Dy-164	Neutron transport
6725	Ho-165	Neutron transport, Gamma production
6837	Er-166	Fission product
6840	Er-167	Fission product
7125	Lu-175	Neutron transport
7128	Lu-176	Neutron transport
7200	Hf	Neutron transport
7225	Hf-174	Neutron transport
7231	Hf-176	Neutron transport
7234	Hf-177	Neutron transport
7237	Hf-178	Neutron transport
7240	Hf-179	Neutron transport
7243	Hf-180	Neutron transport
7328	Ta-181	Neutron transport, Gamma production
7331	Ta-182	Neutron transport
7400	W	Neutron transport, Gamma production
7431	W-182	Neutron transport, Gamma production
7434	W-183	Neutron transport, Gamma production

MAT No. Material Usage

7437	W-184	Neutron transport, Gamma production
7443	W-186	Neutron transport, Gamma production
7525	Re-185	Neutron transport, Covariances
7531	Re-187	Neutron transport, Covariance
7925	Au-197	Standard, Neutron transport, Gamma production, Covariances
8231	Pb-206	Neutron transport, Gamma production, Covariances
8234	Pb-207	Neutron transport, Gamma production, Covariances
8237	Pb-208	Neutron transport, Gamma production, Covariances
8325	Bi-209	Neutron transport, Gamma production, Covariances
9034	Th-230	Neutron transport
9040	Th-232	Neutron transport, Gamma production, Covariances
9131	Pa-231	Neutron transport
9137	Pa-233	Neutron transport
9219	U-232	Neutron transport
9222	U-233	Neutron transport, Gamma production
9225	U-234	Neutron transport
9228	U-235	Neutron transport, Gamma production, Covariances
9231	U-236	Neutron transport
9234	U-237	Neutron transport, Gamma production
9237	U-238	Neutron transport, Gamma production, Covariances
9346	Np-237	Neutron transport, Gamma production
9349	Np-238	Neutron transport
9352	Np-239	Neutron transport
9428	Pu-236	Neutron transport
9431	Pu-237	Neutron transport
9437	Pu-239	Neutron transport, Gamma production
9440	Pu-240	Neutron transport, Gamma production, Covariances
9443	Pu-241	Neutron transport, Gamma production, Covariances
9543	Am-241	Neutron transport, Gamma production, Covariances
9446	Pu-238	Neutron transport
9449	Pu-243	Neutron transport, Gamma production
9452	Pu-244	Neutron transport
9546	Am-242	Neutron transport
9547	Am-242m	Neutron transport, Gamma production

MAT NO.	Material	Usage
---------	----------	-------

9549	Am-243	Neutron transport, Gamma production
9628	Cm-241	Neutron transport
9631	Cm-242	Neutron transport, Gamma production
9634	Cm-243	Neutron transport, Gamma production
9637	Cm-244	Neutron transport, Gamma production
9640	Cm-245	Neutron transport, Gamma production
9643	Cm-246	Neutron transport, Gamma production
9646	Cm-247	Neutron transport, Gamma production
9649	Cm-248	Neutron transport, Gamma production
9752	Bk-249	Neutron transport
9852	Cf-249	Neutron transport
9855	Cf-250	Neutron transport, Gamma production
9858	Cf-251	Neutron transport, Gamma production
9861	Cf-252	Neutron transport, Gamma production
9864	Cf-253	Neutron transport (total, elastic, fission, capture only)
9913	Es-253	Neutron transport (total, elastic, capture only)

Thermal Scattering Law Sublibrary

MAT No	Material	Usage
--------	----------	-------

0001	Water	Thermal scattering law
0007	H in ZrH	Thermal scattering law
0011	Heavy Water	Thermal scattering law
0026	Beryllea	Thermal scattering law
0027	Beryllium O	Thermal scattering law
0031	Graphite	Thermal scattering law
0037	Polyethelene	Thermal scattering law
0040	Benzene	Thermal scattering law
0058	Zr in ZrH	Thermal scattering law

High Energy Evaluations

Neutron Sublibrary

MAT No.	Material	Usage
---------	----------	-------

2631	Fe-56	High energy, Neutron transport, Gamma production
------	-------	--

Proton Sublibrary

MAT No.	Material	Usage
---------	----------	-------

2631	Fe-56	High energy, Proton transport, Gamma production
------	-------	---

Contributors to Documentation

G. M. Hale	Los Alamos National Laboratory
D. Larson	Oak Ridge National Laboratory
S. Pearlstein	Brookhaven National Laboratory
R. E. Schenter	Hanford Engineering Development Laboratory
L. W. Weston	Oak Ridge National Laboratory
R. Q. Wright	Oak Ridge National Laboratory
P. G. Young	Los Alamos National Laboratory

Introduction

Introduction

Responsibility for oversight of the ENDF/B Evaluated Nuclear Data File lies with the Cross Section Evaluation Working Group (CSEWG), which is comprised of representatives from various governmental and industrial laboratories in the United States. Individual evaluations are provided by scientists at several U.S. laboratories, including significant contributions by scientists from all over the world. In addition, ENDF/B-VI includes for the first time complete evaluations for three materials that were provided from laboratories outside the U.S. All data are checked and reviewed by CSEWG, and the data file is maintained and issued by the National Nuclear Data Center at Brookhaven National Laboratory. The previous version of the library, ENDF/B-V, was issued in 1979, and two revisions to the data file were provided in subsequent years, the latest occurring in 1981.

Preparation for Version VI of ENDF/B has been underway since the early 1980's. Since the issue of ENDF/B-V, a large quantity of new experimental data has become available for a variety of nuclear reactions. Additionally, improved nuclear models and theoretical codes have been developed to permit more reliable interpolation and extrapolations of nuclear data into unmeasured regions. Significant extensions and revisions have been made to the data formats available for ENDF/B-VI evaluations, including provisions for incident charged particles and photo-nuclear data by partitioning the ENDF library into sublibraries, new multilevel resonance formulae, and several options for representing energy-angle correlated emission data. The latter development is essential for extending the evaluated data files to higher incident energies, and several evaluations with maximum energies greater than 20 MeV are included in Version VI. Considerable effort has been directed at developing the computational tools required to utilize the new formatting capabilities.

The most comprehensive and thorough analysis ever attempted by CSEWG was performed for the cross sections of standard reactions. This study incorporated cross sections and covariances for the most important heavy element standards (including thermal standards) into a simultaneous analysis, which was then combined with results from detailed coupled-channel R-matrix analyses of the light element standards. Not only were the cross sections for the standard reactions included in the analysis, but also absolute data for other important reactions that are linked to the standards through ratio measurements were also incorporated. Examples of such related data that were included in the simultaneous analysis are the $^{238}\text{U}(n,\gamma)$, $^{238}\text{U}(n,f)$, and $^{239}\text{Pu}(n,f)$ cross sections. Largely because of the standards analysis, we feel that

ENDF/B-VI should be the most internally self-consistent evaluation to date.

A total of 75 new or extensively modified neutron sublibrary evaluations are included in ENDF/B-VI, and are summarized in this document. One incident proton sublibrary is described for Fe⁵⁶. The remaining evaluations in ENDF/B-VI have been carried over from earlier versions of ENDF, and have been updated to reflect the new formats. The release of ENDF/B-VI was carried out between January and June of 1990, with groups of materials being released on "tapes". Table 1 is an index to the evaluation summaries, and includes the material identification or MAT number, the responsible laboratory, and the "tape" number. These evaluations have been released without restrictions on their distribution or use.

Table 1. Summary of Evaluations

Material	MAT	Laboratory	Tape	Page
H-1	125	LANL	100	7
H-2	128	LANL	116	13
He-3	225	LANL	116	15
Li-6	325	LANL	100	18
Li-7	328	LANL	100	33
Be-9	425	LLNL	100	40
B-10	525	LANL	100	48
B-11	528	LANL	100	69
C	600	ORNL	100	78
N-14	725	LANL	116	82
N-15	728	LANL	116	91
O-16	825	LANL	116	101
F-19	925	ORNL	115	120
V	2300	ANL	103	124
Cr-50	2425	ORNL	111	131
Cr-52	2431	ORNL	111	131
Cr-53	2434	ORNL	111	131
Cr-54	2437	ORNL	11	131
Mn-55	2525	ORNL	114	144
Fe-54	2625	ORNL	112	149
Fe-56	2631	ORNL	112	149

Table 1. (Continued)

Material	MAT	Laboratory	Tape	Page
Fe-56 (n*)	2631	BNL	800	168
Fe-56 (p*)	2631	BNL	801	168
Fe-57	2634	ORNL	112	149
Fe-58	2637	ORNL	112	149
Co-59	2725	ANL	103	174
Ni-58	2825	ORNL	113	181
Ni-59	2828	WHC	113	197
Ni-60	2831	ORNL	113	181
Ni-61	2834	ORNL	113	181
Ni-62	2837	ORNL	113	181
Ni-64	2843	ORNL	113	181
Cu-63	2925	ORNL	114	198
Cu-65	2931	ORNL	114	198
Y-89	3925	ANL	103	211
Nb	4125	ANL	116	216
Pd-105	4634	ORNL	103	222
Pd-107	4640	ORNL	103	225
In	4900	ANL	116	228
In-115	4931	ANL	116	234
Cs-134	5528	ORNL	103	238
Ba-134	5637	ORNL	103	241
Ba-135	5640	ORNL	103	245
Ba-136	5643	ORNL	103	249
Ba-137	5646	ORNL	103	253
Nd-147	6040	ORNL	103	256
Pm-147	6149	ORNL	103	260
Sm-147	6234	ORNL	103	263
SM-151	6246	ORNL	103	268
Eu-151	6325	LANL	103	271
Eu-152	6328	ORNL	103	278
Eu-153	6331	LANL	103	285
Eu-154	6334	ORNL	103	291
Eu-155	6337	ORNL	103	298
Ho-165	6725	LANL	103	302

* Data up to 1 GeV incident energy.

Table 1. (Concluded)

Material	MAT	Laboratory	Tape	Page
Er-166	6837	ORNL	103	309
Er-167	6840	ORNL	103	313
Re-185	7525	ORNL	115	317
Re-187	7531	ORNL	115	321
Au-197	7925	LANL	108	325
Pb-206	8231	ORNL	115	336
Pb-207	8234	ORNL	115	336
Pb-208	8237	ORNL	115	336
Bi-209	8325	ANL	108	351
U-235	9228	ORNL	117	356
U-236	9231	WHC	108	376
U-238	9237	ORNL	117	386
Np-237	9346	LANL	117	397
Np-239	9352	ORNL	108	411
Pu-239	9437	LANL	117	415
Pu-240	9440	ORNL	108	434
Pu-241	9443	ORNL	108	446
Am-241	9543	CNDC	108	453
Am-243	9549	ORNL	108	457
Bk-249	9752	CNDC	108	465
Cf-249	9852	CNDC	108	468

ANL Argonne National Laboratory
 BNL Brookhaven National Laboratory
 CNDC Chinese Nuclear Data Center
 LANL Los Alamos National Laboratory
 LLNL Lawrence Livermore National Laboratory
 ORNL Oak Ridge National Laboratory
 WHC Westinghouse Hanford Company

^1_1H

Reference: No Primary Reference
Evaluators: G. M. Hale, D. C. Dodder, E. R. Siciliano, and W. B. Wilson
Evaluated: October 1989
Material: 125
Content: Standard, Neutron transport, Gamma production

File Comments

The ENDF/B-VI cross sections for hydrogen represent the first new evaluation work on n-p scattering since those based on the Hopkins-Breit phase shifts were placed in the file. The new cross sections result from a charge-independent R-matrix analysis of n-p and p-p scattering at energies below 30 MeV that was done by Dodder and Hale.¹ A summary of the channel configuration and data fitting characteristics of the analysis is given in Table 1.

Table 1. 0-30 MeV N-N R-Matrix Analysis

	Channel	ℓ_{max}	a_r (fm)
	n - p	3	3.26
	p - p	3	3.26

Reaction	# Observable Types	# Data Points	χ^2
n-p scattering	3	448	407
p-p scattering	4	388	399
Totals:	7	836	806

| # of parameters = 33 $\Rightarrow \chi^2$ per degree of freedom = 1.004 * | | | |

* Including recent corrections⁵ to the 16.9 Mev n-p analyzing power data of Tornow et al.⁶ reduces the overall chi-square per degree of freedom of the fit to 0.9988.

The R-matrix analysis includes many n-p measurements that were not available at the time of the Hopkins-Breit phase-shift analysis, and gives a representation of

the n-p and p-p data in the 0-30 MeV range that is comparable to or better than that of other recent work.^{2,3} The new analysis also gives predictions for newly measured observables, such as the polarization-transfer data from Karlsruhe⁴, that looks quite reasonable.

The charge independent model used takes the isospin-1 reduced-width amplitudes in the R matrix describing n-p scattering to be identically the same as those describing p-p scattering. The energy eigenvalues in the two systems are taken to differ only by an overall constant Coulomb energy shift. This simple model allows the p-p scattering data to influence the n-p fit. We see in Fig. 1 where measurements of the cross section and analyzing powers for the two reactions are compared, that the data are quite different at the same energy. These differences, coming primarily from Coulomb terms and symmetrization properties of the two systems, are well reproduced by the charge-independent calculation. The calculations also account well for the shape of the n-p angular measurement⁷ shown at 14 MeV.

Two quantities often used to characterize the center of mass n-p angular distribution near 14 MeV are the back-angle cross section, $\sigma(180^\circ)$, and the asymmetry ratio $R = \sigma(180^\circ)/\sigma(90^\circ)$. The ENDF/B-VI evaluation gives for these quantities at $E_n = 14.1$ MeV:

$$\sigma(180^\circ) = 58.89 \pm 0.60 \text{ mb}, \quad R = 1.093 \pm 0.010.$$

The value of R is in agreement with most previous measurements, but disagrees with a recent measurement of Ryves and Kolkowski⁸ ($R = 1.053 \pm 0.015$) that is consistent with the ENDF/B-V value. The ENDF/B-VI values of the back-angle cross section and asymmetry ratio, on the other hand, are in excellent agreement with an evaluation of the 14.1 MeV data that was done in 1982 by Vincour, Bém, and Près-perin.⁹

Elastic cross sections and angular distributions below 30 MeV were determined with the new R-matrix calculation by Hale and Dodder. This calculation gives the ENDF/B-VI elastic scattering cross sections for hydrogen below 26 MeV.

Elastic cross sections and angular distributions above 20 MeV were calculated with the NPSCAT code using phase shifts of Arndt² by Siciliano and Wilson. The angular distributions at 26 MeV agreed well. Below this energy R-matrix angular distributions were used, and at 26 MeV and above phase shift angular distributions were used. Cross sections above 30 MeV were taken from the phase shift work. Between these energies the elastic cross section was taken as follows:

E (eV)	R-matrix	Phase shift	σ
2.6E+07	.36345	.36029	.36345
2.7E+07	.34859		.34811
2.8E+07	.33488	.33104	.33295
2.9E+07	.32227		.31891
3.0E+07	.31080	.30567	.30567

The calculated capture results were merged with available n,γ data by W. B. Wilson. ENDF/B-V (n,γ) data were used below 20 MeV; above this energy, an approximation to the data of M. Bosman et al.¹⁰ was used. The (n,γ) was adjusted to agree exactly with Mughabghab's (1983) value at thermal (0.3326 b), and lower energies were modified according to the $1/v$ law (P. Young, 10/17/89). The total cross section was then summed again to reflect the revised (n,γ).

The scattering radius for File 2 and the photon production files for capture (MF=12 and MF=14) were taken from ENDF/B-V.

References

1. D. C. Dodder and G. M. Hale, to be published (1991), See G. M. Hale & P. G. Young, LANL Report LA-UR 90-1078 (1990).
2. R. A. Arndt, "N-N Phase Shift Analysis," Interactive computer Program SAID, Private Communication (1988).
3. M. M. Nagels, T. A. Rijken, and J. J. deSwaart, "Low Energy Nucleon Nucleon Potential from Regge Pole Theory," Phys. Rev. D17, 768 (1978).
4. H. Klages et al., "Karlsruhe Polarization Transfer Measurements for n-p Scattering," Private Communication from W. Tornow, Duke University (1988).
5. W. Tornow et al., Phys. Rev. C37, 2326 (1988).
6. W. Tornow, P. Lisowski, R. C. Byrd, and R. L. Walter, Phys. Rev. Letters 39, 915 (1977) and Nucl. Phys. A310, 31 (1980).
7. T. Nakamura, J. Phys. Soc. Japan 15, 1359 (1960).

8. T. B. Ryves and P. Kolkowski, "The Differential Cross Section for Neutron-Proton Scattering at 14.5 MeV," Preliminary Draft, National Physical Laboratory, Middlesex, UK (March 1990).
9. J. Vincour, P. Bém, and V. Presperin, "Angular Distribution of Neutron-Proton Scattering at 14.1 MeV," in *Neutron Induced Reactions*, Proceedings of the Europhysics Topical Conference, Smolinice, 413 (1982).
10. M. Bosman et al., Phys. Letters 82b, 242 (1979).

Fig. 1 Comparison of calculated and measured values of differential cross sections ($E_{n,p} \approx 14$ MeV) and analyzing powers ($E_{n,p} \approx 16.5$ MeV) for neutrons and protons incident on ^1H .

Fig. 2 Neutron total cross section for ^1H from R-matrix fit that resulted in the ENDF/B-VI standard $^1\text{H}(n,n)^1\text{H}$ cross section.

${}^2_1\text{H}$

Reference: LA-3271 (1968)
Evaluators: L. Stewart, R. E. MacFarlane, (LASL) A. Horsley
(AWRE)
Evaluated: December 1989
Material: 128
Content: Neutron transport, Gamma production

File Comments

LASL	Eval-Dec89 R. E. MacFarlane
LASL, AWRE	Eval-Nov67 L. Stewart and A. Horsley

Summary of Changes

The ENDF/B-V file for deuterium MAT=1302 was converted for ENDF/B-VI by R. E. MacFarlane (LASL) in December 1989. The reaction cross sections, elastic angular distributions, and photon production data were left unchanged. MF=4 and MF=5 for the (n,2n) reaction were removed and replaced by an MF=6 using an analytic phase-space representation. This is consistent with the original evaluation, but provides a more accurate representation of the energy-angle correlation of (n,2n) neutrons.

Summary of ENDF/B-V Evaluation

MF=2 No resonance parameters given.

MF=3 MT=1 Total cross sections. All data was plotted and compared up to 1967 in LA-3271. Changes were incorporated below 1.5 MeV. The evaluation does not agree with low-energy experiments at the NBS (which are preliminary) but agrees at higher energies. The Davis data show a peculiar drop of a few percent from 3.5 to 9 MeV but agree above and below these energies.

Summary of ENDF/B-V, Continued

- MF=3 MT=2 Elastic cross sections. Data were obtained from integrating n-d and p-d angular distributions. Since the radiative capture is in microbarns, the elastic is essentially equal to the total cross section below the n,2n threshold and the total minus the n,2n above the threshold. Checks and balances were always made. See LA-3271 for the graphical comparisons.
- MT=16 The n,2n cross section. Data were taken from Holmberg and from Catron. See LA-3271. Nothing is known about the cross section above 14 MeV.
- MT=102 Radiative capture cross section. The thermal cross section is 506 microbarns which was extrapolated as $1/v$ up to 1 keV. A curve was drawn above this energy to include measurements on the inverse reaction by Bosch. The 14 MeV value is a factor of 3 lower than Cerineo. See LA-3271 for graphical results.
- MF=4 MT=2 Elastic angular distributions taken from n-d and p-d scattering data. The agreement is consistent with the Van Oers analysis. See LA-3271.
- MF=6 MT=16 Energy-angle energy-angle distributions represented as a 3-body phase space distribution.
- MF=8 MT=102 Radioactive decay information added for tritium production.
- MF=9 MT=102 Multiplicity for production of radioactive tritium.
- MF=12 MT=102 Multiplicity for photon production. A single gamma was assumed to be emitted at all energies. The LP = 2 flag was used to conserve energy. For references, see LA-3271.
- MF=14 MT=102 Angular distributions of the secondary photon.

^3_2He

Reference: No Primary Reference
Evaluators: G. Hale, D. Dodder, and P. Young
Evaluated: May 1990
Material: 225
Content: Standard, Neutron transport

HE-3 FREE ATOM EVALUATION

This file transferred from L. Stewart's ENDF/B-III evaluation with modifications only in the $^3\text{He}(n,p)$ and the elastic cross sections below 5 MeV, and the total cross section below 0.1 MeV.

* * * * *
* * * * *
* * * * *
* Cross section standard He-3 (n,p) *
* * * * *
* The He-3 (n,p) cross section for this material is recommended
* as a standard for neutron energies from thermal to 50 keV.
* * * * *

MF=1

MT=451, Atomic mass = 3.0160.

MF=2

MT=151, Scattering length = 0.2821E-12 cm.

MF=3

MT= 1, Total cross sections --- from .00001 eV to 200 keV.
MT1 taken as sum MT2+MT102+MT103. From 200 keV to 20 MeV, MT1 evaluated using experimental data from Ref.6.

MT= 2, Elastic scattering cross sections --- from .00001 eV to 200 keV, MT2 taken approximately from R-matrix analysis described below. Above 200 keV, obtained

from MT2 = MT1-MT102-MT103-MT104. Note two reactions missing from this evaluation, namely n,n-prime,p and n,2n,2p. Exp. data at 15 MeV indicates non-zero cross sections for these. Included in MT2 this eval.

MT=102, Radiative capture cross sections --- thermal value taken from Ref.17, with assumption of 1/v energy energy variation at all other energies.

MT=103, n-p cross sections --- below 5 MeV taken from an R-matrix calculation of the four-body reactions by Hale and Dodder. $^3\text{He}(n,p)$ data included were from Refs. 5,13,14, and 15. For energies in the 5-20 MeV range, data were used as follows -
 Ref.4 - 5.8 MeV to 11. MeV
 Ref.10- 5.0 MeV to 11. MeV
 Ref.12- 5.0 MeV to 12. MeV
 Ref.16- 14. MeV
 Data extrapolated to 20. MeV.

MT=104, n-d cross sections --- threshold = 4.36147 MeV.
 $Q = -3.2684$ MeV. Evaluation from a detailed balance calculation (Ref.13) and experimental data (Ref.11).

MT=251, Average value of cosine of elastic scattering angle, laboratory system. Obtained from data MF=4, MT=2.

MT=252, Values of Ξ , obtained from data MF=4, MT=2.

MT=253, Values of Γ , obtained from data MF=4, MT=2.

MF=4

MT= 2, Angular distribution of secondary neutrons from elastic scatter. Evaluated from experimental data from Refs.2,7,9,11,16,17 covering incident energies as follows -

Incident energy	References
1.E-5eV	(isotropic)
0.5 MeV	(isotropic)
1.0 MeV	9
2.0 MeV	9
2.6 MeV	11
3.5 MeV	9
5.0 MeV	11
6.0 MeV	9,7(from p+t elastic scatt)
8.0 MeV	11,7(from p+t elastic scatt)
14.5 MeV	16,17(from p+t elastic scatt)

17.5 MeV
20.0 MeV

11
2(from p+t elastic scatt)

References

1. R.Batchelor,R.Aves,and T.H.R.Skyrme,Rev.Sci.Instr.26,1037 (1955).
2. R.A.Vanetsian and E.D.Fenchenko,Soviet Journal of Atomic Energy 2,141(1957).
3. J.N.Bradbury and L.Stewart,Bull.Am.Phys.Soc.3,417(1958).
4. G.F.Bogdanov,N.A.Vlasov,C.P.Kalinin,B.V.Rybakov,L.N.Samoilov,and V.A.Sidorov,JETP(USSR)36,633(1959).
5. J.H.Gibbons and R.L.Macklin,Phys.Rev.114,571(1959).
6. Los Alamos Physics and Cryogenics Groups,Nucl.Phys.12, 291(1959).
7. J.E.Brolley,Jr.,T.M.Putnam,L.Rosen,and L.Stewart,Phys. Rev.159,777(1967).
8. J.E.Perry,Private Communication to Stewart(1960)
9. J.D.Seagrave,L.Cranberg,and J.E.Simmons,Phys.Rev.119, 1981(1960).
10. M.D.Goldberg,J.D.Anderson,J.P.Stoering,and C.Wong,Phys. Rev.122,1510(1961).
11. A.R.Sayers,K.W.Jones,and C.S.Wu,Phys.Rev.122,1853(1961).
12. W.E.Wilson,R.L.Walter,and D.B.Fossan,Nucl.Phys.27,421 (1961).
13. J.Als-Nielsen and O.Dietrich,Phys.Rev.133,B925(1964).
14. R.L.Macklin and J.H.Gibbons,Proceedings of the Internatl. Conf.on the Study of Nucl.Struct.with Neut.,Antwerp(1965).
15. J.H.Gibbons,Private Communication to Stewart(1966).
16. B.Antolkovic,G.Paic,P.Tomas,and R.Rendic,Phys.Rev.159, 777(1967).
17. L.Rosen and W.Leland,Private Communication(1967).
18. S.Mughabghab et al., Neut.Cross Sect., Vol.1, Neut.Reson. Parameters and Thermal Cross Sect., Academic Pr.(1981).

${}^6_3\text{Li}$

Reference: No Primary Reference
Evaluators: G. M. Hale and P. G. Young
Evaluated: April 1989
Material: 325
Content: Standard, Neutron transport, Gamma production

ENDF/VI EVALUATION

G. M. Hale and P. G. Young

MAJOR CHANGES FROM VERSION V OF ENDF/B ARE:

1. Inclusion of the ENDF/B-VI standard (n,t) cross section from the simultaneous standards analysis (Ca85) over the energy range thermal to 1 MeV.
2. Replacement of all major cross sections and elastic angular distributions at energies between 10E-5 eV and 3 MeV with results from the R-matrix analysis performed in conjunction with the simultaneous standards analysis.
3. Revision of the elastic cross sections and angular distributions at energies between 3 and 20 MeV to match recent experimental data, resulting in a general decrease of the elastic cross section in this energy range.
4. Revision of the (n,n')d cross sections to account for recent measurements, resulting in a general increase in the total (n,n')d cross section that tends to offset the decrease in the elastic cross section and maintain about the same total cross section as before.

MF=2 ----- Resonance parameters -----

MT=151 Effective scattering radius = 2.31175E-13 cm.

MF=3 ----- Smooth cross sections -----

The 2200 m/s cross sections are as follows:

MT=1 sigma = 941.6928 barns

MT=2 sigma = 0.67157 barns

MT=102 sigma = 0.03850 barns

MT=105 sigma = 940.9827 barns

MT=1 Total cross section

below 3 MeV, the values are taken from an R-matrix

analysis by Hale, Dodder, and Witte (Ha84) which takes into account data from all reactions possible in Li-7 up to 4 MeV neutron energy. Total cross section data considered in this analysis were those of HA75 and SM77. Between 3 and 20 MeV, the total was taken to be the sum of MT=2,4,24,102,103, and 105, which generally follows the measurements of Sm82, Ke79, Kn77, Go72, and Fo71.

MT=2 Elastic cross section

below 3 MeV, the values are taken from the R-matrix analysis cited for MT=1, which includes the elastic measurements of Sm82 and La61. Above 3 MeV, the curve is a smooth representation of the data of Kn79 and Ba63 up to 7.5 MeV, and of that of Ho79 between 7.5 and 13 MeV. The curve passes through the average of several measurements at 14 MeV, and is extrapolated to 20 MeV using the shape of an optical model calculation.

MT=4 Total inelastic cross section

Sum of MT=51 through MT=81.

MT=24 (n,2n)alpha cross section

passes through the point of Mather and Paine (Ma69) at 14 MeV, taking into account the measurements of As63.

MT=51,52,54-56,58-81 (n,n')d continuum

represented by continuum-level contributions in Li-6, binned in .5-MeV intervals. The energy-angle spectra are determined by a 3-body phase-space calculation, assuming isotropic center-of mass distributions. At each energy, the sum of the continuum-level contributions is normalized to an assumed energy-angle integrated continuum cross section which approximates the difference of the nonelastic sigma and the contribution from the first and second levels in Li-6. The steep rise of the pseudo-level cross sections from their thresholds and the use of fixed bin widths over finite angles produces anomalous structure in the individual cross sections which is especially apparent near the thresholds. Some effort has been made to smooth out these effects, but they remain to some extent.

MT=53 (n,n1)d discrete level cross section

has p-wave penetrability energy dependence from threshold to 3.2 MeV. Matched at higher energies to a curve through fitted legendre coefficients from experimental data of Sa82, Ho79, Sm80, Ho68, Ba63.

MT=57 (n,n2)gamma cross section

is based on the available experimental data, especially that of Ho79, Li80, Sm82, Ho68.

Gradually to 20 MeV, a smooth curve was drawn through data

of Pr69 and Be75.

- MT=102 (n,gamma) cross section
unchanged from version V, which was based on the thermal measurement of Journey (Ju73) and the Pendlebury evaluation (Pe64) at higher energies.
- MT=103 (n,p) cross section
threshold to 9 MeV, based on the data of Ba65. Extended to 20 MeV through the 14 MeV data of Fr54 and Ba53.
- MT=105 (n,t) cross section
below 3 MeV, values are taken from the R-matrix analysis, which includes (n,t) measurements from Re78, La78, Br77, Ov74, and Ba75. Between 3 and 5 MeV, the values are based on Ba75, and at higher energies are taken from the evaluation of Pe64, extended to 20 MeV considering the data of Ke58.

MF = 4-----Angular Distributions-----

- MT=2 Elastic cross section
legendre coefficients determined as follows:
below 4 MeV, coefficients up to l=6 were taken from the R-matrix analysis, which included the measurements La61 and Sm82. Above 4 MeV, the coefficients represent fits to the measurements of Ho68, Ho79, Kn79, Sm82, De73, Ba63, Ab70, and Hy68. Most emphasis was placed on the data of Ho79, Kn79, Sm82. Extrapolation of the coefficients to 20 MeV was aided by optical model calculations.
- MT=24 (n,2n) cross section
lab distributions obtained by integrating over energy the 4-body phase-space spectra that result from transforming isotropic center-of-mass distributions to the laboratory system.
- MT=51-81 (n,n')d cross sections
excitation energy binned data is assumed isotropic in the center of mass reference system. MT = 53 and 57 are real levels. MT = 57 is assumed to be isotropic in the two-body reference system. MT = 53 is given as anisotropic, based on fits of legendre expansions to the experimental data of Ab70, Ba63, Ho68, Ho79, Me65, Hy68, Wo62, Sa82.
- MT=105 (n,t) cross section (to be added)
legendre coefficients obtained from the R-matrix analysis are supplied at energies below 4 MeV. The analysis takes into account (n,t) angular distribution measurements from Kn83, Co82, Dr82, Br77, Ba75, and Ov74.

MF = 5-----Secondary energy distributions-----

MT=24 (n,2n)

lab distributions obtained by integrating over angle the 4-body phase-space spectra that result from transforming isotropic center-of-mass distributions to the laboratory system.

MF = 12-----Gamma-ray multiplicities-----

MT=57 (n,n2)gamma

energy taken from Aj74. Multiplicity assumed to be one.

MT=102 (n,gamma)

energies and transition arrays for radiative capture taken from Ju73, as reported in Aj74. The LP flag was used to describe the MT=102 photons.

MF = 14-----Gamma-ray angular distributions-----

MT=57 (n,n2)gamma

the gamma is assumed isotropic.

MT=102 (n,gamma)

the two high-energy gammas are assumed isotropic. Data on the 477 kev gamma indicates isotropy.

MF=33-----Cross section covariances-----
(to be added later)

The relative covariances for MT=1,2, and 105 below 4 MeV are given in File 33. They are based on calculations using the covariances of the R-matrix parameters in first-order error propagation.

MT=1 Total

relative covariances entered as NC-type sub-subsection, implying that they are to be constructed from those for MT=2 and 105. They are not intended for use at energies above 4 mev.

MT=2,105 Elastic and (n,t)

relative covariances among these two cross sections are entered explicitly as NI-type sub-subsections in the LB=5 (direct) representation at energies below 4 MeV. Although values for the 3.95 - 4.05 MeV bin are repeated in a 4 - 20 MeV bin, the covariances are not intended for use at energies above 4 MeV.

----- References -----

- Ab70 U. Abbondanno, Nuo. Cim. A166, 139 (1970).
- Aj74 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227, 55 (1974).
- Ar64 A. H. Armstrong, J. Gammel, L. Rosen, and G. M. Frye, Nucl. Phys. 52, 505 (1964).
- As63 V. J. Ashby et al., Phys. Rev. 129, 1771 (1963).
- Ba53 M. E. Battat and F. L. Ribe, Phys. Rev. 89, 80 (1953).
- Ba63 R. Batchelor and J. H. Towle, Nucl. Phys. 47, 385 (1963).
- Ba65 R. Bass, C. Bindhardt, and K. Kruger, EANDC(E)-57U (1965).
- Ba75 C. M. Bartle, Proc. Conf. on Nuclear Cross Sections and Technology, Vol. 2, 688 (1975), and private communication (1976). See also Nucl. Phys. A330, 1 (1979).
- Be75 Besotosnyj et al., YK-19, 77 (1975).
- Br71 R. E. Brown, G. G. Ohlsen, R. F. Haglund, and N. Jarmie, Phys. Rev. 16C, 513 (1977).
- Ca85 A. D. Carlson, W. P. Poenitz, G. M. Hale, and R. W. Peele, Nuclear Data for Basic and Applied Science (Santa Fe, N.M.), 1429 (1985).
- Co67 J. A. Cookson and D. Dandy, Nucl. Phys. A91, 273 (1967).
- Co82 H. Conde, T. Andersson, L. Nilsson, and C. Nordborg, Nuclear Data for Science and Technology (Antwerp, Belgium), 447 (1982).
- De73 F. Demanins et al., INFN/BE-73 (1973).
- Dr82 M. Drosch, D. M. Drake, R. A. Hardekopf, and G. M. Hale, LA-9129-MS (1982).
- Dr85 M. Drosch et al., Santa Fe Conf. 1, 145 (1985).
- Fo71 D. G. Foster and D. W. Glasgow, Phys. Rev. C3, 576 (1971).
- Fr54 G. M. Frye, Phys. Rev. 93, 1086 (1954).
- Go72 C. A. Goulding and P. Stoler, EANDC(US)-176U, 161 (1972).
- Ha75 J. A. Harvey and N. W. Hill, Nuclear Cross Sections and Technology (Washington, D.C.), 244 (1975).
- Ha84 G. M. Hale, Nuclear Standard Reference Data (Geel, Belgium) IAEA TECDOC-335, 103 (1984). Describes preliminary analysis.
- Ho68 J. C. Hopkins, D. M. Drake, and H. Conde, Nucl. Phys. A107, 139 (1968), and J. C. Hopkins, D. M. Drake, and H. Conde, LA-3765 (1967).
- Ho79 H. H. Hogue et al., N.S.&E. 69, 22 (1979).
- Ju73 E. T. Jurney, LASL, Private Communication (1973).
- Ke58 R. D. Kern and W. E. Kreger, Phys. Rev. 112, 926 (1958).
- Ke79 J. D. Kellie, G. P. Lamaze, and R. B. Schwartz, Nuclear Cross Sections for Technology (Knoxville, Tn.), 48 (1979).
- Kn77 H. H. Knitter, C. Budtz-Jorgensen, M. Mailly, and R. Vogt, EUR-5726e (1977).
- Kn79 H. D. Knox, R. M. White, and R. O. Lane, N.S.&E. 69, 223 (1979).
- Kn83 H. H. Knitter, C. Budtz-Jorgensen, D. L. Smith, and D. Marletta, N.S.&E. 83, 229 (1983).
- La61 R. O. Lane, A. S. Langsdorf, J. E. Monahan, and A. J. Elwyn, Ann. Phys. 12, 135 (1961).

- La78 G.P.Lamaze, O.A.Wasson, R.A.Schrack, and A.D.Carlson, N.S.&E.
68, (1978).
- Li80 P.W.Lisowski et al., LA-8342 (1980).
- Ma69 D.S.Mather and L.F.Paine, AWRE-O-47/69 (1969).
- Me65 F.Merchez, N.V.Sen, V.Regis, and R.Bouchez, Compt. Rend. 260,
3922 (1965).
- Ov74 J.C.Overley, R.M.Sealock, and D.H.Ehlers, Nucl. Phys. A221,
573 (1974).
- Pe64 E.D.Pendlebury, AWRE-O-60/64 (1964).
- Pr69 G.Presser et al., Nucl.Phys. A131, 679(1969).
- Re78 C.Renner, J.A.Harvey, N.W.Hill, G.L.Morgan, and K.Pusk, Bull.
Am. Phys. Soc. 23, 526 (1978).
- Sa82 E.T.Sadowski, H.Knox, D.A.Resler, and R.O.Lane, BAP 27,624(c5)
(1982).
- Sm77 A.B.Smith, P.Guenther, D.Havel, and J.F.Whalen, ANL/NDM-29
(1977).
- Sm82 A.B.Smith, P.T.Guenther, and J.F.Whalen, Nucl. Phys. A373,
305 (1982).
- Wo62 C.Wong, J.D.Anderson, and J.W.McClure, Nucl. Phys. 33,680
(1962).

$n + {}^6\text{Li}$ Total Cross Section

$n + {}^6\text{Li}$ Elastic Cross Section

$n + {}^6\text{Li}$ Elastic Cross Section

${}^6\text{Li}(n,t){}^4\text{He}$ Cross Section

${}^6\text{Li}(n,n'){}^6\text{Li}^*$ Cross Section $E_x=2.180\text{ MeV}$

SUMMARY DOCUMENTATION FOR ^7Li ENDF/B-VI, MAT = 328

P. G. Young

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. ABSTRACT

A new covariance analysis of $n+^7\text{Li}$ cross section data has been completed for Version VI of ENDF/B. The analysis updates our 1981 work for ENDF/B-V.2 to include new data that has become available since that time and to incorporate cross correlations between different experiments. The bulk of the new measured data consists of some 10 new (or newly revised) tritium-production measurements involving about 70 new data points. The new analysis results in only small changes in the previous evaluation of the tritium-production cross section but significantly reduces the magnitudes of uncertainties due to the more extensive and accurate data base that was used.

II. INTRODUCTION

The major interest in ^7Li for fusion energy applications results from its potential use as a breeding material for tritium.¹ In 1981 a major re-analysis of ^7Li data was completed for Revision 2 of ENDF/B-V.² That analysis resulted in a major change in the $^7\text{Li}(n,n't)$ cross section near 14 MeV, namely, the cross section was decreased ~9% relative to the previous ENDF/B-V evaluation. Since that time, a number of new measurements, mainly of tritium-production cross sections, elastic scattering angular distributions, and neutron-emission spectra, have been completed. Consequently, a new evaluation of $n+^7\text{Li}$ cross section and covariance data was performed for Version VI of ENDF/B to reflect the new information in the experimental data base.

III. NUCLEAR DATA EVALUATION

Analysis Description

As was the case with the ENDF/B-V.2 evaluation, covariance analyses have been performed of each of the major $n + ^7\text{Li}$ cross-section types for which experimental data exist. The GLUCS code system⁴ was utilized to determine evaluated energy-dependent cross sections and covariances for each reaction type from inputted experimental cross sections with their associated uncertainties and correlations. In addition to energy-dependent correlations within individual experiments, cross correlations between different measurements from common flux standards and half life in tritium-counting experiments were included. The results of the GLUCS analysis were combined using the ALVIN code,⁵ under the constraint that all partial reactions sum to the total cross section, with full account being taken of all covariances from the GLUCS analysis.

Using a constant 49-point energy grid, independent covariance analyses were carried out with GLUCS for the following four reactions or combinations of reactions:

1. total cross section;
2. elastic plus (n,n_1') cross section to the first excited state of ^7Li ;
3. $(n,n't)$ tritium-production cross section;

4. (n,2n) plus (n,2nd) plus (n,3np) plus (n,d) cross sections.

Reactions (1)-(4) include all the partial reaction and scattering cross sections that must sum to reaction (1), the total cross section. The data adjustment code, ALVIN, was then used to combine the cross sections and covariances from the independent GLUCS analyses, under the constraint that $\sigma_1 = \sigma_2 + \sigma_3 + \sigma_4$. The results on the 49-point energy grid were smoothed, where necessary, and fit with spline curves for the final evaluated results.

In addition to the above combined analysis, the individual ${}^7\text{Li}(n,n')$ cross sections to the first and second excited states of ${}^7\text{Li}$ were obtained from separate GLUCS analyses of the individual reactions. Because the 0.478-MeV first excited state of ${}^7\text{Li}$ is bound, the experimental data base for the ${}^7\text{Li}(n,n_1)$ reaction consists mainly of $(n,n'\gamma)$ measurements. The second excited state at $E_x = 4.63$ MeV is unbound by 2.16 MeV, and direct measurements of inelastic neutrons are available for the (n,n_2) reaction.

To perform the above analyses, it was necessary to obtain covariance matrices for each experimental data measurement. In many cases, sufficient information was available to infer the correlations in the experimental data, and occasionally the correlation matrices were even provided directly by the experimenters. For several measurements, however, it was necessary to make simple generic assumptions regarding the correlations present in different types of experiments. For example, modern total cross-section measurements were generally assumed to have a normalization uncertainty of the order of 0.3-0.5% due to sample thickness and composition uncertainty. Greater normalization uncertainty was assumed for older measurements. The final GLUCS/ALVIN cross sections were not found to be highly sensitive to the exact assumptions made, although it was observed that significant overestimates of correlations can distort results, especially in energy regions where measured data were scarce.

A simple error-doubling procedure was followed for measurements that differed by more than two standard deviations from trial results from GLUCS. That is, if the results from a particular experiment differed from the GLUCS combination of all other experiments such that χ^2/point was greater than 4, then the uncertainties on all the data from that experiment were doubled. Such a procedure was necessary for some 10 experiments out of the 50 used in the analysis. It should be noted that some 7 of the 10 experiments with doubled errors were reported prior to 1965. The uncertainties on more recent measurements were generally found to be more self consistent.

Experimental Data

All available experimental data for which reasonable error estimates were feasible were included in the GLUCS analyses. A total of some 3400 experimental data points were considered, although the initial 3200 total cross section points were averaged down to about 500 points in order to simplify the analysis. The new experimental data on tritium production⁶⁻¹⁴, completed or revised since the previous ENDF/B-V.2 analysis, are summarized in Table I.* Other new experimental data included in the analysis were the elastic cross section results of Chiba et al.,¹² Shen et al.,¹⁵ Alfimenkov et al.,¹⁶ and Drosge et al.,¹⁷ a new 14-MeV (n,2n) data point from the work of Chiba et al., and new results on the (n,n₂) cross section from Chiba et al., Drosge et al., Schmidt et al.,¹⁸ and Dekempeneer and Liskien.¹⁹

The only experimental data available in the energy range 16-20 MeV are the total and (n,n' γ) cross sections. Therefore, in order to permit an accurate separation of the partial cross sections at these energies, an optical-model analysis was performed covering the energy range 10-20 MeV. The elastic angular distribution measurements of Hogue et al.²⁰ and Shen et al.,¹⁵ together with an average of the total cross section measurements²¹ from

10-20 MeV were fit using the SCATOPT spherical optical model code.²² The results were used to compute elastic cross sections from 15-20 MeV for inclusion in the GLUCS/ALVIN analysis.

Evaluation Results

The total cross section that resulted from the analysis is compared in Fig. 1 with white neutron source measurements²¹ between 2 and 18 MeV. The evaluated curve was obtained by passing a spline curve directly through the ALVIN results on the 49-point energy grid. The resulting curve is virtually indistinguishable from our earlier ENDF/B-V.2 evaluation, which is not surprising as the same total cross section data base was used in both analyses.

The (n,n't) cross sections that resulted from the ALVIN analysis were not as smooth as the total cross section, primarily because of the smaller and less consistent experimental data base that went into the (n,n't) analysis, so some smoothing of those results was necessary. The smoothed results are compared in the left half of Fig. 2 to the experimental (n,n't) data⁶⁻¹⁴ that have been obtained since the ENDF/B-V.2 analysis, as well as to the older measurements²³ (right half of the figure) and to the earlier ENDF/B-V.2 analysis² (dashed curves). Clearly the tritium-production cross section from the present analysis differs only slightly from the 1981 evaluation. The new results lie higher than the earlier analysis between 6 and 10 MeV, fall somewhat lower above 15 MeV, and are within ~1% near 14 MeV. It should be noted, however, that the covariance matrix for the (n,n't) reaction is changed substantially. In particular, the standard deviations are significantly reduced because of the additional data in the analysis. A total uncertainty of about $\pm 2.1\%$ is obtained for the 14-15 MeV region as compared to ~4% for ENDF/B-V.2.

The results for the elastic cross section are compared in Fig. 3 to the available experimental data base²³, and to the ENDF/B-V.2 evaluation. The new analysis represents the experimental data quite well and differs only slightly from the earlier evaluation.

Finally, the ${}^7\text{Li}(n,n_1)$ and ${}^7\text{Li}(n,n_2)$ cross sections that result from the independent GLUCS analyses are compared to experimental data and to ENDF/B-V.2 in Figs. 4 and 5, respectively. The new (n,n₁) results are identical with the earlier evaluation because the same experimental data base was used. The new (n,n₂) evaluation lies higher than ENDF/B-V.2 at neutron energies below 10 MeV and falls lower at higher neutron energies, primarily reflecting the influence of the new Dekempeneer and Liskien¹⁹ data and the fact that a covariance analysis was not used for the (n,n₂) reaction in ENDF/B-V.2.

Additional details are included in the ENDF/B-VI File 1 comment section.

IV. REFERENCES

1. E. T. CHENG, "Nuclear Data Needs for Fusion Energy Development," *Fusion Tech.* **8**, 1423 (1985).
2. P. G. YOUNG, "Evaluation of $n+{}^7\text{Li}$ Reactions Using Variance-Covariance Techniques," *Trans. Am. Nucl. Soc.* **39**, 272 (1981); ENDF/B-V, Rev. 2 data file for ${}^7\text{Li}$ (MAT 1397), described in "ENDF/B-V.2 Summary Documentation," Comps., B. A. MAGURNO and P. G. YOUNG, Brookhaven National Laboratory report BNL-NCS-17541 (ENDF-201, 3rd Ed., Sup. 1, January 1985).
3. J. W. DAVIDSON, D. J. DUDZIAK, J. STEPANEK, C. E. HIGGS, and S. PELLONI, "Analysis of the LBM Experiments at LOTUS," *Fusion Technology*, **10**, 940, November 1986.

4. D. M. HETRICK and C. Y. FU, "GLUCS: A Generalized Least-Squares Program for Updating Cross-Section Evaluations with Correlated Data Sets," Oak Ridge National Laboratory report ORNL/TM-7341 (ENDF-303) (1980).
5. D. R. HARRIS, W. A. REUPKE, and W. B. WILSON, "Consistency Among Differential Nuclear Data and Integral Observations: The ALVIN Code for Data Adjustment, for Sensitivity Calculations, and for Identification of Inconsistent Data," Los Alamos Scientific Laboratory report LA-5987 (December 1975).
6. H. LISKIEN, R. WÖLFE, and S. M. QAIM, "Determination of ${}^7\text{Li}(n,n't){}^4\text{He}$ Cross Sections," *Proc. Int. Conf. Nuclear Data for Science and Technology*, Antwerp, 6-10 Sept. 1982 (Ed, K. H. Böckhoff, D. Reidel Pub. Co., Dordrecht, 1983), p. 349.
7. H. MAEKAWA, K. TSUDA, T. IGUCHI, Y. IKEDA, Y. OYAMA, T. FUKUMOTO, Y. SEKI, and T. NAKAMURA, "Measurements of Tritium Production-Rate Distribution in Simulated Blanket Assemblies at the FNS," Japanese Atomic Energy Research Institute report JAERI-M-83-196 (1983).
8. D. L. SMITH, J. W. MEADOWS, M. M. BRETSCHER and S. A. COX, "Cross Section Measurement for the ${}^7\text{Li}(n,n't){}^4\text{He}$ Reaction at 14.74 MeV," Argonne National Laboratory report ANL/NDM-87 (1984).
9. H. MAEKAWA, K. TSUDA, Y. IKEDA, K. OISHI, and T. IGUCHI, "Measurement of ${}^7\text{Li}(n,n'\alpha){}^3\text{H}$ Cross Section between 13.3 and 14.9 MeV," personal communication of results from the FNS at JAERI and from the University of Tokyo (1986).
10. A. TAKAHASHI, K. YUGAMI, K. KOHNO, N. ISHIGAKI, J. YAMAMOTO, and K. SUMITA, "Measurements of Tritium Breeding Ratios in Lithium Slabs Using Rotating Target Neutron Source," *Proc. 13th Symp. Fusion Technology 1984*, Varese, Italy, 24-28 Sept. 1984 (Pergamon Press, 1984), p. 1325.
11. E. GOLDBERG, R. L. BARBER, P. E. BARRY, N. A. BONNER, J. E. FONTANILLA, C. M. GRIFFITH, R. C. HAIGHT, D. R. NETHAWAY, and G. B. HUDSON, "Measurements of ${}^6\text{Li}$ and ${}^7\text{Li}$ Neutron-Induced Tritium Production Cross Sections at 15 MeV," *Nucl. Sci. Eng.* **91**, 173 (1985).
12. S. CHIBA, M. BABA, H. NAKASHIMA, M. ONO, N. YABUTA, S. YUKINORI, and N. HIRAKAWA, "Double-Differential Neutron Emission Cross Sections of ${}^6\text{Li}$ and ${}^7\text{Li}$ at Incident Neutron Energies of 4.2, 5.4, 6.0 and 14.2 MeV," *J. Nucl. Sci. and Tech.* **22**, 771 (1985).
13. M. T. SWINHOE and C. A. UTTLEY, "An Absolute Measurement of the ${}^7\text{Li}(n,n'\alpha t)$ Reaction Cross Section Between 5 and 14 MeV by Tritium Assaying," *Nucl. Sci. Eng.* **89**, 261 (1985).
14. S. M. QAIM and R. WÖLFLE, " ${}^7\text{Li}(n,n't){}^4\text{He}$ Reaction Cross Section via Tritium Counting," *Nucl. Sci. Eng.* **96**, 52 (1986).
15. G. SHEN, S. WEN, T. HUANG, A. LI, and X. BAI, "Measurements of Differential 14.7-MeV Neutron Scattering Cross Sections of Lithium-7 and Beryllium-9," *Nucl. Sci. Engr.* **86**, 184 (1984).

16. V. ALFIMENKOV, S. BORZAKOV, V. VAN-TKHUAM, J. MAREJEV, L. PIKEL'NER, G. RUBIN, A. KHRYKIN, and E. SHARAPOV, *Jadernaja Fizika* **35**, 542 (1982).
17. M. DROSG, P. LISOWSKI, D. DRAKE, R. HARDEKOPF, and M. MUELLNER, "Double Differential Neutron Emission Cross Sections of ^{10}B and ^{11}B at 6, 10 and 14 MeV, and of ^6Li , ^7Li and ^{12}C at 14 MeV," *Rad. Effects* **92**, 145 (1986).
18. D. SCHMIDT, D. SEELIGER, G. N. LOVCHIKOVA, and A. M. TRUFANOV, "Measurement and Status of Neutron Scattering on ^6Li and ^7Li Between 6 and 14 MeV," *Nucl. Sci. Engr.* **96**, 159 (1987).
19. E. DEKEMPENEER, H. LISKIEN, L. MEWISSEN, and F. POORTMANS, "Double-Differential Neutron-Emission Cross Sections for ^7Li and Incident Neutrons Between 1.6 and 13.8 MeV," *Nucl. Sci. Engr.* **97**, 353 (1987).
20. H. H. HOGUE, P. L. VON BEHREN, D. W. GLASGOW, S. G. GLENDENNING, P. W. LISOWSKI, C. E. NELSON, F. O. PURSER, and W. MORROW, "Elastic and Inelastic Scattering of 7- to 14-MeV Neutrons from Lithium-6 and Lithium-7," *Nucl. Sci. Eng.* **69**, 22 (1979).
21. J. A. HARVEY, Oak Ridge National Laboratory, personal communication through the National Nuclear Data Center, Brookhaven National Laboratory (1978); C. A. GOULDING and P. STOLER, Rensselaer Polytechnic Institute, personal communication through the National Nuclear Data Center, Brookhaven National Laboratory (1971).
22. O. BERSILLON, Bruyeres-le-Chatel, France, personal communication to E. Arthur (1980).
23. Experimental data available from the CSISRS compilation by the National Nuclear Data Center, Brookhaven National Laboratory, Upton, New York.

Table 1. Summary of new ${}^7\text{Li}(n,n't)$ cross section measurements since completion of the 1981 ENDF/B-V.2 evaluation.

Reference	No. Points	Energy Range (MeV)	First Author and Laboratory	Covariance Information
6	26	4.99-16.03	Liskien, Geel	Correlations inferred
7	1	14.9	Maekawa, JAERI	—
8	1	14.74	D.L. Smith, ANL	Correlations with 1981 measurements supplied
9a	6	13.31-14.88	Maekawa, FNS(JAERI)	Correlations estimated
9b	6	13.40-14.79	Maekawa, Tokyo Univ.	Correlations estimated
10	12	13.35-14.83	Takahashi, Osaka Univ.	Correlations estimated
11	1	14.94	Goldberg, LLNL	—
12	3	5.40-14.2	Chiba, Tohoku Univ.	Correlations inferred
13	8	4.57-14.1	Swinhoe, Harwell	Revision of '79 measurements & covariances
14	6	7.945-10.48	Qaim, Jülich	Correlations inferred

Figure 1. Neutron total cross section of ${}^7\text{Li}$. The solid curves are from the present covariance analysis; the points are experimental data.²¹

Figure 2. The ${}^7\text{Li}(n,n't)$ cross section. The solid curves are from the present analysis and the dashed curves are ENDF/B-V.2. The experimental data in the right half²³ of the figure were available for the ENDF/V-V.2 analysis; the experimental data in the left half⁶⁻¹⁴ became available after the ENDF/B-V.2 analysis.

Figure 3. Neutron elastic scattering cross section of ${}^7\text{Li}$. The solid curve is from the present analysis; the points are experimental data.²³

Figure 4. Evaluated and measured²³ cross sections for the ${}^7\text{Li}(n,n'\gamma)$ reaction, corresponding to the ${}^7\text{Li}(n,n_1)$ reaction to the 0.478-MeV first excited state of ${}^7\text{Li}$.

Figure 5. Evaluated and measured^{12,18,19,23} cross sections for the ${}^7\text{Li}(n,n_2)$ reaction to the 4.63-MeV second excited state of ${}^7\text{Li}$. The dashed curve represents the ENDF/B-V.2 evaluation.

${}^9_4\text{Be}$

Reference: No Primary Reference
Evaluators: S. T. Perkins, E. F. Plechaty, and R. J. Howerton
Evaluated: January 1986
Material: 425
Content: Neutron transport, Gamma production

File Comments

The ${}^9\text{Be}$ neutron cross sections for ENDF/B-VI were evaluated by S. T. Perkins, E. F. Plechaty, and R. J. Howerton. Lawrence Livermore National Laboratory, Livermore, Ca., Jan. 1986.

General Comments

The neutron energy range covered extends from .00001 eV to 20 MeV. In addition to elastic scattering which of course is everywhere energetically possible, the following reactions have thresholds at energies less than 20 Mev (UCRL 50400, Vol. 9, (1970)):

Reaction	Threshold (MeV)
n,2n	1.85
n,p	14.26
n,np	18.76
n,c	16.29
n,nd	18.55
n,t	11.60
n,nt	19.65
n, α	0.67
n,n α	2.74
n, γ	<i>exoergic</i>

For the (n,np), (n,nd), and (n,nt) reactions, there are no measurements and since the thresholds are sufficiently high, these cross sections are assumed to be negligible. the (n,n α) reaction is a decay mode for the (n,2n) reaction since ${}^5\text{He}$ is unstable, decaying immediately to a neutron and an alpha particle. The inelastic scattering reaction is also a decay mode for the (n,2n) reaction since the ${}^9\text{Be}^*$ recoil nucleus always decays to the final end products of a neutron and two alpha particles. the selection of the cross sections for elastic scattering, (n,2n), (n,p), (n,d), (n,t), (n, α), (n, γ), and the (n,x γ) reaction is discussed below.

Elastic Scattering Cross Section

The free atom cross section is used for all energies below the upper limit of the molecular binding energy (about 10 eV). This is effectively only the nuclear part of the cross section of a stationary target at zero degrees Kelvin. It is strongly emphasized that the numbers are meaningless in the absence of a proper thermal treatment by either the processing code or the neutronics code which uses these numbers. They are likewise meaningless if the materials for which they are to be used are bound in molecules unless molecular binding is taken into account.

The scattering cross section was taken equal to 6.15 barns (Neutron Cross Sections, Vol. 1, Pt. A, Academic Press (1981), S. F. Mughabghab, M. Divadeenam, and N. E. Holden) from 0.00001 eV to 0.01 MeV. From .01 MeV to the (n,α) threshold at 0.67 MeV, the elastic scattering cross section is equal to the total cross section since the (n,γ) cross section is negligible. The cross section was based on data of Refs. 63, 92, 720, 772, and 1002.

Above the (n,α) threshold, the scattering cross section was taken as the difference between the total and the nonelastic, with the nonelastic being equal to the sum of its parts. Up to 2 MeV, the total cross section was based on the previously mentioned data, and those from Refs. 72, 336 and 1642. From 3.2-4.4 MeV, the Argonne total cross section data was used (A. B. Smith, private communication, 1988). From 4.5-10. MeV the Argonne elastic cross section data was used (A. B. Smith, private communication, 1988). Also above 2. Mev, we relied on the results of Ref. 3088 over the 2.7 Mev resonance, and Ref. 750 and the elastic scattering data from Ref. 4113 and 4473 up to 15 MeV. For incident neutron energies from 15 to 20 MeV, the data from Refs. 107, 673, and 682 were used.

Elastic Scattering Angular Distributions (Normalized Probabilities)

Up to 7 MeV, there are experimental differential scattering data in Refs. 121, 151, 231, 296, 500, 571, 945, 1643, 1645, 1646 and 1647. These results were used to determine the normalized probabilities. In Be below the $(n,2n)$ threshold, the total scattering data are equivalent to those for elastic scattering. The change in shape of the angular distribution going through the scattering resonances was taken into account. From 4.5-10.0 MeV the Argonne differential elastic scattering cross sections were used (A. B. Smith, private communication, 1988). From 6 to 15 MeV, the results of Refs. 4113, 4473, 402, 2585, and 3106 were also used. A smooth extrapolation was made to 20 MeV.

The (n,2n) Evaluation

The evaluation for the ^9Be (n,2n) reaction is described in detail in Nucl. Sci. & Eng. 90, 85 (1985), S. T. Perkins, E. F. Plechaty, and R. J. Howerton. It involved using a Monte Carlo technique and comparing the calculated spectra against recent double differential cross section measurements. At each incident neutron energy, 5,000,000 events were sampled, resulting in 10,000,000 secondary neutrons. The (n,2n) reaction was described by the following four events:

where "*" denotes an excited state, and wide level transitions are considered as required.

Total (n,2n) Cross Section

The total (n,2n) cross section was based on the data from Ref. 683, 763, 3320, 4473, 4871, and 5660. At 14 MeV, the weighted mean of the eight values measured between 1958-1963 was also considered. The resulting curve is similar to that used in ENDF/B-V.

(n,2n) Double Differential Spectra

The calculated double differential spectra were compared to 51 measurements between 3.25 and 15.4 MeV: Ref. 4871 (3.25-15.4 MeV), Ref. 4473 (5.9-14.2 MeV), and Proc. Int. Conf. Nucl. Data for Sci. and Tech., Antwerp, CONF-820906, p. 360 (1982), A. Takahashi et. al.; see also Oktavian Report A-83-01, Osaka Univ., Japan (1983), (14 MeV). Comparisons were made after weighing the calculated results with the experimental resolution function. The final calculated spectra were then smoothed and thinned. This yielded double differential spectra for both the secondary neutrons and the secondary alpha particles.

(n,p) Cross Section

This cross section is entirely for the (n,p₀) reaction and was based on the data from Refs. 913 and 3492. It was smoothly extrapolated from 15.5 to 20 MeV. The proton was assumed isotropic in its center of mass system.

(n,d) Cross Section

This cross section is entirely for the (n,d_0) reaction and was based on the data between 15.5 and 19 MeV reported in Ref. 4841. The deuteron was assumed isotropic in its center of mass system.

(n,t) Cross Section

The cross section for the (n,t) reaction is based on a (n,t_0) and a (n,t_1) component, which proceed through the 0.0 and the 0.477 MeV levels in ${}^7\text{Li}$ respectively. The total (n,t) cross section near 14 MeV was determined from Ref. 496, 3996 and 5722.

The (n,t_1) cross section was based on the work in Ref. 5529 and smoothly extrapolated from 15 to 20 MeV. The resonance near threshold is consistent with structure in the compound system ${}^{10}\text{Be}$. Both tritons were assumed isotropic in their center of mass systems.

(n, α) Cross Section

The cross section was based on the measurements reported in Ref. 160, 494 and 733 up to neutron energies of 8.6 MeV. It was then tied into the four values given at 14 MeV by Refs. 86, 650, 2367, and Nucl. Phys. A227, 330 (1978), J. P. Perroud and C. H. Sellem. It was then extrapolated to 20 MeV. Note that the measured cross section is to ${}^6\text{He}$ (0.0 MeV) which β decays to ${}^6\text{Li}$; higher states in ${}^6\text{He}$ decay to $\alpha + 2n$. The alpha particle angular distribution was taken as isotropic in the center of mass system at threshold. At 14.1 MeV the data from the measurement of Ref. 4351 was used; this distribution was also used at 20 MeV.

(n, γ) Cross Section

The cross section was assumed to be $1/v$ below 100 eV with a 2200 m/sec cross section of 8.6 mb. This is in agreement with the CSEWG Dosimetry and Activation files. It was then extrapolated linearly on a log-log basis to 0.1 mb at 1 keV and then held constant at this value up to 20 MeV.

(n,x γ) Cross Section

γ -rays in ${}^9\text{Be}$ are produced by the (n,γ) and the (n,t) reactions. At thermal energies, Ref. 2415 quotes γ -ray energies of 0.8535, 2.59, 3.368, 3.444, 5.958 and 6.81 MeV

and the corresponding $(n, x\gamma)$ cross sections. The photon spectrum from this work was used at all neutron energies. the multiplicity varied with neutron energy as $M(E) = M(0) \times (E_{cm} + Q) / Q$, where $M(0)$ is the multiplicity at thermal and Q is the (n, γ) Q value. This combination of spectra and multiplicity conserves energy. There are no doubt other levels in ^{10}Be with energies greater than 6.81 MeV excited as the incident neutron energy is increased. The energies of the higher states have not yet been determined so we use this artifact to conserve energy. The component resulting from the (n, t) reaction is equal to the (n, t_1) cross section since its multiplicity is unity.

References

(Reference numbers quoted here refer to data in the LLNL experimental neutron cross section library. See UCRL-50400, Vol. 2 and 3 for comments and indexes to the data).

- 63. Phys. Rev. 80, 1011 (1950) C. K. Bockelman.
- 72. Phys. Rev. 84, 69 (1951) C. K. Bockelman, D. W. Miller, R. K. Adair, H. H. Barschall.
- 86. Phys. Rev. 89, 80 (1953) M. E. Battat, F. L. Ribe.
- 92. Private Communication (1954) P. H. Stelson.
- 107. Phys. Rev. 94, 651 (1954) C. F. Cook, T. W. Bonner.
- 121. Phys. Rev. 98, 669 (1955) J. D. Seagrave, R. L. Henkel.
- 151. Phys. Rev. 104, 1319 (1956) J. R. Beyster, M. Walt, E. W. Salmi.
- 160. Phys. Rev. 106, 1252 (1957) P. H. Stelson, E. C. Campbell.
- 231. ANL-5567 (1956) A. Langsdorf, Jr., R. O. Lane, J. E. Monahan, see also, ANL-5554 p. 22 (1956), Phys. Rev. 107, 1077 (1957), ANL-5567 (Rev.) (1961).
- 296. Phys. Rev. 98, 677 (1955) M. Walt, J. R. Beyster.
- 336. Proc. Phys. Soc. (London) 64, 388 (1951) G. H. Stafford.
- 402. Phys. Rev. 110, 1439 (1959) M. P. Nakada, J. D. Anderson, C. C. Gardner, C. Wong.
- 494. Doklady Akad. Nauk S.S.S.R. 119, 914 (1958) S. S. Vasilev, V. V. Kumarov, A. M. Popova, see also, J. Exptl. Theoret. Phys. (USSR) 33, 527 (1957).

496. Phys. Rev. 112, 1264 (1958) M. E. Wyman, E. M. Fryer, M. M. Thorpe.
500. Phys. Rev. 114, 1584 (1959) J. B. Marion, J. S. Levin, L. Cranberg.
571. ANL-6172 (1960) R. O. Lane, A. S. Langsdorf, Jr., J. E. Monahan, A. J. Elwyn, See also, Ann. Phys. (N.Y. 12, 135 (1961).
650. J. Exptl. Theoret. Phys. (USSR), 40, 1244 (1961), S. A. Myachkova and V. P. Perehygin.
673. Phys. Rev. 120, 521 (1960) J. M. Peterson, A. Bratenahl, J. P. Stoering.
682. Phys. Rev. 123, 209 (1961) D. B. Fossan, R. L. Walter, W. E. Wilson, H. H. Barschall.
720. Private Communication (1962) E. G. Bilpuch, J. A. Farrell, G. C. Kyker, Jr., P. B. Parks, H. Newson.
733. Nucl. Phys. 23, 122 (1961) R. Bass, T. W. Bonner, H. P. Haenni.
750. Private Communication (1967) D. G. Foster, Jr., D. W. Glasgow, See also, HW-73116 (1962), HW-77311 (1963), Phys. Rev. C, 3, 576 (1971), Phys. Rev. C, 3, 604 (1971).
763. Nuclear Phys. 129, 305 (1969) M. Holmberg, J. Hansen, See also, IAEA Conference on Nuclear Data, Paris, Paper CN-23/18 (1966).
772. Compt. Rend. 255, 277 (1962) A. Perrin, G. Surget, C. Thibault, F. Verriere.
913. Phys. Rev. 132, 328 (1963) D. E. Alburger.
945. ORNL-2610 p. 14 (1958) H. C. Cohn, J. L. Fowler, See also, Bull. Amer. Phys. Soc. 3, 305 (1958).
1002. Private Communication (1954) C. T. Hibdon, A. Langsdorf.
1642. Bull. Amer. Phys. Soc. 4, 385 (1959) J. L. Fowler, H. C. Cohn.
1643. Doklady Akad. Nauk S.S.S.R. 158, 574 (1964) G. V. Gorlov, N. S. Lebedeva, V. M. Morozov, See also, Yad. Fiz. 5, 910 (1967).
1645. Private Communication (1960) J. S. Levin, L. Cranberg, See also, WASH-1028 p. 26 (1960), WASH-1029 p. 44 (1960).
1646. Private Communication (1961) D. D. Phillips.
1647. Phys. Rev. 133, 409 (1964) R. O. Lane, A. J. Elwyn, A. Langsdorf, Jr.
2367. Nuclear Phys. 96, 476 (1967) G. Paic, D. Rendic, P. Tomas.

2415. GA-10248 (DASA-2570) (1970) N. C. Rasmussen, V. J. Orphan, T. L. Harper, J. Cunningham, S. A. Ali.
2585. Private Communication (1966) R. Bouchez, See also, IAEA Conference on Nuclear Data, Paris, Paper CN-23/75 (1966).
3088. Conference on Neutron Cross Section Technology, Washington D. C., p. 851 (1968) C. H. Johnson, F. X. Haas, J. L. Fowler, F. D. Martin, R. L. Kernell, H. O. Cohn.
3106. NP-17794 (1968) J. Roturier, See also, Compt. Rend. 260, 4491 (1965), BNL-400, 3rd ed. (1970).
3320. Atomkernenergie 20, 309 (1972) M. Bloser.
3492. Nucl. Sci. and Eng. 54, 190 (1974), R. H. Augustson and H. O. Menlove.
3996. J. Inorg. Nucl. Chem. 37, 1583, (1975), T. Biro, et. al.
4113. Nucl. Sci. & Eng. 68, 38 (1978), H. H. Hogue et. al.
4351. Nucl. Phys. 257, 397 (1976), W. Smolec et. al.
4473. Nucl. Sci. & Eng. 63, 401 (1977), D. M. Drake, G. F. Auchampaugh, E. D. Arthur, C. E. Ragan, and P. G. Young.
4841. Z. Naturforschung 25, 1460 (1970), W. Scobel and M. Bormann.
5529. Nucl. Sci. & Eng. 61, 267 (1976), F. S. Dietrich, L. F. Hansen and R. P. Koopman.
5722. Nucl. Data. for Sci. and Tech., Antwerp, p. 368 (1982), Z. T. Body et. al.

10
5 B

Reference: No Primary Reference
Evaluators: G. M. Hale and P. G. Young
Evaluated: November 1989
Material: 525
Content: Standard, Neutron transport, Gamma production

ENDF/VI EVALUATION
G. M. Hale and P. G. Young

MAJOR CHANGES FROM VERSION V OF ENDF/B ARE:

1. Inclusion of the ENDF/B-VI standard (n,alpha) and (n,alpha1) results from the simultaneous standards analysis (Ca85) over the standard energy range thermal to 100 keV.
2. Replacement of all major cross sections and elastic angular distributions from 10E-5 eV to 1 MeV with results from the R-matrix analysis performed in conjunction with the simultaneous standards analysis.
3. Replaced the total cross section 1-20 MeV with results from a covariance analysis of available data.
4. Revised elastic and inelastic cross sections for low-lying levels incorporating new elastic, inelastic, and (n,xgamma) experimental data. We attempted to better reconcile the inelastic and gamma ray data.
5. Refit all elastic angular distributions from 1-20 MeV with Legendre expansions and incorporated results from new measurements.
6. Fit inelastic neutron angular distributions for first 5 excited states of B-10 with Legendre expansions.
7. Incorporated new (n,t2alpha) cross section data into MT113 and adjusted (n,alpha) cross sections above standard region for better consistency with data as well as other cross sections (esp. total and elastic) determined by data.

*****Note that covariance data will be added at a later date.

MF=2 ----- Resonance parameters -----

MT=151 Effective scattering radius = 4.129038E-13 cm

MF=3 ----- Smooth cross sections -----

The 2200 m/s cross sections are as follows,

MT=1 sigma = 3842.146 barns
 MT=2 sigma = 2.142435 barns
 MT=102 sigma = 0.5 barns
 MT=103 sigma = 0.000566 barns
 MT=107 sigma = 3839.496 barns
 MT=113 sigma = 0.0069993 barns
 MT=600 sigma = 0.000566 barns
 MT=800 sigma = 241.2677 barns
 MT=801 sigma = 3598.228 barns

MT=1 Total cross section

- 0 to 1 mev, calculated from R-matrix parameters obtained from simultaneous standards analysis (Ca85) used to obtain the ENDF/B-VI standard cross sections.
- 1 to 20 mev, covariance analysis of measurements of Di67, Ts62, Fo61, Co52, Au79, and Co54, constrained to match R-matrix fit at 1 mev. GLUCS covariance analysis code (He80) was used in the calculations.

MT=2 Elastic scattering cross section

- 0 to 1 mev, calculated from the R-matrix parameters described for MT=1. Experimental elastic scattering data included in the fit are those of As70 and La71.
- 1 to 6 mev, smooth curve through measurements of La71, Po70, Sa88, and Ho69, constrained to be consistent with total and reaction cross section measurements.
- 6 to 14 mev, smooth curve through measurements of Ho69, Co69, Te62, Va70, Va65, Sa88, and Gl82. Note that the data of Sa88 above 9 MeV were discounted.
- 14 to 20 mev, optical model extrapolation from 14 MeV data.

MT=4 Inelastic cross section

thres.to 20 mev, sum of MT=51-85

MT=51-61 Inelastic cross sections to discrete states

MT=51 Q=-0.717 MeV	MT=55 Q=-4.774 MeV	MT=59 Q=-5.923 MeV
52 -1.740	56 -5.114	60 -6.029
53 -2.154	57 -5.166	61 -6.133
54 -3.585	58 -5.183	

thres.to 20 MeV, based on (n,nprime) measurements of Po70, Co69, Ho69, Va70, Sa88, and Gl82, and the (n,xgamma) measurements of Da56, Da60, Ne70, and Di88, using a gamma-ray decay scheme from analysis of Aj88. Hauser-Feshbach

calculations were used to estimate shapes and relative magnitudes where experimental data were lacking.

MT=62-85 Inelastic cross sections to groups of levels in 0.5-MeV wide bands centered about the Q-values given below (used in lieu of MT=91 and File 5)

MT=62 Q=-6.5 MeV	MT=70 Q=-10.5 MeV	MT=78 Q=-14.5 MeV
63 -7.0	71 -11.0	79 -15.0
64 -7.5	72 -11.5	80 -15.5
65 -8.0	73 -12.0	81 -16.0
66 -8.5	74 -12.5	82 -16.5
67 -9.0	75 -13.0	83 -17.0
68 -9.5	76 -13.5	84 -17.5
69 -10.0	77 -14.0	85 -18.0

thres. to 20 mev, integrated cross section obtained by subtracting the sum of MT=2,51-61,103,104,107, and 113 from MT=1. Cross section distributed among the bands with an evaporation model using a nuclear temperature given by $T=0.9728*\sqrt{EN}$ in MeV, taken from Ir67.

MT=102 (n,gamma) cross section

0 to 1 mev, assumed $1/v$ dependence with thermal value of 0.5 barn.

1 to 20 MeV, assumed negligible, set equal to zero.

MT=103 (n,p) cross section

thres. to 20 MeV, sum of MT=600-605.

MT=104 (n,d) cross section

thres. to 20 MeV, based on Be9(d,n)B11 measurements of Si65 and Ba60, and the (n,d) measurement of Va65.

MT=107 (n,alpha) cross section

0 to 20 MeV, sum of MT=800,801.

MT=113 (n,t2alpha) cross section

0 to 2.3 MeV, based on a single-level fit to the resonance measured at 2 MeV by Da61, assuming $l=0$ incoming neutrons and $l=2$ outgoing tritons. The thermal measurement (7 ± 2 mb) of Ka87 was included in the analysis.

2.3 to 20 MeV, smooth curve through measurements of Fr56, Wy58, Ga88, following general shape of Da61 measurement from 4 to 9 MeV. We assumed that the experimental data of Ga88 supercedes reference Ga85.

MT=600-605 (n,p) cross section to discrete levels from

0 to 20 MeV, crudely estimated from the calculations of Po70 and the (n,xgamma) measurements of Ne70. Cross

section for MT=600 assumed similar to MT=113 below
1 MeV. Gamma-ray decay scheme for Be-10 from Aj88.

MT=800 (n,alpha0) cross section

- 0 to 1 MeV, calculated from the R-matrix parameters described for MT=1. Experimental (n,alpha0) data input to the fit were those of Ma68 and Da61. In addition, the angular distributions of Va72 for the inverse reaction were included in the analysis.
- 1 to 20 MeV, based on Da61 measurements, with smooth extrapolation from 8 to 20 MeV using 14-MeV data of An69. The Da61 data above approximately 2 MeV were renormalized by a factor of approximately 1.4. Note that some of the structure seen in Da61 was expanded to give consistent nonelastic, elastic, and total cross sections when compared with experimental data.

MT=801 (n,alpha1) cross section

- 0 to 1 MeV, calculated from the R-matrix parameters described for MT=1. Experimental (n,alpha1) data included in the fit are those of SC76. In addition, the absolute differential cross-section measurements of Se76 were included in the analysis.
- 1 to 20 MeV, smooth curve through measurements of Da61 and Ne70, with smooth extrapolation from 15 to 20 MeV. The Da61 data above approximately 2 MeV were renormalized by a factor of approximately 1.4. Note that some of the structure seen in Da61 was expanded to give consistent nonelastic, elastic, and total cross sections when compared with experimental data.

MF=4----- Neutron angular distributions -----

MT=2 Elastic angular distributions

- 0 to 1 MeV, calculated from the R-matrix parameters described for MF=1, MT=1. Experimental angular distributions input to the fit for both the elastic scattering cross section and polarization were obtained from available measurements.
- 1 to 14 MeV, smoothed representation of legendre coefficients derived from the measurements of La71, Ha73, Po70, Ho69, Co69, Va69, Va65, Sa88, Gl82, constrained to match the R-matrix calculations at En=1 MeV.
- 14 to 20 MeV, optical model extrapolation of 14-MeV data.

MT=51 Inelastic angular distribution to first level

thres. to 12 mev, fit Legendre expansions to exp. data of

Po70, Gl82, and Sa88.
 12 - 20 MeV, assumed similar distribution as 12 MeV.

MT=52-55 Inelastic angular distribution to first level
 thres. to 12 mev, fit Legendre expansions to exp. data of
 Sa88.
 12 - 20 MeV, assumed similar distribution as 12 MeV.

MT=56-85 Inelastic angular distributions
 thres. to 20 mev, assumed isotropic in center of mass.

MF=12 ----- Gamma ray multiplicities -----

MT=102 Capture gamma rays
 0 to 20 MeV, capture spectra and transition probabilities
 derived from the thermal data of Th67, after slight
 changes in the probabilities and renormalization to the
 energy levels of Aj75. The LP flag is used to conserve
 energy and to reduce significantly the amount of data
 required in the file. Except for the modification due
 to the LP flag, the thermal spectrum is used over the
 entire energy range.

MT=801 0.4776 MeV photon from the (n,alpha1) reaction
 0 to 20 MeV, multiplicity of 1.0 at all energies.

MF=13 ----- Gamma-ray production cross sections -----

MT=4 (n,ngamma) cross section
 thres. to 20 MeV, obtained from MT=51-60 using B-10 decay
 scheme obtained from Aj88.

MT=103 (n,pgamma) cross sections
 thres. to 20 MeV, obtained from MT=601-605 using Be-10
 decay scheme deduced from AJ88.

MF=14 ----- Gamma ray angular distributions -----

MT=4 (n,ngamma) angular distributions
 thres. to 20 MeV, assumed isotropic.

MT=102 (n,gamma) angular distributions
 0 to 20 MeV, assumed isotropic.

MT=103 (n,pgamma) angular distributions

thres. to 20 MeV, assumed isotropic.

MT=801 (n,alpha1/gamma) angular distribution
0 to 20 Mev, assumed isotropic.

----- References -----

- Aj75 F. Ajzenberg-Selove, Nucl. Phys. A248,6 (1975).
Aj88 F. Ajzenberg-Selove, Nucl. Phys. A490,1 (1988).
An69 B. Antolkovic, Nuc.Phys.A139, 10 (1969).
As70 A. Asami and M.C. Moxon, J.Nucl.Energy 24,85 (1970).
Au79 G.Auchampaugh et al., Nucl.Sci.Eng.69,30(1979).
Ba60 R.Bardes and G.E. Owen, Phys.Rev.120,1369 (1960).
Be56 R.L. Becker and H.H. Barschall, Phys.Rev.102,1384 (1956).
Bo51 C.K.Bockelman et al., Phys.Rev. 84,69 (1951).
Bo69 D.Bogart and L.L.Nichols, Nucl.Phys.A125,463 (1969).
Ca85 A.Carlson et al., Nuc.Data for Basic & Applied Science,
Santa Fe, NM (1985) p.1429.
Co52 J.H.Coon et al., Phys.Rev. 88,562 (1952).
Co54 C.F.Cook and T.W. Bonner,Phys.Rev. 94,651 (1954).
Co67 S.A. Cox and F.R. Pontet, J.Nucl.Energy 21,271 (1967).
Co69 J.A. Cookson and J.G.Locke,Nucl.Phys.A146,417(1970).
Co73 M.S. Coates et al., Priv. Comm. to L.Stewart (1973).
Da56 R.B.Day,Phys.Rev.102,767 (1956).
Da60 R.B. Day and M.Walt,Phys.Rev.117,1330 (1960).
Da61 E.A. Davis et al., Nucl.Phys.27,448 (1961).
Di67 K.M. Diment, AERE-R-5224 (1967).
Di88 J.K.Dickens, Proc.Conf. on Nuc.Data for Sci.& Tech.,Mito,
Japan (1988) p.213.
Fo61 D.M. Fossan et al., Phys.Rev. 123,209 (1961).
Fr56 G.M. Frye and J.H. Gammel,Phys.Rev. 103,328 (1956).
Gl82 S.Glendinning, Nuc.Sci.Eng.80,256(1982).
Ha73 S.L.Hausladen, Thesis, Ohio Univ. C00-1717-5 (1973).
He80 D.Hetrick & C.Y.Fu, ORNL/TM-7341 (1980).
Hy69 M.Hyakutake, EANDC(J)-13 (1969) p.29.
Ho69 J.C. Hopkins, Priv. Comm. LASL (1969).
Ir67 D.C.Irving, ORNL-TM-1872 (1967).
Ka87 R.Kavanagh & R.Marcley, Phys.Rev.C36, 1194 (1987).
La71 R.C. Lane et al., Phys.Rev.C4,380 (1971).
Ma68 R.L.Macklin and J.H.Gibbons,Phys.Rev.165,1147 (1968).
Mo66 F.P.Mooring et al.,Nucl.Phys.82,16 (1966).
Ne54 N.G.Nereson,LA-1655 (1954).
Ne70 D.C.Nellis et al., Phys.Rev. C1,847 (1970).
Po70 D.Porter et al., AWRE O 45/70 (1970).
Qa85 S.Qaim et al., Santa Fe Conf. (1985)p.97.
Qa88 S.Qaim et al., Mito Conf. (1988) p.225.
Sa88 E.T. Sadowski, Ph.D thesis, Ohio U., (Nov.,1988).

- Sc76 R.A. Schrack et al., Proc. Icinn(Erda-Conf-760715-p2),1345 (1976).
- Se76 R.M. Sealock and J.C. Overley, Phys.Rev.C13,2149 (1976).
- Si65 R.H.Siemssen et al., Nucl.Phys.69,209 (1965).
- Sp73 R.R. Spencer et al., EANDC(E)147,A1 (1973).
- Te62 K.Tesch, Nucl.Phys.37,412 (1962).
- Th67 G.E. Thomas et al., Nucl.Instr.Meth.56,325 (1967).
- Ts63 K.Tsukada and O.Tanaka,J Phys.Soc.Japan 18,610 (1963).
- Va65 V.Valkovic et al., Phys.Rev. 139,331 (1965).
- Va70 B.Vaucher et al.,Helv.Phys.Acta 43,237 (1970).
- Va72 L.van der Zwan and K.W.Geiger, Nucl.Phys. A180,615 (1972).
- Wi55 H.B. Willard et al., Phys.Rev. 98,669(1955).
- Wy58 M.E. Wyman et al., Phys.Rev.112,1264 (1958).

$n + {}^{10}\text{B}$ Total Cross Section

n + B-10 TOTAL CROSS SECTION

n + B-10 TOTAL CROSS SECTION

n + B-10 ELASTIC CROSS SECTION

n + B-10 ELASTIC CROSS SECTION

B10(n,n')B10* CROSS SECTION, 0.717-MeV Level

$^{10}\text{B}(n,\alpha_0)^7\text{Li}$ Cross Section

B10(n,alpha0)Li7 CROSS SECTION

$^{10}\text{B}(n, \alpha_1)^7\text{Li}$ Cross Section, $E_x=0.478$ MeV

B10(n,alpha1)Li7 CROSS SECTION, Ex=0.478 MeV

SUMMARY DOCUMENTATION FOR ^{11}B ENDF/B-VI, MAT = 528

P. G. Young

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

The fusion energy interest in ^{11}B data results from its potential presence in boron carbide shield walls.¹ The existing ENDF/B-V data file² is actually a pre-1970 United Kingdom evaluation that was converted to ENDF/B format in 1971. A great deal of improved experimental data has become available since that time, notably neutron total cross section, elastic and inelastic scattering angular distributions, and gamma-ray production cross sections and energy distributions. The thrust of the ENDF/B-VI evaluation is to incorporate the new total cross section and elastic/inelastic angular distributions data into the evaluation. Because measurements are lacking at higher energies, optical model and Hauser-Feshbach statistical theory calculations are used to supplement the experimental data base above $E_n = 10$ MeV.

The new evaluation covers the neutron energy range from 10^{-5} eV to 20 MeV. The main new source of neutron total cross-section data for ^{11}B is the 1979 measurements of Auchampaugh et al.,³ which were used in the evaluation from $E_n = 1$ to 14 MeV. At lower energies, the evaluated total cross section is based on the 1970 Lane⁴ measurement and, to a lesser extent, on the 1966 Mooring⁵ results. At higher energies, the 1954 measurements of Cook⁶ are the only data available and were used with optical model calculations to extend the evaluation to 20 MeV.

Figure 1 compares the evaluated total cross section to ENDF/B-V² (dashed curve) and to the available experimental data. As is evident, large discrepancies exist between experimental data represented by the present evaluation and ENDF/B-V, with differences of 40% occurring near 1 and 2.3 MeV and a systematic 8-10% in the range 8-17 MeV. These differences are also evident in the integrated elastic cross section for these neutron energy ranges, with ENDF/B-V lying approximately 30% lower than Version VI near 14 MeV.

The evaluation of the elastic and discrete inelastic cross sections to the first few excited states was done in concert, under the constraint that all cross sections sum to the relatively well-determined total cross section. Integrated cross sections as well as angular distributions were determined for elastic and inelastic scattering by fitting the measured angular distributions with Legendre expansions. The recent measurements of White et al.,⁷ Koehler et al.,⁸ and Glendinning et al.⁹ form the main basis for the evaluation in the MeV region. The evaluated inelastic cross section to the 2.14-MeV state in ^{11}B is shown in Fig. 2, together with ENDF/B-V and representative experimental data. Clearly, there is little resemblance between ENDF/B-V and the experimental data used for the ENDF/B-VI evaluation. There is generally reasonable agreement among the recent scattering measurements.

Additional details are included in the ENDF/B File 1 comment section, which is attached following the figures.

II. REFERENCES

1. E. T. CHENG, "Nuclear Data Needs for Fusion Energy Development," *Fusion Tech.* **8**, 1423 (1985).
2. C. COWAN, ENDF/B-V data file for ^{11}B (MAT 1160), described in "ENDF/B Summary Documentation," R. KINSEY, Comp., Brookhaven National Laboratory report BNL-NCS-17541 (ENDF-201), 1979 (available from the National Nuclear Data Center, Brookhaven National Laboratory, Upton, N.Y.).
3. G. F. AUCHAMPAUGH, S. PLATTARD, and N. HILL, "Neutron Total Cross Section Measurements of ^9Be , $^{10,11}\text{B}$, and ^{12}C from 1.0 to 14 MeV Using the $^9\text{Be}(d,n)^{10}\text{B}$ Reaction as a "White" Neutron Source," *Nucl. Sci. Eng.* **69**, 30 (1979).
4. R. O. LANE, C. E. NELSON, J. L. ADAMS, J. E. MONAHAN, A. J. ELWYN, F. P. MOORING, and A. LANGSDORF, JR., "States in ^{12}B Observed in the Scattering of Neutrons by ^{11}B ," *Phys. Rev. C* **2**, 2097 (1970).
5. F. P. MOORING, J. E. MONAHAN, and C. M. HUDDLESTON, "Neutron Cross Sections of the Boron Isotopes for Energies Between 10 and 500 keV," *Nucl. Phys.* **82** 16 (1966).
6. C. F. COOK and T. W. BONNER, "Scattering of Fast Neutrons in Light Elements," *Phys. Rev.* **94**, 651 (1954).
7. R. M. WHITE, R. O. LANE, H. D. KNOX, and J. COX, "States in ^{12}B from Measurement and R-Matrix Analysis of $\sigma(\theta)$ for $^{11}\text{B}(n,n)^{11}\text{B}$," *Nucl. Phys. A* **340**, 13 (1980).
8. P. E. KOEHLER, H. D. KNOX, D. A. RESLER, and R. O. LANE, "Structure of ^{12}B from Measurement and R-Matrix Analysis of $\sigma(\theta)$ for $^{11}\text{B}(n,n)^{11}\text{B}$ and $^{11}\text{B}(n,n')^{11}\text{B}^*(2.12\text{ MeV})$, and Shell Model Calculations," *Nucl. Phys. A* **394**, 221 (1983).
9. S. G. GLENDINNING, S. EL-KADI, C. E. NELSON, R. S. PEDRONI, F. O. PURSER, R. L. WALTER, A. G. BEYERLE, C. R. GOULD, L. W. SEAGONDOLLAR, and P. TLAMBIDURAI, "Elastic and Inelastic Cross Sections for Boron-10 and Boron-11," *Nucl. Sci. Eng.* **80**, 256 (1982).

$n + {}^{11}\text{B}$ Total Cross Section

Figure 1. Comparison of evaluated and experimental values of the neutron total cross section of ${}^{11}\text{B}$. The solid curve represents the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V,¹⁷ and the points are experimental data as indicated.

$n + {}^{11}\text{B}$ Elastic Cross Section

Figure 2. Comparison of evaluated and experimental values of the elastic scattering cross section of ${}^{11}\text{B}$. See caption of Fig. 1 for details of curves and symbols.

$^{11}\text{B}(n,n')^{11*}$ 4.46-MeV State

$^{11}\text{B}(n,n')^{11*}$ 2.14-MeV State

Figure 3. The $^{11}\text{B}(n,n')$ cross section to the 2.14-MeV first excited state and to the 4.46-MeV second excited state of ^{11}B . See caption of Fig. 1 for details of curves and symbols.

¹¹₅B

Reference: No Primary Reference
Evaluators: P. G. Young
Evaluated: May 1989
Material: 528
Content: Neutron transport, Gamma production

***** GENERAL DESCRIPTION *****

This evaluation is a synthesis of experimental results and theoretical calculations using the Hauser-Feshbach, statistical theory code GNASH (Ar88, Yo77). It replaces a pre-1970 U.K. evaluation that was adapted for ENDF/B in 1971 (Co79). Major emphasis was placed on experimental data where possible, usually using the calculations to estimate shapes and exp. data to normalize the calculations. Spherical optical model calculations were used to obtain particle transmission coefficients, using global potentials for protons and alphas, and the work of Dave et al.(Da83) for the neutron potential. Gilbert-Cameron level density parameters were used and preequilibrium corrections from an exciton model were included. In general, the calculations were required for the gamma and particle emission spectra from charged-particle reactions, as well as for inelastic neutron and gamma emission above the third excited state of B-11, and for all states above an incident neutron energy of about 8 MeV.

***** MF=2 Resonance parameters *****

MT=151 Scattering radius only.

***** MF=3 Thermal cross sections*****

The 1981 evaluation by Mughabghab (Mu81) was used for the 2200 m/s cross sections, as follows:

MT = 1 sigma = 4.8455 b
MT = 2 sigma = 4.8400 b
MT =102 sigma = 0.0055 b

***** MF=3 Smooth neutron cross sections *****

MT= 1 Total cross section. At low energies, evaluation by Mu81a of thermal data used. At higher energies exp. data of Mo66, La70, Ca73, Au79, Co52 & Co54 were used. The high resolution data of Au79 were emphasized. Optical model

calculations with the Da83 potential used to define shape above 14 MeV.

- MT= 2 Elastic cross section. Based on exp. data of La70, Wi55, Po70, Ne73, Wh80, Ko82, Ho69, Gl82, Co70 & Al69. Optical model calculations with potential of Da83 used with renormalization above 14 MeV. Thermal eval. of Mu81a used at low energies.
- MT= 4 (n,nprime)gamma+(n,nprime)continuum. sum of MT=51-60,91.
- MT= 16 (n,2n) cross section. Based entirely on GNASH analysis.
- MT= 22 (n,na) cross section. Based entirely on GNASH analysis.
- MT= 28 (n,np) cross section. Based entirely on GNASH analysis.
- MT=51-54 (n,n'gamma) cross sections. Based on exp. data of Au86, Gl82, Ho69, Po70, Co70, Ko82, Ba85. The Au86 data shapes were used to establish low energy behavior.
- MT=55-60 (n,n'gamma) cross sections. Shapes of excitation curves calculated with GNASH. Absolute magnitudes obtained by renormalizing such that sum of all partials gave reaction, elastic, total x/s consistent with avail. data.
- MT= 91 (n,n'continuum) cross section. Based on GNASH calculation entirely.
- MT=102 (n,gamma) cross section. Adopted from ENDF/B-V.1 except thermal cross section of Mu81a used.
- MT=103 (n,p) cross section. Based on GNASH calculation and exp. data of Sc70a, Fl67, St65, etc.
- MT=105 (n,t) cross section. Similar to ENDF/B-V data, with a smooth curve passing thru exp. data of WY58 at 14.1 MeV.
- MT=107 (n,alpha) cross section. Based on GNASH calculation and exp. data (An79, Sc70b, Ar56).

***** MF=4 Neutron angular distributions *****

- MT= 2 Legendre coefficients obtained by drawing smooth curve through fitted coefficients from measurements listed under MF3/MT2. Data of La60, Gl79, Wh80, Ne73, Ho69, Hy74 emphasized. Optical model calculations used above 14 MeV.
- MT=51-54 Legendre coefficients obtained by fitting exp. data, especially Al69, Hy74, Po70, Ho69, Gl79 & Co69. Smooth curves then passed through fitted coefficients.
- MT=55-60 Isotropy assumed.

***** MF=6 Energy-angle correlated distributions *****

- MT= 16 Neutron and photon distributions are given based on GNASH calculations described above to obtain spectra and multiplicities. Kalbach-Mann (Ka81) systematics (KMS) used for neutron angular distributions. Photon distributions taken as isotropic.
- MT= 22 Neutron and alpha distributions are based on KMS.

Photon distributions assumed isotropic. Multiplicities and spectra based on GNASH calculations.

MT= 28 Neutron and proton distributions are based on KMS. Photon distributions assumed isotropic. Multiplicities and spectra based on GNASH calculations.

MT= 91 Neutron distributions based on KMS and isotropic photon distributions given. Multiplicities and spectra obtained from GNASH analysis.

MT=103 Proton distributions based on KMS and isotropic photon distributions given. Multiplicities and spectra obtained from GNASH analysis.

MT=107 Alpha distributions based on KMS and isotropic photon distributions given. Multiplicities and spectra obtained from GNASH analysis.

***** MF=12 Photon multiplicities *****

MT=102 Radiative capture photon yields obtained from theoretical calculations (Mu81) based on Lane-Lynn theory of direct capture.

***** MF=13 Photon cross sections *****

MT= 4 Gamma ray production cross sections from inelastic scat. Obtained using discrete data (MF=3, MT=51-60) and photon branching ratios (Aj85).

***** MF=14 Photon angular distributions *****

MT= 51 Isotropy assumed at all energies.
 MT=102 Isotropy assumed for all gammas at all energies.

***** MF=33 Neutron cross section covariances *****

To be provided in the future.

***** References *****

Aj85 F.Ajzenberg-Selove, N.P.A433,1(1985).
 Al69 J.Alder et al., Nuc.Phys.147,657(1969).
 An79 B.Antolkovic et al., Nuc.Phys.A235,189(1979).
 Ar56 A.Armstrong et al., Phys.Rev.103,335(1956).
 Ar88 E.Arthur, ICTP Workshop, Trieste (1988) [LA-UR 88-1753].
 Au79 G.Auchampaugh et al., NSE 69,30(1979).
 Au86 G.Auchampaugh & S.Wender, personal communcation (1986).
 Ba85 M.Baba et al., Santa Fe Nuc.Data Conf. (1985)p.223.
 Ca73 J.Cabe et al., CEA-R-4524 (1973).
 Co52 J.Coon, PR 88,562(1952).

Co54 C.Cook & T.Bonner, PR 94,651(1954).
 Co70 J.Cookson et al., NP A146,417(1970).
 Co79 C.Cowan, Summary Documentation ENDF/B-V, ENDF-201 (1979).
 Da83 J.Dave & C.Gould, Phys.Rev. C28, 2212(1983).
 Fl67 F.Flesch et al., OAW 176, 45(1967).
 Gl82 S.Glendingning et al., NSE 80,256(1982).
 Ho69 J.Hopkins et al., NSE 36,275(1969).
 Hy74 M.Hyakutake et al., J.Nuc.Sci.Tech. 11,407(1974).
 Ka81 C.Kalbach & F.Mann, P.R.C23,112(1981).
 Ko83 P.Koehler et al., NP A294,221(1983).
 La70 R.Lane et al., PR C2,2097(1970).
 Mo66 F.Mooring et al., NP 82,16(1966).
 Mu81a S.F.Mughabghab et al., Neutron Res.Parameters and
 Thermal Cross Sect., Academic Press (1981) v1.
 Mu81b S.F.Mughabghab, Proc.Conf.on Nuc.Data Eval.Meth. & Proc.
 BNL-NCS-51363 (1981),V1,p.339.
 Ne73 C.Nelson et al., C00-1717-8 (1973).
 Po70 D.Porter et al., AWRE-0-45-70 (1970).
 Sc70a Schantl, personal communication to NNDC (1970).
 Sc70b W.Scobel et al., ZN A25,1406,(1970).
 St65 J.Strain et al., ORNL-3672 (1965).
 Wh80 R.M.White et al., NP A340,13(1980).
 Wi55 H.Willard et al., PR 98,669(1955).
 Wy58 M.Wyman, PR 112,1264(1958).
 Yo77 P.Young & E.Arthur,LA-6947 (1977).

DESCRIPTION OF EVALUATION FOR NATURAL CARBON PERFORMED FOR ENDF/B-VI†

C. Y. Fu

Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6356, U. S. A.

ABSTRACT

An evaluation of data for neutron induced reactions on natural carbon was performed for ENDF/B-VI and is briefly described. The evaluation is based on R-Matrix fits to measured cross sections for $E_n < 5$ MeV, on least-squares adjustment of the ENDF/B-V data to new experimental information, including KERMA factors, for E_n between 5 and 20 MeV, and on experimental data and theory from 20 to 32 MeV. Evaluated data are given for neutron induced reaction cross sections, angular and energy distributions of the secondary neutrons, and gamma-ray production cross sections and spectra. Uncertainty files are included for the file 3 cross sections. Resonances in ^{13}C below 2 MeV were added. Important improvements to ENDF/B-V were made for the $(n,n'\alpha)$ cross sections. The upper incident energy was extended to 32 MeV, resulting in the addition of cross sections for many more reactions.

1. INTRODUCTION

Major improvements made for ENDF/B-VI carbon are for the energy ranges below 2 MeV (FU90) and above 5 MeV (AX88). For the energy range below 2 MeV, the carbon cross section is an elastic scattering standard. For high-resolution applications, the two small ^{13}C resonances in this energy region may have some effect. Therefore, for ENDF/B-VI, these resonance cross sections and the associated elastic angular distributions were carefully evaluated and combined with the previous R-Matrix results for ^{12}C used for ENDF/B-V. The evaluation (FU78) for the energy range from 2 to 5 MeV was not changed. The evaluation between 5 and 32 MeV was mostly based on the work of Axton (AX88). For the energy range between 5 and 20 MeV, Axton used a least-squares technique to incorporate new data using ENDF/B-V (FU82) as the prior. The evaluation of Axton from 20 to 32 MeV is completely new. Since the evaluations for these three energy ranges have already been documented in detail, the present summary describes only the most important parts of the improvements.

†Research sponsored by the Office of Energy Research, Nuclear Physics, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

In Section 2 the effects of the added resonances of ^{13}C are discussed; Section 3 contains a brief description of the evaluation obtained from R-Matrix fits to experimental data in the energy range below 5 MeV; Section 4 is devoted to the energy range between 5 and 20 MeV; Section 5 the energy range between 20 and 32 MeV; Section 6 summarizes the angular and energy distributions of the secondary neutrons; Section 7 describes the uncertainty files; Section 8 is on needs of important data and possible ways to improve the evaluation.

Much of this information is abstracted from FU78, FU82, FU90, and AX88.

2. INCLUSION OF ^{13}C RESONANCES

The ENDF/B-V differential cross sections for neutron scattering from natural carbon below 2 MeV (FU78), recommended as standards for measurements and based on an R-Matrix analysis for ^{12}C using natural carbon data, are revised to include ^{13}C resonances for high-resolution applications. The recommended ^{13}C cross sections are also based on an R-Matrix analysis (FU90) of the available data. The 0.1529- and 1.736-MeV resonances rise above the natural carbon background by 7% and 1%, respectively. The angular distribution of the elastically scattered neutrons from ^{13}C are generated by the R-Matrix theory and combined with the previous results for ^{12}C to obtain the final recommended data for natural carbon.

Uncertainty information obtained previously (FU78) still appears reasonable and was not changed.

3. CROSS SECTIONS BETWEEN 2 AND 5 MEV

For ENDF/B-IV (and ENDF/B-V), an R-Matrix analysis (FU78) was done for ^{12}C using natural carbon data from 0 to 5 MeV, including polarization data. The results for the energy range between 2 and 5 MeV still appear valid for natural carbon and have been retained for ENDF/B-VI. In this energy range, the capture cross section is nearly negligible, therefore the total cross section and the elastic scattering cross section may be considered the same.

4. CROSS SECTIONS BETWEEN 5 AND 20 MEV

A simultaneous least-squares adjustment of the ENDF/B-V cross sections between 5 and 20 MeV to incorporate new data and KERMA factors was attempted by Axton (AX88). For simplicity, the relative excitation function shape was maintained. This was found to be inadequate for accomodating the newly available $(n,n'3\alpha)$ data (AN86, BR84). These and the older data (reported in MC88) were re-evaluated first and inserted into the ENDF/B-V file for a new fit. The results appear satisfactory and were adopted for ENDF/B-VI.

The $(n,n'3\alpha)$ cross sections between 15 and 20 MeV have been reduced by up to 25%.

5. CROSS SECTIONS BETWEEN 20 AND 32 MEV

The results obtained above for the energy range between 5 and 20 MeV were extended to 32 MeV by Axton (AX88), guided by kinematics and normalization to a few data (MC88). Many new reaction channels become open above 20 MeV. Several of these reactions have no MT assignments in the ENDF/B-VI formats and were merged with those having MT numbers, see File 1 for details.

6. ANGULAR AND ENERGY DISTRIBUTIONS

Angular distributions for all reactions between 5 and 20 MeV were not changed. Angular distributions for elastic scattering from 20 to 32 MeV were evaluated by Axton (AX88) and are based on experimental data (ME84, MC86). Legendre coefficients for the angular distributions for the discrete levels of the (n,n') and $(n,n'3\alpha)$ above 20 MeV were based on linear extrapolation of the ENDF/B-V values below 20 MeV.

Energy distributions for the outgoing neutrons in the continuum part of the $(n,n'3\alpha)$ reaction below 20 MeV were based on a TNG (FU88,SH86) calculation (FU82) and given as evaporation spectra having energy-dependent temperatures. Axton (AX88) did not evaluate the energy distributions in his extension to 32 MeV. For ENDF/B-VI, the energy distributions from 20 to 32 MeV were based on a linear extrapolation of the energy-dependent temperatures from those below 20 MeV.

7. UNCERTAINTY INFORMATION

Uncertainties files are given only for the cross sections in File 3, and not for energy distributions or angular distributions. Fractional and absolute components, correlated only within a given energy interval, are based on least-squares estimates (FU78) of the individual experimental data for $E_n < 2$ MeV and on scatter in experimental data for higher energies. Minor changes were made to the uncertainty estimates in ENDF/B-V to reflect the improvements made and the extension of the upper energy to 32 MeV.

8. DATA NEEDS AND EVALUATION IMPROVEMENTS

ENDF/B-VI for carbon has been extended to 32 MeV. In the extension, most reaction cross sections were based on estimates. Since $(n,n'3\alpha)$ appears to be the largest of all cross sections from 20 to 32 MeV, some measurements for this cross section would help constrain the estimates for other cross sections. Some $(n,n'3\alpha)$ data are available near 20 MeV, but their spread is a factor of two.

Dickens (DI88) has made an independent evaluation of the carbon cross sections up to 80 MeV for detector response calculations. This work should be compared with Axton's evaluation (AX88) adopted for ENDF/B-VI to determine if Dickens' work could be used, perhaps partially, as an improvement.

REFERENCES

- AN86 B. Antolkovic et al., Proc. Int. Conf. Fast Neutron Physics, p. 137, Dubrovnik, Yugoslavia (1986).
- AX88 E. J. Axton, "Report on An Evaluation of KERMA of Carbon and the Carbon Cross Sections," National Bureau of Standards, 1988 (to be published).
- BR84 D. J. Brenner and R. E. Prael, Nucl. Sci. Eng. 88, 97 (1984).
- DI88 J. K. Dickens, "SCINFUL: A Monte Carlo Based Computer Program to Determine a Scintillator Full Energy Response to Neutron Detection for E_n Between 0.1 and 80 MeV: Program Development and Comparisons of Program Predictions with Experimental Data," ORNL-6463 (1988).
- FU78 C. Y. Fu and F. G. Perey, Atomic Data and Nucl. Data Tables 22, 249 (1978).
- FU82 C. Y. Fu, "Summary of ENDF/B-V Evaluations for Carbon, Calcium, Iron, Copper, and Lead and ENDF/B-V Revision 2 for Calcium and Iron," ORNL/TM-8283, ENDF-325 (1982).
- FU88 C. Y. Fu, Nucl. Sci. Eng. 100, 61 (1988).
- FU90 C. Y. Fu, Nucl. Sci. Eng. 106, 489 (1990).
- MC86 J. C. McDonald, Private Communication to E. J. Axton, National Bureau of Standards (1986).
- MC88 V. McLane et al., Neutron Cross Section Curves, Academic Press, 1988.
- ME84 A. S. Meigoni et al., Phys. Med. Biol. 29, 643 (1984).
- SH86 K. Shibata and C. Y. Fu, "Recent Improvements to the TNG Statistical Model Code," ORNL/TM-10093 (1986).

SUMMARY DOCUMENTATION FOR ^{14}N ENDF/B-VI, MAT = 725

P. G. Young, G. M. Hale, and M.B. Chadwick

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

An evaluation of $n + ^{14}\text{N}$ nuclear data was completed for the first issue of ENDF/B-VI in 1990. The evaluation is divided into two parts: (1) a coupled-channel, R-matrix analysis incorporating all available data (including polarization and $p + ^{14}\text{C}$ data) up to the threshold for inelastic scattering ($E_n = 2.5$ MeV), and (2) an assessment and evaluation of all experimental data for the incident neutron energy range $E_n = 2 - 20$ MeV.

The previous ENDF/B evaluation of neutron-induced reactions with ^{14}N was completed in the early 1970's¹ and was part of an intensive effort to improve calculational capabilities for transporting neutrons and γ -rays through the atmosphere. The results of that analysis have been carried over basically unchanged in ENDF/B libraries since that time.² The availability of new and improved experimental data and theoretical techniques, together with the identification of discrepancies between measurements and calculations involving the transport of neutrons through large distances in air, motivated the present effort. The importance of adequately describing properties of the low-energy resonances, particularly as regards the transport of lower energy neutrons, is clear from the pronounced structure in the total cross section. While our R-matrix analysis represents the resonances to 2 MeV, it is important to note that pronounced structure continues to neutron energies of 9 MeV or higher, with a major feature of the data (from the point of view of neutron transport) being the neutron total cross section minimum near 5 MeV.

Major new experiments completed since 1973 include measurements by Chardine et al.³ of neutron elastic and inelastic scattering angular distributions to the lowest five excited states of ^{14}N for incident neutron energies between 7.7 and 13.5 MeV, and elastic scattering measurements by Templon et al.⁴ at three energies between 11 and 17 MeV, by Perey and Dickens⁵ between 4.3 and 8.6 MeV, by Baba et al.⁶ at 14.2 MeV, and by Petler et al.⁷ at 20 and 25 MeV. Important new information on neutron inelastic scattering is provided by the measurements of Nelson et al.,⁸ Auchampaugh and Wender,⁹ and Rogers et al.¹⁰ up to $E_n = 20$ MeV, as well as the extensive double-differential neutron emission measurements by Baba et al.⁶ and Takahashi et al.¹¹ for $E_n \cong 14$ MeV. A comprehensive new measurement of $^{14}\text{N}(n,p)$ and $^{14}\text{N}(n,\alpha)$ cross sections to discrete states by Morgan et al.¹² up to $E_n = 14$ MeV is useful for the experimental data evaluation as well as the R-matrix studies.

Unfortunately, the only major new total cross section measurement,¹³ covering the energy range $E_n = 0.97$ to 5.3 MeV, is in substantial disagreement ($\sim 14\%$) with older, precision measurements and was not included in the ENDF/B-VI evaluation. Consequently, major changes were not made in the evaluated total cross section. However, the new experimental measurements listed above resulted in considerable modification of some of the evaluated data. A comparison of the ENDF/B-VI and ENDF/B-V evaluated elastic scattering cross section between 0 and 6 MeV is given in Fig.

1 and between 6 and 20 MeV in Fig. 2. Note that the cross section has increased by some 9% near $E_n = 14$ MeV. Comparisons of the ENDF/B-VI $^{14}\text{N}(n,n')$ cross sections to the first 2 excited states with the previous ENDF/B-V evaluation are given in the upper halves of Figs. 3 and 4, also including the available experimental data. Gamma-ray production cross sections for emission of prominent γ -rays from these same states are shown in the lower halves of Figs. 3 and 4. To further illustrate the extent of some of the changes, elastic scattering angular distributions from ENDF/B-VI and ENDF/B-V are compared with experimental data at $E_n = 4.92, 11.1$, and 13.5 MeV in Fig. 5. Similarly, new and old inelastic ($E_x = 2.31$ MeV) angular distributions are compared at selected energies in Fig. 6.

Additional details of the evaluation are included in the ENDF File 1 comments.

II. REFERENCES

1. P. G. Young and D. G. Foster, Jr., "An Evaluation of the Neutron and Gamma-Ray Production Cross Sections for Nitrogen," Los Alamos Scientific Laboratory report LA-4725 (1972).
2. P. G. Young and D. G. Foster, Jr., ENDF/B-V data file for ^{14}N (MAT 1275), described in "ENDF/B Summary Documentation," R. Kinsey, Comp., Brookhaven National Laboratory report BNL-NCS-17541 (ENDF-201), 1979 (available from the National Nuclear Data Center, Brookhaven National Laboratory, Upton, N.Y.).
3. J. Chardine, G. Haouat, S. Seguin, and C. Humeau, "Diffusion Elastique et Inelastique de Neutrons sur ^{14}N Entre 7.7 et 13.5 MeV," Commissariat à l'Energie Atomique report CEA-N-2506 (1986).
4. J. A. Templon, J. H. Dave, C. R. Gould, and S. Singkarat, "Neutron Scattering Cross Sections from ^{14}N and ^9Be at 11, 14, and 17 MeV," *Nucl. Sci. Eng.* **91**, 451 (1985).
5. F. G. Perey and J. K. Dickens, "Nitrogen Neutron Elastic and Inelastic Scattering Cross Sections from 4.34 to 8.56 MeV," Oak Ridge National Laboratory report ORNL-4805 (1974).
6. M. Baba, M. Ono, N. Yabuta, T. Kikuti, and N. Hirakawa, "Scattering of 14.1-MeV Neutrons from ^{10}B , ^{11}B , C, N, O, F, and Si," Proc. Int. Conf. on *Nuclear Data for Basic and Applied Science*, Santa Fe, New Mexico, 13-17 May 1985, v.1, p.223.
7. J. S. Petler, M. S. Islam, R. W. Finlay, and F. S. Dietrich, "Microscopic Optical Model Analysis of Nucleon Scattering from Light Nuclei," *Phys. Rev. C* **32**, 673 (1985).
8. R. O. Nelson, S. A. Wender, C. M. Laymon, H. A. O'Brien, R. C. Haight, G. L. Morgan, D. Drake, P. G. Young, M. Drosig, H. Vonach, A. Pavlik, D. C. Larson, P. Englert, and J. Brueckner, "Studies of Higher-Order (n,x γ) Reactions at the WNR Spallation Neutron Source," to be presented at the Washington meeting of the American Physical Society, 16-19 April 1990.
9. G. Auchampaugh and S. Wender, Los Alamos National Laboratory, personal communication (1986).

10. V. C. Rogers, V. J. Orphan, C. G. Hoot, and V. V. Verbinski, "Gamma-ray Production Cross Sections for Carbon and Nitrogen from Threshold to 20.7 MeV," Defense Nuclear Agency report DNA 3495F (1974).
11. A. Takahashi, J. Yamamoto, K. Ohshima, M. Fukazawa, Y. Yanagi, M. Ueda, J. Miyaguchi, S. Kohno, K. Yugami, H. Nonaka, E. Ichimura, H. Sugimoto, and K. Sumita, "Double-Differential Neutron Emission Cross Sections around 14 MeV," OKTAV--87-01 (1987).
12. G. L. Morgan, "Cross Sections for the $^{14}\text{N}(n,p_0)$, (n,α_0) , and (n,α_1) Reactions from 0.5 to 15 MeV," *Nucl. Sci. Eng.* **70**, 163 (1979).
13. J. Bommer, "The Total Neutron Cross Sections of ^{14}N and ^{24}Mg ," *Nucl. Phys.* **A263**, 86 (1976).

$^{14}\text{N}(n,n)$ Elastic Scattering

Figure 1. Neutron elastic scattering cross section for ^{14}N between 0.01 and 6 MeV. The solid curve is the ENDF/B-VI evaluation and the dashed curve is ENDF/B-V.

$^{14}\text{N}(n,n)$ Elastic Scattering

Figure 2. Neutron elastic scattering cross section for ^{14}N between 6 and 20 MeV. The solid curve is the ENDF/B-VI evaluation and the dashed curve is ENDF/B-V.

$$^{14}\text{N}(n,n') E_x = 2.313 \text{ MeV}$$

$$^{14}\text{N}(n,n'\gamma) E_\gamma = 2.313 \text{ MeV}$$

Figure 3. Inelastic neutron and γ -ray cross sections from $^{14}\text{N}(n,n')$ reactions to the first excited state ($E_x = 2.313 \text{ MeV}$) of ^{14}N . The ENDF/B-VI evaluation (solid curve) is compared to the ENDF/B-V evaluation (dashed curve) and to the available experimental data.

$$^{14}\text{N}(n,n') E_x = 3.948 \text{ MeV}$$

$$^{14}\text{N}(n,n'\gamma) E_\gamma = 1.635 \text{ MeV}$$

Figure 4. Inelastic neutron and γ -ray cross sections from $^{14}\text{N}(n,n')$ reactions to the second excited state ($E_x = 3.948 \text{ MeV}$) of ^{14}N . The ENDF/B-VI evaluation (solid curve) is compared to the ENDF/B-V evaluation (dashed curve) and to the available experimental data.

Figure 5. Measured and evaluated $n + {}^{14}\text{N}$ elastic scattering angular distributions at $E_n = 4.92, 11.1, \text{ and } 13.5 \text{ MeV}$. The solid curves are from the ENDF/B-VI evaluation and the dashed curves are ENDF/B-V.

Figure 6. Measured and evaluated inelastic scattering angular distributions for the first excited state of ^{14}N ($E_x = 2.313$ MeV) at $E_n = 10.65$, 12.4 , and 13.5 MeV. The solid curves are from the ENDF/B-VI evaluation and the dashed curves are ENDF/B-V.

SUMMARY DOCUMENTATION FOR ^{15}N ENDF/B-VI, MAT = 728

E. D. Arthur, P. G. Young, and G. M. Hale

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

At energies below $E_n = 5.4$ MeV, the ENDF/B-V evaluation is based on an R-matrix analysis of total and elastic scattering angular distribution measurements and is thought to be reasonably reliable. At higher energies experimental data are more sparse, however, and the ENDF/B-V evaluation relies almost entirely on Hauser-Feshbach statistical theory calculations to complement the available total cross section measurements. Because new nuclear structure information and improved calculational methods have become available since the Version V evaluation, an updated evaluation for $E_n \geq 5.4$ MeV was performed for ENDF/B-VI using the GNASH Hauser-Feshbach statistical theory code.¹

Fig. 1 compares the ENDF/B-V and ENDF/B-VI neutron total cross section evaluations with the experimental data of Zeitnitz² from 6 to 20 MeV. The evaluated elastic cross sections are compared over the same energy range in Fig. 2. The new elastic results are $\approx 20\%$ higher than ENDF/B-V above 11 MeV and are more consistent with measured data for ^{14}N .

The (n,n') and $(n,2n)$ cross sections from the Version V and VI evaluations are compared in Fig. 3. Similarly, the (n,p) and (n,d) evaluated cross sections are compared in Fig. 4 with the very limited experimental data² that are available. The increase in the elastic cross section noted above for ENDF/B-VI was accompanied by significant decreases in the $(n,2n)$, (n,p) , (n,α) , and (n,np) cross sections. Changes were less dramatic in the (n,n') and (n,d) cross sections, although significant modifications did occur for (n,n') reactions to individual levels.

The ENDF File 1 comments are reproduced following the illustrations.

¹ P. G. Young and E. D. Arthur, "Evaluation of Neutron-Induced Reactions on ^{15}N ," in Applied Nuclear Data Research and Development Seminannual Progress Report, Oct. 1, 1982 - Mar. 31, 1983 (compiled by E. D. Arthur, 1983) LA-9841-PR, p.9.

² See references given in ENDF File 1 comments, which are attached.

Figure 1. Neutron total cross section of ${}^{15}\text{N}$ from $E_n = 0$ to 20 MeV. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

$n + {}^{15}\text{N}$ Elastic Cross Section

Figure 2. Evaluated elastic scattering cross sections for ${}^{15}\text{N}$ from 0 to 20 MeV. The solid curve is the ENDF/B-VI evaluation, and the dashed curve is ENDF/B-V.

$^{15}\text{N}(n,2n)^{14}\text{N}$ Cross Section

$n + ^{15}\text{N}$ Total Inelastic Cross Section

Figure 3. Evaluated $^{15}\text{N}(n,n')^{15}\text{N}^*$ and $^{15}\text{N}(n,2n)^{14}\text{N}$ cross sections from threshold to 20 MeV. The solid curve is the ENDF/B-VI evaluation, and the dashed curve is ENDF/B-V.

Figure 4. The $^{15}\text{N}(n,p)^{15}\text{C}$ and $^{15}\text{N}(n,d)^{14}\text{C}$ cross sections from threshold to 20 MeV. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

¹⁵₇N

Reference: No Primary Reference
Evaluators: E. Arthur, P. Young, G. Hale
Evaluated: September 1983
Material 728
Content: Neutron transport, Gamma production

ENDF/B-VI Modifications

Revised spring, summer of 1983 by Young and Arthur to include improved optical model parameters, updated level schemes, improved version of GNASH, and more careful treatment of level density parameters. See Arthur and Young articles in the Oct.1982 - Mar.1983 T-2 progress report (LA-9841-PR, 1983) for details. sections updated are,

MF= 3/ MT=2,4,16,22,28,51-57,91,103,104,105,107

MF=13/ MT=4,16,22,28,103,104,105,107

MF=14/ MT=22,28

MF=15/ MT=4,16,22,28,103,104,105,107

ENDF/B-V Description

The lack of experimental data, with the exception of total cross section measurements, led us to obtain cross sections with a variety of nuclear model codes. We used an energy dependent R matrix analysis to fit the total cross section data of Ze71 for neutron energies from 0.8 to 5.4 mev. As a starting point we used the level scheme reported in Ze71. Also used in the R matrix analysis were the elastic angular distributions of Si62 and Ze71 in the energy range from 1.9 to 3.64 MeV. The parameters of the R-matrix analysis were then used to generate cross sections and to determine legendre coefficients in the range from 1.0E-05 eV to 5.4 MeV. In the energy region from 6 to 20 MeV we used two statistical model codes to generate cross sections. With the first, COMNUC, we computed the angular distributions of elastically scattered neutrons. The legendre coefficients obtained were then joined smoothly

with those from the R-matrix analysis. We also obtained angular distributions of neutrons leading to the first seven excited states of N-15. To generate neutron transmission coefficients for this part of the analysis we used the global optical model parameters of Wilmore and Hodgson as reported by Pe74, and adjusted them to give good agreement with the measured total cross section. The generation of capture, inelastic, and non elastic cross sections as well as neutron, gamma-ray, and charged-particle spectra was done with the statistical model code, GNASH. This newly developed code is based largely on the statistical model formalism of Uh71. Again neutron transmission coefficients were based on the optical model parameters of Wilmore and Hodgson. Transmission coefficients were generated for protons, deuterons, and He4, with the appropriate global forms reported in Pe74. For tritons we used the global parameters for He-3 reported in Pe74 since these were consistent with triton optical model parameters for individual light nuclei as compiled by Pe74. The Gilbert-Cameron level density model was used with standard values for level density parameters (Gi64). To account for direct and semi-direct processes we used the preequilibrium model of Blann and Cline (Cl71) and assumed a ten percent preequilibrium fraction.

MF=2 --- Resonance parameters ---

MT=151 Effective scattering radius = 0.592916E-12 cm

MF=3 --- Smooth cross sections ---

The 2200 m/s cross sections are

MT= 1 sigma = 4.4185 barns

MT= 2 sigma = 4.418 barns

MT=102 sigma = 24.0E-06 barns

MT=1 Total cross section

from 1.0E-05 eV to 5.4 MeV total cross section values are based on the R-matrix analysis described above. From 5.5 to 20 MeV the evaluated cross section is based on the smoothed results of Ze71.

MT=2 Elastic scattering cross section

zero to 20 MeV, subtracted evaluated non-elastic from total to give elastic cross section.

MT=3 Sum of all cross sections except elastic.

MT=4 Sum of MT=51-57, 91 cross sections.

MT=16 (n,2n) cross section

based on smooth curve drawn through results of the statistical model calculations described before.

MT=22 (n,n alpha) cross section

based on smoothed results of statistical model calculation.

MT=28 (n,np) cross section

based on smoothed results of statistical model calculation.

MT=51-57 Inelastic cross section to discrete states

MT=51	Q=-5.27 MeV	MT=55	Q=-7.301 MeV
MT=52	Q=-5.299 MeV	MT=56	Q=-7.566 MeV
MT=53	Q=-6.324 MeV	MT=57	Q=-8.313 MeV
MT=54	Q=-7.155 MeV		

MT=91 (n,n) to continuum
continuum defined to begin at 0.1 MeV above last N-15
discrete level included in calculation, i.e. continuum
begins at 8.413 MeV.

MT=102 (n,gamma) cross section
zero to 0.010 MeV, 1/v from 24.0E-06 barns at thermal
6 to 20 MeV calculated from statistical model with Axel
(Ax62) giant dipole resonance approximation for gamma-
ray transmission coefficients.
.010 to 6 MeV, cross section based on smoothed curve from
region where 1/v dependence ends to region where
calculated values begin.

MT=103 Sum of MT= 700,701, and 718 cross sections.
MT=104 Sum of MT= 720,721,722,and 738 cross sections.
MT=105 Sum of MT=740,741, and 758 cross sections.
MT=107 Sum of MT= 780-785 and 798 cross sections.

MT=700,701 (n,p) cross section to c15 ground and first excited
states.

MT=700	Q=-8.990 MeV	Ex=0. MeV
701	Q=-9.737	Ex=.747

MT=738 (n,d) to continuum states of C-14
continuum defined to begin at 0.85 MeV, just above first
excited state.

MT=720-722 (n,d) cross section to C-14 ground,first,and second
excited states.

MT=720	Q=-7.984 MeV	Ex=0. MeV
MT=721	Q=-14.079	Ex=6.095
MT=722	Q=-14.567	Ex=6.583

MT=740-742 (n,t) cross section to C-13 ground,first,and second
excited states.

MT=738 (n,d) to continuum states of C-15
continuum defined to begin at 6.681 MeV, just above second
excited level in C-14.

MT=740	Q=-9.902 MeV	Ex=0. MeV
741	Q=-12.992	Ex=3.09
742	Q=-13.586	Ex=3.684

MT=780-785 (n,alpha) cross section to B-12 ground and first
five excited states.

MT=780	Q=-7.623 MeV	Ex=0. MeV
781	-8.576	.953
782	-9.297	1.674
783	-10.244	2.621
784	-10.343	2.72

785 -11.013 3.39
 MT=758 (n,t) to continuum states of C-13
 continuum begins at 3.784 MeV, above second excited level
 in C-13.
 MT=798 (n,alpha) to continuum states of B-12
 continuum begins at 3.49 MeV, just above fifth excited
 state in B-12.
 MF=4 --- Secondary neutron and charged particle angular
 distributions.
 MT=2 Elastic angular distributions
 zero to 5.4 MeV based on results obtained from R matrix
 theory analysis.
 6 to 20 MeV based on optical model calculation.
 MT=16 Angular distribution for (n,2n) reaction
 in the absence of data isotropy in the lab system is assumed.
 MT=22 Angular distribution of neutrons from (n,nalpha)
 reaction assumed isotropic in the lab system.
 MT=28 Angular distribution of neutrons from (n,np) reaction
 assumed isotropic in the lab system.
 MT=51-57 Angular distribution calculated using Hauser-
 Feshbach theory.
 MT=718 Angular distribution of protons from (n,p) reaction
 assumed isotropic in the lab system.
 MT=719 Angular distribution of protons from (n,np) assumed
 isotropic in the lab system.
 MT=738 Angular distribution of deuterons from (n,d) assumed
 isotropic in the lab system.
 MT=758 Angular distribution of tritons from (n,t) reaction
 assumed isotropic in the lab system.
 MT=798 Angular distribution of alphas from (n,alpha) reaction
 assumed isotropic in the lab system.
 MT=799 Angular distribution of alphas from (n,n alpha) reaction
 assumed isotropic in the lab system.
 MF=5---Energy distribution of secondary neutrons and charged
 particles---
 MT=16,22,28,91,718,758,and 798 calculated with statistical
 model code.
 MT=719 used for proton spectra from (n,np) reaction.
 MT=799 used for alpha spectra from (n,n alpha) reaction.
 MF=13 --- Photon production cross sections---
 MT=4,16,103,104,105,and 107 include both discrete and
 continuum contributions and were generated with the
 statistical model calculation.
 MF=14--- Photon angular distributions---
 MT= 4 (n,n) to n-15, all isotropic.
 MT=103 (n,p) to C-15, all isotropic.
 MT=104 (n,d) to C-14, all isotropic.
 MT=105 (n,t) to C-13, all isotropic.

MT=107 (n,alpha) to B-12, all isotropic.
MT=16 (n,2n gamma) to N-14, all isotropic.
MF=15 --- Energy distribution of second photon ---
MT=4,16,103,104,105,107 generated using statistical model.

-----References-----

- Aj70 F.Ajzenberg-Selove, Nucl.Phys. A152,1 (1973).
Aj71 F.Ajzenberg-Selove, Nucl.Phys. A166,1 (1971).
Ax62 L.P.Axel, Phys.Rev. 126,671 (1972).
Cl71 C.K.Cline and M.Blann,Nucl.Phys.A172,225 (1971).
Fe67 P.Fessenden and D.R.Maxson,Phys.Rev.158,948 (1967).
Gi65 A.Gilbert and A.G.W.Cameron,Can.J. of Phys.43,1446 (1965).
Pe74 C.M.Perey and F.G.Perey,Atomic and Nucl. Data Tables 13,293
(1974).
Si62 C.P.Sikkema Nucl.Phys. 32,470 (1962).
Uh71 M.Uhl Acta Physica Austria 31,245 (1971).
Ze71 B.Zeitnitz et al.,Nucl.Phys.A166,443 (1971).

¹⁶₈O

Reference: No Primary Reference
Evaluators: G. Hale, Z. Chen, P. Young
Material: 825
Content: Neutron transport, Gamma production

O-16 Free Atom Evaluation

General information:

The ENDF/B-VI evaluation is a complete revision of the earlier ENDF/B-V evaluation, which dates to the mid 1970's. The new evaluation includes extensive new experimental data, a complete new R-matrix analysis below 6.2 MeV, a covariance analysis of the total cross section above 6.2 MeV, and new Hauser-Feshbach statistical and DWBA calculations for the higher energy regions. Especially important new exp. data include the total cross sections of La80, Oh84, Ci80, the differential elastic and inelastic data of Gl82, Bo89, Pe85 and Ba85, the neutron emiss. measurements of Ta83 and Ba85, and the gamma-ray production data of No78, Mo79, and Ne90.

***** MF=2 *****

MT=2 Effective scattering radius = 5.56256E-13 cm

***** MF=3 *****

Thermal (2200 m/s) cross sections

Total	3.88849 b	Radiative capture	0.1900 mb
Elastic	3.88830 b		

MT=1 Total cross section

0.0 to 6.2 mev, essentially calculated from R-matrix parameters obtained by fitting simultaneously almost all available data for reactions ¹⁶O-¹⁶O(n,n)¹⁶O, ¹⁶O-¹⁶O(n,α)¹³C, ¹³C-¹³C(α,α)¹³C at energies below En = 6.2 MeV. Exp. total cross section data for the analysis were the La80 results below En= 3 MeV, and the Ci80 measurements at higher energies, with inserts of the high-resolution data of Jo79.

6.2 to 20.0 MeV, based on a covariance analysis with the GLUCS

Bayes code (He80) of all the major total cross section measurements (Fo71,Pe72,Sc72a,La80,Ci68,Ci80), with structure inserted from the high-resolution Ci80 measurement.

MT=2 Elastic scattering cross section

0.0 to 6.2 MeV, R-matrix analysis that includes all available exp. data, as described above. Elastic angular dist. data used in the analysis includes La60, Ok55, Fo58, Ma62, Fo70, Hu62, Ph61, Jo67, Ki72, Ch61, Dr76 and Fo64.
6.2 to 20 MeV, obtained by subtracting the sum of MT=4,102, 103,104,107 from MT=1. However, adjustments were made to the (n,n') and (n,alpha) reaction cross sects. to enhance agreement with the elastic measurements of Ph61,Ch61,Bu73, Ba63,Mc66,Be67,Ne72,Ki72,Bo89,Gl82,Pe85 and Ba85. The magnitude of these adjustments was generally small and was greatest near structure and near 18 mev (~15 %).

MT=4 Inelastic cross section

thres. to 20 MeV, sum of MT=51-88. note that MT=4 is actually the sum of the (n,n'), (n,np), (n,nalpha), and (n,2n) cross sections because of the use of the excitation energy binning technique to represent MT=51-88.

MT=51-69 Inelastic cross section to discrete states

thres. to 20 MeV, MT=52-55,59,60 are based mainly on (n,ng) data of Ne90,No78,Dr70,Di70,Ny69,C169,Or70,Bu71,Ha59, and the (n,n') data of Bo89, Ki72, Ne72, Ba63, Mc66, Ba85 and Me69. The cross sections for the remaining MT numbers are based on compd. nuc. calculations with the COMNUC and GNASH Hauser-Feshbach statistical codes (Du70,Yo77), & DWBA direct components calculated with DWUCK (Ku76) using the data and parameters from references Cl90 and Is87. Because only very limited gamma ray ang. dist. data is available, inelastic cross sections were obtained from gamma data by multiplying 55 or 125 deg. data by 4π . This leads to increased uncertainty, particularly for the MT=52, J,Pi=3- state. Accordingly, small adjustments were made in the integrated cross sections to enhance agreement with the elastic and total cross section data listed above.
Note that MT= 51 decays by pair production.

MT=70-88 Inelastic cross section to 0.3-MeV bands centered about the given excitation energies, used in lieu of MT=91 and MF=5.

Thres. to 20 mev, integrated cross section adjusted to give a nonelastic x/s in agreement with the evaluated total and elastic. The x/s was divided among bands according to a nuclear temperature calculation using T=2 MeV.

****Please note that much of the cross section to levels above
****9 MeV subsequently results in charged particle emission.
****MT=56-58,61-66,68-70,72-73,75,77-78,80,82-83,85,87,89, are
****flagged as decaying by alpha emission. This choice leads to
****a total alpha emission x/s similar to data of Li52, Da68, Bo66
****below 17 MeV. The remaining MT numbers above MT=66 are
****flagged as being proton emitters.

MT=102 (n,gamma) cross section

1/v variation from 190 microbarns at 2200 m/s, taken from the
compilation of Mu81.

MT=103 (n,p) cross section

thres. to 20 mev, based on a revision of the evaluation of
Sl65, using the experimental data of Bo66, Se62, De62,
Sc72b, and Ma54.

MT=104 (n,d) cross section

thres. to 20 MeV, compd. nuc. react. theory calc. norm. to
single datum point of Li52 at 14 MeV. Taken over intact
from ENDF/B-V.

MT=107 (n,alpha) cross section

thres. to 20 mev, sum of MT=780-783.

MT=800 (n,alpha) integrated cross section to C-13 ground state

0.0 to 6.2 MeV, based on R-matrix analysis described above
under MT=1. Energy scale of Ba72 adjusted to
match total cross sect and resolution corrections appli-
ed to give width and height of peaks consistent with
total. normalization of 0.808 for Ba72 determined from
R-matrix fit and applied. level scheme determined from
R-matrix analysis and Se86.

6.2 to 20 MeV, based on data of Da63, Da68, Si68, Ba73, and
composite of Mc66a, Ma68, and Le68 at 14 MeV. Note that
the Da63 data were renormalized by factor of 1.7 to
bring them into rough agreement with the R-matrix analysis
of the Ba73 (n,a0) data, together with the total and
elastic data in the analysis. Because the Da68 exp. data
were normalized to Da63, the former were also re-
normalized by the factor of 1.7.

MT=801-803 (n,a) cross section to excited levels of C-13.

thres. to 15 mev, used (n,a) data of Da63 and Si68, and the
(n,ag) data of Ne90, No78, Di70, Or70, Cl69, Ny69, En64,
and Bu71.

***** MF=4 *****

MT=2 Elastic angular distributions

0.0 to 6.2 MeV, calculated from R-matrix fit (see MF=3, MT=1).

Measured angular distributions input to the fit were those of Ch61, Fo58, Fo70, Hi58, Hu62, Jo67, Ki72, La60, Li66, Ma62, Ph61, and Jo79. Fits were attempted using new data of Sc77 but those measurements were found to have significant uncertainties and some discrepancies with other data were noted.

6.2 to 20 MeV, smooth curve through coefficients derived from fits to elastic data of Ph61, Ne72, Ch61, Ba63, Be67, Mc66, Ki72, Bu73, Ne72, Gl82, Ba85, Bo89, & Pe85.

MT=51-55 Inelastic angular distributions anisotropic

above 10 MeV, based on fits of the 14-MeV data of Ki72, Ne72, Ba63, Mc66, Ba85, Me69, and especially Bo89.

MT=56-88 Inelastic angular distributions

threshold to 20 mev. assumed isotropic in CM system.

***** MF=12 *****

MT=102 Capture gammas, private communication from Ju64.

Anisotropic secondary gamma ray angular distributions were added for the 6.131 and 6.917 MeV gammas.

These results taken directly from ENDF/B-V.

***** MF=13 *****

MT=4 (n,ng) cross sections

thres. to 15 MeV, based mainly on data of Ne90, No78, Ha59, Di70, Or70, Dr70, Cl69, Lu70, Bu71, Ny69, obtained from MF=3, MT=51-60 using an 0-16 decay scheme from Aj86a.

****Note that the first excited level of o16 at 6.052 mev is
****assumed to decay with emission of two 0.51 MeV gamma rays.

MT=22 (n,nag) cross sections

thres. to 20 MeV, smooth curve based crudely on (n,na) cross section and known levels in C-12, adjusted to agree with composite of 14 MeV data of Cl69, Or70, and data of Ne90.

MT=103 (n,pg) cross sections

thres. to 20 MeV, based on crude division of MF=3, MT=103 x/s among available levels according to $(2J+1)$.

These results were taken from ENDF/B-V.

MT=107 (n,ag) cross sections

thres. to 20 MeV, based mainly on (n,ag) data of Ne90, No78,
Di70, Or70, Cl69, Ny89, En64, Bu71, and (n,a) data of Da63
and Si68.

***** MF=14 *****

Assumed isotropic except for the 6.131-MeV gamma, which is
based on the angular dist. measurement of Dr70, Lu70,
Bu71, and Di70, and the 6.917-MeV gamma, which is based
on the two-angle measurements of Di70. These results
were carried over from ENDF/B-V.

MT=22 (n,ag) angular distributions
assumed isotropic.

MT=103 (n,pg) angular distributions
assumed isotropic.

MT=107 (n,ag) angular distributions
assumed isotropic.

***** MF=33 *****

to be provided at a later date.

----- References -----

- Ad49 R.K.Adair et al., Phys. Rev. 75, 1124 (1949).
Aj86a F.Ajzenberg-Selove, Nucl. Phys. A460,1 (1986).
Aj86b F.Ajzenberg-Selove, Nucl. Phys. A449,1 (1986).
Ba63 R.W.Bauer et al., Nucl. Phys. 47, 241 (1963).
Ba73 J.K.Bair and F.X.Haas, Phys.Rev.C7,1356 (1973).
(n,a0) xsect. obtained by detailed balance from this ref.
Ba85 M.Baba et al., 1985 Santa Fe Nuc.Data Conf. 1, p223.
Be67 P.L.Beach et al., Phys. Rev. 156, 1201 (1967).
Bo66 M.Bormann et al., Proc. IAEA Conf. Nucl. Data,
Paris (1966), p.225.
Bo73 M.Bormann, Zeit.Phys. 258, 285 (1973).
Bo89 G.Borker, W.Mannhart et al., PTB-N-1 (1989) and
personal communication.
Bu71 P.S.Buchanan et al., ORO-2791-32 (1971).
Bu73 W.P.Bucher et al., BRL-R-1652 (1973).
Ch61 L.F.Chase et al., AFSWC-TR-61-15 (1961).
Ci68 S.Cierjacks et al., KFK-1000 (1968).
Ci80 S.Cierjacks et al., Nuc.Inst.Meth. 169, 185 (1980).
Cl69 G.Clayeux and G.Grenier, CEA-R-3807 (1969).
Da63 E.A.Davis et al., Nucl. Phys. 48, 169 (1963).

Da68 D.Dandy et al., AWRE 060/68 (1968).
 De62 J.A.Dejuren et al., Phys. Rev. 127, 1229 (1962).
 Di70 J.K.Dickens+F.G.Perey, Nucl. Sci. Eng. 40, 283 (1970).
 Dr70 D.M.Drake et al., Nucl. Sci. Eng. 40, 294 (1970).
 Dr76 L.Drigo et al., NC/A 31,(1), (1976).
 Du70 C.Dunford, AI-AEC-12931 (1970).
 En64 F.C.Engesser+W.E.Thompson, J. Nucl. Eng. 21, 487 (1967).
 Fo58 J.L.Fowler and H.O.Cohn, Phys.Rev. 109,89 (1958).
 Fo64 J.L.Fowler and C.H.Johnson, ORNL-3582, p.90 (1964).
 Fo70 J.L.Fowler and C.H.Johnson, Phys.Rev.C2,124 (1970).
 Fo71 D.Foster and D.Glasgow, Phys.Rev.C3, 576 (1971).
 Fo73 J.L.Fowler et.al., Phys.Rev. C8,545 (1973).
 Gl82 S.Glendingning et al., Nuc.Sci.Eng. 82, 393 (1982).
 Ha59 H.Hall et al., Nuc.Phys.14, 295 (1959).
 He80 D.Hetrick and C.Y.Fu, ORNL/TM-7341 (1980).
 Hi58 R.W.Hill, Phys.Rev. 109,2105 (1958).
 Hu62 W.Hunzinger and P.Huber, Helv.Phys.Acta 35,351 (1962).
 Is87 M.Islam et al., Nuc.Phys. A464, 395 (1987).
 Jo48 W.B.Jones,Jr., Phys. Rev. 74, 364 (1948).
 Jo67 C.H.Johnson and J.L.Fowler, Phys.Rev. 162,890 (1967).
 Jo73 C.H.Johnson, Phys.Rev. C7,561 (1973).
 Jo79 C.Johson and J.Fowler, 1979 Knoxville Nuc.Data Conf..
 Ju64 E.T.Jurney and H.T.Motz, Bull.Am.Phys.Soc. 9, 176 (1964).
 Ki72 W.E.Kinney and F.G.Perey, ORNL-4780 (1972).
 Ku76 P.D.Kunz, Unpublished Report (1976).
 La60 R.O.Lane et.al., ANL-6172 (1960).
 La80 D.Larson et al., 1980 BNL Conf. on 10-50 MeV Data &
 Personal Communication (1989).
 Le68 B.Leroux et al., Nucl. Phys. A116, 196 (1968).
 Li52 A.B.Lillie, Phys. Rev. 87, 716 (1952).
 Li66 D.Lister and A.Sayres, Phys.Rev. 143,745 (1966).
 Lu70 B.lundberg et al., physica scripta 2, 273 (1970)
 ma54 h.cLmartin, Phys. Rev. 93, 498 (1954).
 Ma62 J.P.Martin and M.S.Zucker, Bull.Am.Phys.Soc. 7,72 (1962).
 Ma68 D.R.Maxson and R.D.Murphy, Nucl. Phys. A110, 555 (1968).
 Mc66 W.J.Mcdonald et al., Nucl. Phys. 75, 353 (1966).
 Mc66a W.N.Mcdicken and W.Jack, Nucl. Phys. 88, 457 (1966).
 Me49 E.Melkonian, Phys. Rev. 76, 1750 (1949).
 Mo79 G.Morgan et al., ORNL-5575 (1979).
 Mu81 S.Mughabghab et al., Neut.Cross Sect.,Vol.1, Neut.Reson.
 Para. and Ther. Cross Sect., Academic Press (1981).
 Ne72 D.O.Nellis and P.S. Buchanan, DNA-2716 (1972).
 Ne90 R.Nelson et al., LANL (WNR), Pers. Communication,3/1990.
 No78 C.Nordborg et al., Nuc.Sci.Eng. 66, 75 (1978).
 Ny71 K.Nyberg, P.S. 4, 165 (1971).
 Ch84 M.Ohkubo et al., Pers.Communication to NNDC, BNL.
 Cl90 N. Olsson et al., Nuc. Phys. A509, 161 (1990).
 Ok55 A.Okazaki, Phys. Rev. 99, 55 (1955).

Or70 V.J.Orphan et al., Nucl. Sci. Eng. 42, 352 (1970).
Pe72 F.Perey et al., ORNL-4823 (1972).
Pe85 J.Petler et al., Phys.Rev. C32, 673 (1985).
Ph61 D.D.Phillips, Private Communication to BNL (1961).
Sc72a R.Schrack, BAPS 17, 555 (1972).
Sc72b M.Schmidt-Hoenow, RCA-17, 142 (1972).
Sc77 I.Schouky et al., KFK-2503.
Se62 K.w.Seemann AND W.E.Moore, KAPL-2214 (1962).
Se86 F.Ajenberg-Selove, Nuc.Phys. A460, 1 (1986).
Si68 Von I. Sick et al., Helv. Phys. Acta 41, 573 (1968).
Sl65 E.L.Slaggie and J.T.Reynolds, KAPL-M-6452 (1965).
Ta83 A.Takahashi et al., OKTAV-A-83-01.
Ya78 T.Yamamoto et al., NST 15, 797 (1978).
Yo77 P.Young and E.Arthur, LA-6947 (1977).

n + O-16 Total Cross Section

n + O-16 Total Cross Section

n + O-16 Elastic Cross Section

n + O-16 Elastic Cross Section

O-16(n,alpha0) Cross Section

O-16(n,alpha0) Cross Section

O-16(n,agamma) EG = 3.684 MeV

O-16(n,n'gamma) EG = 6.917 MeV

n + O-16 ELASTIC ANGULAR DISTRIBUTIONS
E = 9.708 MeV

n + O-16 ELASTIC ANGULAR DISTRIBUTIONS
E = 18.000 MeV

¹⁹₉F

Reference: No Primary Reference
Evaluators: Z. X. Zhao, C. Y. Fu, and D. C. Larson
Evaluated: June 1990
Material: 925
Content: Neutron transport, Gamma production, Covariances

Summary of ENDF/B-VI Evaluation

This is a new evaluation based upon measured data available on a CSISRS data tape obtained from the National Nuclear Data Center (Cs90), and theoretical calculations using the statistical and pre-compound theory code TNG (Sh86, Fu88).

File Information:

MF=2 MT=151 Resonance parameters. Only the scattering length is given (Ko79).

MF=3
MT=1 Smooth neutron cross sections.
Total interaction cross sections.
10⁻⁵ to 10 eV. Sum of the free atom scattering cross section and the capture cross section.
10 eV to 2.0 MeV. Calculated from a set of resonance parameters which were adjusted to fit the high resolution measured total cross sections (La76) by using the SAMMY multilevel R-matrix analysis code (La87).
2.0 MeV to 20 MeV. ENDF/B-V cross sections were used in this energy range.

MT=2 The elastic scattering was obtained by subtracting the non-elastic cross section from total cross section.

MT=3 The non-elastic scattering was obtained from MT3 = MT4 + MT16 + MT22 + MT28 + MT102 + MT103 + MT104 + MT105 + MT107.

MT=4 The total inelastic scattering was obtained by summing MT = 51 to 71 and MT = 91.

MT=16 The (n,2n) reaction was calculated using the TNG code.

MT=22 The (n,nα) + (n,αn) reaction was calculated using the TNG code.

- MT=28 The (n,np) + (n,pn) reaction was calculated using the TNG code.
- MT=51, 52 The inelastic scattering for the first and second levels was obtained from the ratio $cs(ex) / mg(TNG)$. $Cs(ex)$ denotes the measured 110 keV and 197 keV γ -ray production cross sections (Mo74), and $mg(TNG)$ are multiplicities for the 110 keV and 197 keV γ -rays calculated using TNG.
- MT=53, 54... 71 The inelastic scattering for other levels was calculated with TNG and the DWUCK code (Ku72).
- MT=91 Continuum inelastic scattering cross sections were calculated using TNG.
- MT=102 The (n, γ) reaction from 10^{-5} eV to 2.0 MeV was calculated using SAMMY. A set of resonance parameters were used in the calculation for total cross sections but the γ widths were adjusted to fit measured data (Ma63, Ga59). Above 2.0 MeV the capture cross sections are very small so a $1/E$ dependence was used to extrapolate to 20 MeV.
- MT=103 The (n,p) reaction below 8 MeV used ENDF/B-V. Above 8.0 MeV TNG calculated cross sections were used.
- MT=104 The (n,d) reaction was calculated from systematics (Zh88). The excitation function was normalized to 22 mb given by integrating early angular distribution data (Fa68, Re68) at $E_n = 14$ MeV.
- MT=105 The (n,t) reaction was calculated from systematics (Zh88). The excitation function was normalized to spectrum averaged measured data (Qa78).
- MT=107 The (n, α) reaction below 9 MeV remains unchanged from ENDF/B-V. Above 9.0 MeV TNG code results were used.
- MF=4 Angular distributions of secondary neutrons.
- MT=2 The legendre coefficients for elastic scattering were calculated with the GENOA optical model code (Pe67). The compound cross sections and legendre coefficients were calculated using TNG as input. Before doing the calculation, optical model parameters were adjusted to fit the measured angular distributions (Ba85 and Cl70). The same optical model parameters were used in the TNG, DWUCK and GENOA calculations.

- MF=4 MT=51, 52... 71 The inelastic scattering cross sections $cs(d)$, and legendre coefficients $fl(d)$ contributed from direct interaction were calculated using DWUCK and $cs(s)$ and $fl(s)$ were obtained from the statistical TNG calculation using $fl = \frac{cs(d) \times fl(d) + cs(s) \times fl(s)}{cs(d) + cs(s)}$.
- MF=6 Double differential cross sections (DDCS). The DDCS for all sections were obtained based on TNG calculations for the first outgoing neutron only. For the second outgoing neutron and both the first and second outgoing charged particles, the angular distributions were assumed isotropic. Recoil spectra were generated when the calculated particle spectra were converted to the laboratory system.
- MF=8 Radioactive decay data.
- MF=9 Multiplicities for production of radioactive nuclides.
- MF=12 Transition probability arrays for sections 51-71 are based on (Aj87).
- MF=14 Photon angular distributions for sections 51-71. The inelastic scattering is isotropic.
- MF=33 MT=1, 51, 52, 102, 103, 107 Uncertainties for File 3 cross sections. These are the same as in ENDF/B-V but supplemented with a sub-subsection flag LB=8 for every section.
- MT=2, 3 Obtained using "NC-type" sub-subsections for these two derived cross sections.
- MT=4, 16, 22 and 28 Calculated from a covariance matrix of theoretical parameters (several most sensitive parameters were chosen as follows: energy level density parameters for ^{19}F , ^{19}O and ^{16}N ; optical model parameters for ^{19}F : V_0 and W for real and imaginary well depths, r_r and r_w for the radii of the real and imaginary potentials). The covariance matrix for the theoretical parameters were estimated based on the scatter of available measured data about the theoretical values and the systematic errors of the measured data. A sub-subsection of LB=8 are also given for every section.

MF=33 MT=104 105 Calculated from the covariance matrix of the systematic parameters (Zh88). A sub-subsection of lb=8 are also given for every section.

References

- Aj88 F. Ajzenberg-Selove, Nucl. Phys. A475, 1 (1987).
- Ba85 M. Baba et al., Proc. of Conf. on Nucl. Data for Basic and Applied Science, Santa Fe, (1985).
- Cl70 W. Clarke et al., Nucl. Phys. A147, 174 (1970).
- Cs90 EXFOR data on CSISRS tape from the National Nuclear Data Center at Brookhaven National Laboratory (1990).
- Fa68 M. Fazio et al., Nucl. Phys. A111, 255 (1968)
- Fu88 C. Y. Fu, Nucl. Sci. & Eng. 100, 61 (1988).
- Ga59 F. Gabbard et al., Phys. Rev. 114, 201 (1959).
- Ko79 L. Koester et al., Z. Physik, A292, 95 (1979).
- Ku72 P. D. Kunz, "Distored Wave Code DWUCK 72," University of Colorado (1972).
- La76 D. C. Larson et al., ORNL-TM-5612 (1976).
- La87 N. M. Larson, ORNL/TM-9179 (1987).
- Ma63 R. L. Macklin et al., Phys. Rev., C7, 1766 (1963).
- Mo74 G. L. Morgan and J. K. Dickens, ORNL/TM-4823 (1974).
- Pe67 F. G. Perey, Computer Code GENOA, Oak Ridge National Laboratory Unpublished Computer Program (1967).
- Qa78 S. M. Qaim et al., Nucl. Phys. A295, 150 (1978).
- Sh86 K. Shibata and C. Y. Fu, ORNL/TM-10093 (1986).
- Zh88 Z. Zhao and Z. Zhou, Nucl. Sci. & Eng., 99 367 (1988).

Reference: ANL/NDM-105
Evaluators: A. Smith, D. Smith, P. Guenther, J. Meadows, R. Lawson
(ANL), T. Djemil (U of I) and R. Howerton (LLNL).
Evaluated: June 1988
Material: 2300
Content: Neutron transport, Gamma production, Covariances

1. Introduction

This is a comprehensive evaluated file for neutronic applications, particularly those associated with the development of fission and fusion energy systems. Elemental vanadium consists of two isotopes ^{50}V and ^{51}V . This file is nominally an elemental file, but gives little consideration to the very minor isotope ^{50}V (0.25% abundant).

2. Evaluated Resolved Resonance Range

The resonance parameter representation is employed for incident energies from 2.0 to 100 keV using s- and p-wave parameters of $^{50,51}\text{V}$. The resonance parameters were taken directly from Mughabghab et al.¹ Negative energy resonances were not used, and a background was introduced to assure consistency with the thermal values of Ref. 1 and other experimental data. The present resonance evaluation cannot be directly compared with ENDF/B-V as the latter uses a point-wise representation of the resonance region.

3. Evaluated Total Cross Sections

A comprehensive experimental data base was constructed from the literature, as referenced in CINDA² and augmented with values from measurements undertaken by the authors. This evaluation was constructed from the above energy averaged data base using the statistical processing code GMA.³ Below 5 MeV resonance fluctuations were introduced into the evaluation. Ref. 4 was used up to 220 keV, Ref. 5 from 220 to 360 keV, and Ref. 6 from 0.36 to 6.0 MeV.

4. Evaluated Elastic Scattering Cross Sections

The evaluated energy averaged elastic scattering cross sections were derived from the studies of Ref. 7. Below ≈ 4.0 MeV the cross sections were determined from the

difference between the sum of other partial cross sections and the total cross section. Above 4.0 MeV the present evaluation is significantly lower than ENDF/B-V, and by as much as 15 to 20% from 10 to 14 MeV.

Evaluated Inelastic Scattering Cross Sections

5.1 Discrete Inelastic Processes

The discrete inelastic scattering cross sections are the result of the excitation of 25 levels. The excitations and J^π values are taken from Ref. 8. More than half of the data were obtained using direct neutron detection techniques and half were deduced from measurements of γ rays emitted in the $(n,n\gamma)$ process using a number of reference standards. The $(n,n\gamma)$ measurements were systematically larger than those resulting from direct neutron measurements. Most of the experimental results were obtained at incident energies < 5 MeV. Only 2 sets extend to higher energies.^{7,9} The cross sections were calculated with the model and methods of Ref. 7. The calculated results were compared with the experimental data base, combining calculated excitations as necessary to make comparisons with the experimental values. The calculated results were then subjectively adjusted to improve the description of the measured cross sections. The normalizations were confined to the first few levels and were always less than 10%.

5.2 Continuum Inelastic Scattering Processes

The integrated continuum inelastic scattering cross sections were determined from the difference between the non-elastic cross section and the other partial cross sections. Above 10 MeV the (n,np) reaction becomes significant and is not well known. Above 15 MeV there are large uncertainties in the $(n,2n)$ cross sections. Both contribute to large uncertainties in the continuum inelastic cross section.

6. Evaluated Radiative Capture Cross Sections

The data base for capture cross sections is sparse. The information is essentially all derived from activation measurements. The data are best described in the 200 keV to 1.5 MeV energy range, and that region was used for normalizing calculated values using the code ABAREX¹⁰, the optical model potential of Ref. 7, and an adjustment of the $\ell = 0$ strength function to give a description of the experimental values over the range 200 keV to 1.5 MeV. The calculation describes only the compound nucleus process, so a small direct capture component is added at high energies.

7. Evaluated (n,2n) and (n,3n) Reactions

The (n,2n) Q-value is -11.051 MeV and the (n,3n) Q-value is -20.385 MeV. Thus only the former process need be considered.

Experimental information is obtained only using prompt detection and it is sparse. The data of Frehaut et al.¹¹ and the preliminary data of Ref. 12 were used. The Ref. 12 results are $\approx 10\%$ lower than Ref. 11 in the region of overlap near 15 MeV. These data sets give the only experimental indication of the energy dependence above 15 MeV. The data were extrapolated above 15 MeV using the code CADE.¹³ The CADE results are reasonably consistent with the unconfirmed higher energy values of Ref. 12.

8. Evaluated Charged Particle Emitting Reactions

8.1 (n, α) and (n, $n\alpha$) + (n, αn) Reactions

The Q-value of -2.055 MeV for the (n, α) reaction¹⁶ is relatively low, and the process is readily accessible to activation measurements. The evaluation using the assembled data base and the evaluation program GMA³ systematically follows the smooth curve proposed by Kanno et al.¹⁵ The (n, α) cross sections were also calculated, assuming statistical processes, using the code CADE.¹³ The calculated results had a shape similar to the present evaluation but 30% larger in magnitude. In particular, the calculation indicated that the (n, αn) reaction was very small below ≈ 14 MeV, and that there was no gross structure in the cross section. The present evaluation is $\approx 25\%$ smaller in magnitude at the maximum than that of ENDF/B-V, and it does not show the gross structure of ENDF/B-V at ≈ 10 MeV. Furthermore, the present evaluation gives a much better definition of the threshold region due to recent experimental information in that energy range.¹⁶

The threshold for the (n, $n\alpha$) and (n, αn) processes are relatively high ($Q = -10.294$ MeV)¹⁴ The sum of the cross sections was set equal to the difference between the total α production cross section and the above (n, α, γ) cross section. The total α production cross section was determined by calculations made with CADE, and then normalized to the above evaluation below 12 MeV. The calculated (n, αn) cross sections are much larger than the (n, αn) values up to 20 MeV. The calculations indicate that above 16 MeV the cross section rises rapidly, supported by the experimental results of Ref. 17 and the higher energy (above 20 MeV) measured value of Ref. 18.

8.2 (n,p) and (n,np) + (n,pn) Reactions

The Q-value for the (n,p) cross section is -1.684 MeV.¹⁶ The data can be grouped into 3 categories: (1) extensive ANL measurements¹⁹, (2) a single set of measurements

extending to 20 MeV²⁰ and (3) a number of experimental results at ≈ 14.5 MeV. A weighted average of the accepted experimental values near 14.5 MeV was constructed, yielding a cross section $28.6 \pm 5\%$ mb at 14.5 MeV. Below ≈ 13 MeV there is only one set of experimental data (Ref. 19), and it is in agreement with statistical model calculations using CADE. Thus this set of data, complemented by model calculations to interpolate the shape from 10 to 14 MeV, was used to determine the evaluation below the 14.5 MeV region. There is only one set of data above 14.5 MeV²⁰, and it extends from 13 to 20 MeV. However, its magnitude is 10 to 15% larger than the above evaluated 14.5 MeV result. Moreover, the same authors, in a companion study of the (n,α) reaction, obtained results higher than the present evaluation. Therefore the normalization of this data set is suspect. The evaluation simply uses the measured values of Ref. 20 above ≈ 14.5 MeV, normalized to the evaluated 14.5 MeV result outlined above. Above 10 MeV the present evaluation is systematically smaller than ENDF/B-V by $\approx 25\%$, and below 10 MeV, there are considerable differences in both shape and magnitude. These differences are due to the new and much better data available for the present evaluation. Cross section guidelines are set forth in Ref. 19. They are 20 - 30% larger than the present evaluation, again reflecting new and improved experimental data not available at the time. Ref. 19 does make some valuable comparisons between differential and fission spectrum averaged cross sections.

There is no experimental information for the (n,np) and (n,pn) reactions. Calculations were made using the Weisskopf-Ewing statistical model,¹³ with the addition of pre-compound contributions.²¹ The two calculated results agree very well at ≈ 20 MeV (i.e., $\sigma = 143$ mb). The evaluation makes a compromise between the two types of calculated results, adjusted to give a reasonable neutron emission spectrum. The evaluation is characterized by relatively large values near threshold, falling to much smaller values with the onset of the $(n,2n)$ cross section. Near threshold the (n,np) process is the larger contribution to the total process.

8.3 (n,d) and $(n,nd) + (n,dn)$ Reactions

The (n,d) reaction product is stable ^{50}Ti . Experimental results suggest a cross section in the range of 5 to 10 mb in the 14 to 15 MeV region. The most careful measurement appears to be that of Ref. 22 (≈ 7 mb). The evaluation uses the results of CADE statistical calculations normalized to Ref. 22 at 15 MeV. The threshold of the $(n,nd + dn)$ reaction is very high ($Q = -16.776$ MeV)¹¹ and statistical calculations suggest cross sections of far less than 1 mb even at 20 MeV. The results are very uncertain, perhaps by 100% or more, but, due to the high threshold and the small magnitudes, that uncertainty is of little concern in most applications.

8.4 (n,t) and $(n,nt) + (n,tn)$ Reactions

The cross sections for the (n,t) reaction were calculated using CADE. At 20 MeV the calculations indicate a cross section of 0.48 mb consistent with the exper-

imental 30 MeV value of 0.55 mb given in Ref. 23. Ref. 24 suggests a value 1 mb at ≈ 14 MeV, somewhat larger than our calculated result. The present evaluation uses the calculated results and is approximately an order of magnitude smaller than the comparable evaluation of ENDF/B-V. The $(n, nt + tn)$ cross section is less than 1 microbarn at 20 MeV and was ignored. Other minor reactions were included based upon the statistical model calculations.

8.5 Continuum Neutron Emission Spectra

Individual reaction spectra were calculated using the computer codes ALICE²⁵ and CADE.²⁶ ALICE calculates reaction cross sections and the spectra of the emitted particles. As used it employs a hybrid model for pre-compound processes, and the Weisskopf-Ewing evaporation model for compound decay. It does not include a γ -ray channel. CADE performs compound nucleus calculations by the Weisskopf-Ewing formalism. The level density formalism and parameters are those of Brancazio and Cameron.²⁷ γ -ray emission is described by the giant dipole resonance formalism. ALICE seems to give reasonable spectra for the emitted particles and CADE was used to determine how the spectra were to be apportioned among the several reactions. The above spectra are considerably different from those given in ENDF/B-V.

9. Evaluated Photon Production Reactions

The spectrum of photons from the neutron capture reaction was taken from Orphan et al.²⁸ at thermal energy. The same spectrum was used at 20 MeV with the multiplicity adjusted to conserve energy. For the photons associated with inelastic scattering to specific levels CASCADE²⁹ was used. For other reactions the cross sections and spectra were calculated using the R-parameter formalism of Perkins et al.³⁰

References

1. S. F. Mughabghab, Neutron Cross Sections Vol. 1, Part B, Academic Press Inc. New York, (1984); also S. Mughabghab and C. Dunford, private communication (1982).
2. CINDA, A computerized Index to Nuclear Data, IAEA Press (1986)
3. W. P. Poenitz, Brookhaven National Laboratory Report, BNL-NCS-51363 Vol. I 249 (1981); as modified by M. Sugimoto (1987).
4. F. Manero, Nucl. Phys. 65 419 (1965). Data from 3.2 to 5.2 MeV.
5. A. Smith et al., Phys. Rev. C1 581 (1970). Data from 0.1 to 1.45 MeV.

6. S. Cierjacks et al., Karlsruhe Report, KFK-1000 (1970). Data from 0.36 to 20.0 MeV.
7. A. Smith, P. Guenther, and R. Lawson, Argonne National Laboratory Report ANL/NDM-106 (1988).
8. Zhou Chunmei et al., Nuclear Data Sheets 48 111 (1986).
9. F. Perey and W. Kinney, Oak Ridge National Laboratory Report, ORNL-4551 (1970)
10. P. Moldauer, computer code ABAREX, private communication (1982).
11. J. Frehaut and G. Mosinski, CEA Report, CEA-N-1875 (1976) See also proceedings of the 1980 Gaussig Conference.
12. G. Auchampaugh, D. Drake, and L. Veesser, Brookhaven National Laboratory Report, BNL-NCS-60681 (1977).
13. D. Wilmore and P. E. Hodgson, J. Phys. G11 1007 (1985).
14. J. K. Tuli, Nuclear Wallet Cards, National Nuclear Data Center, Brookhaven National Laboratory (1985).
15. I. Kanno, J. Meadows, and D. Smith, Argonne National Laboratory Report, ANL/NDM-86 (1984).
16. I. Kanno, J. Meadows, and D. Smith, Ann. Nucl. Energy, 11 623 (1984).
17. M. Borman et al., Z. Phys. 166 477 (1962).
18. C. Wu et al., Nucl. Phys. A329 63 (1979).
19. D. Smith, J. Meadows, and I. Kanno, Argonne National Laboratory Report, ANL/NDM-85 (1984).
20. M. Borman et al., Z. Phys. 174 1 (1963).
21. V. Weisskopf and D. Ewing, Phys. Rev. 57 472 (1940); code ALICE by M. Blann, Lawrence Livermore National Laboratory Report, UCID-19614 (1982).
22. S. Grimes et al., Phys. Rev. C17 508 (1978).
23. S. Qaim et al., NEANDC Report, NEANDC(E)-232 (1982).
24. C. Khurana and I. Govil, Nucl. Phys. 69 153 (1965).
25. M. Blann, Lawrence Livermore National Laboratory report, UCID-20169 (1984).

26. D. Willmore, Harwell Report, AERE-R-11515 (1984).
27. P. Brancazio and A. Cameron, Can. Jour. Phys. 47 1029 (1969).
28. V. J. Orphan, N. C. Rasmussen, and T. L. Harper, "Line and Continuum γ -ray Yields from Thermal Neutron Capture in 75 Elements," Gulf General Atomic Report, GA-10248/DASA 2570 (1970).
29. W. E. Warren, R. J. Howerton, and G. Reffo, CASCADE Cray program for γ -production from discrete level inelastic scattering, Lawrence Livermore Nuclear Data Group Internal Report, PD-134 (1986), unpublished.
30. S. T. Perkins, R. C. Haight, and R. J. Howerton, Nucl. Sci. and Eng. 57 1 (1975).

DESCRIPTION OF EVALUATIONS FOR $^{50,52,53,54}\text{Cr}$ PERFORMED FOR ENDF/B-VI*

D. M. Hetrick, D. C. Larson, N. M. Larson, and C. Y. Fu
Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6356

ABSTRACT

Isotopic evaluations for $^{50,52,53,54}\text{Cr}$ performed for ENDF/B-VI are briefly reviewed. The evaluations are based on analysis of experimental data and results of model calculations which reproduce the experimental data. Evaluated data are provided for neutron induced reaction cross sections, angular and energy distributions, and for gamma-ray production cross sections associated with the reactions. File 6 formats are used to represent energy-angle correlated emission and recoil spectra. Uncertainty files are included for all File 3 cross sections.

1. INTRODUCTION

Separate evaluations have been done for each of the stable isotopes of chromium. In this report we briefly review the structure of the evaluations, describe how the evaluations were done, and note the major pieces of data considered in the evaluation process. Experimental data references were obtained primarily from CINDA; the data themselves were mostly obtained from the National Nuclear Data Center at Brookhaven National Laboratory and, occasionally, from the literature and reports. The R-Matrix code SAMMY (LA84) was used for the resonance region analysis. The nuclear model code TNG (FU88, SH86), a multistep Hauser-Feshbach code which includes precompound and compound contributions to cross sections, and angular and energy distributions in a self-consistent manner, calculates gamma-ray production, and conserves angular momentum in all steps, was used for these evaluations above the resonance region. Extensive model calculations were performed for each isotope with the goal of simultaneously reproducing experimental data for all reaction channels with one set of parameters. This ensures internal consistency and energy conservation within the evaluation. In the case of the isotopic chromium evaluations, no reaction was deemed to have enough appropriate data for a Bayesian analysis (HE80), so the evaluations depend upon TNG results and hand-drawn curves. That is, in cases where the available data were deemed to be accurate, but in disagreement with the TNG results, a smoothed curve representation through the chosen data sets was used for the evaluation. A similar method was also used for cross sections where resonant structure was felt to be important, but resonance parameters were not included. The final evaluation is thus a combination of SAMMY results, TNG results (used where extrapolation and interpolation was required and where data sets were badly discrepant), and hand-drawn curves.

In Section 2 the resonance parameters are discussed; Section 3 contains a description of the major cross sections included in the evaluation; Section 4 is devoted to angular distributions; and Section 5 to energy-angle correlated distributions. Section 6 describes the uncertainty files.

* Research sponsored by the Office of Energy Research, Division of Nuclear Physics, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

Much of this information is abstracted from Ref. HE87, a report devoted to a description of the calculations for ^{52}Cr , and Ref. SH87, a similar report for ^{53}Cr .

2. RESONANCE PARAMETERS

From 1 keV to about 900 keV, the parameters are derived from recent ORNL (AG84) and Geel (BR86, RO89) transmission data. Unfortunately, the reported resonance parameters were obtained by analyzing individual energy regions of the experimental data independently, with the same nuclear radius and set of dummy resonances not necessarily used for each region. Thus, simply adopting the resonance parameters and generating pointwise cross sections from them yield results which are often in poor agreement with the original experimental data. Some reanalysis was done with SAMMY (LA89) which, using the Bayesian approach and saving the correlation matrices, provides consistent results independent of whether the data are analyzed in regions or all at one time. These adjustments resulted in resonance parameters which give the correct thermal cross section values as well as providing a better reproduction of the experimental data. However, the results still required a background file in File 3 Section 1 to provide the desired reproduction of the experimental data. Further work needs to be done to provide a consistent, accurate set of resonance parameters commensurate with the quality of those in the iron and nickel evaluations. Capture widths are also taken from the above references, with average parameters used where individual parameters are not available. The resonance parameters should be processed with the Reich-Moore formalism, and not single- or multi-level Breit-Wigner formalisms. Details for the resonance regions for each isotope are given in the File 1 comments for each evaluation.

3. CROSS SECTIONS

In this section the evaluation of reaction cross sections is presented; details for each isotope are given in File 1 of each evaluation. For $^{50,53,54}\text{Cr}$ the total cross section from the resonance region to 4 MeV is taken from the isotopic data. From 4 to 20 MeV, the total cross section is evaluated from recent ORELA transmission measurements on a natural sample. This natural data is also used for ^{52}Cr , the major isotope, from the upper end of the resonance region to 20 MeV. The nonelastic cross section is derived by summing the reaction cross sections, while the elastic cross section is derived by subtracting the nonelastic from the total.

Experimental data for cross sections describing inelastic scattering to discrete levels in ^{52}Cr are available for only the first two levels at incident neutron energies >4 MeV. Thus, results of the TNG model calculations, including direct reaction contributions, were used for the evaluation. The calculations were in good agreement with available data, and with a recent measurement of the total inelastic scattering cross section from 1 to 40 MeV (LA85). Figure 1 shows a comparison of the ENDF/B-V and ENDF/B-VI total inelastic cross section with the experimental data. Cross sections for inelastic scattering in the minor isotopes were also taken from the TNG calculations. In all cases a continuum was used to describe inelastic scattering for excitation energies above the discrete levels.

The calculated (n,p) cross sections were modified to be in acceptable agreement with the experimental data for both $^{52,53}\text{Cr}$ from 5 to 9 MeV, and around 14 MeV. Calculated values for ^{50}Cr are in good agreement with the one measured point near 14 MeV. For ^{54}Cr , the evaluated (n,p) cross section includes the (n,pn) contribution as the (n,np) and (n,pn) cross sections were not included in the evaluation. The calculated values of the

total proton emission cross sections for ^{52}Cr are in good agreement with the data of Smith and Meadows (SM80), Grimes et al. (GR79), Colli et al. (CO62), and Barschall (BA82).

Since the (n, α) reaction on both $^{52,53}\text{Cr}$ leads to a stable nucleus, no activation data are available, only alpha production data from integrating measured alpha spectra. For ^{52}Cr the TNG results agree within uncertainties with the data of Paulsen et al. (PA81) up to 10 MeV, but are at the lower limit of the experimental uncertainties at 14 MeV for the data of Grimes et al. (GR79), Dolja et al. (DO73), and Barschall (BA82). Since much of the energy range to 20 MeV has no data, the calculated results were taken for the evaluation. For $^{50,53,54}\text{Cr}$, the calculated results are used, due to the paucity of data. Except for ^{50}Cr , the cross sections are all less than 50 mb.

Available data for the $^{52}\text{Cr}(n, 2n)$ reaction are in disagreement near 14 MeV. The calculated results from TNG provide a reasonable compromise near 14 MeV, and are about 15% larger than the measurement of Bormann et al. (BO68) from 13 to 20 MeV. Since no data are available for the $^{53}\text{Cr}(n, 2n)$ cross section, the calculated results were used in the evaluation. Combining the TNG results for $^{52,53}\text{Cr}$ give results which are in good agreement with the $(n, 2n)$ natural sample data of Auchampaugh et al. (AU77), Frehaut et al. (FR80), and Frehaut and Mosinski (FR75). File 1 comments in the evaluations give information for $^{50,54}\text{Cr}$.

4. ANGULAR DISTRIBUTIONS

Elastic scattering angular distributions in File 4 Section 2 from ENDF/B-V were compared with available data, found to be adequate, and adopted for ENDF/B-VI. Comparisons with experimental data are given in reference (PR76). Angular distribution data for inelastic scattering to discrete levels are given in File 6. They are symmetric, taken from the TNG compound nucleus calculations, except for levels which include a direct interaction term where the angular distributions are a weighted sum of direct interaction and compound nucleus contributions.

5. ENERGY-ANGLE CORRELATED DISTRIBUTIONS

Particle emission spectra are often measured as a function of outgoing particle angle, and this correlation of outgoing angle with measured energy spectra can now be represented in File 6. However, often these distributions are measured only at one incident energy, and we must rely upon model calculations to reproduce the available energy and angular distribution information so it can be calculated for the evaluation at other incident energies. File 6 is extensively used in this evaluation to represent energy-angle correlated data. Energy spectra for all outgoing particles, including photons, are given in this file. Angular distributions are given for the neutron emission spectra for ^{52}Cr only; angular distributions for all other isotopes and particles are assumed to be isotropic. Figure 2 (HE87) shows a comparison of the calculated and measured angular distributions for three outgoing neutron energy bins. Figure 3 illustrates the components of the neutron emission calculated with TNG which sum to give the total emission spectra for ^{53}Cr . Figure 4 (HE87) shows a comparison of the measured and calculated neutron emission spectra around 14 MeV incident neutron energy. Figures 5 and 6 (HE87) show a comparison of the measured and calculated emission spectra for the $^{52}\text{Cr}(n, xp)$ and $^{52}\text{Cr}(n, x\alpha)$ reactions around 14 MeV. These calculated results, at several incident energies between 1 and 20 MeV, have been adopted for the evaluation and put in File 6.

Prior to incorporation in File 6, the neutron and charged particle energy distributions from TNG are input to the RECOIL code (FU85), which converts the energy distributions from the center of mass to the laboratory frame, and calculates the energy spectrum of the heavy recoil nucleus. These tabulated energy distributions in the lab frame are given in File 6, with the neutrons usually having anisotropic angular distributions, and isotropic angular distributions for the charged particles (including the recoil nucleus).

Figure 7 (HE87) shows a comparison of the calculated gamma-ray emission spectrum at 9.5 MeV with the measurement of Morgan and Newman (MO76) for the bin covering incident neutron energies from 9 to 10 MeV. Calculated results from TNG at several incident neutron energies between 1 and 20 MeV were used for the evaluation, with File 6 being used for ease in calculating energy balance.

As an example of the usage of File 6, consider the $^{52}\text{Cr}(n, n\alpha)$ reaction. In Section 6/22, constant yields are given for the outgoing neutron, alpha, and ^{48}Ti residual, and an energy-dependent yield is used for the gamma rays associated with the $(n, n\alpha)$ reaction. Normalized energy distributions at several incident energies are given for each outgoing product, but only the outgoing neutron has a non-isotropic angular distribution. The cross section to be used for normalization is taken from Section 3/22.

Capture gamma-ray cross sections and spectra are obtained from Files 3, 12 and 15 and are based on a combination of experimental data and calculated results.

Energy balance $((E_n + Q))$ must equal sum of all outgoing particle and gamma-ray energies) has been checked for all reactions, energies and isotopes, and is achieved within 1%.

6. UNCERTAINTY INFORMATION

Uncertainty files are given only for the cross sections in File 3, and not for the resonance parameters, energy distributions or angular distributions. Fractional and absolute components, correlated only within a given energy interval, are based on scatter in experimental data and estimates of uncertainties associated with the model calculations. Further details are given in (HE91).

7. DATA NEEDS AND EVALUATION IMPROVEMENTS

Consistent sets of transmission data for each isotope of chromium from 100 eV to 20 MeV would allow improved resonance parameter analyses, which is a problem with the current evaluations. High resolution resonance scattering cross sections for $^{52,53}\text{Cr}$ would allow determination of spins and parities. Capture cross-section data used in these evaluations have large uncertainties since they suffer from the response function difficulties observed by the International Task Force for the 1.15-keV iron resonance, and existing data should be reanalyzed or new data acquired. The $^{50}\text{Cr}(n, p)$ and (n, np) cross sections are large, with only one data point available, and existing evaluations of these cross sections disagree. New data for these reactions would be useful. $^{52}\text{Cr}(n, p)$ would benefit from new data above 10 MeV, where the data are nonexistent or in significant disagreement. $^{53,54}\text{Cr}(n, 2n)$ have large cross sections, and some experimental verification of the evaluations is necessary. As with many of the structural materials in this mass range, the shape of the (n, α) cross section is not well predicted by models, and further experimental data over a wide energy range would be useful for understanding of this problem. Neutron emission data for incident neutron energies other than 14 MeV is necessary to improve nuclear

models presently used for extrapolation. Uncertainties should be provided for important resonance parameters, and angular and energy distributions.

REFERENCES

- AG84 H. M. Agrawal, J. B. Garg, and J. A. Harvey, *Phys. Rev. C* **30**(6), 1880 (1984).
- AU77 G. F. Auchampaugh, D. M. Drake, and L. R. Veecer, *Neutron Cross Section Programs in the Energy Region from 1 to 24 MeV at the LASL Van de Graaff Facilities*, BNL-NCS-50681, pp. 231-246 (May 1977).
- BA82 H. H. Barschall, Europhys. Topical Conf. on Neutron Induced Reactions, Smolenice (June 1982); Proceedings published by Slovak Acad. Sciences as *Physics and Applications*, **10**, 279 (1982).
- BO68 M. Bormann, A. Behrend, I. Richle, and O. Vogel, *Nucl. Phys.* **A115**, 309 (July 1968).
- BR85 A. Brusegan, R. Buyl, F. Corvi, L. Mewissen, F. Poortmans, G. Rohr, R. Shelley, T. Van der Veen, I. Van Marcke, "High Resolution Neutron Capture and Total Cross Section Measurements of ^{50}Cr , ^{52}Cr , and ^{53}Cr ," p. 633 in *Proc. Int. Conf. Nuclear Data for Basic and Applied Science*, Vol. 1, Santa Fe, N.M., May 13-17, 1985.
- CO62 L. Colli, I. Iori, S. Micheletti, and M. Pignanelli, *Nuovo Cimento* **21**, 966 (1962).
- DO73 G. P. Dolja, V. P. Bozhko, V. Ja. Golovnja, A. P. Kljucharev, and A. I. Tutubalin, *Proc. Nejtronnaja Fizika* **3**, 131 Kiev (1973).
- DR78 D. M. Drake, E. D. Arthur, and M. G. Silbert, *Nucl. Sci. Eng.* **15**, 54 (1978).
- FR75 J. Frehaut and G. Mosinski, 5th Int. Seminar on Interaction of Fast Neutrons with Nuclei, Gaussig, (November 1975).
- FR80 J. Frehaut, A. Bertin, R. Bois, and J. Jary, *Status of $(n, 2n)$ Cross Section Measurements at Bruyeres-Le-Chatel*, BNL-NCS-51245, pp. 399-411 (May 1980).
- FU88 C. Y. Fu, *Nucl. Sci. Eng.* **100**, 61 (1988).
- GR79 S. M. Grimes, R. C. Haight, K. R. Alvar, H. H. Barschall, and R. R. Borchers, *Phys. Rev. C* **19**, 2127 (June 1979).
- HE75 D. Hermsdorf, A. Meister, S. Sassonoff, D. Seeliger, K. Seidel, and F. Shahin, Zentralinstitut fur Kernforschung Rossendorf Bei Dresden, Zfk-277 (U) (1975).
- HE80 D. M. Hetrick and C. Y. Fu, *GLUCS: A Generalized Least-Squares Program for Updating Cross Section Evaluations With Correlated Data Sets*, ORNL/TM-7341, ENDF-303 (October 1980).
- HE87 D. M. Hetrick, C. Y. Fu, and D. C. Larson, *Calculated Neutron-Induced Cross Sections for ^{52}Cr from 1 to 20 MeV and Comparisons with Experiments*, ORNL/TM-10417, ENDF-346 (September 1987).
- HE91 D. M. Hetrick, D. C. Larson and C. Y. Fu, *Generation of Covariance Files for the Isotopes of Cr, Fe, Ni, Cu, and Pb in ENDF/B-VI*, ORNL/TM-11763 (February 1991).
- LA85 D. C. Larson, "High-Resolution Structural Material $(n, x\gamma)$ Production Cross Sections for $0.2 < E_n \leq 40$ MeV," p. 71 in *Proc. Conf. on Nuclear Data for Basic and Applied Science*, Santa Fe, New Mexico, Vol. 1, 1985.

- LA89 N. M. Larson, *Updated Users' Guide for SAMMY: Multilevel R-Matrix Fits to Neutron Data Using Bayes' Equations*, ORNL/TM-9179 (August 1984). Also ORNL/TM-9179/R1 (July 1985) and ORNL/TM-9179/R2 (June 1989).
- MO76 G. L. Morgan and E. Newman, *The $\text{Cr}(n, \gamma)$ Reaction Cross Section for Incident Neutron Energies Between 0.2 and 20.0 MeV*, ORNL/TM-5098, ENDF-222 (January 1976).
- PA81 A. Paulsen, H. Liskien, F. Arnotte, and R. Widera, *Nucl. Sci. and Eng.* **78**, 377 (1981).
- PR76 A. Prince, *Evaluation of Chromium Neutron and Gamma Production Cross Sections for ENDF/IV*, BNL-NCS-50593, ENDF-246 (August 1976).
- RO89 G. Rohr, R. Shelley, A. Brusegan, F. Poortmans, and L. Mewissen, *Phys. Rev.* **C39**, 426 (1989).
- SA72 O. A. Salnikov, G. N. Lovchikova, G. V. Kotelnikova, A. M. Trufanov, and N. I. Fetisov, *Differential Cross Sections of Inelastic Scattering Neutrons on Nuclei Cr, Mn, Fe, Co, Ni, Cu, Y, Zr, Nb, W, Bi*, Report Jadernye Konstanty -7, 102 (March 1972).
- SH86 K. Shibata and C. Y. Fu, *Recent Improvements of the TNG Statistical Model Code*, ORNL/TM-10093 (August 1986).
- SH87 K. Shibata and D. M. Hetrick, *Calculated Neutron-Induced Cross Sections for ^{53}Cr from 1 to 20 MeV*, ORNL/TM-10381 (1987).
- SM80 D. L. Smith and J. W. Meadows, *Nucl. Sci. and Eng.* **76**, 43 (September 1980).
- TA65 S. Tagesen and P. Hille, *Oesterr. Akad. Wiss., Math. Naturwiss. Kl., Sitz. Abt.* **2**, 174, 85 (March 1965).
- TA83 A. Takahashi, J. Yamamoto, T. Murakami, K. Oshima, H. Oda, K. Fujimoto, M. Ueda, M. Fukazawa, Y. Yanagi, J. Miyaguchi, and K. Sumita, Oktavian Report A-83-01, Osaka University, Japan (June 1983).
- VA62 D. M. Van Patter, N. Nath, S. M. Shafroth, S. S. Malik, and M. A. Rothman, *Phys. Rev.* **128**, 1246 (1962).
- VO80 H. Vonach, A. Chalupka, F. Wenninger, and G. Staffel, "Measurement of the Angle-Integrated Secondary Neutron Spectra from Interaction of 14 MeV Neutrons with Medium and Heavy Nuclei," *Proc. Symp. on Neutron Cross-Sections from 10 to 50 MeV*, BNL-NCS-51245, Brookhaven National Laboratory (July 1980).

Fig. 1. Comparison of evaluated and experimental total inelastic scattering cross sections for ^{52}Cr .

Fig. 2. Comparison of calculated and experimental neutron production cross sections as a function of angle for several outgoing neutron energy bins.

Fig. 3. Neutron emission spectra from the TNG calculation for ^{53}Cr at 14.5 MeV, used for ENDF/B-VI. Contributions from the various neutron-producing components are shown (they sum to the total). The curve labeled (n, np) includes the (n, pn) component. The (n, n α) and (n, α n) components were also calculated, but fall below the bottom of the plot.

Fig. 4. Neutron emission spectra from ENDF/B-V (line) and ENDF/B-VI (histogram) compared with experimental data. The data of Takahashi et al. (TA83) were taken at 80° , and the other measured data sets shown (HE74, VO80, and SA72) are angle integrated. The data are for natural chromium, and the isotopic evaluations have been combined to form the evaluated curve for ENDF/B-VI.

Fig. 5. Comparison of the ENDF/B-VI proton production spectra for ^{52}Cr compared with experimental data. The measurements were taken at incident energies of 14.8 and 14.1 MeV, the TNG calculation used for the evaluation was for $E_n = 14.5$ MeV. The data of Grimes et al. (GR79, JA77) are angle integrated; the data of Colli et al. (CO62) were taken at 15° .

Fig. 6. Comparison of ENDF/B-VI and experimental alpha production spectra for ^{52}Cr . The measurement was taken at an incident energy of 14.8 MeV, the TNG calculation used for the evaluation was for $E_n = 14.5$ MeV.

Fig. 7. Secondary gamma-ray production cross section versus gamma-ray energy from ENDF/B-VI (incident energy $E_n = 9.5$ MeV) compared with the data of Morgan and Newman (MO76).

⁵⁵₂₅Mn

Reference: No Primary Reference
Evaluator: K. Shibata, JAERI, ORNL
Evaluated: March 1988
Material: 2525
Content: Neutron transport, Gamma production, Covariances

File Comments

MF=1 General information.
 MT=451 Descriptive data and dictionary.

MF=2 Resonance parameters.
 MT=151 The resolved resonance parameters are in the MLBW formalism. The parameters of the lowest four resonances were taken from the work of Macklin /1/. Others were taken from the compilation of Mughabghab et al. /2/ except that the parameters of two negative resonances were adjusted so as to fit to experimental thermal cross sections.
 Resonance region : 10^{-5} eV to 100 keV.
 Scattering radius: 5.15 fm.
 Calculated 2200-m/s cross sections and res. integrals.

	2200-m/s	Res. Integ.
Elastic	2.167 b	-
Capture	13.413 b	11.73 b
Total	15.579 b	-

MF=3 Neutron cross sections.
 MT=1 Total.
 Below 100 keV: No background.
 Above 100 keV: Based on the experimental data /3,4,5/.
 MT=2 Elastic scattering. Defined as the total σ minus the nonelastic cross section.
 MT=3 Non-elastic cross section. Defined as the sum of MT=4, 16, 22, 28, 102, 103, 104, 105, 106 and 107.
 MT=4, 51-79 and 91. Inelastic scattering cross section. Statistical-model calculations were performed using TNG /6/. The precompound process was considered above 5 MeV.

MF=3

Continued. The calculated cross section for MT=51 was multiplied by a factor of 1.2. For the levels of MT=51, 52, 57, 61, 64, 65, 67, and 70, direct process components were taken into account through the DWBA calculations. The optical potential parameters used were as follows - (in units of MeV and fm) /7/ :

$$V = 49.747 - 0.4295 \times E - 0.0003 \times E^2 \quad R0 = 1.287 \quad A0 = 0.56$$

$$WS = 11.2 - 0.09 \times E \quad RS = 1.345 \quad AS = 0.47$$

$$VS0 = 6.2 \quad RS0 = 1.120 \quad AS0 = 0.47$$

The level scheme was taken from Ref. /8/.

No.	Energy(MeV)	Spin-parity
g.s.	0.0	5/2 -
1.	0.126	7/2 -
2.	0.984	9/2 -
3.	1.290	1/2 -
4.	1.292	11/2 -
5.	1.293	1/2 -
6.	1.528	3/2 -
7.	1.884	7/2 -
8.	2.015	7/2 -
9.	2.198	7/2 -
10.	2.215	5/2 -
11.	2.252	3/2 -
12.	2.267	5/2 -
13.	2.312	13/2 -
14.	2.366	5/2 -
15.	2.398	9/2 +
16.	2.427	1/2 +
17.	2.563	3/2 -
18.	2.727	7/2 -
19.	2.753	5/2 -
20.	2.822	9/2 -
21.	2.824	5/2 -
22.	2.873	1/2 -
23.	2.954	3/2 -
24.	2.976	3/2 -
25.	2.992	7/2 -
26.	3.006	3/2 -
27.	3.036	11/2 -
28.	3.038	1/2 -
29.	3.040	3/2 +

Levels above 3.046 mev were assumed to be overlapping.

MF=3 MT=16, 22, 28, 103 and 107. (n,2n), (n,n' α), (n,n'p), (n,p) and (n, α) cross sections were calculated using TNG. Global optical-potential parameters were employed for protons and alpha-particles /9,10/.

MT=102 Radiative capture cross section.
Below 100 keV: resonance parameters given (no background).
Above 100 keV: based on experimental data /11/-/15/.
(Taken from JENDL-2 /20/)

MT=104 (n,d) cross section.
The excitation function of the (n,p) cross section calculated with TNG was used for the (n,d) reaction by shifting the threshold energy. The cross sections were normalized to the experimental datum at 14.1 MeV /16/.

MT=105 (n,t) cross section.
The excitation function of the (n,p) cross section calculated with TNG was used for the (n,t) reaction by shifting the threshold energy. the cross sections were normalized to the experimental datum at 14.7 MeV /17/.

MT=106 (n, ^3He) cross section.
This cross section is based on experimental data /18,19/.

MF=4 MT=2 Elastic angular distributions of secondary neutrons were obtained from optical and statistical-model calculations.

MF=6 Energy-angle distributions.

MT=16 (n,2n) energy distributions of neutrons, recoils and gamma-rays were calculated using TNG. Angular distributions are given for neutrons only.

MT=22 (n,n' α) energy distributions of neutrons, alphas, recoils and gamma-rays were calculated using TNG. Angular distributions are given for neutrons only.

MT=28 (n,n'p) energy distributions of neutrons, protons, recoils and gamma-rays were calculated using TNG. Angular distributions are given for neutrons only.

MT=51 52... 79. Inelastic scattering to discrete levels. Optical and statistical-model calculations were used. The components of the direct-interaction process were added to the levels of MT=51,52,57,61,64,65,67 and 70 using DWBA calculations.

MF=6	MT=91	Inelastic scattering to the continuum levels. Energy distributions of neutrons, recoils and gamma-rays were calculated using TNG. Angular distributions are given for neutrons.
	MT=103	(n,p) distributions. The energy distributions of protons, recoils and gamma-rays were calculated using TNG.
	MT=107	(n, α) distributions. The energy distributions of alphas, recoils and gamma-rays were calculated using TNG.
MF=12		Photon production multiplicities.
	MT=51	52... 79. Branching ratios for the levels are given.
	MT=102	Capture. Calculated with TNG.
MF=14		Photon angular distributions.
	MT=51	52... 79. Assumed to be isotropic.
	MT=102	Capture. Assumed to be isotropic.
MF=15		Photon energy distributions.
	MT=102	Capture. Calculated with TNG.
MF=33		Covariance data.
	MT=1,	2, 3, 4, 16, 22, 28, 51-79, 91, 102, 103, 104, 105, 106, and 107 estimated from experimental data and calculation.

References:

1. Macklin, R. L., Nucl. Sci. Eng., 89, 362 (1985).
2. Mugabghab, S. F., Divadeenam, M. and Holden, N. E., "Neutron Cross Sections", Vol. 1, Part A, Academic Press (1981).
3. Cierjacks, S., Forti, P., Kopsch, D., Kropp, L., Nebe, J. and Unseld, H., "High Resolution Total Cross Sections for Na, Cl, K, V, Mn and Co between 0.5 and 30 MeV", KFK-1000 (1968).
4. Pineo, W. F. E., Divadeenam, M., Bilpuch, E. G., Seth, K. K. and Newson, H. W., Ann. Phys., 84, 165 (1974).
5. Garg, J. B., Rainwater, J. and Havens, Jr., W. W., Nucl. Sci. Eng., 65, 76 (1978).
6. Fu, C. Y., "A Consistent Nuclear Model for Compound and Precompound Reactions with Conservation of Angular Momentum", ORNL/TM-7042 (1980).

7. Fu, C. Y., Private Communication (1985).
8. Zhou Enchen, Huo Junde, Zhou Chunmei, Lu Xiane and Wang Lizheng, Nucl. Data Sheets, 44, 463 (1985).
9. Perey, F. G., Phys. Rev., 131, 745 (1963).
10. Huizenga, J. R. and Igo, G. J., Nucl. Phys., 29, 462 (1962).
11. Garg, J. B., Macklin, R. L. and Halperin, J., Phys. Rev., C18, 2079 (1978).
12. Dovbenko, A. G., Kolesov, V. E., Koroleva, V. P., Tolstikov, V. A., Atom. Energ., 26, 67 (1969).
13. Menlove, H. O., Coop, K. L., Grench, H. A. and Sher, R., Phys. Rev., 163, 1299 (1967).
14. Schwerer, O., Winkler-Rohatsch, M., Warhanek, H. and Winkler, G., Nucl. Phys., A264, 105 (1976).
15. Budnar, M., Cvelbar, F., Hodgson, E., Ludoklin, A., Ivkovic, V., Likar, A., Mihailovic, M. V., Martincic, R., Najzer, M., Perdan, A., Potokar, M. and Ramsak, V., "Prompt Gamma-ray Spectra and Integrated Cross Sections for the Radiative Capture of 14 MeV Neutrons for 28 Natural Targets in the Mass Region from 12 to 208", INDC(YUG)-6 (1979).
16. Colli, L., Iori, I., Micheletti, S. and Pignanelli, M., Nuovo Cimento, 21, 966 (1962).
17. Sudar, S. and Csikai, J., Nucl. Phys., A319, 157 (1979).
18. Diksic, M., Strohal, P. and Slaus, I., Journal Inorg. Nucl. Chem., 36, 477 (1974).
19. Wu, C. H., Woelfle, R. and Qaim, S. M., Nucl. Phys., A329, 63 (1979).
20. Nakagawa, T., "Summary of JENDL-2 General Purpose File", JAERI-M 84-103 (1984).

DESCRIPTION OF EVALUATIONS FOR $^{54,56,57,58}\text{Fe}$ PERFORMED FOR ENDF/B-VI†

C. Y. Fu, D. M. Hetrick, C. M. Perey,
F. G. Perey, N. M. Larson, and D. C. Larson
Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6356, U. S. A.

ABSTRACT

Isotopic evaluations for $^{54,56,57,58}\text{Fe}$ performed for ENDF/B-VI are briefly reviewed. The evaluations are based on analysis of experimental data and results of model calculations which reproduce the experimental data, including data for the isotopes and the natural element. Evaluated data are given for neutron-induced reaction cross sections, angular and energy distributions of secondary particles, and gamma-ray production cross sections associated with the reactions. File 6 formats are used to represent energy-angle correlated data and recoil spectra. Uncertainty files are included for all File 3 cross sections. A detailed description of the evaluation is given for ^{56}Fe and results of calculations for the major reactions are used for evaluations of the minor isotopes, with particular attention paid to inelastic scattering to the low-lying levels in ^{57}Fe .

1. INTRODUCTION

Separate evaluations have been done for each of the stable isotopes of iron. In this report, we briefly review the structure of the evaluations, describe how the evaluations were done, and note the major pieces of data considered in the evaluation process. Experimental data references were obtained primarily from CINDA; the data themselves were mostly obtained from the National Nuclear Data Center at Brookhaven National Laboratory and, occasionally, from the literature and reports. The R-Matrix code SAMMY (LA89) was used for the resonance region analysis. The TNG nuclear model code (FU88, SH86), a multistep Hauser-Feshbach code which includes precompound and compound contributions to cross sections, angular, and energy distributions in a self-consistent manner, calculates gamma-ray production, and conserves angular momentum in all steps, was the primary code used for these evaluations. Extensive model calculations were performed with the goal of simultaneously reproducing experimental data for all reaction channels with one set of parameters. This ensures internal consistency and energy conservation within the

†Research sponsored by the Office of Energy Research, Nuclear Physics, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

evaluation. In the case of reactions for which sufficient data were available, a Bayesian analysis using the GLUCS code (HE80) was frequently done, using ENDF/B-V or the TNG results as the prior. In cases where insufficient data were available for a GLUCS analysis, and the available data were deemed to be accurate, but in disagreement with the TNG results, a smoothed-curve representation of the data was used for the evaluation. A similar method was also used for cross sections where resonant structure was felt to be important, but resonance parameters were not used. The final evaluation is thus a combination of TNG results, GLUCS results, and smoothed curves. Since iron is a very important material in fission and fusion reactors, much effort has gone into its evaluation (and analysis of the evaluations) over the past years, and results of this effort are reflected in the fine tuning done for ENDF/B-VI.

In Section 2 the resonance parameters are discussed; Section 3 contains a description of the major cross sections included in the evaluation; Section 4 is devoted to angular distributions; Section 5 to energy-angle correlated distributions; Section 6 to gamma-ray production cross sections; Section 7 describes the uncertainty files; and Section 8 is on needs of important data and possible ways to improve the evaluation.

Much of this information is abstracted from Refs. FU80, FU82, FU86, and FU88.

2. RESONANCE PARAMETERS

Resonance parameters for ^{56}Fe were taken from an evaluation up to 850 keV by Perey et al. (PE90). Recent high resolution transmission data (PE90) for ^{56}Fe were analyzed with the R-Matrix code SAMMY (LA89). Capture data of (CO84) were used to determine the radiative widths. Elastic scattering data of (PE90) were analyzed with the code RFUNC (PE89) to aid in the spin and parity assignments. An example is shown in Fig. 1 where the measured transmission data and elastic angular distributions jointly determine the spin and parity assignments of several resonances.

Resonance parameters for the other isotopes were mostly based on those compiled by Mughabghab (MU81) but were re-analyzed by the SAMMY code to achieve a better fit to the available data than shown by Mughabghab.

For all isotopes, total cross-section data were fitted well enough that background cross sections were not needed in File 3. However, because many capture resonances could not be seen in the transmission data, background cross sections were used in File 3 so that the sum of the resonance capture cross sections and File 3 data agreed with the available data (MU81). For example, capture resonances in ^{56}Fe included in the SAMMY analysis appeared complete up to 400 keV, but not so between 400 and 850 keV, in which a File 3 cross section was used based on the data of Allen et al. (AL76).

3. CROSS SECTIONS

In this section we briefly describe contents of the files containing cross sections for the more important reactions. The total cross section for the minor isotopes above the resonance region was taken from an ORELA measurement for natural iron. For ^{56}Fe ,

results of an ORELA high-resolution transmission measurement for this isotope (which covered the energy range from 5 keV to 40 MeV and was used for the resonance parameter analysis) were used to 2 MeV, below which the isotopic and natural-iron data were in excellent agreement. From 2 to 20 MeV, the natural-iron transmission data were still used for ^{56}Fe because these data were judged to be more reliable. There are many data sets available for inelastic scattering to the first excited level in ^{56}Fe at 847 keV, including high resolution data showing resonant structure. Below 2.2 MeV the evaluation is based on the high-resolution data of Kinney and Perey (KI77) and Voss (VO72). Above 2.2 MeV several data sets were used, and the evaluated curve was adjusted within experimental uncertainties based on integral results and a consistent nuclear model analysis with TNG. The inelastic scattering cross sections for higher-lying levels in ^{56}Fe are based on a TNG analysis of experimental data and include a direct-reaction contribution to many of the levels. The total inelastic scattering cross section for natural iron, summed from the isotopic evaluations, is compared in Fig. 2 with the data of Larson (LA85) and with ENDF/B-V.

Inelastic scattering to the discrete levels of $^{54,57,58}\text{Fe}$ is also included in evaluations of these isotopes. In particular, inelastic scattering to the first three levels in ^{57}Fe occurs at neutron energies below the first level in ^{56}Fe and is known to be an important energy loss mechanism when neutrons of energies less than 1 MeV impinge upon natural iron. Above the discrete levels, inelastic scattering is represented by a continuum given in File 3 Section 91. Precompound effects are included both in the cross section and the angular and energy distributions.

The $^{56}\text{Fe}(n,p)$ reaction has several data sets available, and a GLUCS analysis was done for 14 correlated reactions which included this reaction. Results of that analysis were used in the evaluation of this cross section. This cross section was also well reproduced by the TNG results. Evaluation of the (n,p) reaction for the minor isotopes was taken from TNG results. There are no measurements of the $^{56}\text{Fe}(n,\alpha)$ cross section via activation, since the product ^{53}Cr is stable. Thus, this cross section is taken from results of the TNG analysis, which is in good agreement with measured alpha production spectra at 14 MeV, and agrees well with measured fission-spectrum averages. Cross sections for the (n,α) reaction on the minor isotopes were also taken from the TNG calculations.

Experimental data for the $(n,2n)$ reaction are available for both ^{56}Fe and natural iron. Figure 3 shows results of the TNG calculations for ^{56}Fe compared with the available data and with earlier evaluations. The TNG calculated result is used for the present evaluation.

Cross sections for important tertiary reactions are given for all four isotopes, based on results of the TNG calculations.

4. ANGULAR DISTRIBUTIONS

High-resolution differential elastic scattering data is available up to 1.23 MeV (KI77) and is used for the evaluation. A Legendre polynomial representation in the center-of-mass system is used. From 1.23 to 20 MeV the evaluation is based on a number of data sets and results of optical model calculations (FU80). Angular distributions are given for inelastic scattering to each level in ^{56}Fe , are taken from the TNG calculations, and most include a

direct interaction component. Inelastic scattering to discrete levels in the minor isotopes were done in the same manner. Angular distributions for inelastic scattering from the continuum and for other neutron-producing reactions are correlated with outgoing energy and given in File 6.

5. ENERGY-ANGLE CORRELATED DISTRIBUTIONS (FILE 6)

Perhaps the major improvement in the iron evaluation for ENDF/B-VI is the use of correlated energy-angle distributions for the outgoing neutrons, obtained from TNG calculations. A revision to ENDF/B-V, incorporating these correlations for inelastic scattering from the continuum (crudely, however, due to format limitations), was an important factor in improving agreement of calculated results with measured values for several reactor problems. ENDF/B-VI uses these results, translated into File 6 formats which allow for more precise representation of the evaluated distributions. Energy-angle correlations in the secondary neutron spectra from the $(n,2n)$, (n,np) , and $(n,n\alpha)$ reactions are also incorporated, using File 6. Variable bin widths (SH86) have been used for the neutron emission spectra to faithfully represent the peak area of the spectra. These correlations are particularly important since it is known that the angular distributions vary strongly with outgoing neutron energy (FU86). Energy-angle correlations were also used for the secondary neutrons from the $(n,\alpha n)$ and (n,pn) reactions. All File 6 evaluated data are taken from the TNG calculations, which have been compared with available data. Figure 4 shows a comparison of the TNG results with available neutron emission data as a function of angle for four outgoing neutron energy bins. Figure 5 shows a comparison of several measured neutron emission spectra with the ENDF/B-VI results, obtained from summing the isotopic contributions.

Charged-particle and recoil spectra are also given in File 6, with isotropic angular distributions assumed. Figure 6 shows a comparison of the measured proton production spectrum for ^{56}Fe of Grimes et al. (GR79) with the evaluation for ENDF/B-VI. Figure 7 shows a similar comparison for the alpha production spectrum. In both cases, the calculated spectra are in good agreement with measured spectra.

As an example of the usage of File 6, consider the $^{56}\text{Fe}(n,n\alpha)^{52}\text{Cr}$ reaction. In File 6/22, constant yields are given for the outgoing neutron, alpha and ^{52}Cr residual, and an energy dependent yield is used for the gamma rays associated with the $(n,n\alpha)$ reaction. Normalized energy distributions are given for each outgoing product, but only the outgoing neutron has a non-isotropic angular distribution. The sum of the average energies for the outgoing neutrons, alpha-particles, heavy recoils, and gamma rays was checked to agree within 1% of the incident neutron energy. The cross section to be used for normalization is taken from File 3/22.

6. GAMMA-RAY-PRODUCTION CROSS SECTIONS

One of the major problems with earlier ENDF/B evaluations was a lack of energy balance between gamma rays and particle production which, between a few MeV and 15

MeV, have comparable average energies. For ENDF/B-VI an effort has been made to alleviate this difficulty. Often the problem has arisen because experimentally measured gamma-ray-production spectra have been used directly in evaluations, without checking for energy balance with other outgoing particles. A typical case existed for iron, where three major sets of gamma-ray production data existed as a function of neutron energy: that of Orphan et al. (OR75), Dickens et al. (DI72), and Chapman et al. (CH76). However, the measured data sets disagreed by as much as a factor of two at some energies. A careful evaluation of the data, based on a TNG analysis which attempted to simultaneously fit data for all neutron-induced reactions as a function of incident energy, showed little room for changing neutron production cross sections, secondary neutron energy distributions, or secondary gamma-ray energy distributions. Results of these calculations produced gamma-ray-production cross sections that, while not in good agreement with any single measurement over the entire energy range, were in reasonable average agreement and removed the energy balance problem. Results from TNG calculations were used for the evaluation of all four isotopes. Some calculated gamma-ray production spectra for ^{56}Fe are compared with the experimental data (DI72, CH76, DR78) in Figs. 8-11. Note these spectra are the sum from all reactions, while in the evaluation, the spectra are broken down into individual ones for each reaction. Therefore the energy balance is maintained for each reaction and the cross section for any reaction can be altered from the calculated values, without upsetting the energy balance, e. g. the $^{56}\text{Fe}(n,p)$ cross section was taken from the GLUCS analysis.

Gamma rays originating from discrete inelastic scattering are given by branching ratios for each inelastic level cross section for exact energy balance. The part of the discrete gamma rays, arising from deexcitations from the continuum inelastic levels, are separately given in section 91. This allows checking of energy balance for section 91. With this separation of gamma-rays produced in sections 51-90 (the discrete inelastic levels) from those produced in section 91 achieved, all reactions have their own gamma-ray production spectra which could not be done with past evaluations since gamma-ray-production spectra were given only for the nonelastic reaction. File 6 was also chosen to represent the continuum gamma-ray production energy distributions to ensure consistency with the neutron and charged-particle distributions. Isotropic angular distributions were assumed for the gamma rays.

Capture gamma-ray yields and distributions are given in Files 12 and 15, respectively, and are based on an old evaluation for natural iron (WH73). This is an area that requires improvement; see Section 8 for details.

7. UNCERTAINTY INFORMATION

Uncertainty files are given only for the cross sections in File 3. They are not given for energy distributions or angular distributions. Fractional and absolute components, correlated only within a given energy interval, are based on scatter in experimental data and estimates of uncertainties associated with the model calculations.

8. DATA NEEDS AND EVALUATION IMPROVEMENTS

The $^{56}\text{Fe}(n,n')$ cross section exciting the 0.847-MeV level up to an incident neutron energy of 5 MeV is known to about 5-10%, but an accuracy of 2% is needed for reactor pressure vessel surveillance dosimetry. The capture cross section data uncertainties may be as much as 25% in the keV region as shown for the 1.15-keV resonance in ^{56}Fe by an International Task Force (P89A). Hence, new high resolution capture data are needed in the resonance region, at least for $^{56,58}\text{Fe}$. $^{58}\text{Fe}(n,\gamma)$ is still being used for reactor dosimetry, but the existing data base used for ENDF/B-VI is very poor. High-quality data are needed for the lowest 10 s-wave resonances, particularly the radiative widths. $^{56}\text{Fe}(n,\alpha)$ requires measurements other than 14 MeV because calculations used for ENDF/B-VI, JENDL-3, and EFF-2 evaluations do not agree below and above 14 MeV. Double differential neutron emission cross sections for natural iron are needed at incident energies other than 14 MeV to benchmark the model calculations.

A known deficiency exists in the capture gamma-ray spectra. These spectra were taken from an old evaluation (WH73). In this evaluation, experimental capture gamma-ray branching ratios for s-wave and p-wave resonances in $^{54,56}\text{Fe}$ were separately fitted and extrapolated by model calculations. The fitted branching ratios, as a function of incident neutron energy, were combined for natural iron using R-Matrix analysis for the resonance capture cross sections. The evaluated data were then grouped in such a way that the characteristics of the capture spectrum of each dominant resonance in $^{54,56}\text{Fe}$ up to 1 MeV were separately represented. This is a good approach because the capture Q-values of the two isotopes differ by 1.7 MeV. These results for natural iron were used in ENDF/B-VI for each isotope of iron by adjusting the multiplicities to conserve energy. This is an area that requires improvement. Newer experimental capture spectra should be analyzed and supplemented by TNG calculations and given separately for each isotope.

Uncertainties should be given for important resonance parameters, and angular and energy distributions.

REFERENCES

- AL76 B. J. Allen et al., Nucl. Phys. A469, 408 (1976).
- AS58 V. J. Ashby, H. C. Catron, and L. L. Newkirk, Phys. Rev. 111, 616 (1958).
- CH76 G. T. Chapman, G. L. Morgan, and F. G. Percy, "A Re-measurement of the Neutron-Induced Gamma-Ray Productions Cross Sections for Iron in the Energy Range 850 keV $< E_n < 20.0$ MeV," Oak Ridge National Laboratory, ORNL/TM-5416 (1976).
- CL72 G. Clayeux and J. Voignier, "Diffusion Non Elastique de Neutrons de 14 MeV sur Mg, Al, Si, S, Ca, Sc, Fe, Ni, Cu, Au, Pb, et Bi," Centre d'Etudes de Limeil CEA-R-4279 (1972).
- CO78 V. Corcalciuc, B. Holmqvist, A. Marcinkowski, and G. A. Prokopets, Nucl. Phys. A307(3), 445 (1978).
- CO84 F. Corvi et al., Proc. Consultants Meeting on Nuclear Data for Structural Materials, Vienna, Austria, 1983, International Nuclear Data Committee Report INDC(NDS)-152 L, 1984 and private communication to C. M. Percy, 1984.

- DI72 J. K. Dickens, G. L. Morgan, and F. G. Perey, "Gamma-Ray Production Due to Neutron Interactions with Iron for Incident Neutron Energies Between 0.8 and 20 MeV: Tabulated Differential Cross Sections," ORNL-4798, Oak Ridge National Laboratory, 1972.
- DR78 D. M. Drake et al., Nucl. Sci. Eng. 40, 294 (1978).
- FR80 J. Frehaut, A. Bertin, R. Bois, and J. Jary, Symp. on Neutron Cross Sections from 10-50 MeV, Brookhaven National Laboratory, May 12-14, 1980.
- FU80 C. Y. Fu and F. G. Perey, "Evaluation of Neutron and Gamma-Ray Production Cross Sections for Natural Iron (ENDF/B-V MAT 1326)," ORNL/TM-7523, ENDF-302, Oak Ridge National Laboratory, 1980.
- FU82 C. Y. Fu, "Summary of ENDF/B-V Evaluations for Carbon, Calcium, Iron, Copper, and Lead and ENDF/B-V Mod-3 for Calcium and Iron," ORNL/TM-8283, ENDF-325, Oak Ridge National Laboratory, 1982.
- FU86 C. Y. Fu and D. M. Hetrick, "Update of ENDF/B-V Mod-3 Iron: Neutron-Producing Reaction Cross Sections and Energy-Angle Correlations," ORNL/TM-9964, ENDF-341 (July 1986).
- FU88 C. Y. Fu, Nucl. Sci. Eng. 100, 61 (1988).
- GR79 S. M. Grimes, R. C. Haight, K. R. Alvar, H. H. Barschall, and R. R. Borchers, Phys. Rev. C19, 2127 (1979).
- HE75 D. Hermsdorf et al., "Differential Neutron Emission Cross Sections from Iron by 14.6-MeV Neutrons," ZfK-277, Institute of Nuclear Physics, University of Dresden (East Germany), 1975.
- HE80 D. M. Hetrick and C. Y. Fu, "GLUCS: A Generalized Least-Squares Program for Updating Cross Section Evaluations with Correlated Data Sets," ORNL/TM-7341, ENDF-303 (October 1980).
- KA72 J. L. Kammerdiener, "Neutron Spectra Emitted by ^{239}Pu , ^{238}U , ^{235}U , Pb, Nb, Ni, Fe, Al, and C Irradiated by 14 MeV Neutrons," UCRL-51232, Lawrence Livermore Laboratory, 1972.
- KI77 W. E. Kinney and F. G. Perey, Nucl. Sci. Eng. 63, 418 (1977).
- LA85 D. C. Larson, "High-Resolution Structural Material (n,x γ) Production Cross Sections for E_n from 0.2 to 40 MeV," p.71, Proc. Int. Conf. Nucl. Data for Basic and Applied Sciences, Santa Fe, May 13-17, 1985.
- LA89 N. M. Larson and F. G. Perey, "Users' Guide for SAMMY: A Computer Model for Multi-Level R-Matrix Fits to Neutron Data Using Bayes' Equations," ORNL/TM-7485 (1980); Updates ORNL/TM-9179 (1984), ORNL/TM-9179/R1 (1985) and /R2 (1989).
- MU81 S. F. Mughabghab et al., Neutron Resonance Parameters and Thermal Cross Sections, Academic Press, 1981.
- OR75 V. G. Orphan, C. G. Hoot, and V. C. Rogers, Nucl. Sci. Eng. 57, 309 (1975).
- PE89 F. G. Perey, "RFUNC - A Code to Analyze Differential Elastic-Scattering Data," ORNL/TM-11112 (1989).
- P89A F. G. Perey, Oak Ridge National Laboratory, private communication (1989).
- PE90 C. M. Perey et al, " ^{56}Fe Resonance Parameters for Neutron Energies up to 850 keV," Oak Ridge National Laboratory, ORNL/TM-11742 (1990).
- QA77 S. M. Qaim and N. I. Molla, Nucl. Phys. A283, 269 (1977).

- SA72 O. A. Salmikov, G. N. Lovchikova, G. V. Kotelnikova, A. M. Trufanov, and N. I. Fetisov, *Jadernye Konstanty* - 7, 102 (1972).
- SH86 K. Shibata and C. Y. Fu, "Recent Improvements of the TNG Statistical Model Code," Oak Ridge National Laboratory, ORNL/TM-10093 (1986).
- TA83 A. Takahashi et al., "Double Differential Neutron Emission Cross Sections, Numerical Tables and Figures (1983)," A-83-01, Intense 14-MeV Neutron Source Facility, Osaka University, 1983.
- VE79 L. R. Veaser, Figure 61 in "Evaluated Neutron-Induced Cross Sections for $^{54,56}\text{Fe}$ to 40 MeV," by E. D. Arthur and P. G. Young, Los Alamo National Laboratory, LA-8626-MS, ENDF-304 (1980).
- VO72 F. Voss, "Measurement and Fluctuation Analysis of γ -Ray Production Cross Sections After Inelastic Neutron Scattering on ^{56}Fe and ^{27}Al between 0.8 and 13 MeV," Karlsruhe Nuclear Research Center, KFK-1611 (1972).
- WH73 J. E. White, C. Y. Fu, and K. J. Yost, *Nucl. Sci. Eng.* 51, 496 (1973).
- WE62 R. Wenusch and H. Vonach, *Anz. Oesterr. Akad. Wiss., Math-Naturwiss. Kl.* 99 1 (1962).

Fig. 1. In the upper graph the theoretical total cross sections, calculated with the parameters of ENDF/B-VI ^{56}Fe , are compared with the data (PE90). Parentheses are used to indicate uncertain spin and parity assignments. The assignments were made using the differential elastic scattering data (PE90) shown in the three lower graphs.

nat-Fe TOTAL INELASTIC SCATTERING CROSS SECTION

Fig. 2. Comparison of the Fe inelastic cross sections of ENDF/B-VI, summed from those in the isotopic evaluations, with ENDF/B-V and with experimental data of Larson (LA85).

Fig. 3. Comparison of the ENDF/B-VI $^{56}\text{Fe}(n,2n)$ cross sections with experimental data and with the natural-iron $(n,2n)$ in ENDF/B-V.

Fig. 4. Comparison of experimental (HE75, KA72, TA83) double differential Fe(n,xn) cross sections for 14.5-MeV incident neutrons and several secondary energy ranges with the TNG calculation used for ENDF/B-VI.

Fig. 5. Comparison of the neutron production spectrum of ENDF/B-VI natural iron, obtained from summing the isotopic components, with measurements.

Fig. 6. Comparison of the proton production spectrum of ENDF/B-VI ^{56}Fe with measurements.

Fig. 7. Comparison of the alpha-particle production spectrum of ENDF/B-VI ^{56}Fe with measurements.

Fig. 8. Comparison of experimental gamma-ray production spectrum at $E_n = 4.25$ MeV for natural iron (CH76) with the TNG calculation used for ENDF/B-VI.

Fig. 9. Comparison of experimental gamma-ray production spectrum at $E_n = 9.25$ MeV for natural iron (CH76) with the TNG calculation used for ENDF/B-VI.

Fig. 10. Comparison of experimental gamma-ray production spectrum at $E_n = 14.1$ MeV for natural iron (DI72) with the TNG calculation used for ENDF/B-VI.

Fig. 11. Comparison of experimental gamma-ray production spectrum at $E_n = 14.1$ MeV for natural iron (DR78) with the TNG calculation used for ENDF/B-VI.

⁵⁶₂₆Fe

Reference: The Astrophysical Journal, 346, 1049-1060 (1989)

Authors: S. Pearlstein (BNL)

Evaluation: August 1989

Material: 2631

Content: Neutron transport, Gamma production

Library: High Energy Library (NLIB=4)

Sublibrary: Incident-Neutron Data (NSUB=10), Incident Proton Data (NSUB=10010)

Content: Neutron transport, Gamma production for NSUB=10
Proton transport, Gamma production for NSUB=10010

1. Introduction

The primary objective of this evaluation is a practical file for radiation transport applications which reasonably summarizes the contemporary physical knowledge. This documentation covers both the neutron-incident and proton-incident data for ⁵⁶Fe in the ENDF/B-VI High Energy library.

2. General Method

The nuclear model code ALICE-P, a modification by S. Pearlstein (Ref. 6) of ALICE85 (Ref. 5), is extensively used. ALICE-P uses parameters in a woods-saxon potential to fit neutron total and particle reaction cross-sections from data and systematics (Refs. 1,14). Tritium and ³He emission are added.

3. Energy Range

The energy range for the neutron induced data is 10⁻⁵eV to 1000 MeV. Below 20 meV the ENDF/B-VI evaluation (Ref. 15) is used. The energy range for the proton-induced data is 1 to 1000 MeV.

4. Total Cross Sections (MT=1, neutron induced data only)

The total cross-section is calculated by ALICE-P.

5. Elastic Scattering Cross Section (MT=2)

The elastic scattering cross-section is calculated by ALICE-P.

Rutherford-nuclear interference is ignored. For protons the elastic scattering cross-section is equal to the total minus the nonelastic cross-section. The elastic scattering angular distributions are based on a diffraction model (Ref. 3) amended for relativistic effects and empirical fits to high energy data.

6. Nonelastic Cross Section (MT=3)

The nonelastic cross section is calculated by ALICE-P. Neutron and proton nonelastic cross-sections are similar at high energies.

7. Spallation Product Yields

The spallation product yields are calculated by ALICE-P. They compare fairly well with experimental data (Refs. 10-13).

8. Neutron Production (MT=201)

The neutron production is calculated by ALICE-P. Double-differential data above 250 MeV are based on systematics (Ref. 4) and are consistent with data (Refs. 7,8). For neutrons below 20 MeV the sum of MT's = 4,2x16,22,28 (Ref. 15) is calculated.

9. Gamma Production (MT=202)

The gamma production is calculated by ALICE-P containing recent modifications (Ref. 9).

10. Proton Production (MT=203)

The proton production is calculated by ALICE-P. For neutrons below 20 MeV the sum of MT's = 28,103 (Ref. 15) is calculated.

11. Deuteron Production (MT=204)

The deuteron production is calculated by ALICE-P. For neutrons below 20 MeV, MT=104 (Ref. 15) is used.

12. Tritium Production (MT=205)

The tritium production is calculated by ALICE-P. For neutrons below 20 MeV, MT = 105 (Ref. 15) is used.

13. ^3He Production (MT=206)

The ^3He production is calculated by ALICE-P. For neutrons below 20 MeV, MT = 106 (Ref. 15) is used.

14. Alpha Production (MT=207)

The alpha production is calculated by ALICE-P. For neutrons below 20 MeV the sum of MT's = 22,107 (Ref. 15) is calculated.

References

1. J. R. Letaw et al, *Astrophys. J. Suppl. Ser.* 51,271 (1983).
2. I. Angeli et al, *Nucl. Phys.* A170,577 (1971).
3. S. Pearlstein, *Nucl. Sci. Eng.* 49,162 (1972).
4. S. Pearlstein, *Nucl. Sci. Eng.* 95,116 (1987).
5. M. Blann et al, *Phys. Rev.* C28,1648 (1983) and LLNL Rep. UCID 20169.
6. S. Pearlstein, *J. Astrophys.* 346,1049 (1989).
7. M. M. Meier et al, *Proc. Santa Fe Conf., Rad. Eff.* II,1415 (1986).
8. D. Filges et al, *Kfk Rep.* Jul-1960 (Nov. 1984).
9. M. Blann et al, *Nucl. Inst. Meth.* A265,490 (1988).
10. B. K. Gupta et al, *Nucl. Phys.* A170,49 (1970).
11. R. Silberberg et al, *Nrl Rep.* 7593 (Dec. 1973).
12. C. J. Orth et al, *J. Inorg. Nucl. Chem.* 28,13 (1976).
13. R. Michel et al, *Nucl. Inst. Meth.* B16,61 (1986).
14. L. Townsend et al, *Rad. Res.* 106,283 (1986).
15. C. Fu et al, *ORNL/TM-9964* (1986) and Updates for ENDF/B-VI.

Graphic Displays

The total and nonelastic cross-sections play an important role in intermediate energy nuclear data. Above 20 MeV for light targets and above a few MeV for heavy targets these cross-sections are smooth and their magnitude systematically well known. The sum of partial reaction cross-sections, even if not individually well known, cannot exceed the nonelastic cross section. The total cross-section for incident charged particles is infinite when Rutherford scattering is integrated over all angles.

As examples of the smooth systematic behavior of neutron induced total and nonelastic cross-sections, comparisons between calculation for ^{56}Fe and experiment for elemental Fe of the total cross-section are shown in Fig. 1(a) and the nonelastic cross-section in Fig. 1(b). As examples of the proton induced data a comparison of calculation and experiment for selected spallation products is given in Fig. 2 and the calculated double-differential neutron emission from 300 meV incident protons is given in Fig. 3. References for experiments given in the figures may be found in the reference publication for these evaluations.

Fig. 1 ^{56}Fe total (a) and nonelastic (b) neutron cross-sections.

Fig. 2 - Comparison of Calculations with experiments of the proton-induced ${}^{56}\text{Fe}$ spallation products ${}^{56}\text{Co}$, ${}^{55}\text{Co}$, ${}^{55}\text{Fe}$, and ${}^{54}\text{Mn}$.

Fig. 3 - Angular-dependent neutron emission spectrum from 300 MeV protons on ${}^{56}\text{Fe}$.

⁵⁹
²⁷Co

Reference: ANL/NDM-107
Evaluators: A. Smith, D. Smith, P. Guenther, J. Meadows, R. Lawson
(ANL), R. Howerton (LLNL) and M. Sugimoto (JAERI).
Evaluated: July 1989
Material: 2725
Content: Neutron transport, Gamma production, Covariances

1. Introduction

The objective of this file is a quantitative neutronic data base for fission and fusion energy applications. Particular attention is given to dosimetry related cross sections. Elemental cobalt is readily available, and provides good dosimetry energy coverage with a single sample through (n, γ) (low-energy), (n,p) and (n, α) (moderate-energy), and (n,2n) (high-energy) reactions.

2. Evaluated Resolved Resonance Range

The present evaluation uses the resonance parameters of Mughabghab¹ to describe the neutron elastic scattering and capture cross sections up to incident neutron energies of 100 keV. A small background was introduced to assure continuity at the matching energy.

3. Evaluated Total Cross Sections

A comprehensive total cross section data base was assembled from the literature as referenced in CINDA.² This data was assembled with values obtained from measurements by the authors³ and from the results of comprehensive measurements by Harvey.⁴ The evaluation above 100 keV was constructed from the above energy averaged data base using the statistical processing code GMA.⁵ Below 5 MeV resonance fluctuations were introduced into the evaluation with the aid of Ref. 4. The fine resolution results of Harvey had to be averaged over narrow energy ranges to reduce the number of energy points in the evaluation to meet ENDF requirements. The experimental data base is now relatively good up to $\approx$ 16 MeV. Additional experimental data would be desirable above that energy.

4. Evaluated Elastic Scattering Cross Sections

The experimental data base relies heavily on the extensive new work of A. B. Smith et al.³ undertaken in support of this evaluation. This data base adequately describes the cross sections in the 200 keV to 10 MeV energy range. Ref. 3 gives detailed attention to the fundamental and applied aspects of the optical-statistical model used to describe the observed elastic scattering. The calculated values were very slightly ($\approx 1\%$) adjusted so that the concurrently calculated total cross sections agreed with the evaluation. The present evaluation is significantly different from ENDF/B-V.

5. Evaluated Inelastic Scattering Cross Sections

5.1 Discrete Inelastic Processes

The present evaluation uses level information taken from Ref. 6., and the calculated inelastic excitation cross sections of Ref. 3. The nuclear model described in Ref. 3 includes direct excitation of 5 of the first 7 levels assuming a weak coupling model. The calculated inelastic cross sections were renormalized slightly to agree with experimental observations. Generally the present evaluation results in 10 - 20% more aggregate inelastic cross section than that of ENDF/B-V and a different shape above 3 MeV.

5.2 Continuum Inelastic Scattering Processes

Continuum cross sections start at 3 MeV. Their magnitudes were determined from the difference between the non-elastic cross sections and the sum of the other partial cross sections. Individual spectra were calculated using the computer codes ALICE⁷ and CADE.⁸

6. Evaluated Radiative Capture Cross Sections

Neutron capture in ^{59}Co results in either 10.5 minute activity of the 2+ state or 5.3 year activity of the ground state (5+).⁹ The latter activity determines the capture cross section essentially. The long half life makes activity measurements tedious, so they don't exist. Direct γ -ray detection methods give knowledge of the capture cross section. The works of Refs. 10 and 11 span the significant energy range, and there are results of direct detection measurements (Refs. 12 and 13) near 14 MeV. The present evaluation relies on calculations using the dipole model of the code ABAREX¹¹ and the optical potential of Ref. 3. A small direct capture component was added at high energies. Above 1 MeV the ENDF/B-V values are considerably larger than the present work, and larger than any experimental result.

7. Evaluated (n,2n) and (n,3n) Reactions

A wealth of data exists for the (n,2n) reaction. This data was used along with the specific 14 MeV evaluation carried out by Evain et al.¹⁵ The least squares evaluation code GMA⁴ was used in the present work. ENDF/B-V was accepted as an a priori input guess to GMA. It provided a shape used to shift the experimental data to the selected grid point energies required by GMA, and it supplemented the available data in regions where measured values were sparse or nonexistent, in order to stabilize the least squares procedures in GMA. The magnitude of ENDF/B-V lies above the general trend of the experimental data, and consequently was reduced everywhere by 7%. This reduction factor was chosen so that the adjusted ENDF/B-V yielded reasonable shape and normalization agreement with the consistent data sets available from Paulsen and Liskien¹⁶ and Frehaut et al.¹⁷ Taken together, these two sets of data span most of the energy span of the evaluation. The present evaluation results fall noticeably below ENDF/B-V (e.g. by $\approx 8\%$ around 16 MeV) over the entire energy range. The uncertainties range from as much as 60% near threshold to as little as 2% in the vicinity of 14 MeV. The results of this evaluation agree very well with that of Evain et al.¹⁵

The threshold of the (n,3n) reaction is 19.353 MeV. There have been no experimental results reported below 21 MeV, and at that energy the cross section is very small, 21 ± 11 mb.¹⁸ In view of the very high threshold energy and the small cross sections the (n,3n) process was not considered in the present evaluation.

8. Evaluated (n, α) Reactions

The ⁵⁹Co (n, α) reaction is negligible below 4 MeV. There exists an extensive data base for this reaction.² The ENDF/B-V values are based on an outmoded 1970 evaluation by Bresesti et al.¹⁹ The details of the evaluation of this reaction has been reported by D. Smith in ANL/NDM-77. The ENDF/B-V evaluation was used as an a priori data set in an adjustment procedure which incorporated differential cross sections obtained at ANL,²⁰ a detailed evaluation from 13 MeV to 15 MeV,²¹ and the results of recent 14 MeV measurements also made at ANL.²² The present evaluation is in better agreement with fission spectrum averaged data^{23,24} than is ENDF/B-V.

9. Evaluated (n,p) Reactions

It became evident that the present evaluation could not be based upon the experimental data. In particular in the 10 to 13 MeV region where the (n,2n) and (n,p) processes compete there is no experimental information. The code GMA⁵ was employed to merge the experimental data and the subjective model information. Below 2.5 MeV CADE was used to calculate a shape which was normalized to ENDF/B-V at

3 MeV where the evaluation represents the data of Smith and Meadows well. Above 13 MeV calculations were made with ALICE⁷ normalized to the measured values in the vicinity of 14 MeV, where the experimental data are both definitive and in reasonable agreement. Data between 10 and 13 MeV were taken from a smooth curve joining the lower (< 10 MeV) and higher (> 13 MeV) regions. The final evaluation is reasonably consistent with the shape predicted by Williams et al.²⁵ The evaluated results of Evian et al.¹⁵ differ significantly from the present values in the vicinity of 14 MeV. The evaluation of Evian carried a large error due to the wide scatter of the experimental results available in 1984.

10. Other Charged Particle Emitting Reactions

The reactions considered appear in Table 1.²⁶ The available experimental data is sparse so the evaluations are heavily dependent upon nuclear model calculations. CADE⁸ which employs the Weisskopf-Ewing assumption,²⁷ and ALICE,⁷ a statistical code which includes pre-compound contributions were used extensively.

Table 1

Q-values and Reaction Products for ⁵⁹Co(n,x) Processes

Reaction	Q-value(MeV)	Product Nucleus	Half Life	Decay Mode
(n,np+pn)	-7.364	⁵⁸ Fe	Stable	
(n,nd+dn)	-15.183	⁵⁷ Fe	Stable	
(n,nt+tn)	-16.573	⁵⁶ Fe	Stable	
(n,n α + α n)	-6.942	⁵⁵ Mn	Stable	
(n,2p)	-12.904	⁵⁸ Mn	65.3 s	β^-
(n,p α + α p)	-8.764	⁵⁵ Cr	3.497 m	β^-
(n,2 α)	-7.566	⁵² V	3.75 m	β^-
(n,d)	-5.139	⁵⁸ Fe	Stable	
(n,t)	-8.927	⁵⁷ Fe	Stable	
(n, ³ He)	-11.600	⁵⁷ Mn	1.45 m	β^-

10.1 (n,n α) + (n, α n) Reaction

This reaction leads to the production of stable ⁵⁵Mn; however, the cross section can be deduced experimentally by comparing the total measured α -production with the production of ⁵⁶Mn via the (n, α) reaction, since no other α -production reactions are of any importance below 20 MeV. Several comparable (n, α x) cross sections have

been reported² and reviewed by Kneff et al.²⁸ Kneff employed mass spectrometric methods to measure helium gas accumulations in pure cobalt samples irradiated with 14.8 MeV neutrons. They measured 40 ± 3 mb for the total α -production cross section. Subtracting 30 mb for the (n,α) cross section yields (10 ± 3) mb. The CADE calculation gave 6.4 mb at 14.8 MeV in fair agreement. The present evaluation was generated by renormalizing the CADE results to the experimental value at 14.8 MeV, as indicated above. The comparable ENDF/B-V cross sections are considerably smaller throughout the energy range, and do not show the broad maximum of the present evaluation near 17 MeV.

10.2 $(n,np) + (n,pn)$ Reaction

This reaction is of significant concern because both experimental and theoretical studies indicate that this process provides a significant fraction of the total proton emission yield at energies of interest for fusion applications.

Most available data has been deduced by the detection of emitted protons at 14.1 MeV. Interpretation of the data is difficult. Derived cross sections appear to be in the range 11 to 60 mb. The CADE and ALICE codes were used in combination to obtain the energy dependent cross sections to 20 MeV. An uncertainty of more than a factor of two is very possible.

10.3 Balance of Charged Particle Emitting Reactions

One data set has been reported for the (n,d) reaction, namely the results of Colli et al.^{29,30} at 14 MeV. Calculated results were in agreement with this measurement and were accepted without alteration.

The (n,t) reaction is of interest because it is the principal tritium producing reaction in cobalt. The present evaluation employs the results of CADE renormalized to agree with the recent relatively precise data of Qaim et al.^{31,32}

The remaining reaction evaluations were all based entirely on nuclear model calculations. There are no comparable files in ENDF/B-V.

11. Evaluated Photon production Reactions

The spectrum of photons from neutron capture was taken from Orphan et al.³³ The same spectrum was used at 20 MeV with the multiplicity adjusted to conserve energy. CASCADE³⁴ was used to determine the energy dependent cross sections for photons resulting from de-excitation of levels excited by inelastic scattering. For all other reactions the R-parameter formalism of Perkins et al.³⁵ was used.

References

1. S. F. Mughabghab, Neutron Cross Sections Vol. 1, Part B, Academic Press Inc. New York, (1984); also S. Mughabghab and C. Dunford, private communication (1982).
2. CINDA, Computerized Index to Nuclear Data, IAEA Press, Vienna (1987).
3. A. B. Smith, P. T. Guenther, R. D. Lawson, and J. F. Whalen, Argonne National Laboratory Report, ANL/NDM-101 (1987). Also Nucl. Phys. A483 50 (1988).
4. J. A. Harvey, Private communication (1986). Data available at the National Nuclear Data Center.
5. W. P. Poenitz, Brookhaven National Laboratory Report, BNL-NCS-51363 Vol. I 249(1981); as modified by M. Sugimoto (1987).
6. P. Anderson, L. Ekstrom, and J. Lyttkens, Nucl. Data Sheets 39 641 (1983) Values given on page 654 used.
7. M. Blann, Lawrence Livermore National Laboratory Report, UCID-20169 (1984).
8. D. Willmore, Harwell Report, AERE-R-11515 (1984).
9. J. Carre and R. Vidal, CEA Report, R2486 (1964).
10. R. Spencer and R. Macklin, Nucl. Sci. and Eng. 61 346 (1976).
11. A. Paulsen, Z. Phys. 205 226 (1967).
12. F. Rigaud et al., Nucl Phys. A173 551 (1974).
13. M. Budnar et al., INDC Report, INDC(YUG) 6 (1979).
14. P. Moldauer, computer code ABAREX, private communication (1982).
15. B. P. Evain et al., Argonne National Laboratory Report, ANL/NDM-89 (1985).
16. A. Paulsen and H. Liskien, J. Nucl. Energy A/B19 907 (1965).
17. J. Frehaut et al., "Proc. Symp. on Neut. Cross Sections from 10-50 MeV, Vol 1," p 399, Brookhaven National Laboratory Report, BNL-NCS-51245 (1980).
18. L. R. Veaser et al., Phys. Rev. C16 1792 (1977).
19. A. Bresesti et al., Nucl. Sci. and Eng. 40 331 (1970).

20. J. W. Meadows, D. L. Smith, and R. D. Lawson, *Ann. Nucl. Energy* 14 603 (1987).
21. R. Spencer and H. Beer, *Bull. Am. Phys. Soc.* 19 574 (1974).
22. J. Meadows, D. Smith, M. Bretscher, and S. Cox, *Ann. Nucl. Energy* 14 489 (1987).
23. D. L. Smith Argonne National Laboratory Report, ANL/NDM-77 (1982).
24. W. Mannhart and A. Fabry, NEANDC(W)-262/U, Vol. 5, p. 58 (1985).
25. J. R. Williams et al., *Proc. Int'l. Conf. on Nucl. Data for Basic and Applied Science*, Santa Fe, Gordon and Breach Publishing Company, New York (1985).
26. J. K. Tuli, Nuclear Wallet Cards, National Nuclear Data Center, Brookhaven National Laboratory (1985).
27. V. F. Weisskopf and D. E. Ewing, *Phys. Rev.* 57 472 (1940).
28. D. W. Kneff, B. M. Oliver, H. Farrar IV, and L. R. Greenwood, *Nucl. Sci. Eng.* 92 491 (1986).
29. L. Colli, I. Iori, S. Micheletti, and M. Pignanelli, *Nuovo Cimento* 20, 94 (1961).
30. L. Colli, I. Iori, S. Micheletti, and M. Pignanelli, *Nucl. Phys.* 46, 73 (1963).
31. S. M. Qaim, R. Woelfe, and H. Liskien, Report INDC(EUR)-13, p. 23, IAEA, Vienna (1980).
32. S. M. Qaim, R. Woelfe, and H. Liskien, *Phys. Rev.* C25, 203 (1982).
33. V. J. Orphan, N. C. Rasmussen, and T. L. Harper, "Line and Continuum γ -ray Yields from Thermal Neutron Capture in 75 Elements," Gulf General Atomic Report, GA-10248/DASA 2570 (1970).
34. W. E. Warren, R. J. Howerton, and G. Reffo, CASCADE Cray program for γ -production from discrete level inelastic scattering, Lawrence Livermore Nuclear Data Group Internal Report, PD-134 (1986), unpublished.
35. S. T. Perkins, R. C. Haight, and R. J. Howerton, *Nucl. Sci. and Eng.* 57 1 (1975).

DESCRIPTION OF EVALUATIONS FOR $^{58,60,61,62,64}\text{Ni}$ PERFORMED FOR ENDF/B-VI*

D. C. Larson, C. M. Perey, D. M. Hetrick, and C. Y. Fu
Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6356

ABSTRACT

Isotopic evaluations for $^{58,60,61,62,64}\text{Ni}$ performed for ENDF/B-VI are briefly reviewed. The evaluations are based on analysis of experimental data and results of model calculations which reproduce the experimental data. Evaluated data are given for neutron induced reaction cross sections, angular and energy distributions, and for gamma-ray production cross sections associated with the reactions. File 6 formats are used to represent energy-angle correlated data and recoil spectra. Uncertainty files are included for all File 3 cross sections.

1. INTRODUCTION

Separate evaluations have been done for each of the stable isotopes of nickel. In this report, we briefly review the structure of the evaluations, describe how the evaluations were done, and note the major pieces of data considered in the evaluation process. Experimental data references were obtained primarily from CINDA; the data themselves were mostly obtained from the National Nuclear Data Center at Brookhaven National Laboratory and, occasionally, from the literature and reports. The R-Matrix code SAMMY (LA89) was used for the resonance region analysis. The TNG nuclear model code (FU88, SH86), a multistep Hauser-Feshbach code which includes precompound and compound contributions to cross sections and angular and energy distributions in a self-consistent manner, calculates gamma-ray production, and conserves angular momentum in all steps, was the primary code used for these evaluations. Extensive model calculations were performed with the goal of simultaneously reproducing experimental data for all reaction channels with one set of parameters. This ensures internal consistency and energy conservation within the evaluation. In the case of reactions for which sufficient data were available, a Bayesian analysis using the GLUCS code (HE80) was frequently done, using ENDF/B-V (DI79) or the TNG results as the prior. In cases where insufficient data were available for a GLUCS analysis, and the available data were deemed to be accurate, but in disagreement with the TNG results, a smoothed curve representation through the data was used for the evaluation. A similar method was also used for cross sections where resonant structure was felt to be important, but resonance parameters were not included. The final evaluation is thus a combination of TNG results (used where extrapolation and interpolation was required and where data sets were badly discrepant), GLUCS results (used where sufficient data existed to do an analysis), and smoothed curves.

In Section 2 the resonance parameters are discussed; Section 3 contains a description of the major cross sections included in the evaluation; Section 4 is devoted to angular distributions; and Section 5 to energy-angle correlated distributions. Section 6 describes the uncertainty files.

* Research sponsored by the Office of Energy Research, Division of Nuclear Physics, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

The TNG calculations for $^{58,60}\text{Ni}$ are documented and extensively compared with data in (HE87). File 1 for each evaluation should be referred to for additional details.

2. RESONANCE PARAMETERS

Resonance parameters for ^{58}Ni from 10^{-5} eV to 810 keV were taken from a recent SAMMY analysis (PE88) of ORELA transmission, scattering, and capture data. Sixty-two $\ell = 0$ and 410 $\ell > 0$ resonances were identified and are included, using the Reich-Moore formats. Resonance parameters for ^{60}Ni cover the energy range from 10^{-5} eV to 450 keV and were also taken from a SAMMY analysis of ORELA transmission and capture data (PE83). Thirty $\ell = 0$ and 227 $\ell > 0$ resonances were identified and included in the ^{60}Ni evaluation. For the $^{61,62,64}\text{Ni}$ evaluations, the resonance parameters were taken from the compilation of Mughabghab (MU81).

In each case SAMMY was used to adjust negative energy dummy resonances to give the correct thermal cross sections. As noted in File 1 comments given in the evaluations, no File 3 background cross sections are used from thermal to the end of the resonance region; the cross sections are given directly by the resonance parameters.

3. CROSS SECTIONS

In this section we briefly describe the contents of the files containing cross sections for the more important reactions. The total cross section for ^{58}Ni above the resonance region was taken from a high-resolution measurement (PE88) up to 20 MeV. For ^{60}Ni the total cross section above the resonance region was also taken from isotopic data. For the minor isotopes the total cross section above the resonance region was taken from a high-resolution measurement of natural nickel by Larson (LA83). The nonelastic cross section is derived by summing the individual reaction cross sections, while the elastic cross section is derived by subtracting the nonelastic from the total. Capture cross sections are given by the resonance parameters, and renormalized TNG results are used from the end of the resonance region to 20 MeV.

Cross sections for inelastic scattering to discrete levels in $^{58,60}\text{Ni}$ were taken from the model calculations (HE87). Direct interaction contributions were included for many of the levels. Agreement with experimental data is generally favorable; however, the experimental uncertainties are often rather large. Figures 1 and 2 show a comparison of the evaluated results with experimental data for the total inelastic scattering cross section for $^{58,60}\text{Ni}$, respectively. For $^{61,62,64}\text{Ni}$ the cross sections for the lowest few levels were included from the calculations, and a continuum was used to represent the remainder of the inelastic scattering cross section.

Abundant data are available to define the $^{58,60}\text{Ni}(n,p)$ reaction cross sections. Figure 3 shows a comparison of the available data, and the ENDF/B-V and ENDF/B-VI results for the $^{58}\text{Ni}(n,p)$ cross section. The evaluated $^{58}\text{Ni}(n,p)$ cross section was partially taken from a Bayes' simultaneous analysis of several correlated cross sections (FU82), and other experimental data (see File 1 of the ^{58}Ni evaluation for details). The $^{60,61}\text{Ni}(n,p)$ cross sections were evaluated from data and TNG results. The $^{62,64}\text{Ni}(n,p)$ cross sections were taken from the TNG calculations. Data for the (n,α) reactions are sparse, and the evaluations are mainly based on calculated (occasionally renormalized) results, which compare with available experimental data. Total proton and alpha emission cross sections were also taken from the TNG and GLUCS calculations and for $^{58,60}\text{Ni}$ agreed well with the

integrated data at 14 MeV of Grimes et al. (GR79) and Kneff et al. (KN86), and with the data of Qaim et al. (QA84) at lower energies.

There is abundant cross section data for the $^{58}\text{Ni}(n, 2n)$ reaction, but no data for the $(n, 2n)$ cross section on any of the other isotopes. Results of the TNG model calculations were in good agreement with the available $(n, 2n)$ data, as well as the neutron emission spectra for natural Ni; thus results of the model calculations were used for the $(n, 2n)$ cross sections for all of the isotopes except $^{58}\text{Ni}(n, 2n)$, for which the evaluation by Pavlik and Winkler (PA83) was adopted. It should be noted that the $(n, 2n)$ cross sections are large for the minor isotopes $^{61,62,64}\text{Ni}$.

Cross sections for all other significant tertiary reactions are given for each isotopic evaluation. In particular, $^{58}\text{Ni}(n, np + n, pn)$ has a large cross section, and the evaluation is based on a renormalized TNG calculation. There is very little data for this reaction on the other isotopes. See the detailed descriptions in Ref. (HE87) for $^{58,60}\text{Ni}$, and File 1 comments in each evaluation.

4. ANGULAR DISTRIBUTIONS

Elastic-scattering angular distributions from ENDF/B-V (DI79) were reviewed and found to be in good agreement with experimental data and are retained for ENDF/B-VI as Legendre coefficients in File 4/2.

Disagreements in experimental angular distribution data sets for inelastic scattering to discrete levels are often outside rather large uncertainties. Model calculations including direct interaction and compound reaction contributions were compared with available data and used for the evaluations. These data are also entered as Legendre coefficients in File 6/51-90 in the $^{58,60}\text{Ni}$ evaluations for as many levels as discrete information is available. Only the few lowest levels were used for the minor isotopes, and isotropic angular distributions were assumed.

5. ENERGY-ANGLE CORRELATED DISTRIBUTIONS (FILE 6)

Often neutron, proton, alpha, and gamma-ray emission spectral data are measured as a function of outgoing particle angle, and this correlation of outgoing angle with measured spectra can now be represented in File 6. However, generally these distributions have only been measured at one or at most a few incident energies, thus we rely upon the TNG model calculations to reproduce the available data as a function of outgoing energy and angle, and then extrapolate to other incident neutron energies. Figure 4 illustrates the components of the neutron emission calculated with TNG which sum to give the total emission spectra for ^{58}Ni . Figure 5 shows a comparison of the experimental data with the calculated results for the natural $\text{Ni}(n, xn)$ cross section, and Figure 6 (HE87) shows a comparison of the measured and calculated angular distributions for three outgoing neutron energy bins. These calculated energy-angle distributions have been taken from the TNG calculations and entered in File 6 for the $^{58,60}\text{Ni}$ evaluations for a number of incident energies between 1 and 20 MeV. Isotropic energy-angle distributions are assumed for the minor isotope evaluations, also contained in File 6. Cross sections associated with these distributions are given in File 3.

Figures 7 and 8 show comparisons of ENDF/B-VI with experimental data for the $^{58}\text{Ni}(n, xp)$ and $^{60}\text{Ni}(n, x\alpha)$ reactions near 14 MeV, respectively. These energy distributions, with isotropic angular distributions assumed, have been entered in File 6. Recoil

spectra for the heavy residual nuclei have also been included in File 6. Since the angular distributions are given as isotropic, File 5 could have been used for all charged particle spectra with the exception of the recoil spectra, but for ease of energy balance and KERMA calculations, a consistent File 6 usage is desirable. Cross sections associated with these distributions are given in File 3.

Prior to incorporation in File 6, the neutron and charged particle energy distributions from TNG are input to the RECOIL code (FU85), which converts the energy distributions from the center of mass to the laboratory frame, and calculates the energy spectrum of the heavy recoil nucleus. These tabulated energy distributions in the lab frame are given in File 6, with the neutrons usually having anisotropic angular distributions, and isotropic angular distributions for the charged particles (including the recoil nucleus).

File 6 was also chosen to represent the gamma-ray production energy distributions, for consistency with the neutron and charged particle distributions. Isotropic angular distributions were used for the gamma rays. Figure 9 (HE87) shows a comparison of measured gamma-ray spectra around 14 MeV with the TNG calculation at 14.5 MeV. Note that without use of the calculated results, a significant amount of cross section below about 1-MeV gamma-ray energy would be missing. Calculated distributions are given in File 6 for several incident neutron energies from 1 to 20 MeV. Cross sections associated with these distributions are given in File 3.

Capture gamma-ray cross sections and spectra are obtained from information given in Files 3 (cross section), 12 (multiplicities), and 15 (spectral shapes), and are based on a combination of experimental data and calculation.

As an example of the usage of File 6, consider the $^{58}\text{Ni}(n, n\alpha)$ reaction. In Section 6/22, constant yields are given for the outgoing neutron, alpha and ^{54}Fe residual, and an energy dependent yield is used for the gamma rays associated with the $(n, n\alpha)$ reaction. Normalized energy distributions at several incident energies are given for each outgoing product, but only the outgoing neutron has a non-isotropic angular distribution. The cross section to be used for normalization is taken from Section 3/22. With the information given in Files 3 and 6, direct computation of heating, KERMA, etc. is now possible.

Energy balance ($(E_n + Q)$ must equal sum of all outgoing particle and gamma-ray energies) has been checked for all reactions, energies and isotopes, and is achieved within 1%.

6. UNCERTAINTY INFORMATION

Uncertainty files are given only for the cross sections in File 3 and not for the resonance parameters, energy distributions or angular distributions. Fractional and absolute components, correlated only within a given energy interval, are based on scatter in experimental data and estimates of uncertainties associated with the model calculations. Details of this work can be found in (HE91).

7. DATA NEEDS AND EVALUATION IMPROVEMENTS

The resonance region for $^{58,60}\text{Ni}$ is in good shape, but high-resolution transmission data for $^{61,62,64}\text{Ni}$ would improve evaluations for these materials. The capture cross-section data uncertainties may be as much as 25% for materials in this mass region, as shown for the 1.15-keV resonance in ^{56}Fe by an International Task Force. Thus, new high-resolution capture data are needed in the resonance region for at least $^{58,60}\text{Ni}$, and preferably for all

isotopes. Capture spectra at selected energies from thermal through the resonance region would be useful to improve the evaluations. The $^{58}\text{Ni}(n, np)$ reaction has a large cross section with existing data mainly around 14 MeV but discrepant. New data are needed at energies from 10 to 14 MeV and up to 20 MeV. The $^{60}\text{Ni}(n, np)$ reaction also has a large cross section; however, no data are available to verify the model calculations. The $(n, 2n)$ cross sections are large for $^{60,61,62,64}\text{Ni}$, but few data are available except for one discrepant point at 14.8 MeV for ^{60}Ni , and two points for ^{64}Ni . Further experimental guidance is necessary to verify the model calculations. Neutron emission cross-section data are needed at incident energies other than around 14 MeV to benchmark the model calculations. Uncertainties should be given for important resonance parameters, and angular and energy distributions.

REFERENCES

- BR71 W. Breunlich and G. Stengel, *Z. Naturforsch.* **A 26**, 451 (March 1971).
- CL72 G. Clayeux and J. Voignier, Centre d' Etudes de Limeil, CEA-R-4279 (1972).
- CO62 L. Colli, I. Iori, S. Micheletti, and M. Pignanelli, *Nuovo Cimento* **21**, 966 (1962).
- DI73 J. K. Dickens, T. A. Love, and G. L. Morgan, *Gamma-Ray Production From Neutron Interactions with Nickel for Incident Neutron Energies Between 1.0 and 10 MeV: Tabulated Differential Cross Sections*, ORNL/TM-4379 (November 1973). (Title has error; should read 1.0 and 20 MeV.)
- DI79 M. Divadeenam, *Ni Elemental Neutron Induced Reaction Cross-Section Evaluation*, Report BNL-NCS-51346, ENDF-294, (March 1979).
- FI84 R. Fischer, G. Traxler, M. Uhl, and H. Vonach, *Phys. Rev.* **C30**, 72 (1984).
- FU82 C. Y. Fu and D. M. Hetrick, "Experience in Using the Covariances of Some ENDF/B-V Dosimetry Cross Sections: Proposed Improvements and Addition of Cross-Reaction Covariances," p. 877 in *Proc. Fourth ASTM-EURATOM Symp. on Reactor Dosimetry*, Gaithersburg, Maryland, March 22-26, 1982, U.S. National Bureau of Standards.
- FU85 C. Y. Fu and D. M. Hetrick, unpublished, code available from authors.
- FU88 C. Y. Fu, *Nucl. Sci. Eng.* **100**, 61 (1988).
- GR79 S. M. Grimes, R. C. Haight, K. R. Alvar, H. H. Barschall, and R. R. Borchers, *Phys. Rev.* **C19**, 2127 (June 1979).
- HE75 D. Hermsdorf, A. Meister, S. Sassonoff, D. Seeliger, K. Seidel, and F. Shahin, Zentralinstitut Fur Kernforschung Rossendorf Bei Dresden, Zfk-277 (U) (1975).
- HE87 D. M. Hetrick, C. Y. Fu, and D. C. Larson, *Calculated Neutron-Induced Cross Sections for $^{58,60}\text{Ni}$ from 1 to 20 MeV and Comparisons with Experiments*, ORNL/TM-10219 (ENDF-344) (June 1987).
- HE80 D. M. Hetrick and C. Y. Fu, *GLUCS: A Generalized Least-Squares Program for Updating Cross Section Evaluations with Correlated Data Sets*, ORNL/TM-7341, ENDF-303 (October 1980).
- HE91 D. M. Hetrick, D. C. Larson and C. Y. Fu, *Generation of Covariance Files for the Isotopes of Cr, Fe, Ni, Cu, and Pb in ENDF/B-VI*, ORNL/TM-11763, (February 1991).
- JO69 B. Joensson, K. Nyberg, and I. Bergqvist, *Ark. Fys.* **39**, 295 (1969).

- KN86 D. W. Kneff, B. M. Oliver, H. Farrar IV, and L. R. Greenwood, *Nucl. Sci. Eng.* **92**, 401-524 (1986).
- LA83 D. C. Larson, N. M. Larson, J. A. Harvey, N. W. Hill, and C. H. Johnson, *Application of New Techniques to ORELA Neutron Transmission Measurements and Their Uncertainty Analysis: The Case of Natural Nickel From 2 keV to 20 MeV*, ORNL/TM-8203, ENDF-333, Oak Ridge National Laboratory, Oak Ridge, Tenn. (October 1983).
- LA85 D. C. Larson, "High-Resolution Structural Material ($n, x\gamma$) Production Cross Sections for $0.2 < E_n \leq 40$ MeV," *Proc. Conf. on Nucl. Data for Basic and Applied Science, Santa Fe, New Mexico* Vol. 1, 71 (1985).
- LA89 N. M. Larson, *Updated Users' Guide for SAMMY: Multilevel R-Matrix Fits to Neutron Data Using Bayes' Equations*, ORNL/TM-9179 (August 1984). Also ORNL/TM-9179/R1 (July 1985) and ORNL/TM-9179/R2 (June 1989).
- MA69 S. C. Mathur, P. S. Buchanan, and I. L. Morgan, *Phys. Rev.* **86**, 1038 (October 1969).
- MU81 S. F. Mughabghab, M. Divadeenam, and N. E. Holden, *Neutron Cross Sections, Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part A, Z=1-60*, Academic Press (1981).
- PA83 A. Pavlik and G. Winkler, *Evaluation of the $^{58}\text{Ni}(n,2n)^{57}\text{Ni}$ Cross Sections*, IAEA Report INDC(AUS)-9/L (1983).
- PE70 F. G. Perey, C. O. LeRigoleur, and W. E. Kinney, *Nickel-60 Neutron Elastic- and Inelastic-Scattering Cross Sections from 6.5 to 8.5 MeV*, ORNL-4523 (April 1970).
- PE83 C. M. Perey, J. A. Harvey, R. L. Macklin, and F. G. Perey, *Phys. Rev.* **C27**, 2556 (June 1983).
- PE88 C. M. Perey, F. G. Perey, J. A. Harvey, N. W. Hill, N. M. Larson, and R. L. Macklin, $^{58}\text{Ni} + n$ Transmission, Differential Elastic Scattering and Capture Measurements and Analysis from 5 to 813 keV, ORNL/TM-10841 (ENDF-347) (September 1988).
- QA84 S. M. Qaim, R. Wolffe, M.M. Rahman, and H. Ollig, *Nucl. Sci. Eng.* **88**, 143-153 (1984).
- SA72 O. A. Salnikov, G. N. Lovchikova, G. V. Kotelnikova, A. M. Trufanov, and N. I. Fetisov, *Differential Cross Sections of Inelastic Scattering Neutrons on Nuclei Cr, Mn, Fe, Co, Ni, Cu, Y, Zr, Nb, W, Bi*, Report Jadernye Konstanty -7, 102 (March 1972).
- SH86 K. Shibata and C. Y. Fu, *Recent Improvements of the TNG Statistical Model Code*, ORNL/TM-10093 (August 1986).
- TA83 A. Takahashi, J. Yamamoto, T. Murakami, K. Oshima, H. Oda, K. Fujimoto, M. Ueda, M. Fukazawa, Y. Yanagi, J. Mizaguchi, and K. Sumita, *Oktavian Report A-83-01*, Osaka University, Japan (June 1983).
- TO67 J. H. Towle and R. O. Owens, *Nucl. Phys.* **A100**, 257 (1967).
- VO80 H. Vonach, A. Chalupka, F. Wenninger, and G. Staffel, "Measurement of the Angle-Integrated Secondary Neutron Spectra from Interaction of 14 MeV Neutrons with Medium and Heavy Nuclei," *Proc. Symp. on Neutron Cross-Sections from 10 to 50 MeV*, BNL-NCS-51245, Brookhaven National Laboratory (July 1980).
- VO89 H. Vonach and M. Wagner, "Neutron Activation Cross-Sections of ^{58}Ni and ^{60}Ni for 8-12 MeV Neutrons," *Proc. of a Specialists' Meeting on Neutron Activation Cross*

Sections for Fission and Fusion Energy Applications, NEANDC-259'U', Argonne National Laboratory (September 13-15, 1989).

XI82 S. Xiamin, W. Yongshun, S. Ronglin, X. Jinqiang, and D. Dazhav, *Proc. Int. Conf. on Nuclear Data for Science and Technology, Antwerp, 373 (Sept. 6-10, 1982).*

Fig. 1. Comparison of ENDF/B-V and ENDF/B-VI with experimental total inelastic scattering cross-section data for ^{58}Ni .

Fig. 2. Comparison of ENDF/B-V and ENDF/B-VI with experimental total inelastic scattering cross-section data for ^{60}Ni .

Fig. 3. Comparison of $^{58}\text{Ni}(n,p)$ experimental data with ENDF/B-V and ENDF/B-VI.
(See Ref. HE87 for references.)

Fig. 4. Neutron emission spectra for ^{58}Ni from ENDF/B-VI at 14.5 MeV. Contributions from the various neutron-producing components are shown (they sum to the total). The curves labeled (n, np) and (n, nα) include the (n, pn) and (n, αn) components, respectively.

Fig. 5. Neutron emission spectra from ENDF/B-V (line) and ENDF/B-VI (histogram) compared with experimental data. The data of Clayeux and Voignier (CL72) and Mathur et al. (MA69) were taken at 90° , the data of Takahashi et al. (TA83) were taken at 80° , and the other measured data sets shown (HE75, VO80, and SA72) are angle integrated. The data are for natural nickel, and the isotopic evaluations have been combined to give the ENDF/B-VI result.

Fig. 6. Comparison of ENDF/B-VI with experimental neutron production cross sections as a function of angle for several outgoing neutron energy bins. This information was not previously available in ENDF/B.

Fig. 7. Comparison of ENDF/B-VI proton production spectra for ^{58}Ni with experimental data. The measurements were taken at incident energies of 14.8 and 14.1 MeV; ENDF/B-VI taken from the TNG calculation was for $E_n = 14.5$ MeV. The data of Grimes et al. (GR79, HA77) are angle integrated; the data of Colli et al. (CO62) were taken at 15° . This information was not previously available in ENDF/B.

Fig. 8. Comparison of ENDF/B-VI and experimental alpha production spectra for ^{60}Ni . The measurements were taken at incident energies of 14.8 and 14.1 MeV and are angle integrated; the TNG calculation was for $E_n = 14.5$ MeV. This information was not previously available in ENDF/B.

Fig. 9. Secondary gamma-ray production cross section versus gamma-ray energy from the TNG calculation (incident energy $E_n = 14.5$ MeV) compared with the data of Dickens et al. (DI73).

⁵⁹₂₈Ni

Reference: No Primary Reference
Evaluator: F. M. Mann
Evaluated: January 1983
Material: 2828
Content: Activation

File Comments

This file contains activation cross sections for ⁵⁹Ni, and includes smooth MF=3 cross sections for capture MT=102, proton production MT=103, and α production MT=107.

The evaluation uses a line shape based upon the resonance parameters from the compilation of S. F. Mughabghab up to 10 keV.¹ The smooth cross sections are also based on Hauser-Feshbach calculations which agree with ⁵⁶Fe (α, n_0) measurements by R. W. Kavanagh (Cal Tech).²

References:

1. S. F. Mughabghab, M. Divadeenam and N. E. Holden, "Neutron Cross Sections," Vol. 1A, Academic Press, New York (1981).
2. R. W. Kavanagh, California Institute of Technology, Private Communication (1982).

DESCRIPTION OF EVALUATIONS FOR $^{63,65}\text{Cu}$ PERFORMED FOR ENDF/B-VI*

D. M. Hetrick, C. Y. Fu and D. C. Larson
Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6356

ABSTRACT

Isotopic evaluations for $^{63,65}\text{Cu}$ performed for ENDF/B-VI are briefly reviewed. The evaluations are based on analysis of experimental data and results of model calculations which reproduce the experimental data. Evaluated data are given for neutron-induced reaction cross sections, angular and energy distributions, and for gamma-ray production cross sections associated with the reactions. File 6 formats are used to represent energy-angle correlated data and recoil spectra. Uncertainty files are included for all File 3 cross sections.

1. INTRODUCTION

Separate evaluations have been done for the two stable isotopes of copper. In this report we briefly review the structure of the evaluations, describe how the evaluations were done, and note the major pieces of data considered in the evaluation process. Experimental data references were obtained primarily from CINDA; the data themselves were mostly obtained from the National Nuclear Data Center at Brookhaven National Laboratory and, occasionally, from the literature and reports. The TNG nuclear model code (FU88, SH86), a multistep Hauser-Feshbach code which includes precompound and compound contributions to cross sections and angular and energy distributions in a self-consistent manner, calculates gamma-ray production, and conserves angular momentum in all steps, was the primary code used for these evaluations. Extensive model calculations were performed with the goal of simultaneously reproducing experimental data for all reaction channels with one set of parameters. This ensures internal consistency and energy conservation within the evaluation. In the case of reactions for which sufficient data were available, a Bayesian analysis using the GLUCS code (HE80) was frequently done, using ENDF/B-V or the TNG results as the prior. In cases where insufficient data were available for a GLUCS analysis and the available data were deemed to be accurate, but in disagreement with the TNG results, a smoothed curve representation through the data was used for the evaluation. A similar method was also used for cross sections where resonant structure was felt to be important, but resonance parameters were not included. The final evaluation is thus a combination of TNG results (used where extrapolation and interpolation was required and where data sets were badly discrepant), GLUCS results (used where sufficient data existed to do a statistical analysis), and smoothed curves.

In Section 2 the resonance parameters are discussed; Section 3 contains a description of the major cross sections included in the evaluation; Section 4 is devoted to angular distributions; and Section 5 to energy-angle correlated distributions. Section 6 describes the uncertainty files. Further details of each evaluation are given in the File 1 comment sections.

* Research sponsored by the Office of Energy Research, Division of Nuclear Physics, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

The TNG calculations performed for this work are documented and extensively compared with experimental data in (HE84).

2. RESONANCE PARAMETERS

Resonance parameters for $^{63,65}\text{Cu}$ are taken from the compilation of Mughabghab (MU81). They describe the energy range from 10^{-5} eV to 153 keV for ^{63}Cu and 10^{-5} eV to 149 keV for ^{65}Cu , however the fit to the data above 100 keV is rather poor, so the resonance region stops at 99.5 keV for both isotopes. Average capture widths are used for neutron energies above about 50 keV. A smooth background cross section is included to provide the correct thermal cross sections. The resonance parameters should be processed with the Reich-Moore formalism. These evaluations would benefit from a better analysis of the resonance region data.

3. CROSS SECTIONS

This section contains a brief discussion of the cross-section files in the evaluations for $^{63,65}\text{Cu}$. The total cross section above the resonance region to 1.12 MeV was taken from the isotopic experimental data of Pandey (PA77). From 1.12 to 20 MeV, natural data of Perey (PE77) and Larson (LA80) was used in the absence of isotopic data. The nonelastic cross section was derived by summing the individual reaction cross sections. The elastic cross section was derived as the difference between the total and elastic cross sections.

Cross sections for inelastic scattering to discrete levels are taken from the model calculations, which included a direct interaction component and generally are in good agreement with the available experimental data. A continuum was used to represent the inelastic scattering cross section for excitation energies above the discrete levels. Comparisons with experimental data are shown in (HE84).

The $^{63}\text{Cu}(n, p)$ reaction has very little data, but the calculated result agrees with the data of Qaim and Molla (QA77) and Allan (AL61). The available data for this reaction is confusing, and the situation is discussed in (FU82a). The $^{63}\text{Cu}(n, \alpha)$ reaction has much data and is a common dosimetry cross section. The evaluated cross section for this reaction is taken from the results of a generalized least-squares (GLUCS) analysis (FU82) of twelve dosimetry reactions, which included ratio data and covariance information. The $^{65}\text{Cu}(n, p)$ cross section has abundant data and is adequately compromised by the TNG calculations, which are used for the evaluation. The $^{65}\text{Cu}(n, \alpha)$ cross section is small, and the experimental data are inconsistent. The calculated results are used for the evaluation.

The $^{63,65}\text{Cu}(n, 2n)$ cross sections are well defined by experimental data, and the results of a GLUCS analysis were used for the evaluation. Other tertiary reaction cross sections with data are reproduced by the TNG calculations and are included in each evaluation. $^{63}\text{Cu}(n, np)$ is the only tertiary reaction with a cross section larger than 80 mb.

The capture cross sections for $^{63,65}\text{Cu}$ are defined by the resonance parameters and a smooth background below 100 keV, and by experimental data above the resonance region. Guided by experimental data and the TNG calculations, a smooth line was drawn through the data from 100 keV to 20 MeV and used for the evaluations.

4. ANGULAR DISTRIBUTIONS

Elastic scattering angular distributions were obtained from an optical potential derived by fitting experimental angular distribution data for $^{63,65,\text{nat}}\text{Cu}$ with GENOA (PE67). A compound elastic term was included for neutron energies below 5 MeV. Since very little difference was observed between the experimental data for ^{63}Cu and ^{65}Cu , one potential was derived and used for both evaluations. Figures 1 and 2 show a comparison of the calculated and experimental data for $E_n = 8.05$ and 14.5 MeV. A description of the data sets used, the optical model analysis and final parameters, and comparisons with experimental data are given in (HE84). The angular distributions are represented as Legendre coefficients and given in File 4/2. In the resonance region, the angular distributions can be derived from the Reich-Moore resonance parameters. Angular distributions for inelastic scattering to excited levels and the continuum are given as Legendre coefficients in File 6. They are taken from the TNG and DWUCK analyses, and comparisons with data are shown in (HE84).

5. ENERGY-ANGLE CORRELATED DISTRIBUTIONS (FILE 6)

Neutron emission spectra, as a function of outgoing energy and angle, are given in File 6. For copper, the measurements of Morgan et al. (MO79) give the outgoing neutron spectra at one angle for several incident neutron energies between 1 and 20 MeV, while the measurements of Hermsdorf et al. (HE75), Vonach et al. (VO80), Salnikov et al. (SA75), and Takahashi et al. (TA83) give the outgoing spectra at several angles but only near 14.5-MeV incident energy. Such complementary measurements allow a good determination of the model parameters for the calculations and, thereby, reliable interpolation and extrapolation to energies where there are no data. For these reasons, as well as ensuring energy conservation, results from the model codes, expressed in File 6 formats, were used for the evaluations. The angular distributions were expressed in terms of Legendre coefficients, while the energy distributions were expressed as tabulated probability distributions. Figure 3 illustrates the components of the neutron emission calculated with TNG which sum to give the total emission spectra for ^{63}Cu . Figure 4 shows the neutron emission data of Morgan et al. (MO79) compared with ENDF/B-V and ENDF/B-VI for the incident neutron-energy bin from 9 to 10 MeV. Figure 5 shows several sets of neutron emission data around 14.5 MeV, compared with ENDF/B-V and ENDF/B-VI. The data of Takahashi et al. (TA83) became available after the evaluation was done but are found to be in good agreement with the evaluation.

Proton and alpha emission spectra for both isotopes are available (GR79) at an incident energy of 14.8 MeV. The calculations are in excellent agreement with the measured spectra, including reproducing the observed sub-coulomb emission of protons. Figure 6 shows a comparison of the measured data for proton emission from ^{63}Cu with ENDF/B-VI. However, the observed sub-coulomb emission of alphas is not as well reproduced by the TNG calculations. Figure 7 shows a comparison of the measured data for ^{63}Cu alpha emission, compared with the ENDF/B-VI results.

Prior to incorporation in File 6, the neutron and charged particle energy distributions from TNG are input to the RECOIL code (FU85), which converts the energy distributions from the center of mass to the laboratory frame, and calculates the energy spectrum of the heavy recoil nucleus. These tabulated energy distributions in the lab frame are given in File 6, with the neutrons usually having anisotropic angular distributions, and isotropic angular distributions for the charged particles (including the recoil nucleus).

Gamma-ray production spectra were also calculated as part of the TNG calculations, and compared with data sets of Rogers et al. (RO77), Morgan (MO79), Dickens et al. (DI73), and Chapman (CH76) (see Ref. HE84). Figure 8 shows a comparison of the measured data of Dickens et al. with the TNG results around 14-MeV incident energy. Note that without the use of the calculated results, a significant amount of cross section below 700-keV gamma-ray energy would not be accounted for due to gamma rays from the $(n, 2n)$ reaction. Since calculated results are generally used for the evaluation, energy conservation is ensured. Sections of File 6 were used to represent the gamma-ray emission spectra for the individual reactions, and isotropic angular distributions were assumed. The cross sections for the gamma-ray production are given in corresponding sections of File 3.

As an example of the usage of File 6, consider the $^{65}\text{Cu}(n, n\alpha)$ reaction. In Section 6/22, constant yields are given for the outgoing neutron, alpha and ^{61}Co residual, and an energy dependent yield is used for the gamma rays associated with the $(n, n\alpha)$ reaction. Normalized energy distributions are given for each outgoing product, but only the outgoing neutron has a non-isotropic angular distribution. The cross section to be used for normalization is taken from Section 3/22.

Capture gamma-ray cross sections and spectra are obtained from Files 3, 12 and 15, and are based on a combination of experimental data and calculation.

Energy balance $((E_n + Q)$ must equal sum of all outgoing particle and gamma-ray energies) has been checked for all reactions, energies and isotopes, and is achieved within 1%.

6. UNCERTAINTY INFORMATION

Uncertainty files are given for all cross sections in File 3, but not for the resonance parameters, energy distributions or angular distributions. Fractional and absolute components, correlated within a given energy interval, are based on scatter in experimental data and estimates of uncertainties associated with the model calculations (HE91).

7. DATA NEEDS AND EVALUATION IMPROVEMENTS

High-resolution transmission measurements for both isotopes are needed from 100 eV to 20 MeV to allow a detailed resonance parameter analysis. Presently available data do not have adequate resolution. The $^{63}\text{Cu}(n, p)$ reaction has only one reliable data point, at 14.8 MeV, and would benefit from data at lower energies. The $^{65}\text{Cu}(n, p)$ reaction has more data, but the data sets are discrepant and the data base would benefit from further, careful measurements. The $^{63}\text{Cu}(n, np)$ cross section is large and has only discrepant data available. Capture data should be checked for response function problems similar to those for the 1.15-keV resonance in ^{56}Fe ; new data may be needed if the hardness of the capture spectra is significantly different from resonance to resonance. Uncertainties should be provided for important resonance parameters as well as angular and energy distributions.

REFERENCES

- AL61 D. L. Allan, *Nuclear Physics* **24**, 274 (April 1961).
- CH76 G. T. Chapman, *The $\text{Cu}(n, x\gamma)$ Reaction Cross Section for Incident Energies Between 0.2 and 20.0 MeV*, ORNL/TM-5215 (1976).

- CO58 J. H. Coon, R. W. Davis, H. E. Felthausen, D. B. Nicodemus, *Phys. Rev.* **111**, 250 (1958).
- DI73 J. K. Dickens, T. A. Love, and G. L. Morgan, *Gamma-Ray Production Due to Neutron Interactions with Copper for Incident Neutron Energies Between 1.0 and 20.0 MeV: Tabulated Differential Cross Sections*, ORNL-4846 (1973).
- FU80 C. Y. Fu, "A Consistent Nuclear Model for Compound and Precompound Reactions with Conservation of Angular Momentum," p. 757 in *Proc. Int. Conf. Nuclear Cross Sections for Technology*, Knoxville, TN, Oct. 22-26, 1979, NBS-594, U.S. National Bureau of Standards. also, ORNL/TM-7042 (1980).
- FU82 C. Y. Fu and D. M. Hetrick, "Experience in Using the Covariance of Some ENDF/B-V Dosimetry Cross Sections: Proposed Improvements and Addition of Cross-Reaction Covariances," p. 877 in *Proc. Fourth ASTM-EURATOM Symp. on Reactor Dosimetry*, Gaithersburg, Md., March 22-26, 1982, U.S. National Bureau of Standards.
- FU82a C. Y. Fu, *Summary of ENDF/B-V Evaluations for Carbon, Calcium, Iron, Copper, and Lead and ENDF/B-V Revision 2 for Calcium and Iron*, ORNL/TM-8283 (ENDF-325), (1982).
- FU88 C. Y. Fu, *Nucl. Sci. Eng.* **100**, 61 (1988).
- FU85 C. Y. Fu and D. M. Hetrick, unpublished, code available from authors.
- GR79 S. M. Grimes, R. C. Haight, K. R. Alvar, H. H. Barschall, and R. R. Borchers, *Phys. Rev.* **C19**, 2127 (1979).
- HE75 D. Hermsdorf, A. Meister, S. Sassonoff, D. Seeliger, K. Seidel, and F. Shahin, Zentralinstitut Fur Kernforschung Rossendorf Bei Dresden, Zfk-277 (U), (1975).
- HE80 D. M. Hetrick and C. Y. Fu, *GLUCS: A Generalized Least-Squares Program for Updating Cross Section Evaluations with Correlated Data Sets*, ORNL/TM-7341, ENDF-303 (October 1980).
- HE84 D. M. Hetrick, C. Y. Fu, D. C. Larson, *Calculated Neutron-Induced Cross Sections for $^{63,65}\text{Cu}$ from 1 to 20 MeV and Comparisons with Experiments*, ORNL/TM-9083, ENDF-337 (August 1984).
- HE91 D. M. Hetrick, D. C. Larson and C. Y. Fu, *Generation of Covariance Files for the Isotopes of Cr, Fe, Ni, Cu, and Pb in ENDF/B-VI*, ORNL/TM-11763, (February 1991).
- HO69 B. Holmqvist and T. Wiedling, Atomic Energy Company, Studsvik, Nykoping, Sweden, Report AE-366 (1969).
- LA80 D. C. Larson, *ORELA Measurements to Meet Fusion Energy Neutron Cross Section Needs*, BNL-NCS-51245, Brookhaven National Lab. (July 1980)
- MO79 G. L. Morgan, *Cross Sections for the $\text{Cu}(n, xn)$ and $\text{Cu}(n, x\gamma)$ Reactions Between 1 and 20 MeV*, ORNL-5499, ENDF-273 (1979).
- MU81 S. F. Mughabghab, M. Divadencam, and N. E. Holden, *Neutron Cross Sections, Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part A, Z=1-60*, Academic Press (1981).
- PA77 M. S. Pandey, J. B. Garg, and J. A. Harvey, *Phys. Rev.* **C15**, 600 (February 1977).
- PE67 F. G. Perey, Computer code GENOA, Oak Ridge National Laboratory, unpublished (1967).

- PE77 F. G. Perey, private communication, 1977.
- QA77 S. M. Qaim and N. I. Molla, *Nucl. Phys.* **A283**, 269 (June 1977).
- RO77 V. C. Rogers, D. R. Dixon, C. G. Hoot, D. Costello, and V. J. Orphan, *Nucl. Sci. Eng.* **62**, 716 (1977).
- SA75 O. A. Salmikov, G. N. Lovchikova, G. V. Kotelnikova, A. M. Trufanov, N. I. Fetisov, *Energy Spectra of Inelastically Scattered Neutrons for Cr, Mn, Fe, Co, Ni, Cu, Y, Zr, Nb, W, and Bi*, IAEA Nuclear Data Section, Karntner Ring 11, A-1010 Vienna (July 1974).
- SH86 K. Shibata and C. Y. Fu, *Recent Improvements of the TNG Statistical Model Code*, ORNL/TM-10093 (1986).
- TA83 A. Takahashi, J. Yamamoto, T. Murakami, K. Oshima, H. Oda, K. Fujimoto, M. Ueda, M. Fukazawa, Y. Yanagi, J. Miyaguchi, and K. Sumita, Oktavian Report A-83-01, Osaka University, Japan (June 1983).
- VO80 H. Vonach, A. Chalupka, F. Wenninger, and G. Staffel, "Measurement of the Angle-Integrated Secondary Neutron Spectra from Interaction of 14 MeV Neutrons with Medium and Heavy Nuclei," *Proc. Symp. on Neutron Cross Sections from 10 to 50 MeV*, BNL-NCS-51245, Brookhaven National Laboratory (July 1980).

Fig. 1. Comparison of final optical-model fit with elastic scattering data of Holmqvist and Wiedling (HO69) for Cu at 8.05 MeV.

Fig. 2. Comparison of final optical-model fit with elastic scattering data of Coon et al. (CO58) for Cu at 14.5 MeV.

Fig. 3. Neutron emission spectra for ^{63}Cu from ENDF/B-VI at 14.5 MeV. Contributions from the various neutron-producing components are shown (they sum to the total). The curves labeled (n, np) and n, nα) include the (n, pn) and (n, αn) components, respectively.

Fig. 4. Neutron emission spectra from ENDF/B-V (line) and ENDF/B-VI (histogram) compared with the data of Morgan (M079). The data are for natural copper, and the isotopic evaluations have been combined to give the ENDF/B-VI results.

Fig. 5. Neutron emission spectra from ENDF/B-V (line) and ENDF/B-VI (histogram) compared with experimental data. The data of Morgan (MO79) and Takahashi et al. (TA83) were taken at 130° , while the other data sets shown (HE75, VO80, SA72) are angle integrated. The data are for natural copper, and the isotopic evaluations have been combined to give the ENDF/B-VI result.

Fig. 6. Comparison of ENDF/B-VI proton production spectra for ^{63}Cu with experimental data. The measurement was taken at an incident energy of 14.8 MeV; ENDF/B-VI taken from the TNG calculation was for $E_n = 14.5$ MeV. This information was not previously available in ENDF/B.

Fig. 7. Comparison of ENDF/B-VI with experimental alpha production spectra for ^{65}Cu . The measurement was taken at an incident energy of 14.8 MeV; ENDF/B-VI taken from the TNG calculation was for $E_n = 14.5$ MeV. This information was not previously available in ENDF/B.

Fig. 8. Secondary gamma-ray production cross section versus gamma-ray energy from the TNG calculation (incident energy $E_n = 14.5$ MeV) compared with the data of Dickens et al. (DI73).

Reference: ANL/NDM-94
Evaluators: R. Howerton (LLNL), A. Smith and D. Smith (ANL)
Evaluated: January 1986
Material: 3925
Content: Neutron Transport, Gamma production, Covariances

1. Introduction

Elemental yttrium is monoisotopic and magic in neutron number ($N = 50$). It lies at the end of a prominent fission product decay chain with chain yields varying from approximately 6% for ^{232}Th fission to 1.2% for ^{240}Pu fission. As such, its neutronic properties are a consideration in the optimization of FBR and similar nuclear energy systems. The primary reference for this evaluation is ANL/NDM-94, by A. B. Smith, D. L. Smith, P. Rousset, R. D. Lawson, and R. J. Howerton (1986).

2. Evaluated Resolved Resonance Range

This file employs the resonance parameter representation up to 150 keV. The resonance parameters were taken from S. F. Mughabghab et al.¹ The bound resonance of this compilation was deleted, and background cross sections were introduced in a manner as to ensure the correct thermal cross section values as given in Ref. 1.

3. Evaluated Total Cross Sections

The evaluated total cross sections were deduced from experimental values. The data base was assembled from the literature as referenced in CINDA and the files of the National Nuclear Data Center. At low energies (less than 600 keV) there are large fluctuations reflecting partially resolved underlying resonance structure. Where possible self shielding corrections were made. The cross sections were derived from the data base using the rigorous statistical model of Poenitz.² Fluctuations were smoothed by fitting the evaluated data set with a simple optical model calculation. Below 600 keV several measurements, such as in Refs. 3 and 4, show the large and partially resolved resonance structure. These were incorporated in the evaluation by normalizing the fluctuating values to the energy averaged evaluation. The present total cross sections are qualitatively very different from ENDF/B-V values. The relative shape of the ENDF/B-V evaluation seems inconsistent with any known physical interpretation.

4. Evaluated Elastic Scattering Cross Sections

From one to ten MeV the evaluated cross sections are based upon the experimental values of Ref. 5 through 8. Below 1 MeV the elastic scattering cross sections are essentially equivalent to the total cross sections with only a small difference due to radiative capture. Above 10 MeV the cross sections were extrapolated to 20 MeV using the model of Ref. 5.

5. Evaluated Inelastic Scattering Cross Sections

5.1 Discrete Inelastic Processes

The discrete inelastic scattering cross sections extend up to 3.2 MeV, assuming the energies, spins, and parities given in Refs. 6 and 7. The cross sections were largely based upon the experimental results of Refs. 6, 7, and 8. The experimental results were interpolated using the statistical model and optical potential of Refs. 5 and 7. The agreement between measured and calculated values was very good, and thus the calculations were used for the evaluation. The uncertainties associated with the evaluated quantities vary from approximately 5%, for the prominent excitations, to 20+ % for levels which are weakly excited.

5.2 Continuum Inelastic Scattering Processes

The continuum inelastic cross sections extend from 3.2 MeV to 20 MeV. Neutron emission was assumed isotropic. For the present evaluation the continuum inelastic cross section is the difference between the evaluated non-elastic cross section and the sum of the other partial cross sections.

6. Evaluated Radiative Capture Cross Sections

The experimental data base is not particularly definitive. The evaluation primarily relies upon the recent prompt detection data of Refs. 9 - 11. The evaluation is an interpolation of the measured quantities using the code ABAREX.¹² ABAREX adjusts the s-wave strength function to achieve a best fit to the data. A small direct capture component was calculated at high energies consistent with Ref. 13. The ENDF/B-V evaluation is approximately a factor of two larger than this evaluation, and is inconsistent with all recent experimental results.

7. Evaluated (n,2n) and (n,3n) Reactions

The threshold for the (n,3n) reaction is above 20 MeV and thus the process is ignored. The threshold for the (n,2n) reaction is 11.469 MeV. The majority of the

measured values were obtained using activation techniques. No comparison can be made with ENDF/B-V as the latter file does not contain the reaction. The present evaluation is consistent with the data of Philis.¹⁴

8. Evaluated Charged Particle Emitting Reactions

8.1 (n,p) and (n,np) Reactions

Primarily, the experimental data of Bayhurst and Prestwood¹⁵ and the total hydrogen production at 15 MeV reported by Haight et al. in Ref. 16 was used. The energy dependence has been estimated by E. Arthur using multiple step Hauser-Feshbach theory.¹⁷ That prediction is consistent with the available experimental evidence and with other calculational estimates. Therefore, the (n,p) cross section given by Arthur was taken for the evaluation without renormalization. The present evaluation assumes that the experimental total hydrogen production results reported by Haight, and the relative energy dependence predicted by Arthur are representative of the (n,np) process. With this assumption the predictions of Arthur were multiplied by 1.47 to obtain the present evaluation. ENDF/B-V has no comparative cross sections.

8.2 (n, α) and (n,n α) Reactions

The experimental data base is very limited and confined to the (n, α) reaction. The total helium production cross sections of Haight¹⁶ are a reasonable check of the (n, α) cross section. The present evaluation relies on the calculated values of Arthur¹⁷ to obtain the energy dependent shapes and the relative intensities of the (n, α) and (n,n α) cross sections. The calculations were normalized (upwards of 30%) to bring them into good agreement with Haight.¹⁶ There is no comparable ENDF/B-V file.

8.3 Minor (n,x) Reactions

The remaining (n,x) reactions are generally small and have relatively high thresholds. They are included for completeness, though they will have very little effect upon most neutronic applications.

The experimental knowledge of the (n,d) reaction is confined to the single 15 MeV direct particle detection result of Haight.¹⁶ The present evaluation uses calculations¹⁸ to guide the energy dependent shape and normalizes the calculated result to the measured value of Haight. The (n,nd) threshold is at approximately 16 MeV, and has been ignored.

There have been a few measurements of the (n,t) reaction near 14 MeV, all in the micro-barn range. The (n,t) reaction has been qualitatively included in the evaluation,

while the (n,nt) reaction is ignored as the threshold is ≈ 18 MeV.

Several other minor (n,x) processes are qualitatively included for completeness.

9. Evaluated Photon Production Reactions

For capture the spectral measurements of V. Orphan et al.¹⁹ were used. Photon production and spectra were obtained through a multi-step process. The resulting incident neutron energy dependent available photon energies for each reaction and the reaction cross sections were combined using the R-parameter method of Ref. 20 to obtain γ ray spectra and production cross sections.

10. Summary Comments

In a number of sensitive areas the present file is very different from that of ENDF/B-V. The differences may have a strong impact on some applications. The present file is reasonably supported by the newer and more accurate experimental information.

References

1. S. F. Mughabghab, Neutron Cross Sections Vol. 1, Part B, Academic Press Inc. New York, (1984); also S. Mughabghab and C. Dunford, private communication (1982).
2. W. P. Poenitz, Brookhaven National Laboratory Report, BNL-NCS-51363 Vol. I 249(1981); as modified by M. Sugimoto (1987).
3. J. Whalen and J. Meadows, Argonne National Laboratory Report, ANL-7310 (1968). Data from 0.047 to 20 MeV.
4. H. Newson et al., Phys. Rev. 105 1981 (1957). Data from 0.01 to 0.07 MeV.
5. R. Lawson, P. Guenther, and A. Smith, Phys. Rev. C34 1599 (1986).
6. C. Buttz-Jørgenson, P. Guenther, A. Smith, and J. Whalen, Argonne National Laboratory Report, ANL/NDM-79 (1982)
7. C. Buttz-Jørgenson, P. Guenther, J. Whalen, W. McMurray, M. Renan, I. van Heerden and A. Smith, Z. Phys. A319 47 (1984).
8. F. Perey and W. Kinney, Oak Ridge National Laboratory Report, ORNL-4552 (1970).

9. W. Poenitz, Argonne National Laboratory Report, ANL-83-4 (1983).
10. J. Boldeman et al., Phys. Rev. 120 556 (1960).
11. S. Joly et al., Bull. Am. Phys. Soc. 24 87 (1979). Also National Bureau of Standards Publication, NBS-594 (1979).
12. P. Moldauer, computer code ABAREX, private communication (1982).
13. I. Bergqvist et al., Nucl. Phys. A295 256 (1978).
14. C. Philis, CEA Report, CEA-R-4636 (1975).
15. B. Bayhurst and R. Prestwood, J. Inorg. Nucl. Chem. 23 173 (1961).
16. R. Haight et al., Phys. Rev. C23 700 (1981).
17. E. Arthur, Los Alamos National Laboratory Report, LA-7789-MS (1979).
18. M. Blann, Private Communication (1985).
19. V. J. Orphan, N. C. Rasmussen, and T. L. Harper, "Line and Continuum γ -ray Yields from Thermal Neutron Capture in 75 Elements," Gulf General Atomic Report, GA-10248/DASA 2570 (1970).
20. S. T. Perkins, R. C. Haight, and R. J. Howerton, Nucl. Sci. and Eng. 57 1 (1975).

⁹³₄₁Nb

Reference: ANL/NDM-88, ANL/NDM-117
Evaluators: A. Smith, D. Smith, L. Geraldo, and R. Howerton (LLNL).
Evaluated: February 1985 (March 1990, Dosimetry)
Material: 4125
Content: Neutron Transport, Gamma production, Covariances

1. Introduction

The evaluated nuclear data file for niobium extending over the energy range from 10^{-11} MeV to 20 MeV is suitable for comprehensive neutronic calculations. It is particularly suited for calculations dealing with fusion energy systems. The evaluation is referenced in ANL/NDM-88, by A. B. Smith, D. L. Smith (ANL), and R. J. Howerton (LLNL) (1985). The file, converted to ENDF/B-VI, provides dosimetry information as referenced by D. L. Smith and L. P. Geraldo in ANL/NDM-117 (1990).

2. Evaluated Resolved Resonance Range

The file employs the resonance parameter representation to 8 keV. The resonance parameters were taken from S. F. Mughabghab et al.¹ Small background contributions were added to the file 3 total, elastic, and capture cross sections to be consistent with Ref. 1, and to provide a reasonably smooth interface with the energy averaged cross sections at 8 keV.

3. Evaluated Total Cross Sections

This portion of the evaluation extends from 8 keV to 20 MeV. The experimental data base was assembled from files at the National Nuclear Data Center, and from the literature referenced in CINDA. The evaluated result fluctuated depending upon the details of the input data. These fluctuations were smoothed by χ^2 fitting a conventional optical model to the evaluated cross sections. At high energies above 15 MeV the present evaluation is slightly lower than ENDF/B-V. That is a region where recent data has a relatively large effect.

4. Evaluated Elastic Scattering Cross Sections

From 1 to 10 MeV the elastic scattering evaluation explicitly relies upon the experimental results of A. Smith et al.^{2,3} Together with the total cross section and other explicitly measured partial cross sections they define the experimentally poorly-known inelastic continuum cross sections over a wide energy range. The model given in ANL/NDM-70³ was used to extrapolate the measurements to lower energies. The extrapolation is consistent with the measured values of D. Reitmann et al.⁴ Above 10 MeV the evaluation is based on Ref. 5 and the experimental results of Ref. 3. Over the range from one to ten MeV where the evaluation is based on careful measurements the elastic uncertainty is 3%. Elastic scattering distributions are explicitly derived from the experimental values over the 1 - 10 MeV range.

5. Evaluated Inelastic Scattering Cross Sections

5.1 Discrete Inelastic Processes

The evaluation uses 23 excited levels extending to 2.0 MeV taken from Ref. 6. The calculated cross sections were compared with the experimental (n,n') values, grouped to comparable resolutions where necessary, and normalized to the experimental values to obtain the evaluated cross sections. This method was successful to excitations of approximately 1.5 MeV, but for higher energy excitations the normalizations became unreasonably large. Above excitations of 1.9 MeV the evaluation is based entirely upon experimental observation.

5.2 Continuum Inelastic Scattering Processes

The evaluation is consistent with the fragmentary experimental information below the (n,2n) threshold as given in Refs. 7, 8, and 9. The compound nucleus contribution is largely absorbed in the (n,2n) process above 10 MeV and the cross section at higher energies is largely due to pre-compound processes. Fluctuation structure, observed experimentally, is not included in the present evaluation.

6. Evaluated Radiative Capture Cross Sections

The experimental data base was assembled from files at the National Nuclear Data Center, and from the literature. The reported experimental data were renormalized to ENDF/B-V standards. The curve is in good agreement with the recent high resolution measurements of R. Macklin et al.¹⁰ The evaluation is also in good agreement with ENDF/B-V.

7. Evaluated (n,2n) and (n,3n) Reactions

The experimental data is based primarily on L. Veaser et al.¹¹ and J. Frehaut et al.¹² The most comprehensive measurements were made using the tank technique. Below 12 MeV the experimental results are well represented by the evaluation of Philis and Young.¹³ Above 14 MeV there are the recent and comprehensive results of Ref. 11. The present evaluation is generally 10 to 15% larger than ENDF/B-V. The neutron emission spectrum was represented by a simple Maxwellian of the form $\sqrt{E} \times \exp -E/T$. The "temperature" T was adjusted to give a good representation of the measured and calculated 14 MeV emission spectrum.

The (n,3n) reaction has a high threshold (≈ 16.9 MeV) and a small cross section. There appears to be only one experimental data set, (Ref. 11) and the evaluation is a subjectively constructed curve through these few experimental values. The estimated uncertainties are large, 15 - 20% near 20 MeV, and they increase as the energy decreases. The present evaluation is considerably different from ENDF/B-V.

8. Evaluated Charged Particle Emitting Reactions

More than 35 of these processes are energetically available in the bombardment of niobium with neutrons of less than 20 MeV. Most are of no consequence for neutronic analysis for which this file is intended. For special purposes the user is encouraged to consult an activation file, such as that maintained at LLNL.¹¹ The present evaluation considers the reactions shown in table 1. The Q values have been taken from Ref. 14.

Table 1

Q-values for Charged Particle Emitting Reactions

Reaction	Q-value (MeV)
(n,p)	+0.690
(n,np)	-6.042
(n, α)	+4.918
(n,n α)	-1.938
(n,d)	-3.817
(n,nd)	-12.452
(n,t)	-6.195
(n,nt)	-13.395
(n, ^3He)	-7.720
(n,n ^3He)	-15.660

8.1 (n,p) and (n,np) + (n,pn) Reactions

The residual products do not lend themselves to activity measurements. The total proton production at 15 MeV has been measured by Grimes et al.¹⁵ to be 51 ± 8 mb. Pre-compound processes have been shown by P. Young to be significant.¹⁶ Calculated results were normalized by a factor of 1.23 to give agreement with the observed total hydrogen production cross section given by Grimes at 15 MeV. The (n,p) cross section is qualitatively consistent with ENDF/B-V values.

8.2 (n, α) and (n,n α) + (n, α n) Reactions

The (n, α) cross section is reasonably defined by experiments to 20 MeV. See Refs. 17 through 20. Production of helium at 15 MeV has been reported by Grimes et al.¹⁵ and Haight.²¹ The lower energy cross sections follow the calculations of Strohmaier.²² The (n, α) cross section and the measured total helium production imply a (n,n α) cross section of approximately 5.5 mb at 15 MeV in agreement with the calculated results of Ref. 16. Therefore the calculations of Ref. 16 were used for the present (n,n α) evaluation.

8.3 (n,d) and (n,nd) + (n,dn) Reactions

The evaluation employs a simple barrier penetration calculation and a normalization to the measured gas production value.¹⁵ These reactions are not given in ENDF/B-V.

8.4 (n,t) and (n,nt) + (n,tn) Reactions

The evaluation is based on calculations of M. Blann²³ and a measured experimental data base.^{24,25} There are no comparable ENDF/B-V files.

9. Evaluated Photon Production Reactions

The spectrum of neutrons from the capture reaction was taken from Orphan.²⁶ A multiple step process was used to derive photon production cross sections and spectra. The resulting total photon energy and the cross sections for the reactions were combined using the R-parameter method of Perkins et al.²⁷

10. Activation of ^{93m}Nb Dosimetry

The production of the isomer ^{93m}Nb by the (n,n') process is routinely employed for neutron dosimetry applications. This reaction is the first excited state of ⁹³Nb at

30.82 keV. The half life of ^{93m}Nb is 16.1 years and the decay is by isomeric transition with almost 100% internal conversion.

Apparently the only formally published direct experimental result is that of Ryves and Kolkowski at 14.68 MeV.²⁸ Strohmaier et al.^{22,29} generated an evaluation based on model calculations. The calculated cross section of 34.3 mb for the 13.92 - 14.93 MeV range agrees well with the experimental value of 36.5 ± 3.0 mb reported in Ref. 28. Strohmeier's results were used above 700 keV. Model calculations were performed for the evaluation below 700 keV. In this region the cross section is based entirely upon neutron excitation of the first excited level (the isomeric level) of Nb, in competition with radiative capture. The two independent evaluations were joined at approximately 700 keV.

References

1. S. F. Mughabghab, Neutron Cross Sections Vol. 1, Part B, Academic Press Inc. New York, (1984); also S. Mughabghab and C. Dunford, private communication (1982).
2. A. Smith et al., Argonne National Laboratory Report, ANL/NDM-70 (1982)
3. A. Smith et al., Bull. Am. Phys. Soc. 29 637 (1984).
4. D. Reitmann et al., Nucl. Phys. 48 593 (1963).
5. A. Smith et al., to be published.
6. I. van Heerden et al., Z. Phys. 260 9 ((1973).
7. O. Salnikov et al., Jadernye Konstanty 7 102 (1972).
8. N. Birjukov et al., Yadernaya Fizika 19 1190 (1974).
9. D. Thompson, Phys. Rev. 129 1649 (1965).
10. R. Macklin et al., Nucl. Sci. Eng. 59 12 (1976). Data corrected as per private communication from the authors.
11. L. Veesser et al., Phys. Rev. C16 1792 (1977).
12. J. Frehaut and G. Mosinski, private communication. Data available from the National Nuclear Data Center, Brookhaven National Laboratory (1984).
13. C. Philis and P. Young, CEA Report CEA-R-4676 (1975).
14. M. A. Gardner and R. J. Howerton LLNL Report UCRL-50400, Vol. 18 (1978). These data have been extensively revised, but no new documentation has been issued. The data are available upon request from R. J. Howerton.

15. S. Grimes et al., Phys. Rev. C17 508 (1978).
16. P. Young, Los Alamos Report, LA-10069-PR (1984).
17. E. Bramlitt and R. Fink, Phys. Rev. 131 2649 (1963).
18. H. Blosser et al., Phys. Rev. 110 531 (1958).
19. B. Bayhurst and R. Prestwood, Jour. Inorg. Nucl. Chem. 23 173 (1961).
20. H. Tewes et al., Lawrence Livermore Laboratory Report, UCRL-6028-T (1960).
21. R. Haight, National Bureau of Standards Publication, NBS-SP-594 (1979).
22. B. Strohmeier, Ann. Nucl. En. 9 397 (1982).
23. M. Blann, Private Communication. (1984).
24. S. Sudar and J. Csikai, Nucl. Phys. A319 157 (1979).
25. S. Qaim, Private Communication. Data available from the National Nuclear Data Center (1980).
26. V. J. Orphan, N. C. Rasmussen, and T. L. Harper, "Line and Continuum γ -ray Yields from Thermal Neutron Capture in 75 Elements," Gulf General Atomic Report, GA-10248/DASA 2570 (1970).
27. S. T. Perkins, R. C. Haight, and R. J. Howerton, Nucl. Sci. and Eng. 57 1 (1975).
28. T. Reeves and P. Kolkowski, Jour. Phys. G7 529 (1981).
29. B. Strohmeier et al., Physics Data 13 2(1980).

¹⁰⁵Pd
46

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: October 1989
Material: 4634
Content: Fission product

File Comments

ORNL	Eval-Oct89 R. Q. Wright
HEDL	Eval-Feb80 R. E. Schenter and F. Schmittroth
HEDL	Eval-Feb80 F. M. Mann, D. L. Johnson, G. Neely
RCN	Eval-Feb80 H. Gruppelaar

Summary of Changes

The ¹⁰⁵Pd evaluation was modified for ENDF/B-VI by R. Q. Wright in October 1989. The resolved resonance range was revised and extended to 1 KeV. The MLBW formalism was used for this re-evaluation. The highest energy resonance included is 1084.3 eV. The resonance parameters are taken from Ref. 1. The thermal capture cross section for this evaluation is 20.0 barns, which is 43% higher than the ENDF/B-V value. The capture resonance integral is 111.7 barns, which is 13.5% higher than the ENDF/B-V value.

The evaluation was also revised between 1 keV and 1 MeV. Total and elastic cross sections have been increased below 50 keV. The capture cross section has been reduced by about 3 to 10 percent between 1 keV and 1 MeV. The elastic cross section was increased by a very small amount in the range 50 keV to 1 MeV, in order to offset the reduction in the capture cross section. The total cross section is unchanged above 50 keV relative to the ENDF/B-V evaluation.

The revised capture cross section follows the eye guide shown on page 381 of Ref. 2. The capture cross section at 30 keV is 1220 mb which is in good agreement with the value given in Ref. 1, 1190 mb.

The 2200 m/s capture cross section, barns.

(from resonance parameters) = 20.0

computed capture resonance integral

0.5 - 1000 eV = 101.3

above 1000 eV = 10.4

Total = 111.7

References:

1. S. F. Mughabghab, M. Divadeenam, and N. E. Holden, "Neutron Cross Sections," Vol 1A, Academic Press, New York (1981).
2. V. McLane, C. L. Dunford, and P. F. Rose, "Neutron Cross Sections," Vol. 2, Academic Press, New York (1988).

Summary of ENDF/B-V Evaluation

MF=2 MT=151 Resonance parameters from new BNL-325 Ref.(3).

MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.

MF=3 MT= 2 Elastic cross section from $\sigma_t - \sigma_r - \sigma_{in}$ for $E > E_{hi}$.

MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3. Refs.(5, 6).

MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$. The energy region above resonance region was updated by combining available integral and differential data using the generalized least squares adjustment code FERRET (HEDL-TME 77-51)

Summary of ENDF/B-V (Continued)

- MF=4 MT=2 Angular distributions were calculated from the Moldauer potential.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 The evaporation spectrum (LF=9) parameters were obtained using the NCAP code Ref. (2).

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL 325, 3ed., Vol 1 (June 1973).
4. P. A. Moldauer, Nuc. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford, (Private Communication).

¹⁰⁷
₄₆Pd

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: December 1989
Material: 4640
Content: Fission product

File Comments

ORNL	Eval-Dec89 R. Q. Wright
HEDL	Eval-Feb80 R. E. Schenter and F. Schmittroth
HEDL	Eval-Feb80 F. M. Mann, D.L. Johnson, G. Neely
RCN	Eval-Feb80 H. Gruppelaar

Summary of Changes

The resolved resonance range is revised and extended to 1 keV. The MLBW formalism is used for this re-evaluation. The highest energy resonance included is 1082 eV. The resolved resonance parameters are taken from Macklin (Ref. 1). Γ_γ is taken to be constant at 0.125 eV (from Singh et al., Ref. 2). The thermal capture cross section for this evaluation is 2.07 barns, which is 80% lower than the ENDF/B-V value. No measurement of the thermal capture cross section has been reported. In this evaluation, the thermal capture is computed from the positive resonances; a bound level is not included. The capture resonance integral, 110.8 barns, is in excellent agreement with the value given by Macklin (Ref. 1), which is 108.1 ± 4.3 barns. The revised capture resonance integral is 45% higher than the ENDF/B-V value.

The cross sections are also revised for energies above 1 keV. The total and elastic cross sections have been increased below 100 keV and in the range from 1 to 10 MeV. The inelastic cross sections (MT=4 and MT=91) are revised between 2 and 7 MeV. The revised capture cross section follows the data of Macklin (Ref. 1) between 3 and 600 keV. Macklin's data is also shown in Ref. 3 (see page 381). Compared to ENDF/B-V, the revised evaluation is higher below 400 keV and lower above 400 keV. The capture cross section at 30 keV is 1400 mb. From 1 to 10 MeV, the capture cross section has about the same shape as the ENDF/B-V evaluation but the magnitude is 20-50% lower.

The 2200 m/sec capture cross section, barns

(from resonance parameters) = 2.07

computed resonance integral

0.5 eV - 1 keV = 99.4

above 1 keV = 11.4

Total = 110.8

References:

1. R. L. Macklin, "Neutron Capture Measurements on Fission Product Pd-107," Nucl. Sci. and Eng. 89, 79-86 (1985).
2. U. N. Singh, R. C. Block, and Y. Nakagome, Nucl. Sci. and Eng. 67, 54 (1978)
3. V. McLane, C. L. Dunford, and P. F. Rose, "Neutron Cross Sections," Vol. 2, Academic Press, New York (1988).

Summary of ENDF/B-V Evaluation

MF=2 MT=151 No resonance parameters given except for AP.

MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.

MF=3 MT= 2 Elastic cross section from $\sigma_t = \sigma_e + \sigma_m$ for $E > E_{hi}$, and from $4\pi a^2$ for $E < E_{hi}$.

MF=3 MT= 4, 51,52,,,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).

MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref.(3) for $E < E_{hi}$. The energy region above the resonance region was updated by combining available integral and differential data using the generalized least squares adjustment code FERRET (HEIDL-TME 77-51).

Summary of ENDF/B-V (Continued)

- MF=4 MT=2 Angular distribution calculated from the Moldauer Potential.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 Evaporation spectrum (LF=9) parameters were obtained using the NCAP code Ref. (2).

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. E. Clayton, AAEC/TM 619 (Sept 1972).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford, (Private Communication).

^{nat}
49 **In**

Reference: ANL/NDM-115
Evaluators: A. Smith, S. Chiba, D. Smith, J. Meadows, P. Guenther,
R. Lawson (ANL), and R. Howerton (LLNL)
Evaluated: February 1990
Material: 4900
Content: Neutron transport, Gamma production, Covariances

1. Introduction

Indium has been used in nuclear applications (primarily as a dosimeter) for a half century; it is employed in superconductors, appears as a fission product, and has a large (n,2n) cross section making it a good multiplier. The element consists of two isotopes ¹¹³In (4.3%) and ¹¹⁵In (95.7%). Owing to ENDF format considerations the evaluation of the ¹¹⁵In(n,n')^{115m}In reaction was not included in this general purpose file for elemental indium. Consequently it has been placed in a special ¹¹⁵In file intended for dosimetry purposes (Mat = 4931).

2. Evaluated Resolved Resonance Range

Resonance parameters appropriate to the two isotopes are used to describe the neutron interactions with indium up to 2 keV. The parameters are taken from Mughabghab¹ with small changes in the scattering radius to agree with experiment.

3. Evaluated Total Cross Sections

The evaluation is based upon 23 citations obtained from the NNDC.² The average age of the data is about 25 years, with only 4 citations in the last decade. Some of the data were clearly inconsistent with the body of information, and were not used. The accepted data sets were averaged over 100 keV intervals to 1 MeV, 200 keV intervals from 1 - 2 MeV, and larger intervals above 2 MeV. Subjective estimates were made for noted systematic differences. The energy averaged data base was evaluated using the statistical procedures of the GMA code.³ The two combined isotopic evaluations of ENDF/B-V differ by $\approx 10\%$ or so with the present evaluation.

4. Evaluated Elastic Scattering Cross Sections

The energy averaged neutron elastic scattering cross sections extend from 2 keV to 20 MeV. Up to 15 MeV they are based on the detailed study of differential elastic scattering described by A. B. Smith et al. in Refs. 4 & 5. Above 15 MeV the model described in Ref. 5 was used to extrapolate the cross sections to 20 MeV. There are large differences (factors of 2 at 20 MeV) from ENDF/B-V. These differences also imply large differences in the non-elastic cross sections of the two files.

5. Evaluated Inelastic Scattering Cross Sections

5.1 Discrete Inelastic Processes

Primary attention was given to the excitation of discrete levels in ^{115}In . These have been carefully studied in a cooperative experimental program.¹ The low energy model reasonably matches the higher energy model of Ref. 5 at an energy of several MeV. Sixteen levels of ^{115}In were considered up to excitations of ≈ 1.5 MeV, with excitation energies and J^π values taken from Ref. 4. The cross sections were calculated using the optical statistical model⁵ with results essentially identical with those given in Ref. 4 and supported by experimental results. For completeness the same method was used to determine the discrete inelastic scattering cross sections of the minor ^{113}In isotope. In this case 12 excited levels below 1.5 MeV were used with the excitations and J^π values from Ref. 6.

5.2 Continuum Inelastic Scattering Processes

Above 1.5 keV the continuum inelastic scattering cross section rises rapidly to large values exceeding 2 barns. The evaluation determines the continuum inelastic scattering cross section from the difference between the non-elastic cross section and the other partial cross sections. Below 10 MeV the major contribution is from the discrete inelastic scattering cross section, and above the $(n,2n)$ cross section rises rapidly with a complimentary sharp decrease in the continuum inelastic scattering which falls to ≈ 200 mb at 20 MeV. Above 16 MeV the $(n,3n)$ cross section becomes a factor as well. The inelastic scattering cross sections of the present evaluation are grossly different from those given in ENDF/B-V. Below 10 MeV the two evaluations differ by $\approx 20\%$. At higher energies the differences are even larger, amounting to 500% at 20 MeV.

The continuum neutron spectra emitted as a result of the inelastic scattering process were estimated from experimental measurements below 8 MeV.⁷ Above 8 MeV the individual spectra were calculated using the computer code ALICE⁸ and CADE⁹. The parameters of ALICE were adjusted so that the ratios $(n,n')/(n,2n)$ and $(n,3n)/(n,2n)$ agreed with the values obtained in the evaluation; then the spectra

associated with each component of the individual reactions were calculated using the methods described in Ref. 10.

6. Evaluated Radiative Capture Cross Sections

The data base consisted of measured values available at the National Nuclear Data Center. These data were primarily obtained using prompt detection techniques with some activation results. The data scatter is large, the majority of measurements are below 100 keV, and the cross section is relatively large (i.e., 200 mb) up to more than an MeV. The evaluation is based on a single giant dipole resonance calculation employing the model of Ref. 11 with the S_0 strength function adjusted to obtain what was subjectively judged to be a "best" description of the measured values. The estimated uncertainties are quite large; $\approx 10 - 15\%$ up to 100 keV and $15 - 25\%$ from 100 keV to 2 MeV. The ENDF/B-V values are generally much smaller. Only one data set supports the ENDF/B-V evaluation, and then only over a limited range.

7 Evaluated (n,2n) and (n,3n) Reactions

Experimental knowledge of the (n,2n) cross section is based on activation measurements. For both indium isotopes the primary activity is due to the decay of a metastable state. The evaluation is primarily based upon the experimental data supported by statistical model calculations using CADE.⁹ The isomer activation ratio m/g is $\approx 4.5(\pm 15\%)$ at 14 MeV. It was assumed that this ratio was constant throughout the energy range. The evaluated $^{115}\text{In}(n,2n)$ cross sections were constructed from the $^{115}\text{In}(n,2n)^{116m}\text{In}$ evaluated cross sections. The evaluation assumes that the $^{115}\text{In}(n,2n)$ cross sections are equivalent to those of the element with a slightly lower (~ 0.8 MeV) threshold than the $^{113}\text{In}(n,2n)$ reaction.

Only one measurement of the In (n,3n) has been reported.¹² It involves only the 2.8 day activity from the $^{113}\text{In}(n,3n)^{111}\text{In}$ reaction. A reasonable extrapolation of that data gives an ^{113}In cross section of ≈ 120 mb at 20 MeV. The ^{115}In (n,3n) threshold is ≈ 0.81 MeV lower than that of the ^{113}In (n,3n) reaction, and due to the rapid increase of the cross section with energy it is reasonable to expect the $^{115}\text{In}(n,3n)$ cross section to be 400 to 500 mb at 20 MeV. Calculations using ALICE and CADE predict somewhat lower cross sections. The evaluated (n,3n) cross sections are based upon the difference between the experimentally based (n,2n) cross section and the general energy dependent trend of the reaction cross section. They are somewhat larger than suggested by the above experimental evidence, but less than the prediction of calculations. It is impossible to compare the present evaluation with the two ENDF/B-V isotopic files as the latter do not contain these reactions.

8 Evaluated Charged Particle Emitting Reactions

In the present evaluation the interactions with the prominent isotope ^{115}In are considered. See table 1. below. The respective Q values were taken from Ref. 13.

Table 1

Q-values for Charged Particle Emitting Reactions

Reaction	Q-value (MeV)
(n,p)	-0.666
(n,np)	-6.811
(n,d)	-4.587
(n,nd)	-13.627
(n,t)	-7.370
(n,nt)	-13.914
(n, ^3He)	-9.362
(n, n ^3He)	-17.853
(n, α)	+2.726
(n,n α)	-3.740

All the energetically allowed processes were calculated using CADE with the addition of a pre-compound component determined using the code ALICE. The calculated results were compared with available experimental information and adjusted where judged appropriate, to obtain evaluated quantities. The experimental data base is very weak, however much of the evaluation is based solely on statistical calculations.

8.1 (n,p) and (n,np) Reactions

The experimental data base is limited to nine measurements all near 14 MeV. The cross section resulting in the activation of the ground state has been measured 6 times with various results. Ignoring two exceptional values the cross section seems to be between 4 and 5 mb at 14 MeV. A single measurement of the cross section for the excitation of the metastable state at 14.8 MeV gives 7.7 ± 1.2 mb. Thus the fragmentary experimental evidence suggests an (n,p) cross section of 10 – 15 mb at 14 – 15 MeV. The calculations indicate that the cross section is largely due to pre-compound processes, and near 14 MeV the ALICE result was ≈ 14 mb in reasonable agreement with the experimental evidence. The Alice results have been used without renormalization for the (n,p) and (n,np) reactions.

8.2 (n, α) and (n,n α) Reactions

The $^{115}\text{In}(n,\alpha)$ process results in ^{112}Ag which has a 3.14 hour activity and can be reasonably measured. The results are closely grouped between 2.5 to 3.0 mb at ≈ 14 MeV, with an average of 2.7 mb at 14.25 MeV. The CADE and ALICE results were much smaller than the experimental values in the 14 MeV region, possibly due to not including pre-compound processes. The data was renormalized to the experimental values near 14 MeV and the same normalization factor was used to obtain the (n,n α) evaluation from the calculations.

9. Evaluated Photon Production Reactions

The spectrum of photons from the neutron capture reaction was taken from the work of Orphan et al.¹⁴ at thermal energy. The same spectrum was used at 20 MeV with the multiplicity adjusted to conserve energy.

For photons associated with the inelastic scattering to specific levels Warren's code CASCADE¹⁵ which incorporates the method used in Reffo's BRANCH code¹⁶ was used.

For all other reactions the photon production cross sections and spectra were calculated using the R-parameter formalism of Perkins et al.¹⁷ Since the ENDF/B-VI formats and procedures allow for secondary charged particle distributions in File 5 only if there is a single secondary particle, the file was translated to the ENDL format where energy distributions for all secondaries can be represented. The R(U) values were taken from the "global" values of Ref. 17.

References

1. S. F. Mughabghab, Neutron Cross Sections Vol. 1, Part B, Academic Press Inc. New York, (1984); also S. Mughabghab and C. Dunford, private communication (1982).
2. National Nuclear Data Center, Brookhaven National Laboratory, Upton, New York 11973.
3. W. P. Poenitz, Brookhaven National Laboratory Report, BNL-NCS-51363 Vol. I 249(1981); as modified by M. Sugimoto (1987).
4. A. Smith, P. Guenther, J. Whalen, I. Van Heerden and W. McMurray, J. Phys G11 125 (1985)
5. S. Chiba, P. T. Guenther, R. D. Lawson, and A. B. Smith, Argonne National Laboratory Report, ANL/NDM-116 (1990)

6. C. Lederer and V. Shirley, eds., Table of Isotopes, 7th Edition, John Wiley and Sons Inc. New York (1978).
7. P. Guenther, Report to the IAEA Coordinated Research Program on the Measurement and Analysis of Double-Differential Neutron Emission Spectra in (p,n) and (α ,n) Reactions (1989).
8. M. Blann, Lawrence Livermore National Laboratory Report, UCID-20169 (1984)
9. D. Wilmore, Harwell Report AERE-R-11515 (1984).
10. P. Guenther et al., Argonne National Laboratory Report, ANL/NDM-107 (1988)
11. P. Moldauer, Private Communication (1982).
12. H. Liskien, Nucl. Phys. A118 379 (1968).
13. R. Howerton, Tabulation of Q-values, Informal LLNL report.
14. V. J. Orphan, N. C. Rasmussen, and T. L. Harper, "Line and Continuum γ -ray Yields from Thermal Neutron Capture in 75 Elements," Gulf General Atomic Report, GA-10248/DASA 2570 (1970).
15. W. E. Warren, R. J. Howerton, and G. Reffo, CASCADE Cray program for γ -production from discrete level inelastic scattering, Lawrence Livermore Nuclear Data Group Internal Report, PD-134 (1986), unpublished.
16. G. Reffo, IDA - A modular system of nuclear model codes for the calculation of cross sections for nuclear reactors, Centro Ricerche Energia, Bologna, unpublished (1980).
17. S. T. Perkins, R. C. Haight, and R. J. Howerton, Nucl. Sci. and Eng. 57 1 (1975).

¹¹⁵₄₉In

Reference: ANL/NDM-115
Evaluators: R. E. Schenter and F. Schmittroth, Activation
S. Chiba, and D. L. Smith, Dosimetry
Evaluated: March 1990
Material: 4931
Content: Activation, Dosimetry

File Comments

ANL Eval-Jan90 S. Chiba and D. L. Smith
HEDL Eval-Feb84 R. E. Schenter and F. Schmittroth

* * * * * * * * * * * *

The ¹¹⁵In file was updated at ANL by S. Chiba, D. L. Smith, and A. B. Smith in January 1990. The dosimetry reaction ¹¹⁵In(n,n')^{115m}In was revised extensively.

Summary of Changes

The production of the isomer ^{115m}In by the (n,n') process is routinely employed for neutron dosimetry applications. This isomer is the first-excited state of the isotope ¹¹⁵In (336 keV excitation energy). The reaction threshold energy is 339 keV. The isotopic abundance of ¹¹⁵In in natural indium is 95.7%. The half life of ^{115m}In is 4.486 hours. The decay modes are - β⁻ (5 percent) and Isomeric Transition (95.0%). The number of Decay 336-keV γ-rays emitted per disintegration of ^{115m}In is 0.459.

The documentation for the ¹¹⁵In(n,n')^{115m}In dosimetry reaction is provided by A. B. Smith et al. Report ANL/NDM-115, Argonne National Laboratory (1999).¹

The available differential data was assembled from the literature as determined from CINDA and CSISRS. A total of 32 experimental data sets (147 data points) were included in the present evaluation. Nuclear model calculations were performed with the code ABAREX² to determine the theoretical cross section shape close to threshold. The evaluation itself was carried out with the least squares adjustment code GMA as described by W. Poenitz in 1981³ and later revised by M. Sugimoto (1987) and S. Chiba in 1990.⁴ The earlier evaluation of D. L. Smith in ANL/NDM-26 was used to establish an *a priori* cross section shape.

The present evaluation tends to be a few percent larger than ENDF/B-V. Mannhart has evaluated the available experimental integral data (averaged over a ²⁵²Cf

spontaneous fission spectrum) and obtained 197.6 mb ($\pm 1.4\%$).⁵ Using Mannhart's spectral data the present evaluation gives 189.6 mb ($\pm 2.2\%$). This leads to a C/E = 0.96. In this respect the present evaluation represents a significant improvement over the earlier evaluation.

References

1. A. B. Smith, S. Chiba, D. L. Smith, J. W. Meadows, P. T. Guenther, R. D. Lawson, and R. J. Howerton, ANL/NDM-115, Argonne National Laboratory (1990).
2. ABAREX, "A Spherical Optical Model Code", P. Moldauer, Private Communication (1983), and as revised by R. D. Lawson (1986).
3. W. P. Poenitz, Brookhaven National Laboratory Report, BNL-NCS-51363 Vol. I 249 (1981); as modified by M. Sugimoto (1987).
4. S. Chiba, P. T. Guenther, R. D. Lawson, and A. B. Smith, ANL/NDM-116, Argonne National Laboratory (1990)
5. W. Mannhart, "Reactor Dosimetry: Methods, Applications, and Standardization." H. Farrar IV and E. Lippincott, Eds., American Society for Testing Materials, ASTM STP-1001, Philadelphia, p. 340 (1989).

* * * * *

Summary of Previous Evaluation

MF=1 MT=451	Atomic Mass from Ref (1).
MF=2 MT=151	Evaluation of Resolved Resonance Parameters is based on new BNL-325, Ref (2).
MF=3 MT=51	The evaluation of the 4.486 hour isomer is based entirely on reported experimental data. The documentation is available as ANL/NDM-26 by D. L. Smith. References 10 through 25, listed below, were used in this evaluation.

Summary of Previous Evaluation (Continued)

MF=3 MT=102 Version-V unresolved region contains adjusted data. See ANL documentation. The radiative neutron capture to the ^{110m}In (54 min.) state was evaluated. For $E > E_{hi}$, the evaluation is based on experimental data, Ref.(3 - 7) and theoretical calculations, Ref.(8, 9). For $E < E_{hi}$, a $1/v$ component was added to give the correct 2200 m/s cross section to the 54 min. state (the 2.2 sec. state cross section was included). The radiative capture to the 2.2 sec. state of ^{116}In was included as part of the capture to the 54 min. state for both thermal and fast energies. The results were divided by 0.79 to give the total capture cross section in File 3. In 1984 R. Schenter added File 9 with multiplicity 0.79, and modified the total capture width in File 2 to be $\Gamma_\gamma = \Gamma_\gamma^m/0.79$. File 9 combined with File 3 is required to produce the capture for the 54 min. isomeric state.

The 2200 m/sec capture cross section (to the 54 min. state) computed from the resonance parameters is 166.4 barns. The computed resonance integral is 2587.3 barns.

References

1. A. H. Wapstra and N. B. Gove, Nuclear Data Tables, Vol. 9, Part 1(1971).
2. S. F. Mughabghab and D. I. Garber, BNL-325, 3rd ed., Vol.1 (1973).
3. H. A. Grench and H. O. Menlove, Phys. Rev. **165**, 1298 (1968).
4. H. O. Menlove, et al., Phys. Rev. **163** 1299 (1967).
5. S. A. Cox, Phys. Rev. **133**, B378 (1964).
6. A. E. Johnsrud et al., Phys. Rev. **116**, 927 (1959).
7. G. Peto et al., J. Nucl. En. **21**, 797 (1967)..
8. F. Schmittroth, HEDL-TME 71-106 (August 1971).
9. F.Schmittroth, HEDL-TME 73-79, ENDF-195 (November 1973).
10. D. L. Smith et.al., ANL/NDM-14, (1975).

11. D. C. Santry and J. P. Butler, *Can. J. Phys.* 54, 757 (1975).
12. K. Kobayashi et.al., *J. Nuc. En.* 27, 741 (1973).
13. A. Paszit and J. Osikai, *Sov. J. Nuc. Phys.* 15, 232 (1972).
14. J. K. Temperly and D. E. Barnes, BRL-1491 (1970).
15. P. Decowski et al., INR-1197 Poland (1970).
16. I. Kimura et al., *J. Nuc. Sci. Tech. Japan* 6, 485 (1969).
17. R. C. Barrall et al., AFWL-TR-68-134, (1969).
18. H. Roetzer, *Nucl. Phys.* A109, 694 (1968).
19. B. Minetti and A. Paquaretti, *Z. Phys.* 217, 83 (1968).
20. H. A. Grench and H. O. Menlove, *Phys. Rev.* 165, 1298 (1968).
21. H. O. Menlove et al., *Phys. Rev.* 163, 1308 (1967).
22. W. Nagel and A. H. W. Aten Jr., *Physica* 31, 1091 (1965).
23. A. A. Abel and C. Goodman, *Phys. Rev.* 93, 197 (1954).
24. H. C. Martin and B. C. Diven, *Phys. Rev.* 93, 199 (1954).
25. S. G. Cohen, *Nature* 161, 475 (1948).

¹³⁴₅₅Cs

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, and F. Schmittroth
Evaluated: December 1988
Material: 5528
Content: Fission product

File Comments

HEDL Eval-Apr74 R. E. Schenter and F. Schmittroth
ORNL Eval-Dec88 R. Q. Wright

Summary of Changes

The ENDF/B-V ¹³⁴Cs evaluation, MAT 9663, has been revised below 180 ev. The revised evaluation has been assigned MAT 5528 in order to differentiate it from the original evaluation. In the revised evaluation, resolved resonance parameters are used to define the total, elastic, and capture cross sections below 180 ev. Above 180 ev the evaluation is unchanged from ENDF/B-V.

The resolved resonance parameters are taken from Ref. (1). It should be noted that the 42.13 eV level given in Ref. (1) must be assigned to ¹³⁵Cs (See Ref. 2). The MLBW (LRF=2) representation was used with the smooth background cross sections set to zero in the resonance region. The largest contribution to the thermal capture cross section (almost 100%) is from the bound level at -14 eV.

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	139.64
computed resonance integral		
(from resonance parameters)	=	53.27
above 180 ev	=	24.79
Total	=	78.06

References:

1. S. F. Mughabghab, M. Divadeenam, and N. E. Holden, "Neutron Cross Sections," Vol 1A, Academic Press, New York (1981).
2. H. G. Priesmeyer, "Low Energy Neutron Cross Section Measurements of Radioactive Fission Product Nuclides," Proc. Specialists Mtg. on Neutron Cross Sections of Fission Product Nuclei, Bologna, Italy, Dec. 12-14, 1979, NEANDC(D)-209, 1, p77 (1980).

Summary of ENDF/B-V Evaluation

Comment cards for are for the ENDF/B-IV evaluation which was translated into ENDF/B-V formats by F. M. Mann and R. E. Schenter (HEDL) in January 1980 as MAT 9663.

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0.0
(from 1/V component)	=	140.0
Total	=	140.0

computed resonance integral = 212.9

MF=2 MT=151 No resonance parameters given except for AP.

MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.

MF=3 MT= 2 Elastic cross section from $\sigma_t = \sigma_e + \sigma_m$ for $E > E_{hi}$, and from $4\pi a^2$ for $E < E_{hi}$.

MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs.(5, 6).

Summary of ENDF/B-V (Continued)

- MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{th}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{th}$. The low energy capture was also adjusted to give a resonance integral (to within 1σ) of Ref. (7).
- MF=4 MT=2 Angular distribution assumed isotropic.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT=91 Evaporation spectrum (LF=9) parameters were obtained using the NCAP code Ref. (2).

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL 325, 3ed, Vol. 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).
7. P. Ribon and J. Krebs, Bologna Panel Report (April 1974).

¹³⁴Ba
56

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: December 1988
Material: 5637
Content: Fission product

File Comments

ENDF/B-VI MAT 5637 Evaluated by R. Q. Wright (ORNL)
ENDF/B-V MAT 9684 Evaluated by R. E. Schenter and F. Schmittroth (HEDL)
File converted to ENDF-6 Format by the NNDC

Summary of Changes

The ¹³⁴Ba evaluation, MAT 9684, was revised by R. Q. Wright, June 1988. The new evaluation is assigned MAT No. 5637. The resolved resonance parameters for MAT 5636 are from Ref. 1 ($E_R = 2071.8$ eV). The bound level at -104 eV has $\Gamma_n = 0.347$ eV and $\Gamma_\gamma = 0.114$ eV; this choice gives the desired value of 1.98 b for the thermal capture cross section. Values of Γ_γ not given in Ref. 1 are set to 0.120 eV. The value for the scattering radius is 0.61725 fm (unchanged). The highest energy resonance included is 1892.0 eV.

In File 3 total, elastic, and capture cross sections are set to zero in the resolved resonance range (10^{-5} to 2071.8 eV.)

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	1.98
(from 1/v component)	=	0.00
total	=	1.98

computed resonance integral = 24.11

References:

1. S. F. Mughabghab, "Neutron Cross Sections" Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part A: Z=1-60, Academic Press (1981).

Summary of ENDF/B-V Evaluation

- MF=2 MT=151 Resonance parameters from BNL-325, Ref. (3).
- MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.
- MF=3 MT= 2 Elastic cross section from $\sigma_t = \sigma_e + \sigma_{in}$ for $E > E_{hi}$.
- MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs.(5, 6).
- MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$. The calculated resonance integral agrees (to within 1σ) with the value given in Ref. (3).
- MF=4 MT=2 Angular distribution assumed isotropic.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 Evaporation spectrum (LF=9) parameters obtained using NCAP code Ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0.485
(from $1/v$ component)	=	1.673
Total	=	2.158

computed resonance integral = 23.897

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL-325, 3ed, Vol. 1 (June 1973).
4. P. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

						$^{134}_{56}\text{Ba } \sigma_{n,\gamma}$
78	ORL	Musgrove+	78HARWELL, 449	15	3.500 keV to 0.175 MeV	$^6\text{Li } \sigma_{n,t}$

¹³⁵₅₆Ba

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: December 1988
Material: 5640
Content: Fission product

File Comments

ENDF/B-VI MAT 5640 Evaluated by R. Q. Wright (ORNL)
ENDF/B-V MAT 9685 Evaluated by R. E. Schenter and F. Schmittroth (HEDL)
File converted to ENDF-6 Format by the NNDC

Summary of Changes

The ¹³⁵Ba evaluation, MAT 9685 was revised by R. Q. Wright, June 1988. The new evaluation is assigned MAT No. 5640. The resolved resonance parameters are from Ref. 1 ($E_{hi}=1650.0$ eV). The bound level at -51 eV has $\Gamma_n = 0.1824$ eV and $\Gamma_\gamma = 0.140$ eV; this choice gives the desired value of 5.81 b for the thermal capture cross section. Values of Γ_γ not given in Ref. 1 are set to 0.150 eV. The value for the scattering radius is 0.61880 fm (unchanged). The highest energy resonance included is 1621.0 eV.

In File 3 total, elastic, and capture are set to zero in the resolved resonance range (10^{-5} to 1650 eV).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	5.81
(from 1/v component)	=	0.00
total	=	5.81

computed resonance integral	=	99.34
-----------------------------	---	-------

References:

1. S. F. Mughabghab, "Neutron cross sections" Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part A: Z=1-60, Academic Press (1981).

Summary of ENDF/B-V Evaluation

- MF=2 MT=151 Resonance parameters from BNL-325, Ref. (3).
- MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.
- MF=3 MT= 2 Elastic cross section from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$.
- MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).
- MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$. The calculated resonance integral agrees (to within 1σ) with the value given in Ref.(3).
- MF=4 MT=2 Angular distributions assumed isotropic.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 Evaporation spectrum (LF=9) parameters obtained using NCAP code Ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	2.133
(from 1/v component)	=	3.681
Total	=	5.814

computed resonance integral = 100.555

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Shmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL 325, 3ed, Vol. 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

$^{135}_{86}\text{Ba } \sigma_{n,\gamma}$						
74	ORL	Musgrove+	AAEC/E-327	15	3.500 keV to 0.175 MeV	$^6\text{Li } \sigma_{n,t}$

¹³⁶Ba
56

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: December 1988
Material: 5643
Content: Fission product

File Comments

ENDF/B-VI MAT 5643 Evaluated by R. Q. Wright (ORNL)
ENDF/B-V MAT 9687 Evaluated by R. E. Schenter and F. Schmittroth (HEDL)
File converted to ENDF-6 Format by the NNDC

Summary of Changes

The ¹³⁶Ba evaluation, MAT 9687 was revised by R. Q. Wright, June 1988. The new evaluation is assigned MAT No. 5643. The resolved resonance parameters are from Ref. 1 ($E_{hi}=3177.2$ eV). The bound level at -250 eV has $\Gamma_n = 0.759$ eV and $\Gamma_\gamma = 0.125$ eV; this choice gives the desired value of 0.41 b for the thermal capture cross section. Values of Γ_γ not given in Ref. 1 are set to 0.125 eV. The value for the scattering radius is 0.62032 fm (unchanged). The highest energy resonance included is 1644.0 eV.

In File 3 total, elastic, and capture cross sections are set to zero in the resolved resonance range (10^{-5} to 3177.2 eV).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0.41
(from 1/v component)	=	0.00
total	=	0.41

computed resonance integral = 1.72

Reference

1. S. F. Mughabghab, "Neutron cross sections" Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part A: Z=1-60, Academic Press (1981).

Summary of ENDF/B-V Evaluation

MF=2 MT=151 Resonance parameters from BNL-325 Ref. (3).

MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.

MF=3 MT= 2 Elastic cross section from $\sigma_t = \sigma_e + \sigma_{in}$ for $E > E_{hi}$.

MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).

MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$.

MF=4 MT=2 Angular distributions assumed isotropic.

MF=4 Non-elastic angular distributions assumed isotropic.

MF=5 MT= 91 Evaporation spectrum (LF=9) parameters obtained using NCAP code Ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0.020
(from $1/v$ component)	=	0.390
Total	=	0.410

computed resonance integral = 1.958

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL 325, 3ed, Vol. 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

$^{136}_{56}\text{Ba } \sigma_{n,\gamma}$

78	ORL	Musgrove+	78HARWELL, 449	15	3.500 keV to 0.175 MeV	$^6\text{Li } \sigma_{n,t}$
71	AUA	Stroud+	AAEC/PR-34P, 9	1	80.00 mb at 30.00 keV	$^{197}\text{Au } \sigma_{n,\gamma}$

¹³⁷₅₆Ba

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: December 1988
Material: 5646
Content: Fission product

File Comments

ENDF/B-VI MAT 5646 Evaluated by R. Q. Wright (ORNL)
ENDF/B-V MAT 9689 Evaluated by R. E. Schenter and F. Schmittroth (HEDL)
File converted to ENDF-6 Format by the NNDC

Summary of Changes

The ¹³⁷Ba evaluation, MAT 9689 was revised by R. Q. Wright, June 1988. The new evaluation is assigned MAT No. 5646. The resolved resonance parameters are from Ref. 1 ($E_{hi} = 1947.5$ eV). The bound level at -26 eV has $\Gamma_n = 0.081$ eV and $\Gamma_\gamma = 0.083$ eV; this choice gives the desired value of 5.10 b for the thermal capture cross section. Values of Γ_γ not given in Ref. 1 are set to 0.080 eV. The value for the scattering radius is 0.62184 fm (unchanged). The highest energy resonance included is 1737.0 eV.

In File 3 total, elastic, and capture cross sections are set to zero in the resolved resonance range (10^{-5} to 1947.5 eV).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	5.10
(from 1/v component)	=	0.00
Total	=	5.10

computed resonance integral	=	3.92
-----------------------------	---	------

Reference

1. S. F. Mughabghab, "Neutron cross sections" Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part A: Z=1-60, Academic Press (1981).

Summary of ENDF/B-V Evaluation

- MF=2 MT=151 Resonance parameters from BNL-325 Ref. (3).
- MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.
- MF=3 MT= 2 Elastic cross section from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$.
- MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).
- MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$. The calculated resonance integral agrees (to within 1σ) with the value given in Ref. (3).
- MF=4 MT=2 Angular distributions assumed isotropic.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 Evaporation spectrum (LF=9) parameters obtained using NCAP code Ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0.071
(from $1/v$ component)	=	5.030
Total	=	5.101

computed resonance integral = 4.949

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL 325, 3ed Vol 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).

¹⁴⁷₆₀Nd

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, and F. Schmittroth
Evaluated: December 1988
Material: 6040
Content: Fission product

File Comments

HEDL Eval-Apr74 R. E. Schenter and F. Schmittroth
ORNL Eval-Dec88 R. Q. Wright

Summary of Changes

The ENDF/B-V ¹⁴⁷Nd evaluation, MAT 9768, has been revised below 35 eV. The revised evaluation has been assigned MAT No. 6040 in order to differentiate it from the original evaluation. In the revised evaluation, resolved resonance parameters are used to define the total, elastic, and capture cross sections below 35 eV. Above 35 eV the evaluation is unchanged from ENDF/B-V. The resolved resonance parameters are taken from Ref. (1). The MLBW (LRF=2) representation was used with the smooth background set to zero in the resonance region. The largest contribution to the thermal capture cross section (about 98%) is from the bound level at - 5 eV. The thermal capture cross section is higher than the ENDF/B-V value by about a factor of 9. The capture resonance integral is slightly lower.

The 2200 m/s capture cross section, barns

(from resonance parameters) = 439.6

computed resonance integral

(from resonance parameters) = 431.3

above 35 eV = 144.0

Total = 575.3

Reference

1. S. F. Mughabghab, M. Divadeenam, and N. E. Holden, "Neutron Cross Sections," Vol. 1A, Academic Press, New York (1981).

Summary of ENDF/B-V Evaluation

- MF=2 MT=151 No resonance parameters given except AP.
- MF=3 MT= 1 Total cross section calculated using Moldauer Potential from Ref. (4) for $E > E_{hi}$.
- MF=3 MT= 2 Elastic cross section from $\sigma_t = \sigma_e + \sigma_{in}$ for $E > E_{hi}$, from $4\pi a^2$ for $E < E_{hi}$.
- MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).
- MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs.(1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref.(3) for $E < E_{hi}$. The low energy capture was also adjusted to give the resonance integral (to within 1σ) of Ref. (7).
- MF=4 MT=2 Angular distributions assumed isotropic.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 Evaporation spectrum (LF=9) parameters were obtained using the NCAP code ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0.0
(from $1/v$ component)	=	49.0
Total	=	49.0

computed resonance integral = 647.8

This file was translated into ENDF-5 format by F. M. Mann and R. E. Schenter (HEDL) in January 1980.

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. E. Clayton, AAEC/TM 619 (Sept 1972).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).
7. P. Ribon and J. Krebs, Bologna Panel Report (April 1974).

¹⁴⁷₆₁Pm

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: April 1989
Material: 6149
Content: Neutron transport, Fission product

File Comments

ORNL	Eval-Apr89 R. Q. Wright
HEDL	Eval-Feb80 R. E. Schenter and F. Schmittroth
HEDL	Eval-Feb80 F. M. Mann, D. L. Johnson, G. Neely
RCN	Eval-Feb80 H. Gruppelaar
BNL	Eval-Oct74 A. Prince

The Pm-147 evaluation, Mat 9783, was revised by R. Q. Wright in April 1989. The new evaluation is assigned Mat. No. 6149. ¹⁴⁷Pm is an isotope of considerable importance to reactor neutron economy. This is due to its effect on the growth, during reactor operation, of ¹⁴⁹Sm, which is a very serious reactor poison. For this reason it is important to have accurate values of the ¹⁴⁷Pm thermal capture cross section and capture resonance integral. ¹⁴⁷Pm has a half-life of 2.62 years and decays to ¹⁴⁷Sm.

Summary of Changes

The resolved resonance parameters were taken from Ref. 1 ($E_{hi} = 300.0$ eV). Two bound levels at -22.1 and -8.88 eV are used in this evaluation. The contribution from the bound levels to the thermal capture cross section is 83.5 b. The other resonance contribution is 84.9 b. Thus, the thermal capture cross section is 168.4 barns, which is about 8 % lower than the ENDF/B-V evaluation. In addition, the total cross section is 190.5 b, which is about 5 % below some old experimental values. Values of Γ_γ not given in Ref. 1 are set to 0.067 eV. The value for the scattering radius is 0.83E-12 cm. The upper limit of the resolved resonance range is increased from 58.078 to 300.0 eV, and the highest energy resonance included is 316.5 eV.

Unresolved resonance parameters were added to the file. The unresolved range is 300 eV to 20 keV. The unresolved parameters are based on $D_0 = 3.6$ eV and $S_0 = 3.1$.

Total, elastic, and capture cross sections were set to zero in the resolved and unresolved resonance ranges (1.0E-05 eV to 20 keV).

The 2200 m/s capture cross section, barns.

(From resonance parameters)	=	168.4
(From 1/v component)	=	0.0
Total	=	168.4

Computed resonance integral	=	2197
-----------------------------	---	------

References:

1. S. F. Mughabghab, "Neutron cross sections: Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B: Z=61-100," Academic Press (1984).

Summary of ENDF/B-V Evaluation

MF=2 MT=151	Resonance parameters from new BNL-325 Ref. (3).
MF=3 MT= 1	Total cross section calculated with a deformed potential from Ref. (4) for $E > E_{hi}$.
MF=3 MT= 2	Elastic cross section from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$.
MF=3 MT= 4,	51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5,6). The level scheme data is from the Nuclear Data Tables and S. Igarasi(Japan) private communication.
MF=3 MT= 16(n,2n), 17(n,3n), 22(n,nd), 28(n,np), 103(n,p), 104(n,d), 105(n,t), 106(n, ³ He), 107(n, ⁴ He)	calculated using the THRESH code Ref. (7).

Summary of ENDF/B-V (Continued)

- MF=3 MT=102 The neutron capture was evaluated using COMNUC-3 and NCAP in Refs. (1,2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E > E_{hi}$. The energy region above the resonance region was updated by combining available integral and differential data using the generalized least squares adjustment code FERRET (HEDL-TME 77-51).
- MF=4 MT=2 The angular distribution was calculated from the Moldauer potential.
- MF=4 Non-elastic energy distributions assumed isotropic.
- MF=5 MT=16, 17,22,28,91 Energy distributions of secondary neutrons given as a histogram using calculations of nuclear temperature from reference (11).

References

1. T. Tamura, Computer program JUPITOR I for coupled-channel calculations, ORNL-4152 (1967).
2. F. Schmittroth, HEDL TME 73-79(Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL 325, 3ed., Vol. 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47(1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (private communication).
7. S. Pearlstein, Jour. Nucl. Energy 27, 81 (1973).
8. H. Baba and S. Baba, JAERI 1183 (1969).
9. G. Lautenbach, RCN-191 (1973).
10. S. M. Zakharova et al., INDC (CCP)-27/1.
11. A. Gilbert and A. G. W. Cameron, Can. J. Phys. 43, 1446 (1965).

¹⁴⁷₆₂Sm

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, F. M. Mann, A. Prince,
Others
Evaluated: April 1989
Material: 6234
Content: Neutron transport, Fission product

File Contents

ORNL	Eval-Apr89 R. Q. Wright
HEDL	Eval-Feb80 R. E. Schenter and F. Schmittroth
HEDL	Eval-Feb80 F. M. Mann, D. L. Johnson, G. Neely
RCN	Eval-Feb80 H. Gruppelaar
BNL	Eval-Oct74 A. Prince

The ¹⁴⁷Sm evaluation, MAT 9806, was revised by R. Q. Wright in February 1989. The new evaluation is assigned MAT No. 6234. ¹⁴⁷Sm is a naturally occurring isotope, with an abundance of 15 %. Actually ¹⁴⁷Sm is radioactive with a half-life of about 1.06×10^{11} years and decays by alpha decay to ¹⁴³Nd. ¹⁴⁷Sm is also produced by the radioactive decay of ¹⁴⁷Pm, hence it is also a fission product.

Summary of Changes

The resolved resonance parameters were taken from Ref. 1 ($E_{hi} = 1000.0$ eV). The contribution from the bound level to the 0.0253 eV capture cross section is 35.5 barns. Other resonances contribute 21.5 barns. Thus, the thermal capture cross section is 57.0 barns. Values of Γ_γ not given in Ref. 1 are set to 0.069 eV. The value for the scattering radius is 0.83. The upper limit of the resolved resonance range is increased from 401.88 to 1000.0 eV. The highest energy resonance included is 1050.0 eV.

Unresolved resonance parameters were added to the file. The unresolved range extends from 1 keV to 30 keV. The unresolved parameters are based on $D0 = 5.7$ eV and $S0 = 4.8$, see Ref. 1.

The capture cross section for the MAT 6234 evaluation is lower than ENDF/B-V (MAT 9806) and also lower than the data of Mizumoto (1981), but higher than the data of Macklin (1986) by about 1 to 5 percent. See Ref. 2, p. 514 for a plot of the capture data of Mizumoto and Macklin. The MAT 6234 capture cross section is compared with the data of Macklin in Table 1.

Table 1. ^{147}Sm Capture Cross Section (barns)

E (keV)	Macklin	MAT 6234	pcd
3- 4	4.35	4.40	1.1
4- 6	3.12	3.28	5.1
6- 8	2.37	2.48	4.6
8-10	1.94	2.02	4.1
10-15	1.52	1.58	3.9
15-20	1.19	1.23	3.4
20-30	0.962	0.968	0.62
30-40	0.777	0.780	0.39
40-60	0.645	0.648	0.47
60-80	0.546	0.548	0.37
80-100	0.484	0.490	1.24
100-150	0.425	0.426	0.24
150-200	0.3545	0.3556	0.31
200-300	0.3059	0.3056	-0.10
300-400	0.2623	0.2636	0.50
400-500	0.2459	0.2473	0.57
500-600	0.2454	0.2445	-0.37
600-700	0.2403	0.2401	-0.08

pcd = percent difference (MAT 6234 - Macklin)/Macklin

In File 3 the elastic and capture cross sections are set to zero in the resolved and unresolved range (10^{-5} eV to 30 keV). The 30-700 keV capture is based on the data of Macklin (1986). From 700 keV to 2 MeV, the capture cross section is reduced to match the data of Macklin at 700 keV. The MAT 6234 capture is about 30 percent lower than ENDF/B-V between 50 keV and 1 MeV. Above 2 MeV the MAT 6234 capture is unchanged from the ENDF/B-V evaluation.

The total cross section above 70 keV is unchanged from the ENDF/B-V evaluation, and the elastic cross section above 30 keV was increased slightly to offset the reduction in the capture cross section up to 2 MeV.

The (n,α) cross section has been revised below 230 eV. The cross section is based on the alpha widths given in Ref. 1. The thermal cross section is 0.623 mb, which is in good agreement with the Ref. 1 value $(0.58 \pm 0.06 \text{ mb})$. The (n,α) cross section is unchanged above 230 eV.

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	57.0
(from $1/v$ component)	=	0.0
Total	=	57.0

computed resonance integral = 790.0

References:

1. S. M. Mughabghab, "Neutron cross sections" Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B: Z=61-100, Academic Press (1984).
2. V. McLane, C. L. Dunford, and P. F. Rose, "Neutron Cross Sections," Vol. 2, Academic Press, New York (1988).

Summary of ENDF/B-V Evaluation

MF=2 MT=151 Resonance parameters from BNL-325 Ref.(3).

MF=3 MT= 1 Total cross section calculated with a deformed potential from Ref. (4) for $E > E_{hi}$.

MF=3 MT= 2 Elastic cross section from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$.

MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).

Summary of ENDF/B-V (Continued)

MF=3 MT=4,	51,52,...,91 Continued. The level scheme data is from the nuclear data tables and S. Igarasi (Japan), Private Communication.
MF=3 MT= 16(n,2n),	17(n,3n), 22(n,nd), 28(n,np), 103(n,p), 104(n,d), 105(n,t), 106(n, ³ He), 107(n, ⁴ He) calculated using the THRESH code Ref. (7).
MF=3 MT=102	Neutron capture was evaluated using COMNUC-3 and NCAP in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref.(3) for $E < E_{hi}$. The energy region above the resonance region was updated by combining the available integral and differential data using the generalized least squares adjustment code FERRET (HEDL-TME 77-51). The low energy capture also also adjusted to give a resonance integral (to within 1σ) of Ref.(3).
MF=4 MT=2	Angular distributions assumed isotropic.
MF=4	Non-elastic angular distributions assumed isotropic.
MF=5 MT=16,	17,22,28,91 The energy distributions of secondary neutrons are given as a histogram using calculations of nuclear temperature from Ref. (11).

References

1. T. Tamura, Computer Program JUPITOR I for Coupled-Channel Calculations, ORNL-4152 (1967).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL-325, 3ed, Vol. 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).
7. S. Pearlstein, Jour. Nucl. Energy 27, 81 (1973).

8. H. Baba and S. Baba, JAERI 1183 (1969).
9. G. Lautenbach, RCN-191 (1973).
10. S. M. Zakharova et al., INDC (CCP)-27/1.
11. A. Gilbert and A. G. W. Cameron, Can. J. Phys. 43, 1446 (1965).

¹⁵¹₆₂Sm

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: March 1989
Material: 6246
Content: Neutron transport, Fission product

File Contents

ORNL	Eval-Mar89 R. Q. Wright
HEDL	Eval-Feb80 R. E. Schenter and F. Schmittroth
HEDL	Eval-Feb80 F. M. Mann, D. L. Johnson, G. Neely
RCN	Eval-Feb80 H. Gruppelaar
BNL	Eval-Oct74 A. Prince

The ¹⁵¹Sm evaluation, MAT 9810, was revised by R. Q. Wright in August 1988. The new evaluation is MAT=6246. ¹⁵¹Sm has a half-life of 90 yr., and it is a significant reactor poison.

Summary of Changes

The resolved resonance parameters were taken from Ref. 1 ($E_{hi} = 300.0$ eV). The contribution from the bound level to the 0.0253 eV capture cross section is 14976 b. Other resonances contribute 224 barns. Thus, the thermal capture cross section is 15200 barns. Values of Γ_γ not given in Ref. 1 are set to 0.092 eV. The value for the scattering radius is 0.83 fm. The upper limit of the resolved resonance range has been increased from 6.941 to 300.0 eV, and the highest energy resonance included is at 295.7 eV. The resolved resonance range has been significantly improved in the new evaluation with 121 resolved resonance parameter sets, including one bound level, as against ENDF/B-V with only 8 resonances.

In File 3 the total, elastic, and capture cross sections are set to zero in the resolved resonance range (10^{-5} to 300.0 eV).

The 2200 m/s capture cross section, barns

(from resonance parameters) = 15200
(from 1/v component) = 0.0
Total = 15200

computed resonance integral = 3435

Reference:

1. S. F. Mughabghab, "Neutron Cross Sections" Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B: Z = 61-100, Academic Press (1984).

Summary of ENDF/B-V Evaluation

MF=2 MT=151	Resonance parameters from BNL-325 Ref. (3).
MF=3 MT= 1	Total cross section calculated with a deformed potential from Ref. (4) for $E > E_{hi}$.
MF=3 MT= 2	The elastic cross section was obtained from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$.
MF=3 MT= 4,	51,52,...,91 Inelastic cross sections were calculated using COMNUC-3, Refs. (5, 6). The level scheme data was taken from the Nuclear Data Tables and S. Igarasi (Japan) Private Communication.
MF=3 MT= 16(n,2n), 17(n,3n), 22(n,nd), 28(n,np), 103(n,p), 104(n,d), 105(n,t), 106(n, ³ He), 107(n, ⁴ He)	calculated using the THRESH code Ref (7).
MF=3 MT=102	Neutron capture was evaluated using COMNUC-3 and NCAP in Refs. (1, 2) for $E > E_{hi}$. A 1/v component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$.

Summary of ENDF/B-V (Continued)

MF=3 MT=102 Continued. The energy region above the resonance region was updated by combining available integral and differential data using the generalized least squares adjustment code FERRET (HEDL-TME 77-51). The low energy capture was also adjusted to give a resonance integral (to within 1σ) of Ref. (3).

MF=4 MT=2 Angular distributions assumed isotropic.

MF=4 Non-elastic angular distributions assumed isotropic.

MF=5 MT=16, 17,22,28,91 Energy distributions of secondary neutrons are given as a histogram using calculations of nuclear temperature from Ref. (11).

References

1. T. Tamura, Computer Program JUPITOR I for Coupled-Channel Calculations, ORNL-4152 (1967).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL-325, 3ed, Vol 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (Jul 1970).
6. C. L. Dunford (Private Communication).
7. S. Pearlstein, Jour. Nucl. Energy 27, 81 (1973).
8. H. Baba and S. Baba, JAERI 1183 (1969).
9. G. Lautenbach, RCN-191 (1973).
10. S. M. Zakharova et al., INDC (CCP)-27/1.
11. A. Gilbert and A. G. W. Cameron, Can. J. Phys. 43, 1446 (1965).

SUMMARY DOCUMENTATION FOR ^{151}Eu ENDF/B-VI, MAT = 6325

P. G. Young

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

The ENDF/B-VI evaluation for ^{151}Eu combines results from a new theoretical analysis¹ above the resonance region with the previous ENDF/B-V resonance parameter evaluation. The theoretical analysis utilizes a deformed optical model to calculate neutron transmission coefficients and cross sections, a giant-dipole-resonance model to determine gamma-ray transmission coefficients, and Hauser-Feshbach statistical theory to calculate partial reaction cross sections.

II. NUCLEAR MODEL CALCULATIONS

The Hauser-Feshbach statistical-theory calculations were performed with the COMNUC and GNASH reaction theory codes, using neutron transmission coefficients from the coupled-channel optical model analysis.² The total neutron cross section for natural europium that resulted from the deformed optical model calculations is compared to experimental data in Fig. 1. The COMNUC calculations include width-fluctuation corrections, which are important at lower energies, and the GNASH calculations incorporate preequilibrium effects, which become significant at higher energies. COMNUC was used to calculate all cross sections below $E_n = 8$ MeV, whereas GNASH was used for calculations above 8 MeV and for all continuous spectral calculations. Both codes utilize the Gilbert and Cameron level density formulation and the Cook tabulation of level density parameters.² A maximum amount of experimental information concerning discrete energy levels was incorporated into the calculations, and the constant temperature part of the Gilbert and Cameron level density was matched to the discrete level data for each residual nucleus in the analysis.

III. EVALUATION RESULTS

Resolved resonance parameters from ENDF/B-V are used to represent the cross sections from 10^{-5} eV to 98.81 eV, with some adjustment made to the background cross sections to improve agreement with thermal and resonance integral data. From 98.81 eV to 1 keV, average resonance parameters from Version V are used to specify the cross sections. Above 1 keV, the smooth cross sections were calculated from the theoretical analysis described above, as were the secondary angular and energy distributions. Exceptions to this are the (n,p), (n,d), (n,t), (n, ^3He), and (n, α) cross sections, which were taken directly from ENDF/B-V. See the attached ENDF File 1 comment section for additional details and for references.

The $^{151}\text{Eu}(n,\gamma)$ cross section from ENDF/B-VI is compared to the ENDF/B-V evaluation and to a selection of experimental data in Fig. 2. Also shown in Fig. 2 is the (n, γ) cross section calculated using a second level density option in the Hauser-Feshbach statistical theory calculations.

¹ R. L. Macklin and P. G. Young, "Neutron Capture Cross Sections of ^{151}Eu and ^{153}Eu from 3 to 2200 keV," *Nucl. Sci. Eng.* **95**, 189 (1987).

² See the ENDF/B File 1 comment section (attached) for references.

Figure 1. Comparison of experimental values of the neutron total cross section of ^{nat}Eu with coupled-channel optical model calculations. The solid curve represents the optical model results, which closely approximates the ENDF/B-VI evaluation, and the points are experimental data.

Figure 2. Comparison of evaluated and experimental values of the $^{151}\text{Eu}(n,\gamma)$ cross section. The solid curve is the ENDF/B-VI evaluation, which utilizes a Gilbert-Cameron temperature/Fermi gas level density in the calculations. The dashed curve represents calculations using a back-shifted Fermi gas level density model.

¹⁵¹₆₃Eu

Reference: No Primary Reference
Evaluators: P. G. Young and E. D. Arthur
Evaluated: April 1986
Material: 6325
Content: Neutron transport, Gamma production

Resolved Resonance Region (From ENDF/B-V, S. Mughabghab)

File 2 Resonance parameters

The resolved resonance parameters recommended in BNL-325, Vol 1(1) third edition are adopted. Where spin values are not determined, assignments are made randomly in order to satisfy the (2J+1) level spacing law and the J-independence of strength function. The resolved positive energy resonances contribute 1430.5b to the thermal capture cross section. Parameters of a bound level are derived to fit a measured capture cross-section of 9200+100 b (Ref 1).

The thermal cross sections are

capture	= 920.0 b
scattering	= 6.3 b
total	= 920.6 b

Unresolved resonance parameters (from ENDF/B-V with upper energy lowered from 10 to 1 kev).

The unresolved resonance region, 98.81 eV to 1 keV is described by average resonance parameters obtained from Ref.1. The average radiative width for s-wave resonances was increased from 91.17 mv to 98 mv in order to fit measurements in the energy region of 100 eV to 10 keV.

References

- (1) S.F.Mughabghab and D.I.Garber, Brookhaven National Laboratory report BNL-325, Vol 1, 3rd. ed. 1973.

Energy range above the resonance region.

The evaluation above 10 keV is based on a detailed theoretical analysis utilizing the available experimental data. Coupled channel optical model calculations with the ECIS code (Ra70) were used to provide the total, elastic, and inelastic cross sections to the first 3 members of the ground state rotational band, as well as neutron elastic and inelastic angular distributions to the rotational levels. The ECIS code was also used to calculate neutron transmission coefficients. Hauser-Feshbach statistical theory calculations were carried out with the GNASH (Ar88, Yo77) and COMNUC (Du70) code systems, including preequilibrium contributions. Systematics were used to obtain parameters for the exciton preequilibrium model, with small adjustments made to improve agreement with available exp. data. The Gilbert-Cameron level density model was used to supplement available experimental information on low-lying levels (Gi65). The Brink-Axel model (Br55, Ax62) was used to calculate gamma-ray transmission coefficients, using gamma-ray strength function results compiled by Mughabghab (Mu84).

A description of the calculations is given in Mc87.

*****MF=3 Smooth Cross Sections*****

MT=1 Neutron Total Cross Section. 0.01 to 20 MeV, based on coupled-channel optical calculations, which were optimized to the available experimental data (Mc88).
 MT=2 1.E-11 to 20 MeV, based on subtraction of MT=4,16,17,102, 103,104,105,106,107 from MT=1. This corresponds closely to using the results of the coupled-channel optical and Hauser-Feshbach model calculated elastic x/s.
 MT=4 Sum of MT=51-91
 MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=51-56,58-64,66-67 Thres.to 8.0 MeV, Compound nucleus reaction theory calculations using the COMNUC code (Du70) and including width fluctuation corrections. Transmission coefficients from cc optical model calculations used.
 MT=57,65 Thres. to 20 MeV, Coupled-channel optical model calculations plus compound-nucleus contributions.
 MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=103 (n,p) cross section retained from ENDF/B-V.
 MT=104 (n,d) cross section retained from ENDF/B-V.
 MT=105 (n,t) cross section retained from ENDF/B-V.
 MT=106 (n,He3) cross section retained from ENDF/B-V.
 MT=107 (n,He4) cross section retained from ENDF/B-V.

*****MF=4 Neutron Angular Distributions*****

MT=2 Elastic scattering angular distribution based on ECIS coupled-channel calculations, with a compound elastic component from COMNUC included below 8 MeV.

MT=16 (n,2n) distributions assumed isotropic in the laboratory system.

MT=17 (n,3n) distributions assumed isotropic in the laboratory system.

MT=51-56,58-64,66-67 Thres.to 8.0 MeV, Compound nucleus reaction theory calculations using the COMNUC code (Du70) and including width fluctuation corrections. Transmission coefficients from cc optical model calculations used.

MT=57,65 Thres. to 20 MeV, Coupled-channel optical model calculations plus compound-nucleus contributions.

MT=91 (n,n'continuum) distributions assumed isotropic in the laboratory system.

*****MF=5 Neutron Energy Distributions*****

MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc. Tabulated laboratory distributions given.

MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc. Tabulated laboratory distributions given.

MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc. Tabulated laboratory distributions given.

*****MF=12 Photon Multiplicities*****

MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc. Note that photons from (n,gn') reactions are included in MF=12,MT=102 but not in MF=3,MT=102, which causes the multiplicities at higher energies to become somewhat large.

*****MF=13 Photon Production Cross Sections*****

MT=4 GNASH Hauser-Feshbach statistical/preequilibrium calc.

MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.

MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****MF=14 Photon Angular Distributions*****

MT=4 Isotropy assumed.

MT=16 Isotropy assumed.

MT=17 Isotropy assumed.

MT=102 Isotropy assumed.

*****MF=15 Photon Energy Distributions*****

MT=4 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****References*****

Ar88 E.D.Arthur, LA-UR-88-382 (1988).
Ax62 P.Axel, Phys.Rev.126, 671 (1962).
Br55 D.M.Brink, D.Ph. Thesis, Oxford (1955).
Du70 C.L.Dunford, AI-AEC-12931 (1970).
Mc87 R.Macklin and P.G.Young, Nuc.Sci.Eng.95, 189(1987).
Mc88 V.McLane et al., Neutron Cross Sections V2 (Acad.Pr.1988).
Mu84 S.F.Mughabghab, Neutron Cross Sections V1 (Acad.Pr.1986).
Ra70 J.Raynal, IAEA SMR-9/8 (1970).
Yo77 P.G.Young & E.D.Arthur, LA-6947 (1977).

¹⁵²₆₃Eu

Reference: No Primary Reference
Evaluators: R. Q. Wright, H. Takahashi
Evaluated: December 1988
Material: 6328
Content: Neutron transport, Fission product

File Comments

ORNL	Eval-Dec88 R. Q. Wright
BNL	Eval-Dec73 H. Takahashi

The ENDF/B-V ¹⁵²Eu evaluation, MAT 1292, has been revised below 90.0 keV. The revised evaluation has been assigned MAT No 6328 in order to differentiate it from the original evaluation.

Summary of Changes

The ¹⁵²Eu evaluation, MAT 1292, was originally done for ENDF/B-IV in December 1973. No experimental data were available (other than the thermal capture cross section, 2300 ± 1000 barns). Resolved resonance parameters (fictitious) were generated using the procedure described in the MAT 1292 file 1 comments. The resolved resonance range extended from 0.93470 to 61.5 eV.

For this revision, resolved resonance parameters (based on experimental data) are taken from Ref. (1) and are used to define the total, elastic, and capture cross sections for energies between 0.00001 and 10.8 eV. The original resonance parameters in the energy range 10.8 to 61.5 eV are modified as follows, (relative to the ENDF/B-V parameters):

E_0	Same as MAT 1292 (ENDF/B-V)
Γ_n	$\Gamma_n = 0.3 \times \Gamma_n / (2.0 \times g)$
Γ_γ	$\Gamma_\gamma = 0.160$ eV
Γ_t	$\Gamma_t = \Gamma_n + \Gamma_\gamma$

The average reduced neutron width is 4.19048×10^{-4} , the average gamma width is 1.59821×10^{-1} , and the strength function is 2.83842×10^{-4} . The MLBW formalism is used. The largest contribution to the thermal capture cross section (about 98%) is from the bound level at -0.1 eV. The thermal capture cross section is higher than the ENDF/B-V value by a factor of about 5.5. The capture resonance integral is lower than the ENDF/B-V value by about 37%.

In the unresolved resonance range: (upper limit is 3 keV) the unresolved resonance parameters are based on the data given in Ref. 1:

Average Γ_γ	0.160 eV
d_0	0.25 eV
s_0 (lower than Ref. 1)	2.8200×10^{-4}

The File 3 changes are as follows: elastic and capture backgrounds (MF=3) are zero below 3 keV, and the capture cross section at 30 keV is 5196 mb. Capture is lower below about 5 keV and is unchanged above 15 keV. The elastic cross section is lower than ENDF/B-V below 90 keV. The total cross section was revised to agree with the sum of the partial cross sections.

The 2200 m/s capture cross section, barns

(from resonance parameters) = 12819

computed resonance integral

0.5 - 10.8 eV	=	1660
10.8 - 61.5 eV	=	387
Above 61.5 eV	=	279
Total	=	2326

Reference:

1. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Part B: Z=61-100, Academic Press, New York (1984).

Summary of ENDF/B-V Evaluation

This material contains the evaluated results for ^{152}Eu . No experimental data, except for a few reactions, data are available for the isotopes of Eu, so that the evaluations were mostly carried out using nuclear model calculations.

The resolved resonance parameters were made by taking into account their statistical properties for level spacing and reduced neutron width fluctuation. The method used in this calculation was similar to the procedure used by Cook.¹ However, instead of using a Monte Carlo calculation, the level spacing and the reduced neutron width fluctuations were determined by using the statistical properties of ^{151}Eu . The average values of these quantities were determined in a similar way to the procedure used by Barr et al.² That is, the ratios of the average values for odd-even nuclei to those for odd-odd nuclei were estimated from their neighboring nuclei. These ratios were multiplied by the values of ^{151}Eu to obtain the ones for ^{152}Eu . The gamma ray widths were taken as constant values for all resonances. The 108 resonances were assigned between 0.01 eV and 61.704 eV. The thermal neutron capture cross section has been measured by Hayden et al.³ and Walker.⁴ The preliminary draft of BNL-325 recommends a value 2300 ± 1000 barns for the ^{152}Eu ground state. The parameters of the lowest resonances were adjusted so that the calculated thermal neutron capture cross sections agreed with the values recommended in BNL-325.⁵

Unresolved resonance parameters were given in the energy region from 61.5 eV to 10 keV. As mentioned above, Barr and Devaney² evaluated the unresolved resonance parameters by studying the change of these parameters from odd-odd nuclei to odd-even nuclei in ^{175}Lu , ^{176}Lu , ^{180}Ta and, ^{181}Ta . The unresolved resonance parameters were estimated from the BNL-325 values.

Between 10000 eV and 2.5 MeV, the total cross sections were calculated using the optical model ABACUS-2 code.⁶ The optical parameters used in the calculation were taken from the study of ^{151}Eu and ^{153}Eu .⁷ Above 2.5 MeV, the total cross sections were assumed to be the same as the experimental values of natural europium measured by Foster.⁸

The elastic scattering cross sections in the energy range above the unresolved resonance region were obtained by subtracting the non elastic cross section from the evaluated total cross section.

The nonelastic scattering cross section was calculated by summing up all cross sections except the elastic scattering cross section.

The inelastic scattering cross sections were given as total (MT = 4), discrete level excitation cross sections (MT = 51...) for the first 5 levels, and a continuum level excitation cross section (MT = 91). The level scheme is taken from Refs. (9, 10, 11, 12, and 13). Since no experimental data are available for the individual level excitation cross sections, they were calculated using the COMNUC-3 code^{11,15} for energies up to 3 MeV. Above 3 MeV in neutron energy, the inelastic scattering is mostly the

excitation of the continuum of levels, so that the inelastic scattering cross section for discrete level excitation above this energy was neglected, and the inelastic scattering cross section for continuum level excitation was calculated using the cascade calculation from the GROGI-3 code.¹⁶ The level density parameters for the continuum of levels were taken from Cook's data¹⁸ for deformed nuclei using the Gilbert-Cameron formula.¹⁹

For the (n,p) and (n,np) cross section (MT = 103, 28), no experimental values were available, so they were calculated using nuclear model codes. For the (n,p) reaction, the semi-empirical statistical model code THRESH²¹ was used, but the evaluation of ¹⁵¹Eu and ¹⁵³Eu⁷ indicated that the cross sections around 14 MeV calculated by this code were small compared to the experimental values. Thus, the calculated cross sections were normalized by the factors obtained for ¹⁵¹Eu. The (n,np) cross sections were calculated using the GROGI-3 code.

The (n, α) and (n,nd) cross sections (MT = 107, 22) were obtained in a similar manner to the (n,p) and (n,np) reactions.

The (n,2n), and (n,3n) cross sections (MT = 16, 17) were calculated using the GROGI-3 code. The optical model parameters mentioned previously were used.

The (n,d), (n,t), and (n,³He) reaction cross sections (MT=104, 105, and 107) were adapted from calculations using THRESH.

The radiative capture cross sections (MT = 102) at low energies were calculated from resonance parameters as previously discussed, and are presented as smooth cross sections. The cross sections between 100 eV and 10 keV were obtained from the unresolved resonance parameters. For energies higher than 10 keV, the cross sections were evaluated from the COMNUC-3 calculations. These calculations were similar to the ones for ¹⁵¹Eu and ¹⁵³Eu.⁷ That is, we assumed Moldauer's Q value to be zero, and the correlation correction factor due to the degrees of freedom associated with the open channel was taken into account in the calculation. From 3 MeV to 20 MeV, the capture cross section was obtained from GROGI-3 for compound processes, by Cvelbar's formula²² based on Lane and Lynn²³, and Brown's²⁴ formula for direct and semi-direct reactions.

The elastic scattering (MT = 2) and the angular distribution of secondary neutrons in File 4 were calculated using ABACUS-2 (NABAK, a PDP-10 computer version).⁶ Legendre coefficients were calculated using CHAD (NUCHAD, on the PDP-10).²⁵ Since the elastic scattering due to the nuclear compound process is small in the energy range above 3 MeV, the angular distributions of elastic scattering neutrons

were calculated by taking only the shape elastic scattering into account above 3 MeV.

Inelastically scattered neutrons, for the $(n,2n)$, $(n,3n)$, (n,np) , and $(n,n\alpha)$ reactions (MT = 51, ..., 91, MT=16, 17, 22, 23) were assumed to be isotropic in the center of mass system.

The energy distribution of secondary neutrons from the $(n,2n)$, $(n,3n)$, and continuum (n,n') reactions, (MT = 16, 19, and 91) were assumed Maxwellian with an effective temperature obtained from the Weiskopf formula.²⁶

The file was translated into the ENDF-5 format by F. M. Mann and R. E. Schenter (HEDL) in January 1979.

References

1. J. L. Cook, AAEC/TM-549 (1969).
2. D. W. Barr and J. H. Devaney, LA-3643 (1967).
3. R. J. Hayden, et. al., Phys. Rev 75, 1500 (1949).
4. W. H. Walker, AECL-3037, Part I (1969).
5. S. F. Mughabghab, and D. I. Garber, BNL-325, Third Edition, Vol 1. (1973).
6. E. H. Auerbach, BNL-6562 (1962).
7. H. Takahashi, "Evaluation of the Neutron and Gamma-Ray Production Cross Sections of ^{151}Eu and ^{153}Eu ," BNL-19455 (ENDF-213) (1974)
8. D. G. Foster Jr., and D. W. Glasgow, PNWL unpublished data (1966).
9. T. Lewise and R. Gratzner, Nucl. Phys. A162, 145 (1971).
10. A. Faessler, Nucl. Phys. 59, 1977 (1964).
11. L. V. Groshev et al., Nucl. Data Table A5, 1 (1968).
12. D. J. Horen et al., "Nuclear Level Scheme A = 45 through A = 257 From Nuclear Data Sheets," Academic Press Inc., New York (1973)
13. C. Lederer, J. Hollander and I. Perlman, "Table of Isotopes," Sixth Edition (1967).
14. C. L. Dunford, Private Communication (COMNUC-3 code) (1971).

15. C. L. Dunford, AI-AEC-12931 (1970).
16. H. Takahashi, "GROGI-3", (Modified from GROG-2. See Ref. 17)
17. J. Gilat, BNL-50246 (T-580) (1969).
18. J. Cook, H. Ferguson and A. Musgrove, AAEC/TM-392 (1967).
19. A. Gilbert and A. Cameron, Can. J. Phys. 43, 1446 (1965).
20. T. Tamura, Rev. Mod. Phys. 37, 679 (1965).
21. S. Pearlstein, J. Nucl. Ener. 27, 81 (1973).
22. F. Cvelbar et. al., NIJS Report T-529 (1968).
23. A. M. Lane and J. E. Lynn, Geneva Conference on Peaceful Uses of Atomic Energy, 15, 4 (1958).
24. G. E. Brown, Nucl. Phys. 57, 339 (1964).
25. R. F. Berland, NAA-SR-11231 (1965).
26. A. Weinberg and E. Wigner, "The Physical Theory of Neutron Chain Reactors," University of Chicago Press (1959)

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

$^{152}_{83}\text{Eu } \sigma_{n,\text{tot}}$

77	IJ1	Vertebnyj+	YF 26 (6), 1137	19	23.40 mv to 0.192 eV	
78	IJ1	Razbudej+	YFI-22, 19	18	6.500 mv to 0.168 eV	

SUMMARY DOCUMENTATION FOR ^{153}Eu ENDF/B-VI, MAT = 6331

P. G. Young

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

The ENDF/B-VI evaluation for ^{153}Eu combines results from a new theoretical analysis¹ above the resonance region with the previous ENDF/B-V resonance parameter evaluation. The theoretical analysis utilizes a deformed optical model to calculate neutron transmission coefficients and cross sections, a giant-dipole-resonance model to determine gamma-ray transmission coefficients, and Hauser-Feshbach statistical theory to calculate partial reaction cross sections.

II. NUCLEAR MODEL CALCULATIONS

The Hauser-Feshbach statistical-theory calculations were performed with the COMNUC and GNASH reaction theory codes, using neutron transmission coefficients from the coupled-channel optical model analysis.² The total neutron cross section for natural europium that resulted from the deformed optical model calculations is illustrated in the previous section on ^{151}Eu . The COMNUC calculations include width-fluctuation corrections, which are important at lower energies, and the GNASH calculations incorporate preequilibrium effects, which become significant at higher energies. COMNUC was used to calculate all cross sections below $E_n = 8$ MeV, whereas GNASH was used for calculations above 8 MeV and for all continuous spectral calculations. Both codes utilize the Gilbert and Cameron level density formulation and the Cook tabulation of level density parameters.² A maximum amount of experimental information concerning discrete energy levels was incorporated into the calculations, and the constant temperature part of the Gilbert and Cameron level density was matched to the discrete level data for each residual nucleus in the analysis.

III. EVALUATION RESULTS

Resolved resonance parameters from ENDF/B-V are used to represent the cross sections from 10^{-5} eV to 97.22 eV, with some adjustment made to the background cross sections to improve agreement with thermal and resonance integral data. From 97.22 eV to 1 keV, average resonance parameters from Version V are used to specify the cross sections. Above 1 keV, the smooth cross sections were calculated from the theoretical analysis described above, as were the secondary angular and energy distributions. Exceptions to this are the (n,p), (n,d), (n,t), (n, ^3He), and (n, α) cross sections, which were taken directly from ENDF/B-V. See the attached ENDF File 1 comment section for additional details and for references.

The $^{153}\text{Eu}(n,\gamma)$ cross section from ENDF/B-VI is compared to the Version V evaluation and to a selection of experimental data in Fig. 1. Also shown in Fig. 1 is the (n, γ) cross section calculated with a second level density option in the Hauser-Feshbach statistical theory calculations.

¹ R. L. Macklin and P. G. Young, "Neutron Capture Cross Sections of ^{151}Eu and ^{153}Eu from 3 to 2200 keV," *Nucl. Sci. Eng.* **95**, 189 (1987).

² See the ENDF/B File 1 comment section (attached) for references.

Figure 1. Comparison of evaluated and experimental values of the $^{153}\text{Eu}(n,\gamma)$ cross section. The solid curve is the ENDF/B-VI evaluation, which utilizes a Gilbert-Cameron temperature/Fermi gas level density in the calculations. The dashed curve represents calculations using a back-shifted Fermi gas level density model.

¹⁵³₆₃Eu

Reference: No Primary Reference
Evaluators: P. G. Young and E. D. Arthur
Evaluated: April 1986
Material: 6331
Content: Neutron transport, Gamma production

Resolved resonance region (from ENDF/B-V, S. Mughabghab)

File 2 resonance parameters.

Resonance parameters recommended in BNL-325(1973) (Ref.1) were adopted in this evaluation. Spin assignment of one resonance at 2.457 eV is determined. For the other resonances, spin assignments were made randomly in order to satisfy spin independence of strength function and the 2J+1 law of level density. Recent data on the measurement of the thermal cross section of Eu-153 brought out the problems with the accurate determination of this cross section. (Ref2-6). These problems are related to the lack of very accurate knowledge of Eu-151 content in Eu-153 samples and previous inaccurate value of half life of Eu-154. (16years). The half life of Eu-154 is presently known as 8.2+0.3 years. After correcting the data for Eu-151 impurity and half life of Eu-154, a weighted average value 300b is derived for the thermal cross section of Eu-153. The positive energy resonances contribute 58b to the thermal cross-section. The difference is accounted for by a negative energy resonance with a spin of J=2 as derived from thermal spectra measurement (Ref.1).

The thermal cross sections are

capture	=	312.0 b
scattering	=	9.0 b
total	=	321.0 b

Unresolved resonance parameters (from ENDF/B-V with upper energy lowered from 10 to 1 kev).

The unresolved energy region from 97.22 eV to 1 keV is represented by average resonance parameters as recommended in Ref.1

S(0)	=	2.50	E-04
Gamma width	=	95.8	mv
Average spacing	=	1.37	ev

References

- 1) S.F. Mughabghab and D.I. Garber, BNL-325 Third edition Vol-1, 1973.
- 2) M.C. Moxon, D.A.J. Endacott, and J.E. Jolly, Annals Nucl. Ener. 3, 399 (1976).
- 3) J.I. Widder, Nucl. Sc. Eng., 60, 53 (1976).
- 4) J.I. Kim, E.M. Gryntakis, H.J. Born, Radiochimica Acta, 22, 20, (1975).
- 5) V.P. Vertebny et al, Proc 1st Neutron Physics Conf, Kiev, Part 2, 239, (1973).
- 6) V.F. Razbudej, A.F. Fedorova, A.V. Muravitskij, INDC(CCP)-100/U, 23 (1977).
- 7) W. Stoffl, D. Rabenstein, K. Schreckenbach, and T. von Egidy, Z. Physik A282, 97 (1977).

 Energy range above the resonance region.

The evaluation above 10 keV is based on a detailed theoretical analysis utilizing the available experimental data. Coupled channel optical model calculations with the ECIS code (Ra70) were used to provide the total, elastic, and inelastic cross sections to the first 3 members of the ground state rotational band, as well as neutron elastic and inelastic angular distributions to the rotational levels. The ECIS code was also used to calculate neutron transmission coefficients. Hauser-Feshbach statistical theory calculations were carried out with the GNASH (Ar88, Yo77) and COMNUC (Du70) code systems, including preequilibrium contributions. Systematics were used to obtain parameters for the exciton preequilibrium model, with small adjustments made to improve agreement with available exp. data. The Gilbert-Cameron level density model was used to supplement available experimental information on low-lying levels (Gi65). The Brink-Axel model (Br55, Ax62) was used to calculate gamma-ray transmission coefficients, using gamma-ray strength function results compiled by Mughabghab (Mu84).

A description of the calculations is given in Mc87.

*****MF=3 Smooth Cross Sections*****

- MT=1 Neutron Total Cross Section. 0.01 to 20 MeV, based on coupled-channel optical calculations, which were optimized to the available experimental data (Mc88).
- MT=2 1.E-11 to 20 MeV, based on subtraction of MT=4,16,17,102, 103,104,105,106,107 from MT=1. This corresponds closely

to using the results of the coupled-channel optical
and Hauser-Feshbach model calculated elastic x/s.

MT=4 Sum of MT=51-91
MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=51,56 Thres. to 20 MeV, Coupled-channel optical model
calculations plus compound-nucleus contributions.
MT=52-55,57-60 Threshold to 8.0 MeV, Compound nucleus reaction
theory calculations using the COMNUC code (Du70) and
including width fluctuation corrections. Transmission
coefficients from cc optical model calculations used.
MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=103 (n,p) cross section retained from ENDF/B-V.
MT=104 (n,d) cross section retained from ENDF/B-V.
MT=105 (n,t) cross section retained from ENDF/B-V.
MT=106 (n,He3) cross section retained from ENDF/B-V.
MT=107 (n,He4) cross section retained from ENDF/B-V.

*****MF=4 Neutron Angular Distributions*****

MT=2 Elastic scattering angular distribution based on ECIS
coupled-channel calculations, with a compound elastic
component from COMNUC included below 8 MeV.
MT=16 (n,2n) distributions assumed isotropic in the laboratory
system.
MT=17 (n,3n) distributions assumed isotropic in the laboratory
system.
MT=51,56 Thres. to 20 MeV, Coupled-channel optical model
calculations plus compound-nucleus contributions.
MT=52-55,57-60 Threshold to 8.0 MeV, Compound nucleus reaction
theory calculations using the COMNUC code (Du70) and
including width fluctuation corrections. Transmission
coefficients from cc optical model calculations used.
MT=91 (n,n'continuum) distributions assumed isotropic in the
laboratory system.

*****MF=5 Neutron Energy Distributions*****

MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
Tabulated laboratory distributions given.
MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
Tabulated laboratory distributions given.
MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc.
Tabulated laboratory distributions given.

*****MF=12 Photon Multiplicities*****

MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.
Note that photons from (n,gn') reactions are included
in MF=12,MT=102 but not in MF=3,MT=102, which causes
the multiplicities at higher energies to become
somewhat large.

*****MF=13 Photon Production Cross Sections*****

MT=4 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****MF=14 Photon Angular Distributions*****

MT=4 Isotropy assumed.
MT=16 Isotropy assumed.
MT=17 Isotropy assumed.
MT=102 Isotropy assumed.

*****MF=15 Photon Energy Distributions*****

MT=4 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****References*****

Ar88 E.D.Arthur, LA-UR-88-382 (1988).
Ax62 P.Axel, Phys.Rev.126, 671 (1962).
Br55 D.M.Brink, D.Ph. Thesis, Oxford (1955).
Du70 C.L.Dunford, AI-AEC-12931 (1970).
Mc87 R.Macklin and P.G.Young, Nuc.Sci.Eng.95, 189(1987).
Mc88 V.McLane et al., Neutron Cross Sections V2 (Acad.Pr.1988).
Mu84 S.F.Mughabghab, Neutron Cross Sections V1 (Acad.Pr.1986).
Ra70 J.Raynal, IAEA SMR-9/8 (1970).
Yo77 P.G.Young & E.D.Arthur, LA-6947 (1977).

¹⁵⁴₆₃Eu

Reference: No Primary Reference
Evaluators: R. Q. Wright, H. Takahashi
Evaluated: May 1989
Material: 6334
Content: Neutron transport, Fission product

File Comments

ORNL	Eval-May89 R. Q. Wright
BNL	Eval-Dec73 H. Takahashi

The ENDF/B-V ¹⁵⁴Eu evaluation, MAT 1293, has been revised below 10 keV. The revised evaluation has been assigned MAT No. 6334 in order to differentiate it from the original evaluation.

Summary of Changes

The ¹⁵⁴Eu evaluation, MAT 1293, was originally done for ENDF/B-IV in December 1973. No experimental data were available (other than the thermal capture cross section, 1500 ± 400 barns). Resolved resonance parameters (fictitious) were generated using the procedure described in the MAT 1293 File 1 comments. The resolved resonance range extended from 0.88962 to 60.0 eV.

For this revision, resolved resonance parameters (based on experimental data) are taken from Ref. (1) and are used to define the total, elastic, and capture cross sections for energies between 0.00001 and 27.8 eV. The original resonance parameters in the energy range 27.8 to 63.0 eV are modified as follows, (relative to the ENDF/B-V parameters):

E_0	Same as MAT 1293 (ENDF/B-V)
Γ_n	$\Gamma_n = \Gamma_n / (2.0 \times g)$ (to keep same value of $2g \times \Gamma_n$)
Γ_γ	$\Gamma_\gamma = \Gamma_\gamma \times 1.3125$ (to get average width = 0.126)
Γ_t	$\Gamma_t = \Gamma_n + \Gamma_\gamma$

The average reduced neutron width is 4.44540×10^{-1} , the average gamma width is 1.25926×10^{-1} , and the strength function is 2.19309×10^{-1} . The MLBW (LRF = 2) formalism is used, and $E_{hi} = 63.0$ eV.

The upper limit of the unresolved resonance range is 10 keV. The unresolved resonance parameters are based on the data given in Ref. 1:

Average Γ_γ	0.1260 eV
D0 (Ref. 1 has 0.92 eV)	0.9752 eV
S0 (Not given in Ref. 1)	2.5709×10^{-1}

The File 3 changes are as follows: elastic and capture backgrounds (MF=3) are zero below 10 keV, and the elastic cross section at 10 keV was reduced to 13.8 barns. The capture cross section at 30 keV is 2920 mb (unchanged). The total cross section was revised by small amounts at 21 points between 8.7 and 11.8 MeV to agree with the sum of the partial cross sections.

The thermal capture cross section is lower than the ENDF/B-V value by about 10%. The capture resonance integral is lower than the ENDF/B-V value by about 47%.

The 2200 m/s capture cross section, barns

(from resonance parameters) = 1352

computed resonance integral

0.5 - 27.8 eV	=	984
27.8 - 63.0 eV	=	145
Above 63.0 eV	=	216
Total	=	1345

Reference:

1. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Part B: Z=61-100, Academic Press, New York (1984).

Summary of ENDF/B-V Evaluation

This material contains the evaluated results for the neutron cross sections of ^{151}Eu . No experimental data, except for a few reactions, are available for the isotopes of Eu,

so that the evaluations were mostly carried out using nuclear model calculations.

The resolved resonance parameters were made by taking into account their statistical properties for level spacing and reduced neutron width fluctuations. The method used in this calculation was similar to the procedure used by Cook.¹ However, instead of using a Monte Carlo calculation, the level spacing and the reduced neutron width fluctuation are determined by using the statistical properties of ^{153}Eu . The average values of these quantities are determined in a similar way to the procedure used by Barr et al.² that is, the ratios of the average values for odd-even nuclei to those for odd-odd nuclei were estimated from their neighboring nuclei. These ratios were multiplied to the values of ^{153}Eu to obtain the ones for ^{154}Eu . The gamma ray widths were taken as constant values for all resonances.

The 96 resonances were assigned between 0.01 eV and 59.732 eV. The thermal neutron capture cross section has been measured by Hayden et al.³ and Walker.⁴ The preliminary draft of BNL-325 recommends those values as 1500 ± 400 barns for ^{154}Eu ($T_{1/2} = 8$ years). The parameters of the lowest resonances were adjusted so that the calculated thermal neutron capture cross sections agreed with the values recommended in BNL-325.⁵

The unresolved resonance parameters were given in the energy region from 60.0 eV to 10 keV. As mentioned above, Barr and Devaney² evaluated the unresolved resonance parameters by studying the change of these parameters from odd-odd nuclei to odd-even nuclei in ^{175}Lu , ^{176}Lu , ^{180}Ta , and ^{181}Ta . The unresolved resonance parameters were estimated by using the BNL-325 values.

Between 10000 eV and 2.5 MeV, the total cross sections were calculated using the ABACUS-2 code⁶ The optical parameters used in the calculation were taken from the study of ^{151}Eu and ^{153}Eu .⁷ Above 2.5 MeV, the total cross sections were assumed to be the same as the experimental values of natural europium measured by Foster.⁸

The elastic scattering cross sections in the energy above the unresolved resonance energy range were obtained by subtracting the non-elastic cross section from the evaluated total cross section.

The nonelastic scattering cross section was calculated by summing up all cross sections except the elastic scattering cross section.

The inelastic scattering cross sections were given as total ($MT = 4$), discrete level excitation cross sections ($MT = 51\dots$) for the first 5 levels, and a continuum level excitation cross section ($MT = 91$). The level scheme for these discrete level is taken

from Refs. (9, 10, 11, 12, and 13). Since no experimental data are available for the individual level excitation cross sections, they were calculated using the COMNUC-3 code^{14,15} for energies up to 3 MeV. Above 3 MeV, the inelastic scattering is mostly the excitation of the continuum of levels, so that the inelastic scattering cross section for discrete level excitation above this energy was neglected and the inelastic scattering cross section for continuum level excitation was calculated using the cascade calculation of GROGI-3.¹⁶ The level density parameters for the continuum of levels were taken from Cook's data¹⁸ for deformed nuclei using the Gilbert-Cameron formula.¹⁹

For the (n,p) and (n,np) cross section (MT = 103, 28) no experimental values were available, so that they were calculated using nuclear model codes. For the (n,p) reaction, the semi-empirical statistical model code THRESH²¹ was used, but the evaluation of ¹⁵¹Eu and ¹⁵³Eu⁷ indicated that the cross sections around 14 MeV calculated using this code were too small compared to the experimental values. Thus, the calculated cross sections were normalized by the factors obtained for ¹⁵¹Eu. The (n,np) cross sections were calculated using GROGI-3.

The (n, α) and (n,nd) cross sections (MT = 107, 22) were obtained in a similar manner to the (n,p) and (n,np) reactions.

The (n,2n) and (n,3n) cross section (MT = 16, 17) were calculated using the GROGI-3 code. The optical model parameters mentioned previously were used.

The (n,d), (n,t), and (n,³He) reaction cross sections (MT=104, 105, and 107) were calculated using THRESH.

The radiative capture cross sections at low energy (MT = 102) were calculated from the resonance parameters discussed above, and are presented as the smooth cross sections. The cross sections between 100 eV and 10 keV were obtained from the unresolved resonance parameters. For energies higher than 10 keV, the cross sections were evaluated using COMNUC-3. The calculation was done similarly to the ones for ¹⁵¹Eu and ¹⁵³Eu.⁷ That is, Moldauer's Q value was assumed to be zero, and the correlation correction factor due to the degrees of freedom associated with an open channel was taken into account in the calculation. From 3 MeV to 20 MeV, the capture cross section was obtained using GROGI-3 for compound processes, by Cvelbar's formula²² based on Lane and Lynn²³ and Brown's²⁴ formula for direct and semi-direct reactions.

The elastic scattering (MT = 2) and the angular distribution of secondary neutrons in File 4 were calculated using ABACUS-2 (NABAK, the PDP-10 version)⁶ and

the legendre coefficients were calculated using CHAD (NUCHAD, the PDP-10 version)²⁵ were given. Since the elastic scattering due to the nuclear compound process is small in the energy range above 3 MeV, the angular distribution of elastic scattering neutrons was calculated by taking only the shape elastic scattering into account above 3 MeV.

Inelastically scattered neutrons, for the (n,2n), (n,3n), (n,np), and (n,n α) reactions (MT=51, ..., 91, MT=16, 17, 22, and 23) were assumed to be isotropic in the center of mass system.

The energy distribution of secondary neutrons from the (n,2n), (n,3n), and (n,n') continuum reactions (MT=16, 17, and 91) were assumed as maxwellian with an effective temperature obtained using the Weiskopf formula.²⁶

The file was translated into the ENDF-5 format by F. M. Mann and R. E. Schenter (HEDL) in December 1978.

References

1. J. L. Cook, AAEC/TM-549 (1969).
2. D. W. Barr and J. H. Devaney, LA-3643 (1967).
3. R. J. Hayden, et. al., Phys. Rev. 75, 1500 (1949).
4. W. H. Walker, AECL-3037, Part I (1969).
5. S. F. Mughabghab, and D. Garber, BNL-325 Third Edition Vol 1. (1973).
6. E. H. Auerbach, BNL-6562 (1962).
7. H. Takahashi, "Evaluation of the Neutron and Gamma-Ray Production Cross Sections of ¹⁵¹Eu and ¹⁵³Eu" BNL-19455 (ENDF-213) (1974).
8. D. G. Foster Jr., and D. W. Glasgow, PNWL unpublished data (1966).
9. T. Lewis and R. Gratzner, Nucl. Phys. A162, 145 (1971).
10. A. Faessler, Nucl. Phys. 59, 1977 (1964).
11. L. V. Groshev et al., Nucl. Data Table A5, 1 (1968).
12. D. J. Horen et al., "Nuclear Level Scheme A=45 through A=257," Academic Press Inc., New York (1973).

13. C. Lederer, J. Hollander and I. Perlman, "Table of Isotopes," Sixth Edition (1967).
14. C. Dunford, Private Communication (COMNUC-3 code) (1971).
15. C. Dunford, AI-AEC-12931 (1970).
16. H. Takahashi, GROGI-3 (Modified from GROGI-2, see Ref. 17)
17. J. Gilat, BNL-50246 (T-580) (1969).
18. J. Cook, H. Ferguson and A. Musgrove, AAEC/TM-392 (1967).
19. A. Gilbert and A. Cameron, Can. J. Phys. 43, 1446 (1965).
20. T. Tamura, Rev. Mod. Phys. 37, 679 (1965).
21. S. Pearlstein, J. Nucl. Ener. 27, 81 (1973).
22. F. Cvelbar, et. al., NIJS Report T-529 (1968).
23. A. M. Lane and J. E. Lynn, Geneva Conference on Peaceful Uses of Atomic Energy, 15, 4 (1958).
24. G. E. Brown, Nucl. Phys. 57, 339 (1964).
25. R. F. Berland, NAA-SR-11231 (1965).
26. A. Weinberg and E. Wigner, "The Physical Theory of Neutron Chain Reactors," University of Chicago Press (1959).

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

$^{154}\text{Eu } \sigma_{n,\text{tot}}$						
76	IJI	Razbudej+	YFI-22, 19	1	1.530 kb	at thermal

¹⁵⁵₆₃Eu

Reference: No Primary Reference
Evaluators: R. Q. Wright, A. Prince, and R. E. Schenter
Evaluated: December 1988
Material: 6337
Content: Neutron transport, Fission product

File Comments

ORNL	Eval-Dec88 R. Q. Wright
INEL	Eval-Dec79 A. Prince, R. E. Schenter
BNL,HEDL	Eval-Oct74 A. Prince and R. E. Schenter

The ENDF/B-V ¹⁵⁵Eu evaluation, MAT 9832, has been revised below 35 eV. The revised evaluation has been assigned MAT No. 6327 in order to differentiate it from the original evaluation.

Summary of Changes

In the revised evaluation, resolved resonance parameters are used to define the total, elastic, and capture cross sections below 35 eV. Above 35 eV the evaluation is unchanged from ENDF/B-V. The resolved resonance parameters are taken from Ref. (1). The MLBW (LRF=2) representation was used with the smooth background set to zero in the resonance region.

The largest contribution to the thermal capture cross section and to the capture resonance integral is from the first resonance at 0.603 eV which has a peak capture cross section (at 300° K) of about 102,000 barns. The thermal capture cross section is slightly lower than ENDF/B-V, but the capture resonance integral is higher than the ENDF/B-V value by about a factor of 12.

The 2200 m/s capture cross section, barns

(from resonance parameters) == 3941

computed resonance integral

(from resonance parameters) == 22927

Above 35 eV == 272

Total == 23199

Reference:

1. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Part B: Z=61-100, Academic Press, New York (1984).

Summary of ENDF/B-V Evaluation

The ^{155}Eu file was translated into ENDF-5 format by F. M. Mann and R. E. Schenter (HEDL) in January 1980.

MF=2 MT=151	No resonance parameters given except AP.
MF=3 MT= 1	Total cross section calculated with a deformed potential from Ref. (4) for $E > E_{hi}$.
MF=3 MT= 2	Elastic cross section from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$, from $4\pi a^2$ for $E < E_{hi}$.
MF=3 MT= 4,	51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6). The level scheme data was taken from the Nuclear Data Tables and S. Igarasi (Japan), Private Communication.
MF=3 MT= 16(n,2n),	17(n,3n), 22(n,nd), 28(n,np), 103(n,p), 104(n,d), 105(n,t), 106(n, ^3He), 107(n, ^4He) calculated using the THRESH code Ref (7).
MF=3 MT=102	Neutron capture evaluated using COMNUC-3 and NCAP in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$.
MF=4 MT=2	Angular distributions calculated from the Moldauer potential.
MF=4	Non-elastic angular distributions assumed isotropic.
MF=5 MT=16,	17, 22, 28, 91 Energy distributions of secondary neutrons given as a histogram using calculations of nuclear temperature from Ref. (11).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	0
(from 1/v component)	=	4040
Total	=	4040

computed resonance integral = 1856

References

1. T. Tamura, Computer Program JUPITOR I for Coupled-Channel Calculations, ORNL-4152 (1967).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL-325, 3ed, Vol 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).
7. S. Pearlstein, Jour. Nucl. Energy 27, 81 (1973).
8. H. Baba and S. Baba, JAERI 1183 (1969).
9. G. Lautenbach, RCN-191 (1973).
10. S. M. Zakharova et al., INDC (CCP)-27/1.
11. A. Gilbert and A. G. W. Cameron, Can. J. Phys. 43, 1446 (1965).

Yr	Lab	Author	Reference	Points	Range	Standard
$^{155}_{63}\text{Eu } \sigma_{n,\gamma}$						
82	IJI	Vertebnyj+	YK 5/49, 16	1	3.950kb	at thermal

SUMMARY DOCUMENTATION FOR ^{165}Ho ENDF/B-VI, MAT = 6325

P. G. Young and E. D. Arthur

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

The ENDF/B-VI evaluation for ^{165}Ho is based on a new theoretical analysis¹ above the resonance region, which was combined with the previous ENDF/B-V resonance parameter evaluation. The theoretical analysis utilizes a deformed optical model to calculate neutron transmission coefficients and cross sections, a giant-dipole-resonance model to determine gamma-ray transmission coefficients, and Hauser-Feshbach statistical theory to calculate partial reaction cross sections.

II. NUCLEAR MODEL CALCULATIONS

The Hauser-Feshbach statistical-theory calculations were performed with the COMNUC and GNASH reaction theory codes.² The COMNUC calculations include width-fluctuation correction which are important at lower energies, and the GNASH calculations incorporate preequilibrium effects, which become significant at higher energies. COMNUC was used to calculate all cross sections below $E_n = 8$ MeV, whereas GNASH was used for calculations above 8 MeV and for all continuous spectral calculations. Both codes utilize the Gilbert and Cameron level density formulation and the Cook tabulation of level density parameters.² A maximum amount of experimental information concerning discrete energy levels was incorporated into the calculations, and the constant temperature part of the Gilbert and Cameron level density was matched to the discrete level data for each residual nucleus in the the analysis.

III. EVALUATION RESULTS

Resolved resonance parameters from ENDF/B-V are used to represent the cross sections from 10^{-5} eV to 151.92 eV, with some adjustment made to the background cross sections to improve agreement with thermal and resonance integral data. Above 152 eV, the cross sections are joined smoothly to results from the theoretical analysis described above. Above 10 keV, all cross sections were obtained from the theoretical calculations, as were the secondary angular and energy distributions. See the attached ENDF File 1 comment section for additional details and for references.

The evaluated ^{165}Ho total neutron cross section is compared to experimental data and to ENDF/B-V in Fig. 1. Similarly, the evaluated $^{165}\text{Ho}(n,\gamma)$ cross section is compared to the Version V evaluation and to a selection of experimental data in Fig. 2. Finally, evaluated total γ -ray emission spectra that come from the above theoretical analysis are compared to experimental data at $E_n = 4$ and 420 keV in Fig. 3. The spectra are dominated obviously by radiative capture at both energies.

¹ P. G. Young, "Reaction Theory Calculations of $n + ^{165}\text{Ho}$ Reactions," in *Applied Nuclear Science Research and Development Progress Report*, June 1, 1985 - Nov. 30, 1985 (Cp. E. D. Arthur and A. D. Mutschlecner, 1986) LA-10689-PR, p. 53.

² See the ENDF/B File 1 comment section (attached) for references.

$n + {}^{165}\text{Ho}$ Total Cross Section

Figure 1. Comparison of evaluated and experimental values of the neutron total cross section of ${}^{165}\text{Ho}$. The solid curve represents the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points are experimental data as indicated.

$^{165}\text{Ho}(n,\gamma)^{166}\text{Ho}$ Cross Section

Figure 2. Comparison of evaluated and experimental values of the $^{165}\text{Ho}(n,\gamma)$ cross section. See caption of Fig. 1 for details of curves and symbols.

Figure 3. Total gamma-ray emission spectra from $n + {}^{165}\text{Ho}$ reactions obtained from the ENDF/B-VI evaluation compared with experimental data for $E_n = 4$ and 420 keV.

¹⁶⁵₆₇Ho

Reference: No Primary Reference
Evaluators: P. G. Young and E. D. Arthur
Evaluated: April 1988
Material: 6725
Content: Neutron transport, Gamma production

Resolved Resonance Range 1.0E-5 to 151.92 eV.

MF=2 MT=151 Resonance parameters from old BNL-325 Ref.(1).
from ENDF/B-V Schenter and Schmittroth evaluation.

References

1. S.F. Mughabghab and D.I. Garber, BNL-325,3ed,Vol 1(June 1973).

2200m/s capture cross section, barns.

(from resonance parameters) =	20.7446 b
(from 1/v component) =	43.9554 b
total =	64.700 b

Energy range above the resonance region.

The evaluation above 10 keV is based on a detailed theoretical analysis utilizing the available experimental data. Coupled channel optical model calculations with the ECIS code (Ra70) were used to provide the total, elastic, and inelastic cross sections to the first 3 members of the ground state rotational band, as well as neutron elastic and inelastic angular distributions to the rotational levels. The ECIS code was also used to calculate neutron transmission coefficients. Hauser-Feshbach statistical theory calculations were carried out with the GNASH (Ar88, Yo77) and COMNUC (Du70) code systems, including preequilibrium contributions. Systematics were used to obtain parameters for the exciton preequilibrium model, with small adjustments made to improve agreement with available exp. data. The Gilbert-Cameron level density model was used to supplement available experimental information on low-lying levels (Gi65). The Brink-Axel model (Br55,Ax62) was used to calculate gamma-ray transmission coefficients, using gamma-ray strength function results compiled by Mughabghab (Mu84). A description of the calculations is given in Yo86.

*****MF=3 Smooth Cross Sections*****

MT=1 Neutron Total Cross Section. 0.01 to 30 MeV, based on coupled-channel optical calculations, which were optimized to the available experimental data (Mc88).
 MT=2 0.030 to 30 MeV, based on subtraction of MT=4,16,17,37, and 102 from MT=1. Note that this corresponds exactly to using the results of the coupled-channel optical and Hauser-Feshbach model results.
 MT=4 Sum of MT=51-91
 MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=51,52 Thres. to 30 MeV, Coupled-channel optical model calculations plus compound-nucleus contributions.
 MT=53-63 Threshold to 8.0 MeV, Compound nucleus reaction theory calculations using the COMNUC code (Du70) and including width fluctuation corrections. Transmission coefficients from cc optical model calculations used.
 MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****MF=4 Neutron Angular Distributions*****

MT=2 Elastic scattering angular distribution based on ECIS coupled-channel calculations, with a compound elastic component from COMNUC included below 8 MeV.
 MT=16 (n,2n) distributions assumed isotropic in the laboratory system.
 MT=17 (n,3n) distributions assumed isotropic in the laboratory system.
 MT=37 (n,4n) distributions assumed isotropic in the laboratory system.
 MT=51,52 Thres. to 30 MeV, Coupled-channel optical model calculations plus compound-nucleus contributions.
 MT=53-63 Threshold to 8.0 MeV, Compound nucleus reaction theory calculations using the COMNUC code (Du70) and including width fluctuation corrections. Transmission coefficients from cc optical model calculations used.
 MT=91 (n,n'continuum) distributions assumed isotropic in the laboratory system.

*****MF=5 Neutron Energy Distributions*****

MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc. Tabulated laboratory distributions given.
 MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.

Tabulated laboratory distributions given.
 MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 Tabulated laboratory distributions given.
 MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 Tabulated laboratory distributions given.

*****MF=12 Photon Multiplicities*****

MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 Note that photons from (n,gn') reactions are included
 in MF=12,MT=102 but not in MF=3,MT=102, which causes
 the multiplicities at higher energies to become
 somewhat large.

*****MF=13 Photon Production Cross Sections*****

MT=4 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****MF=14 Photon Angular Distributions*****

MT=4 Isotropy assumed.
 MT=16 Isotropy assumed.
 MT=17 Isotropy assumed.
 MT=37 Isotropy assumed.
 MT=102 Isotropy assumed.

*****MF=15 Photon Energy Distributions*****

MT=4 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calc.
 MT=102 GNASH Hauser-Feshbach statistical/preequilibrium calc.

*****References*****

Ar88 E.D.Arthur, LA-UR-88-382 (1988).
 Ax62 P.Axel, Phys.Rev.126, 671 (1962).
 Br55 D.M.Brink, D.Ph. Thesis, Oxford (1955).
 Du70 C.L.Dunford, AI-AEC-12931 (1970).
 Mc88 V.McLane et al., Neutron Cross Sections V2 (Acad.Pr.1988).
 Mu84 S.F.Mughabghab, Neutron Cross Sections V1 (Acad.Pr.1986).
 Ra70 J.Raynal, IAEA SMR-9/8 (1970).
 Yo77 P.G.Young & E.D.Arthur, LA-6947 (1977).
 Yo86 P.G.Young, LA-10689-PR (1986) p.53.

¹⁶⁶Er
68

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, and F. Schmittroth
Evaluated: December 1988
Material: 6837
Content: Fission product

File Comments

HEDL Eval-Apr74 R. E. Schenter and F. Schmittroth
ORNL Eval-Dec88 R. Q. Wright

The ¹⁶⁶Er evaluation, MAT 9875, was revised by R. Q. Wright in March 1989. The new evaluation is assigned MAT No. 6837.

Summary of Changes

The resolved resonance parameters for MAT 6837 are taken from Ref. 1 ($E_{li} = 2007.9$ eV). The bound level at -40.4 eV has $\Gamma_n = 0.4887$ eV and $\Gamma_\gamma = 0.092$ eV. This choice gives the desired value for the thermal capture cross section, 19.60 b. Values of Γ_γ not given in Ref. 1 are set to 0.092 eV. The value for the scattering radius is 0.81 (from Ref. 1), and the highest energy resonance included is at 2128.9 eV.

In File 3 the total, elastic, and capture cross sections are set to zero in the resolved resonance range (1.0×10^{-5} to 2007.9 eV). The elastic cross section at 2007.9 eV is reduced to 17.2 b. and has been generally reduced for energies up to 100 keV.

The elastic cross section at 2 keV is based on the data of Vertebnyi et al. (see Ref. 2, page 564). The capture cross section is revised in the energy range 3 - 200 keV. The capture cross section is based on the data of Kononov et al. and is 40 to 50% higher than ENDF/B-V in the energy range 10 - 100 keV. See Ref. 2, page 561 for a plot of the Kononov data ($\Delta 78$ FEI Ko). The new evaluation is slightly higher than the "eye guide" for the range 15 to 50 keV; the capture cross section is 600 mb at 30 keV. The capture cross section is unchanged from ENDF/B-V above 200 keV.

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	19.60
(from 1/v component)	=	0.00
Total	=	19.60

computed resonance integral = 99.20

References:

1. S. F. Mughabghab, "Neutron cross sections," Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B: Z = 61-100, Academic Press (1984).
2. V. Mclane, C. L. Dunford, and P. F. Rose, "Neutron Cross Sections," Vol. 2, Academic Press, New York (1988).

Summary of ENDF/B-V Evaluation

MF=2 MT=151 Resonance parameters from BNL-325 Ref. (3).

MF=3 MT= 1 Total cross section calculated using the Moldauer potential from Ref. (4) for $E > E_{hi}$.

MF=3 MT= 2 The elastic cross section was obtained from $\sigma_t = \sigma_e + \sigma_{in}$ for $E > E_{hi}$.

MF=3 MT= 4, 51,52,...,91 Inelastic cross sections calculated using COMNUC-3 Refs. (5, 6).

MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A 1/v component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$.

MF=4 MT=2 Angular distributions assumed isotropic.

MF=4 Non-elastic angular distributions assumed isotropic.

MF=5 MT= 91 Evaporation spectrum (LF = 9) parameters obtained using NCAP code Ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters	=	1.67
(from 1/v component)	=	33.33
Total	=	35.00

computed resonance integral = 141.12

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL-325, 3ed, Vol 1 (June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

$^{166}_{88}\text{Er } \sigma_{n,\text{tot}}$

77	LIN	Djumin+	77KIEV 2, 74	1	5.270 b	at 14.20 MeV
65	IFU	Vertebnyj+	65ANTWERP, 572(186)	121	11.60 mv	to 0.113 eV

$^{166}_{88}\text{Er } \sigma_{n,\gamma}$

78	FEI	Kononov+	YF 27, 10	38	6.500 keV	to 0.335 MeV $^{197}\text{Au } \sigma_{n,\gamma}$
74	FEI	Shorin+	YF 19, 5	32	5.280 keV	to 68.20 keV $^{197}\text{Au } \sigma_{n,\gamma}$

¹⁶⁷Er
68

Reference: No Primary Reference
Evaluators: R. Q. Wright, R. E. Schenter, Others
Evaluated: December 1988
Material: 6840
Content: Fission product

File Comments

HEDL Eval-Apr74 R. E. Schenter and F. Schmittroth
ORNL Eval-Dec88 R. Q. Wright

THE ¹⁶⁷Er evaluation, MAT 9876, was revised by R. Q. Wright in March 1989. The new evaluation is assigned MAT No. 6840.

Summary of Changes

The resolved resonance parameters were taken from Ref. 1 ($E_{hi} = 500.0$ eV). In all, 113 resonances are included, up to $E_0 = 518.9$ eV. Thirty-one resonances did not have values given for "J", 12 are assigned to $J = 3$ and the remaining 19 to $J = 4$. Values of Γ_γ not given in Ref. 1 are set to 0.089 eV. The value for the scattering radius is 0.79 fm (from Ref. 1), and the highest energy resonance included is 518.90 eV.

In the unresolved resonance range unresolved parameters originally from Ref. 2 were modified as follows:

Unresolved resonance range	500.0 to 10000 eV
Scattering radius	0.79 (same as resolved range)
Γ_γ	0.089 eV (same as resolved range)
Average reduced neutron widths	All increased by 20%

The impact of this change was to increase capture by about 10% at all energies. The elastic cross section was increased by about 20% at 500 eV and by about 10% at 10000 eV. The resulting value for capture at 10000 eV as computed from the

unresolved parameters is 2.70 b, which agrees well with the value from File 3 (2.724 b).

In file 3 the total, elastic, and capture cross sections are set to zero in the resolved and unresolved resonance ranges (1.0×10^{-5} to 10000 eV). The elastic cross section at 10000 eV is reduced to 14.6 b; and other reductions were made in the elastic cross section up to 100 keV.

In 1989 the capture cross section was revised in the energy range 10 - 370 keV. The capture is based on the data of Shorin et al. and is 25 to 30% higher than ENDF/B-V in the energy range 10 - 100 keV. See Ref. 3, page 562 for a plot of the Shorin data (Δ 74 FEI Sh). The capture cross section is 1500 mb at 30 keV. The total and elastic cross sections were also changed to insure that the total is the sum of the partials (10 - 370 keV).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	657.8
(from 1/v component)	=	0.0
Total	=	657.8

computed resonance integral = 2991

References:

1. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B: Z = 61 - 100, Academic Press (1984).
2. J. Hardy, Jr., Bettis Atomic Power Laboratory, "ENDF/B Data Sets for ^{166}Er and ^{167}Er ," Personal communication to Lester Petrie, ORNL, April 11, 1988.
3. V. McLane, C. L. Dunford, and P. F. Rose, "Neutron Cross Sections," Vol. 2, Academic Press, New York (1988).

Summary of ENDF/B-V Evaluation

MF=2 MT=151 Resonance parameters from BNL-325 Ref. (3).

MF=3 MT= 1 Total cross section calculated using Moldauer potential from Ref. (4) for $E > E_{thr}$.

Summary of ENDF/B-V, Continued

- MF=3 MT= 2 Elastic cross section from $\sigma_t - \sigma_c - \sigma_{in}$ for $E > E_{hi}$.
- MF=3 mt= 4, 51,52,,.,91 Inelastic cross sections calculated using COMNUC-3 refs. (5, 6).
- MF=3 MT=102 Neutron capture evaluated using methods (NCAP code) in Refs. (1, 2) for $E > E_{hi}$. A $1/v$ component was added to give the 2200 m/s cross section of Ref. (3) for $E < E_{hi}$.
- MF=4 MT=2 Angular distributions assumed isotropic.
- MF=4 Non-elastic angular distributions assumed isotropic.
- MF=5 MT= 91 Evaporation spectrum (LF = 9) parameters obtained using NCAP code Ref. (2).

The 2200 m/s capture cross section, barns

(from resonance parameters)	=	655.71
(from $1/v$ component)	=	14.29
Total	=	670.00

computed resonance integral = 2977.00

References

1. F. Schmittroth and R. E. Schenter, HEDL TME 73-63 (Aug 1973).
2. F. Schmittroth, HEDL TME 73-79 (Nov 1973).
3. S. F. Mughabghab and D. I. Garber, BNL-325, 3ed, Vol 1 June 1973).
4. P. A. Moldauer, Nucl. Phys. 47 (1963) 65.
5. C. L. Dunford, AI-AEC-12931 (July 1970).
6. C. L. Dunford (Private Communication).

Yr	Lab	Author	Reference	Points	Range	Standard
----	-----	--------	-----------	--------	-------	----------

$^{167}_{88}\text{Er } \sigma_{n,\text{tot}}$

77	LIN	Djumin+	77KIEV 2, 74	1	3.520b	at 14.20MeV
65	IFU	Vertebnyj+	65ANTWERP, 572(186)	121	11.60mv	to 0.113eV
60	BNL	Moller+	NSE 8, 183	86	0.206eV	to 0.889eV

$^{167}_{88}\text{Er } \sigma_{n,\gamma}$

74	FEI	Shorin+	YF 19, 5	31	6.040keV	to 68.20keV $^{197}\text{Au } \sigma_{n,\gamma}$
----	-----	---------	----------	----	----------	--

¹⁸⁵Re
⁷⁵

Reference: No Primary Reference
Evaluators: L. W. Weston, P. G. Young, Others
Evaluated: March 1990
Material: 7525
Content: Neutron transport, Covariances

File Comments

ORNL	Eval-Mar90 L. W. Weston
LANL	Eval-Mar90 P. G. Young
GE-NMPO	Eval-Jan68 W. B. Henderson, J. W. Zwick

The previous version of this file was evaluated by W.B. Henderson and J. W. Zwick at GE-NMPO, in January 1968. For the ENDF/B-VI evaluation Files 2, 3 and 33 were extensively revised.

MF=1 MT=453 Radioactive Decay Data. Decay constants were derived from the half lives of the ground state (1). Daughters are from Ref (1) except for ¹⁸⁶Os which was made a daughter of ¹⁸⁶Re, lacking a branching ratio specification, since 95 % of the decays go that way (2).

MF=2 MT=151 Resonance parameters. The resolved resonance parameters were taken directly from the evaluation of Mughabghab, Ref (3). The resolved resonance region extends from 10⁻⁵ to 2000 eV. Values calculated at 0.0253 eV:

Total Cross Section = 121.0 barns

Capture Cross Section = 112.2 barns

See the note in MF=3 regarding the scattering radius. The unresolved resonance parameters were derived primarily from the resolved resonance parameters (Ref. 3) with some input from the fit to the MF=3 smooth cross sections. Unresolved resonance parameters are to be used only for calculating self shielding factors. The unresolved resonance parameters extend from 2 to 35 keV.

- MF=3 Smooth Cross Sections. The smooth cross sections from 2 to 125 keV were derived from a fit to the data of R. L. Macklin and P. G. Young, Ref (4), on natural rhenium using the code FITACS by F. H. Froehner, Ref (5). The isotopic separation was done with average resonance parameters derived from previous resolved and unresolved data. Capture from this fit extends to 400 keV. Above 125 keV the model code calculations of P. G. Young, Ref (4), were used for the smooth cross sections. A lower scattering radius (7.9 fm) and p-wave strength function (.40) than evaluated in Ref. 3 was necessary to fit the data above 2 keV. See Ref. 4 for comparison with experimental data.
- MF=4 MT=2 Elastic Secondary Angular Distributions. The transfer matrix and legendre coefficients were computed using the code CHAD (6).
- MF=5 MT=4, 16 and 17 Secondary energy distributions. Discrete level and continuum (n, n', γ) cross sections to 1.5 MeV were obtained from ABACUS-NEARREX calculations (Ref. 7-9). Above 1.5 MeV a continuum treatment was used. Each continuum cross section was treated as a Maxwellian with $T = \sqrt{E/a}$ and $a = (25 \text{ MeV})^{-1}$.

References

1. D. Goldman and J. Roesser, Chart of the Nuclides, 9th Ed., Knolls Atomic Power Laboratory 11/66.
2. "Nuclear data," Section B, Vol.1, No. 2, Academic Press, June 1966.
3. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B, Brookhaven National Laboratory, National Nuclear Data Center 1984.
4. R. L. Macklin and P. G. Young, Nucl. Sci. & Eng. 97, 239, 1987.
5. F. H. Froehner, Kernforschungszentrum Karlsruhe, Private Communication, 1986.
6. R. Berland, NAA-SR-11231, Dec. 31, 1965. BNWL Version, Modified by GE-NMPO.

7. ABACUS-NEARREX, Undocumented Optical Model Code System, Version of 2/5/66, Brookhaven National Laboratory.
8. E. H. Auerbach, ABACUS-II, BNL Informal Report BNL-6562 (1962).
9. P. A. Moldauer et al., "NEARREX, A Computer Code for Nuclear Reaction Calculations," Argonne National Laboratory, ANL-6978 (1984), See also P. A. Moldauer, Rev. Mod. Phys. 36, 1079 (1964).

Fig. 1. Comparison of ENDF/B-VI (solid line) with ENDF/B-V (dashed line) for ^{185}Re . Top curves are total cross section. Elastic scattering curves are next to top. Inelastic scattering curves have a threshold just above 100 keV. The capture cross curves are the lowest.

¹⁸⁷
⁷⁵Re

Reference: No Primary Reference
Evaluators: L. W. Weston, P. G. Young, Others
Evaluated: March 1990
Material: 7531
Content: Neutron transport, Covariances

File Comments

ORNL	Eval-Mar90 L. W. Weston
LANL	Eval-Mar90 P. G. Young
GE-NMPO	Eval-Jan68 W. B. Henderson, J. W. Zwick

The previous version of this file was evaluated by W. B. Henderson and J. W. Zwick at GE-NMPO in January 1968. For the ENDF/B-VI evaluation Files 2, 3 and 33 were extensively revised.

MF=1 MT=453 Reaction Branching Ratios. Decay constants were derived from the half lives of the ground state (1). Daughters are from Ref (1) except ¹⁸⁶Os which was made a daughter of ¹⁸⁶Re, lacking a branching ratio specification, since 95 % of the decays go that way (2).

MT=457 Radioactive Decay Data. Radioactive decay data were prepared for the evaluation in January 1974 by C. W. Reich (ANC). Q-values were from the 1973 revision of the Wapstra-Gove mass tables. Half-lives were from N. E. Holden, Chart of the Nuclides and Private Communication January (1974). Also see W. B. Ewbank, Nuclear Data B1, No. 2, 23 (1966). Note: A first-forbidden, unique shape correction was considered in deriving E_{β} for ground-state beta transitions.

MF=2 MT=151 Resonance Parameters. Resolved resonance parameters are directly from the evaluation of S. F. Mughabghab, Ref (3). The resolved energy range spans 10^{-5} to 2000 eV. Values calculated at 0.0253 eV:

Total cross section	=	86.6 barns
Capture cross section	=	76.7 barns

- MF=2 MT=151 Continued. See note in MF=3 concerning the scattering radius. The unresolved resonance parameters were derived primarily from the resolved resonance parameters (Ref. 3) with some input from the fit to the MF=3 smooth cross sections. Unresolved resonance parameters are to be used only for calculating self shielding factors. The unresolved parameters extend from 2 to 35 keV.
- MF=3 Smooth Cross Sections. The smooth cross sections from 2 to 125 keV were derived from a fit to capture data on natural rhenium by R. L. Macklin and P. G. Young, Ref (4), using the code FITACS, by F. H. Froehner, Ref (5). The isotopic separation was done with average resonance parameters determined from previous resolved and unresolved data. Above 125 keV the model code calculations of P. G. Young, Ref (4), were used for smooth cross sections. A lower scattering radius (7.9 fm) and p-wave strength function (.28) than evaluated in Ref. 3 was necessary to fit data above 2 keV. See Ref. 4 for comparison with experimental data.
- MF=4 MT=2 Elastic Secondary Angular Distributions. The transfer matrix and legendre coefficients were computed using the CHAD code (6).
- MF=5 MT=4, 16 and 17 Secondary energy distributions. Discrete level and continuum(n,n', γ) cross sections to 1.5 MeV were obtained from ABACUS-NEARREX calculations. (Ref. 7-9) Above 1.5 MeV everything was treated as a continuum. Each continuum cross section was specified as Maxwellian with $T = \sqrt{E/a}$ and $a = (25 \text{ MeV})^{-1}$.

References

1. D. Goldman and J. Roesser, Chart of the Nuclides, 9th Ed., Knolls Atomic Power Laboratory 11/66.
2. "Nuclear Data," Section B, Vol. 1, No. 2, Academic Press, June 1966.
3. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B, Brookhaven National Laboratory, National Nuclear Data Center (1984).

4. R. L. Macklin and P. G. Young, Nucl. Sci. & Eng. 27, 239, 1987.
5. F. H. Froehner, Kernforschungszentrum Karlsruhe, Private Communication, 1986.
6. R. Berland, NAA-SR-11231, Dec. 31, 1965. BNWL version, modified by GE-NMPO.
7. ABACUS-NEARREX, Undocumented Optical Model Code System, Version of 2/5/66, Brookhaven National Laboratory.
8. E. H. Auerbach, ABACUS-II, BNL Informal Report BNL-6562 (1962).
9. P. A. Moldauer et al., "NEARREX, A Computer Code for Nuclear Reaction Calculations, Argonne National Laboratory Report ANL-6978 (1964), See also P. A. Moldauer, Rev. Mod. Phys. 36, 1079 (1964).

Fig. 1. Comparison of ENDF/B-VI (solid line) with ENDF/B-V (dashed line) for ^{187}Re . Top curves are total cross section. Elastic scattering curves are next to top. Inelastic scattering curves have a threshold just above 100 keV. The capture cross curves are the lowest.

SUMMARY DOCUMENTATION FOR ^{197}Au ENDF/B-VI, MAT = 7925

P. G. Young and E. D. Arthur

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

The ENDF/B-VI evaluation for ^{197}Au combines results from a new theoretical analysis¹ above the resonance region with the ENDF/B-V resonance parameter evaluation and with the ENDF/B-VI standard cross section analysis. The analysis involves use of a deformed optical model to calculate neutron transmission coefficients, a giant-dipole-resonance model and experimental data to determine gamma-ray transmission coefficients, and Hauser-Feshbach statistical theory to calculate partial reaction cross sections. Particular emphasis was given to obtaining gamma-ray strength functions that are consistent with spectral measurements of gamma-ray emission between $E_n = 0.2$ and 20 MeV by Morgan and Newman,² while at the same time requiring agreement with (n,γ) and (n,xn) cross section data.

II. THEORETICAL ANALYSIS

The deformed optical model parameterization by Delaroche² was utilized in the analysis. The coupled-channel code ECIS was used with the lowest three states of the ^{197}Au ground-state rotational band coupled in the calculations ($J^\pi = 3/2^+, 5/2^+, 7/2^+$ at $E_x = 0, 279, 548$ keV, respectively). Neutron transmission coefficients were calculated to 20 MeV with ECIS and were collapsed to a form dependent only on incident neutron energy and orbital angular momentum for use in the Hauser-Feshbach calculations.

The Hauser-Feshbach statistical-theory calculations were performed with the COMNUC and GNASH reaction theory codes.² The COMNUC calculations include width-fluctuation corrections, which are important at lower energies, and the GNASH calculations incorporate preequilibrium effects, which become significant at higher energies. COMNUC was used to calculate all cross sections below $E_n = 3$ MeV, whereas GNASH was used for calculations above 3 MeV and for all continuous spectral calculations. Both codes utilize the Gilbert and Cameron level density formulation and the Cook tabulation of level density parameters.² A maximum amount of experimental information concerning discrete energy levels was incorporated into the calculations, and the constant temperature part of the Gilbert and Cameron level density was matched to the discrete level data for each residual nucleus in the analysis. Gamma-ray transmission coefficients were calculated from E1 and M1 strength functions. The shapes of the E1 strength functions were determined for $E_\gamma < 8$ MeV by matching trial and error calculations of gamma-ray spectra from $^{197}\text{Au}(n,\gamma)$ reactions with the data of Morgan and Newman.² Above $E_\gamma = 8$ MeV, the empirically determined E1 strength functions were joined to a

¹ P. G. Young and E. D. Arthur, "Analysis of $n + ^{197}\text{Au}$ Cross Sections for $E_n = 0.01 - 20$ MeV," Proc. Int. Sym. on Capture Gamma-Ray Spectroscopy and Related Topics, Knoxville, Tenn., Sept. 10-14, 1984 (Ed. S. Raman, 1985), AIP Conf. Proc. No. 125, p. 530.

² See references in ENDF/B File 1 comments (attached).

giant-dipole resonance shape. A giant dipole resonance shape was also utilized for the M1 strength functions.

III. EVALUATION RESULTS

Resolved resonance parameters from ENDF/B-V are used to represent the cross sections from 10^{-5} eV to 5 keV. From 5 keV to 2.5 MeV, the radiative capture cross section from the ENDF/B-VI simultaneous standards analysis is utilized in the evaluation. Except for the neutron total cross section, all other smooth cross sections were calculated from the theoretical analysis described above, as were the secondary angular and energy distributions. The evaluated total cross section was obtained from a covariance analysis of the available experimental data using the GLUCS analysis code. See the attached ENDF File 1 comment section for additional details and references.

The evaluated ^{197}Au total neutron cross section is compared to experimental data and to ENDF/B-V in Fig. 1. Similarly, the evaluated $^{197}\text{Au}(n,\gamma)$ cross section is compared to the Version V evaluation and to a selection of experimental data in Fig. 2, and comparisons of $^{197}\text{Au}(n,xn)$ calculated and measured cross sections are given in Fig. 3. Finally, evaluated total γ -ray emission spectra that come from the above theoretical analysis are compared to experimental data at $E_n = 0.8$ and 11.0 MeV in Fig. 4. Note that the spectrum at 0.8 MeV is dominated by the $^{197}\text{Au}(n,\gamma)$ reaction, whereas both $(n,n'\gamma)$ and $(n,2n\gamma)$ reactions are important in the 11-MeV spectrum.

$n + {}^{197}\text{Au}$ Total Cross Section

Figure 1. Comparison of evaluated and experimental values of the neutron total cross section of ${}^{197}\text{Au}$. The solid curve represents the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points are experimental data as indicated.

Figure 2. Comparison of evaluated and experimental values of the $^{197}\text{Au}(n,\gamma)$ cross section. See caption of Fig. 1 for details of curves and symbols.

Figure 3. Comparison of the evaluated $^{197}\text{Au}(n,2n)^{196}\text{Au}$ and $^{197}\text{Au}(n,3n)^{195}\text{Au}$ cross sections with experimental data and with ENDF/B-V. See caption of Fig. 1 for details of curves and symbols.

Figure 4. Total gamma-ray emission spectra obtained from the ENDF/B-VI evaluation compared with experimental data for $E_n = 0.8$ and 11 MeV.

¹⁹⁷
⁷⁹Au

Reference: LA-10069-PR
Evaluator: P. G. Young
Evaluated: January 1984
Material: 7925
Content: Standard, Neutron transport, Gamma production, Co-variances

***** SUMMARY *****

A new evaluation of all neutron and gamma-ray data above the resonance region is joined with the ENDF/B-V resolved resonance region evaluation and with the Version VI standard cross section for the (n,gamma) reaction below a neutron energy of 2.5 MeV.

***** GENERAL DESCRIPTION *****

P.G.Young and E.D.Arthur

The new evaluation for Files 3,4,5,12,13,14,15 is based on statistical theory, Hauser-Feshbach, preequilibrium calculations with the COMNUC and GNASH codes (Ref 1,2). Deformed optical potential of DeLaroche and ECIS coupled-channel code were used to calculate neutron transmission coefficients and total and elastic elastic cross sections (Ref 3,4). Gamma-ray strength functions were obtained by fitting Morgan n,xg data (Ref 5) at 0.4 and 6.5 MeV. Calculated results were used for all major reactions except total cross section. For total, the theoretical cross section was used as a prior in covariance analysis of experimental data using GLUCS code (Ref 6). More details on experimental data used are given below and in the main reference for the evaluation (Ref 7).

***** MF=2 RESONANCE PARAMETERS *****

MT=151 Resolved resonance parameters given from 1.0E-05 eV to 2 keV based on Ref 8 and references therein and a bound level. Some of the resonance spin assignments from Ref 9. From 2 to 4.827 keV the parameters are based on Macklin et al and Hoffman et al normalized data. See Refs 10 and 11.
Thermal cross sections are as follows:
capture = 98.71 b
scattering = 6.84 b

total = 105.55 b

The absorption resonance integral is 1559 b.

***** MF=3 SMOOTH NEUTRON CROSS SECTIONS *****

MT= 1 Total cross section. Based on GLUCS covariance analysis using deformed optical model calculation as the prior and experimental data from Refs 12-22, 29 for fitting.

MT= 2 Elastic cross section. Difference of MT=1 and sum of all nonelastic cross sections. Closely approximates theoretical results.

MT= 4 Inelastic cross section. Sum of MT=51-63, 91.

MT= 16 (n,2n) cross section. Theoretical calculation used.
In good agreement with exp. below 23 MeV. See Refs 23-25.

MT= 17 (n,3n) cross section. Theoretical calculation used.
In good agreement with exp. at all energies (Refs 24,25).

MT= 37 (n,4n) cross section. Theoretical calculation used.
In reasonable agreement with data of Ref 25.

MT=51-63 (n,nprime) cross sections to levels. Except for MT=53 and 56, all are from compound-nucleus calculations with the COMNUC code. MT=53 and 56 also include direct reaction components from ECIS calculations (MT53 and 56 are the 5/2+ and 7/2+ members of the ground state rotational band) and extend to 30 MeV. MT=51,52,54,55,57-63 are zeroed above 6 MeV.

MT= 91 Inelastic continuum cross sections from GNASH theoretical calculations. Includes (n,gn) component from 0.1 to 2.0 MeV. Conventional (n,ng) continuum starts at 1.2236 MeV. Q-value has no significance except corresponds to thres.

MT=102 (n,gamma) cross section. Below 2.5 MeV, adopted the ENDF/B-VI standard cross section (Ref.30,31) down to the resonance region. At higher energies, the theoretical calculations were adjusted to agree with experimental data. A semi-direct component normalized to an average of experimental data at 14 MeV was included above En = 6 MeV.
At higher energies, used theoretical calculations, which agree reasonably with available exp. data. Above 5 MeV, calculation includes semi-direct component normalized to average of 14 MeV data.

MT=103 (n,p) cross section. Adopted ENDF/B-V with smooth extrapolation to 30 MeV. Based on exp data of Ref 26.

MT=107 (n,alpha) cross section. Adopted ENDF/B-V with smooth extrapolation to 30 MeV. Based on data of Ref 26.

***** MF=4 NEUTRON ANGULAR DISTRIBUTIONS *****

MT= 2 Elastic scattering. Legendre coefficients obtained by combining ECIS direct reaction calculations with COMNUC compound nucleus results.

MT= 16 (n,2n) angular distribution. Used Kalbach-Mann (Ref 27).
 Semi-empirical shape averaged over the emitted neutron
 spectrum at each incident neutron energy.
 MT= 17 (n,3n) angular distribution. Same comment as MT=16.
 MT= 37 (n,4n) angular distribution. Same comment as MT=16.
 MT=51-63 (n,nprime) level angular distributions. Legendre coef-
 -ficients obtained from COMNUC compound nucleus calculations.
 For MT=53 and 56, ECIS direct reaction results were combined
 with the compound nucleus calculations.
 MT= 91 (n,nprime) continuum. Same comment as for MT=16.

***** MF=5 NEUTRON ENERGY DISTRIBUTIONS *****

MT= 16 (n,2n) tabulated distribution from GNASH calculations.
 MT= 17 (n,3n) tabulated distribution from GNASH calculations.
 MT= 37 (n,4n) tabulated distribution from GNASH calculations.
 MT= 91 (n,nprime) continuum tabulated distribution obtained from
 GNASH calculations.

***** MF=8 RADIOACTIVE DECAY DATA *****

MT= 16 Decay data for the 10 hour metastable sixth excited state
 in Au-196. ENDF/B-V data adopted without change.

***** MF=10 RADIOACTIVE NUCLIDE CROSS SECTIONS *****

MT= 16 Production cross section for the 10-hour metastable sixth
 excited state of Au-196 through (n,2n) reactions. ENDF/B-V
 data adopted, with smooth extrapolation to 30 MeV.

***** MF=12 PHOTON MULTIPLICITIES *****

MT=102 (n,gamma) yield at low energies obtained by requiring
 energy conservation with MF=15, MT=102 results. Beginning
 near 10 keV, GNASH results used.

***** MF=13 PHOTON CROSS SECTIONS *****

MT= 4 Gamma-ray production cross sections obtained from GNASH
 calculations for continua regions and from COMNUC for
 discrete levels. ECIS was used to calculate direct react-
 tion contributions for 3rd and 6th levels of Au-197.
 MT= 16 Gamma-ray production cross sections obtained from GNASH
 calculations at all incident neutron energies.
 MT= 17 gamma-ray production cross sections obtained from GNASH
 calculations at all incident neutron energies.
 MT= 37 gamma-ray production cross sections obtained from GNASH
 calculations at all incident neutron energies.

***** MF=14 PHOTON ANGULAR DISTRIBUTIONS *****

MT= 4 Photons from inelastic scattering assumed isotropic.
 MT= 16 Photons from (n,2n) reactions assumed isotropic.
 MT= 17 Photons from (n,3n) reactions assumed isotropic.
 MT= 37 Photons from (n,4n) reactions assumed isotropic.
 MT=102 Photons from (n,gamma) reactions assumed isotropic.

***** MF=15 PHOTON ENERGY DISTRIBUTIONS *****

MT= 4 Inelastic scattering photon tabulated distributions
 obtained from GNASH calculations for continua regions and
 from COMNUC for discrete levels. Direct contributions for
 MT=53 and MT=56 obtained from ECIS calculations.
 MT= 16 (n,2n) photon tabulated distributions obtained from
 GNASH calculations.
 MT= 17 (n,3n) photon tabulated distributions obtained from
 GNASH calculations.
 MT= 37 (n,4n) photon tabulated distributions obtained from
 GNASH calculations.
 MT=102 (n,gamma) tabulated thermal distribution obtained from
 experimental data of Ref 28. Thermal spectrum linearly inter-
 polated to GNASH calculation at 10 keV. GNASH results used
 at higher energies.

***** MF=33 NEUTRON CROSS SECTION COVARIANCES *****

MT= 1 Total cross section covariance from GLUCS analysis.

***** REFERENCES *****

1. C.L.Dunford. AI-AEC-12931(1970).
2. P.G.Young, E.D.Arthur, LA-6947 (1977).
3. J.P.DeLaroche, Harwell Conference (1978)p.366.
4. J.Raynal, IAEA SMR-9/8 (1970).
5. G.L.Morgan, E.Newman, ORNL-TM-4973 (1975).
6. D.M.Hetrick, C.Y.Fu, ORNL/TM-7341 (1980).
7. P.G.Young, E.D.Arthur, LA-10069-PR (1984)p.12.
8. S.F.Mughabghab and D.I.Garber BNL-325,3rd edn, vol 1(1973).
9. A.Lottin and A.Jain, Conf on Nuclear Structure Study with
 Neutrons, Budapest,1972 p34 and private communication.
10. R.Macklin et al. Phys. Rev/C 11,1270(1975) and private
 communication.
11. M.M. Hoffman et al. 71 Knoxville Conf., p868 (1971).
12. W.Poenitz et al., Nuc.Sci.Eng. 78, 333(1981).
13. D.G.Foster Jr., D.Glasgow, Phys.Rev. C3, 576(1971).
14. K.K.Seth,Phys.Letters,16,306(1965).

15. S.C.Snowdon, Phys.Rev. 90, 615(1953).
16. J.F.Whalen, ANL-7210,16(1966).
17. N.Nereson, Phys.Rev. 94, 1678(1954).
18. A.Bratenahl et al., Phys.Rev. 110, 927(1958).
19. J.P.Conner,Phys.Rev.109,1268(1958).
20. J.H.Coon,Phys.Rev.88,562(1952).
21. J.M.Peterson,Phys.Rev.120,521(1960).
22. E.G.Bilpuch,private communication(1959).
23. J.Frehaut et al, Proc. 10-50 MeV Conf, BNL-NCS-51245 (1980)
page 399.
24. L.R.Veesser et al, Phys.Rev. C16, 1792(1977).
25. B.P.Bayhurst et al, Phys.Rev. C12, 451(1975).
26. R.J.Prestwood and B.P.Bayhurst,Phys.Rev.121,1438(1961).
27. C.Kalbach and F.Mann, BNL-NCS-5/245,p.689 (1980).
28. V.J.Orphan et al, GA-10248 (1970).
29. D.C.Larson, Proc. 10-50 MeV Conf, BNL-NCS-51245 (1980) p.277.
30. A.Carlson et al., Nuc.Data for Basic & Applied Science,
Santa Fe, NM (1985) p.1429.
31. W.Poenitz, ANL-West, personnal communication (1989).

DESCRIPTION OF EVALUATIONS FOR $^{206,207,208}\text{Pb}$ PERFORMED FOR ENDF/B-VI†

C. Y. Fu, D. C. Larson, and N. M. Larson
Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6356, U. S. A.

ABSTRACT

An evaluation of data for neutron induced reactions on $^{206,207,208}\text{Pb}$ was performed for ENDF/B-VI and is briefly described. The evaluation is based on experimental data guided by model calculations. Evaluated data are given for neutron induced reaction cross sections, angular and energy distributions of the secondary neutrons, recoil spectra, and gamma-ray production cross sections and spectra. File 6 formats are used to represent energy-angle correlated data for the outgoing neutrons. Uncertainty files are included for all File 3 cross sections. New data are available for (n,2n) cross sections and energy-angle correlated neutron emission spectra. Resonance parameters, absent from the previous evaluations, have been added. Serious energy imbalance problems in ENDF/B-V have been completely removed by using isotopic evaluations, by using calculated gamma-ray production spectra instead of adopting experimental data directly, and by using the File 6 formats.

1. INTRODUCTION

The previous major evaluation for natural lead was done in 1970-1971 for ENDF/B-III and documented in detail (FU75). The gamma-ray production cross sections and spectra were mainly based on model calculations because few data were available at the time. Later, a major measurement for the gamma-ray production cross sections and spectra for incident neutron energies from 0.6 to 20 MeV became available (CH75). It was believed these data were more reliable than the calculated results and were adopted for ENDF/B-IV. The adoption of measured gamma-ray production spectra in the evaluation, without first checking for its consistency with the particle emission spectra, could have caused an energy imbalance. However, the seriousness of this energy imbalance was not fully understood until ENDF/B-V was released and checked by MacFarlane (MA84). Therefore, a major goal of ENDF/B-VI was to solve this problem by making sure the evaluated neutron emission spectrum and gamma-ray production spectrum for each incident neutron energy,

†Research sponsored by the Office of Energy Research, Nuclear Physics, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

each reaction, and each isotope are consistent. The sum of the average energies of all reaction products, including the heavy recoils, now agrees within 1% of the incident neutron energy plus the Q-values of the reaction.

The improvements to neutron emission spectra, (n,2n) and (n,3n) cross sections resulted from the availability of new data (TA83, FR80) and from improvements to the model code TNG (FU88, SH86) used for the earlier evaluation. The new compilation of resonance parameters by Mughabghab (MU81) and the advances in the R-Matrix code SAMMY (LA89) greatly facilitated the evaluation of resonance parameters.

In Section 2 the resonance parameters are discussed; Section 3 contains a description of the major cross sections included in the evaluation; Section 4 is devoted to angular distributions; Section 5 to energy-angle correlated distributions; Section 6 to gamma-ray production cross sections and spectra; Section 7 describes the uncertainty files; and Section 8 describes important data needs and possible ways to improve the evaluation.

Part of this information is abstracted from FU75 and FU82.

2. RESONANCE PARAMETERS

Point cross sections were used in all previous ENDF/B evaluations for natural lead in the resonance energy region. For ENDF/B-VI resonance parameters were added for ^{206}Pb , ^{207}Pb , and ^{208}Pb and were mostly based on those compiled by Mughabghab (MU81) and re-fitted using SAMMY (LA89) to high-resolution isotopic data of Horen et al. for ^{206}Pb (HO81), ^{207}Pb (HO78), and ^{208}Pb (HO86). Other data shown in (MU81), including the thermal values, were also considered in the fits.

For $^{206,207}\text{Pb}$, the total cross section was fitted so well that no background was needed in File 3; there are negative entries for the elastic cross section in the resonance region because $^{206}\text{Pb}(n,n')$ and $^{207}\text{Pb}(n,\alpha)$ have contributions in this region. The total and elastic scattering cross sections in the resonance region for ^{208}Pb required a negative background.

3. CROSS SECTIONS

The total cross sections from the upper resonance energy to 20 MeV were taken from (HO81) for ^{206}Pb and (HO78) for ^{207}Pb . For ^{208}Pb , (HO86) was used only up to 2 MeV and from 2 to 20 MeV, the natural lead data evaluated before (FU75) were judged to be more reliable and were used. This is an area that requires improvement; see Section 8 for details.

The cross section values for the (n,n') continuum, (n,2n), and (n,3n) reactions were calculated by TNG but the (n,2n) cross sections were adjusted to agree with the shape of the available (n,2n) data (FR80). Figures 1 to 3 compare the isotopic data for the three isotopes (FR80), respectively, with the calculated and the adjusted values for ENDF/B-VI. The natural lead results, summed from the isotopic evaluations for ENDF/B-VI, are compared with the available data (FR80, IW86, TA86) in Fig. 4. It should be noted that the data of (FR75) shown in (FU82) are different from the data of (FR80) shown here in Fig. 4, the latter being smaller. The (n,2n) results of Takahashi et al. are likely too large,

approaching the evaluated nonelastic cross section at 14.1 MeV. Older (n,2n) data near 14 MeV are represented by the ENDF/B-V value, hence not shown.

The charged particle emission cross sections (n,p), (n,t) and (n, α) have been introduced for activation purposes. These cross sections are very small and were mostly based on TNG calculations.

The discrete inelastic cross sections were based on the new TNG calculations using the same parameters and direct interaction contributions as before (FU75), therefore changed little.

The capture cross sections above the resonance region were based on the new calculation and agree with a few data points (MC88). The new calculation for the capture cross sections has a direct/semi-direct component, resulting in a peak for the cross section near 14 MeV.

4. ANGULAR DISTRIBUTIONS

Angular distributions for the elastic and discrete inelastic cross sections were not changed (FU75), the latter having been separated for each isotopic file. The angular distributions for the (n,n') continuum, (n,2n), and (n,3n) reactions are correlated with the energy distributions and are described below.

5. ENERGY-ANGLE CORRELATED DISTRIBUTIONS (FILE 6)

Since lead is a likely neutron multiplier in fusion reactors, the energy and angular distributions for secondary neutrons must be carefully considered. For ENDF/B-V, the angular distribution of neutrons emitted from the continuum was assumed to be isotropic. However, experimental neutron emission data (KA72, HE75, TA83) shown in Fig. 5 exhibit strong anisotropic scattering and it is clear that the angular distribution is a function of the outgoing neutron energy. The major upgrade of the lead evaluation for version VI is to incorporate realistic energy-angle correlated distributions for neutrons emitted from the (n,n') continuum, (n,2n), and (n,3n) reactions.

Figure 6 compares the angle-integrated neutron emission data of Takahashi et al. (TA83) with the TNG result for $E_n = 14.1$ MeV. The data show a compound component at low outgoing energies with an abrupt change to a nearly flat component near the outgoing energy of 6 MeV. This flat component in the neutron spectra is believed to be due to a direct reaction that involves many high-lying discrete levels and is rather difficult to model. Reproduction of this effect was achieved by incorporating a constant level density in TNG whose excitation is tied to the lowest exciton component in the precompound stage. This technique gave a good fit to the angle integrated spectrum (see Figure 6) as well as angular distributions for the emitted neutrons in good agreement with measured data (see Figure 5). However, this method of simulating the continuum direct reaction has not been tested for other incident neutron energies. It might be expected to work for incident neutron energies lower than 14 MeV because the precompound component decreases with decreasing incident neutron energy and this direct component should get

less important. Anyway, it was used for the entire incident energy range where continuum neutrons are emitted. Further theoretical work is in progress to provide better modeling of this problem in TNG.

Correlated energy-angle distributions for the outgoing neutrons from the (n,n') continuum, $(n,2n)$, and $(n,3n)$ reactions are given in File 6, along with the corresponding recoil spectra and gamma-ray production spectra. The latter are described below.

6. GAMMA-RAY PRODUCTION CROSS SECTIONS AND SPECTRA

For all previous ENDF/B evaluations for natural lead, gamma-ray production cross sections and spectra were given for the nonelastic reaction, which is the sum of all reaction cross sections. This practice prevented a precise check of energy conservation. As in all other TNG-based evaluations for ENDF/B-VI ($^{50,52,53,54}\text{Cr}$, $^{54,56,57,58}\text{Fe}$, $^{58,60,61,62,64}\text{Ni}$, and $^{63,65}\text{Cu}$), gamma-ray production cross sections and spectra are given for each isotope and each reaction after checking against experimental data for the natural element. The new approach ensured energy conservation for each reaction, allowing adjustment of the calculated cross sections to experimental data without upsetting the energy balance. Calculated gamma-ray production spectra are compared in Figs. 7-9 with the data of Chapman and Morgan (CH75) for three incident neutron energies, respectively.

Capture gamma-ray spectra were obtained from the calculation. The new TNG code makes use of experimental s-wave branching ratios in the capture gamma-ray cascades, thus assuring a good fit to the capture spectra for thermal neutrons. The calculated results are represented in File 12 for the multiplicities and File 15 for the energy distributions. File 6 is not used because recoil energy from capture gamma-rays is very small and there is no known application in the case of lead. Even without the recoil spectra, energy conservation in the capture event is automatically guaranteed.

7. UNCERTAINTY INFORMATION

Uncertainty files are given only for cross sections in File 3, and not for the resonance parameters, energy distributions or angular distributions. Fractional and absolute components, correlated only within a given energy interval, are based on scatter in experimental data and estimates of uncertainties associated with the model calculations.

8. DATA NEEDS AND EVALUATION IMPROVEMENTS

Double differential neutron emission cross sections for each isotope are needed at incident energies lower than 14 MeV to benchmark the model calculations. Isotopic data are needed because the $(n,2n)$ thresholds of the three major isotopes are significantly different. The method for approximating the continuum direct component currently used in the TNG code (see Section 5) needs further development and testing. Note that this component is particularly large for the lead isotopes.

The total cross sections for ^{208}Pb from 2 to 20 MeV were taken from the ENDF/B-V evaluation for natural lead because the ^{208}Pb data (HO86) in this energy range were found to be poor. Upon reflection, these cross sections should have been taken as the difference between the natural lead data and the sum of the ^{206}Pb and ^{207}Pb data to avoid double counting of some of the resonances. However, new high resolution data for ^{208}Pb are now available, obtained as part of an effort to measure the polarizability of the neutron. These new data should be analyzed and incorporated in the ^{208}Pb evaluation.

An evaluation for ^{204}Pb should be made. Even though this isotope has a natural abundance of about 1%, the capture resonances in this isotope between 10 and 30 keV could contribute as much as 30% to the natural lead cross section in this energy range.

REFERENCES

- CA91 R. F. Carlton et al., "R-Matrix Analysis of an ORELA Measurement of the $n + ^{208}\text{Pb}$ Total Cross Section from 78 to 1700 keV," Submitted to April 1991 Meeting of the American Physical Society, Washington, D. C.
- CH75 G. T. Chapman and G. L. Morgan, "The $\text{Pb}(n, x\gamma)$ Reaction for Incident Neutron Energies Between 0.6 and 20 MeV," ORNL/TM-4822, Oak Ridge National Laboratory (1975).
- FR75 J. Frehaut and G. Mosinski, "Measurement of $(n, 2n)$ and $(n, 3n)$ Cross Sections at Incident Energies Between 8 and 15 MeV," 5th Int. Symp. on Interaction of Fast Neutrons with Nuclei, Gaussig, DDR, 1975.
- FR80 J. Frehaut, "Status of $(n, 2n)$ Cross Section Measurements at Bruyeres-Le-Chatel," p. 399 in Symposium on Neutron Cross Sections from 10 to 50 MeV, edited by M. R. Bhat and S. Pearlstein, Brookhaven National Laboratory (1980).
- FU75 C. Y. Fu, Atomic Data and Nucl. Data Tables 16, 409 (1975).
- FU82 C. Y. Fu, "Summary of ENDF/B-V Evaluations for Carbon, Calcium, Iron, Copper, and Lead and ENDF/B-V Revision 2 for Calcium and Iron," ORNL/TM-8283, ENDF-325 (1982).
- FU88 C. Y. Fu, Nucl. Sci. Eng. 100, 61 (1988).
- HE75 D. Hermsdorf et al., "Differential Neutron Emission Cross Sections by 14.6-MeV Neutrons," Institute of Nuclear Physics, University of Dresden (1975).
- HO78 D. J. Horen et al., Phys. Rev. C12, 722 (1978).
- HO81 D. J. Horen et al., Phys. Rev. C24, 1961 (1981).
- HO86 D. J. Horen et al., Phys. Rev. C34, 429 (1986).
- IW86 S. Iwasaki et al., Rad. Eff. 92, 191 (1986).
- KA72 J. L. Kammerdiener, "Neutron Spectra Emitted by Pb Irradiated by 14 MeV Neutrons," UCRL-51232, Lawrence Livermore Laboratory (1972).
- LA89 N. M. Larson and F. G. Perey, "User's Guide for SAMMY: A Computer Model for Multi-Level R-Matrix Fits to Neutron Data Using Bayes' Equations," ORNL/TM-7485 (1980); Updates ORNL/TM-9179 (1984), ORNL/TM-9179/R1 (1985) and /R2 (1989).
- MC88 V. McLane et al., Neutron Cross Section Curves, Academic Press, 1988.

- SH86 K. Shibata and C. Y. Fu, "Recent Improvements of the TNG Statistical Model Code," ORNL/TM-10093 (1986).
- TA83 A. Takahashi et al., "Double Differential Neutron Emission Cross Sections, Numerical Tables and Figures (1983)," A-83-01, Intense Neutron Source Facility, Osaka University, 1983.
- TA86 A. Takahashi, "Nuclear Data for Fusion Blanket Neutronics," p. 190 in Proc. 1985 Seminar on Nuclear Data, edited by T. Asami and M. Mizumoto, JAERI-M 86-080, Japan Atomic Energy Research Institute (1986). Also Rad. Eff. 92, 59 (1986).

Fig. 1. Comparison of experimental $^{206}\text{Pb}(n,2n)$ data (FR80) with ENDF/B-VI and the TNG calculation.

Fig. 2. Comparison of experimental $^{207}\text{Pb}(n,2n)$ data (FR80) with ENDF/B-VI and the TNG calculation.

Fig. 3. Comparison of experimental $^{208}\text{Pb}(n,2n)$ data (FR80) with ENDF/B-VI and the TNG calculation.

Fig. 4. Comparison of experimental (n,2n) data for natural lead (IW86,TA86,FR80) with ENDF/B-V and ENDF/B-VI, the latter obtained from summing the isotopic evaluations.

Fig. 5. Comparison of experimental Pb(n,xn) spectrum at 14.1 MeV as a function of angle for several secondary energy ranges (TA83, HE75, KA72) with results of the TNG calculation used for ENDF/B-VI.

Fig. 6. Comparison of angle-integrated neutron emission spectrum of Takahashi et al. (TA83) at 14.1 MeV for natural lead with ENDF/B-V and results of the TNG calculation used for ENDF/B-VI.

Fig. 7. Comparison of experimental gamma-ray production spectrum at $E_n = 4.25$ MeV for natural lead (CH75) with the TNG calculation used for ENDF/B-VI.

Fig. 8. Comparison of experimental gamma-ray production spectrum at $E_n = 9.5$ MeV for natural lead (CH75) with the TNG calculation used for ENDF/B-VI.

Fig. 9. Comparison of experimental gamma-ray production spectrum at $E_n = 14$ MeV for natural lead (CH75) with the TNG calculation used for ENDF/B-VI.

²⁰⁹₈₃Bi

Reference: ANL/NDM-109
Evaluators: A. Smith, D. Smith, P. Guenther, J. Meadows, R. Lawson
(ANL), R. Howerton (LLNL), and M. Sugimoto (JAERI)
Evaluated: August 1989
Material: 8325
Content: Neutron transport, Gamma production, Covariances

1. Introduction

The primary objective of this evaluation is a practical file for neutronic applications. The evaluation reasonably summarizes the contemporary physical knowledge.

2. Evaluated Resolved Resonance Range

The resolved resonance region was described by resonance parameters taken from Mughabghab¹ up to 0.1 MeV. The ENDF/B-V evaluation for bismuth has no resonance parameters for comparison.

3. Evaluated Total Cross Sections

Above 0.1 MeV the present evaluation uses a pointwise representation with detailed resonance fluctuations to 2 MeV. The energy averaged data base was evaluated using the statistical procedures of the code GMA.² Into the several MeV region, detailed partially resolved resonance structure has been reported, particularly in Refs. 3 and 4. Below 2 MeV the best resolution data appear to be from reference 3. The evaluation is qualitatively consistent with ENDF/B-V. The present work has far more detail at lower energies due to the introduction of new experimental information.

4. Evaluated Elastic Scattering Cross Sections

Below 0.1 MeV the evaluated cross sections follow from the resonance parameters. Above 0.1 MeV the evaluation is explicitly based upon Refs. 5 and 6. The present elastic scattering evaluation is in good agreement with ENDF/B-V.

5. Evaluated Inelastic Scattering Cross Sections

5.1 Discrete Inelastic Processes

The experimental data base for discrete inelastic scattering was assembled from the files of the NNDC. The evaluation used the potential of Ref. 5, nineteen excited levels (Ref. 7), and the optical statistical model. Above 3.1 MeV the calculation assumed a continuum of levels given by a modified Gilbert and Cameron⁸ statistical representation as defined in Ref. 5. At energies higher than 4 MeV the measurements of Ref. 9 were used and were significantly larger than those calculated. There was reasonable agreement between calculated and measured $(n,n\gamma)$ cross sections near thresholds. The present evaluation has far more detail than ENDF/B-V and contains 13 more excited states.

5.2 Continuum Inelastic Scattering Processes

The magnitude of the continuum inelastic scattering cross section was defined by the difference between the total cross section and the other partial cross sections. It is similar to that given in ENDF/B-V. The neutron spectra emitted as a result of continuum inelastic scattering was calculated using the methods given in ANL/NDM-105 (1988)¹⁰ and verified against the direct measurements of Ref. 11.

6. Evaluated Radiative Capture Cross Sections

Radiative capture measurements for bismuth are sparse. Moreover fluctuations lead to large variations in the experimental results below 200 keV. With the uncertain data base, calculations were relied upon. A simple dipole model given by Moldauer¹² normalized to the S_0 strength function given by Mughabghab¹ was used. The present evaluation is generally larger than that of ENDF/B-V by 25-50%.

7. Evaluated $(n,2n)$ and $(n,3n)$ Reactions

The experimental data base consists of individual measurements at 14 MeV and two comprehensive data sets.^{13,14} J. Frehaut¹¹ gives a good coverage from threshold to 14 MeV and L. Veaser¹³ above 14 MeV. The data base was supplemented with statistical calculations using the code CADE.¹⁵ The evaluation is consistent with the more precise of the isolated 14 MeV experimental results. The evaluated $(n,2n)$ cross sections are similar to ENDF/B-V.

There are only 2 data sets relevant to the $(n,3n)$ cross sections, a single value near threshold¹⁶, and five values from L. Veaser between 16 and 20 MeV. CADE results are somewhat lower in the 16-18 MeV region. The present evaluation is a compromise between experimental values of Ref. 13 and the CADE calculated results.

8. Evaluated Charged Particle Emitting Reactions

A number of reactions are energetically possible. See table 1. All the reactions are greatly inhibited by the coulomb barrier, and as a consequence the cross sections are small. Experimental information is sparse to non-existent, and the evaluation relies on model calculations using CADE.

Table 1

Q-values for Charged Particle Emitting Reactions

Reaction	Q-value (MeV)
(n,p)	+0.138
(n,np)	-3.798
(n,d)	-1.573
(n,nd)	-8.941
(n,t)	-2.685
(n,nt)	-9.424
(n, ^3He)	-4.087
(n,n ^3He)	-10.931
(n, α)	+9.648
(n,n α)	+3.144

8.1 (n,p) and (n,np) Reactions

CADE results were normalized to three consistent experimental sections. The cross sections of the present evaluation are approximately a factor of two smaller than ENDF/B-V near 14 MeV. No experimental information is available for the (n,np) reaction, thus the evaluation is based entirely upon the CADE calculations normalized by the same factor used for the (n,p) evaluation.

8.2 (n,d) and (n,nd) Reactions

The evaluations rely entirely upon the CADE calculations. The cross section behavior is consistent with experimental evidence by S. Quaim et al.¹⁷ at ≈ 22 MeV which suggests that the sum of the (n,d) and (n,nd) cross sections are less than 1 mb below 20 MeV. There are no comparable ENDF/B-V files.

8.3 (n,t) and (n,nt) Reactions

There is experimental evidence for a total tritium cross section at 22 MeV of less than a few mb.¹⁷ This is consistent with the CADE results which were used for the present evaluation.

8.4 (n, α) and (n,n α) Reactions

There are a few experimental results near 14 Mev with cross sections varying from 0.5 to 1.2 mb. These are relatively consistent with the CADE results. The calculated results were slightly renormalized to improve the agreement with these cross sections. There may be some pre-compound contributions which were not taken into account. The results are qualitatively similar to ENDF/B-V.

9. Evaluated Photon Production Reactions

The spectrum of neutrons from the capture reaction was taken from Orphan.¹⁸ For photons associated with inelastic scattering to specific levels the code CASCADE¹⁹ was used. For all other reactions the photon production cross sections and spectra were calculated using the R-parameter formalism of Perkins et al.²⁰

References

1. S. F. Mughabghab, Neutron Cross Sections Vol. 1, Part B, Academic Press Inc. New York, (1984); also S. Mughabghab and C. Dunford, private communication (1982).
2. W. P. Poenitz, Brookhaven National Laboratory Report, BNL-NCS-51363 Vol. I 249 (1981); as modified by M. Sugimoto (1987).
3. J. Harvey, private communication, data at NNDC (1985).
4. S. Cierjacks et al., Kernforschungszentrum-Karlsruhe report, KFK-1000 (1968).
5. A. Smith, P. Guenther, and R. Lawson, Argonne National Laboratory Report ANL/NDM-100 (1987).
6. R. D. Lawson, P. T. Guenther, and A. B. Smith, Phys. Rev. C36 1298 (1987).
7. C. Lederer and V. Shirley, eds., Table of Isotopes, 7th Edition, John Wiley and Sons Inc. New York (1978).
8. A. Gilbert and A. Cameron, Can. J. Phys. 43 1446 (1965).

9. S. Chiba and A. Smith, to be published.
10. A. Smith, D. Smith, P. Guenther, J. Meadows, R. Lawson, R. Howerton, T. Djemil, and B. Micklich, Argonne National Laboratory Report, ANL/NDM-105 (1988).
11. P. T. Guenther, to be published.
12. P. Moldauer, computer code ABAREX, private communication (1982).
13. L. Veaser et al., Proc. Inter. Conf. on the Interaction of Neutrons with Nuclei 2 1351 (1976).
14. J. Frehaut et al., Proc. Kiev Conf. (1975).
15. D. Wilmore, Harwell Report AERE-R-11515 (1984).
16. Lawrence Livermore National Laboratory Table of Q-values, available from one of the authors (RJH).
17. S. Qaim, R. Wolfe, and G. Stocklin, Nucl. Chem. 36 3639 (1974), and also Nucl. Phys. A295 150 (1978).
18. V. J. Orphan, N. C. Rasmussen, and T. L. Harper, "Line and Continuum γ -ray Yields from Thermal Neutron Capture in 75 Elements," Gulf General Atomic Report, GA-10248/DASA 2570 (1970).
19. W. E. Warren, R. J. Howerton, and G. Reffo, CASCADE Cray program for γ -production from discrete level inelastic scattering, Lawrence Livermore Nuclear Data Group Internal Report, PD-134 (1986), unpublished.
20. S. T. Perkins, R. C. Haight, and R. J. Howerton, Nucl. Sci. and Eng. 57 1 (1975).

²³⁵₉₂U

Reference: No Primary Reference
Evaluators: L. W. Weston, P. G. Young, W. P. Poenitz, Others
Evaluated: April 1989
Material: 9228
Content: Neutron transport, Gamma production, Covariances

File Comments

1. Principal Evaluators

Thermal parameters:

Standards Committee of CSEWG.

Resolved Resonance Region: (0-2250 eV)

U.TENN - L. C. Leal and R. B. Perez

ORNL - G. deSaussure, N. M. Larson, and R. Q. Wright

LANL - M. S. Moore

Unresolved Resonance Region and File 3 below 100 keV.

Capture Cross Sections above 100 keV:

ANL - W. P. Poenitz

ORNL - L. W. Weston

Fission Cross Section Above 100 keV:

Standards Committee of CSEWG.

ANL - W. P. Poenitz

Model Calculations and Fits above 100 keV:

LANL - P. G. Young, R. E. MacFarlane, and E. D. Arthur

Covariance Files:

ORNL - R. W. Peelle

2. Neutron Yields and Nubar

MF=1 MT=452 Total Nubar - sum of MT=455 and 456.
MT=455 Delayed Neutron Yields. England (En89).
MT=456 Prompt Neutron Yields. At neutron energies below 1 keV, taken from the evaluation of Frehaut (renormalized to match the the ²³⁵U thermal standard), which indicates a constant value. Above 1 keV, a new covariance analysis of all available experimental data was performed.

MF=1 MT=456 Continued. This analysis was performed using the GLUCS analysis code (He80). The data were obtained from the NNDC and were renormalized using the ENDF/B-VI standards. A smooth curve was drawn through the results of the covariance analysis.

3. Analysis of the Resonance Region.

3.1 Introduction

The ^{235}U neutron cross sections are described with Reich-Moore type resolved resonance parameters up to 2250 eV (20). The resolved range is divided into 10 regions, each described by its own set of resonance parameters as shown in the table below.

Subdivisions of the Resolved Resonance Region.

Energy region (eV)	Number of resonances
0- 110	236
110- 300	384
300- 500	272
500- 750	361
750-1000	303
1000-1250	368
1250-1500	334
1500-1750	302
1750-2000	280
2000-2250	502

The resonance parameters were obtained by fitting experimental data with the resonance analysis code SAMMY (1). The partial cross section measurements were all renormalized to the 2200 m/s values of the ENDF/B-VI standards committee (2). The energy scale of all the data sets were aligned on the energy scale of the 80 meter flight path transmission measurement of Harvey et al. (3). The length of that flight path has been measured with great accuracy (4).

3.2 Thermal Region

In the thermal region the following measurements were given most weight: the transmission measurement of R. R. Spencer et al. (5); the recent fission cross section measurements of R. Gwin et al. (6), of Schrack (7) and Wagemans et al. (8);

the ENDF/B-V capture cross section below 0.5 eV renormalized at 2200 m/s to the value proposed by the ENDF/B-VI standards committee (2). A value of 9.938 fm was obtained for the effective scattering radius from the consistent fit of these data. In the following table, the 2200 m/s values of the cross section computed with the parameters of this evaluation, using the Leal-Hwang method (9), are compared to the values proposed by the ENDF/B-VI standards committee:

2200 m/s Values of the Cross Sections (b).			
Cross section	** this evaluation **		ENDF/B-VI Standards
	At 0 K	At 300 K	
total	698.42	698.22	698.67 ± 1.71
scattering	15.48	15.52	15.46 ± 1.06
absorption	682.94	682.70	683.22 ± 1.34
fission	584.18	583.98	584.25 ± 1.11
capture	98.76	98.72	98.96 ± 0.74

3.3 The Energy Region 0-110 eV

In the resonance region up to 110 eV, the following measurements were given most weight: the transmission measurements of Harvey et al. (3); the fission measurements of Gwin et al. (6), of Schrack (7), of Wagemans and Deruytter (10) and of Weston and Todd (11); the spin-separated fission cross section data of Moore et al. (12) from the analysis of the polarized neutron polarized target measurement of Keyworth et al. (13). The following tables provide a comparison of the fission and capture cross sections integrated over several energy intervals as obtained from different data sets and from this evaluation:

Comparison of Fission Integrals (b ev).					
Interval (eV)	This Evaluation	Schrack (7)	Gwin + (6)	Wagemans + (10)	Weston + (11)
0.0206-0.06239	19.16	19.18	19.26	19.26 ± 0.08	
7.8-11.0	246.02	239.41	247.4	$246. \pm 2.5$	
0.5- 10.0	404.	397.	406.	406.	
10.0- 50.0	1805.	1796.	1838.5	1838.	
50.0-100.0	1582.	1586.	1632.	1647.5	1601.9
100.0-110.0	187.	186.	183.	190.6	188.

Comparison of Capture Integrals (b eV).

Range (eV)	This eval.	Desaussure et al. (14)	Perez et al. (15)
0.5- 10.0	224.9	231.6	(not measured)
10.0- 50.0	1162	1178	1252
50.0-100.0	685	721	747
100.0-110.0	152	158	176

3.4 The Energy Region 110-2250 eV

Above 110 eV, the resonances cannot be fully resolved. The resonance parameters provided in this evaluation represent well the high resolution thick sample transmission measurements of Harvey et al. (3) and the fission measurement of Weston and Todd (11, 18). The assignment of spins to the resonance structures is also roughly consistent with the spin-separated fission data of Moore et al. (12). In the following table, the fission cross sections averaged over decimal intervals between 100 and 2250 eV from this evaluation is compared with values from ENDF/B-V (16) and with values proposed by the ENDF/B-VI standards committee (2).

Comparison of Average Evaluated Fission Cross Sections (b).

Energy Interval (eV)	This Eval.	ENDF/B-V	Proposed Standard	Schrack (7)	Weston + (18)
100- 200	20.64	20.71	21.14 ± 0.09	20.91	20.48
200- 300	20.04	20.21	20.67 ± 0.10	20.05	20.15
300- 400	12.80	12.90	13.14 ± 0.07	13.21	12.86
400- 500	13.43	13.46	13.79 ± 0.07	13.83	13.53
500- 600	14.95	14.86	15.19 ± 0.08	14.63	14.77
600- 700	11.49	11.35	11.47 ± 0.06	11.46	11.27
700- 800	10.88	10.90	11.14 ± 0.06	10.79	10.79
800- 900	8.32	8.02	8.25 ± 0.04	7.82	7.95
900-1000	7.22	7.34	7.53 ± 0.04	7.18	7.30
1000-2000	7.12	7.20	7.35 ± 0.04		7.04

In the following table the fission and capture resonance integrals obtained from this evaluation are compared to the values obtained for ENDF/B-V and to the values evaluated from integral measurement (19). The few percent contributions above 2250 eV for this evaluation were estimated to be equal to the values obtained from ENDF/B-V.

The ratio of the capture resonance integral to the fission resonance integral, 0.477, is lower than the value 0.513 ± 0.015 obtained from direct measurements (17).

Comparison of Fission and Capture Resonance Integrals (b).

E (eV)	***** Fission *****		***** Capture *****	
	This eval.	ENDF/B-V	This eval.	ENDF/B-V
0.5- 5	85.06	85.27	25.50	24.66
5- 50	109.09	111.68	74.67	78.49
50- 110	25.36	25.74	11.36	11.87
110- 300	20.81	20.93	8.46	10.67
300- 500	6.70	6.73	2.50	2.95
500- 750	5.23	5.29	1.83	1.98
750- 1000	2.46	2.37	1.18	1.27
1000- 1250	1.94	1.83	0.94	0.85
1250- 1500	1.32	1.33	0.52	0.49
1500- 1750	0.95	0.99	0.53	0.37
1750- 2000	0.89	0.91	0.34	0.39
2000- 2250	0.60	0.65	0.37	0.30
2250 - 20 MeV	(18.20)	18.20	(4.68)	4.68
0.5 - 20 MeV	278.61	281.92	132.88	138.97
From reference 19.		275 ± 5		144 ± 6

3.5 References for Resolved Resonance Region

1. N. M. Larson, ORNL/TM-9719/R1, (1985); also N. M. Larson and F. G. Perey, "Resonance Parameter Analysis with SAMMY," Int. Conf. Nuclear Data for Science and Tech., May 30 - June 3, 1988, Mito, Japan.
2. A. Carlson et al., "Results of the ENDF/B-VI Standards Evaluation," Private Communication 31 Aug 1987.

3. J. A. Harvey et al., "High-resolution Neutron Transmission Measurements on ^{235}U , ^{238}U , and ^{239}Pu ," Inter. Conf. Nucl. Data for Science and Technology, May 30 - June 3, 1988, Mito, Japan.
4. D. C. Larson and N. M. Larson, ORNL/TM-9097 (1985).
5. R. R. Spencer et al. Nucl. Sci. Eng. 96, 318 (1987).
6. R. Gwin et al., Nucl. Sci. Eng. 88, 37 (1984).
7. R. C. Schrack, "Measurement of the ^{235}U (n,f) Reaction from Thermal to 1 keV," Inter. Conf. Nuclear Data for Science and Technology, May 30 - June 3, Mito, Japan.
8. C. Wagemans et al., "Subthermal Fission Cross section Measurements for ^{233}U , ^{235}U , and ^{239}Pu ," Inter. Conf. Nuclear Data for Science and Technology, May 30 - June 3, 1988, Mito, Japan.
9. L. C. Leal and R. N. Hwang, Trans. American Nuclear Society 55, 1-341 (1987).
10. C. Wagemans and A. J. Deruytter, p499 in Proc. of Nuclear Data for Basic and Applied Science, Santa Fe, New Mexico, May 13 - 17, 1985 Vol. 1 (1986).
11. L. W. Weston and J. H. Todd, Nucl. Sci. Eng. 88, 567 (1984).
12. M. S. Moore et al., Phys. Rev. C18, 1328 (1978).
13. G. A. Keyworth et al., Phys. Rev. Letters 31, 1077 (1973).
14. G. deSaussure et al., ORNL/TM-1804 (1967).
15. R. B. Perez et al., Nucl. Sci. Eng. 53, 46 (1973).
16. M. R. Bhat, BNL-NCS-51184 (ENDF-248) (1980).
17. J. Hardy Jr., " ^{235}U Resonance Fission Integral and Alpha Based on Integral Measurements," ENDF-300, Section B.1 (1979).
18. L. W. Weston and J. H. Todd, Private Communication (1988). These measurements done on a flight path of 80 m. have a much better resolution than those of ref. 11 and have been used above 110 eV.
19. S. F. Mughabghab, "Neutron Cross Sections," Vol. 1, Part B (1984).
20. L. C. Leal, "Resonance Analysis and Evaluation of the ^{235}U Neutron Induced Cross Sections," ORNL/TM-11547 (1990).

3.6 Unresolved Resonance Region Analysis

The unresolved resonance region was derived by a FITACS (code developed by Fritz Froehner) fit by L. W. Weston to the standards committee recommendation for the fission cross section and new capture evaluation based on newer alpha measurements (see ANL-83-4 supplement). These results were then fit with URES (code developed by Ed Pennington) so ENDF would reproduce the cross sections. The unresolved resonance region extends from 2.25 to 25 keV and is to be used only for self shielding calculations. Dilute cross sections are taken from File 3 which shows experimentally observed structure carried over from ENDF/B-V up to 100 keV.

4.0 Remaining File Evaluations

4.1 Smooth Cross Sections MF=3

P. G. Young, R. E. MacFarlane and E. D. Arthur (LANL) performed model calculations in support of ENDF/B-VI. The evaluation above 100 keV is based on a detailed theoretical analysis utilizing the available experimental data. Coupled channel optical model calculations with the ECIS code (Ra70) were used to provide the total, elastic, and inelastic cross sections to the first 3 members of the ground state rotational band, as well as neutron elastic and inelastic angular distributions to the rotational levels. The ECIS code was also used to calculate neutron transmission coefficients. Hauser-Feshbach statistical theory calculations were carried out with the GNASH (Ar88, Yo77) and COMNUC (Du70) code systems, including preequilibrium and fission. DWBA calculations were performed with the DWUCK code (Ku70) for several vibrational levels, using $B(E\ell)$ values inferred from (d, d') data on ^{231}U , ^{235}U , ^{238}U , as well as coulomb excitation measurements. A weak coupling model (Pe69) was used to apply the ^{231}U and ^{238}U results to states in ^{235}U . A preliminary description of the analysis was given at the Mito conference (Yo88).

MT=1 Sum of partial cross sections from 2.25 to 100 keV. In the range 0.10 to 20 MeV, obtained from a covariance analysis of available experimental data, using an initial or prior cross section from the coupled-channel optical model analysis. Experimental data used include Fo71, Ve80, Bo71, Po81, Gr73, Sc74, Po83, Pe60, Wh65, Ca73, Ga60, and Br58. The GLUCS code was used for analysis (He80).

MT=2 Unchanged from version 5 from 2.25 to 100 keV. From 0.12 to 20 MeV, based on subtraction of MT=4, 16, 17, 18, 37, and 102 from MT=1.

- MT=4 Sum of MT =51 ... 91.
- MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calculation.
- MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calculation.
- MT=18 2.2 to 120 keV average values from simultaneous evaluation of W. P. Poenitz with structure carried over from version 5. The renormalizaion was over 3 ranges per decade. From 0.10 to 20 MeV the standards evaluation by the CSEWG standards committee was used.
- MT=19 (n,f) first-chance fission cross section. Ratio of first-chance to total fission obtained from GNASH calculations.
- MT=20 (n,nf) second-chance fission cross section. Ratio of second-chance to total fission obtained from GNASH calculations.
- MT=21 (n,2nf) third-chance fission cross section. Ratio of third-chance to total fission obtained from GNASH calculations.
- MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calculation.
- MT=38 (n,3nf) fourth-chance fission cross section.
- MT=51, 52,54,55,57,58,60-62,64-72,74-76 Threshold to 6.0 MeV, compound-nucleus reaction theory calculations with width fluctuation corrections using the COMNUC code.
- MT=53, 50,59,63 Threshold to 20 MeV, coupled-channel optical model calculations ($9/2^-$, $11/2^-$, $13/2^-$, $15/2^-$ members of the ground-state rotational band), plus compound-nucleus contributions from COMNUC calculations.
- MT=73, 77-84 Threshold to 20 MeV, distorted wave born approximation calculations with the DWUCK code. These levels are composites of $\ell=2$ and $\ell=3$ vibrational states. The $\ell=2$ states are MT=78-80, 82 and the $\ell=3$ states are MT=73, 77, 81, 83 and 84.
- MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calculation. Note that the MT=77-84 vibrational states lie in the MT=91 continuum region.

MT=102 2.25 to 1000 keV capture is based on newer alpha measurements (see ANL-83-4 supplement) and fission from standards committee recommendations. Most of the structure below 100 keV is from fission. Between one and 20 MeV it is based on a renormalized COMNUC/GNASH calculation, with a semi-direct component added above a few MeV.

4.2 Neutron Angular Distributions MF=4

- MT=2 Elastic scattering angular distributions, are based on ECIS coupled-channel calculations, with a compound elastic component from COMNUC included below 6 MeV.
- MT=51, 52,54,55,57,58,60-62,64-72,74-76 Threshold to 6.0 MeV, compound-nucleus reaction theory calculations with width fluctuation corrections using the COMNUC code.
- MT=53, 56,59,63 Threshold to 20 MeV, coupled-channel optical model calculations ($9/2^-$, $11/2^-$, $13/2^-$, $15/2^-$ members of the ground-state rotational band), plus compound-nucleus contributions from COMNUC calculations.
- MT=73, 77-84 Threshold to 20 MeV, distorted wave Born approximation calculations using the DWUCK code.

4.3 Neutron Energy Distributions MF=5

- MT=18 Composite neutron energy distributions from fission. Based on calculations by D. Madland (Ma88) using the Madland-Nix formalism. The calculations include the first-, second-, and third-chance fission neutron components. The calculations end at 15 MeV; the 20-MeV spectrum is simply a duplication of the 15-MeV spectrum. A tabulated distribution law (LF=1) is used.
- MT=455 From a study by T. England (En89).

4.4 Correlated Energy-Angle Distributions MF=6

- MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calculation. Neutron distributions only.
- MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calculation. Neutron distributions only.

MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calculation.
Neutron distributions only.

MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calculation.
Neutron distributions only.

4.4 References

- Ar88 E. D. Arthur, LA-UR-88-382 (1988).
- Bo72 K. Boeckhoff et al., J. Nucl. En. 26, 91 (1972).
- Br58 A. Bratenahl et al., Phys. Rev. 110, 927 (1958).
- Ca73 J. Cabe et al., CEA-R-4524 (1973).
- Du70 C. L. Dunford, AI-AEC-12931 (1970).
- En89 T. R. England et al., LA-11151-MS(88), LA-11534-T(89); LAURA-88-4118,
and M. C. Brady & T. R. England, Nucl. Sci. & Eng. 103, 129 (1989).
- Fo71 D. Foster & D. Glasgow, Phys. Rev. C3, 576 (1971).
- Ku70 P. D. Kunz, DWUCK: A Distorted-Wave Born Approximation Program,
Unpublished Report.
- Fr86 J. Frehaut, NEANDC(E)-238/1 (1986).
- Gr73 L. Green et al., USNDC-9 (1973) p.170.
- He80 D. Hetrick & C. Y. Fu, ORNL/TM-7341 (1980).
- Pe60 J. Peterson et al., Phys. Rev. 120, 521 (1960).
- Pe69 R. J. Peterson, Ann. Phys. 53, 40 (1969).
- Po81 W. Poenitz et al., Nucl. Sci. Eng. 78, 333 (1981).
- Po83 W. Poenitz et al., ANL-NDM-80, 1983.
- Ra70 J. Raynal, IAEA SMR-9/8 (1970).
- Sc74 R. Schwartz et al., Nucl. Sci. Eng. 54, 322 (1974).
- Ut66 C. Uttley et al., Paris Conf. (1966) V-1, p165.
- Ve80 V. Vertebnyj et al., YFI 16, 8 (1973).
- Wh65 W. Whalen et al., ANL-7110 (1965) p.15.

Yo77 P. G. Young & E. D. Arthur, LA-6947 (1977).

Yo88 P. G. Young & E. D. Arthur, Nucl. Data for Science and Technology, Mito
Conference (1988) p.603.

Fig. 1 ETA for ^{235}U below 1 eV. Solid line is ENDF/B-VI calculated from resonance parameters, dashed line is ENDF/B-V and the points are preliminary Geel data by H. Weigman. The evaluation was influenced by the preliminary measurements of M. Moxon of Harwell as well as previous measurements.

Fig. 2. Comparison of the transmission data of Harvey et al.³ with calculations using the resonance parameters. The two upper curves have been displaced upward by 0.5 for clarity.

Fig. 3 Comparison of the fission cross-section data of Blons (J. Blons, Nucl. Sci. Eng. 51, 130 (1973)) and of Weston and Todd¹¹ with calculations using the resonance parameters.

Fig. 4 Dilute fission cross section (line) for ^{235}U from 2 to 100 keV as compared to the ENDF Standards Committee recommendation (points). Structure is from ENDF/B-V but renormalized to Standards Committee recommendation.

Fig. 5 Alpha, the ratio of capture to fission, for ENDF/B-VI (line) as compared to recent measurements. Circles are the data of Corvi (F. Corvi et al., ANL-83-4(1982)) and the triangles are the data of Muradyan (G. V. Muradyan et al., Nuclear Cross Sections for Technology, Knoxville (1979)).

Fig. 6 The ENDF/B-VI total cross section for ^{235}U (line) compared to recent measurements.

Fig. 7 The ^{235}U (n,xn) cross sections (lines) as compared to ENDF/B-V and measurements.

Fig. 8 Nubar for ^{235}U compared with ENDF/B-V and measurements.

Fig. 9 The ratio of the fission cross section evaluation of ^{235}U for ENDF/B-VI to that for ENDF/B-V. The ranges over which the structure in ENDF/B-V was renormalized for ENDF/B-VI to agree with the simultaneous fit by Poenitz are obvious below 100 keV.

Reference: No Primary Reference
Evaluators: F. M. Mann and R. E. Schenter
Evaluated: October 1989
Material: 9231
Content: Neutron transport

File Comments

WHC Eval-Oct89 F. M. Mann & R. E. Schenter (File 2 +)
BNL Eval-Jul78 M. Divadeenam (ν)
HEDL Eval-Apr78 F. M. Mann & R. E. Schenter (Fast (n,f))

MF=1 General Information.

MT=452 Conde + Holmberg's ν -bar. Data was renormalized to ²⁵²Cf with $\nu = 3.75$.

MT=455 M. C. Brady & T. R. England, Nucl. Sci. & Eng. 103, 129 (1989). Also MF=5.

MT=458 Energy from fission based on Sher (Ref. 1).

MF=2 Resonance parameters.

MT=151 Resolved parameters. The resolved resonance region (10^{-5} eV to 1.5 keV) has been reevaluated using the new data of Macklin and Alexander (Ref. 2). As their data begin at 20 eV, the first resonances from the evaluation of Mughabghab (Ref. 3) (-9.7 and 5.45 eV) were used. At higher energies, the values of $g\Gamma_n \Gamma_a / \Gamma_t$ from Macklin and Alexander were combined with resonance parameters evaluated by Mughabghab. For ²³⁶U, Mughabghab's evaluation is based on the measurements of Carraro and Brusegan (Ref. 4), Mewissen et al. (Ref. 5) and Carlson et al. (Ref. 6). The procedure used for each resonance was to use the $g\Gamma_n$ from Mughabghab to infer a Γ_a using Macklin's area data and to infer a $g\Gamma_n$ if a Γ_a was present in Mughabghab's evaluation.

- MF=2 MT=151 Continued. The resonance energies and fission widths (which are quite small) were taken from Mughabghab's evaluation. In those cases where Macklin could directly infer $g\Gamma_n$ or Γ_a , these values were used. The resonance parameters inferred from Macklin's data were then weighted with Mughabghab's values to achieve the final set. Average parameters are given in table 1. Because of missing s-wave resonances, the resolved resonance region was stopped at 1.5 keV. Values of ℓ for each resonance were obtained by using a Bayesian analysis. Very few p-wave resonances were seen. The thermal capture and fission cross sections as calculated from these parameters are 5.13 and 0.047 barns as compared to Mughabghab's evaluation of 5.11 ± 0.21 and 0.07 barns. The corresponding resonance integrals are 338 and 7.77 barns, compared to 360 ± 15 and 7.8 ± 1.6 barns.
- MT=151 Unresolved parameters. The unresolved resonance region (1.5 keV to 100.0 keV) is described by average resonance parameters (s-wave) obtained from the resolved resonance region evaluation and from adjustments to fit (p and d waves) of recent absorption and capture measurements (Ref. 2, 7-9). The fitting procedure used the FERRET data analysis code (Ref. 10). A competitive width was included which was taken to have the same energy dependence as ^{238}U in ENDF/B-V. Table 2 gives the final set of unresolved parameters.
- MF=3 Smooth cross sections (100 keV - 20.0 MeV)
- MT=1 The total cross section was obtained from Klepatskij et al. (Ref. 20).
- MT=2 The elastic cross section was obtained by subtracting inelastic, capture, etc. cross sections from the total.
- MT=4 Inelastic from ^{238}U from version V, Ref. 13.
- MT=16 The (n,2n) is from Ref. 11 and 12, and the Q-value, Ref. 14.
- MT=17 The (n,3n) is from Ref. 11 and 12, and the Q-value, Ref. 15.
- MT=18 Fission. Above 100 keV the data of Behrens et al. (Ref. 16 and 17) was used, normalized to ^{235}U (n,f) of ENDF/B-VI.
- MT=19 Same as MT=18 below (n,nf) threshold, thereafter constant.

Table 1. Average Resolved Resonance Parameters for ^{238}U

Description	82 s Wave Resonances	35 p Wave Resonances
Average total width	0.070 eV	0.022 eV
Average reduced neutron width	0.001967 eV	0.0130 eV
Average gamma width	0.0191 eV	0.0202 eV
Average fission width	0.00034 eV	0.00035 eV
Average level spacing	18.36 eV	42.09 eV
Strength function	0.0001085	0.0001063

Table 2. Unresolved Resonance Parameters for ^{238}U

ℓ	J	ν_x	ν_n	ν_g	ν_f	D	Γ_n^0	Γ_x	Γ_f
0	.5	2.0	1.0	0.0	1.0	18.36	.1992-2	.191-1	.34-3
1	.5	1.0	1.0	0.0	1.0	18.36	.4957-2	.240-1	.34-3
1	1.5	2.0	1.0	0.0	1.0	9.18	.2479-2	.240-1	.34-3
2	1.5	1.0	1.0	0.0	1.0	9.18	.1836-2	.190-1	.34-3
2	2.5	2.0	1.0	0.0	1.0	6.12	.1224-2	.190-1	.34-3

Table 2a Γ_x Parameters Versus Energy
(All ℓ and J Values)

Energy eV	x
1500 - 45000	0.0
50000	0.203-4
60000	0.333-3
70000	0.120-2
80000	0.276-2
90000	0.512-2
100000	0.837-2

MF=3	MT=20	MT=20 is the difference of MT=18 and MT=19 until the (n,2nf) threshold, thereafter a constant.
	MT=21	MT=21 is the difference of MT=18 and MT=19 and 20.
	MT=51,	52... 91 From ^{238}U version V (Ref. 13).
	MT=102	Neutron capture. Above the unresolved energy region (100 keV to 20 MeV) multigroup capture cross section values were obtained by subtracting the previous ENDF/B-V fission cross section from the total absorption cross section of Macklin and Alexander. A smooth cross section curve was obtained by combining the Macklin and Alexander results with recent Russian data (Ref. 7, 8, and 8a) and the older Barry (Ref. 9a) measurements. The combination was made using the FERRET code. The shape of the capture curve above 1.0 MeV was assumed to be nearly the same as ^{238}U
MF=4		Angular distributions.
	MT=2	The differential elastic cross section is the same as for ^{232}Th , (Ref. 18).
MF=5		Energy distributions.
	MT=16	The (n,2n) energy distribution is described by a Maxwellian.
	MT=17	The (n,3n) energy distribution is described by a Maxwellian.
	MT=18,	19, 20 For these MT's the neutron energy distribution is given by a simple fission spectrum plus a Maxwellian.
	MT=91	An evaporation temperature was taken from Gilbert and Cameron (Ref. 19).

References

1. R. Sher and C. Beck, EPRI NP-1771 plus revision 1/83, and Personal Communication to B. A. Magurno 2/83.
2. R. L. Macklin and C. W. Alexander, Nucl. Sci. & Eng. 104, No. 3, p.258 (1990).
3. S. F. Mughabghab, BNL-325, Vol. 1 (1984).
4. G. Carraro and A. Brusegan, Nucl. Phys., A257, 333 (1976).
5. L. Mewissen et al., NBS Special Publ. 425, p.729 (1975).

6. A. D. Carlson et al., Nucl. Phys., A141, 577 (1976).
7. L. E. Kazakov et al., Jadernye Konstanty, 2, 44 (1985).
8. A. A. Bergmann et al., Atomnaya Energiya, 52, 406 (1982); also Soviet Atomic Energy, 52, 403 (1982).
- 8a. A. N. Davletshim et al., Atomnaya Energiya, 58, 183 (1985); also Soviet Atomic Energy, 58, 216 (1985).
9. A. D. Carlson et al., Nucl. Phys., A141, 577-591 (1970).
- 9a. J. F. Barry, Proc. Phys. Soc. 78, 801 (1961).
10. F. Schmittroth, Nucl. Sci. & Eng., 72, 19-34 (1979).
11. M. K. Drake and A. N. Nichols, GA-8135 (1967).
12. Parker, AWRE-O-30/64 (1964).
13. W. Poenitz et al., ANL/NDM-32 (1977).
14. Maples et al., UCRL-16964 (1966).
15. R. J. Howerton et al., UCRL-14000 (1964).
16. J. W. Behrens, G. W. Carlson, and R. W. Bauer, Nucl. Cross Sections and Technology, NBS-425 p.591 (1975).
17. F. M. Mann and R. E. Schenter, Trans. Amer. Nucl. Soc. 23 546 (1976).
18. M. K. Drake and A. N. Nichols, GA-6404 (1966)
19. A. Gilbert and A. G. W. Cameron, Can. J. Phys. 43 1446 (1965).
20. A. B. Klepatskij et al., INDC(CCP)-295/1 (1989).

U-236 TOTAL CROSS SECTION

U-236 CAPTURE CROSS SECTION

U-236 CAPTURE CROSS SECTION

U-236 INELASTIC CROSS SECTION

U-236 FISSION CROSS SECTION

²³⁸₉₂U

Reference: No Primary Reference
Evaluators: L. W. Weston, P. G. Young, R. E. MacFarlane,
W. Poenitz, Others
Evaluated: November 1989
Material: 9237
Content: Neutron transport, Gamma production, Covariances

1. Summary of ENDF/B-VI Evaluation

1.1 Principal Evaluators

Resolved Resonance Region: (1.0×10^{-5} to 10,000 eV)

ORNL – G. DeSaussure, D. K. Olsen, R. Macklin

HARWELL – M. C. Moxon, M. G. Sowerby

Unresolved Resonance Region: (10 TO 149 keV)

KFK – F. H. Froehner

ANL – W. P. Poenitz

Reactions Above 149 keV:

Model Code Calculations: (LANL) – P. G. Young and R. E. MacFarlane

Fission: (ANL) – W. P. Poenitz

Nu-bar Delayed and Delayed Neutron Spectra:

Kaiser and Carpenter (Ka78)

Gamma Production Files:

LLNL – R. J. Howerton

Uncertainty Files:

ORNL – L. W. Weston

1.2 Evaluation Above the Unresolved Resonance Region

Above the unresolved resonance region, new evaluations were performed of the neutron total, (n,2n), (n,3n), (n,4n), (n,f), (n,ni), (n,2nf), (n,3nf), and (n, γ) cross sections as well as prompt nubar. The elastic and inelastic data from ENDF/B-V were carried over for Version VI.

To provide the new data, coupled channel optical model calculations were performed with the ECIS code (Ra70) for the lowest 3 members of the ²³⁸U ground state rotational band. These calculations were used to provide initial (prior) values for a covariance analysis of the total cross section and to provide neutron transmission coefficients for nuclear reaction theory calculations with the GNASH (Ar88, Yo77) and

COMNUC (Du70) Hauser-Feshbach statistical/fission/preequilibrium codes. These theory calculations were used to provide the MF=6 neutron distributions from the (n,2n), (n,3n), and (n,4n) reactions as well as prior values for covariance analyses of the cross sections for those reactions. Additionally, the above analyses plus DWBA calculations were used to check the ENDF/B-V evaluation of elastic and inelastic scattering. While some differences were found, the earlier work was generally found to be reliable, and it was decided to carry over the ENDF/B-V data because of the effort taken to match experimental data, both at lower energies and at 14 MeV.

1.3 File Entries

MF=1	Descriptive and Nubar Information
MT=452	Total Nubar. Sum of MT=455 and 456.
MT=455	Delayed Neutron Yields. Kaiser and Carpenter (Ka78).
MT=456	Prompt Neutron Yields. Taken from Frehaut (Fr86).
MF=2	Resonance Region
MT=151	<p>The resolved resonance region determines the thermal cross sections. The thermal capture is that adopted by the Standards Committee of CSEWG (2.7081 with an uncertainty of 0.0095 barns). The thermal (2200 m/s) constants are:</p> <p style="padding-left: 40px;">Total Cross Section: 12.068 barns</p> <p style="padding-left: 40px;">Capture Cross Section: 2.709 barns</p> <p>The resolved resonance region has been extended to 10 keV. The evaluation was from the work of D. K. Olsen (ORNL), G. deSaussure (ORNL), Roger Macklin (ORNL), M. C. Moxon (Harwell), and M. G. Sowerby (Harwell). In particular see Gd78 and Ol86.</p> <p>The unresolved resonance region extends from 10 to 149 keV and was evaluated by F. H. Froehner (KFK) with his code FITACS which does simultaneous fits to the experimental data (Fr89). The unresolved resonance region is to be used only for self shielding calculations and File 3 should be used to determine the dilute cross sections.</p>
MF=3	Smooth Cross Sections.
MT=1	Neutron Total Cross Sections. From 10 to 149 keV the evaluation is based on a FITACS fit to the experimental data made by F. H. Froehner (KFK).

- MF=3 MT=1 Smooth Cross Sections, Continued. From 0.25 to 20 MeV, the evaluation is based on a covariance analysis of the available experimental data, especially Fo71, Sc74, Po81, Po83, Ha73, Br58, Ca73, Pe60, Wh71, Ba65, Ut66, Sh79, and Li79. Similar data exclusions as used in the covariance analysis by Smith et al. (Sm82) were incorporated but additional data were included. A prior cross section for the analysis was taken from the coupled-channel analysis, described above. The GLUCS code system was used for the covariance analysis (He80).
- MT=16 The (n,2n) Cross Section. From threshold to 20 MeV the (n,2n) cross section is based on a covariance analysis of the available data (Ba66, Fr80a, Ko80, Yo78, Ma69, Ma72, Ka79, La73, Ro57, Ph56, Pe61, Ve78, An58, and Ry80), similar to MT=1. The prior cross section was obtained from a GNASH analysis, with approximately 20% uncertainty.
- MT=17 The (n,3n) Cross Section. From threshold to 20 MeV, the (n,3n) cross section is based on a covariance analysis similar to MT=16. Experimental data used in the analysis were Fr80b, Wh62, Ro57, Ma69, Al61, and Ve78.
- MT=18 Fission Cross Section (total). In the energy range 0.01 TO 0.3 MeV the evaluation was taken from Di80 and Sl77 by L. W. Weston, and from 0.3 to 20 MeV, the evaluation was taken directly from the simultaneous standards analysis (Ca85, Po89).
- MT=19 First-Chance Fission (n,f) Cross Section. From 0.3 to 20 MeV, the evaluation is based on the ratio of the first-chance to the total fission from ENDF/B-V using the present MT=18 sigma.
- MT=20 Second-Chance Fission (n,nf) Cross Section. From 5.0 to 20 MeV, the evaluation is based on the ratio of second-chance to total fission from ENDF/B-V using the present MT=18 sigma.
- MT=21 Third-Chance Fission (n,2nf) Cross Section. From 12 to 20 MeV, the evaluation is based on the ratio of third-chance to total fission from ENDF/B-V using the present MT=18 sigma.
- MT=37 The (n,4n) Cross Section. From threshold to 20 MeV the evaluation is taken directly from the GNASH analysis described above.

- MF=3 MT=38 Fourth-Chance Fission (n,3nf) Cross Section. From 18 to 20 MeV the evaluation is based on the ratio of fourth-chance to total fission from ENDF/B-V using the present MT=18 sigma.
- MT=102 Radiative Capture Cross Section. From 10 to 149 keV the evaluation by F. H. Froehner (KFK) was used. The evaluation is based on a FITACS fit to the experimental data. From 0.15 to 20 MeV the evaluation is taken directly from the simultaneous standards analysis (Ca85, Po89).
- MF=6 Correlated Energy-angle Distributions.
- MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calculations. Neutron distributions only.
- MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calculations. Neutron distributions only.
- MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calculations. Neutron distributions only.

1.4 References

- Al61 K. Allen et al., J. Nuc. En. 14, 100 (1961).
- An58 G. Antropov et al., A. E. 5, 456 (1958).
- Ar88 E. D. Arthur, LA-UR-88-382 (1988).
- Ba66 D. Barr, (LANL) Personal Communication to R. Howerton (1966).
- Ba65 R. Batchelor et al., Nuc. Phys. 65, 236 (1965).
- Br58 A. Bratenahl et al., Phys. Rev. 110, 927 (1958).
- Ca73 J. Cabe et al., CEA-R-4524 (1973).
- Ca85 A. Carlson et al., Nuc. Data for Basic & Applied Science, Santa Fe, NM (1985) p.1429.
- DI80 F. C. Difillippo et al., PR/C 21, 1400.
- Du70 C. L. Dunford, AI-AEC-12931 (1970).
- Fo71 D. Foster & D. Glasgow, Phys. Rev. C3, 576 (1971).
- Fr80a J. Frehaut et al., Nucl. Sci. Eng. 74, 29 (1980).
- Fr80b J. Frehaut et al., BNL-NCS-512457 (1980) p399.

- Fr86 J. Frehaut, NEANDC(E) 238/L (1986).
- Fr89 F. H. Frohner et al., "The Unresolved Resonance Range of ^{238}U ," Nucl. Sci. Eng. 103, 119 (1989).
- Gd78 G. deSaussure et al., "Calculation of the ^{238}U Neutron Cross Sections for Incident Neutron Energies up to 4 keV," ORNL/TM-6152, ENDF-257 (1978).
- Ha73 S. Hayes et al., Nucl. Sci. Eng. 50, 243 (1973).
- He80 D. Hetrick & C. Y. Fu, ORNL/TM-7341 (1980).
- Ka78 R. Kaiser & S. Carpenter (ANL-West) Personal Communication.
- Ko80 N. Kornilov et al., ZFK-410 (1980) p68.
- La73 J. Landrum et al., Phys. Rev. C8, 1938 (1973).
- Li79 P. Lisowski et al., (LANL WNR measurement) Personal Communication (1979).
- Ma69 D. Mather, AWRE-O-47 (1969).
- Ma72 D. Mather et al., AWRE-O-72 (1972).
- Ol86 D. K. Olsen, "Resolved Resonance Parameters for ^{238}U from 1 to 10 keV," Nucl. Sci. Eng. 94, 102 (1986).
- Pe60 J. Peterson et al., Phys. Rev. 120, 521 (1960).
- Pe61 J. Perkin, J. Nuc. En. 14, 69 (1961).
- Ph56 J. Phillips, AERE-NP/R-2033 (1956).
- Po81 W. Poenitz et al., Nucl. Sci. Eng. 78, 333 (1981).
- Po83 W. Poenitz et al., ANL-NDM-80, 1983.
- Po89 W. Poenitz, (ANL-West) Personal Communication (1989).
- Ra70 J. Raynal, IAEA SMR-9/8 (1970).
- Ro57 L. Rosen et al., LA-2111 (1958).
- Ry80 T. Ryves, J. Phys. G 6, 771 (1980).
- Sc74 R. Schwartz et al., Nucl. Sci. Eng. 54, 322 (1974).
- Sh78 R. Shamu et al., Personal Communication, 1978.
- SL77 R. Slovacek et al., Nucl. Sci. Eng. 62, 455.

- Sm82 A. Smith et al., ANL/NDM-74 (1982).
- Ut66 C. Uttley et al., Paris Conf. (1966) Vol 1, p165.
- Ve78 L. Veesser & E. Arthur, Harwell Nuclear Data Conference (1978) p1054.
- Wh62 P. White et al., J. Nucl. En. A/B 16, 261 (1962).
- Wh71 J. Whalen et al., ANL-7710 (1971) p9.
- Yo77 P. G. Young & E. D. Arthur, LA-6947 (1977).
- Yo78 Chou You-Pu, HSJ-77091 (1978).

2. Description of Carryover from ENDF/B-V

2.1 ENDF/B-V Evaluators

Principal Evaluators:

ANL - E. M. Pennington, A. B. Smith, W. P. Poenitz

Gamma Production Files:

LLNL - R. J. Howerton

Nu-bar Delayed and Delayed Neutron Spectra:

ANL-West - R. Kaiser and S. Carpenter

2.2 Carryover File Description

- MF=3 Smooth Cross Sections. The cross sections above 45 keV were evaluated by A. Smith. More details are provided under each reaction type below. Most of the evaluation above 45 keV is described in ANL/NDM-32.
- MT=2 Elastic Scattering Cross Section. Elastic scattering cross sections and angular distributions were evaluated using data referenced in ANL/NDM-32. Coupled-channel optical model calculations assisted in the analysis (Ref. 9). Below about 1 MeV, the elastic cross section was determined for consistency with the total and partial non-elastics. Above 1 MeV, the total and elastic cross sections determine the non-elastic. The angular distributions are entirely experimental below 1.8 MeV, and calculated at higher energies. At most energies the ENDF/B-V elastic cross section is smaller than ENDF/B-IV.

MF=3 MT=4, 51... 77, and 91. Total inelastic, Inelastic to 27 levels, and Continuum Inelastic Cross Sections. The evaluation is a correlation of theory (Ref. 9) and experiment. Experimental data referenced in ANL/NDM-32 are considered. Levels at higher energies are composites of actual levels. The total inelastic cross section is generally much higher than in ENDF/B-IV, except near threshold and over a region below 1 MeV. The individual levels have tails extending to much higher energies than in ENDF/B-IV, resulting in a much smaller continuum cross section. Thus inelastic scattering at high incident energies leads to higher average final energies in ENDF/B-V than in ENDF/B-IV. This tends to counteract the effect on the flux spectrum of having higher total inelastic in ENDF/B-V than in ENDF/B-IV. In performing the inelastic evaluation, one revision was made in the first draft (Ref. 10) to give improved agreement with calculations for ZPR-6-7. However, no changes outside the estimated uncertainties were made.

MF=4 Angular Distributions.
 MT=2 Elastic. The elastic angular distributions are expressed as $\mathcal{F}(\ell)$ coefficients in the center of mass system at 31 energies. They were obtained using data referenced in ANL/NDM-32 above 45 keV. Coupled channel optical model calculations assisted in the analysis. Below 45 keV ENDF/B-IV was used. The 20 MeV $\mathcal{F}(\ell)$ were modified in order to avoid negative excursions caused by the restriction to NL=20.

MT=18, 19, ... 21, 38, and 91. Angular distributions were taken as isotropic in lab system.

MT=51, 52... 77. Inelastic Levels. The angular distributions for the inelastic levels are given as probabilities in the C.-of- M. system. They were calculated with the model described in Ref. 9. Comparisons with experiment are given in ANL/NDM-32. In ENDF/B-IV all but the four highest pseudo levels were assumed isotropic.

MF=5 Secondary Energy Distributions.

MT=18 Total Fission. The total fission spectrum is an energy-dependent watt spectrum (LF=11) with parameters chosen to give the same average energy as the combination of MT=19,20,21,38.

MF=5	MT=19	First Chance Fission. An energy-dependent Watt spectrum was determined using methods similar to those of Ref.15. A Watt spectrum has more neutrons at intermediate energies, and fewer at low and high energies than a Maxwellian of the same average energy.
	MT=20, 21, 38.	Second, 3rd, and 4th Chance Fission. A combination of a Watt spectrum and 1,2, or 3 LF=9 laws for MT=20, 21, and 38, respectively, was used.
	MT=91	Inelastic Continuum. LF=1 by A. Smith.
	MT=455	Delayed Neutron Spectra. See Ref.1.
MF=12		Photon Production Multiplicities.
	MT=18	Fission, and MT=102 (n, γ) multiplicities are given.
MF=13		Photon Production Cross Sections.
	MT=3	Non-elastic. Photon production cross sections are given above the inelastic threshold.
MF=14		Photon Angular Distributions.
	MT=3, 18, 102.	Photon angular distributions are isotropic.
MF=15		Photon Energy Spectra.
	MT=3, 18, 102.	Continuous photon energy spectra are present. Description of the evaluation of MF=12-15 is included in ANL/NDM-32.

2.3 References

1. R. E. Kaiser and S. G. Carpenter (ANL-West), Private Communication (March 78). Data inserted into file at BNL by R. Kinsey 4/18/78.
2. S. A. Cox, ANL/NDM-5 (1974).
3. C. Besant et al., Sem. Fast Pulsed Reactors CONF-760111 (1976).
4. F. Manero and V. Konshin, Atomic Energy Review 10 No. 4 (1972).
5. M. Soleilhac, Revised data received from National Nuclear Data Center.
6. B. Nurpeisov et al., Soviet Atomic Energy Translation 807 (March 1976).
7. G. deSaussure et al., PNE 3, 87 (1979).
8. D. Olsen et al., Nucl. Sci. Eng. 62, 479 (1977).

9. P. Guenther, D. Havel and A. Smith, ANL-NDM-22 (1976).
10. E. Pennington, W. Poenitz and A. Smith, Transactions American Nuclear Society 26, 591 (1977).
11. W. P. Poenitz, ANL/NDM-45 (1979).
12. W. P. Poenitz and A. B. Smith, Eds. ANL-76-90 (1976).
13. R. Slovacek et al., Nucl. Sci. Eng. 62, 455 (1977).
14. M. Caner, M. Segev and W. Yiftah, Nucl. Sci. Eng. 59, 395 (1976).
15. E. Kujawski and L. Stewart, Transactions American Nuclear Society 24, 453 (1976).
16. L. W. Weston, Personal Communication to B.A. Magurno November 12, 1982.
17. R. Sher and C. Beck, EPRI NP-1771/81 + Rev. 1/83 plus Personal Communication to B. A. Magurno (NNDC) 2/83.

Fig. 1 Comparison of the Froehner evaluation (line) of the capture cross section of ^{238}U which was accepted for ENDF/B-VI with the recommendation of the Standards Committee (points).

Fig. 2 The ENDF/B-VI evaluation of the fission cross section of ^{238}U (line) and the Standards Committee recommendation (points).

SUMMARY DOCUMENTATION FOR ^{237}Np ENDF/B-VI, MAT = 9346

P. G. Young

Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM 87545

I. SUMMARY

In 1984, a theoretical analysis and interim evaluation of $n+^{237}\text{Np}$ reactions up to 5 MeV was performed at Los Alamos.¹ The evaluation was comprised roughly of the resonance and low-energy region data of Derrien,² the Los Alamos analysis from 10 keV to 5 MeV,¹ and ENDF/B-V from 5 to 20 MeV.³ For ENDF/B-VI, the theoretical analysis was extended to 20 MeV and a revised evaluation was performed covering the energy range 10^{-5} eV to 20 MeV.

Essentials of the new evaluation are given in the sections that follow. Some general features are:

1. The resonance parameters and low-energy data ($E_n < 8$ keV) are taken from a revision⁴ by Derrien of his 1980 evaluation² but are essentially identical to the earlier work.
2. Our previous theoretical analysis was extended to 20 MeV. Results from the new analysis are used to replace all the ENDF/B-V data that was incorporated in our 1984 revision, that is, the $(n,n'$ continuum), $(n,2n)$, and $(n,3n)$ cross sections and energy distributions. The new evaluation includes correlated energy-angle distributions through use of Kalbach⁵ systematics and the new ENDF/B File 6 formats.⁶ Additionally, distorted-wave Born approximation calculations for vibrational levels are included in the (n,n') data, which results in a more realistic neutron emission spectrum. Adjustments were made in our calculation of the $(n,2n)$ cross section near threshold for experimental data.
3. The ENDF/B-V evaluation of (n,f) cross sections above $E_n=1$ MeV have been replaced by our evaluation of new measurements^{7,8} as well as the existing data base. The new evaluation was also used as an input for our theoretical analysis at higher energies.
4. The prompt nubar evaluation was revised slightly to better agree with experimental data,^{9,10} after renormalization of the latter for new ENDF/B-VI standards. The delayed nubar values were updated with the best estimates now available,¹¹ which will also be used for ENDF/B-VI.

II. THEORETICAL ANALYSIS

The primary purpose for performing the theoretical analysis was to provide data on the reactions and energy ranges where little or no experimental data exist. In the case of $n+^{237}\text{Np}$, there were no experimental total cross section data available above $E_n=14$ keV when the analysis began, virtually no elastic or inelastic scattering data, only fragmentary information on $(n,2n)$ reactions, and no experimental data on $(n,3n)$ reactions or secondary neutron energy distributions. The situation for radiative capture was somewhat better as

several measurements exist below $E_n = 2$ MeV, although there are significant discrepancies among some of the measurements. The reaction that is best described experimentally is fission, as new fission ratio measurements have recently been completed at WNR⁷ and Argonne,⁸ and prompt nubar measurements have been made over much of the energy range of interest here. Therefore, the main function of the theoretical analysis was to provide the total, elastic, inelastic, (n,2n), and (n,3n) cross sections, as well as the angular and energy distributions of secondary neutrons.

To summarize the analysis briefly, coupled-channel deformed optical model calculations were performed with the ECIS code¹² over the incident neutron energy range from 0.001 to 20 MeV. As described earlier,¹ an optical model potential based on the work of Lagrange¹³ was chosen for our calculations, with some modification for the present analysis to improve the calculations above 10 MeV and to make the potential consistent with those used in analyses of ^{235,238}U and ²³⁹Pu experimental data for ENDF/B-VI.¹⁴ The coupled-channel calculations are used in the present analysis to obtain total, elastic, and (n,n') cross sections to the first and third excited levels of ²³⁷Np, which are members of the ground state rotational band, and to provide neutron transmission coefficients for Hauser-Feshbach statistical theory calculations.

The Hauser-Feshbach statistical calculations were performed with the COMNUC¹⁵ and GNASH¹⁶ codes. Both codes include a double-humped fission barrier model, using uncoupled oscillators for the barrier representation in GNASH and coupled or uncoupled oscillators in COMNUC. The COMNUC calculations include width-fluctuation corrections, which are needed at lower energies, whereas GNASH provides the preequilibrium corrections that are required at higher energies. Accordingly, COMNUC was used in the calculations below the threshold for second chance fission (approximately 5 MeV), utilizing fairly strongly damped coupled oscillators. The GNASH code was employed at higher energies, using uncoupled oscillators for second and higher chance fission. Fission transition state spectra were assumed identical within each compound system and were constructed by taking known (or calculated) energy levels and compressing their spacing by a factor of 2. As usual, Gilbert and Cameron¹⁷ phenomenological level density functions were used to represent continuum levels at ground-state deformations, appropriately matched to available experimental level data. Multiplicative factors were applied to the level density functions to account for enhancements in the fission transition-state densities at barriers due to increased asymmetry conditions.

Distorted-wave Born approximation calculations were performed with the DWUCK code¹⁸ to estimate the energy dependence of direct (n,n') reactions to vibrational states in ²³⁷Np. Because of the paucity of data on ²³⁷Np, the deformation parameters needed for normalizing the DWUCK calculations were estimated from systematics. In particular, the required B(E2) and B(E3) values were assumed to be similar to those determined in analyses for ^{235,238}U and ²³⁹Pu.¹⁴ All the $\ell=2$ and $\ell=3$ vibrational strength was placed into two fictitious states near $E_x = 1$ MeV.

III. DESCRIPTION OF EVALUATED DATA BASE

A. Total Cross Section

The total cross section below 8 keV is taken from the evaluation of Derrien,² which is also being used for the JEF-2 data file.⁴ From 8 keV to approximately 1 MeV, the coupled-channel deformed optical model calculations described above are used directly. Above ≈ 1 MeV, results from a covariance analysis of new measurements from WNR¹⁹ were utilized for the evaluation. The ENDF/B-VI evaluation of σ_{tot} is compared with the ENDF/B-V.²³ evaluation and with the Los Alamos measurements¹⁹ in Fig. 1.

B. Elastic Cross Section

The elastic cross section at all energies is obtained from the difference of the total and nonelastic cross sections. Below 8 keV it comes from the Derrien evaluation,² and at higher

energies it is determined mainly by the coupled-channel calculations (reaction cross section) and by the Los Alamos total cross section data.¹⁹ The ENDF/B-VI results are compared with the ENDF/B-V.2 and JEF-2 evaluations in Fig. 2. Significant differences occur among the various evaluations, with the spread exceeding 10% at some energies.

C. Fission Cross Section

To have confidence in our predictions of (n,n') and (n,xn) cross sections, it is essential that the theoretical analysis reasonably reproduce the measured (n,f) cross section. A comparison is given in Fig. 3 of our theoretical fit (dashed curve) to a selection of recently measured (n,f) cross sections as well as to our ENDF/B-VI evaluation (solid curve). The theoretical curve is seen to agree to roughly $\pm 5\%$ or better with the recommended values at all energies. To obtain the best possible estimate of the other reactions, the difference between the calculated and evaluated (n,f) cross sections was distributed proportionately among the compound nucleus reactions occurring at a given energy. Because of the dominance of direct and preequilibrium processes in (n,n') reactions above a few MeV, most of this difference was distributed to the (n,2n) and (n,3n) cross sections.

The present evaluation is compared with all of the more recent (n,f) cross section measurements in Figs. 4 and 5, as well as the ENDF/B-V.2 and JEF-2 evaluations.

D. Radiative Capture Cross Section

The radiative capture cross section is left unchanged from our 1984 evaluation.¹ The values below 5 MeV are taken from Derrien's 1980 evaluation,² which is identical to JEF-2⁴ at those energies. Above 5 MeV, the present evaluation is taken from ENDF/B-V.2. The results are compared in Fig. 6 to the available experimental data and to the various data evaluations.

E. Inelastic Neutron Cross Sections

The total inelastic neutron cross sections from the ENDF/B-V.2, JEF-2, and ENDF/B-VI evaluations are compared in the lower half of Fig. 7. The only evaluation that does not include direct reaction effects is ENDF/B-V.2, which is the reason those data fall essentially to zero at 10 MeV.

Discrete (n,n') cross sections are included for 31 excited states of ²³⁷Np. The $J^\pi = 7/2^+$ and $9/2^+$ first- and third- excited states are members of the $K = 5/2$ ground-state rotational band and include coupled-channel as well as compound nucleus contributions. The remaining discrete-state cross sections through the 29th excited state are entirely from compound nucleus processes and were calculated with the COMNUC code. The 30th and 31st excited states ($E_x = 0.984$ and 1.013 MeV, respectively) are vibrational states representing $l=2$ and $l=3$ transitions and were obtained from DWBA calculations, as described earlier. The cross sections for all purely compound-nucleus states are zeroed in the evaluation for neutron energies above 6 MeV.

F. Cross Sections for (n,xn) Reactions

The (n,2n), (n,3n), and (n,4n) cross sections result mainly from the theoretical analysis, after adjustment for the calculated/measured fission cross section difference as described above under item III. Near threshold, the (n,2n) cross section was adjusted to improve agreement with experiment. The ENDF/B-VI results for the (n,2n) cross section are compared to the ENDF/B-V.2 and JEF-2 evaluations in the upper half of Fig. 7. The ENDF/B-V.2 values for the (n,2n) reaction are significantly lower than our present results at most energies. The present ENDF/B-VI values are somewhat closer to Derrien's evaluation,⁴ although significant differences are evident.

In 1986, Arthur²⁰ obtained an estimate of the total $^{237}\text{Np}(n,2n)$ cross section by evaluating the measured $(n,2n)$ data for excitation of the 22.5-h (short-lived) metastable state of ^{236}Np , and then applying calculations of the long- and short-half-life components of the reaction by M. Gardner and D. Gardner²¹ of Livermore. Arthur's results, which are entirely independent of the present analysis, fall within $\pm 10\%$ of the present evaluation at most energies.

Because of the lack of knowledge about the structure of ^{238}Np , the present analysis only provides a good determination of the total $(n,2n)$ cross section, that is, it does not determine accurately the splitting of the $(n,2n)$ cross section into the long- and short-lived components. It is possible to estimate the individual components, however, by using the calculations of Gardner and Gardner, which utilized detailed structure calculations, together with our calculation of the total $(n,2n)$ cross section.

G. Nubar

The evaluation of delayed nubar (ENDF/B MF=1, MT=455) was updated to include England's latest values,¹¹ which are part of his ENDF/B-VI evaluation. Prompt nubar (MF=1, MT=456) in our 1984 evaluation,¹ which was based entirely on calculations using the Madland-Nix formalism,²² was adjusted slightly in the present work to agree better with the available experimental data base. The measurements were renormalized where appropriate for consistency with ENDF/B-VI standards,²³ that is, $\bar{\nu}_{\text{sf}}(^{252}\text{Cf}) = 3.7676 \pm 0.0049$. A comparison of the ENDF/B-VI prompt nubar curve for $n + ^{237}\text{Np}$ with the experimental data and with the Madland-Nix calculation (dashed curve) is given in Fig. 8.

H. Fission Neutron Spectra

The fission neutron spectra in the revised evaluation remain unchanged from our 1984 evaluation. The distributions are based on calculations using Madland-Nix theory²² and are represented as tabulated distributions (ENDF/B Law 12). Only the composite distribution is given, but it includes the individual first-, second-, and third-chance fission components.

I. Discrete Elastic and Inelastic Neutron Angular Distributions

The elastic and inelastic neutron angular distributions for discrete states are all given as Legendre expansions. The elastic (MT=2) and inelastic distributions for the 1st and 3rd excited states (MT=51 and 53) were obtained by summing the coupled-channel optical model (ECIS) and the compound nucleus (COMNUC) calculations. The distributions for the 2nd and 4th through the 29th excited states result from pure compound nucleus calculations. The angular distributions for the 30th and 31st excited states represent $l=2$ and $l=3$ vibrational state contributions and were obtained from distorted-wave Born approximation calculations with the DWUCK code.

J. Continuum Inelastic and (n,xn) Energy-Angle Distributions

Correlated energy-angle distributions are given in the new evaluation for continuum inelastic, $(n,2n)$, $(n,3n)$, and $(n,4n)$ neutrons (MT = 91, 16, 17, and 37, respectively). The data are represented using the new ENDF/B-VI File 6 format and make use of the option for use of Kalbach⁵ systematics to specify angular distributions as functions of emitted neutron energy. All the energy distributions and preequilibrium ratios, which are required parameters for the Kalbach distributions, were obtained from the GNASH calculations. The RECOIL code²⁴ was used to extract the neutron distributions for individual reactions from the GNASH outputs.

REFERENCES

1. E. D. Arthur, D. G. Madland, and P. G. Young, "Calculation and Evaluation of $n + {}^{237}\text{Np}$ Cross Sections," in Applied Nuclear Science Research and Development Semiannual Progress Report (Cp. E. D. Arthur and A. D. Mutschlecner) LA-10288-PR (1985) p. 13.
2. H. Derrien, J. P. Doat, E. Fort, and D. Lafond, "Evaluation of ${}^{237}\text{Np}$ Neutron Cross Sections in the Energy Range from 10^{-5} eV to 14 MeV," INDC(FR) - 42/L (1980).
3. F. Mann, ENDF/B-V.2 data file for ${}^{237}\text{Np}$ (MAT 1337), described in "ENDF/B Summary Documentation," B. A. Magurno and P. G. Young, Comp., Brookhaven National Laboratory report BNL-NCS-17541 (ENDF-201, Supplement 1), 1985 (available from the National Nuclear Data Center, Brookhaven National Laboratory, Upton, N.Y).
4. H. Derrien, personal communication, June 1989.
5. C. Kalbach, "Systematics of Continuum Angular Distributions: Extensions to Higher Energies," Phys. Rev. C37, 2350 (1988); C. Kalbach and F. M. Mann, "Phenomenology of Continuum Angular Distributions. I. Systematics and Parameterization," Phys. Rev. C23, 112 (1981).
6. P. F. Rose and C. L. Dunford, "ENDF-102: Data Formats and Procedures for the Evaluated Nuclear Data File, ENDF/B," preliminary draft, May, 1988.
7. P. W. Lisowski, J. L. Ullmann, S. J. Balestrini, A. D. Carlson, O. A. Wasson, and N. W. Hill, "Neutron-Induced Fission Cross-Section Ratios for ${}^{232}\text{Th}$, ${}^{235,238}\text{U}$, ${}^{237}\text{Np}$, and ${}^{239}\text{Pu}$ from 1 to 400 MeV," Int. Conf. on Nucl. Data for Science and Technology, Mito, Japan, May 30 - June 3, 1988 (Ed. S. Igarasi, Saikon Publ. Co., Ltd., 1988) p. 97.
8. J. W. Meadows et al., Nucl. Sci. Eng. 85, 271 (1983); J. W. Meadows et al., Ann. Nucl. En. 15,421 (1988).
9. V. V. Malinovsky, V. G. Vorob'eva, B. d. Kuz'minov, V. M. Piksaikin, N. N. Semenova, S. M. Solov'ev, and P. S. Soloshenkov, "Discrepancy of the Results of $\bar{\nu}_p$ Measurements in the Fission of ${}^{237}\text{Np}$ Nuclei by Neutrons," Atom. Energiya 54, 208 (1983).
10. J. Frehaut, A. Bertin, and R. Bois, "Mesure de $\bar{\nu}_p$ et $\bar{E}_\gamma$ Pour La Fission de ${}^{232}\text{Th}$, ${}^{235}\text{U}$ et ${}^{237}\text{Np}$ Induite Par des Neutrons d'Energie Comprise Entre 1 et 15 MeV," Int. Conf. on Nucl Data For Science and Tech., Antwerp, Netherlands, Sept. 6-10, 1982, p. 78.
11. T. R. England, personal communication (June, 1989).
12. J. Raynal, "Optical-Model and Coupled-Channel Calculations in Nuclear Physics," IAEA SMR-9/8, Int. At. En. Agency (1970).
13. G. Haouat, Ch. Lagrange, J. Lachkar, J. Jary, Y. Patin, and J. Sigaud, "Fast Neutron Scattering Cross Sections for Actinide Nuclei," Int. Conf. on Nuclear Cross Sections for Technology, Knoxville, Tennessee (Oct. 22-26, 1979) p. 672.

14. For example, see P. G. Young and E. D. Arthur, "Calculation of $^{235}\text{U}(n,n')$ Cross Sections for ENDF/B-VI," Int. Conf. on Nucl. Data for Science and Technology, Mito, Japan, May 30 - June 3, 1988 (Ed. S. Igarasi, Saikon Publ. Co., Ltd., 1988) p. 603.
15. C. L. Dunford, "A Unified Model for Analysis of Compound Nucleus Reactions," AI-AEC-12931, Atomics Int. (1970).
16. P. G. Young and E. D. Arthur, "GNASH: A Preequilibrium Statistical Nuclear-Model Code for Calculation of Cross Sections and Emission Spectra," Los Alamos Scientific Laboratory report LA-6947 (Nov. 1977); E. D. Arthur, "The GNASH Preequilibrium-Statistical Model Code," LA-UR-88-382 (1988).
17. A. Gilbert and A. G. W. Cameron, "A Composite Nuclear-Level Density Formula with Shell Corrections," Can. J. Phys. 43, 1446 (1965).
18. P. D. Kunz, "DWUCK - A Distorted-Wave Born Approximation Program," unpublished.
19. E. D. Arthur, " $^{237}\text{Np}(n,2n)$ Values," Los Alamos Internal Memo T-2-M-1701 to Steve Becker, Group X-2, April 2, 1986.
20. P. W. Lisowski, Los Alamos National Laboratory, personal communication, 1990.
21. M. Gardner and D. Gardner, Lawrence Livermore National Laboratory, personal communication to E. D. Arthur, 1986, and P. G. Young, 1989.
22. D. G. Madland and J. R. Nix, "New Calculation of Prompt Fission Neutron Spectra and Average Prompt Neutron Multiplicities," Nucl. Sci. Eng. 81, 213 (1982).
23. A. D. CARLSON, W. P. POENITZ, G. M. HALE, and R. W. PEELLE, "The Neutron Cross Section Standards Evaluation for ENDF/B-VI," *Proc. Int. Conf. on Nucl. Data for Basic and Applied Science*, Santa Fe, N. M., 13-17 May 1985, V2, p. 1429.
24. R. E. MacFarlane, personal communication, 1989.

Figure 1. Measured and evaluated $n + {}^{237}\text{Np}$ total cross section between 0.1 and 20 MeV. The solid curve represents the ENDF/B-VI evaluation, and the dashed curve is ENDF/B-V.

Figure 2. Comparison of evaluated elastic scattering cross sections for $n + {}^{237}\text{Np}$ interactions. The solid curve represents the ENDF/B-VI evaluation, the dotted curve is a preliminary JEF-2 evaluation, and the dashed curve is ENDF/B-V.

$^{237}\text{Np}(n,f)$ Cross Section

Figure 3. Comparison of calculated, evaluated, and measured $^{237}\text{Np}(n,f)$ cross sections from 0.4 to 20 MeV. The solid curve represents the ENDF/B-VI evaluation, and the dashed curve is from the GNASH theoretical analysis.

Figure 4. The $^{237}\text{Np}(n,f)$ cross section from 0 to 6 MeV. The solid curve represents the ENDF/B-VI evaluation, the dotted curve is a preliminary JEF-2 evaluation, and the dashed curve is ENDF/B-V.

$^{237}\text{Np}(n,f)$ Cross Section

Figure 5. The $^{237}\text{Np}(n,f)$ cross section from 6 to 20 MeV. The solid curve represents the ENDF/B-VI evaluation, the dotted curve is a preliminary JEF-2 evaluation, and the dashed curve is ENDF/B-V.

$^{237}\text{Np}(n,\gamma)^{238}\text{Np}$ Cross Section

Figure 6. The $^{237}\text{Np}(n,\gamma)^{238}\text{Np}$ radiative capture cross section from 0.02 to 20 MeV. The solid curve represents the ENDF/B-VI evaluation, and the dashed curve is ENDF/B-V.

Figure 7 Comparisons of the $^{237}\text{Np}(n,n')^{237}\text{Np}^*$ and $^{237}\text{Np}(n,2n)^{236}\text{Np}$ cross sections from various data evaluations. The solid curve represents the ENDF/B-VI evaluation, the dotted curve is a preliminary JEF-2 evaluation, and the dashed curve is ENDF/B-V.

Figure 8. Measured and evaluated values of prompt $\bar{\nu}$ for $n + {}^{237}\text{Np}$ fission reactions. The solid curve represents the ENDF/B-VI evaluation, and the dashed curve is from a 1984 Los Alamos evaluation,¹ which is based on calculations using Madland-Nix theory.²²

²³⁹₉₃Np

Reference: No Primary Reference
Evaluator: R. Q. Wright
Evaluated: December 1988
Material: 9352
Content: Neutron transport

File Comments

ENDF/B-VI MAT 9352 Revised by R. Q. Wright (ORNL)
Converted from JENDL - 2 Evaluation, MAT 2932 (See below)

Summary of Changes

The JENDL-2 ²³⁹Np evaluation, MAT 2932, has been revised below 4.0 eV by R. Q. Wright (ORNL) in March 1987. In this energy range the capture cross section, MF = 3, MT = 102, is given by:

$$\sigma_c(E) = 77.0 \times (E_0/E)^{1/2}$$

where E is the energy in eV and $E_0 = 0.0253$ eV.

This change was made in order for the thermal capture cross section to be in agreement with the value given in Ref. 1. The total cross section was modified to be in agreement with the sum of the elastic and the revised capture cross sections.

In addition, the total cross sections at 6.2533 and 7.5000 MeV were increased by about 0.1% so that they would be in agreement with the sum of the partial cross sections at these energies.

The format was changed from ENDF/B-IV to ENDF/B-V. The NNDC converted the file to ENDF-6 format.

Reference

1. S. F. Mughabghab, "Neutron Cross Sections" Vol 1, Neutron Resonance Parameters and Thermal Cross Sections, Part B: Z = 61-100 Academic Press, (1984).

Summary of JENDL Evaluations

History

- 76-03 The evaluation for JENDL-1 was performed by Y. Kanda (Kyushu University) and the JENDL-1 compilation group. Details are given in Ref. /1/.
- 83-03 JENDL-1 data were adopted for JENDL-2 and extended to 20 MeV. MF=5 was revised.
- 84-01 Comment data were added.

File Information

- MF=1 MT=451 Descriptive data and dictionary.
- MT=452 Number of neutrons per fission taken from the ENDF/B-IV ^{237}Np data.

- MF=2 MT=151 No resonance parameters were given.

2200-m/sec Cross Sections and Calculated Resonance Integrals

	2200 m/sec	Res. Integ.
elastic	10.50 b	-
capture	37.00 b	445. b
fission	0.0 b	7.06 b
total	47.50 b	-

- MF=3 Neutron cross sections below 4.0 eV.
- MT=1 The total cross section is sum of partial cross sections.
- MT=2 The constant cross section of 10.5 barns for elastic scattering was assumed from $\sigma = 4\pi(0.147 \times a^{1/3})^2$.
- MT=18 Fission assumed to be zero barns.
- MT=102 Capture in the form of $1/v$ was assumed. The 2200-m/sec cross section was adopted from the experimental data by Stoughton and Halperin /2/.

MF=3 Neutron cross sections above 4.0 eV.
 MT=1 Total σ calculated with the optical and statistical model code CASTHY /3/. Optical potential parameters were obtained by Ohta and Miyamoto /4/ using the total cross section of ^{239}Pu .

$$\begin{array}{llll} V = 45.87 - 0.2E & W_i = 0.06 & W_s = 14.1 & V_{so} = 7.3 \text{ (MeV)} \\ r = 1.27 & r_i = 1.27 & r_s = 1.302 & r_{so} = 1.27 \text{ (fm)} \\ a_0 = 0.652 & a_i = 0.315 & a_s = 0.98 & a_{so} = 0.652 \text{ (fm)} \end{array}$$

MT=2 Elastic scattering calculated with CASTHY /3/.
 MT=4, 51-58,91 Inelastic scattering calculated with CASTHY /3/. The level scheme was adopted from the Nuclear Data Sheets Vol.6.

No.	Energy (MeV)	Spin-Parity
g.s.	0.0	5/2 +
1	0.03114	7/2 +
2	0.07112	9/2 +
3	0.07467	5/2 -
4	0.11766	11/2 +
5	0.1230	7/2 -
6	0.17305	9/2 -
7	0.2414	11/2 -
8	0.320	13/2 -

Levels above 430 keV were assumed to be overlapping. In the calculation the capture, fission, (n,2n) and (n,3n) cross sections were considered as competing processes.

MT=16, 17 (n,2n) and (n,3n) cross sections were calculated using Pearlstein's method /5/.

MT=18 Fission was estimated from the ^{237}Np fission cross section by normalization with the neutron separation energies.

MT=102 The capture cross section below 100 keV was calculated from $\sigma = 435/\sqrt{E}$ barns. Above 100 keV, the shape of the experimental data for ^{237}Np was adopted from Nagle et al. /6/ and normalized to 1.4 barns at 100 keV.

MF=4 Distributions of secondary neutrons.

MT=2 Calculated with the CASTHY code /3/.

MT=51, 52-58 Isotropic in the center-of-mass system.

MT=16, 17,18,91 Isotropic in the laboratory system.

MF=5 Energy distributions of secondary neutrons.
MT=16, 17,91 Evaporation Spectrum.
MT=18 Maxwellian fission spectrum estimated from $Z^{2/A}$ systematics /7/.

References

1. Igarasi S. et al. : JAERI 1261 (1979).
2. Stoughton R. W. and Halperin J. : Nucl. Sci. Eng., 6, 100 (1959).
3. Igarasi S. : J. Nucl. Sci. Technol., 12, 67 (1975).
4. Ohta M. and Miyamoto K. : J. Nucl. Sci. Technol., 10, 583 (1973).
5. Pearlstein S. : Nucl. Sci. Eng., 23, 238 (1965).
6. Nagel R. J. et al. : 1971 Knoxville Conf., 259 (1971).
7. Smith A. B. et al. : ANL/NDM-50 (1979).

SUMMARY DOCUMENTATION FOR ^{239}Pu ENDF/B-VI, MAT = 9437

P. G. Young, L. W. Weston,* and W. P. Poenitz†

Theoretical Division
Los Alamos National Laboratory
Los Alamos, New Mexico 87545

I. SUMMARY

The ENDF/B-VI evaluation for ^{239}Pu is based on a new evaluation of the available experimental data above the resonance region,¹ together with a new analysis of the resolved and unresolved resonance regions by Derrien and de Saussure². The evaluation also makes use of results from the ENDF/B-VI standards analysis at thermal energies and above the resonance region, although minor adjustments were made to those results.

The resolved resonance data were obtained from a multi-level Reich-Moore analysis that simultaneously fit transmission, fission, absorption, and capture data. The resolved resonances extend to an incident neutron energy of 2 keV, and the unresolved region continues with average parameters to an incident energy of 30 keV. Prompt $\bar{\nu}$ in the resonance region was obtained from the evaluation of Fort et al.¹ after minor renormalization for consistency with ENDF/B-VI standards.

Above the resonance region, a combination of experimental data evaluation and theoretical calculations are used to represent the ^{239}Pu data. Covariance analyses were performed of the complete experimental data base for the total and fission cross sections and for prompt $\bar{\nu}$. These results were used directly (after minor smoothing) for the evaluated total cross section and for $\bar{\nu}_p$. Similarly, the results from the covariance analysis were used directly for $\sigma_{n,f}$ above $E_n \approx 2$ MeV. Between 50 keV and 1.25 MeV, the fission cross section was obtained from the ENDF/B-VI standards analysis, after renormalization by a factor of 1.007 (about 1 standard deviation) to improve consistency with integral data. Below 50 keV, $\sigma_{n,f}$ was matched smoothly with the results of the resonance analysis, which required a decrease in the standards result by 4 % at 50 keV.

The elastic, inelastic, and (n,xn) cross sections, angular distributions, and energy spectra above the resonance range were obtained from a theoretical analysis that was optimized to match the total and fission cross sections. The inelastic and (n,xn) continuum data are represented in energy-angle correlated arrays using the ENDF/B-6 File 6 format. The theoretical analysis involved coupled-channel optical model, Hauser-Feshbach statistical theory, Moldauer width-fluctuation, fission theory, and preequilibrium calculations over the energy range $E_n = 0.001 - 20$ MeV.

II. EVALUATION DETAILS

A selection of illustrative figures is shown on the pages that follow. Included in the figures are comparisons of the ENDF/B-V.2 and ENDF/B-VI evaluations of the total, (n,f), (n,n'), and (n,2n) with experimental data. Similar comparisons are given for $\bar{\nu}_p$, for the ratio of the $^{239}\text{Pu}(n,f)$ and $^{235}\text{U}(n,f)$ cross sections, and for a small selection of elastic scattering angular distributions. A more detailed summary of the evaluated data files is included in the ENDF File 1 descriptive comments, reproduced in the section following the figures.

* Oak Ridge National Laboratory, Oak Ridge, Tennessee

† Argonne National Laboratory, Idaho Falls, Idaho

¹ P. G. Young and R. E. MacFarlane, "Evaluation and Testing of n + ^{239}Pu Data for ENDF/B-VI in the keV and MeV Energy Region," submitted to Int. Conf. on Nucl. Data for Sci. and Tech., 13-17 May 1991, Jülich.

² See references in the ENDF/B File 1 comment section that follows.

n + Pu-239 Total Cross Section

Figure 1. Neutron total cross section of ^{239}Pu between 0.03 and 20 MeV. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

Pu-239/U-235 (n,f) Cross Section Ratio

Figure 2. Ratios of the $^{239}\text{Pu}(n,f)$ and $^{235}\text{U}(n,f)$ cross sections between 0.02 and 20 MeV. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

Pu-239(n,f) Cross Section

Figure 3. The $^{239}\text{Pu}(n,f)$ cross section between 0.03 and 16 MeV. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

n + Pu239 Nubar Prompt

Figure 4. Comparison of measured and evaluated values of prompt $\bar{\nu}$ for neutron-induced reactions with ${}^{239}\text{Pu}$. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

Pu239(n,2n) Cross Section

Pu239(n,n') Cross Section

Figure 5. The $^{239}\text{Pu}(n,n')^{239}\text{Pu}^*$ and $^{239}\text{Pu}(n,2n)^{238}\text{Pu}$ reactions between threshold and 20 MeV. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

Figure 6. Sample comparisons of the evaluated $n + {}^{239}\text{Pu}$ elastic scattering angular distributions with experimental data. The solid curve is the ENDF/B-VI evaluation, the dashed curve is ENDF/B-V, and the points represent experimental data.

²³⁹₉₄Pu

Reference: No Primary Reference
Evaluators: P. Young, L. Weston, W. Poenitz
Evaluated: April 1989
Material: 9437
Content: Neutron transport, Gamma production, Covariances

SUMMARY OF ENDF/B-VI EVALUATION
P.G.Young, L.W.Weston, and W.P.Poenitz

1. Resonance region evaluation performed by H.Derrien and G.de Saussure ORNL/TM-10986 (January, 1989). NEW PARAMETER SET EXTENDING TO 2 KEV SUBSTITUTED NOV. 89.
2. Prompt nubar evaluation obtained from work of E.Fort (Nuc. Sci.Eng.99, 375 (1988).
3. Delayed nubar evaluation (En89).
4. Energy range above unresolved resonances (0.03 - 20 MeV) was evaluated by P.Young, using results from W.Poenitz, G.Hale, R.Peelle, and A.Carlson from the simultaneous standards evaluation (Santa Fe Conf.(1985)p.1429), and with contributions from R.MacFarlane and L.Weston. More details are given below, and full documentation will be issued later.

THERMAL REGION

The Reich-Moore resonance parameters have been obtained from the analysis of several transmission, fission, absorption and capture experimental data(1,4,5,7,10). This set of resonance parameters also considered the 1988 fission data of Weston and Todd. The thermal region was then refitted and the following were obtained for the 0.0253 cross sections.

	Evaluation	Proposed
	293K	Standard values(barn)(11)

Fission	747.08	747.99+-1.87
Capture	271.39	271.43+-2.14
Scattering	8.00	7.88+-0.97
Total	1025.79	1027.30+-5.00

The experimental fission cross sections were renormalized to the value of 748.0 barns at 0.0253 eV in agreement with the proposed standard value and with the up-dated absolute value of Deruyter (12). The other experimental absolute value is the 1025.0 ± 6.0 barns obtained by Spencer(10) for the total cross section. Fitting the renormalized experimental fission, capture and absorption data and Spencer experimental transmission over the energy range 0.02 eV to 7.0 eV one obtains, at 0.0253 eV, 270.17 barns for the capture cross-section and 747.19 barns for the fission cross-section, in very good agreement with the proposed standard values.

The Weston and Todd 1988 fission data, obtained on a 80 m flight path with a resolution comparable to the Harvey transmission resolution, were included in the SAMMY fit experimental data base. Despite the difficulties encountered with a quite large residual background in the new Weston data, the new set of resonance parameters has improved compared to the previous one. Due to the improvement of the resolution in the fission experiment more resonances were identified in the high energy range of the data and the fission widths are more accurate.

A Reich-Moore SAMMY fit of Harvey transmission data and Weston 1988 fission data was performed in the energy range 1 keV to 2 keV. Preliminary results are given in the file. A background contribution in file 3 must be added. Further analysis should be performed to obtain more accurate sets of resonance parameters in 1 to 2 keV energy range. However, the present set of data should be most useful for the calculation of the self shielding in this energy range.

The scattering cross-section is 0.91 barn larger than the proposed standard. It corresponds to a radius of 9.46 ± 0.20 fm obtained in the analysis of the transmission data up to 1 keV. To obtain the the proposed standard value one should use a radius of 9.11 fm.

One should also note that the 293 degrees K cross-sections calculated at 0.0253 eV depend on the way the Doppler broadening calculation is performed. For instance using a Gaussian broadening function will give a fission cross-section about 2.5 barns larger than the one obtained from the accurate calculation which conserves the $1/v$ cross section. This difference is of the same order of magnitude as the accuracy on the proposed standard fission cross section. The values given in the above table were obtained from an accurate calculation method (special SAMMY option for the thermal energy range).

The following table shows the experimental cross sections averaged over two energy intervals and compared to the calculated values:

Average cross-sections(barn)

References(1-10) 0.02 to 0.06 eV 0.02 to 0.65 eV

	exp	calc (293K)	exp	calc (293K)
Gwin71 fission	631.41		843.71	
Gwin76 fission	631.41		838.39	
Gwin84 fission(*)	631.41	631.45(+0.01%)	837.18	839.01(+0.22%)
Deruyter70 fission	631.41		859.43	
Wagemans80 fission	631.41		862.56	
Gwin71 capture	243.84	242.84(-0.41%)	524.75	518.13(-1.26%)
Gwin76 absorpt(*)	875.90	874.29(-0.18%)	1359.96	1357.14(-0.21%)
Spencer84 tot(*)	883.20	882.99(-0.02%)	1361.69	1368.8 (+0.52%)

(*)These data had the largest weight in the thermal fit. The values between the parentheses give the percentage deviation between the calculated data and the experimental data.

The value of 631.4 barns for all the average experimental fission cross-sections in the energy range 0.02 eV to 0.06 eV corresponds to the renormalization of all the fission experiments to 748. $\pm$ 1. barns at 0.0253 eV. ORNL data are consistent within 0.8% over the 0.3 eV resonance. Deruyter and Wagemans data are about 2.5 % larger and were not included in the fit. When normalized on the standard value at 0.0253 eV, Gwin76 absorption agrees with the absorption obtained from Spencer total within 0.7% over the resonance. The present evaluation is essentially the result of a consistent analysis of all the available ORNL data with a larger weight on Gwin 1984 fission, Gwin 1976 absorption and Spencer transmission data.

THE RESOLVED RESONANCE REGION

The Reich-Moore resonance parameters are given in the energy range up to 2 keV. Four negative energy resonances and six fictitious resonances above 2 keV are used to represent the effect of the external region resonances. A constant value of 9.41 fm could be used for the scattering radius R' for the calculation of the cross-section in the entire energy range analysed.

The set of resonance parameters is essentially the result of a Bayesian Reich-Moore analysis(SAMMY)(13) of the Harvey transmission data(9) and Gwin and Weston 1984 fission data(7,8). The Harvey data were taken on a 80 m flight path at liquid nitrogen temperature with a resolution good enough to separate more than 80 % of the resonance up to 1 keV. Blons 1973 data(3), which have better resolution than 84 Weston data, were used to identify narrow fis-

sion resonances in the high energy region. A preliminary correlated fit on Harvey, Weston and Blons data, allowing the adjustment of the normalization coefficients and of the background corrections, has shown that no such adjustment was necessary to have consistency between Harvey data and Weston data. Blons needed a large re-adjustment of the background and of the normalisation. The final fit was performed by using only Harvey, Gwin 1984($E < 30$ eV) and Weston 1984($E > 10$ eV) as input to SAMMY.

If one compares the most recent experimental fission data, in the resolved resonance region above 100 eV, one finds that Weston data are the smallest and Blons data the largest. The following table shows the proposed ENDF/B-VI standard values(11) and the values obtained from the resonance parameters, averaged in the same energy intervals:

Cross-section in barns				
Energy (keV)	Calcul (293K)	Weston84	Blons73	Proposed Standard
0.1-0.2	18.159	18.095(-3.1%)	18.93(+1.4%)	18.66+-0.13
0.2-0.3	17.318	17.441(-2.7%)	17.79(-0.5%)	17.88+-0.12
0.3-0.4	8.116	8.130(-3.6%)	8.91(+5.7%)	8.43+-0.06
0.4-0.5	9.318	9.337(-2.5%)	9.71(+1.5%)	9.57+-0.07
0.5-0.6	14.944	15.170(-2.6%)	15.51(-0.3%)	15.56+-0.11
0.6-0.7	4.337	4.192(-6.4%)	4.63(+3.8%)	4.46+-0.04
0.7-0.8	5.340	5.385(-4.5%)	5.94(+5.5%)	5.63+-0.04
0.8-0.9	4.636	4.765(-4.5%)	5.11(+2.6%)	4.98+-0.04
0.9-1.0	8.050	8.165(-1.7%)	8.57(+3.3%)	8.30+-0.07
0.1-1.0	10.024	10.075(-3.1%)	10.57(+1.7%)	10.39

The values between parentheses are the percentage deviation from the standard data. The calculated cross-sections agree within 0.5% on average with Weston values and are about 3.5% smaller than the proposed standard values. They are only 1.8% smaller than ENDF/B-V over the energy range 0.1 to 1.0 keV. The authors of the present evaluation have the feeling that many of the experimental fission cross-sections suffer from an underestimation of the experimental background leading to a systematic overestimate of the cross-section. The authors of the ENDF/B-VI standard evaluation have probably not considered the background problems in this way. The very small errors they obtained are due to a statistical processing of a large amount of experimental data. Blons fission cross-sections are an example of data in which the underestimation of the experimental background could be very important (about 3 barns at 40 eV and 0.4 barn at 1 keV).

Gwin 1971 and Gwin 1976 absorption data(1,4) were not included in the R-matrix analysis of the resolved resonance region. Accurate absorption cross-sections should be calculated by the parameters obtained from the analysis of the transmission data. The following table shows average calculated values of the capture, absorption and alpha compared to Gwin 1976 data:

Cross-section in barns					

Energy(eV)	Calc. values (293K)			Gwin 1976 data	

	capt	absorp	alpha	absorp	alpha
7.3- 16.0	79.08	194.99	0.68	208.00	0.74(*)
16.0- 37.5	21.12	44.83	0.89	46.50	0.89(*)
37.5- 50.0	47.90	68.18	2.36	83.15	2.96(*)
50.0-100.0	34.56	91.59	0.61	92.84	0.63
100.0-200.0	15.78	33.94	0.87	33.66	0.87
200.0-300.0	16.01	33.33	0.92	34.69	0.94
300.0-400.0	9.83	17.94	1.21	18.31	1.16
400.0-500.0	4.07	13.39	0.44	13.56	0.44
500.0-600.0	11.12	26.06	0.74	26.54	0.72
600.0-700.0	6.51	10.84	1.50	11.57	1.54
700.0-800.0	5.01	10.35	0.94	10.52	0.97
800.0-900.0	3.78	8.42	0.82	9.30	0.82
900.0-999.9	5.39	13.44	0.67	13.23	0.70

(*) Gwin 1971 data

Below 500 eV, the calculated absorption values agree quite well with Gwin 1976 experimental data and are on average 2.9% lower above 500 eV. The calculated alpha values are on average 1.7% lower above 500 eV.

***** UNRESOLVED RESONANCE REGION*****

The average resonance parameters are given in the energy range 2 keV to 30 keV for 74 energy points. They were obtained by using the Cadarache statistical code FISINGA to fit the gross structure of the Saclay experimental total cross sections(17) below 4 keV and of selected experimental fission cross sections normalized to the ENDF/B-VI standard evaluation(11). Above 4 keV no high resolution total cross section data are available; average total cross sections were calculated to be consistent with the statistical parameters obtained in the resolved resonance region(14) and with the optical model parameters of Lagrange and Madland(15) obtained by fitting the experimental data in the

high energy range. A value of 9.46 fm was used for the effective radius. The values obtained for alpha are consistent with the experimental data.

The competitive width is not used for the inelastic scattering cross section. In each energy point of the unresolved region, the neutron width corresponds only to the elastic scattering cross section. The inelastic scattering cross section should be found in File 3.

The cross sections obtained at OK by processing the evaluated file by NJ0Y-87.1, are given in the following table, 'fiss' for the fission values and 'capt' for the capture values:

Energy(keV)	Cross sections (barn)		Energy(keV)	Cross sections (barn)	
	fiss	capt		fiss	capt
2.050	1.879	3.137	11.750	1.812	1.042
2.150	3.119	3.315	12.250	1.900	0.968
2.250	2.691	2.800	12.750	1.864	0.947
2.350	3.436	3.331	13.250	1.858	0.917
2.450	4.280	2.456	13.750	1.715	0.942
2.550	2.725	2.754	14.250	1.492	0.948
2.650	3.103	3.425	14.750	1.797	0.854
2.750	4.169	2.010	15.250	1.883	0.797
2.850	4.126	2.077	15.750	1.697	0.843
2.950	3.362	3.710	16.250	1.801	0.782
3.050	3.017	1.998	16.750	1.628	0.824
3.150	4.896	1.934	17.250	1.498	0.819
3.250	3.954	2.277	17.750	1.862	0.701
3.350	1.710	2.166	18.250	1.711	0.736
3.450	2.198	2.572	18.750	1.632	0.748
3.550	2.214	1.885	19.250	1.738	0.694
3.650	2.394	2.948	19.750	1.743	0.677
3.750	3.067	1.624	20.500	1.672	0.679
3.850	3.556	2.122	21.500	1.646	0.661
3.950	2.931	2.397	22.500	1.472	0.697
4.125	2.114	2.270	23.500	1.632	0.619
4.375	2.509	2.129	24.500	1.636	0.597
4.625	2.772	1.715	25.500	1.547	0.607
4.875	1.980	2.186	26.500	1.628	0.562
5.125	2.406	1.916	27.500	1.544	0.572
5.375	2.153	1.953	28.500	1.568	0.549
5.625	2.294	1.807	29.500	1.609	0.521

Average values of cross sections compared to the ENDF/B-VI standard evaluation (11) and alpha values compared to some experimental data are given in the following table:

Energy (keV)	* Cross section (barns) *				***** alpha *****			
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
2- 3	3.284	3.304	2.894	3.31	0.881	1.000	1.108	1.028
3- 4	2.992	3.000	2.213	2.20	0.740	0.720	0.895	0.820
4- 5	2.394	2.383	2.073	2.07	0.866	0.870	0.821	0.837
5- 6	2.266	2.301	1.863	1.91	0.822	0.820	0.867	0.834
6- 7	2.006	2.008	1.677	1.63	0.836	0.790	0.816	0.793
7- 8	2.134	2.054	1.409	1.34	0.660	0.640	0.630	0.605
8- 9	2.207	2.216	1.245	1.23	0.564	0.540	0.575	0.530
9-10	1.867	1.864	1.136	1.05	0.608	0.550	0.617	0.569
1-10	2.628	2.622	2.014	2.06	0.767	0.752	0.806	0.768
10-20	1.762	1.764	0.876	0.85	0.497	0.480	0.466	0.498
20-30	1.597	1.595	0.606	0.58	0.379	0.350	0.373	0.388

- (1) Fission cross section , present evaluation (OK)
- (2) Fission cross section , ENDF/B-VI standard (11)
- (3) Capture cross section, present evaluation (293K)
- (4) Capture cross section, Gwin et al.1976 (4)
- (5) Alpha value , present evaluation (293K)
- (6) Alpha value from Gwin et al. 1976 (4)
- (7) Alpha value from Sowerby-Konshin evaluation 1971 (16)
- (8) Average alpha value from experimental data

The fission and capture resonance integrals at OK are compared to ENDF/B-V data in the following table:

Energy range(ev)	Fission(barn)		Capture(barn)		

	ENDF/B-V present		ENDF/B-V present		
0.5 - 5.0	86.02	85.71	32.31	28.65	
5.0 - 10.0	26.03	25.08	20.14	19.06	
10.0 - 50.0	100.25	96.87	78.66	77.19	
50.0 - 100.0	40.32	40.47	27.23	25.93	
100.0 - 301.0	19.98	19.68	19.52	17.95	
301.0 -1000.0	10.15	10.05	8.54	8.35	

0.5 -1000.0	282.76	277.85	186.30	177.13	

By adding the ENDF/B-V value above 1 keV on obtain from the present evaluation:

Ri fission	297.22 barns
Ri capture	184.93 barns

the corresponding values from the ENDF/B-V evaluation are:

Ri fission 302.13 barns

Ri capture 194.10 barns

References

- 1-R.Gwin et al.,Nucl.Sci.Eng.,45,25(1971)
- 2-A.J.Deruyter et al.,J.Ncl.Ener.,26,293(1972)
- 3-J.Blons, Nucl.Sci.Eng.,51,130(1973)
- 4-R.Gwin et al.,Nucl.Sci.Eng.,59,79(1976)
- 5-R.Gwin et al.,Nucl.Sci.Eng.,61,116(1976)
- 6-W.Wagemans,Annals of Nucl.Ener.7,9,495(1980)
- 7-R.Gwin et al.,Nucl.Sci.Eng.,88,37(1984)
- 8-L.W.Weston et al.,Nucl.Sci.Eng.88,567(1984)
- 9-J.A.Harvey ,Private communication(1985)
- 10-R.R.Spencer et al.,Nucl.Sci.Eng.,96,318(1987)
- 11-A.Carlson et al.,Preliminary results of the ENDF/B-VI standard evaluation (Sept 8 1987)
- 12-A.J.Deruyter,J.Nucl.Ener.,26,293(1972)
- 13-N.M.Larson et al.,ORNL/TM-7485,ORNL/TM-9179,ORNL/TM-9719/R1
- 14-H.Derrien and G.de Saussure, ORNL-TM-10986(1988)
- 15-Ch.Lagrange and D.G.Madland,Phys.Rev.C,33,5(1986)
- 16-M.G.Sowerby et al.,At. Energy Rev.,10,4,453,IAEA,Vienna(1972)
- 17-H.Derrien,thesis,Orsay Serie A,1172(1973)

ENERGY REGION 0.03 TO 20 MEV

Principal LANL evaluators: P.G.Young, R.E.MacFarlane, E.D.Arthur

The evaluation above 10 keV is based on a detailed theoretical analysis utilizing the available experimental data. Coupled channel optical model calculations with the ECIS code (Ra70) were used to provide the total, elastic, and inelastic cross sections to the first 7 members of the ground state rotational band, as well as neutron elastic and inelastic angular distributions to the rotational levels. The ECIS code was also used to calculate neutron transmission coefficients. Hauser-Feshbach statistical theory calculations were carried out with the GNASH (Ar88, Yo77) and COMNUC (Du70) code systems, including preequilibrium and fission. DWBA calculations were performed with the DWUCK code (Ku70) for several vibrational levels, using B(E1) values inferred from (d,d') data on Pu238 and Pu240, as well as Coulomb excitation measurements. A weak coupling model (Pe69) was used to apply the Pu238 and Pu240 results to

states in Pu239.

This analysis is an extension of the calculations used for the ENDF/B-V.2 evaluation, described in reference Ar82.

*****MF=1 Descriptive and Nubar Information*****

MT=452 Total Nubar. Sum of MT=455 and 456.

MT=455 Delayed Neutron Yields. England (En89).

MT=456 Prompt Neutron Yields. From 10⁻⁵ to 650 eV, based on the evaluation of Fort (Fo88), after minor renormalization for consistency with CSEWG standards. Fort's values were multiplied by 1.000411 from 10⁻⁵ to 10 eV to fall within 1/2 std. deviation of Pu239 thermal nubar value, with the factor varying linearly to 1.00282 at 500 eV for consistency with Cf252 nubar. Fort's eval. (renormalized) was used intact below 62.3 eV but higher energy data were thinned with a thinning criterion of 0.036%, a factor of 5 less than the standard deviation of the CSEWG standard value of Pu239 thermal nubar. Above 650 eV, the evaluation is based on a covariance analysis of all available exp. data from the CSISRS data file at the NNDC, BNL, using the GLUCS analysis code (He80). A smooth curve was passed through the GLUCS results with structure removed. All the exp. data were renormalized to ENDF/B-VI standards prior to the covariance analysis.

*****MF=3 Smooth Cross Sections*****

MT=1 Neutron Total Cross Section. 0.03 to 20 MeV, based on coupled-channel optical calculations and the exp. data of Po81, Sh78, Po83, Sc74, Fo71, Sm73, Na73, Pe60, Ca73, Li90. The exp. and theoretical results were combined through a covariance analysis with the GLUCS code system (He80). The covariance analysis results, which agreed well with Derrien's (De89) unresolved resonance analysis at 30 keV, were smoothly joined to those results between 30 and 50 keV.

MT=2 0.030 to 20 MeV, based on subtraction of MT=4,16,17,18,37, and 102 from MT=1.

MT=4 Sum of MT=51-91

MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc.

MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc.

MT=18 The Pu239(n,f) cross section that resulted from the simultaneous standards analysis for Vers.VI was renormalized by a factor of 1.007 and used directly from 50 keV to 1.25 MeV. Between 30 and 50 keV, the evaluation was matched smoothly to the Derrien

unresolved resonance parameter analysis, requiring a reduction of about 4% near 30 keV. Above 1.25 MeV, the evaluation is based on a new covariance analysis of all available ratio and absolute Pu239(n,f) data, including new measurements (Li88,Me88) that were not available for the simul. standards analysis. The GLUCS code was used for the covariance analysis, and a smooth curve was drawn through the analysis results. The new covariance analysis agrees reasonably with the simultaneous standards results (both cross section and standard deviation) at energies below the newer exp. results. At higher energies, the effect of the new data on the analysis is to raise the (n,f) cross section somewhat at higher energies, particularly near 9 MeV (about 4%) and above 15 MeV (few %).

- MT=19 (n,f) first-chance fission cross section.
Ratio of first-chance to total fission obtained from GNASH calculations.
- MT=20 (n,nf) second-chance fission cross section.
Ratio of second-chance to total fission obtained from GNASH calculations.
- MT=21 (n,2nf) third-chance fission cross section.
Ratio of third-chance to total fission obtained from GNASH calculations.
- MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calc.
- MT=38 (n,3nf) fourth-chance fission cross section.
Ratio of fourth-chance to total fission obtained from GNASH calculations.
- MT=51-55,57 Thres. to 20 MeV, coupled-channel optical model calculations ($3/2^+$ to $13/2^+$ members of the $K=1/2$ ground state rotational band) using the ECIS code. Compound nucleus contributions, obtained from COMNUC calculations, are also included.
- MT=56,58-69,71,72,74-77 Threshold to 6.0 MeV, Compound nucleus reaction theory calculation with width fluctuations, using the COMNUC code.
- MT=70,73,78-81 Threshold to 20 MeV, distorted wave Born approximation calculations with the DWUCK code for $l=2$ and $l=3$ vibrational states. The $l=2$ states are MT=78,80 and the $l=3$ states are MT=70,73,79, and 81. Compound nucleus contributions were included in the data for MT = 70 and 73.
- MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc.
Note that the MT=78-81 vibrational states lie in the MT=91 continuum region.
- MT=102 0.030-20 MeV, obtained using the values of alpha (ratio of capture to fission cross sections) from ENDF/B-V.2, together with MT=18 from the present evaluation. Between

40 and 100 MeV, structure in the cross section was smoothed out. Below 50 keV, the results were smoothly joined to the unresolved res. result at 30 keV, requiring an increase of 0.2% at 30 keV.

*****MF=4 Neutron Angular Distributions*****

MT=2 Elastic scattering angular distribution based on ECIS coupled-channel calculations, with a compound elastic component from COMNUC included below 6 MeV.
MT=51-55,57 Thres. to 20 MeV, Coupled-channel optical model calculations plus compound-nucleus contributions.
MT=56,58-69,71,72,74-77 Threshold to 6.0 MeV, Compound nucleus reaction theory calculation with width fluctuations, using the COMNUC code.
MT=70,73,78-81 Thres. to 20 MeV, Distorted wave Born approximation calculations with DWUCK code. Compound-nucleus contributions are included for MT=70 and 73.

*****MF=5 Neutron Energy Distributions*****

MF=18 Composite neutron energy distributions from fission. Based on calculations by D.Madland (Ar84) using Madland-Nix formalism. The calculations include the first-, second-, and third-chance fission neutron components. These data are the same as were used for Revision 2 of ENDF/B-V. Parameters for the calculation were adjusted to give the same average fission neutron energy at thermal as ENDF/B-V.0. Tabulated data (LF=1) used.
MT=455 Tal England (En89).

*****MF=6 Correlated Energy-Angle Distributions*****

MT=16 GNASH Hauser-Feshbach statistical/preequilibrium calc. Updated Kalbach-Mann systematics used for specifying neutron distributions (Ka87). Only neutrons given.
MT=17 GNASH Hauser-Feshbach statistical/preequilibrium calc. Updated Kalbach-Mann systematics used for specifying neutron distributions (Ka87). Only neutrons given.
MT=37 GNASH Hauser-Feshbach statistical/preequilibrium calc. Updated Kalbach-Mann systematics used for specifying neutron distributions (Ka87). Only neutrons given.
MT=91 GNASH Hauser-Feshbach statistical/preequilibrium calc. Updated Kalbach-Mann systematics used for specifying neutron distributions (Ka87). Only neutrons given.

*****MF=12,13,14,15 Photon-Production Data*****

All photon-production data were carried over from ENDF/B-V.2, MAT=1399. Data are given for MF=12, MT=4,18,102; MF=13, MT=3; MF=14, MT=3,4,18,102; and, MF=15, MT=3,4,18,102.

*****References*****

- Ar84 E.Arthur et al., Nuc.Sci.Eng.88,56(1984).
- Ar88 E.D.Arthur, LA-UR-88-382 (1988).
- Ca73 J.Cabe et al., CEA-R-4524 (1973).
- Ca85 A.Carlson et al., Nuc.Data for Basic & Applied Science, Santa Fe, NM (1985) p.1429.
- De89 H.Derrien & G.de Saussure, ORNL-10986 (1989).
- Du70 C.L.Dunford, AI-AEC-12931 (1970).
- En89 T.R.England et al,LA11151-MS(1988),LA-11534-T(1989); LAUR-88-4118 to be published in NSE(1989).
- Fo71 D.Foster & D.Glasgow, Phys.Rev.C3,576(1971).
- Fo88 E.Fort et al., Nuc.Sci.Eng.99,375(1988).
- Fr86 J.Frehaut, NEANDC(E) 238/L (1986).
- He80 D.Hetrick & C.Y.Fu, ORNL/TM-7341 (1980).
- Ka87 C.Kalbach, LA-UR-87-4139 (1987) to be pub.in Phys.Rev.C!
- Ku70 P.D.Kunz, DWUCK: A Distorted-Wave Born Approximation Program, unpublished report.
- Li88 P.Lisowski et al., Nuc.Data for Sci.& Tech.,Mito Conf.,p97.
- Li90 P.Lisowski, Pers.Communication of WNR data taken in 1985.
- Me88 J.Meadows, Ann.Nucl.Energy 15, 421 (1988).
- Na73 K.Nadolny et al., USNDC-9 (1973)p.170
- Pe60 J.Peterson et al., Phys.Rev.120, 521(1960).
- Pe69 R.J.Peterson, Ann.Phys. 53, 40 (1969).
- Po81 W.Poenitz et al., Nuc.Sci.Eng.78, 333(1981).
- Po83 W.Poenitz et al., ANL-NDM-80, 1983.
- Ra70 J.Raynal,IAEA SMR-9/8 (1970).
- Sc74 R.Schwartz et al., Nuc.Sci.Engr.54,322(1974).
- Sh78 R.Shamu et al., personal communication, 1978.
- Sm73 A.Smith et al., J.Nuc.En.27,317(1973).
- Yo77 P.G.Young & E.D.Arthur, LA-6947 (1977).
- Yo88 P.G.Young & E.D.Arthur, Nuc.Data for Sci.& Tech., Mito Conference (1988) p.603.

²⁴⁰₉₄Pu

Reference: ORNL/TM-10386, ENDF-243 (1987)
Evaluators: L. W. Weston, E. D. Arthur, Others
Evaluated: August 1986
Material: 9440
Content: Neutron transport, Gamma production, Covariances

File Comment

²⁴⁰Pu revised for ENDF/B-VI by L. W. Weston.

MF=1 MT=452 Delayed nubar unchanged from ENDF/B-IV, Ref 1, 2, and 3. The prompt ratios to Cf are from Frehaut, Ref.4. The same value as used in version V. The nubar prompt relative to Cf assumed as 3.741.

MT=456 See MT=452.

MT=458 Energy release/fission from Sher/83 Ref. 5.

MF=2 MT=151 Resolved region 0.0. to 5.7 keV. The resolved resonance region extends to zero energy. The room temperature cross sections at 0.0253 ev are:

total	=	288.6 barns
scattering	=	0.96 barns
fission	=	0.064 barns
capture	=	287.6 barns

The resonance at 1.056 eV was evaluated from Ref. 6, 7 and 8. From 20 to 665 eV the neutron and radiation widths are weighted averages of Weigmann et al., Ref 9, Moxon, Ref 10, and Hockenbury, Ref 11, taken from a Weigmann review, Ref 9 and unchanged from version V. The neutron widths above 665 eV are from Ref 9 and extended to 5.7 keV for version VI. The fission widths are revised over the full energy region. They are taken from Weston, Ref 12, Migneco, Ref 13, and Auchampaugh, Ref 14, as a weighted average of values. The unresolved region extends from 5.70 to 40 keV. The parameters are from FITACS, Ref 15, and URES, Ref 16, and fits to Hockenbury, Ref 11, Weston, Ref 17, and Wisshak, Ref 18, average capture.

MF=3	MT=1	Background cross sections 2680 - 5700 eV due to missed resonances. A FITACS fit to Gwin, Ref 19, and Poenitz, Ref 20, from 5.7 to 500 keV. Above 500 keV from Lagrange, Ref. 21.
	MT=2	Difference between MT=1 and sum of other cross sections.
	MT=4,	51 ... 59, 91 Inelastic scattering is from model calculations of Lagrange and Jary, Ref 21, which were adapted for use in ENDF/B by E. D. Arthur, LANL.
	MT=16,	17 (n,2n) and (n,3n) taken from Lagrange, Ref 21, and corrected after phase 1 review by the Chinese 11/89.
	MT=18	No smooth fission background in the resonance region. From 40 to 100 keV from Weston, Ref 12 and Knitter, Ref 22. From 100 keV to 1 MeV from a ratio to ^{235}U by Behrens, Ref 23, and same as version V, in agreement with Weston, Ref 24, and Kari, Ref 25. From 1 to 20 MeV it is a compromise of Behrens, Ref 23, Gilboy, Ref 26, Savin, Ref 27, Weston, Ref 24, Kari, Ref 25, and Meadows, Ref 28.
	MT=102	Background from 2680 - 5700 eV due to missed resonances. No smooth cross sections in the unresolved range. The evaluation is based on Hockenbury, Ref 11, Weston, Ref 17, and Wisshak, Ref 18, from 40 keV to 300 keV. ENDF/B-IV is used above 300 keV.
MF=4	MT=2,	16-18, 51-69, and 91 Angular distributions are from the ENDF/B-V evaluation of ^{242}Pu by Madland and Young, Ref 29.
MF=5	MT=455	Delayed neutron secondary energy distribution from M. C. Brady & T. R. England, Nucl. Sci. & Eng. 103, 129 (1989). See also T. R. England et al., LA-11151-MS(1988), LA-11534-T(1989), and LAURA-88-4118.
MF=12		13, 14, and 15 The γ -ray files were evaluated by Hunter and Stewart (LANL) in 1972 and are described in LA-4901. These data were input in version III and have survived through versions IV and V. They have not been changed for version VI. Below 1.09 MeV, however, gamma ray multiplicities were used in MF=12 for radiative capture, inelastic scattering and fission.

MF=12 Continued... Therefore current improvements in this energy range are reflected in the γ -ray production cross sections calculated for version VI. MF=12 includes MT=4 contributions for discrete gammas from inelastic scattering. Above 1.09 MeV, MF=13 is used throughout. All γ rays are assumed isotropic.

Note: The uncertainty files are unchanged from Rev. 2 of version V.

MF=32 MT=151 The resonance parameter error file extends from 0.5 to 105 eV. The errors are based on difference in measurements because the quoted errors are not consistent in Ref 9, 10 and 11. The parameters of the 1 eV resonance are highly correlated.

MF=33 MT=18 From revision to ENDF/B-V, Dec. 1982, by L. W. Weston, Ref 30. LB=8 was added 11/89.

MT=102 See MT=18. LB=8 was added 11/89.

References:

1. S. A. Cox, "Delayed Neutron Data-review and Evaluation," ANL/NDM-5, Argonne National Laboratory (1974).
2. R. E. Hunter, L. Stewart and T. J. Hirons LA-5172 (June,1973).
3. R. E. Hunter and L. Stewart LA-4901 (July,1972).
4. J. Frehaut et al., Proc. Second All-Union Conference on Neutron Physics, Part 3, 153 (1974).
5. R. Sher and C. Beck, EPRI NP-1771/81 + Rev 1/83 + Personal Communication to B. A. Magurno 2/83.
6. M. Lounsbury, R. W. Durham and G. C. Hanna, "Measurements of α and of Fission Cross Section Ratios for ^{233}U , ^{235}U and ^{239}Pu at Thermal Energies," p.287, Nucl. Data for Reactors, IAEA Conference (1970).
7. H. I. Liou and R. E. Chrien, "Neutron Cross Sections and Doppler Effect of the 1.056 eV Resonance in ^{240}Pu ," IAEA Consultants Meeting on U and Pu Isotope Resonance Parameters, September 28, 1981, INDC(NDS)-129, Vienna p.438.
8. R. R. Spencer et al., "Neutron Total Cross Sections of ^{240}Pu Below 6 eV and the Parameters of the 1.056 eV Resonance," Proceedings Intl. Conf. on

Nuclear Data for Basic and Applied Science, May 13 - 17, 1985, Santa Fe.
Gordon & Breach, New York p.581.

9. H. Weigmann and J. P. Theobald, J. Nucl. Energy 26, 643 (1972). Also W. Kolar and K. H. Bockhoff, J. Nucl. Energy 22, 299 (1968). Also H. Weigmann, G. Rohr, and F. Poortmans, Proc. Conference Resonance Parameters of Fertile Nuclei and ^{239}Pu , Saclay, 219, NEANDC(E) 163U (1974).
10. M. Moxon, See Ref 9.
11. R. W. Hockenbury, W. R. Moyer, and R. C. Block, Nucl. Sci. & Eng. 49, 153 (1972).
12. L. W. Weston and J. H. Todd, Nucl. Sci. & Eng. 88, 567 (1984).
13. E. Migneco and J. P. Theobald, Nucl. Phys. A112, 603 (1968).
14. G. F. Auchampaugh and L. W. Weston, Phys. Rev. C12, 1850 (1975).
15. F. H. Froehner, Kernforschungszentrum, Karlsruhe, Private Communication (1982).
16. E. M. Pennington, Argonne National Laboratory, Private Communication (1973).
17. L. W. Weston and J. H. Todd, Nucl. Sci. & Eng. 63, 143 (1977).
18. K. Wisshak and F. Kaeppler, Nucl. Sci. & Eng. 66, 363 (1978).
19. R. Gwin, Oak Ridge National Laboratory, Private Communication (1985).
20. W. P. Poenitz, J. P. Whalen and A. B. Smith, Nucl. Sci. & Eng. 78, 333 (1981).
21. C. H. Lagrange and J. Jary, "Coherent Optical and Statistical Model Calculations of Neutron Cross Sections for ^{210}Pu and ^{212}Pu Between 10 keV and 20 MeV," NEANDC(E) 198 "I", and INDC(Fr) 30/1, Bruyeres-le-Chatel, France (1978).
22. C. Budtz-Jorgensen and H. H. Knitter, Nucl. Sci. & Eng. 79, 380 (1981).
23. J. W. Behrens, J. C. Browne and G. W. Carlson, UCID-17047, III (1976).
24. L. W. Weston and J. H. Todd, Nucl. Sci. & Eng. 84, 248 (1983). Reactor Applications, BNL-50991, Brookhaven National Laboratory (1979).
26. W. B. Gilboy and G. Knoll, KFK-450, Kernforschungszentrum (1966).
27. M. V. Savin et al., INDC(CCP)-8/U (1970).

28. J. W. Meadows, Nucl. Sci. & Eng. 79, 223 (1981).
29. D. G. Madland and P. G. Young, "Evaluation of $n + {}^{242}\text{Pu}$ Reaction from 10 keV to 20 MeV," Proc. Nuclear Data for Pu and Am for Reactor Applications, p.189, BNL-50991 (1978).
30. L. W. Weston Personal Communication to B. A. Magurno Nov.12, 1982.

Fig. 1. Example of the subthreshold fission cross section as defined by the resolved resonance parameters. Points are data from Weston and Todd¹².

Fig. 2. Fit and evaluation of the total cross section from 4 to 500 keV. The dashed line is ENDF/B-V. The circles are the data of Gwin and the pluses are Poenitz et al.²⁰.

Fig. 3. Fit and evaluation of the capture cross section of ^{240}Pu (solid line). The circles are the data of Weston and Todd¹⁷ and the pluses and triangles are the data of Wissihak and Kaeppler.¹⁸ The dashed line is the ENDF/B-V evaluation. The cusp at 43 keV is due to the first inelastic scattering level.

Fig. 4. The evaluation of the fission cross section of ^{240}Pu from 40 keV to 20 MeV as compared to ENDF/B-7 (dashed line).

Fig. 5. The evaluation (solid line) of the total cross section from 0.1 to 20 MeV as compared to ENDF/B-V (dashed line). The pluses are the data of Gwin¹⁹ and the circles are the data of Poenitz et al.²⁰

Fig. 6. The evaluation (solid line) of the total inelastic scattering and the inelastic scattering from the first two levels for ^{240}Pu . The dashed lines are the corresponding ENDF/B-V evaluations.

Fig. 7. The accepted evaluation (solid line) by Lagrange and Jary²¹ of the (n,2n) and (n,3n) cross sections for ^{240}Pu . The dashed line is the corresponding ENDF/B-V evaluation.

Reference: No Primary Reference
Evaluators: L. W. Weston, R. Q. Wright, H. Derrien, Others
Evaluated: October 1988
Material: 9443
Content: Neutron transport, Gamma production, Covariances

1. Introduction

^{241}Pu was revised for ENDF/B-VI by L. W. Weston. The changes were the adoption of the recommended values of the standards committee for the thermal constants; the adoption of the Reich-Moore resonance parameter evaluation of the resolved resonance region by H. Derrien of Cadarache, France; renormalization of capture above 300 eV by 14 % downward as recommended by H. Derrien. The totals above 100 keV were made the same as ^{239}Pu . The capture was revised downward above 1.5 MeV.

MF=1 MT=452 A thermal nubar of 2.9453 recommended by the Standards committee. Prompt ratios to Cf above .47 MeV from Frehaut, Ref. 1. Nubar prompt relative to Cf = 3.7676.
MT=455 Tabulation of ν delayed (Ref. 2) except at thermal.
MT=456 See MT=452.
MT=458 See Sher Ref. 3.

MF=2 MT=151 Resolved resonance region to 300 eV, using a Reich-Moore representation.

2. Status of the Resonance Region Evaluation, H. Derrien, March 1988

2.1 Thermal Range

The cross section values proposed by the ENDF/B-VI Standards Evaluation group have been used as reference at 0.0253 eV. ¹ All the experimental data considered in the present evaluation have been renormalized to this reference in the energy range 0.02 eV to 0.03 eV. The experimental data analysed are the total cross sections from Young et al. ⁵ and from Simpson et al. ⁶, the fission cross sections from Wagemans et al. ⁷ and from Weston et al. ⁸, and the capture cross sections from Weston et al. ⁸. The cross section values are given in the following table:

	ENDF/B-VI Proposed Standard at 0.0253 eV (barns)	Energy Range 0.02 - 0.03 eV ^a (barns)	Calculated 0 ^a K At 0.0253 eV ^b (barns)
Fission	1012.68 ± 6.58	1023.9	1011.74
Capture	361.29 ± 4.95	366.0	362.97
Scattering	12.17 ± 2.62		11.16
Total	1386.14 ± 8.64	1402.1	1385.88

^a Average cross-sections for the renormalization of the experimental data to the proposed standard values.

^b R-matrix calculation from the evaluated data set of resonance parameters using the program NJOY-87.0. The calculated values are in very good agreement with the proposed standard at 0.0253 eV.

A correlated R-matrix fit⁹ was performed on Young total, Simpson total and Wagemans fission in the energy range 0.001 to 3 eV. The renormalized Seppi fission data¹⁰ were also considered as a possible reference point in the energy range 0.002 to 0.005 eV. The cross-sections integrated over the 0.27 eV resonance, in the energy range 0.02 eV to 0.45 eV, are shown on the following table:

	Cross Section Experimental (barn . eV)	Cross Section Calculated (barn . eV)	Deviation %
Young total	455.01	456.19	+0.3
Simpson total	458.97	459.12	+0.0
Wagemans fission	326.03	327.79	+0.5
Weston fission	334.18	327.80	- 1.9
Weston capture	137.43	126.49	- 8.6

The difference between the Young and Simpson total data is nearly the same on the experimental and on the calculated data; it is due to a different experimental resolution function. There is a severe discrepancy on the Weston capture data which was already discussed by Weston et al. in the ENDF/V-V evaluation.¹¹ The shape of the Weston fission cross section in the energy range below 0.03 eV differs from that of Wagemans. Normalizing the Weston and Wagemans data in the energy range

0.02 to 0.03 eV results in a discrepancy of about 2% over the 0.26 eV resonance. The same remark applies to the Weston absorption data which should be normalized to the Simpson or Young absorption data over an energy range including the 0.26 eV resonance.

2.2 The Resolved Resonance Range

The Harvey-Simpson transmission data¹² which were obtained in 1972 from sample cooled down to nitrogen temperature were analysed in the energy range 0.3 to 300 eV along with Blons¹³, Migneco¹¹ and Weston⁸ fission data. A correlated SAMMY Reich-Moore fit was performed with 50 or 100 eV range correlation matrices. Fictitious resonances (negative energy resonances and resonances above 300 eV) were used to take into account the effect of the external range in such way that the cross sections in the range thermal to 300 eV could be reproduced by the set of resonance parameters without the use of a file 3. A radius $r' = 9.50$ fermi was used.

A comparison of all the available fission data in the resolved energy range was made by Weston et al.⁸ showing that large discrepancies exist among the data. These discrepancies could be due to back-ground correction effects or other experimental effects leading to local normalization errors. The SAMMY fits have been performed to take into account the eventual local renormalization and back-ground corrections. The set of resonance parameters obtained is expected to represent the cross sections with about 5% accuracy at least for the fission cross-sections. Comparisons between calculated and experimental integrated fission cross-sections are shown on the following table:

Energy range eV	Cross sections (barn-eV)		
	Average ^a	Weston	Calculated (NJOY 300 ° K)
3.0- 4.9	350.8	370.6	359.5
4.9- 8.0	876.8	891.7	836.1
8.0- 9.0	237.6	243.8	234.7
9.0- 12.0	311.1	321.4	292.0
12.0- 14.0	284.8	286.3	279.6
14.0- 17.4	928.0	953.1	922.3
17.4- 20.0	143.8	146.2	139.9
20.0- 30.0	795.3	866.1	843.1
30.0- 40.0	452.0	492.7	487.0
40.0- 50.0	390.9	436.6	408.6

Energy Range eV	Cross sections (barn-eV)		
	Average ^a	Weston	Calculated (NJOY 300 °K)
50.0- 60.0	173.1	176.1	175.4
60.0- 70.0	559.1	591.1	574.2
70.0- 80.0	278.3	270.4	262.6
80.0- 90.0	685.9	733.1	730.7
90.0-100.0	279.2	278.5	285.5
100.0-200.0	2660.0	2686.5	2626.4
200.0-300.0	2780.0	2861.1	2827.2

Energy range eV	Cross sections (barns)		
	Average ^a	Weston	Calculated
3.0-300.0	41.03	42.44	41.36

^a Average over all available fission data. See reference (8).

The cross sections calculated by the set of resonance parameters are in good agreement with the average data from all the available experimental values. The agreement is even perfect if one considers the average values over the entire energy range. The 3% difference observed in the Weston data should go away after a renormalization over the 0.26 eV resonance in Wagemans data.

The following table shows the calculated capture cross-sections and alpha values compared to Westons⁸ experimental data:

Energy range eV	Calculated capture (NJOY 300 °K)	Calculated alpha	Experimental alpha	Deviation %
10.- 20.	74.88	0.511	0.559	9.4
20.- 30.	15.48	0.184	0.213	15.7
30.- 40.	9.86	0.202	0.216	6.9
40.- 50.	5.38	0.132	0.184	39.4
50.- 60.	2.15	0.123	0.198	61.0
60.- 70.	12.94	0.225	0.279	24.0
70.- 80.	14.26	0.543	0.572	5.3

Energy range eV	Calculated capture (NJOY 300 °K)	Calculated alpha	Experimental alpha	Deviation %
80.- 90.	23.39	0.320	0.337	5.3
90.-100.	5.76	0.202	0.207	2.5
100.-200.	5.81	0.221	0.268	21.3
200.-300.	6.60	0.233	0.264	13.3
10.-300.		0.244	0.279	14.3

On average the alpha values calculated from the resonance parameters are about 14% smaller than the experimental values given by Weston.⁸ A tentative SAMMY fit was performed on the Weston capture in the energy range 3 eV to 20 eV only, in correlation with the fission experimental data and the transmission data. A renormalization of 15% for the capture was obtained in this energy range. The absorption experimental data were normalized by Weston to the absorption obtained from Kolar total cross sections.¹⁵ The present evaluation, which was mainly based on the Harvey-Simpson transmission data for the determination of the neutron widths, gives widths which are 8% on average smaller than those obtained from the Kolar transmission data. Renormalizing the Weston absorption data to values calculated from the new set of neutron widths should give more realistic values of the experimental absorption cross sections.

The unresolved resonance parameters start at 300 eV. They are the same as ENDF/B-V except that the capture has been reduced by 14%.

3. Remainder of File

MF=3	MT=1	Totals were adjusted to compensate changes in capture.
	MT=2	Unchanged from ENDF/B-IV.
	MT=18	The evaluation uses ENDF/B-V above 40.2 keV and is based upon Behrens, Ref.16, Kappelar, Ref. 17, Szabo, Ref. 18, and ratios to ²³⁵ U using the ENDF/B-V evaluation of ²³⁵ U.
	MT=102	Renormalized downward by 14% from version V.
MF=5	MT=16	Revised by R. J. Howerton for version V.
	MT=17	Revised by R. J. Howerton for version V.

MF=5 MT=18 The fission neutron energy distribution is given as a simple fission spectrum plus a Maxwellian (AWRE O-101/64)

MT=91 Revised by R. J. Howerton for version V.

MT=455 Delayed neutron secondary energy distributions. See ref. 13.

MF=12 MT=18 Used data of Peele and Maienschein, Ref 19, for ^{235}U thermal fission.

MT=102 Used ^{238}U spectrum adjusted for multiplicity and energy conservation.

MF=13 MT=3 Calculated by R. J. Howerton.

MF=15 MT=3 Calculated by R. J. Howerton.

MT=18 Calculated by R. J. Howerton. Used Peele and Maienschein data, Ref 19, and methods described in Ref 20.

MT=102 Calculated by R. J. Howerton.

MF=33 MT=18, 102 Replaced by L. Weston in December 1982. Ref. 21.

References

1. J. Frehaut et al., CEA-R-4626 (1974).
2. M. C. Brady and T. R. England, Nucl. Sci. & Eng. 103, 129 (1989). See also T. R. England et al., LA-11151-MS (1989), LA-11534-T (1989) and LAURA-88-4118 (1988).
3. R. Sher and C. Beck, EPRI NP-1771/81 + Rev. 1/83.
4. Proposed data by the ENDF/B-VI Standards Working Group.
5. T. B. Young and J. R. Smith, WASH-1093 60.
6. O. D. Simpson and R. P. Shuman, Nucl. Sci. & Eng. 11, 111 (1961).
7. C. Wagemans and A. J. Deruyter, Nucl. Sci. & Eng. 60, 1 (1976).
8. L. W. Weston and J. H. Todd, Nucl. Sci. & Eng. 65, 454 (1978) and Nucl. Sci. & Eng. 68, 125 (1978).
9. H. Derrien and G. DeSaussure, "R-Matrix Analysis of the ^{241}Pu Neutron Cross Sections in the Energy Range Thermal to 300 eV," ORNL/TM-11123 (1989).

10. E. J. Seppi et al., HW-55879 (1958).
11. L. W. Weston and R. Q. Wright, NBS Special Publication 594, p. 464 (1980).
12. J. Harvey and O. D. Simpson, Unpublished (1972).
13. J. Blons and H. Derrien, Journal de Physique 37, 659 (1976).
14. E. Migneco, Helsinki (1970). Vol. 1, page 437.
15. W. Kolar and G. Carraro, Proc. Third Conference on Neutron Cross Sections & Technology, Knoxville, Tennessee, CONF-710301, Vol 2, p. 707 USAEC (1971).
16. J. W. Behrens and G. W. Carlson, UCRL-51925 (1975).
17. F. Kappelar and E. Pfletschinger, Nucl. Sci. & Eng., 51, 124 (1973).
18. I. Szabo et al., Symp. Neutron Standards, Argonne National Laboratory 257 (1970).
19. R. W. Peele and F. C. Maienschein, Nucl. Sci. & Eng. 40, 485 (1970).
20. S. T. Perkins, R. C. Haight, and R. J. Howerton, Nucl. Sci. & Eng. 57, 1 (1975).
21. L. W. Weston Personal Communication to B. A. Magurno, November (1982).

²⁴¹₉₅Am

Reference: No Primary Reference
Evaluators: Zhou Delin, Gu Fuhua and Others
Evaluated: February 1988
Material: 9543
Content: Neutron transport, Gamma production, Covariances

Summary of ENDF/B-VI Evaluation

History

The original evaluation was performed by Gu Fuhua, Yu Baosheng, Zhou Delin, Zhuang Youxiang, Shi Xiangjun, Yan Shiwei, Wang Cuilan, and Zhang Jengshang under contracts with the IAEA/NDS and the IAE, 1985.¹ The file was revised by Zhou Delin, Yu Baosheng, Liu Tong, Shi Xiangjun, and Yan Shiwei in 1988.² The capture cross sections and resonance parameters have been reviewed and reevaluated by Zhou Delin et al. 1988.³

File Information

MF=1	General information.
MT=451	Comments and Dictionary.
MT=452	Total number of neutrons per fission. It is the sum of the delayed neutrons (MT = 455) and prompt neutrons per fission (MT = 456).
MT=455	T. R. England +, LA-11151, LA-11534, and LAURA-88-4118.
MT=456	Number of prompt neutrons per fission. Data are taken from Y. Kikuchi et al. ¹
MF=2 MT=151	Resonance Parameters. The SLBW formula was used for resolved resonance parameters in the 1.0×10^{-5} to 150 eV energy region. Resolved parameters were recommended from the analysis of several data sets.

File Information, Continued

It was noticed that the parameters of the first several well defined resonances (0.308, 0.576, 1.276, 1.928, 2.372, and 2.358 eV resonances for example) given by Derrien et al.⁵ (Transmission), Kalebin et al.⁶ (Transmission) as well as Weston et al.⁷ (Absorption, normalized to a thermal capture cross section of 582 b) respectively are in excellent agreement with each other. This means that the resonance parameters for the full resonance energy region can be combined by using random error weight averaging without taking into account the absorption measurement with 8% systematic error.⁷ The recommended values for most resonances were averaged using equal weights. An exception was for the random errors of Weston et al.'s measurements and the errors of Kalebin et al.'s measurements which are much larger than those of Derrien et al. In such cases the average was properly made with unequal weights.

Unresolved resonance parameters were defined in the 150 eV to 30 keV energy region. Kikuch's parameters¹ were used as input data and adjusted to fit the capture cross sections (evaluated on the bases of Weston et al., Vanpraet et al.⁸, Wisshak et al.⁹, and Derrien et al.'s measurements) and the fission cross section of Dabbs¹⁰ (averaged over the proper energy range). Recommended unresolved resonance parameters were obtained.

At thermal energy a value of 3.15 b was adopted for the fission cross section, and a capture cross section 620 ± 13 b was adopted. A weighted averaged value of the following data was adopted:

- Kalebin: 625 ± 20 b. ($= 640 \text{ b} - 3.15 \text{ b} - 11.5 \text{ b}$, Transmission);
- Dovbenko et al.: 654 ± 104 b. (Capture)¹¹;
- Harbour et al.: 612 ± 25 b. (Capture, deduced from an analysis by Story, 1978. Quoted from Lynn et al.)¹²;
- Pomerance: 625 ± 35 b. (Quoted from Lynn et al.)¹²;
- Wisshak et al.: 625 ± 35 b. (Isomeric ratio, indirectly)¹³;

File Information, Continued

- Weston et al.: 582 ± 50 b. (A relative measurement of absorption normalized to a thermal capture cross section of 582 b. It may be considered to be normalized equivalently to Derrien's measurement of the first several resonances. The 8% systematic uncertainty in the energy region > 0.2 eV must be added to the thermal value)³.
- MF=3 Neutron Reaction Cross Sections.
 - MT=1, 2, 4, 16, 17, 51-65, 91, & 102 Total, elastic scattering, inelastic scattering, (n,2n), (n,3n) and capture cross sections were calculated using optical model theory. (The parameters have been adjusted to fit the Phillips' data.¹⁴) Hauser-Feshbach statistical theory with width fluctuation corrections, exciton, and evaporation models were used.¹
 - MT=18 Fission cross sections in the energy region greater than 1 keV were evaluated by Gu Fuhua et al.¹ and revised by Zhou Delin considering Dabbs' new measurement. Below 1 keV Dabbs' measurement has also been used for an unresolved resonance parameter adjustment.³
- MF=4 Angular distributions of secondary neutrons.
 - MT=2, 51-65 Angular distributions of elastic and inelastic scattering were calculated using optical models and Hauser-Feshbach theory.
 - MT=16, 17, 18, and 91 Angular distributions of (n,2n), (n,3n), fission, and inelastic scattering (continuum part) reactions were assumed isotropic in the Center of Mass (for 16, 17 & 18) and Laboratory frame (for 91).
- MF=5 Energy distributions of secondary neutrons.
 - MT=16, 17, 18, and 91 Calculated with Hauser-Feshbach and evaporation models taking preequilibrium processes into account. For MT=18 a maxwellian distribution was specified. The parameters $T(E)$ are estimated from Hu Jimin et al.¹⁵

File Information, Concluded

- MF=9 MT=102 Data for the isomeric ratio has been revised based on a new evaluation² and the shape of a theoretical calculation.¹³
- MF=12 13, 14, and 15 Photon data calculated by R. J. Howerton (Personal Communication) From MF=2, 3, and 5.
- MF=32 33 Covariances. File 32 and 33 of ENDF/B-V, Rev. 2 have been adopted. However, the covariances of MT=151 of File 32 and the covariances of MT=18 of file 33 have been modified to match the present evaluations.

References

1. Gu Fuhua et al., "Neutron Data Evaluation for ²⁴¹Am," 1985. (Unpublished).
2. Thou Delin et al., "Neutron Data Evaluation for ²⁴¹Am," 1988. (Unpublished).
3. Zhou Delin, "On the Capture Data Evaluation for ²⁴¹Am," 1988. (To be Published).
4. Kikuch Y., JAERI-M 82-096, 1982.
5. Derrien H. et al., WASH 75, p.637, 1975.
6. Kalebin S. et al., AE 40, 373 (1976).
7. Weston L. et al., Nucl. Sci. Eng. 61, 356 (1976).
8. Vanpraet et al., Santa Fe (1985), p.493, 1985.
9. Wisshak K. et al., Nucl. Sci. Eng., 76, 148 (1980).
10. Dabbs J. et al., Nucl. Sci. Eng., 83, 22 (1983).
11. Dovbenko A. et al., LA-TR-71-74, 1971.
12. Lynn J. et al., "Progress in Nuclear Energy," Vol 5, p.255, 1980.
Harbour R. et al., Nucl. Sci. Eng., 50, 364 (1973).
Pomerance H. ORNL-1879, p.50, 1955.
13. Wisshak K. et al., Nucl. Sci. Eng., 81, 396 (1982).

²⁴³₉₅Am

Reference: No Primary Reference
Evaluators: L. W. Weston, F. M. Mann, R. E. Schenter, R. J. Howerton, Others
Evaluated: October 1988
Material: 9549
Content: Neutron transport, Gamma production

File Comments

ORNL	Eval-Oct88 L. W. Weston
HEDL	Eval-Apr78 F. M. Mann and R. E. Schenter (fast)
LLNL	Eval-Apr78 R. J. Howerton (gamma production)

Summary of Evaluation

MF=1	General information
MT=452	Nubar. The thermal value was computed from the semi-empirical work of Gordeeva and Smirenkin (Ref. 1) as revised by Manero and Konshin (Ref. 2). The energy dependence is from the work work of R. J. Howerton (Ref. 3).
MT=455	Delayed neutron yields. Taken from M. C. Brady (ORNL) and T. R. England, Nucl. Sci. & Eng. <u>103</u> , 129 (1989).
MT=458	Energy from fission based on Sher (Ref. 4).
MF=2	Resonance parameters (0 to 42 keV)
MT=151	Resolved resonances. Two hundred and nineteen resolved resonances plus one bound level are included based upon the total cross section measurements of Simpson et al. (Ref. 5). The thermal capture is based on Mughabghab's evaluation (Ref. 6). The thermal fission cross section is based on Wagemans measurement (Ref. 7).

Summary of Evaluation, Continued

MF=2	MT=151	At thermal energy the total cross section is 84 barns, the capture cross section is 75 barns and the fission cross section is 74 millibarns. Resolved region - 0 to 250 eV. The fission widths are based on Knitter's measurements (Ref. 8). The unresolved resonance parameters are based on the evaluation of Froehner et al. (Ref. 9) and Weston & Todd (Ref. 10). Unresolved region - 250 eV to 42 keV.
MF=3		Smooth cross sections (42 keV to 20 MeV).
	MT=1	Total. Sum of partial cross sections.
	MT=2	Elastic cross sections. The elastic cross sections are based upon optical model calculations (Ref. 11) above 0.65 MeV, and on average resonance parameters below.
	MT=4	Inelastic cross section. The inelastic cross section above 100 keV is based on statistical model calculations to 17 excited levels plus the continuum (Ref. 11).
	MT=16	(n,2n) based on statistical model calculations (Ref. 11).
	MT=17	(n,3n) based on statistical model calculations (Ref. 11).
	MT=18	Fission is based on the data of Seeger et al. (Ref. 12), Knitter (Ref. 8), Wisshak (Ref. 13), and Behrens (Ref. 14).
	MT=19	Same as MT=18 until (n,nf) threshold, after which the cross section is constant.
	MT=20	Is (MT=18) - (MT=19).
	MT=37	(n,4n) is based on statistical model calculations (Ref. 11).
	MT=51, 52... 67, and 91	Above 100 keV the evaluation is based on statistical model calculations to 17 excited levels plus the continuum (Ref. 11).
	MT=102	Capture based on FITACS (Ref. 15) fit to Wisshak (Ref. 13) and Weston (Ref. 10) for data below 200 keV. At energies above 200 keV the evaluation is based on statistical model calculations.
MF=4		Secondary neutron angular distributions.

Summary of Evaluation, Concluded

- MF=4 MT=2 The elastic angular distributions were supplied by H. Alter (Atomics International). They were composed of a mixture of measured data for ^{235}U , ^{238}U , and ^{239}Pu .
 MT=51, 52... 67, and 91 Assumed isotropic.
- MF=5 Secondary neutron energy distributions.
 MT=16 Based on parameters of Gilbert and Cameron (Ref. 16).
 MT=17 Same reference as MT=16.
 MT=18 The fission spectrum has a Maxwellian density with the temperature based on Terrell's prescription (Ref. 17). The thermal value of ν was used to determine the temperature.
 MT=19, 20 Same as MT=18.
 MT=37 91 Same reference as for MT=16.
 MT=455 Brady et al., Nucl. Sci. & Eng. 103, 129 (1989).
- MF=8 Radioactive decay data.
 MT=102 Decay data was used from ENDF/B-V MAT numbers 7544 and 7554.
 MT=454 A ^{241}Am yield curve was used (Ref. 13 and 19).
- MF=9 MT=102 Based on statistical model calculations (Ref. 11).
- MF=12, 13, 14, and 15 Photon production files taken from the evaluations of R. J. Howerton documented in UCRL 50400, Vol. 15, part A (methods) Sept 75 and part B (curves) Apr 76. The files were extended to the energy range 1.0×10^{-5} eV to 20 MeV and merged to this evaluation at the Brookhaven National Laboratory by R. Kinsey of the National Nuclear Data Center.

References

1. L. Gordeeva and G. Smirenkin, Sov. Atomic En. 14 (1963) 562.
2. F. Manero and V. Konshin, At. En. Rev. 10 (1972) 637.
3. R. Howerton, Nucl. Sci. Eng. 46 (1971) 42.
4. R. Sher and C. Beck, EPRI NP-1771/81 + Rev 1/83, also personal communication to B. A. Magurno (BNL) 2/83.

5. O. Simpson, F. Simpson, J. Harvey, G. Slaughter, R. Benjamin, and C. Ahlfeld, Nucl. Sci. Eng. 55 (1974) 273.
6. S. F. Mughabghab, Neutron Cross Sections, Vol 1, p 95-9, National Nuclear Data Center (1984).
7. C. Wagemans, P. Schillebeeckx and J. P. Bocquet, Nucl. Sci. Eng. 101 (1989) 276.
8. H. Knitter and C. Budtz-Jorgensen, Nucl. Sci. Eng. 99, (1988) 1.
9. F. H. Froehner, B. Goel, U. Fisher and H. Jahn, Proc. Nuc. Data for Sci. and Tech., p. 211, Antwerp, Belgium, 1982.
10. L. W. Weston and J. H. Todd, Nucl. Sci. Eng. 91, (1985) 444.
11. F. M. Mann and R. E. Schenter Trans. Am. Nuc. Soc. 23 (1976) 546, and HEDL TME-77-54 (1977).
12. P. A. Seeger, LA-4420 (1970).
13. K. Wisshak and F. Kaeppler, Nucl. Sci. Eng. 85, (1983) 251.
14. J. W. Behrens and J. C. Browne, Nucl. Sci. Eng. 77, (1981) 44.
15. Code by F. H. Froehner, Karlsruhe, Private Communication (1982).
16. A. Gilbert and A. G. W. Cameron, Can. J. Phys. 43 (1965) 1446.
17. J. Terrell, Phys. and Chem. of Fission, Vol 2, IAEA (1965).
18. J. G. Cuninghame, J. Inorg. Nucl. Chem. 4 (1957) 7.
19. R. R. Richard et. al., Trans. Am. Nuc. Soc. 6 (1963) 2.

Fig. 1. The total cross section of ^{243}Am from .002 to 20 MeV for ENDF/B-VI (solid line) and ENDF/B-V (dashed line).

Fig. 2. The capture cross section of ^{243}Am from 2 to 300 keV. The solid line is ENDF/B-VI and the dashed line is ENDF/B-V. Circles are the data of Weston and Todd¹⁰ and the other points are the data of Wisshak and Kaeppler.

Fig. 3. The fission cross section of ^{243}Am from .002 to 20 Mev. The solid line is ENDF/B-VI and the dashed line is ENDF/B-V. The circles are the data of Behrens and Browne¹⁴ and the pluses and histograms are the data of Knitter and Budtz-Jorgensen.⁸

Fig. 4. The fission cross section of ^{243}Am from 0 to 10 MeV. The solid line is ENDF/B-VI and the dashed line is ENDF/B-V. The circles are the data of Behrens and Browne¹⁴ and the pluses are the data of Knitter and Budtz-Jorgensen.⁸

²⁴⁹₉₇Bk

Reference: No Primary Reference
Evaluators: Zhou Delin and Others
Evaluated: June 1986
Material: 9752
Content: Neutron transport

Summary of ENDF/B-VI Evaluation

History

This evaluation was performed by Zhou Delin et al. under a contract between the IAEA/NDS and the CNDC. For a complete discussion of this evaluation refer to the report "Evaluation of Neutron Nuclear Data for ²⁴⁹Bk." ¹

File Information

MF=1 General information.
 MT=451 Comments and Dictionary.
 MT=452 Total number of neutrons per fission. It is the sum of
 delayed neutrons (MT=455) and prompt neutrons per
 fission (MT=456).
 MT=455 Delayed neutron data. From calculations made by
 Y. Kikuchi et al. ²
 MT=456 Number of prompt neutrons per fission. From calculations made by Qiu Xijun et al. ³

MF=2 MT=151 Resonance Parameters. The MLBW formalism was used for the resolved resonance parameters in the 1.0×10^{-5} to 60 eV energy range. Recommended resonance parameters were mainly based on the measurements of Benjamin et al. ⁴, and Anufriev et al. ⁵ Also consulted were the evaluations of Y. Kikuchi et al. ² and Mughabghab. ⁶ Unresolved resonance parameters were selected from parameters recommended by Y. Kikuchi et al. ²

File Information, Continued

MF=3	Neutron Reaction Cross Sections.
MT=1,	2, 4, 16, 17, 51-68, 91, and 102 Total, elastic scattering, inelastic scattering, (n,2n), (n,3n) and capture cross sections calculated using optical model theory. Hauser-Feshbach statistical theory with width fluctuation corrections, exciton, and evaporation models were used.
MT=18	Fission cross sections were evaluated on the bases of measured data by M. Silbert ⁷ , E. Fomushkin ⁸ and I. Vorotnikov. ⁹
MT=103,	107 (n,p), and (n, α) Cross Sections. The excitation function of the (n,p) and (n, α) reactions were calculated using systematics in an evaporation model including pre-equilibrium effects.
MF=4	Angular distributions of secondary neutrons.
MT=2,	51-68 Angular distributions of elastic and inelastic scattering were calculated using optical models and Hauser-Feshbach theory.
MT=16,	17, and 18 Assumed isotropic in the center-of-mass system.
MT=91	Assumed isotropic in the laboratory system.
MF=5	Energy distributions of secondary neutrons.
MT=16,	17, 18, and 91 Calculated with optical, exciton and evaporation models; and given in tabulated distribution form.
MT=18	Maxwellian fission spectrum. The temperatures were estimated from Hu Jimin et al. ¹⁰

References

1. Zhou Delin et al., "Evaluation of Neutron Nuclear Data for ²⁴⁹Bk," (1986), Internal Report.
2. Y. Kikuchi et al., JAERI-M 85-138 (1985).
3. Qiu Xijun et al., HSJ-78235 (LLJS), Internal Report (1978).
4. R. Benjamin et al., Nucl. Sci. Eng., 85, 261 (1983).
5. V. Anufriev et al., AE, 55, 285 (1983).

6. S. Muughabghab, "Neutron Cross Sections," Vol.1, Neutron Resonance Parameters and Thermal Cross Sections (1984).
7. M. Silbert, Nucl. Sci. Eng., 63, 198 (1977).
8. E. Fomshkin et al., Sov. J. Nucl. Phys., 14, 41 (1972).
9. I. Vorotnikov et al., Sov. J. Nucl. Phys., 10, 419 (1970).
10. Hu Jimin et al., HSJ-78221 (LLJS), Internal Report (1978).

Reference: No Primary Reference
Evaluators: Zhou Delin, Su Zhongdi and Others
Evaluated: April 1989
Material: 9852
Content: Neutron transport

Summary of ENDF/B-VI Evaluation

History

This evaluation was performed by Su Zhongdi and Zhang Jin et al. under contracts between the IAEA/NDS and the CNDC. (For a complete discussion of this evaluation refer to the report "Evaluation of Neutron Nuclear Data for ²⁴⁹Cf,"¹) and revision by Zhou Delin and Liu Tong.

File Information

MF=1 General information.
 MT=451 Comments and Dictionary.
 MT=452 Total number of neutrons per fission. It is the sum of delayed neutrons (MT=455) and prompt neutrons per fission (MT=456).
 MT=455 Delayed neutron data. T. R. England +, LA-11151, LA-11534, and LAURA-88-4118.
 MT=456 Number of prompt neutrons per fission. Taken from a calculation from Qiu Xijun et al.³

MF=2 MT=151 Resonance Parameters. The MLBW formalism was used for the resolved resonance parameters in the 1.0×10^{-5} to 70 eV range. Recommended resonance parameters were mainly based on the measurement of Benjamin et al.⁴ and Anufriev et al.⁵ Also consulted were the evaluations of Mughabghab⁶ and Y. Kikuchi et al.² Unresolved resonance parameters were obtained on the bases of recommended resolved resonance parameters from the evaluation of Y. Kikuchi et al.²

File Information, Continued

MF=3	Neutron Reaction Cross Sections.
MT=1,	2, 4, 16, 17, 51-65, 91, and 102 Total, elastic scattering, inelastic scattering, (n,2n), (n,3n) and capture cross sections calculated using optical, Hauser-Feshbach statistical theory with width fluctuation corrections, exciton, and evaporation models.
MT=18	Fission cross sections were evaluated on the bases of measured data by M. Silbert ⁷ , E. Fomushkin ⁸ and I. Vorotnikov. ⁹
MT=103,	107 (n,p) and (n, α) Cross Sections. The excitation function of the (n,p) and (n, α) reactions were calculated by Zhao Zhixiang using systematics in an evaporation model including pre-equilibrium effects. ¹⁰
MF=4	Angular distributions of secondary neutrons.
MT=2,	51-65 Angular distributions of elastic and inelastic scattering were calculated using the optical model.
MT=16,	17, 18 Assumed isotropic in the Center of Mass system.
MT=91	Assumed isotropic in the Laboratory system.
MF=5	Energy distributions of secondary neutrons.
MT=16,	17, 18, and 91 Calculated with optical, exciton and evaporation models, and given in tabulated distribution form.
MT=18	Maxwellian fission spectrum temperatures estimated from Hu Jimin et al. ¹¹

References

1. Su Zhongdi et al., "Report of Evaluation of Neutron Nuclear Data for ²⁴⁹Cf" (1986), To be Published.
2. Y. Kikuchi et al., JAERI-M 85-138 (1985).
3. Qiu Xijun et al., HSJ-78235 (LLJS), Internal Report (1978).
4. R. Benjamin et al., Nucl. Sci. Eng., **85**, 261 (1983).
5. V. Anufriev et al., AE, **55**, 285 (1983).
6. S. Mughabghab, "Neutron Cross Sections," Vol.1, Neutron Resonance Parameters and Thermal Cross Sections (1984).

7. M. Silbert, Nucl. Sci. Eng., 63, 198 (1977).
8. E. Fomshkin et al., Sov. J. Nucl. Phys., 14, 41 (1972).
9. I. Vorotnikov et al., Sov. J. Nucl. Phys., 10, 419 (1970).
10. Zhao Zhixiang et al., "Systematics of Excitation Functions for (n, charged particle) Reactions," Masters Thesis, Institute of Atomic Energy, Beijing, China, 1985.

Appendix

ENDF/B-VI Changes in Release 1

The following errors have been detected in the initial release of ENDF/B-VI and are corrected in release 1.

MAT	Error
0125 (^1H)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
0225 (^3He)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
0325 (^6Li)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
0325 (^6Li)	MF = 3, MT = 53, LR should be 32 and Q should be -1.4737 MeV
0525 (^{10}B)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
0525 (^{10}B)	MF = 3, MT = 57, LR should be 0
0525 (^{10}B)	MF = 3, MT = 62,64,68,70,71,73,74,76,77,79,80,81,83,84, LR should be 35, and Q should be -5.934 MeV
0525 (^{10}B)	MF = 3, MT = 65,78, LR should be 28 and Q should be -6.585 MeV
0525 (^{10}B)	MF = 3, MT = 55, LR should be 22 and Q should be -4.46 MeV
0625 (^{nat}C)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
1125 (^{23}Na)	Mf = 32, MT = 151, NER should be 1, not 0
2425 (^{50}Cr)	MF = 6, MT = 51-56, LCT should be =2
2431 (^{52}Cr)	Remove elastic transformation matrix
2434 (^{53}Cr)	MF = 6, MT = 51-63, LCT should be =2
2437 (^{54}Cr)	MF = 6, MT = 51-54, LCT should be =2
2625 (^{54}Fe)	Remove elastic transformation matrix
2631 (^{56}Fe)	Remove elastic transformation matrix
2634 (^{57}Fe)	MF = 6, MT = 51-55, LCT should be =2
2637 (^{58}Fe)	MF = 6, MT = 51-52, LCT should be =2
2828 (^{58}Ni)	Correct capture widths for 58.7 and 439.52 keV resonances

MAT	Error
2831 (^{60}Ni)	Revise resonance region and remove elastic transformation matrix
2834 (^{61}Ni)	MF = 6, MT = 51-58, LCT should be =2
2837 (^{62}Ni)	MF = 6, MT = 51-54, LCT should be =2
2843 (^{64}Ni)	MF = 6, MT = 51-52, LCT should be =2
4000 (^{nat}Zr)	MF = 5, MT = 91, U should = -Q = 2.821 not 2.871
4125 (^{93}Nb)	All SMODS=0 in the directory should be =1.
5025 (^{112}Sn)	MF = 3, MT = 102, interpolation code for low energy should be log-log (5)
5031 (^{114}Sn)	MF = 3, MT = 102, interpolation code for low energy should be log-log (5)
5728 (^{139}La)	Incorrect evaluation converted to ENDF/B-VI
6040 (^{147}Nd)	MF = 2, MT = 151, fictitious J-values used in MLBW representation
6149 (^{147}Pm)	MF = 2, MT = 151, fictitious J-values used in MLBW representation
6246 (^{151}Sm)	MF = 2, MT = 151, fictitious J-values used in MLBW representation
6337 (^{155}Eu)	MF = 2, MT = 151, fictitious J-values used in MLBW representation
7243 (^{180}Hf)	MF=2, MT=151, unresolved region, L=1, second J-value should be 1.5 not 1.0
7400 (^{nat}W)	MF=3, MT=1,2,102, have unequal energy values at several discontinuities
7400 (^{nat}W)	MF=14, MT=4, NK(N1) should be 1 not 198
7925 (^{197}Au)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
7925 (^{197}Au)	MF=3, MT=102, Q should be 6.51238 MeV
8234 (^{207}Pb)	correction to resonance region
9228 (^{235}U)	Add uncertainties from 1990 CESWG Standards Report to file 1 comments
9228 (^{235}U)	MF=2 MT=151 Revised resonance parameters to give "drooping" η below .1 eV.
9228 (^{235}U)	Correct for minor glitch in file 3 cross sections at 100 keV.

MAT	Error
9228 (^{235}U)	Update $\bar{\nu}$ covariances.
9228 (^{235}U)	Covariance file MF=33 should be removed, version V
9237 (^{238}U)	Covariance files MF=33, MT = 1,2,18,102 should be removed
9346 (^{237}Np)	Remove MF=4, MT=19,20,21,38
9440 (^{240}Pu)	Covariance file MF=33, MT=18 should be removed
9443 (^{241}Pu)	Covariance file MF=33 should be removed, version V
9546 (^{242}Am)	The delayed fission neutron spectrum should be removed. The spectrum is for the metastable target.
9547 (^{242m}Am)	The delayed fission neutron spectrum should be added. Erroneously added to MAT 9546
9861 (^{252}Cf)	The delayed fission neutron spectrum should be removed. The spectrum is for spontaneous fission.
0031 (Graphite)	D. Mathews corrections missing

END

**DATE
FILMED**

5 / 7 / 92