2-18-92 JSD

MSRC

1991 Annual Report

Molecular Science Research Center

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

This document is printed on recycled paper.

Molecular Science Research Center

1991 Annual Report

**M. L. Knotek
and the Staff of the MSRC**

March 1992

**Prepared for
the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830**

**Pacific Northwest Laboratory
Richland, Washington 99352**

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

EB

Contents

Opening Remarks

<i>M. L. Knotek</i>	vii
---------------------------	-----

Chemical Structure and Dynamics 1

1. Cluster Ions as Models for Bulk-Phase Chemistry: The Solvation Process <i>S. E. Barlow, T. M. Orlando, and D. Ray</i>	2
2. Structure and Dynamics of Size-Selected Cluster Ions <i>K. W. Brown and D. Ray</i>	3
3. Photofragment Imaging: The Photodissociation of Bromomethane, Bromoethane, and Bromoethanol <i>D. W. Chandler, J. W. Thomas, Jr., and W. P. Hess</i>	5
4. A New Approach to the Vibrational Spectroscopy of Clusters: The ZEKE Spectrum of the Phenol-Water Complex <i>S. D. Colson, O. Dopfer, G. Reiser, K. Müller-Dethlefs, and E. W. Schlag</i>	6
5. Ions at Surfaces <i>B. Maschhoff, J. P. Cowin, B. Ellison, J. Biesecker, and T. Orlando</i>	9
6. Spectroscopy and Dynamics of Molecular Clusters as a Route to Understanding Solvent Influences in Chemical Reactions <i>R. Knochenmuss, D. Ray, and W. Hess</i>	10
7. Surface Chemistry of Chlorinated and Fluorinated Hydrocarbons Via Photons and Hot Electrons <i>M. Kwini, M. Iedema, and J. P. Cowin</i>	11
8. Molecular Dynamics at the Water/Solid Interface <i>M. Kwini, M. Iedema, and J. P. Cowin</i>	12
9. Single-Molecule Chemistry <i>S. A. Joyce, L. Bumm, and J. P. Cowin</i>	14
10. Photodissociation Dynamics of Molecules and Clusters <i>W. P. Hess and S. P. Sapers</i>	15
11. Apparatus for Time-Resolved Spectroscopy <i>D. M. Friedrich, M. O. Trulsen, and A. G. Joly</i>	16
12. Time-Resolved Fluorescence as a Probe of Molecular Dynamics <i>G. R. Holtom</i>	16
13. Stimulated Cluster-Surface Chemistry <i>T. M. Orlando and S. E. Barlow</i>	17
14. Spectroscopy of Small Cluster Molecules in Low-Temperature Beams <i>S. W. Sharpe, T. A. Hu, and E. L. Chappell</i>	18
15. Spectroscopic Studies of the Gas-Catalytic Surface Interface <i>R. S. Smith and S. D. Colson</i>	21

Environmental Dynamics and Simulation	25
1. Rational Design of Metal Selective Liquids	
<i>B. P. Hay</i>	25
2. Molecular-Scale Investigations of Alkali Halides	
<i>L. E. Eary, R. L. Erikson, and C. J. Hosteller</i>	26
3. Synchrotron Development	
<i>J. E. Amonette and S. V. Mattigod</i>	27
 Macromolecular Structure and Dynamics	 29
1. Progress in the Development of a Gigahertz NMR Spectrometer	
<i>T. S. Tenforde</i>	30
2. NMR Analysis of DNA Intrinsic Bending Modes	
<i>M. A. Kennedy and B. R. Reid</i>	32
3. Large Molecule Mass Spectrometry	
<i>R. D. Smith, J. A. Loo, A. L. Rockwood, H. R. Udseth,</i> <i>R. R. Ogorzalek-Loo, and B. E. Winger</i>	36
 Materials and Interfaces	 43
1. Surface Structure and Chemistry of Oxides and Minerals	
<i>D. R. Baer, D. L. Blanchard, and M. H. Engelhard</i>	43
2. Design of Selective Layers for Chemical Microsensors	
<i>J. Janata</i>	46
 Theory, Modeling, and Simulation	 47
1. Molecular Theory and Modeling	
<i>B. C. Garrett, L. X. Dang, T. H. Dunning, Jr., A. C. Hess, M. I. McCarthy,</i> <i>M. M. Marino, R. P. McRae, G. K. Schenter, S. S. Xantheas, G. Haynes,</i> <i>R. Poshusta, G. C. Schatz, G. Voth, and W. C. Ermler</i>	47
2. Materials Theory and Modeling	
<i>A. C. Hess, J. P. LaFemina, D. L. Lessor, M. I. McCarthy, T. J. Godin,</i> <i>A. J. Skinner, J. C. White, G. K. Schenter, D. R. Baer, C. B. Duke,</i> <i>D. L. Blanchard, A. S. Gibson, and R. Haydock</i>	56
3. Biomolecular Modeling and Simulation	
<i>R. L. Ornstein, G. E. Arnold, M. B. Bass, M. D. Paulsen, et al.</i>	63
4. Molecular Science Software Group	
<i>R. A. Bair, D. F. Feller, M. W. Feyereisen, R. A. Kendall, M. A. Thompson</i> <i>M. A. Stave, R. J. Littlefield, M. M. Szczesniak, and G. Chalasinski</i>	68

Computing and Information Sciences 77

1. Computer Systems Engineering	
<i>J. O. Schroeder, J. W. Brothers, D. M. DeVaney, D. R. Jones,</i>	
<i>R. J. Littlefield, M. J. Peterson</i>	78
2. Distributed Computing Systems	
<i>J. O. Schroeder, W. S. Eggers, D. R. McGee</i>	80
3. Distributed Computing Systems Testbed	
<i>T. L. Harper, J. O. Schroeder</i>	81
4. Scientific Networking	
<i>M. J. Peterson, T. A. Harper</i>	83
5. Data Storage and Archival	
<i>D. M. DeVaney</i>	84
6. Database Management Systems	
<i>D. M. DeVaney</i>	87
7. Self Describing Data	
<i>R. S. Barga, D. M. DeVaney</i>	88
8. Scientific Laboratory Notebook	
<i>D. M. DeVaney, A. Schur, M. L. Yim</i>	90
9. Visual Science Computing	
<i>D. R. Jones, J. J. Thomas</i>	91
10. Graphical User Interface Development	
<i>D. R. Jones, G. Chin, J. J. Thomas</i>	93
11. Software Development for Visual Tools	
<i>D. R. Jones, E. N. Thornton, J. J. Thomas</i>	95
12. High-Performance Computing	
<i>R. J. Littlefield, G. S. Thomas, G. I. Fann, A. B. Shiftlet</i>	96
13. Data Acquisition System Development	
<i>G. A. Anderson</i>	99
14. Electronics and Computer Interfaces	
<i>K. D. Steele, G. A. Anderson</i>	99

Appendix A - MSRC Staff and Associates A.1

MSRC Management	A.1
MSRC Staff	A.2
MSRC Associates	A.22

Appendix B - 1991 Publications and Presentations B.1

Publications	B.1
Manuscripts Not Yet Published	B.5
Presentations	B.10

Appendix C - Activities	C.1
Collaborations	C.1
Colloquia Speakers During 1991	C.6
MSRC Technical Advisory Panel	C.8
Tours and Visitors During 1991	C.9
 Appendix D - Acronyms and Abbreviations	 D.1

Opening Remarks

During 1991, the Molecular Science Research Center (MSRC) experienced solid growth and accomplishment and the Environmental, and Molecular Sciences Laboratory (EMSL) construction project moved forward. We began with strong programs in chemical structure and dynamics and theory, modeling, and simulation, and both these programs continued to thrive. We also made significant advances in the development of programs in materials and interfaces and macromolecular structure and dynamics, largely as a result of the key staff recruited to lead these efforts.

The EMSL Construction Project team also made excellent progress during the first half of the year. Contract negotiations with the architect/engineering firm of Zimmer Gunsul Frasca Associates (ZGF) of Portland, Oregon, were completed, and the kick-off meeting for the definitive design effort was held at Pacific Northwest Laboratory (PNL) in February. Over the following months, ZGF staff developed a work plan for the facility design as well as a Quality Assurance Plan, both of which were approved.

In addition, ZGF completed the main Phase I document, the Definitive Design Program Document, after discussing operational and facility needs with all PNL user scientists. Based on this information, ZGF modified the original building concept and developed the Definitive Design Program Document.

If there was one pervasive activity for the past year, however, it was to strengthen the role of the EMSL in the overall environmental restoration and waste management (ER/WM) mission at Hanford. Since Secretary of Energy Watkins identified Hanford as the "flagship" of the U.S. Department of Energy's (DOE) ER/WM effort in October 1990, we have waged an aggressive campaign to heighten awareness of the critical role the EMSL can play in providing the scientific and technical breakthroughs that will be necessary to accomplish cleanup in a timely, cost-effective manner. These extended activities involved not only MSRC and EMSL staff but all PNL scientific and technical staff engaged in ER/WM programs.

Our primary goal was to frame a comprehensive definition of the technological voids and gaps that could

impede cleanup at Hanford and to describe the roles that basic and applied research will play in meeting these needs. Our efforts were put to the test on June 11-12, 1991, when DOE's EMSL Working Group met to review and evaluate the EMSL project and research program scope. The objective of the review was to determine whether the EMSL project and programs would support Hanford site-specific technology and facility needs for directed research.

The Working Group included representatives from Sandia National Laboratories, Texas A&M University, Vanderbilt University/Clemson University, and CH2M Hill, and was supported by personnel from DOE Headquarters and the DOE Richland Field Office (RL). The summary of the group's draft report included the following statements, which were characterized as "consensus conclusions."

- "The EMSL proposal, as presented, contains a satisfying array of basic and applied research efforts. The Working Group believes that the proposal complements the required strategy. Moreover, the proposal is 'robust' in that it is broad and flexible enough to accommodate future changes in scope and direction."
- "The Working Group concluded that the EMSL proposal is sensible and flexible enough to enable DOE to proceed with the project. There is a very low risk of expending funds that are not properly directed toward solving some of the significant Hanford Site problems."

Following this review, James Decker, who was Director of the DOE Office of Energy Research at that time, and Leo Duffy, Director of the Office of Environmental Restoration and Waste Management, signed the Memorandum of Understanding related to construction and operation of the EMSL.

The exercise of focusing the programs of the EMSL on ER/WM needs strengthened the contextual framework for EMSL research and highlighted several previously unidentified needs. As a result, we established major sensor development and advanced processing technology efforts as part of the EMSL program and

clarified the long-term goals and focus of the EMSL and the MSRC in light of DOE's ER/WM programs. These efforts have strengthened MSRC research program directions and provided key linkages that tie molecular science research to other programs within PNL and at universities and industrial laboratories.

In addition, we increased the number of collaborations, memoranda of understanding, and other cooperative ventures that have been established with universities, DOE laboratories, and industry. These extended agreements serve as guidelines for focusing MSRC, EMSL, and PNL research; as media for transmitting DOE's research needs to the larger scientific community; and as vehicles for transferring the output of the organization to its users.

This year has also been a very important and critical in the intellectual development of the MSRC as we recruited key leaders and brought them up to speed in their research programs. Dr. Jiri (Art) Janata, formerly a professor in the Department of Materials Science and Engineering at the University of Utah, accepted the position of Associate Director for the Materials and Interfaces program. One of his priorities has been to recruit staff, and he has already been joined by Drs. Bruce C. Bunker and Michael A. Henderson from Sandia National Laboratories and the University of Pittsburgh, respectively.

Dr. Gary P. Drobny, associate professor of chemistry at the University of Washington, has been placed under contract to head the MSRC's nuclear magnetic resonance (NMR) program, which includes studies of the structure and dynamics of biological molecules and solid-state NMR. Eventually, Dr. Drobny will transition into the management role for the development of an ultrahigh-field (gigahertz) NMR system that can be used in combination with advanced mass spectrometers to perform structural analyses of large macromolecules such as enzymes to be used for bioremediation of contaminants at DOE sites. The NMR program was also joined during the past year by Dr. Michael A. Kennedy, who became the second PNL scientist to win the distinguished Alexander Hollaender postdoctoral fellowship awarded by the DOE.

Ms. Janice R. Lewis, staff scientist in PNL's Applied Physics Center, is the new acting manager of the Computing and Information Sciences program. Dr. Richard J. Kouzes, formerly a senior research

physicist and lecturer at Princeton University, has been assigned to the group full-time along with one post-doctoral appointee. A number of other staff from the Applied Physics Center are also supporting the group's efforts.

During 1991, five scientists joined the Chemical Structure and Dynamics program staff: Dr. Stephan E. Barlow, University of Colorado, Denver; Drs. Stephen A. Joyce and Thomas M. Orlando, Sandia National Laboratories; Dr. Robin (Rod) S. McDowell, Los Alamos National Laboratory; and Mr. Martin J. Iedema, a Ph.D. candidate and former high school science teacher. Seven new postdoctoral appointees were assigned to the group through the NORCUS (Northwest College and University Association for Science) program at PNL. In addition, Dr. Miles J. Dresser, Associate Professor of Physics at Washington State University, spent eight weeks with the group this summer.

The Theory, Modeling, and Simulation program added six full-time staff members in 1991: Dr. L. René Corrales, University of Texas at Austin; Dr. Lien X. Dang, IBM Almaden Research Center, San Jose; Dr. Martin W. Feyereisen, University of Utah, Salt Lake City; Dr. Anthony C. Hess, until recently a postdoctoral appointee with the MSRC; Dr. Maureen I. McCarthy, formerly a postdoctoral fellow at the Hebrew University in Jerusalem, Israel; and Dr. Mark A. Thompson, Brookhaven National Laboratory.

Four new postdoctoral appointees and two visiting scientists are also participating in the program. Dr. Walter C. Ermler, a professor at the Stevens Institute of Technology in Hoboken, New Jersey, was awarded a 1-year appointment; and Dr. Ronald D. Poshusta, professor with the Chemistry Department at Washington State University, received a 9-month appointment.

As a result of this aggressive recruiting program, the MSRC staff now totals 121. This number includes 31 full-time scientists, 22 administrative personnel, 39 researchers assigned from other PNL organizations, 28 postdoctoral appointees, and 1 doctoral candidate. By the end of the next fiscal year, we expect our staffing level to increase to about 180, including new associate directors for our Macromolecular Structure and Dynamics and Computing and Information Sciences programs.

During the past year, staff members contributed to or wrote nearly 90 publications and made approximately 100 presentations at national and international events. About 55 manuscripts are awaiting publication. The Colloquia Series, which began in 1989, also flourished. This year the series brought 121 distinguished scientists to PNL and the MSRC to share their knowledge and insight. They represented universities, research laboratories, and industry in the United States, Canada, Czechoslovakia, England, Germany, Italy, Japan, and Switzerland. We also welcomed hundreds of visitors from academia, government, and industry and sponsored the following workshops:

- Environmental Materials and Interfaces
- FTICR Design for Large Molecule Studies
- Parallel Eigensolver
- Large Molecule Activation and Disassociation
- Parallel Computing in Chemical Physics.

In 1992, we are planning to offer workshops on

- Scientific Information Management
- Research Needs for Environmental Restoration
- Parallel Chemical Physics
- Visualization for Computational Chemistry

- Remote and *In-Situ* Optical Analyses
- Reactions at Environmental Interfaces
- NMR Imaging for Materials Science.

We are extremely pleased with our progress and program development over the last year. The Appendixes to this report contain biographic information about our full-time staff and associates, lists of publications, names of colloquia speakers and Technical Advisory Panel members, and tours and visitors who were hosted at the MSRC during 1991.

We have an opportunity to create a unique scientific resource at PNL -- a resource that can have an important impact on a problem that is critical to the DOE and the nation. With the coming year, we face the fundamental challenge of completing the EMSL design activities and beginning construction. We hope next year's report will announce that EMSL construction has begun.

Michael L. Knotek
Manager, Molecular Science Research Center

Chemical Structure and Dynamics

Chemical Structure and Dynamics

The Chemical Structure and Dynamics program is developed around several initiatives in response to the need for a fundamental molecular-level understanding of chemistry required to meet the environmental and waste management needs of the U.S. Department of Energy (DOE). These needs are very broad, ranging from understanding health effects to the cleanup of contaminated sites. The Chemical Structure and Dynamics program in combination with other programs within the Molecular Science Research Center (MSRC) and at Pacific Northwest Laboratory (PNL) constitutes the broad range of capabilities required to mount a multilevel attack on the key scientific and technology issues. Specific Chemical Structure and Dynamics program elements are designed to focus on the DOE initiatives in health effects, processing and containment of mixed (chemical and radioactive) wastes, contaminated soil and groundwater, and advanced analytical and characterization methods. The impact of the work is also expected to extend into many other fields essential to U.S. technological and economic strength.

Of vital importance to the establishment of this new program has been the development of the scientific staff. The past year has been particularly rewarding in this respect with the addition of the following scientists: Dr. Stephen E. Barlow (ion chemistry and mass spectroscopy), Dr. Stephen A. Joyce (surface microscopy), Dr. Bruce D. Kay (surface kinetics), Dr. Robin S. (Rod) McDowell (structural spectroscopy and Chemical Structure and Dynamics Program Manager), and Dr. Thomas M. Orlando (cluster and surface reactions). Our research staff now includes thirteen research scientists and about the same number of students, postdoctoral fellows, and other staff.

The overall research philosophy of the Chemical Structure and Dynamics program is built around the established relationship between structure and function that is the underpinning of molecular science. The intent is to describe chemical processes at the molecular level in terms of the initial and final structures, including measurements of the controlling energetics, rates, mechanisms, and real-time evolution of these structures.

The structural studies provide immediate insight into the nature of the condensed phase and are key to the

direct comparison with theory. Structures provide well defined targets for testing theoretical assumptions. This approach is particularly important in dealing with the complex problems associated with the description of the molecule/substrate interface. Without good structural data, it will be very difficult to evaluate the quality of the potential functions that are needed to describe the molecular dynamics. The structural program is built around a combination of laser-based optical methods as well as atomic force and scanning tunneling surface microscopies.

Real-time measurements of the motions and chemical reactions on model substrates will have two direct consequences. First, they will provide new insights that will help shape the description of these phenomena in terms of molecular processes. Second, they will provide the data necessary for the evaluation of molecular dynamics theories that are being developed to describe molecular processes in terms of the kinematics of atomic nuclei on potential energy surfaces (PES).

Another program initiative relies on the synthesis and characterization of microscopic models of condensed-phase chemistry. Atomic and molecular clusters are a form of matter whose properties lie outside the realm of general chemical experimental science. Thus, their study not only provides a quantitative basis for comparison with theory, but is also the source of a microscopic understanding of condensed-phase structures and dynamics. At the heart of this activity is a Cluster Structure and Dynamics Research Facility (CSDRF) that was designed, in part, as a result of workshop attended by a group of 50 scientists. The workshop report discusses the relationship between cluster chemistry and pollution problems and outlines the basic requirements for developing the CSDRF as an effective collaborative user facility.

The combination of state-of-the-art lasers and atomic resolution surface-mapping tools, such as scanning tunneling microscopy (STM) and atomic force microscopy (AFM), will be used to study interfacial chemistry and dynamics on extended structures. With this combination of tools, scientists can for the first time directly observe chemical processes on realistic chemical surfaces rather than on idealized single-crystal models. This initiative, in concert with the studies of

molecular clusters, provides a new window through which we can observe new aspects of condensed-phase chemistry and physics and, thus, expands our understanding of the chemistry of materials science in ways that are expected to be far reaching.

The initiatives described above illustrate elements of a broad program, including structure and dynamics studies of molecules in isolation, condensed in microstructures, in solution, and on extended surfaces. This multilevel, multidisciplinary approach allows for the dissection of complicated systems with the anticipation that the study of the component parts will lead to a clearer understanding of complex chemical processes in natural and other complex media.

1. Cluster Ions as Models for Bulk-Phase Chemistry: The Solvation Process

S. E. Barlow, T. M. Orlando, and D. Ray

Gas-phase chemistry has made such progress in the last few decades that its methods are generally preferred, when available, for probing the basic structure, thermochemistry, and reaction dynamics of molecules. However, the connection between such results and bulk-phase chemistry in liquids and at interfaces has remained more tenuous. Progress is now being made in applying the techniques of gas-phase studies to cluster materials.⁽¹⁻⁴⁾ Areas of particular interest include solvent-solute interactions,⁽¹⁾ crystal formation,⁽²⁻⁴⁾ and cluster chemistry.^(3,4)

Although the ionic nature of salts has been recognized for nearly two centuries, direct observation of solvation has proven elusive. We are developing instrumentation to study solvation, one solvent molecule at a time. In addition to providing valuable insight into a number of important questions surrounding solvation, these experiments will serve to develop techniques that will be of general value in the longer-range mission of the Environmental and Molecular Sciences Laboratory (EMSL) at PNL.

A Laser Ablation Source (LAS) that provides a pulse of clusters (cations, anions, and neutrals) is under construction at the MSRC.⁽⁵⁾ A schematic diagram of the LAS is shown in Figure 1.1. The ion beam is sent down a time-of-flight (TOF) tube to a gate that allows

Figure 1.1. LAS-2 System. Laser ablation source with TOF/MS and RF trap for solvation studies of microcrystalline salts.

only ion clusters of the desired mass to charge ratio to enter the radio-frequency (RF) trap and be captured. This scheme will allow us to build up a large number of mass-identified trapped clusters for experimental purposes. Alternatively, the gate and trap can be floated at the TOF potential and the entire ion beam analyzed directly by standard TOF/mass spectrometry.

This experiment represents the application of previously developed technologies in a new and unique combination. Previous experimental work at other laboratories has demonstrated that laser ablation sources can produce small fragments of ionic crystals.^(3,4) Other work has demonstrated the feasibility of collecting externally produced ions in an RF quadrupole trap as well as using these devices as chemical reactors.⁽⁶⁾

Initial experiments will focus on two questions. First, is it possible to observe the reactions of the type:

where M is a metal and X is an anion? That is, can we observe the formation of a solvation sphere? The second and more general question is whether charged

cluster ions are applicable for investigating the chemistry of neutral systems. Some work with silicon clusters indicates that this is indeed the case,⁽³⁾ but we need to investigate more cases to determine the range of validity of the hypothesis. In principle, one anticipates that when the particle size exceeds the electrostatic capture cross section (ion-dipole and ion-induced dipole), the perturbation these long-range forces bring to the molecular interactions will become negligible. The LAS will allow us to investigate this important question by producing a variety of both cations and anions.

We should also be able to investigate the important phenomena of multiply-ionized-solvated species, such as Ca^{2+} , Sr^{2+} , and Fe^{3+} . For example, by seeding helium with water in the pulsed valve, we should be able to produce clusters of singly-ionized strontium. Then, by directing a laser through the stored ion cloud, we can observe the change in the second ionization potential as a function of solvation.

Should these experiments be successful, we would then follow up with more detailed photoionization measurements. Preliminary work can also be conducted to examine the thermochemistry of solvation shells.^(7,8)

The LAS will provide a convenient and very versatile platform for examining many fundamental solvation phenomena with unprecedented resolution and speed. The focus of investigations can be rapidly redirected for survey work and may prove useful for identifying fruitful paths for more detailed investigations by other techniques.

References

1. S. E. Barlow, T. Dang, and V. M. Bierbaum, "Hydrogen Deuterium Exchange Reactions of Methoxy-Methanol Clusters," *J. Amer. Chem. Soc.* **112**, 6832 (1990).
2. E. C. Honea, M. L. Homer, and R. L. Whetten, "Excess-Electron and Excess-Hole States of Charged Alkali Halide Clusters," *Int. J. Mass Spectrom. Ion Proc.* **102**, 213 (1990).
3. M. F. Jarrold, "Nanosurface Chemistry on Size-Selected Silicon Clusters," *Science* **252**, 1085 (1991).
4. J. L. Elkind, F. D. Weiss, J. M. Alford, R. T. Laaksonen, and R. E. Smalley, "Fourier Transform Ion Cyclotron Resonance Studies of H_2 Chemisorption on Niobium Cations," *J. Chem. Phys.* **88**, 5215 (1988).
5. S. Maruyama, L. R. Anderson, and R. E. Smalley, "Direct Injection Supersonic Cluster Beam Source for FTICR Studies of Clusters," *Rev. Sci. Instr.* (in press); see also W. R. Gentry, in *Atomic and Molecular Beam Methods*, G. Scoles, ed., Ch. 3, Oxford, New York, 1988.
6. R. E. March, and R. J. Hughes, *Quadrupole Storage Mass Spectrometry*, John Wiley & Sons, Inc., New York, 1989.
7. E. Kochanski, and E. Constantin, "Theoretical and Experimental Studies of the System $\text{Ca}^+(\text{H}_2\text{O})_n$ for $n=1$ to 10," *J. Chem. Phys.* **87**, 1661 (1987).
8. M. H. Shen, and J. M. Farrar, "Absorption of Size-Selected Solvated Metal Cations: Electronic States, Symmetries and Orbitals in $\text{Sr}^+(\text{NH}_3)_{1,2}$ and $\text{Sr}^+(\text{H}_2\text{O})_{1,2}$," *J. Chem. Phys.* **94**, 3322 (1991).

2. Structure and Dynamics of Size-Selected Cluster Ions

K. W. Brown and D. Ray

The synthesis of clusters in molecular beams affords the opportunity to create model systems that exhibit a rich variety of chemical phenomena. The application of techniques of optical spectroscopy to size-selected cluster beams facilitates rigorous studies of fundamental molecular processes in these model systems. We are currently beginning projects to study solvation, reaction, and adsorption in size-selected clusters with up to 100 constituents. This range of cluster sizes corresponds to a molecule or reactive system in solution surrounded by up to two solvation shells, an important size range for many-body effects on molecular processes.

Our apparatus is nearing completion. The instrument, a tandem TOF mass spectrometer, allows a wide variety of action spectroscopies to be performed on mass selected cluster ions. For example, vibrational and electronic spectra can be obtained via photo-absorption spectroscopy, allowing determination of

cluster size effects on the electronic structure of an ionic chromophore. Photoelectron spectroscopy of inorganic oxide cluster anions is also possible, yielding thermochemical and structural data on these important model species. Some of this frequency-domain data can also be interpreted within a time-dependent framework to yield information on dynamical behavior.

Experiments involving the photodissociation of molecules within clusters are planned. Hydrogen halide ions, HX^+ , will be used to examine the effects of sequential solvation on predissociation dynamics. Upon ultraviolet excitation, HF^+ predissociates to form H^+ and F , while HCl^+ and HBr^+ dissociate into H and X^+ . Surrounding these reactants with a solvent cluster of selected size and composition (e.g., H_2O) and monitoring the ionic photoproducts offers a direct measurement of the effects of selective solvation on the predissociation dynamics. Since the charge is localized on different products in this series of predissociation reactions (i. e., $H + X^+$ or $H^+ + X$), the photoproduct distributions as a function of photodissociation energy will reflect the differing dynamics of H and H^+ interacting with the surrounding microsolvant.

Recent experiments⁽¹⁻³⁾ exploring the effects of solvation on the dynamics of chemical reactions indicate the feasibility of real-time measurements of intracuster dynamics, and furthermore suggest that such measurements may provide qualitatively new information on chemical dynamics in a microsolvant environment. Time-resolved measurements of reactive dynamics in mass-selected clusters will be made in our apparatus using ultrafast, time-resolved absorption, emission, and photoelectron spectroscopies. The goal of these experiments will be to provide new, detailed dynamical information on chemical reactions in these model systems.

The new technique of ultrafast, time-resolved photoelectron spectroscopy is under development. Initial experiments will measure the evolution of the electronic wave function of a dissociating molecule. Theory suggests that the change in electronic structure from reactant-like to product-like is abrupt. Understanding this evolution is fundamental to understanding solvent effects on photoinitiated chemical reactions. In these experiments, photodissociation will be initiated by an ultrafast laser pulse, and photo-

detachment will be made by a second ultrafast laser pulse at a later time. The kinetic energy of the detached electrons will be analyzed by a TOF electron energy analyzer. The evolution of the electronic wave function will be monitored by measuring the kinetic energy and angular distribution of the photodetached electrons as a function of the time delay between the pump and probe pulses. This experiment will provide the first direct probe of this fundamental process.

Ultrafast, time-resolved spectroscopies (e.g., photoabsorption and photoelectron) will be used to study the dynamics of dissociation and photoinduced charge-transfer processes in clusters. Reaction [e.g., proton transfer from $HCCl_2^+$ to Ar , from H_2O^+ to N_2O or CO_2 , electron transfer in $KrCO_2^+(\text{solvent})_n$ or $ArCO_2^+(\text{solvent})_n$ clusters] will be initiated by a tunable ultrafast pump pulse and interrogated, at a later time, by a second tunable ultrafast probe pulse. In the photoabsorption experiments, the probe pulse can be tuned to monitor the appearance of products, the disappearance of reactants, or the transient intermediate species. Measuring the probe-induced photofragmentation as a function of delay between the pump and probe pulses will map the dynamics in real time. Ultrafast, time-resolved photoelectron spectroscopy will be used to study the breaking of chemical bonds in a microsolvant environment. The photodissociation of a diatomic molecule (e.g., I_2^+) interacting with selected numbers of solvent molecules in a mass-selected cluster will be studied to explore the effects of the solvent on the transition from reactant-like to product-like electronic structure.

References

1. D. Ray, N. E. Levinger, J. M. Papanikolas, and W. C. Lineberger, *J. Chem. Phys.* **91**, 6533 (1989); J. M. Papanikolas, J. R. Gord, N. E. Levinger, D. Ray, V. Vorsa, and W. C. Lineberger, *J. Phys. Chem.* **95**, 8028 (1991).
2. J. J. Breen, L. W. Peng, D. M. Wilberg, A. Heikal, P. Cong, and A. H. Zewail, *J. Chem. Phys.* **92**, 805 (1990).
3. J. Steadman, and J. A. Syage, *J. Chem. Phys.* **92**, 4630 (1990); J. A. Syage, and J. Steadman, *J. Chem. Phys.* **95**, 2497 (1991).

3. Photofragment Imaging: The Photodissociation of Bromomethane, Bromoethane, and Bromoethanol

David W. Chandler,^(a) John W. Thomas Jr.,^(b)
and Wayne P. Hess

Photolysis of brominated hydrocarbons is routinely used to produce radicals for kinetic studies, but the details of this process are incompletely understood.⁽¹⁾ We have photolyzed three brominated organics and measured the kinetic energy distributions of the bromine fragments. Bromomethane, bromoethane, and bromoethanol in a supersonic molecular beam were photolyzed at 205 nm, producing neutral fragments in various translational, rotational, vibrational, and electronic states. The velocity of the bromine atoms in ground ($^2P_{3/2}$) or excited ($^2P_{1/2}$) states was determined by a variation of the resonance-enhanced, multiphoton-ionization, time-of-flight (REMPI-TOF) technique called photofragment imaging.^(1,2) The ionized bromine atoms are detected by a position-sensitive ion detector consisting of a microchannel plate/phosphor screen/charge-coupled-device (CCD) camera combination at the end of the TOF tube. By measuring the arrival time and distance the photofragments have moved off-axis, we obtain the speed of the fragments. With the geometry used, the three-dimensional (3-D) spatial distribution of the selected photofragment is uniquely determined by taking the inverse Abel transform of the two-dimensional (2-D) image recorded by the detector.⁽¹⁾

Excitation in the first absorption continuum of these compounds results in subpicosecond photodissociation of the C-Br bond. Therefore, the angular distribution of the bromine reflects the orientation of the transition moment in the parent molecule. For bromomethane, excitation is primarily to the 3Q_0 state;⁽³⁻⁵⁾ this state correlates to the formation of Br $^2P_{1/2}$ and is coupled to the ground state via a parallel transition moment. Absorption via a parallel transition leads to Br $^2P_{1/2}$ with a predominantly $\cos^2\theta$ angular distribution, where θ is measured with respect to the polarization vector of the photolysis light. Absorption to the

1Q state is via a perpendicular transition and correlates to formation of Br $^2P_{3/2}$ with a dominant $\sin^2\theta$ distribution. Excited-electronic-state curve crossing as the molecules dissociate mixes the states.

The results indicate that the fastest Br atoms correlate to the bond dissociation energy of the C-Br bond in the photodissociation of bromomethane, but not in the photodissociation of bromoethane or bromoethanol. This result suggests that the ethyl and hydroxyethyl fragments are produced in rotationally excited states. In the latter cases, the impulse is not directed at the center of mass of the radical fragment, and substantially more rotational energy is expected to be deposited into the radical fragment. Because ground-vibrational-state methyl fragments are formed in the photodissociation of bromomethane, a Franck-Condon argument would indicate that comparable amounts of vibrational-ground-state ethyl and hydroxyethyl radicals are formed in the dissociation of their brominated analogues. Estimates of Br kinetic energies support this explanation.

The velocity distributions for bromomethane reflect the different dissociation channels (Br and Br*) leading to different amounts of excitation in the methyl fragment. From our data we can determine the amount of translational energy in each fragment, but are unable to determine the partitioning of the remainder of the energy between the vibrational and rotational degrees of freedom in the organic radical.

References

1. D. W. Chandler, M. H. M. Janssen, S. Stolte, R. N. Strickland, J. W. Thoman Jr., and D. H. Parker, *J. Phys. Chem.* **94**, 4839 (1990).
2. J. W. Thoman Jr., D. W. Chandler, D. H. Parker, and M. H. M. Janssen, *Laser Chem.* **9**, 27 (1988).
3. R. K. Sparks, K. Shobatake, L. C. Carlson, and Y. T. Lee, *J. Chem. Phys.* **75**, 3838 (1981).
4. G. N. A. van Veen, T. Baller, A. E. DeVries, and N. J. A. van Veen, *J. Chem. Phys.* **87**, 405 (1984).
5. M. D. Barry and P. A. Gorrry, *Mol. Phys.* **52**, 461 (1984).

(a) Combustion Research Facility, Sandia National Laboratories, Livermore, California.

(b) Department of Chemistry, Williams College, Williamstown, Massachusetts.

4. A New Approach to the Vibrational Spectroscopy of Clusters: The ZEKE Spectrum of the Phenol-Water Complex^(a)

Steven D. Colson, Otto Dopfer,^(b)G. Reiser,^(b)
K. Müller-Dethlefs,^(b)and E. W. Schlag^(b)

Developing accurate potential energy functions for the description of hydrogen bonding can have significant impact on the validity of molecular dynamics calculations of complex processes, ranging from biomolecular activity to those processes governing contaminant transport in subterranean systems. One goal of this work is to provide experimental data for use in the quantitative evaluation and further development of potential energy functions for such applications. It is part of an ongoing project that includes close collaborations with theorists, using phenol-water complexes as model systems. The subject of this report is the vibrational spectrum of the phenol-H₂O cation complex, which is obtained by optical-optical double resonance threshold ionization spectroscopy with detection of the zero kinetic energy electrons (ZEKE). The ZEKE signal only appears when the total photon energy is resonant with a level in the ion.⁽¹⁻³⁾ This work represents a new application of this method to molecular cluster vibrational spectroscopy. The results include a well resolved vibrational spectrum from which the six H-bonding vibrational frequencies can be derived. This method provides vibrational frequencies with a precision that is normally limited by the laser resolution and/or by intrinsic properties of the molecular cluster (a few wavenumbers in this case). This is in direct contrast to the usual photoelectron methods used to obtain vibrational spectra of ions, where the width of the PES peaks (≥ 50 cm⁻¹) limits the resolution.

The one-to-one phenol water complex is an ideal case for the development of the ZEKE method for obtaining vibrational spectra of cluster ions. The electronic spectrum of the intermediate (neutral) state shows large shifts with the addition of each water molecule^(4,5) (Figure 4.1). Thus, through the use of

Figure 4.1. S₁-S₀ absorption features for the phenol-(H₂O)_n complexes: (A-C) the phenol-H₂O origin band, H₂O-wag, and O-H-O-stretch respectively; (D) the phenol-(H₂O)₂ origin region; (E) the phenol-(H₂O)₃ origin band.

threshold ionization spectroscopy, the spectra of the phenol-(H₂O)_n ($n=1-3$) complexes have been definitively identified⁽⁵⁾ and can be used as intermediate levels with little interference from other complexes. The $n=1$ complex shows activity due to H-bonding vibrations that are also well resolved from absorptions by other complexes⁽⁴⁻⁶⁾ and can be used as intermediate levels to modify the vibrational activity in the ZEKE spectra.

The spectra shown in Figure 4.2 were obtained by a pump-probe, two-laser technique with detection of the ZEKE produced by resonant ionization. This technique (Figure 4.3) uses a pump laser to prepare an excited state of the complex and a probe laser for ionization of the excited species. For Figure 4.2a, the pump laser was tuned to be resonant with the vibrationless origin [peak (A) in Figure 4.1] of the lowest excited singlet state of the phenol-water complex (formed by the supersonic expansion of phenol and water in an argon carrier gas). The probe laser wavelength was then scanned through the threshold ionization region of the excited complex. ZEKE electrons

(a) This work resulted from an Alexander von Humboldt Senior Science Award and was conducted at Munich.

(b) Institut für Physikalische und Theoretische Chemie der Technischen Universität, München, Germany.

Figure 4.2. Pump-probe ZEKE spectra of the phenol-H₂O complex via excitation of the (a) zero, (b) O-H-O stretch, and (c) H₂O-wag modes or the intermediate singlet excited state, respectively. The energy scale is that of the probe (ionization) laser.

are produced by ionization at threshold and at above-threshold energies where the probe laser is resonant with rovibrational energy levels in the ion. In this way,

Figure 4.3. The pump-probe threshold ionization spectra are taken by first exciting the selected species to the desired vibronic intermediate state with the pump laser. The probe laser is then scanned through the energies required for formation of the ion at the ionization threshold or at higher thresholds reached by preparation of the ion in excited rovibronic states.

the ionization energy and the rovibrational energies of the ion are revealed by the spectrum of the ZEKE electron intensity as a function of the probe laser energy.

The spectra shown in Figures 4.2b and 4.2c were taken respectively by exciting the O-H-O-stretch and the H₂O-wag modes in the intermediate state (peaks C and B in Figure 4.1.) This change in the intermediate state changes the Franck-Condon factor for the ionization step and results in the observed changes in vibronic activity in the ZEKE spectrum. For an allowed transition such as in this case, one generally expects to see an enhancement in the relative intensity of the corresponding mode in the ZEKE spectrum. This led to the assignment of the peaks labeled 1¹ and 4¹ respectively as the stretch (240 cm⁻¹) and wag (328 cm⁻¹) excited modes in the ion. A weak progression (seen repeatedly in scans of this spectrum) is also attributed to the water torsional motion (5¹ = 84 cm⁻¹). The other three modes attributed to the bonding of water to phenol show up weakly in Figure 2b and can be seen more easily in Figure 4.4, which shows smaller segments of the spectra, scanned more slowly to give improved detail and reduced noise. Two phenol ring modes ($\nu_{6a} = 517 \text{ cm}^{-1}$ and $\nu_{12} = 812 \text{ cm}^{-1}$) are also seen clearly in the spectra in Figure 4.2.

Figure 4.4. Pump-probe ZEKE spectra of the phenol-H₂O complex via excitation of the (a) zero and (b) O-H-O stretch vibrational levels in the intermediate singlet state.

Other than for ν_1 and ν_4 , the above assignments were based upon observed and calculated isotope effects. The spectrum of the d₃-phenol-water complex is shown in Figure 4.5. The frequencies of the six hydrogen-bonding modes were calculated for the proto and deuterioisotopes using two standard potential functions. Since these functions have been derived for ground state complexes, the main utility of the calculations is to give the isotope effects and a crude estimate of the frequency order. The observed and predicted isotope effects are internally consistent.

These results establish a benchmark for the development and evaluation of bonding theories (via comparisons with *ab initio* calculations) and semiempirical potential energy functions describing the bonding in cationic hydrogen bonded complexes. Even without the aid of calculations, the results provide direct insight into the bonding in the ion. In the cases where there are known values of the vibrational frequencies in the ground or excited singlet state of the neutral cluster, the corresponding frequencies in the ion are significantly higher. This is not unexpected since the

Figure 4.5. Pump-probe ZEKE spectra of the d_3 -phenol-water complex via excitation of the zero vibrational level in the intermediate singlet state.

binding energy in the ion is roughly three times that in the neutral complex. The possibility for proton transfer in the cluster ion (or other large changes in bonding and structure) suggests the existence of at least two tautomers that may have similar energies. The relatively strong origin and sparsity of the observed spectra clearly show that the ionization leads to a tautomer of geometry very similar to that of the ground state. Otherwise, one expects to see a very dense and complex spectrum corresponding to large changes in vibrational quantum numbers.

References

1. M. Sander, L. A. Chewter, K. Müller-Dethlefs, and E. W. Schlag, *Phys. Rev. A* **36**, 4543 (1987).
2. W. Habenicht, R. Baumann, K. Müller-Dethlefs, and E. W. Schlag, *Ber. Bunsenges. Phys. Chem.* **92**, 414 (1988).
3. G. Reiser, W. Habenicht, K. Müller-Dethlefs, and E. W. Schlag, *Chem. Phys. Lett.* **152**, 119 (1988).
4. A. Oikawa, H. Abe, N. Mikami, and M. Ito, *J. Phys. Chem.* **87**, 5083 (1983).
5. R. J. Lipert, and S. D. Colson, *J. Chem. Phys.* **89**, 4579 (1988).
6. R. J. Lipert, and S. D. Colson, *J. Phys. Chem.* **93**, 135 (1989).
7. S. L. Anderson, L. Goodman, K. Krogh-Jespersen, A. G. Ozkabak, R. N. Zare, and C. Zheng, *J. Chem. Phys.* **82**, 5329 (1985).
8. G. Reiser, O. Dopfer, R. Lindner, G. Henri, K. Müller-Dethlefs, E. W. Schlag, and S. D. Colson, *Chem. Phys. Lett.* **181**, 1 (1991).
9. J. R. Reimers, and R. O. Watts, *Chem. Phys.* **85**, 83 (1984).

5. Ions at Surfaces

B. Maschhoff,^(a) J. P. Cowin, B. Ellison,^(b)
J. Biesecker,^(b) and T. Orlando

Ionic species and processes at surfaces are important, yet understanding them is made difficult by a lack of structural information and adequate sources. In 1991, we made a major step toward improving this situation by constructing a new system, and establishing a crucial collaboration.

The new system has equipment to measure the structure of ions at surfaces and complex interfaces. Electron-stimulated desorption, ion angular distributions (ESDIAD) will explore multilayer water structures and ion-water clustering on surfaces. Ion scattering spectroscopy will determine the identity of the outermost layer of atoms exposed to the vacuum. X-ray photoemission will identify the ionic charge state of adsorbates and the chemical environment. This new capability will be complemented by more sensitive TOF capabilities than now exist, and by the ability to explore desorbing ions (plus or minus), or neutrals, including internal state spectroscopy.

A crucial feature of the new system is a versatile ion source, which is capable of producing intense beams of positive or negative molecular ions for deposition at very low energies. This source is being built by our University of Colorado collaborators, who are experts at negative ion generation. Together, the system

(a) Northwest Colleges and Universities Association for Science (NORCUS) postdoctoral appointee.

(b) University of Colorado collaborators.

provides the ability to recreate the complete structure of the electrochemical double layer and measure its properties.

The first experiment we are undertaking in this system is a time resolved study of the lifetime of metastable ions at surfaces, like Cl_2^- . A first laser pulse (picoseconds or nanoseconds) will move an electron from the substrate to an adsorbed Cl_2 . A second delayed pulse will either remove the electron, or place this bound anion in an internally repulsive state. We will observe desorbed electrons, Cl^- , Cl or Cl_2 as a function of pulse delay and wavelength. Spacer layers between the surface and the Cl_2 will allow us to change the lifetime of the intermediate anion from probably femtoseconds to milliseconds.

6. Spectroscopy and Dynamics of Molecular Clusters as a Route to Understanding Solvent Influences in Chemical Reactions

R. Knochenmuss, D. Ray, and W. Hess

This part of the MSRC effort is aimed at using laser spectroscopic methods in both the frequency and time domains to study clusters of solvent molecules around or on prototype solutes. Since these clusters can be synthesized in sizes spanning the range from isolated molecules to bulk-like microcrystals or droplets, it is expected that molecular and bulk properties can be explicitly correlated via the intermediate-sized clusters. Effects of solvation on reactivity over this size range will be probed by initiating photochemical reactions in the clusters with laser pulses. The reaction of primary immediate interest is one of the most fundamental and widespread in all chemistry, the transfer of protons between molecules.

Initial work in this direction has been performed^(1,2), but much remains to be learned. The strategy is to combine as many methods as possible to obtain a comprehensive picture of the events that occur in solvent-induced or -influenced reactions, and their time evolution, with the goal of understanding the underlying governing principles. To this end, a variety of instruments are being constructed and/or made suitable for combination in a multifaceted experiment.

The central instrument is a molecular beam machine in which the relevant clusters can be synthesized. This

instrument is nearly ready for use. The remaining critical part is a valve that can be heated to a few hundred degrees Celsius, and is corrosion resistant. Such valves have been recently constructed and are being installed.

Initial cluster studies will utilize nanosecond lasers to perform resonant two-photon ionization (R2PI), for obtaining cluster-selective absorption spectra, and laser-induced fluorescence (LiF) and fluorescence excitation spectroscopies. For R2PI experiments a Bruker TOF-1 reflectron-type mass spectrometer will be used. This instrument is operational, and been interfaced to an appropriate data acquisition system. Variants on the simplest R2PI experiment will include 2-color methods for fragmentation control, isomer- and phase-specific R2PI, threshold ionization spectroscopy, and R2PI detected hole-burning. The LiF/excitation part of the apparatus is under construction and will initially be capable of measuring dispersed fluorescence with nanosecond time resolution.

Also using the nanosecond lasers, unconventional spectroscopic methods will be applied and developed for use on the clusters. Methods under consideration are coherent anti-Stokes Raman spectroscopy; degenerate four-wave mixing, either alone or in conjunction with stimulated emission pumping; resonance-enhanced Raman multiresonant four wave mixing; and infrared-optical double resonance.

Time-domain experiments depend on the availability of high-power lasers that have pulsewidths of a few picoseconds. Low-power Nd:YAG-pumped dye lasers of this type are in operation; a 20-Hz regenerative amplifier is available to increase the pulse energies to appropriate levels, and will be set up near the cluster source for a variety of experiments.

First experiments will combine fast excitation with high-speed detection systems to measure the temporal characteristics of the fluorescence into the subnanosecond range. Afterward, a variety of pump-probe type experiments are planned, which should begin to provide information about the dynamics and nature of the proton-transfer reaction coordinate. Many of these experiments take advantage of the fact that the second photon in an R2PI experiment can be delayed in time with respect to the first, which has initiated the proton transfer reaction. The change in spectral properties with this time delay then give information about the evolution of the reaction event. Structural information

on the clusters will be available through rotational coherence spectroscopy, and intramolecular vibrational relaxation can be studied if quantum beats prove detectable. Picosecond versions of the multiwave, nanosecond experiments may also prove valuable, again using time delays between pulses to probe the dynamics of the reaction.

Finally, collaborations with the MSRC theoreticians are expected to yield *ab initio*, semiempirical and molecular dynamics results that will assist in energetic, structural, and dynamical interpretation of experiments.

References

1. For cluster studies of naphthols see: R. Knochenmuss, and S. Leutwyler, *J. Chem. Phys.*, **91**, 1268 (1989); and T. Droz, R. Knochenmuss, and S. Leutwyler, *J. Chem. Phys.*, **93**, 4520 (1990), and references therein.
2. For studies of phenol clusters see: R. J. Lipert, and S. D. Colson, *J. Chem. Phys.*, **93**, 3801 (1989); C. Jouvot et al., *J. Phys. Chem.*, **94**, 5041 (1990); and references therein.

7. Surface Chemistry of Chlorinated and Fluorinated Hydrocarbons Via Photons and Hot Electrons

M. Kwini,^(a) M. Iedema, and J. P. Cowin

Surface reactions play a critical role in environmental problems in the soil, and on atmospheric dust, sols, and ices. The chlorinated and fluorinated hydrocarbons are especially important, as many of the former are persistent, prevalent groundwater contaminants, while both are important sources of stratospheric halogens that now threaten our ozone layer. Crucial to the destruction of these molecules is the cleavage of the halogen bond. These molecules can be cleaved both by electron rearrangement via ultraviolet photon adsorption, or by electron attachment, in the gas phase. Our recent studies⁽¹⁾ for adsorbed molecules show how intertwined these two mechanisms can be.

(a) NORCUS graduate student at PNL, from the University of California at Santa Barbara, funded by the National Science Foundation (NSF).

If a photon excites a substrate electron, as in Figure 7.1, this electron could be resonant with the antibonding affinity orbital of an adsorbed chlorinated molecule. The anion formed should dissociate, similar to dissociative electron attachment in the gas phase. The electron transfer mediated fragmentation should occur at wavelengths where the molecule shows no direct transitions. Thus, we should by wavelength studies be able to isolate the electron transfer mediated fragmentation from any direct process. Also the direct and indirect processes should produce different fragment kinetic energies.

Figure 7.2 shows the results for CH_3Cl adsorbed on a clean Ni(111) surface, upon irradiation.⁽²⁾ Shown are the arrival times at the mass spectrometer of the ejected CH_3 radicals. For 193-nm laser irradiation we see that two kinds of methyl radicals are produced. The fast 1.5 eV methyls come from the directly excited molecule, and the slower 0.6 eV methyls come from the dissociative electron attachment process. At 248 nm, the molecule is not directly excitable, and correspondingly only the 0.6-eV methyl peak is observed. Note the results for irradiating increasing thicknesses of CH_3Cl . Only the topmost layer of CH_3Cl can contribute to the narrow TOF features. Note that as expected, after 8 to 15 layers, the hot

Figure 7.1. Electron transfers quenching or promoting cleavage.

Figure 7.2. CH_3 radicals photoejected from CH_3Cl -covered nickel.

electrons can no longer reach the topmost layer, evidenced at high coverages by the elimination of the slower of the two peaks. We also find that, due to quenching, both the direct and indirect channels are dramatically reduced in the first 1 to 2 monolayers from the surface. The antibonding electron hops to the substrate faster before dissociation occurs.

The above results⁽³⁾ gave, for the first time, a clear, conclusive demonstration of the role that electron transfers play in laser surface chemistry, both in promotion and inhibition. We are now beginning a study to probe the photodestruction of chlorinated hydrocarbons seen to occur when contaminated water is passed over TiO_2 in the presence of sunlight. Researchers at Sandia National Laboratories in Livermore and at the Solar Energy Research Institute

have demonstrated that this photocatalytic process is a practical means of cheaply treating water using solar collectors to concentrate the light. We hope to develop an understanding about the mechanisms for this process, concerning which there is much debate.

References

1. E. P. Marsh, M. R. Schneider, T. R. Gilton, F. L. Tabares, W. Meier, and J. P. Cowin, *Phys. Rev. Lett.* **60**, 2551 (1988).
2. P. Avouris, and R. E. Walkup, *Ann. Rev. Phys. Chem.* **40**, 1 (1989); P. Nordlander, and J. C. Tully, *Phys. Rev. Lett.* **61**, 990 (1988).
3. E. P. Marsh, T. L. Gilton, W. Meier, M. A. Schneider, and J. P. Cowin, *Phys. Rev. Lett.* **61**, 2535 (1988); E. P. Marsh, F. L. Tabares, M. R. Schneider, T. L. Gilton, W. Meier, and J. P. Cowin, *J. Chem. Phys.* **92**, 2004 (1990).
4. M. R. Schneider, C. P. Dehnhostel, T. L. Gilton, and J. P. Cowin, *J. Appl. Phys.* (submitted).

8. Molecular Dynamics at the Water/Solid Interface

M. Kwini,^(a) M. Iedema, and J. P. Cowin

Reactions at the water/solid interface are crucial to understanding both soil and atmospheric environmental reactions, many of which are photostimulated and/or involve an electron transfer. Our water-layer model systems provide a uniquely direct approach to these problems, and to the general field of electrochemistry. Monolayer-by-monolayer deposition using molecular beams gives the opportunity to create highly structured, layered surfaces. This process is done at sufficiently low temperature to prevent gross vertical diffusion.

In a previously completed study of electron transmission⁽¹⁾, we used the molecule CH_3Cl as a "detector" of electrons. In a separate study,⁽²⁾ we showed that 248-nm photons caused CH_3Cl to dissociate only via the photo-stimulated hot substrate electrons, causing dissociative electron attachment. By placing spacer

(a) NORCUS graduate student at PNL, from the University of California at Santa Barbara, funded by the National Science Foundation.

layers of water between the surface and the CH_3Cl , we can measure the distance that the electrons are able to propagate through the water (Figure 8.1).

At 50 K, we deposited on a Ni(111) surface various thicknesses of water, upon which was deposited one monolayer of CH_3Cl . We then irradiated the surface with a low-power, 248-nm excimer laser. We measured the ability of photoexcited electrons with energies near the vacuum level to propagate through the water by observing either the yield of CH_3 in the normal direction, or by measuring via thermal desorption spectrometry the decrease in the surface CH_3Cl (Figure 8.2). The sharp increase for one monolayer of water spacer is due to the water preventing the molecular anion's electron from returning to the substrate on the time scale of the fragmentation. After peaking at two monolayers of water, the signal then decayed roughly exponentially with the $1/e$ distance of about two monolayers. This phenomenon is indicative of a very strong interaction of the electrons with the water spacer. The distance behavior in Figure 8.2 measured by the loss of CH_3Cl is roughly in agreement with the fragment TOF yield.

Figure 8.1. Ultrahigh vacuum (UHV) water/solid electron transfer model system.

Figure 8.2. Electron transmission versus water depth.

Our measurement of the mean free path of low-energy electrons in water is not the first.⁽³⁾ However, it is unique in that it measures the transmission in the chemically important energy regime, with well known structures, and is extremely flexible.

In related work, which has been submitted for publication⁽⁴⁾, we performed a preliminary study of femto-second solvation dynamics. The energy of the methyl fragment is about 0.6 eV, much faster than the 0.2 eV seen for gas-phase dissociative electron attachment. This extra energy comes from that part of the solvation that can occur on the 20-femtosecond time scale of the fragmentation.

By placing a single monolayer of CH_3Cl on hexane, CH_3Cl , or water, the energy of the ejected CH_3 ranges from 0.4 to 0.7 eV. We are doing further studies, and beginning a theoretical collaboration, to uncover the mechanisms of this femtosecond solvation.

References

1. T. L. Gilton, C. Dehnhostel, M. R. Schneider, and J. P. Cowin, *J. Chem. Phys.* **91**, 1937 (1989).
2. E. P. Marsh, T. L. Gilton, W. Meier, M. A. Schneider, and J. P. Cowin, *Phys. Rev. Lett.* **61**, 2535 (1988).
3. R. N. Barnett, U. Landman, and A. Nitzen, *J. Chem. Phys.* (in press).
4. M. Kwini, T. Gilton, M. Iedema, and J. P. Cowin, *J. Phys. Chem.* (submitted).

9. Single-Molecule Chemistry

S. A. Joyce, L. Bumm,^(a) and J. P. Cowin

Several groups have demonstrated the ability of the scanning tunneling microscope (STM) to observe single molecules, interpret their chemistry, and even arrange the molecules on an atomic scale (Figure 9.1). This is an ideal approach for unraveling the complex chemistry that environmental restoration entails. The most important topics to explore are the role of defect sites in surface reactions of insulators, the details of charge transfer processes at surfaces, and how surface bonding affects adsorbate chemistry. Applications include contaminant chemistry on soil particles and molecular-specific sensors.

Crucial are scanning surface probes with capabilities to control the surface temperature, to deliver reagents *while observing*, and to stimulate the surface reactions using thermal and laser excitation means. Scanning probes required are STMs in ultrahigh vacuum; a "users" scanning force microscopy of insulators in fluids; scanning optical probes of isolated adsorbates, especially on insulators; and cryogenic and high temperature STMs.

Several staff members are involved in a coordinated way. Scanning optical and atomic force microscopy projects are to begin in FY 1992, led by other scientists in the MSRC. Progress in 1991 occurred in both STM studies of charge transfer processes at interfaces, and in cryogenic STM.

(a) NORCUS postdoctoral appointee.

Figure 9.1. Schematic of atomic scale imaging of adsorbates and defects with a scanning probe microscope.

Dr. Stephan A. Joyce, a highly experienced surface scientist, was hired to participate in this research. He will play a pivotal role in facilitating all the scanning surface probe work, and will personally develop the cryogenic STM system. Postdoctoral appointee, Lloyd A. Bumm, also began experiments using the commercial ultrahigh vacuum room temperature STM.

The cryogenic STM is essential for studying weakly adsorbed molecules, which are commonly encountered in aqueous or gaseous interfaces with solids. At room temperature these adsorbates diffuse from site to site on the surface so fast that they are virtually invisible to the STM. At 20 K or below, we will be able to determine the preferred binding sites of adsorbates on complex surfaces, and to watch reactions induced by lasers. Design principles are currently being explored. After consultation with the several other groups in the world pursuing cryogenic STM, design and construction should commence in January 1992.

The room-temperature STM can observe adsorbates covalently bonded to the surface. During FY 1991, the optical arrangement, which will bring in light to just

the STM tip region, was designed, as well as other improvements to the system. These are being assembled.

The first experiment we will do is to explore back-transfer-mediated quenching of surface reductions. Charge transfer reactions (reduction or oxidation) can be greatly inhibited if before the charge transfer can be "consummated" via bond cleavage or other permanent step, the transferred electron hops back to where it came from. Our experience with macroscopic probes of back transfer processes suggests the rate should be proportional to the densities of states, and thus extremely sensitive to whether the molecule is adsorbed at a step edge, a terrace, or an isolated point defect. By observing many different adsorbate molecules, and gathering simple statistics on the average photon flux needed to cause a successful photostimulated charge transfer reaction, we will infer the site dependencies of back transfer. Adsorbates of choice will be molecules like $\text{CH}_2\text{ClCH}_2\text{CH}_2\text{S}$.

10. Photodissociation Dynamics of Molecules and Clusters

W. P. Hess and S. P. Sapers

The photodynamics of molecules and molecular clusters will be investigated using a newly constructed photofragment imaging apparatus. This device is designed to image the velocity distributions of atomic or molecular photofragments. The design includes access ports for LiF and photoelectron spectroscopy. A variety of cluster sources for creating neutral and ionic clusters (positive and negative) from both volatile and refractive samples will also be constructed.

An initial project for the imaging apparatus involves the generation of silicon oxide clusters to be used to model the microscopic properties of sand. The spectroscopy and dynamics of silicon oxide clusters must be determined prior to using these species as geochemical models. Silicon dioxide molecules have been produced by a laser ablation-type source and the feasibility of producing larger silicon/oxide structures with ablation sources will be investigated. This preliminary project involves the development of ablation sources to produce silicon oxide clusters, mass spectroscopy to determine cluster sizes, and optical spectroscopy to determine the structure of the smaller clusters. These

studies will determine the suitability of silicon oxide clusters for modeling geochemistry and are also of basic interest.

Later studies will investigate the properties of silicon oxide clusters combined with water and other solvent molecules. The bond strength of clusters bonded to water and other solvent molecules is important in understanding the diffusion rate of pollutants from a spill site. A measure of that bonding may be obtained from photodecomposition studies. The method involves creating a cluster consisting of a covalently bonded silicon oxide molecule in contact (by Van der Waals forces) with a model solvent molecule. The Van der Waals bond is then broken by exciting vibrational modes in the cluster or solvent molecules. We will optically excite the clusters using direct infrared absorption or stimulated Raman pumping. By exciting the available vibrational modes in the cluster or the solvent molecule, the energetic threshold for bond dissociation may be bracketed. It will be important to produce cold, well-defined cluster molecules for these studies.

Photodissociation of mass selected Van der Waals molecules will be studied using photofragment imaging. A study of the metastable decay of benzene clusters has recently been reported by Sengteller, Selzle, and Schlag.⁽¹⁾ Their article describes a method for stopping ions that is suitable for studying the dissociation rate of optically excited Van der Waals molecules. In this technique, mass selected ionic clusters may be stopped at a space fixed point for tens of microseconds. Dynamical studies within this time window may then be performed. Following photodissociation of ionic molecules or clusters the photofragment angular distribution may be imaged. The fragmentation pattern is indicative of the detailed dissociation dynamics. The extent of cluster fragmentation following ionization is an open question that may be studied by this technique.

Reference

1. S. Sengteller, H. L. Selzle, and E. W. Schlag, "Study of the Metastable Decay of Single Size Benzene Clusters by an Ion Stopping Technique," *Z. Naturforsch.* **45A**, 169 (1990).

11. Apparatus for Time-Resolved Spectroscopy

D. M. Friedrich, M. O. Trulson, and A. G. Joly

In order to study the structural evolution of molecules in excited electronic states undergoing intramolecular charge transfer or proton transfer, a unique high-efficiency spectrometer for time-resolved ultraviolet (UV) resonance Raman studies has been constructed. The spectrometer consists of a single-stage, 1-m, Czerny-Turner monochromator interfaced at its exit plane to a 2-m, folded-path spatial filter. This arrangement permits efficient rejection of light at the excitation laser wavelength and allows efficient collection of the dispersed Raman spectrum to within 100 cm^{-1} of the laser line in the backscattering collection mode in the quartz UV. In many experiments, the use of this spatial filter eliminates the need for double or triple monochromator configurations, which have considerably lower throughput. The entrance to the spatial filter is fitted with an interchangeable mask and slit assembly, enabling use of the instrument either as a spectrograph with a multichannel detector or as a scanning spectrometer coupled through an exit slit to a photomultiplier. This spectrograph is currently being used to record UV resonance Raman spectra of 4-(dimethylamino)benzonitrile (DMABN), a model compound for twisted intramolecular charge transfer processes, and 3-hydroxy-2-methyl-4-pyrone, which is the central ring structure of a class of compounds showing excited-state intramolecular proton transfer (flavones and chromones).

An ultrafast laser system for excited state dynamical studies became fully operational in 1991. This system consists of two amplified dye lasers with a repetition rate of 540 Hz and pulse widths of 0.1 ps and 0.5 ps. In addition, a second, lower-repetition-rate ultrafast system was also installed. Amplified 20-Hz, 0.5-ps pulses will be used to study dynamics in molecular cluster beams. The unamplified output from the dye laser in this second system will be used in the time-correlated, single-photon counting fluorescence lifetime apparatus described in the next section.

12. Time-Resolved Fluorescence as a Probe of Molecular Dynamics

G. R. Holtom

Technologies for measuring dynamics of fluorescent materials have been under steady improvement at many laboratories for three decades. Applications include biophysical measurements such as energy transfer reactions, measurements of molecular size, binding of deoxyribonucleic acid (DNA), and determination of free versus hindered motions in molecular subunits. Useful capabilities for solution-phase chemistry include solvation dynamics and twisted intramolecular charge transfer (TICT) processes. Gas-phase experiments with molecular beams are used to study aggregation, solvation, and state-to-state energy transfer. A basic technology for fluorescence measurements is time-correlated single-photon counting (TCSPC). This technique is simple, versatile, and sensitive, and permits studies over perhaps 4 orders of magnitude in time, with a resolution approaching 10 ps.

The most general technology uses a cavity-dumped pulsed dye laser to excite the sample. Fluorescence at the desired wavelength is selected by a monochromator and detected by a photomultiplier. Timing electronics create an analog signal that is stored and digitized by a multichannel analyzer for computer analysis and viewing. With excitation provided by a mode-locked YAG laser operating at the second or third harmonic (green or UV output), the dye laser covers the entire visible region and extends into the near infrared. Power is high enough to frequency double into the far UV region, so that the total excitation region spans 210 nm to more than 800 nm. Excitation in the region from 210 nm to 280 nm provides capabilities not available in many laboratories: small-molecule studies; direct excitation of unmodified DNA bases; and selective excitation of the fluorescent amino acids tyrosine and phenylalanine as well as tryptophan; all are possible in the deep UV region. Since the laser operates at megahertz repetition rates, the energy per pulse is small, thereby minimizing undesirable degradation of the sample.

Data collection is simple and usually requires from 1 to 10 minutes per sample, so that a substantial number of experiments can be carried out in a day. Analysis of data can be done on line, with preliminary information available in a matter of minutes. This setup is useful because problems can be identified before a great deal of time is expended on collecting a large data set.

Two basic kinds of information are available. One experiment measures emission strength as a function of time, yielding population information. Since only electronically excited states can fluoresce, emission intensity is directly proportional to excited-state population. This mode is useful for measuring state-to-state energy flow and kinetics.

In another type of experiment, the decay of anisotropy is determined from the polarization of the emitting state. If the sample is excited with polarized light, then radiation emitted parallel to the excitation differs from radiation emitted perpendicular to the excitation. Changes in the emission dipole are measured, which can be due to energy transfer to a new state or more often to overall rotation of the molecule which, in turn, causes diffusion of the direction of the molecular dipole.

In either case, the raw data is not directly useful for quantitatively extracting molecular information. One usually fits a model to the raw data, making iterative adjustments to the parameters of the model until the calculated curve matches the experimental results. A composite response of the excitation source (dye laser pulse width), the detector response, and electronics contributions is contained in the instrument function. Usually it is important to model data even in the time scales where the molecular response is important (10 to 100 ps). In order to handle this exactly, a convolution integral must be performed. The details of this are not important except to note that useful experimental results are not carried by the raw data but instead by analytic functions that model the raw data.

Work on improving time-resolved fluorescence analysis is progressing on several fronts: simplifying the operation of laser systems, automating data collection, and improving the user interface or software. Adding models for nonexponential decays will be done on an as-needed basis when required for new experiments.

13. Stimulated Cluster-Surface Chemistry

T. M. Orlando and S. E. Barlow

Investigating the physical and chemical properties of clusters of metallic, nonmetallic, semiconductor, and insulator materials is currently one of the most active frontiers of chemistry and physics⁽¹⁻⁴⁾. Much of the interest stems from the fact that these nanosystems may provide an avenue that bridges the gap between the gas and condensed phase. Since relatively small clusters have large surface-to-volume ratios and large step and defect densities, it is reasonable to expect them to be highly reactive. In fact, there are several studies which clearly demonstrate variations in cluster reactivity patterns with size^(1,2) and in some cases with electronic structure.⁽³⁾ However, for sufficiently large clusters, bulk-like properties are expected to emerge and thus these nanomaterials (nanocrystals) may have chemical properties that are intermediate between compact clusters and bulk surfaces. These nanocrystals may therefore provide a new way to systematically study surface mediated chemistry or interphase reactions. We intend to study the nanomaterial regime where convergence to bulk-like properties is expected to occur. Particular emphasis will be placed on studying thermal and *stimulated* reactions on nanocrystals of either ices (such as water), or oxides and minerals. These studies are expected to provide insight to environmental problems involving heterogeneous reactions on frozen particles in the upper atmosphere or on subsurface soil particles.

Stimulated reactions on bulk surfaces are well known and are generally caused by the impingement of energetic particles such as photons or electrons. Desorption,⁽⁵⁾ dissociative ionization, reactive scattering⁽⁶⁾ and ion-molecule reactions⁽⁷⁾ between coadsorbates can be initiated by localized electronic excitations in the adsorbate-substrate complex. For example, Figure 13.1 shows the translational energy of the NO₂ desorbate that was produced⁽⁶⁾ during electron-beam irradiation of O₂ coadsorbed with NO on Pt(111). A detailed description of the physics involved in these types of stimulated reactions is presently lacking. This is particularly true with respect to final state interactions that involve reneutralization of ions on or in close proximity to the surface. We intend to probe stimulated radical and ion-molecule reactions on bulk and nanocrystal surfaces using low-energy ion beams and sensitive laser techniques such as resonance enhanced multiphoton ionization spectroscopy. This

Figure 13.1. Time-of-Flight E_{trans} distribution of the NO(d) stimulated reaction product measured by monitoring the NO($v=5$) and O(3P_2) photofragments. The inset shows the excitation energy threshold for NO₂ formation.

approach should help provide an understanding of chemical reactions that occur in radiation environments.

Bombardment of multilayer ice surfaces and solid state substrates with energetic electrons and ions generally results in sputtering and erosion of the surface. Therefore, controlled sputtering can be used as a method of producing large quantities of heteronuclear neutral and charged gas-phase clusters. Despite the usefulness of sputtering phenomena, controlling this process is non-trivial since the detailed mechanisms involved in sputtering are not well understood. This is largely due to the fact that the particle-surface interaction involves a complex superposition of momentum transfer (in the case of ion impingement), charge transfer (harpooning and neutralization), as well as electronic and chemical effects. Efforts to develop and optimize ion-beam sputtering using RF driven magnetron discharge devices and electron beam sputtering using magnetic confined plasma sources are currently under way. Methods of generating solvated cations and/or dications using electrospray technology are also under development for future studies on ion-exchange mechanisms.

A radiofrequency guided low energy ion-beam machine that will incorporate these cluster sources has been designed and will serve as the prototype instrument for nanocrystal reaction studies. This apparatus will be significantly improved as part of the Environmental and Molecular Sciences Laboratory mission by installing an ion flow tube (DRIFT) system that should allow shape (geometrical) selection of the nanocrystal beam. The critical roles of geometry and electronic structure on reactivity can then be systematically probed.

References

1. S. A. Ruatta, L. Hanley, and S. L. Anderson, *J. Chem. Phys.* **91**, 226 (1989).
2. M. Jarrold, *Science*, **252**, 1085 (1991).
3. E. K. Parks, B. J. Winter, T. D. Klotz, and S. J. Riley, *J. Chem. Phys.* **94**, 1882 (1991).
4. J. L. Elkind, J. M. Alford, F. D. Weiss, R. T. Laaksonen, and R. E. Smalley, *J. Chem. Phys.* **87**, 2397 (1987).
5. A. R. Burns, E. B. Stechel, and D. R. Jennison, *Phys. Rev. Lett.* **58**, 250, (1987).
6. T. M. Orlando, A. R. Burns, E. B. Stechel, and D. R. Jennison, *J. Chem. Phys.* **93**, 9197, (1990).
7. L. Sanche and L. Parenteau, *Phys. Rev. Lett.* **59**, 1436, (1987)

14. Spectroscopy of Small Cluster Molecules in Low-Temperature Beams

S. W. Sharpe, T. A. Hu, and E. L. Chappell

Small, weakly bonded clusters of molecules and atoms play an important role in experimental and theoretical chemical physics research. These systems are used to elucidate chemical reactions on a microscopic level, where the orientation and quantum states of the reactants are fully characterized.^(1,2) Clusters provide a unique opportunity to study energy transfer and internal vibrational relaxation (IVR) processes.^(3,4) In

addition, small clusters (i.e., $\text{H}_2\text{O}-\text{H}_2\text{O}$) are implicated in the atmospheric radiation balance and may play a role in global warming.⁽⁵⁾

Regardless of the chemical identity, it is the hydrogen-bond and van der Waals or dispersive forces that are primarily responsible for the characteristics of these clusters. Fundamental, quantitative knowledge about the nature and consequences of these long-range intermolecular forces can be obtained from the spectroscopy of such clusters, in which the weak, anisotropic interactions that dictate dynamics and geometry are often exposed in the form of rich and varied signatures.

Researchers have a number of techniques at their disposal, which can be used to examine small clusters. Although mass spectrometry, electronic spectroscopy, and electron diffraction are all powerful techniques, microwave and infrared (IR) spectroscopies are unsurpassed in their ability to obtain high spectral resolution and accurate data. One particularly promising IR technique for exploring the $2400\text{--}300\text{ cm}^{-1}$ region utilizes rapidly tunable diode lasers and pulsed, slit, supersonic expansions. This fast-scan technique, developed by DePiante et al.⁽⁶⁾ and Sharpe et al.⁽⁷⁾, offers rapid data acquisition, high sensitivity, high resolution, continuous coverage and most importantly, access to the majority of vibrational motions including intramolecular bending modes.

A second-generation molecular beam machine is currently being completed and will be used in conjunction with the rapid scan technique. This beam machine consists of an expansion chamber that houses the source, followed by two differentially pumped chambers (Figure 14.1). Presently the machine is configured for simultaneously conducting direct IR absorption and mass spectroscopic analysis. Provisions have also been made for incorporating a liquid-helium-cooled bolometer. The source is a novel pulsed, slit supersonic nozzle, which is schematically represented in Figure 14.2. The effluent generated via a slit source has a number of properties that can be used advantageously in high-resolution IR spectroscopic studies. These advantages include Doppler widths on the order of 20 MHz, rotational temperatures below 10 K, and a collision-free environment. By combining a slit nozzle source with a tunable IR diode laser, samples that are either spectrally too congested or thermodynamically too unstable to study at room temperature become amenable to spectroscopic investigation.

Figure 14.1. The molecular beam machine consists of three chambers. The first chamber contains the source and is pumped by an 18,000 liter/sec diffusion pump. The second chamber is contained within the first chamber and pumped by a 2,500 liter/sec diffusion pump. The third chamber is also pumped by a 2,500 liter/sec diffusion pump and can house either a quadrupole mass spectrometer or a bolometer.

Current research involves the production and detailed study of water clustered with water, carbon dioxide, argon, and several small organic molecules. These studies are motivated by the need for accurate data on the intermolecular forces between atmospheric gases. In addition, equipment and techniques are being developed to produce and study the ubiquitous silicon dioxide molecule and small clusters of SiO_2 . Presently, there is a conspicuous absence of experimental data concerning the structure and force field of SiO_2 . It is thought that gas-phase SiO_2 is a linear molecule, being isoelectronic with CO_2 .⁽⁸⁾ This conclusion is also inferred from the electrostatic molecular beam deflection experiments of Kaufman et al.⁽⁹⁾ With the completion of this new beam machine, it should be possible to perform high resolution IR spectroscopic analysis of SiO_2 and small clusters containing SiO_2 .

A typical experiment consists of expanding a 1:99 mixture of $\text{N}_2\text{O}:\text{Ar}$ at a backing pressure of 0.8 atmosphere through the slit nozzle. The output of a tunable diode laser is then multipassed through the supersonic effluent, and the attenuation of laser power is measured as a function of laser frequency. This results in a direct absorption spectrum of the type displayed in Figure 14.3. The $\text{N}_2\text{O}:\text{Ar}$ cluster was chosen as a first

Figure 14.2. A novel design pulsed, slit supersonic nozzle is used as the source. The supersonic effluent emanates from a 11.4 cm by 150-micron slit. Three solenoids, operating in parallel, are used to momentarily lift a Viton seal from the high pressure side of the slit.

system for study for three reasons: 1) existing spectra of this system allows us to compare the performance of our apparatus with those obtained in other laboratories; 2) nitrous oxide is a greenhouse gas, and as such the IR spectra of the isolated and clustered species are relevant to atmospheric modeling programs; and 3) there is general interest in the physics community concerning the weak forces that bond these types of clusters.

References

1. C. Wittig, S. Sharpe, and R. A. Beaudet, *Acc. Chem. Res.* **21**, 341 (1988).
2. C. Jouvet, and B. Soep, *Chem. Phys. Lett.* **96**, 426 (1983).
3. R. E. Miller, *J. Phys. Chem.* **90**, 3301 (1986).
4. G. E. Ewing, *J. Phys. Chem.* **91**, 4662 (1987).
5. J. T. Houghton, *The Physics of Atmospheres*, Cambridge University Press, 1986.
6. A. DePiante, E. J. Campbell, and S. J. Buclow, *Rev. Sci. Instr.* **60**, 858 (1989).
7. S. W. Sharpe, Y. P. Zeng, C. Wittig, and R. A. Beaudet, *J. Chem. Phys.* **92**, 943 (1990).
8. V. H. Schnockel, *Z. anorg. allg. Chem.* **460**, 37 (1980).
9. M. Kaufman, J. Muentner, and W. Klemperer, *J. Chem. Phys.* **47**, 3365 (1967).

Figure 14.3. High-resolution rovibrational spectra of the nitrous oxide dimer. The upper spectrum was taken by tuning the diode laser over the region corresponding to the ν_3 intramolecular stretching fundamental vibration of N_2O . The lower spectrum was taken in a spectral region corresponding to the combination vibration involving the intramolecular stretch (ν_3) plus one quantum of intermolecular bending. The lower spectrum is classically forbidden and is about 500 times weaker than the upper spectrum, but note the generally high quality of the spectrum furnished by our instrumentation.

15. Spectroscopic Studies of the Gas-Catalytic Surface Interface

R. S. Smith and S. D. Colson

Catalytic chemistry has numerous applications in the chemical industry, in environmental processes, and in the basic understanding of chemical reactions. Our research has focused on studying the interaction of gas-phase molecules with catalytically active surfaces. The gas-surface interface is extremely important for determining reaction mechanisms involved in heterogeneous catalysis. In the past year, we have studied two problems: the catalytic oxidative coupling of methane, and the catalytic decomposition of 1,1,1-trichloroethane. The first problem involves the economical and efficient production of petrochemicals and liquid fuels from methane feedstocks. The process is important because methane is an abundant resource and can be obtained by environmentally safe methods. The second problem involves determining the decomposition mechanism of a widely used chlorocarbon solvent. The process may have applications for the safe disposal of such solvents and for enhancing decomposition on surfaces found in nature.

Although the two problems are chemically different, the reaction mechanism for both processes involves the interaction of gas-phase molecules with catalytic surfaces. Our approach to the study of gas-phase molecules present above the catalytic surface involves the use of a microreactor that quenches the reaction via a supersonic expansion into a vacuum. The expansion effectively stops the reaction because of rapid cooling followed by the greatly reduced collision rates inherent with this type of expansion. As a result, even very reactive species exist long enough to be ionized (a variety of techniques can be used) and, thus, detected using laser ionization mass spectrometry (LIMS). The primary advantage of this reactor is the capability for the direct detection of reactive intermediates and radicals. Another advantage of this design is that LIMS allows for the simultaneous detection of products and reactive-intermediates. The use of laser ionization gives the flexibility and sensitivity to detect small amounts of transient species. Furthermore, the small size of the microreactor allows for rapid temperature changes, making experiments over a wide range of temperatures feasible. Experimental results over a wide range of temperatures are necessary to understand the complicated kinetics involved in the process. We have recently modified the apparatus by adding a mass-flow controller and residual gas analyzer (RGA).

These new features give us more control over the gas flow rates and the ability to detect all reaction products (including H_2O , CO , and CO_2).

Our experimental design has been used to study the oxidative coupling^(1,2) of methane over 1% $\text{Sr/La}_2\text{O}_3$. The formation of methyl radicals is believed to be an important intermediate reaction. The first step in the proposed mechanism is the production of methyl from the reaction of methane with the catalytic surface. This step can be followed by the desorption of methyl into the gas phase leading to the formation of higher molecular weight hydrocarbons via radical-radical combination and/or radical-substrate reactions. In many previous experiments, the direct detection of the transient and unstable radical species was not possible. Our experiments are able to directly measure the formation of methyl radicals and other reaction products. Figure 15.1 is a plot of the intensity versus temperature and shows yields of methyl radicals and olefin products. The figure shows that the olefin production and the methyl radical production do not have the same temperature dependence. These data suggest that the formation of gas-phase methyl radical species is not the rate-determining step for the production of olefins.

To further investigate the reaction mechanism we have done experiments using a 50/50 gas mixture of CH_4/CD_4 . A mechanism involving a simple gas-phase methyl radical recombination is not expected to result in isotopically scrambled ethylene products.

Figure 15.2 shows some data from these experiments. The data show the production of all possible ethylene isotopes. This result suggests other reaction possibilities including surface recombination of methyl radicals or the reaction of intermediate species with the catalytic surface. We are presently working to further elucidate this reaction mechanism.

Another project involves determining the reaction mechanism for the catalytic decomposition of 1,1,1-trichloroethane over high surface area alumina (Al_2O_3).⁽³⁾ The research has been done in collaboration with Todd H. Ballinger and John T. Yates, Jr., at the University of Pittsburgh. The data reveal different reaction mechanisms at different temperatures. Figure 15.3 shows the reaction products as a function of temperature. Between 400-600 K the CH_3CCl_3 decomposition produces primarily CH_2CCl_2 and HCl , possibly through a 1,2 elimination mechanism. At

Figure 15.1. Oxidative coupling product distributions as a function of temperature.

Figure 15.2. Ethylene products from oxidative coupling of a 50:50 CH₄/CD₄ gas mixture.

Figure 15.3. Reactant and product intensities versus temperature for the decomposition of 1,1,1-trichloroethane over alumina.

temperatures above 600 K, the yield of CH_2CCl_2 decreases and at very high temperatures, HCl is the only gas-phase product. Analysis of the used catalyst reveals an increase in the amount of carbon and chlorine on the catalyst. These data suggest that at high temperatures the CH_2CCl_2 is being further decomposed into surface carbon, surface chlorine, and HCl. Presently, surface IR spectroscopy is being used to confirm these surface species. Continued work on this problem will involve the determination of decomposition mechanisms for other chlorocarbons and the use of other oxide catalysts.

References

1. E. E. Gulcicek, L. D. Pfefferle, and S. D. Colson, "Multiphoton Ionization Detection of Methyl Radicals from Catalytic Oxidation of Methane," *J. Chem. Phys.* **94**, 7069 (1990).
2. R. S. Smith, S. D. Colson, E. E. Gulcicek, and L. D. Pfefferle, "Methane Oxidative Coupling in a Quenched Reactor: Intermediates and Olefin Products," *J. Chem. Phys.* (submitted).
3. R. S. Smith, S. D. Colson, T. H. Ballinger, and J. T. Yates, "Catalytic Decomposition of 1,1,1-trichloroethane Over Alumina" (manuscript in preparation).

Environmental Dynamics and Simulation

Environmental Dynamics and Simulation

The Environmental Dynamics and Simulation program combines expertise from PNL's Earth and Environmental Sciences Center, programs funded through the Environmental Science Research Center, and the DOE's Integrated Demonstration Programs. The Environmental Dynamics and Simulation program addresses research related to the restoration of contaminated subsurface regions. Activities in the program are designed to develop the molecular-level understanding of contaminant behavior obtained from all of the MSRC program areas into technologies and approaches that can be applied to environmental remediation and restoration problems. A multi-disciplinary approach will use new experimental results and theoretical analyses to provide detailed descriptions of chemical reactions in complex mixtures and at solid-liquid interfaces. Resulting models will predict the fate and transport of contaminant species and provide a scientific basis for remediation schemes.

1. Rational Design of Metal Selective Liquids

B. P. Hay

The remediation of the many types of wastes and sites in the DOE complex will require the ability to tailor separations chemistry to widely varying conditions. Improved technologies are recommended for the separation of radionuclides and/or toxic metals from a variety of matrices such as contaminated soil, contaminated groundwater, stored low-level waste, stored high-level waste, stored mixed waste, and future waste streams from uranium and plutonium processing. A promising strategy for the improvement of metal ion separations is to enhance the performance of existing methods such as solvent extraction, liquid membranes, and chromatography. Metal-specific ligands are frequently a key component of these methods, serving as the extractant, the transfer agent, and the binding site, respectively. However, available ligands do not always exhibit the degree of selectivity needed to apply these methods to existing waste forms. Dramatically improved separations are possible when ligands are

altered to provide increased specificity for the target ion(s), stronger binding affinity for the target ion(s), or both. Unfortunately, the design criteria for tailoring a ligand for a specific task are often either difficult to implement, unavailable, or both.

We are interested in the development and application of molecular models that will allow for the rational design of ligands with improved selectivity and binding characteristics. To this end, we are investigating the use of molecular mechanics for the analysis of steric interactions and conformational strain that occurs within the ligand upon complexation of a metal ion. Current research efforts focus on the class of ligands containing neutral oxygen donor atoms, as they are among the few types of ligands known to exhibit selectivities for cesium and strontium, two radionuclides of concern to the DOE. Graphical representations of the molecular structure of a ligand that is currently being studied as a possible sequestering agent for the selective extraction of strontium from acid aqueous solutions are shown in Figures 1.1 and 1.2.

Figure 1.1. Two-dimensional representation of the molecular structure of cis, syn, cis-dicyclohexyl-18-crown-6. This ligand has been used as a sequestering agent for the selective extraction of strontium from acidic aqueous solution.

Figure 1.2. Three-dimensional representation of a potassium complex that contains the cis, syn, cis-dicyclohexyl-18-crown-6 ligand (generated from a published crystal structure).

2. Molecular-Scale Investigations of Alkali Halides

L. E. Eary, R. L. Erikson, and C. J. Hostettler

The objective of this research is to investigate relationships between molecular-scale interactions and the macroscopic properties of selected alkali crystals and melts. The proposed work involves an examination of how well periodic quantum methods and statistical mechanics methods perform together in calculating thermodynamic and structural properties involving solid-state phase transitions, melting behavior, and phase equilibria in binary alkali halide systems.

Several computational chemistry methods offer the ability to calculate the structural and thermodynamic properties of solids and liquids. Recent advances in *ab initio* quantum methods have involved the extension of computations beyond isolated clusters to periodic crystalline systems. Empirical and semiempirical statistical mechanics methods such as energy minimization, Monte Carlo, and molecular dynamics have also been used to study properties of solids, liquids, and

glasses, including the alkali halides as well as simple oxides and silicates. However, the use of statistical mechanics methods has been limited by the availability of data used to parameterize potential functions. Empirical potential function parameters are often adequate for use in computing structural details such as bond distances and angles to several percent or better; however, obtaining thermodynamic information including reaction enthalpies, and energies of solid-solid and solid-liquid phase transitions has not been as successful.

Because the thermodynamic properties of solids, liquids, and the solid-water interface act to control the distribution and behavior of chemicals in the environment, an evaluation of how well we can predict thermodynamic properties is needed. The choice to examine the properties of the alkali halides was made because simple cubic structures are easiest to model, and a large body of experimental data exists for the B1/B2 phase transition, melting equilibria, structure, and thermodynamics of molten salts. There also is a considerable body of literature regarding molecular dynamics calculations involving alkali halides; however, there have been few periodic quantum calculations carried out for the alkali halides. Finally, the energy differences of the phase transitions for alkali halides are well understood and are of the same order of magnitude (a few to tens of kJ/mole) as many environmentally relevant chemical reactions (e.g., sorption of metals onto oxide surfaces). Thus, the alkali halides provide a convenient vehicle for examining conceptual approaches on how computational methods can be combined to improve our ability to calculate thermodynamic quantities.

Thermodynamic and structural properties of a selected set of alkali halides (LiF, NaF, NaCl) are being calculated using an approach that combines the methods of *ab initio* quantum mechanics and statistical mechanics. Energy-conformation relationships for alkali halide crystals computed using a quantum method for periodic systems will be used to constrain the derivation of pairwise interaction parameters for potential energy functions used by the statistical mechanics models. Energetic and structural differences accompanying solid-state phase transitions and melting equilibria in unary and binary alkali halide systems will be calculated and compared to experimental data. The comparison of the predicted and experimentally measured properties will allow an evaluation of our ability

to compute thermodynamic properties from first principles in a well known and simple system. Such information will be useful in future extensions of this work involving more complicated oxide and hydroxide structures and the interaction of their surfaces with water.

We have started by performing calculations on crystalline sodium chloride by developing an extended basis set. An algorithm was developed to optimize the orbital exponents of an uncontracted basis set for isolated Na and Cl ions. The algorithm, which uses the downhill simplex method in multidimensions, has been adapted to perform the optimization of the Gaussian exponents. This algorithm has been modified and installed on the Sun SpareStation II as an automated driver program that iteratively adjusts the Gaussian exponents and then operates appropriate portions of the CRYSTAL code to achieve a minimization of the calculated Hartree-Fock energy. The result is an optimized set of Gaussian exponents that give a minimum total energy or most stable molecular orbital configuration for the system being investigated. Using our results for the uncontracted basis set optimized only for the isolated Na and Cl ions, we have calculated the enthalpy of formation of crystalline NaCl to within 5% of the calorimetric value using CRYSTAL. We are currently in the process of developing a contraction scheme for the extended basis set and reoptimizing the outer orbital exponents in the NaCl crystal field.

3. Synchrotron Development

J. E. Amonette and S. V. Mattigod

An important aspect of the Environmental Dynamics and Simulation program is the development of expertise in the application of synchrotron radiation to geochemical problems. Synchrotron sources offer very intense, highly collimated, tunable, and essentially monochromatic radiation in the x-ray, UV, and IR spectral regions.

Research incorporating synchrotron techniques can be done on samples under conditions approaching those in the environment. Spectroscopic and diffraction studies can be performed with much higher resolution than by conventional methods. Kinetic studies with resolutions on the order of micro and nanoseconds are possible.

Staff have been participating in the Pacific Northwest Consortium proposal for the Advanced Photon Source at Argonne National Laboratory and have submitted proposals for exploratory work at the National Synchrotron Light Source at Brookhaven National Laboratory.

Macromolecular Structure and Dynamics

Macromolecular Structure and Dynamics

The principal goal of the Macromolecular Structure and Dynamics program is the application of advanced spectroscopic techniques in the analysis of the structure and dynamics of biological macromolecules. Leading-edge nuclear magnetic resonance (NMR) and mass spectrometry (MS) instruments are being designed and developed for high-resolution measurements on the primary, secondary, and tertiary structures of large proteins and nucleic acids in solution. Other applications of these instruments include analysis of trace components in complex mixtures, microimaging of physical and biological structures, measurements of solute flow profiles in heterogeneous materials, such as soil; analysis of gas-phase and solution-phase chemical dynamics; and studies on interfacial adsorption processes and ligand-receptor interactions.

During the past decade NMR spectroscopy has become a widely used technique for determining the 3-D conformation of biological macromolecules in solution, both in their native states and in perturbed states resulting from the interaction of environmental pollutants, drugs, and other substances. Included in the NMR program plan is the development of a 1,000-MHz (gigahertz) spectrometer that will enable a complete structural analysis to be made of macromolecules with masses as large as 40 to 50 kilodaltons. For example, the gigahertz NMR spectrometer is expected to provide new capabilities for investigating the structures of biodegradative enzymes and a variety of large proteins that are involved in the regulation of cell and tissue growth in normal and cancerous states. Many of these molecules cannot be characterized structurally using presently available NMR spectrometers that operate at frequencies up to 600 MHz.

During the past year Dr. Gary P. Drobny was appointed as the NMR Program Head. Dr. Drobny is a Professor of Chemistry at the University of Washington, and will continue part time in this capacity while heading the NMR Program in the MSRC. He will lead the development of the gigahertz spectrometer and oversee the deployment of conventional instruments for both solution and solid-state NMR research in the EMSL. Progress during the last year in the development of the gigahertz NMR spec-

trometer is described in Section 1 of this chapter by Dr. Thomas S. Tenforde, the Acting Manager of the Macromolecular Structure and Dynamics Program. Another new staff member in the NMR Program is Dr. Michael A. Kennedy, who was awarded a distinguished Alexander Hollaender postdoctoral fellowship by DOE in 1991 for NMR studies on DNA structure. His NMR research during the past year on intrinsic DNA bending modes is summarized in Section 2 of this chapter.

The versatility of MS as a tool for studies on macromolecular structure has been greatly increased during the past decade as a result of the introduction of electrospray ionization (ESI). This technique converts molecules to multiply charged ions with mass-to-charge ratios that are within the range that can be analyzed by MS. The use of an ESI interface has extended by two to three orders of magnitude the size range of molecules that can be studied by MS techniques. In addition, chromatographic and electrophoretic separation techniques have been successfully interfaced with MS instruments for analysis of chemical species in complex mixtures. Another major advance has been the introduction of photodissociation and collisional dissociation techniques in conjunction with MS, thereby enabling sequencing to be done on large biomolecular structures. Mass spectrometry is now widely applicable to the analysis of biomolecules with masses into the megadalton range, and is useful for quantifying trace components in complex mixtures and for studies on the interaction of biomolecules with extremely small quantities of toxic substances.

This year's annual report describes advances in several aspects of the MS program headed by Dr. Richard D. Smith. Recent developments include the design and initial work in fabricating a 7-tesla ion cyclotron resonance MS instrument that will be used for extremely high-resolution measurements of mass-to-charge ratios and for the sequencing of biopolymers. Other research reported in Section 3 of this chapter includes MS studies on the gas-phase dynamics of large molecules, ion-molecule reactions, and ion-ion interaction studies in a flow reactor system. Also discussed are the results of MS studies on noncovalent binding interactions of biomolecules.

1. Progress in the Development of a Gigahertz NMR Spectrometer

T. S. Tenforde

The development of an NMR spectrometer operating at a proton frequency of 1,000 MHz (gigahertz) has been motivated primarily by the need for increased resolution and sensitivity in structural studies on large macromolecules in solution. The present generation of NMR instruments, which operate at frequencies up to 600 MHz, are capable of characterizing the 3-D structures of biomolecules with molecular weights approaching 20 kilodaltons. When used in combination with specific isotope labeling for heteronuclear measurements, a gigahertz spectrometer is projected to double the size range of proteins and nucleic acids for which solution structures can be characterized by NMR.

Development of the gigahertz NMR spectrometer is being approached in five subtasks.

1. Measurement of properties such as critical current density and voltage-current characteristics of several candidate superconducting materials for the 23.5 tesla magnet, fabrication and testing of a prototype magnet, and selection of the optimum superconducting materials for the full-scale magnet system.
2. Design and fabrication of the full-scale spectrometer magnet.
3. Development of the radiofrequency system and operating console.
4. Selection and acquisition of the data analysis computer and workstations.
5. System integration and performance tests.

The total time required for completion of this project is expected to be 4 to 5 years.

During the past year, initial activities on Subtask 1 were undertaken. Based on responses to a request for proposals (RFP) issued by PNL, two contractors were selected following peer reviews of their proposals and site visits by PNL staff and outside consultants. Final contract negotiations are expected to be completed by

early 1992, with work on Subtask 1 being initiated at that time. An RFP for Subtask 2 will be issued in 1992 with the goal of selecting a contractor for design and fabrication of the 23.5 tesla magnet by the end of the year. Work on Subtask 3 is being initiated in 1992 through a subcontract with Dr. Gary Drobný.

In Subtask 1, the research on superconducting materials will focus on defining the critical current density (J_c) and voltage-current relationships for long lengths (approximately 1 km) of candidate superconducting materials selected on the basis of preliminary data obtained in tests with short segments (about 1 m in length). Table 1.1 summarizes the advantages and disadvantages of four classes of candidate materials that have been considered for the high-field coils of the gigahertz NMR spectrometer. Of these materials, only the first type of superconductor, composed of either tantalum-doped or titanium-doped niobium tin, is available commercially in sufficiently long lengths to wind coils for the gigahertz NMR spectrometer. The contractors conducting the tests of superconducting materials in Subtask 1 will therefore focus on measuring the physical properties of these ternary niobium-tin compounds in long lengths. For these measurements the conductors will be wound as prototype solenoidal coils and tested in a background magnetic field exceeding 20 tesla.

As shown in Figure 1.1, J_c in short lengths of tantalum-doped niobium tin conductor exceeds 5,000 A/cm² at a field level of 23.5 tesla and an operating temperature of 1.8 K (data provided by Dr. John E. C. Williams of the Massachusetts Institute of Technology Francis Bitter National Magnet Laboratory). This value of J_c is the minimum required for construction of superconducting coils that can reach the target field level of 23.5 tesla. A primary goal in Subtask 2 will be to test the properties of long lengths of this material under the severe transverse and axial stresses that result from magnetostrictive forces in a high magnetic field environment. Tantalum-doped and titanium-doped niobium tin obtained from four major manufacturers will be characterized in these pilot studies. The information gained from these tests will be essential for designing the magnet system of the gigahertz NMR spectrometer to achieve the desired field level of 23.5 tesla with a uniformity of one part in 10⁹ under conditions where the superconducting coils are subjected to forces as high as 250 megapascals.

Table 1.1. Properties of superconducting materials at field levels exceeding 20 tesla.

Material	Advantages	Disadvantages
(Ni,Ta) ₃ Sn or (Ni,Ti) ₃ Sn	<ul style="list-style-type: none"> • Multifilamentary wire and tape form available in long lengths with uniform properties • Moderate reaction temperature required (~700°C) • Production processes and costs are established • Superconducting joints have been fabricated 	<ul style="list-style-type: none"> • J_c falls rapidly above 22 tesla and is sensitive to strain at high field levels
Nb ₃ Al or Nb ₃ (Al,Ge)	<ul style="list-style-type: none"> • Pilot production tests with tape conductors indicate that J_c ~ 2 x 10⁴ A/cm² at 23 tesla and 4.2 K • Good strain tolerance • Moderate reaction temperature required (~700°C) 	<ul style="list-style-type: none"> • High J_c demonstrated only for modest lengths (50-100 m) of tape conductor • Cost of conductor in >1000-m length is unknown
Chevreil phase materials PbMo ₆ S ₈	<ul style="list-style-type: none"> • Both tape and wire conductors are under development • Good strain tolerance 	<ul style="list-style-type: none"> • High reaction temperature required (>1050°C) • Processing technology for lengths >1 m with uniform properties has not been demonstrated • J_c is very anisotropic • Cost of conductor in lengths >1000 m is unknown
Bi2212 or Bi2223	<ul style="list-style-type: none"> • High J_c at fields >22 tesla 	<ul style="list-style-type: none"> • Relatively high reaction temperature required (>800°C) • J_c is extremely anisotropic • Weak links exist at grain boundaries • Process does not yet exist for producing lengths >1 m with uniform properties • Cost of conductor in lengths >1000 m is unknown

Figure 1.1. Critical current density versus magnetic field for tantalum-doped (4 at.%) niobium tin conductor. A significant increase in critical current density at field levels exceeding 18 tesla is achieved by operation at 1.8 versus 4.2 K. The tantalum-doped niobium tin material is an excellent candidate for the inner set of conductors in the gigahertz NMR spectrometer's magnet, provided that a 1.8 K He-II operating temperature is maintained with a Joule-Thomson refrigerator incorporated into the cryostat.

2. NMR Analysis of DNA Intrinsic Bending Modes

M. A. Kennedy and B. R. Reid^(a)

NMR spectroscopy has become a major technique for analysis of the structure and dynamics of DNA and ribonucleic acid (RNA) in solution.⁽¹⁻³⁾ Oligonucleotides with molecular weights up to 10 kilodaltons (about 14 base pairs) are now routinely analyzed by NMR. Refinement of the 3-D structure of a biopolymer predicted from NMR data is commonly carried out by the "distance geometry" procedure.⁽⁴⁾ This algorithm generates a structure completely within

(a) M. A. Kennedy is a NORCUS postdoctoral fellow who holds an Alexander Hollaender fellowship awarded in 1991 by the Department of Energy. B. R. Reid is a Professor of Chemistry and Biochemistry at the University of Washington in Seattle, Washington.

distance space using proton-proton distances determined primarily from standard Nuclear Overhauser Enhancement (NOE) NMR experiments. For those regions of the biomolecule that do not contain protons with which to constrain the predicted structure, coupling constants determined from 2-D NMR experiments can be used to define a range of satisfactory torsion angles that can be converted into distance information. Although data acquisition has become routine, analysis of the NMR data and structure refinement procedures are still in a continuous state of evolution. For example, as larger oligonucleotides containing more than 14 base pairs are studied by NMR (which becomes feasible with increasingly larger field levels), new problems must be addressed such as accounting for anisotropic diffusion effects in the NOE buildup rates.⁽⁵⁾ At present, local structure determination by NMR techniques appears to be reliable, but global structure determination is less accurate and is a sensitive function of the refinement procedure used. Recently, progress has been made in analyzing the global structures of large DNA oligomers composed of as many as 16 base pairs. This report describes NMR research carried out during the past year on intrinsic DNA curvature in the A-T rich regions of DNA oligomers.

Intrinsic DNA Bending

There is considerable evidence that DNA is intrinsically bent in certain A-T rich regions. The first recognition of DNA bending was based on evidence from polyacrylamide gel electrophoresis studies, which showed that the migration of A-T rich DNA fragments is significantly retarded.⁽⁶⁻⁸⁾ The retarded migration pattern was suggested to be the result of coherently phased microbending over many turns of the DNA helix. In support of this idea, it was observed that retarded migration occurs only with DNA fragments that are properly phased [i.e., ligated at a multiple of the number of base pairs constituting a single turn of the DNA helix (approximately 10 base pairs)]. Several investigators have since proposed theoretical models to explain intrinsic DNA bending,⁽⁹⁻¹¹⁾ and several NMR investigations have explored the origin of this phenomenon.⁽¹²⁻¹⁷⁾

NMR spectroscopy has been used to demonstrate that in a run of adenines [e.g., d(A₃T₃) in DNA oligomers], the distance from the adenine H2 proton at position (n) to the deoxyribose H1' proton at position (m + 1) is gradually shortened from more than 4.5 Å in conventional B-type DNA to about 3.8 Å in what is termed B'-type DNA. This proton-proton distance provides an indirect measure of the DNA minor groove width. The B'-type DNA is therefore a region of DNA in which the minor groove width is compressed relative to ideal B-type DNA. A consequence of the narrowing of the minor groove is the introduction of tilt in the base pairs that are normally perpendicular to the helix axis, and a concomitant intrinsic bending of the DNA molecule.

Analogous DNA oligomers in which the order of the purines and pyrimidines is reversed [e.g., d(T₃A₃)], when ligated in the phased manner described above, show no signs of retardation in gel electrophoresis studies.⁽¹⁸⁾ Two possible explanations are the following: 1) the polymers containing d(T₃A₃) are linear with no B' regions or local bending, or 2) there are regions of bending or B' DNA, but the combination of helical twist and local bending fails to produce coherent bending over a larger region of the DNA molecule. Reid and coworkers have recently provided evidence in support of the latter explanation.

Sequence Context Effects in DNA Bending

It has also been observed that the width of the minor groove at the B to B' transition depends on the identity of the flanking base pairs that lead in to the run of adenine residues.⁽¹²⁾ Specifically, the minor groove width compression begins at 3.8 Å following a purine-purine lead in, compared to 4.5 Å or greater following a pyrimidine-pyrimidine lead in. These results are based on studies of oligomers with 12 or fewer base pairs. In order to investigate the role of sequence context effects in intrinsic DNA bending, we have initiated a study in which the flanking region leading in to an adenine run is varied (e.g., purine-purine, purine-pyrimidine, pyrimidine-purine, and pyrimidine-pyrimidine). Structural information at the terminal one or two base pairs of DNA duplexes is usually disregarded due to uncertainty in the distance

measurements resulting from motional dynamics that differ from the motion in the relatively stable interior of the DNA duplex. Therefore, in order to investigate a DNA duplex structure that contains either d(A₃T₃) or d(T₃A₃), including at least four flanking base pairs on both sides, we have designed hexadecamers that will allow the sequence context region of interest to be well within the stable region of the duplex. Specifically, we have synthesized and conducted NMR studies on two hexadecamers, d(CGAGGTTTAAACCTCG)₂ and d(CGTCCAAATTTGGACG)₂.

Figure 2.1 shows the region of an NOE spectrum for d(CGAGGTTTAAACCTCG)₂ that contains cross-peaks with information on the coupling of aromatic base protons to deoxyribose H1' protons. An analysis of this spectrum leads to the following observations. The A9 adenine H2 proton to T26 deoxyribose H1' proton cross-strand cross-peak intensities are not detectable, which is indicative of B-type DNA at the d(TpA) step in the center of the molecule. The second adenine, A10, has observable but weak cross-strand cross-peaks, and the last adenine, A11, has strong cross-strand cross-peaks and is thus indicative of B'-type DNA. The adenine-H2 proton relaxation times shown in Figure 2.1 are consistent with conclusions based on NOE spectral data. These observations lend support to the idea that DNA duplexes containing oligo-d(T_nA_n) runs do indeed contain regions of B'-type DNA, and are not simply linear molecules. The second sequence that we have studied, d(CGTCCAAATTTGGACG)₂, contains a CC lead in to the run of adenines. The cross-strand NOE cross-peaks for the first adenine in the run are, again, not observable, which is indicative of normal B-type DNA (data not shown). The second and third adenines in the run show progressively stronger cross-strand NOE cross-peaks. These results are indicative of compression of the minor groove.

In summary, we have confirmed the presence of a B to B' transition in a DNA duplex containing a d(T₃A₃) run that includes a d(TTpA) step, and in a duplex containing a d(A₃T₃) run that includes a d(CCpA) step. For these two dipyrimidine flanking sequences, the B to B' junction begins with a minor groove width of greater than 4.5 Å. The detailed structural features of the B to B' transition region are currently under investigation.

Figure 2.1. Contour plot of 500-MHz NOE spectral region showing coupling of aromatic base protons H2, H6, and H8 to deoxyribose H1' protons in the hexadecamer, $d(CGAGGTTTAAACCTCG)_2$. The NOE mixing time was 150 msec. Complete sequential assignments of proton-proton dipole couplings have been made, as indicated by enumeration of the cross-peaks 1 to 16 in the NOE spectrum. The boxed areas indicate regions where (n)-adenine-H2 to (m+1)-deoxyribose-H1' cross-peaks are expected to occur in B-type DNA. The one-dimensional aromatic base proton NMR spectrum is shown at the top of the NOE contour plot, and the adenine H-2 proton T1 relaxation times are shown as an inset. The narrow regions in the hexadecamer cross-strand separation represent segments of B'-type DNA.

References

1. J. M. Van de Ven, and C. W. Hilbers, "Nucleic Acids and Nuclear Magnetic Resonance," *Eur. J. Biochem.* **178**, 1-38 (1988).
2. G. M. Clore, and A. M. Gronenborn, "Determination of Three-Dimensional Structures of Proteins and Nucleic Acids in Solution by Nuclear Magnetic Resonance Spectroscopy," *Clin. Rev. Biochem. and Mol. Biol.* **24**, 479-564 (1989).
3. W. J. Metzler, C. Wang, D. B. Kitchen, R. M. Levy, and A. Pardi, "Determining Local Conformational Variations in DNA," *J. Mol. Biol.* **214**, 711-736 (1990).
4. G. Crippen, *Distance Geometry and Conformational Calculations*, John Wiley & Sons, Inc., New York (1980).
5. A. Wang, S. Kim, P. F. Flynn, and B. R. Reid, "Orientation-Dependent Analysis of NOE Intensities: Application to DNA Oligonucleotides," *J. Mag. Res.* (in press).
6. T. Maniatis, A. Jeffrey, and H. van deSande, "Chain Length Determination of Small Double- and Single-Stranded DNA Molecules by Polyacrylamide Gel Electrophoresis," *Biochem.* **14**, 3787-3794 (1975).
7. J. C. Marini, R. Wiesberg, and A. Landy, "The Isolation of Restriction Fragments Containing the Primary and Secondary (gel T) Bacterial ATT Sites of Phage Lambda," *Virology* **83**, 254-270 (1977).
8. N. C. Stellwagen, "Anomalous Electrophoresis of DNA Restriction Fragments on Polyacrylamide Gels," *Biochem.* **22**, 6186-6193 (1983).
9. V. B. Zhurkin, "Sequence-Dependent Bending of DNA and Phasing of Nucleosomes," *J. Biomol. Struct. Dyn.* **2**, 785-804 (1985).
10. L. E. Ulanovsky, and E. N. Trifonov, "Estimation of Wedge Components in Curved DNA," *Nature* **326**, 720-722 (1987).
11. H. S. Koo, H. M. Wu, and D. M. Crothers, "DNA Bending at Adenine-Thymine Tracts," *Nature* **320**, 501-506 (1986).
12. M. Katahira, H. Sugeta, and Y. Kyogoku, "A New Model for the Bending of DNAs Containing the Oligo(dA) Tracts Based on NMR Observations," *Nucl. Acids Res.* **18**, 613-618 (1990).
13. M. Katahira, H. Sugeta, Y. Kyogoku, S. Fujii, R. Fujisawa, and K. Tomita, "One- and Two-Dimensional NMR Studies on the Conformation of DNA Containing the Oligo(dA)oligo(dT) Tract," *Nucl. Acids Res.* **16**, 8619-8632 (1988).
14. A. Kintanar, R. E. Klevit, and B. R. Reid, "Two-Dimensional NMR Investigation of a Bent DNA Fragment: Assignment of the Proton Resonances and Preliminary Structure Analysis," *Nucl. Acids Res.* **15**, 5845-5862 (1987).
15. M. H. Sarma, G. Gupta, and R. H. Sarma, "Structure of a Bent DNA: Two-Dimensional NMR Studies on d(GAAAATTTTC)₂," *Biochem.* **27**, 3423-3432 (1988).
16. V. P. Chuprina, O. Yu. Federoff, and B. R. Reid, "New Insights Into the Structure of A_n Tracts and B'-B' Bends in DNA," *Biochem.* **30**, 561-568 (1991).
17. B. Celda, H. Widmer, W. Leupin, W. J. Chazin, W. A. Denny, and K. Wuthrich, "Conformational Studies of d(AAAAATTTTT)₂ Using Constraints from Nuclear Overhauser Effects and from Quantitative Analysis of the Cross-Peak Fine Structures in Two-Dimensional ¹H Nuclear Magnetic Resonance Spectra," *Biochem.* **28**, 1462-1471 (1989).
18. P. J. Hagerman, "Sequence-Directed Curvature of DNA," *Nature* **321**, 449-450 (1986).

3. Large Molecule Mass Spectrometry

R. D. Smith, J. A. Loo, A. L. Rockwood, H. R. Udseth, R. R. Ogorzalek-Loo, and B. E. Winger

Current research activities encompass three areas involving 1) the development of methods and techniques for the characterization of large biomolecules relevant to bioremediation and health effects research; 2) the development of advanced mass spectrometric instrumentation providing ultrahigh sensitivity, extended molecular weight range, and enhanced structural information; and 3) the chemical physics of large molecule energy transfer, solvation, and reaction processes. Although these three research activities appear superficially disparate, they are highly interactive and mutually beneficial. For example, mass spectrometric analysis of large biomolecules requires that the mechanism for transfer to the gas phase, desolvation, activation, and ion-molecule chemistry relevant to these species be understood at a level sufficient to guide the development of advanced instrumentation.⁽¹⁻⁴⁾ The sensitivity and the ability to structurally characterize large molecules is ultimately limited by the capabilities of advanced instrumentation as well as the fundamental processes noted above. During 1991, substantial progress was made in each of these areas, which has helped elucidate details of planned EMSL instrumentation, a proposed ultra-trace bioanalytical facility, and highlighted the need for a more fundamental understanding of large molecule processes in the gas phase.

Physical Chemistry of Large Molecule Processes

The ability to transfer large molecules to the gas phase, thereby extending to the megadalton regime, based on ESI provides an opportunity to study large molecule processes in the absence of solvent effects and other perturbations encountered in the condensed phases.⁽¹⁻³⁾ In particular, it is now possible to study biopolymers in the gas phase at essentially any extent of solvation, or to use selective ion-molecule processes to probe molecular structure.⁽³⁾ The fundamental studies of isolated molecules in the gas phase can increase our understanding of large molecule reactivity and interactions with smaller chemically important species, providing a basis for evaluating the nature and function of large molecule processes in, for example, enzymatic reactions that might be useful for bioremediation.

Ion-molecule reactions involving multiply charged species produced by ESI appear to have potential for exhibiting structurally specific reactivity. As molecular ion charge state increases, one expects an increased proclivity for proton transfer (and other reaction processes). The most likely sites to be protonated in solution and their relative basicities can differ substantially from rankings based upon the relative proton affinities of amino acid residues in the gas phase. In the gas phase, one expects sites of high proton affinity to be protonated first (i.e., the preferred sites for lower charge sites).⁽⁵⁾ A factor of considerable importance to ESI-MS is the role of increased Coulombic repulsion between charge sites as net charge increases. For example, Coulombic effects can lower "effective proton affinities" of the residues and, depending upon ion structure, will influence the probability of protonation at a specific site. Indeed, estimates incorporating Coulombic effects suggest that protonation of adjacent residues will dramatically lower barriers for the processes of removing protons from the ion or redistributing them within the ion.

In the case of melittin (a small polypeptide of 26 amino acid residues), there are five highly basic residues in addition to the amino terminus. We have been able to deduce the likely locations of charge sites in the gas phase. In solution, the high dielectric constant of the solvent allows adjacent residues to be protonated, and only very small Coulombic effects might be anticipated. However, as solvent is removed, Coulombic effects will increase. In the case of melittin, which has four adjacent highly basic residues, it is probable that protons migrate to other sites, perhaps by a solvent mediated process prior to complete desolvation. An early collision-induced dissociation (CID) study⁽⁶⁾ for melittin indicated that four protonated sites can exist in the y_{13} fragment that encompasses four adjacent highly basic arginine and lysine residues. However, fragmentation in the highly basic region is not observed in spite of the very strong Coulombic contribution expected if all charges reside on the four highly basic residues.⁽⁶⁾ Apparently protons have shifted to other residues, probably tryptophan and glutamine, that also exist on the y_{13} fragment. Intramolecular proton transfer to tryptophan and glutamine residues would greatly increase ion stability by reducing the Coulombic energy. In general, higher charge states have a greater tendency for proton transfer with reagents such as 1,6-diaminohexane. Intermediate charge states are more amenable to clustering with

such reagents, while lower charge states are relatively unreactive.⁽³⁾ These observations can be ascribed to the exothermicity, near thermoneutrality, and endothermicity of the respective proton transfer reactions. As charge increases, proton transfer becomes favored over clustering.

Reactions for highly protonated proteins that would be highly endothermic for the singly protonated species can occur. Using a flow reactor, we have observed that the higher charge states of cytochrome c ($M_r = 12,360$) undergo slow proton transfer reactions with water (170 kcal/mol) at near atmospheric pressure and temperatures greater than 100°C.⁽⁷⁾ The maximum charge state shifted from 19⁺ to 13⁺ after reaction. It appears the resulting charge state distribution may be substantially dependent upon the manner in which evaporation/desolvation is conducted.

Gas-phase proton transfer reactions between multiply charged protein ions and water, which would be expected to be highly endothermic based on proton affinity values for the individual amino acid residues, proceed due to Coulombic destabilization phenomena. The reactivity of the various amino acid residues in a gas-phase multiply protonated protein ion differ from the independent amino acids due to the contribution of Coulombic repulsion. Each proton transfer reaction, therefore, serves to lower the reactivity (or increase the "short range" or intrinsic proton affinity) of the remaining protonated sites.⁽⁵⁾

Ion-ion reactions are another class of interactions that can be studied by ESI-MS.⁽⁸⁾ A straightforward approach to these studies incorporates a capillary inlet fabricated in the shape of a "Y." This approach enables simultaneous operation of two ion sources in an arrangement merging the streams of oppositely charged ions at near atmospheric pressure. Reactions employing several combinations of ion sources have been explored, including +discharge/-ESI, -discharge/-ESI, and +ESI/-ESI modes of operation.

When multiply protonated proteins, $(M+nH)^{n+}$, react with negatively charged species generated by a discharge, the maxima in the cation charge state distributions shift to lower charge, presumably because of proton transfer. The ion intensity maximum in the ESI mass spectrum of horse heart myoglobin shifted from $(M+19H)^{19+}$ to $(M+14H)^{14+}$ after reaction. In horse heart cytochrome c the maximum shifted from the 15⁺ to an 11⁺ charge state. In these ion-ion

reactions, greater attenuation of the higher charge states is commonly observed. Studies employing dual ESI sources to react $(M+nH)^{n+}$ cations with $(M'-mH)^{m-}$ anions produced by ESI have displayed qualitatively similar behavior, in particular, shifting the charge distributions to lower charge states.

Strong evidence of a single step charge inversion process from ion-ion reactions has also been obtained in dual ESI source experiments.⁽⁸⁾ Protonated adenosine 5'-monophosphate (AMP) and fluorescein molecules, $(M+H)^+$, were observed as products of the reactions of highly charged macroions $(M'+nH)^{n+}$ with species generated in the negative electrospray ionization of either AMP or fluorescein. The $(M+H)^+$ products most likely arise from charge inversion of $(M-H)^-$ anions. Protonated fluorescein was observed as a product of reactions with multiply protonated myoglobin and cytochrome c, and to a smaller extent with melittin, which is not as highly protonated. Protonated fluorescein was not observed when the reaction was attempted with serine or bradykinin ions, which are even less protonated than melittin, or with positively charged species of lower proton affinity from an air discharge. Figure 3.1 shows positive-ion mass spectra obtained while electrospraying myoglobin in the positive-ion mode, without a negative electrospray (top) and with ESI of fluorescein negative ions (bottom). Production of protonated AMP molecules was also only observed for reactions with highly protonated reactants.

In addition to possible analytical applications, these techniques hold promise for future investigations of proton transfer and reactivity with gas-phase macroions. Such reaction processes would be advantageous when gas densities must be very low, and may even contribute to reducing the limitations imposed by space charge effects. Of particular interest will be future studies of gas-phase ion-ion reactions at low pressure, especially using ion trapping instrumentation.

Advanced Mass Spectrometric Instrumentation

The method of ion cyclotron resonance (ICR) mass spectrometry has broad potential for the ultrasensitive characterization of large molecules. Ion cyclotron resonance combines nondestructive ion detection and the capability of very high resolution. The ICR method traps ions for an extended time period, and

Figure 3.1. (Top) Electrospray ionization mass spectrum for myoglobin obtained in the positive-ion mode of operation. The higher charge states of myoglobin are evident above m/z 600. (Bottom) Mass spectrum obtained after merging a gas flow containing fluorescein negative ions (produced by -ESI) with the +ESI myoglobin system shown in the top spectrum. The major differences in the spectrum are attributed to partial attenuation of the higher charge states of myoglobin and the appearance of positively charged fluorescein (at m/z 333) due to a single-step charge inversion process.

allows numerous sequential m/z measurements to be conducted, a capability often referred to as MS^n . The ICR instrument consists of an ion storage cell in a uniform magnetic field. Ions formed in the cell, or introduced from an external ion source, can be trapped for extended time periods due to the combination of magnetic and electric fields. The primary motion of a charged particle in a uniform magnetic field involves a circular path perpendicular to the magnetic field strength. In a 7-tesla magnetic field, for example, ions of m/z 750 (which we anticipate to use for DNA molecular ions) would have a cyclotron frequency of about 143 kHz. In principle, the length of time ions can be trapped, the resolution (and measurement accuracy), and the sensitivity all improve as pressure is decreased and the magnetic field is increased. Important elements of the MSRC activities include advancement of external ion sources and injection methods and improved methods for fast acquisition and handling of high resolution Fourier transform ion cyclotron resonance (FTICR) data.

A new 7-tesla ICR instrument designed to obtain ultrahigh vacuum (10^{-10} to 10^{-11} torr) while operating with an external ESI source has been developed at PNL. The ICR incorporates a RF quadrupole beam deflection region and integral cryopumping to obtain the desired UHV. The quadrupole ion beam deflector prevents the intense neutral molecular beam generated from the ESI source from being propagated towards the ultrahigh vacuum region and the ICR cell. A novel feature of this instrument is the incorporation of two high-speed shutters that open only during ion injection. This design greatly enhances differential pumping and allows the desired ultrahigh vacuum to be obtained. An integral cryopump, similar to that designed for one of our quadrupole mass spectrometers, provides high effective pumping speeds ($>100,000$ L/s) and an extremely low ultimate pressure in the ICR cell region. This device consists of two concentric cylinders that enclose the flight path to the ICR cell, the outer cooled to 77 K and the inner to 15 K. An ultrahigh vacuum turbomolecular pump supplements the cryopump to provide effective pumping for hydrogen and helium gases.

Previously there has been little success using ICR for the analysis of large molecules. Coupling the ESI process with ICR has now clearly demonstrated that the high sensitivity and resolution achievable with ICR can also be realized for large molecular species. In conjunction with Professor F. W. McLafferty of Cornell University, preliminary high resolution and tandem mass spectrometry (i.e., MS^n) has been successfully demonstrated. For the protein bovine ubiquitin, resolutions of 50,000 to 70,000 were readily obtained, clearly separating the isotopic contributions, even when using conventional broadband detection. Figure 3.2 demonstrates the potential to conduct collisionally induced dissociation of one charge state of ubiquitin, $(M+10H)^{10+}$, and then detect its dissociation products. High resolution spectra for three of the dissociation products are shown. The mass measurement accuracy possible with ICR, coupled with the ability to directly determine charge state based upon resolution of the isotopic distribution, allows determination of product charge state and confident assignment of the dissociation products. These results show

that there are no fundamental limitations for large molecule measurements by ICR.

Characterization of Noncovalent Biomolecular Interactions

Mass spectrometric methods have now been demonstrated to be applicable to molecules with molecular weights extending to the megadalton range. Previously, we have demonstrated that collisional dissociation techniques allow information to be obtained on the chemical structure of large molecules, potentially providing information on biomolecule sequence, post translational modifications of proteins, covalent adducts (e.g., biomarkers), and other structural details. An important extension of recent research has been using ESI-MS to probe noncovalent interactions of large molecules.⁽³⁾ The ability to generalize such a capability would allow selective interactions, biomolecule function, sequestration, etc., in solution to be directly probed. In addition, the potential would exist

Figure 3.2. High Resolution ESI-ICR obtained for collisionally-induced dissociation of the $(M+10H)^{10+}$ molecular ion of bovine ubiquitin. Insets show detail for three dissociation products.

to (at least qualitatively) gain information on higher-order, 3-D structures of biomolecules.

Information on higher order protein structure (secondary, tertiary, and quaternary) is of fundamental biological importance. Protein conformation in solution depends in a complex fashion upon the intramolecular forces dictated by amino acid sequence. The native state of a protein is typically folded into a well defined 3-D structure by relatively weak intramolecular forces (e.g., hydrogen bonding). Protein function in biological systems generally involves interactions of specific structural conformations essential for activity. X-ray crystallography, NMR, and various spectroscopic methods have been used as probes of such conformation changes. Mass spectrometry potentially offers greater speed and sensitivity in qualitatively probing higher-order structure.

The charge distribution observed for proteins in the gas phase by ESI-MS is related to the solution-phase structure and chemistry. The details of this relationship are unclear since it is uncertain how ESI and subsequent gas-phase processes, such as proton transfer during desolvation, may alter the initial solution-phase charge distribution. As first demonstrated at PNL, protein charging depends dramatically upon higher-order protein structure.⁽⁹⁾ Reductive cleavage of disulfide bonds may affect the higher-order structure of the protein. The native conformation may have several basic amino acid sites effectively "buried" in a globular structure and not as readily protonated. Repulsive Coulombic forces are also greater for the more compact native structure, and charge sites may be more labile in the gas phase (e.g., facile proton transfer). Disulfide-cleavage reactions allow the protein to "relax" into more extended conformations, allowing greater protonation in solution of such buried sites. This behavior may be carried over into the gas phase, as demonstrated by comparisons of spectra for native disulfide-containing proteins, and their disulfide-reduced forms. In the case of disulfide-containing proteins, it is clear that disulfide-reduction will alter both the solution-phase and the gas-phase structure, and implicitly, the extent of charging. However, a 3-D protein structure is more generally governed not by disulfide bridging, but by much weaker noncovalent interactions. Changes in protein environment, such as changes in pH, temperature, addition of reagents such as guanidine hydrochloride or urea, or other solvent conditions may cause denaturation.

It is also well established that the use of organic solvents, such as those common to ESI methods, can cause protein denaturation. Increasing the organic solvent composition shifts the charge state distribution toward higher charge (lower m/z).⁽¹⁰⁾ The native globular form dominates (producing lower charge states in the mass spectrum) for ubiquitin in a completely aqueous solution. At low methanol concentrations, the charge distribution is distinctly bimodal, probably as a result of a mixture of globular and denatured ubiquitin in solution. Addition of approximately 40% v/v methanol is required to eliminate the bimodal charge state distribution (i.e., maximizing the contribution from the denatured state). Only 20% v/v of other solvents such as acetone, acetonitrile, and isopropanol is needed to produce the same effect. Available results suggest that multiple conformation states can be monitored using the ESI mass spectrum. While the charge state distribution from ESI is at least partially governed by the solution-phase conformation, it is uncertain how liquid-phase structure is related to the relevant gas-phase structure. Even if the charge-state distribution is "locked in" by solution phase structure, the actual gas-phase, 3-D structure may yet be altered due to the heating required to complete desolvation, perhaps assisted by Coulombic forces once the protein is stripped of the mediating effects of the dielectric solvent.

Evidence is increasing that relatively weak noncovalent bonds can influence the gas-phase structure of multiply charged proteins.⁽³⁾ A relatively common observation in polypeptide mass spectra is the appearance of noncovalently bound dimers, particularly for samples prepared at high concentration. Generally, such contributions to ESI spectra are relatively small, particularly since ESI interface conditions are selected to provide sufficient excitation for desolvation. Under such conditions, one would generally expect noncovalently associated species to dissociate. As shown by the myoglobin positive-ion ESI spectra at near neutral pH and the negative ion spectra at higher pH, the presumably noncovalently associated heme is retained for a substantial fraction of the molecular ions. Both the negative-ion spectra of A-chain insulin and the positive-ion spectra of B-chain insulin show abundant dimer-ion peaks under certain conditions. MS/MS studies of the suspected dimer species generally yield the expected monomers, at both higher and lower m/z . Figure 3.3 (top) shows ESI mass spectra for the protein carbonic anhydrase, demonstrating the typical

Figure 3.3. (Top) Electrospray ionization mass spectrum for the protein carbonic anhydrase (MW 29,021) showing the distribution of multiply protonated molecules. (Bottom) Mass spectrum obtained for the collisionally-induced dissociation of the dimer, $(2M+51H)^{51+}$, of carbonic anhydrase at m/z 1140, which yields a distribution of monomer charge states.

distribution of molecular-ion charge states as well as small concentrations of dimers evident above values of m/z of approximately 1,000. Figure 3.3 (bottom) shows the CID spectrum for the 51+ charge state of the carbonic anhydrase dimer species, which produces monomer products of both higher and lower m/z than the parent dimer species. As expected, there is reasonable symmetry of the monomer product charge states about the parent (dimer) species. ESI-MS should prove useful to probe other noncovalent binding interactions important in many biological systems. On the basis of such observations, the assumption that the structure of large biopolymer ions from ESI will necessarily be "extended" in the gas phase are premature. Furthermore, the extent to

which ion formation affects the mass spectrum and the MS^n spectrum is not yet well understood. Investigation of gas-phase structure remains an important subject and will be necessary to realize the analytical potential of mass spectrometry for biochemical applications.

References

1. J. A. Loo, C. G. Edmonds, and R. D. Smith, "Primary Sequence Information from Electrospray Ionization Tandem Mass Spectrometry of Intact Proteins," *Science*, **248**, 201-204 (1990).

2. R. D. Smith, J. A. Loo, C. G. Edmonds, C. J. Barinaga, and H. R. Udseth, "New Developments in Biochemical Mass Spectrometry: Electrospray Ionization," *Anal. Chem.*, **62**, 882-899 (1990).
3. R. D. Smith, J. A. Loo, R. R. Ogorzalek-Loo, M. Busman, and H. R. Udseth, "Principles and Practice of Electrospray Ionization-Mass Spectrometry for Large Polypeptides and Proteins," *Mass Spectrom. Rev.* (submitted).
4. J. A. Loo, C. G. Edmonds, H. R. Udseth, R. R. Ogorzalek-Loo, and R. D. Smith, "Electrospray Ionization Mass Spectrometry and Tandem Mass Spectrometry of Large Biomolecules," *Experimental Mass Spectrometry* (in press).
5. A. L. Rockwood, M. Busman, and R. D. Smith, "Coulombic Effects in the Dissociation of Large Highly Charged Ions," *Int. J. Mass Spectrom. Ion. Proc.*, **111**, 103-129 (1991).
6. C. J. Barinaga, C. G. Edmonds, H. R. Udseth, and R. D. Smith, "Sequence Determination for Large Multiply Charged Peptide Molecular Ions by Electrospray Ionization-Tandem Mass Spectrometry," *Rapid Commun. Mass Spectrom.*, **3**, 160-164 (1989).
7. B. E. Winger, K. J. Light-Wahl, and R. D. Smith, "Gas-Phase Proton Transfer Reactions Involving Multiply Charged Cytochrome c Ions and Water under Thermal Conditions," *J. Amer. Soc. Mass Spectrom.* (submitted).
8. R. R. Ogorzalek-Loo, H. R. Udseth, and R. D. Smith, "Evidence of Charge Inversion in the Reaction of Singly Charged Anions with Multiply Charged Macrolons," *J. Phys. Chem.*, **95**, 6412-6415 (1991).
9. J. A. Loo, C. G. Edmonds, H. R. Udseth, and R. D. Smith, "Effect of Reducing Disulfide Containing Proteins on Electrospray Ionization Mass Spectra," *Anal. Chem.*, **62**, 693-698 (1990).
10. J. A. Loo, R. R. Ogorzalek-Loo, H. R. Udseth, C. G. Edmonds, and R. D. Smith, "Solvent-Induced Conformational Changes of Polypeptides Probed by Electrospray Ionization Mass Spectrometry," *Rapid Commun. Mass Spectrom.*, **5**, 101-105 (1991).

Materials and Interfaces

Materials and Interfaces

The development of a fundamental understanding of interfacial structure and chemistry is critical to a number of environmental issues, including the design of efficient and effective environmental remediation and restoration technologies. This understanding will guide the development of chemical sensor and fuse devices, materials for the catalytic decomposition of chemical waste, specialized separation membranes, containment and barrier materials, and models of subsurface contaminant transport.

Research in the Materials and Interfaces program focuses on 1) developing a microscopic understanding of the interfacial chemistry of environmentally relevant materials, such as alkaline earth oxides, mineral oxides, and clays, by studying connections between well characterized surfaces in idealized environments and real interfaces under complex conditions, and 2) applying existing sensor technologies to environmental contamination problems and designing and characterizing new chemically selective materials that can be used in chemical sensor and fuse devices. These programs are to be spread over four research groups: Chemical Sensors, Interphase Chemistry, Materials Synthesis, and Surface Structure and Chemistry. Research has begun in two areas: 1) the surface structure and chemistry of oxides and minerals, and 2) the design of chemically-selective thin-film polymers for use in chemical sensors. Several new staff have recently arrived, including Dr. Jiri Janata, the Associate Director for Materials and Interfaces. A major goal of the coming year is to recruit and hire additional staff needed to get establish programs in all four of our research groups.

1. Surface Structure and Chemistry of Oxides and Minerals

D. R. Baer, D. L. Blanchard, and
M. H. Engelhard

A microscopic understanding of the interactions between subsurface contaminants and mineral surfaces is a critical component of the formulation of accurate, predictive, macroscopic, field-scale models of subsurface contaminant transport. The development of this understanding, however, hinges upon a detailed knowledge of the atomic and electronic structure of

mineral surfaces, along with the binding sites, structures, and energies associated with adsorption. "Unfortunately, the state of mineral surface chemistry lags significantly behind several aspects of surface science in other disciplines..."⁽¹⁾ Difficulties associated with both specimen properties and preparation (i.e., the lack of cleavage surfaces, small crystal size, and the insulating nature of the materials) as well as the lack of a significant motivation to examine these materials have contributed to the neglect of the area by most of the surface science community. However, environmental concerns have been bringing these materials to the forefront, and some of the critical needs in this area are identified in the report of a multidisciplinary workshop *Environmental Materials and Interfaces* held in the fall of 1990.⁽²⁾ Work over the past year has focused on the structure and chemistry of oxide (MgO , SnO_2) and calcite (CaCO_3) surfaces. In addition, a working group has been established to identify issues important to both environmental and materials scientists and to suggest research projects that will help to resolve those issues.

Oxide Surfaces

One of the difficulties associated with the study of insulator surfaces using charged particle beams, such as with low-energy-electron-diffraction (LEED), is the problem of specimen charging. A study of the $\text{MgO}(001)$ surface using beam-modulated LEED allowed for a quantitative determination of the surface relaxation via LEED intensity analysis.⁽¹⁾ This surface relaxes via a rumpling (i.e., a differential relaxation of the surface anion and cation in the direction perpendicular to the surface) of the surface layer and a contraction of the first interlayer spacing. The results of the LEED intensity analysis were in quantitative agreement with the results of some previous LEED studies and with the results of *ab initio* and tight-binding total energy computations.

The structure of the rutile-structure $\text{SnO}_2(111)$ surface is also currently being examined via LEED in collaboration with a group headed by Professor L. L. Erskine at the University of Texas at Austin. The (111) surface occurs naturally on crystals grown by the vapor phase reaction method and appears to be considerably more stable than the (110) and (100) surfaces

examined previously. A series of LEED I-V data has been collected for the (111) surface and is being analyzed to determine the detailed structure of this surface. When complete, this study will be the first detailed examination of the surface structure of a rutile oxide.

Calcite Surfaces

Calcite is a common soil constituent, a frequent component of alkaline waste sludges, and is found in carbonate-containing groundwater.⁽⁴⁾ Calcite is also known to react with metallic cations,⁽⁴⁻⁷⁾ often found in industrial waste and, thereby, strongly influences the concentration and transport of these metals in the environment. Previous research has shown that the transition metal cations that adsorb onto the calcite surface from solution can be grouped into two different classes: those that readily desorb, and those that stick to the surface. A study comparing the interactions of a metal from each of the two classes (Co, which desorbs, and Mn, which stays at the surface) with the calcite cleavage surface has been undertaken.

X-ray photoelectron spectroscopy (XPS) has been used to examine the calcite surface (both with and without Co and Mn) for changes in metal chemistry when exposed to gaseous and liquid water, and to determine the desorption rates of the metals in pure water. Significant differences in the rates of desorption of the two metals have been observed as shown in Figure 1.1. Atomic force microscopy (AFM) has also been used to characterize the cleavage surfaces and to look at the dissolution of the calcite under water. An AFM image of a cleaved calcite surface under water is shown in Figure 1.2. This surface is very flat compared to the "ideal" single crystal surfaces studied for most metals. The XPS analysis has shown that the calcite surface, after water exposure, is nearly identical to the freshly cleaved surface.

Environmental Materials Working Group

During the past year, several efforts have been initiated by the Environmental Materials Working Group, which includes scientists from PNL's Earth and Environmental Sciences Center, Materials and Chemical Sciences Center, and the MSRC. These areas of exploration include attempts to 1) create a thin film clay layer on a conducting substrate; 2) develop an understanding of mineral phase precipitation from

Figure 1.1. Measurement of metal coverage on calcite [fraction of a monolayer (ML) or a function of time in solution].

Figure 1.2. AFM image of a cleaned calcite surface under water.

solutions on organic substrates, and 3) use IR spectroscopy to study sorption on single crystal mineral surfaces in solution. Explorations into the materials and surface science issues important to understanding the waste stored in tanks at Hanford is planned.

Initial attempts have been made to create a thin-film clay electrode to serve as a model material. Initial efforts include 1) adsorption of a clay suspension on a self-assembled monolayer surface, and 2) laser ablation of clay components onto a substrate subjected to further processing. This effort also includes extended x-ray absorption fine-structure (EXAFS) spectroscopy studies related to precipitation from solution and Fourier transform (FT) IR spectroscopy and electrochemical measurements of adsorption from solution. A set of EXAFS measurements on several standard compounds, two initial films, and several solution precursors have been taken and analysis will continue.

Finally, several collaborative research projects were undertaken during the past year in the effort to develop a user community for the MSRC and, eventually, the EMSL. These efforts include comparison of fractoemission studies with XPS and electron stimulated desorption (ESD) measurements on calcite surfaces in collaboration with Professor J. Tom Dickinson at Washington State University; the investigation of the electronic structure of layered compounds via XPS with Dr. David Cleary, also at Washington State University; and the exploration of thin-film ceramic synthesis processes with Professor David Johnson at the Material Science Institute of the University of Oregon.

References

1. M. F. Hochella, Jr., "Mineral-Water Interface Geochemistry", M. F. Hochella, Jr., and A. F. White, eds., *Reviews in Mineralogy*, **23** (1990).
2. "Environmental Materials and Interfaces," PNL-(CONF-9009424), Pacific Northwest Laboratory, Richland, Washington, 1991.
3. D. L. Blanchard, D. L. Lessor, John P. La Femina, D. R. Baer, W. K. Ford, and T. Guo, *J. Vac. Sci. Tech.* **A9**, 1814 (1991).
4. D. R. Baer, D. L. Blanchard, M. L. Engelhard, and J. M. Zachara, *Surf. and Interf. Analysis*, **17**, 25 (1991).
5. J. M. Zachara, J. A. Kittrick, and J. B. Harsh, *Geochim. Cosmochim. Acta* **52**, 2281 (1988).

6. L. S. Dake, D. R. Baer, and J. M. Zachara, *Surf. and Interf. Analysis* **14**, 71 (1989).
7. J. M. Zachara, C. E. Cowan, and C. T. Resch, *Geochim. Cosmochim. Acta* **55**, 1549 (1991).

2. Design of Selective Layers for Chemical Microsensors

J. Janata

Organic semiconductors can be prepared in many different ways and can yield a great variety of chemically and physically selective layers for gas and liquid sensing. The electrochemical mode of their preparation is particularly suitable for applications in which the selective layer has to be deposited on a small or geometrically complicated area.⁽¹⁾ These layers can be used in all major transduction sensor groups, particularly in mass and in electrochemical sensors.^(2,3) The selectivity of the layers is derived from the specific interactions of the guest molecules with the polymer or from the interactions with specific binding sites incorporated in the matrix. It has been shown that the formation of charge transfer complexes between gases or organic vapors and the semiconducting matrix results in measurable change of work function of the material. Such materials can be incorporated in chemical microsensors based on metal-insulator-semiconductor structures. The theory supporting these experimental observations has been developed.⁽⁴⁾

References

1. M. Josowicz, and J. Janata, "Electroactive Polymers in Chemical Sensors," *Electroactive Polymers*, B. Serosati, ed., Chapman and Hall Publishers (in print).
2. P. Topart, and M. Josowicz, *J. Phys. Chem.* (submitted).
3. J. Langmaier, and J. Janata, *Anal. Chem.*, **64**, 523-527 (1992).
4. J. Janata, *Anal. Chem.*, **63**, 2546-2550 (1991).

Theory, Modeling, and Simulation

Theory, Modeling, and Simulation

Theoretical and computational molecular science will play a critical role in developing molecular-level descriptions of chemical, physical, and biological processes in nature and other complex environments. Theory provides the conceptual framework for the analysis and interpretation of experimental data. Modeling and simulation provide quantitative information that may be expensive, difficult, or even impossible to obtain in the laboratory.

Modeling molecular processes that occur in the environment requires a fundamental understanding of molecules and their interactions in isolation (gas phase) and in liquids, on surfaces, and at interfaces (condensed media). Research in the Theory, Modeling, and Simulation program involves the development of new theoretical methods and modeling and simulation techniques appropriate for the molecular systems found in complex environments as well as the application of state-of-the-art methods to important molecular problems in such systems. The latter program combines efforts devoted to rigorous studies of fundamental molecular processes (diffusion, reaction, solvation, absorption, etc.) in model systems with efforts devoted to the simulation of the complex multi-atom, multispecies, multiphase molecular systems found in the environment.

The Theory, Modeling, and Simulation program contains three research activities: Molecular Theory and Modeling, Materials Theory, and Biomolecular Modeling and Simulation. The Molecular Theory and Modeling activity focuses on understanding elementary molecular processes in both the gas and condensed phases. The Materials Theory and Modeling activity extends theoretical models of surfaces and materials to oxide, carbonate, and clay mineral systems. The Biomolecular Modeling and Simulation activity focuses on the development and application of techniques to model biochemical systems.

The computing demands of research in the Theory, Modeling, and Simulation program are substantial and will grow dramatically over the next few years as new projects involving molecular systems of increasing size and complexity are undertaken, new staff are added to the program, and extensive collaborative projects are

initiated. To provide an optimal response to the computing needs of this effort, the Theory, Modeling, and Simulation program, in collaboration with the Computing and Information Sciences program, will develop and manage the Molecular Science Computing Facility (MSCF) as a collaborative research facility in the EMSL. The computer hardware in the MSCF will provide state-of-the-art high performance computational and visualization capabilities for molecular simulations, while enabling the evaluation and exploitation of promising new computer technologies. To provide the software needed to make maximal use of the hardware in the MSCF, the Theory, Modeling, and Simulation program contains three research and development activities: high performance computing, graphics and visualization; and molecular databases.

1. Molecular Theory and Modeling

B. C. Garrett, L. X. Dang, T. H. Dunning, Jr.,
A. C. Hess, M. I. McCarthy, M. M. Marino,
R. P. McRae, G. K. Schenter, and
S. S. Xantheas; G. Haynes,^(a) R. Poshusta,^(b)
G. C. Schatz,^(c) G. Voth,^(a) W. C. Ermler^(d)

Molecular Theory and Modeling activities are focused on *ab initio* theoretical studies of molecular processes in the gas phase, in liquids, and at gas-solid and liquid-solid interfaces. The research objectives of this program are threefold:

- To apply available *ab initio* techniques to study molecular processes involved in the chemistry of the environment, considering both gas- and condensed-phase processes.
- To extend current *ab initio* techniques to treat molecular systems of future interest and to develop new techniques for addressing problems that are computationally intractable at present.

(a) University of Pennsylvania, Philadelphia, Pennsylvania.

(b) On sabbatical leave; permanent address: Washington State University, Pullman, Washington.

(c) Northwestern University, Evanston, Illinois.

(d) On sabbatical leave; permanent address: Stevens Institute of Technology, Hoboken, New Jersey.

- To aid in the development of more accurate simulation techniques for describing molecular processes occurring in the complex systems found in the environment.

The main emphasis of the research in Molecular Theory and Modeling is on molecular processes in the condensed phase that are important in groundwater and soil chemistry, but current applications also include studies of molecular processes important in atmospheric chemistry.

Molecular Processes in Liquids

Solvation and reaction in aqueous media play key roles in the migration and behavior of contaminants in the subsurface environment. Cluster approaches are being used to address detailed questions about the effect of solvation on molecular structure, properties, and reactivity, with an emphasis on the chemistry of anions. Molecular dynamics simulations using model potentials derived from experimental and/or computational data extend our knowledge of these systems into bulk condensed phases. Molecular dynamics methods are also being evaluated for calculating liquid state properties, such as IR spectra, transport coefficients, and equilibrium properties. Work is also in progress to extend gas-phase theories to study the effects of solvation on chemical reactions.

Modeling Negative Ion-Water Clusters

As noted above, one approach to modeling solvation effects is to simulate the extended solute-solvent system by a finite cluster of molecules. This approach makes full use of the experience gained in modeling conventional molecular systems, but substantial computing power is required to model the larger clusters of interest. On the other hand, clusters of molecules, which represent an aggregated state of matter with properties ranging between the gas and condensed phases, are themselves of intrinsic interest. These species have been identified as key participants in important chemical and physical processes occurring in the earth's atmosphere (e.g., aerosol formation). A more complete understanding of cluster structure and reactivity would not only form the basis for improved models of solute-solvent systems, but would also provide insights into this novel form of matter.

During this past year, we studied the energetics of hydride ion solvation by modeling small water clusters, $\text{H}^-(\text{H}_2\text{O})_n$ with $n = 1-4$. *Ab initio* calculations performed on the $(\text{H}_2\text{O})_2$ and $\text{H}^-(\text{H}_2\text{O})$ clusters indicated that 1) electron correlation is important for the accurate description of these systems, and 2) structural and accurate energetic information can be deduced at the second-order perturbation theory (MP2) level with the new augmented correlation consistent basis sets.⁽¹⁾ For example, the binding energy of the water dimer is computed to be -5.3 kcal/mol, which is in excellent agreement with the experimentally estimated value of -5.4 ± 0.2 kcal/mol.⁽²⁾ The optimized geometry of the dimer at this level is also in good agreement with the experimentally deduced structure (e.g., the calculated $\text{R}_{\text{O-O}}$ is 2.92 Å compared to a measured⁽³⁾ value of 2.976 Å). Therefore, we optimized the geometry of the larger hydride-water clusters at this level of theory.

In contrast to earlier assumptions, we find that the clusters adopt asymmetric structures. The asymmetric structure of the clusters is a result of balancing the strong ion-water interactions with the weaker, but still significant, water-water interactions. The optimal geometries for the $\text{H}^-(\text{H}_2\text{O})_n$ clusters with $n = 1-3$ are shown in Figure 1.1. For $n = 2$ (3) we find that the optimal geometry resembles a cyclic structure in which the two (three) water molecules are connected through hydrogen bond(s). The calculated incremental energy $\Delta E_n = E_n - (E_{n-1} + E_{\text{water}})$ corresponding to successive addition of water molecules is -17.4, -14.8 and -13.1 kcal/mol, respectively. Calculations on $\text{H}^-(\text{H}_2\text{O})_4$ are in progress.

In order to refine the energetics of the larger clusters, we are currently performing more accurate perturbation theory calculations with larger augmented correlation consistent polarized valence basis sets at the previously optimized geometries.

A study of water clusters of the F^- ion has been initiated. For $\text{F}^-(\text{H}_2\text{O})_2$, we find a cyclic structure similar to the one observed for the hydride ion.

The $\text{H}^- + \text{H}_2\text{O} \rightarrow \text{OH}^- + \text{H}_2$ Reaction

As noted above, we have been studying the interaction of the hydride ion, H^- , with water as a simple prototype of anion solvation. However, H^- also rapidly reacts with water forming OH^- and H_2 ; the reaction is

Figure 1.1. Optimal structures for the $\text{H}^+(\text{H}_2\text{O})_n$ clusters ($n=1-3$). Structure 3a corresponds to a local minimum (7.9 kcal/mol above the global minimum, Structure 3b).

exothermic⁽⁴⁾ by 10.1 ± 0.9 kcal/mol. The MP2 calculations coupled with the new basis sets specifically developed for correlated calculations⁽¹⁾ were used to determine the optimal structures and energies of all critical points on the reaction path. The resulting schematic diagram of the ground-state⁽¹⁾ A' potential energy surface is given in Figure 1.2. The calculations predict the $\text{H}^+\cdots\text{H}_2\text{O}$ complex to be bound by 17.0 kcal/mol relative to $\text{H}^+ + \text{H}_2\text{O}$, which is in good agreement with the estimate of 17.3 ± 1.2 kcal/mol from experimental studies.⁽⁵⁾ Note that in Figure 1.2, a second stable complex resembling the structure $\text{OH}^+\cdots\text{H}_2$ exists. This complex lies only 0.6 kcal/mol higher in energy than the $\text{H}^+\cdots\text{H}_2\text{O}$ minimum. The barrier for their interconversion is of the order of 2 kcal/mol.

Figure 1.2. Structures and energies of the stable structures and transition states for the $\text{H}^+(\text{H}_2\text{O})$ complex (experimental estimates are given in parentheses).

The above information provides insights into the isotropic exchange reaction:⁽⁶⁾

$$k = 3 \times 10^{-11} \text{ cm}^3/\text{molec} \cdot \text{sec}$$

The model proposed to explain the rapid rate of this reaction involves 1) formation of a stable $\text{HO}^+\cdots\text{D}_2$ complex, 2) deuteron transfer to form the $\text{HOD}^+\cdots\text{D}^+$ complex, 3) rotation of the HOD to form $\text{DOH}^+\cdots\text{D}^+$, 4) proton transfer to form $\text{DO}^+\cdots\text{HD}$, and 5) dissociation to $\text{DO}^+ + \text{HD}$. Steps 1 and 2 and Steps 4 and 5 are consistent with the results summarized in Figure 1.2. The complexes in Steps 1 and 5 lie below the $\text{OH}^+ + \text{D}_2$ limit, as does the transition state connecting the complexes in Steps 2 and 4. For Step 3 to be rapid, the transition state for the hydrogen atom interchange process must also lie below $\text{OH}^+ + \text{D}_2$. Our calculations place the transition state for this step at 3.7 kcal/mol, over 2 kcal/mol below the dissociation limit and, thus, provide strong support for the proposed mechanism.

A similar investigation of the $\text{H}^+(\text{H}_2\text{O})_2$ potential energy surface is under way.

Development of Accurate Intermolecular Potentials Using Molecular Dynamics: Solvation of Li^+ and F^- Ions in Polarizable Water

It has been recognized for some time that ionic clusters play an important role in many chemical and physical processes, especially groundwater and soil chemistry⁽⁷⁾ but also in atmospheric chemistry and combustion. Consequently, a quantitative description of clusters and solutions has been the focus of many recent theoretical and simulation studies.⁽⁸⁻¹³⁾ Computer simulations by Monte Carlo⁽¹⁴⁾ and molecular dynamics⁽¹⁵⁾ methods have become powerful tools in attempts to understand the properties of these systems.

As part of our ongoing research on the development of accurate intermolecular potentials for water-water and ion-water interactions, we have been investigating the solvation of Li^+ and F^- ions in clusters and in solutions. Sufficient experimental data^(16,17) and *ab initio* calculations^(18,19) exist on the thermodynamics and structural properties of these systems to guide us in developing accurate models for ion-water interactions. The goal of this study is to describe ion-water interactions in both clusters and solution by a set of accurate intermolecular potentials.^(8,9)

In Table 1.1, the calculated and corresponding experimental⁽⁸⁾ enthalpies of the clusters are presented. In all cases, we obtained good agreement with the experimental results. Correct prediction of successive H_2O binding energies for the ion-water complexes for $n \geq 3$ is encouraging, since we did not make any further adjustment of the ion-water potential parameters for these cases. Snapshots of water molecules around the Li^+ and F^- ions are shown in Figure 1.3. The results indicate that the water molecules are distributed nearly symmetrically around the Li^+ ion, while for the F^- ion the water molecules are clustered in the same hemisphere. For $n=4$, the water molecules form a distorted tetrahedron structure around the Li^+ ion.

With the potentials developed in the above cluster studies, we carried out molecular dynamics simulations of aqueous solutions of Li^+ and F^- . The calculated and the experimental⁽²⁰⁾ enthalpies are tabulated in Table 1.2. The calculations reproduce the observed solvation enthalpies very well.

Table 1.1. Calculated and experimental enthalpies of binding (kcal/mol) of ion- $(\text{H}_2\text{O})_n$ clusters at 300 K.

Ion	n	Molecular Dynamics	Exptl ^(a)
Li^+	1	-33.8 ± 1.3	-34.0
	2	-61.5 ± 1.7	-59.8
	3	-80.9 ± 2.7	-80.5
	4	-97.9 ± 2.3	-96.9
	5	-108.1 ± 2.8	-110.8
	6	-120.2 ± 2.7	-122.9
F^-	1	-23.5 ± 1.1	-23.3
	2	-42.1 ± 1.8	-39.9
	3	-56.2 ± 2.8	-53.6
	4	-69.6 ± 2.2	-67.1
	5	-80.8 ± 2.4	-80.3
	6	-91.8 ± 3.4	--

(a) Reference 16.

In summary, accurate, nonadditive intermolecular potentials for the Li^+ -water and F^- -water interactions in the gas phase and the liquid phase were developed. Future applications of these models will include the solvation of ions at water liquid-vapor interfaces, free energy of solvation, interionic potentials of mean force, and related thermodynamics properties of ionic solutions. Work along this line is in progress.

Properties of Liquid Water and Carbon Dioxide from Constant Pressure-Constant Temperature Molecular Dynamics

Although most experiments are controlled by fixing temperature and pressure, most molecular dynamics simulations are controlled by fixing the number of particles, volume, and energy (NVE). Recent advances in simulation methods now allow molecular dynamic simulations for constant pressure and constant temperature (NPT).^(21,22) It has been shown that these methods properly reproduce equilibrium ensemble averages,⁽²²⁾ but their utility for calculating ensemble averages of time-dependent quantities, such as velocity autocorrelation functions for complex fluids, remains a question. We have recently performed systematic comparisons of the microcanonical (NVE) and

(a) $\text{F}^- (\text{H}_2\text{O})_n$

(b) $\text{Li}^+ (\text{H}_2\text{O})_n$

Figure 1.3. Snapshots of water around the ions taken from molecular dynamics simulations with $n = 1, 2, 3$, and 4.

isothermal-isobaric (NPT) ensembles for molecular dynamic simulations of two prototypical molecular fluids: liquid water and liquid carbon dioxide.

Table 1.2. Averaged solvation enthalpies (kcal/mol) for the Li^+ and F^- ions at 300 K.

Ion	Molecular Dynamics	Exptl ^(a)
Li^+	-128 (-107.5) ^(b)	-130
F^-	-114 (-93.5) ^(b)	-116

(a) Reference 20.

(b) Value before Born correction.

The NPT-molecular dynamics simulations were done using the Andersen-Nosé algorithm in which the dynamics are computed for an extended Hamiltonian that has explicit dependence on all the real system coordinates, two new variables (a velocity scaling factor and the volume), and their conjugate momenta. The dynamics of this effective Hamiltonian system constrain the average temperature and pressure of the real variables to externally controlled values. Masses for the two scaling degrees-of-freedom enter as parameters. Although the equilibrium properties of the system are independent of the values chosen for these masses, the dynamics may not be insensitive. Values of the mass parameters were chosen so that the time scales for fluctuations in the velocity scaling factor and the volume were short compared to the total simulation time, thus ensuring sufficient averaging, but long compared to the molecular time scales of the liquid.

Simulations were performed for both rigid and flexible models of water and carbon dioxide. The potentials were taken from the literature.⁽²³⁾ In the NPT-molecular dynamic simulations, the external temperature was set to 298 K for all systems, and the external pressure was set to 1 atm for water and 400 atm for carbon dioxide.

Equilibrium properties, such as pair distribution functions, were virtually independent of the ensemble in which they were calculated. Some time-dependent quantities also showed remarkable insensitivity to the ensemble. Figure 1.4 shows the velocity autocorrelation function for flexible water and Figure 1.5 shows the power spectrum (fourier transform of the dipole autocorrelation function) for bulk water. The results for the two ensembles are nearly identical. Diffusion coefficients for the two ensembles are compared in

Figure 1.4. Velocity autocorrelation functions for the hydrogens on flexible water are plotted for both the NPT- and NVE-molecular dynamics simulations.

Figure 1.5. Power spectrum for flexible water obtained by FT of the dipole correlation function are plotted for both the NPT- and NVE-molecular dynamic simulations.

Table 1.3. Again, the results from the two ensembles agree quite well. For the NPT ensemble, the time scales for a change in sign of the velocities of the scaling variables are on the order of 0.5 to 2.0 ps. This is much larger than the time scale for fluctuations in the velocity autocorrelation function; therefore, it is not surprising that the results for the two ensembles agree well. Even in cases where the time scale of the dynamical process is not significantly faster than the temperature and pressure fluctuation, such as for the diffusion constant, reliable estimates of the dynamical properties can sometimes be made using the NPT ensemble.

Table 1.3. Computed diffusion coefficients ($\text{cm}^2 \text{s}^{-1}$).

Ensemble	System		
	Rigid Water	Flexible Water	Rigid CO_2
NPT	3.5×10^{-5}	1.3×10^{-5}	1.2×10^{-4}
NVE	3.4×10^{-5}	1.5×10^{-5}	1.2×10^{-4}

Dynamic Solvent Effects on Activated Chemical Reactions

Concepts and methods used to describe reactions occurring in the gas phase are finding increased utility for reactions in solution. There has been great progress over the last decade in developing quantitative methods for predicting rates of gas-phase reactions. In the present work, we extend these methods to activated reactions in solution. In previous work, we have examined the validity of gas-phase methods for treating solution-phase reactions within a classical mechanical framework. The present study investigates methods for incorporating quantum mechanical effects into these methods for application to solution-phase reactions.

The constant proximity of solvent molecules can alter both the interaction energies of reacting solute molecules and the dynamics of the reactions through both localized and collective interactions. Dynamical effects of the solvent molecules can involve localized interactions, such as direct catalytic participation of solvent molecules in the reaction and caging effects. These effects can be taken into consideration within gas-phase theories by explicitly treating some subset of the solvent molecules, such as the first solvation shell. It is more difficult to see how gas-phase theories can be applied to dynamical solvent effects that involve

collective interactions, such as solvent-solute collisions that can be described in terms of frictional forces. These latter effects are the subject of the present work.

The effect of solvent friction and stochastic random forces on activated chemical reaction rates was first treated by Kramers⁽²⁴⁾ using a classical Langevin equation. Grote and Hynes⁽²⁵⁾ generalized this result to include memory effects in the solvent friction to form a Generalized Langevin Equation (GLE) description of the motion. Based on the developments of Zwanzig,⁽²⁶⁾ in which a formal correspondence between a GLE description and an effective harmonic bath Hamiltonian was derived, Pollak⁽²⁷⁾ recently showed that rates obtained from a GLE analysis may be obtained from a direct Transition State Theory (TST) analysis of Zwanzig's effective Hamiltonian. Whereas the Kramers and Grote-Hynes theories are restricted to classical mechanical treatment, quantum effects can be included in TST in the normal *ad hoc* manner.

Techniques for calculating rates of chemical reactions based on variational transition state theory (VTST), including important quantum mechanical effects, have been successfully applied to a variety of gas-phase systems.⁽²⁸⁾ In this work, we extend these techniques to the condensed phase by using the Zwanzig effective Hamiltonian. Namely, the minimum energy path in the solute plus the effective solvent bath system is obtained, and variational dividing surfaces are chosen from planes perpendicular to the minimum energy path. Quantum mechanical effects are included by quantizing bound degrees-of-freedom and including multidimensional corrections to quantized motion along the reaction path motion. The influence of each of these effects may be seen from an analysis of the collinear $\text{H}+\text{H}_2$ reaction in a typical rare gas solvent. The reactive flux $Q_R k(\eta^*)$ is compared to its value at zero coupling in Figure 1.6. The parameter η^* is a dimensionless integrated friction representing the strength of the coupling of the effective solvent bath to the solute. We see that both quantizing bound degrees-of-freedom and including quantum mechanical tunneling effects have profound effects on the computed rates. Whereas a purely classical theory predicts decreases in the reaction rate of factors of five at the highest friction, including quantum mechanical effects can cause increasing friction to depress the reactive flux by several orders-of-magnitude.

Figure 1.6. Dependence of reactive flux on integrated friction for the model reaction $\text{H}+\text{H}_2$ in solution. Classical TST is compared with VTST with quantized bound modes, and with quantized bound modes and a multidimensional semiclassical adiabatic tunneling (SAG) correction.

Although these methods have been tested extensively for gas-phase reactions, the validity of these methods for solution-phase reactions needs to be established. A critical test of these methods for a model of a reaction in solution is presented in Figure 1.7. The results of our analysis for the reactive rate of a single solute degree-of-freedom coupled to a single effective solvent bath degree-of-freedom are compared to a centroid density calculation of Voth and Haynes and an exact quantum dynamics calculation by Schatz, which are unpublished. Although the centroid density method⁽²⁹⁾ is based upon very different approximations than the VTST calculations, both agree well with each other, as well as with the accurate results.

Figure 1.7. Comparison of reactive fluxes computed by VTST/SAG, the centroid density method, and accurate quantum dynamics. Absolute fluxes are computed as a function of integrated solvent friction for the model of an Eckart barrier coupled to a single solvent bath coordinate.

Molecular Processes at Gas-Solid and Liquid-Solid Interfaces

Molecular processes at liquid-solid interfaces are critical in determining the binding of contaminants to soil minerals. A prerequisite for modeling dynamical processes that occur at mineral interfaces is an accurate determination of the adsorbate/surface interaction potentials. *Ab initio* computation of these interaction potentials has elucidated the nature of these weak molecule/surface interactions and provided a means of predicting the behavior of certain classes of species on ionic surfaces.

The interfacial properties of molecular adsorbates on the surface of MgO was investigated using periodic Hartree-Fock methods.⁽³⁰⁾ An extensive set of calculations were done to probe the behavior of Cl₂ on the (001) face of MgO.⁽³¹⁾ The Cl₂ molecules were aligned perpendicular to a slab of MgO, over either a magnesium or oxygen site on the surface. The entire system is periodic in two dimensions with the finite thickness of the slab (three layers) being set such that the interlayer atoms resembled those found in the bulk. The basis set for the solid has an optimized split valence form. The Cl₂/MgO binding energy was calculated as a function of the Cl₂ basis set and surface coverage. [Each coverage calculation required the explicit treatment of a unit cell containing two molecules (i.e., 52 atoms, 760 basis functions for a 1:8 dilution).] The total energy of the adsorbate approaching a Mg site on the surface indicates a purely repulsive interaction. Over the O atom, on the other hand, the molecule is bound at 1:8 dilution with

Figure 1.8. Total energy of interaction between the net of Cl₂ molecules and the MgO (001) surface as a function of distance. Adsorbate is aligned perpendicular to the surface plane over either a Mg (solid line) or O (dashed line) site. Calculations are for two-sided adsorption with a surface coverage of 1:4.

a binding energy of 4.1 kcal/mol (Figure 1.8). The charge density and density deformation maps indicated that Cl_2 binding involves very little charge transfer (only 0.03|e|) between the surface and the molecule. The calculated electrostatic potential and the electric field maps revealed that the interaction is dominated by the electrostatic interaction between the molecule and the surface. A simple electrostatic description of the attractive forces in the binding energy can account for the preference of the adsorbate to bind over oxygen and not over magnesium. The total and projected density of states indicates that the higher energy valence states in the slab are dominated by contributions from the oxygen 2s and 2p orbitals. These states overlap and mix with the states arising from the 3s and 3p orbitals of the Cl_2 net. This mixing is much less between chlorine and magnesium states.

Calculations involving a series of diatomic adsorbates showed similar behavior, indicating that the general features of adsorbate/surface interactions for many systems can be accurately described by electrostatic forces. Work is currently in progress to understand the nature of the interaction of more complex adsorbates (i.e., NH_3 , H_2O , CCl_4) on the MgO (001) surface. These *ab initio* calculations are providing the essential components necessary for future simulations of the dynamical processes occurring at mineral interfaces.

References

1. T. H. Dunning, Jr., *J. Chem. Phys.* **90**, 1007 (1989).
2. J. Reimers, R. Watts, and M. Klein, *Chem. Phys.* **64**, 95 (1982).
3. J. A. Odutola, and T. R. Dyke, *J. Chem. Phys.* **72**, 5062 (1980).
4. D. Betowski, J. D. Payzant, G. I. McKay, and D. K. Bohme, *Chem. Phys. Lett.* **31**, 321 (1975).
5. M. Henchman, and J. F. Paulson, *Radiat. Phys. Chem.* **32**, 417 (1988).
6. M. Meot-Ner, J. R. Loyd, E. P. Hunter, W. A. Agosta, and F. H. Field, *J. Am. Chem. Soc.* **102**, 4672 (1980); J. J. Grabowski, C. H. DePuy, and V. M. Birebaum, *J. Am. Chem. Soc.* **105**, 2565 (1983).
7. "Clusters: A New State of Matter," *LBL Research Review* **16**, 2 (1991).
8. J. Caldwell, L. X. Dang, and P. A. Kollman, *J. Am. Chem. Soc.* **112**, 9145 (1990).
9. L. X. Dang, J. E. Rice, J. Caldwell, and P. A. Kollman, *J. Am. Chem. Soc.* **113**, 2481 (1991).
10. L. Perera, and M. L. Berkowitz, *J. Chem. Phys.* **95**, 7556 (1991).
11. P. Cieplak, T. Lybrand, and P. A. Kollman, *J. Chem. Phys.* **87**, 6393 (1990).
12. G. Corongiu, M. Migliore, and E. Clementi, *J. Chem. Phys.* **89**, 4629 (1989).
13. L. A. Curtiss, J. W. Hautman, and A. Rahman, *J. Chem. Phys.* **86**, 2319 (1987); J. B. Foresman, and C. L. Brooks, *J. Chem. Phys.* **87**, 5892 (1987).
14. N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller, *J. Chem. Phys.* **21**, 1087 (1953).
15. B. J. Alder, and A. Wainwright, *J. Chem. Phys.* **31**, 459 (1959).
16. I. Dzidic, and P. Kebarle, *J. Phys. Chem.* **74**, 1466 (1970); M. Arhadi, R. Yamdagni, and P. Kebarle, *J. Phys. Chem.* **74**, 1475 (1970).
17. R. L. Wooding, and J. L. Beauchamp, *J. Am. Chem. Soc.* **100**, 501 (1978).
18. H. Kistenmacher, H. Popkie, and E. Clementi, *J. Chem. Phys.* **61**, 799 (1974).
19. J. E. Del Bene, M. J. Frisch, K. Raghavachari, J. A. Pople, and P. V. R. Schleyer, *J. Phys. Chem.* **87**, 73 (1983); R. L. Wooding, F. A. Houle, and W. A. Goddard III, *Chem. Phys.* **14**, 461 (1976).
20. H. L. Friedman, and C. V. Krishnan. *Water: A Comprehensive Treatise*, Franks, ed., Vol. 6, p. 1, Plenum Press, New York, 1973.
21. H. C. Andersen, *J. Chem. Phys.* **72**, 2384 (1980).
22. S. Nosé, *J. Chem. Phys.* **81**, 511 (1984).

23. W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, and M. L. Klein, *J. Chem. Phys.* **79**, 926 (1983); D. F. Coker, R. E. Miller, and R. O. Watts, *J. Chem. Phys.* **82**, 3554 (1985); C. S. Murthy, K. Singer, and I. R. McDonald, *Mol. Phys.* **44**, 135 (1981); C. S. Murthy, S. F. O'Shea, and I. R. McDonald, *Mol. Phys.* **50**, 531 (1983); G. C. Schatz, and M. J. Redmon, *Chem. Phys.* **58**, 195 (1981).
24. H. A. Kramers, *Physica* **7**, 284 (1940).
25. R. F. Grote, and J. T. Hynes, *J. Chem. Phys.* **73**, 2715 (1980); **77**, 3736 (1982).
26. R. Zwanzig, *J. Stat. Phys.* **9**, 215 (1973).
27. E. Pollak, *J. Chem. Phys.* **85**, 865 (1986).
28. See, for example, A. D. Isaacson, D. G. Truhlar, S. N. Rai, R. Steckler, G. C. Hancock, B. C. Garrett, and M. J. Redmon, *Computer Phys. Comm.* **47**, 91 (1987) and references therein.
29. G. A. Voth, D. Chandler, and W. H. Miller, *J. Chem. Phys.* **91**, 7749 (1989).
30. C. Pisani, R. Dovesi, and C. Roetti, *Hartree-Fock Ab Initio Treatment of Crystalline Systems*, Springer-Verlag, New York, 1988; R. Dovesi, C. Pisani, C. Roetti, and V. R. Saunders, Program Crystal, Quantum Chemistry Program Exchange, Program #577, Indiana University, Department of Chemistry.
31. M. I. McCarthy, and A. C. Hess, "Periodic Hartree-Fock Studies of the Properties of the Cl₂/MgO (001) Interface," *J. Chem. Phys.* (in press).

2. Materials Theory and Modeling

A. C. Hess, J. P. LaFemina, D. L. Lessor, M. I. McCarthy, T. J. Godin,^(a) A. J. Skinner,^(a) J. C. White,^(a) G. K. Schenter, D. R. Baer, C. B. Duke,^(b) D. L. Blanchard,^(a) A. S. Gibson,^(c) and R. Haydock^(d)

The development of accurate, predictive models of contaminant flow in the groundwater and the design of effective remediation technologies to remove or immobilize contaminants from the subsurface environment

depend on understanding of subsurface chemical reactions and their mechanisms at the molecular level. Unfortunately, little is even known about the microscopic structure of the mineral surfaces on which these reactions occur.⁽¹⁾ A more basic problem confronting the design of efficient and effective remediation strategies is the need for sensitive and selective chemical sensors and fuses to detect and characterize environmental contaminants.⁽²⁾

Physics and Chemistry of Oxides and Silicates

One of the goals of the materials theory program is to obtain a detailed microscopic understanding of the surface structure and chemistry of metal oxides and thin film polymeric materials needed to begin solving these problems.

Structure of Oxide and Silicate Surfaces

Tight-binding, total-energy models have been enormously successful in predicting and elucidating the mechanism of semiconductor surface reconstructions. This model was recently extended to include cubic (MgO), cristobalite (SiO₂), and tridymite (SiO₂) oxides. In addition, a unified tight-binding, total-energy model was developed to describe the surface properties of the wurtzite, zincblende, and rocksalt phases of the catalytically important ZnO. These studies have extended the concept that the chemistry of oxide cleavage surfaces reflects the interplay of both topology (via the ability to undergo bond-length-conserving relaxations) and local chemical bonding (via the participation of *p* electrons in bonding versus non-bonding surface state bands).⁽³⁾ More importantly, they illustrate that the surface reconstructions of oxide materials follow the same simple physical and chemical principles first elucidated for semiconductor surfaces.⁽⁴⁾ These principles can be summarized as follows. The creation of the surface creates dangling bonds and dangling bond charge density in an energetically unfavorable configuration. Therefore, a driving force exists at the surface to redistribute this charge density so as to create an "insulating" surface

(a) NORCUS Postdoctoral Research Associate.

(b) Xerox Webster Research Center, 800 Phillips Road, Webster, New York.

(c) Joint MSRC-University of Oregon Materials Science Institute Postdoctoral Research Associate.

(d) Materials Science Institute, University of Oregon, Eugene, Oregon.

(i.e., pair up the dangling bond electrons and open a gap between the occupied and unoccupied surface states). This can be done by forming new bonds at the surface, either between surface atoms or between the surface atoms and adsorbates. Alternatively, the dangling bond charge density can be accommodated structurally, by rearranging the surface atoms so that charge transfer takes place between surface atoms, or electronically, through strong electron correlation. There is a tradeoff, however, between the drive to saturate the dangling bonds and create an insulating surface, and the amount of strain that can be accommodated by the surface and subsurface layers of the material. It is this balance between the energy lowering due to the reduction of dangling bonds and the energy cost due to induced strain that dominates the nature of the surface reconstruction. Consequently, surfaces with the ability to undergo approximately bond-length-conserving relaxations [e.g., SiO_2 , ZnO (zincblende), and ZnO (wurtzite)] exhibit large atomic motion surface relaxations, while those whose topology precludes such motion [e.g., MgO and ZnO (rocksalt)] exhibit small atomic motion relaxations.⁽³⁾

These studies have also identified an additional driving force for surface relaxation that is unique to oxide materials, namely the surface-state lowering produced by the rehybridization of O nonbonding p orbitals.⁽³⁾ For materials in which the oxygen atoms possess lone-pair electrons, the anionic surface states associated with the unrelaxed surface lie in the top of the nonbonding electron valence band, rather than in the gap. Consequently, rehybridization of the occupied nonbonding orbitals into occupied bonding hybrids provides an energy stabilization sufficient to drive surface relaxation or reconstruction. As before, however, the atomic motion accompanying the rehybridization must be allowed by the surface topology through approximately bond-length conserving motions. If not, the energy destabilization arising from the local strain environments will quickly offset the rehybridization energy gain and halt the relaxation.⁽⁴⁾ For the close-packed $\text{ZnO}(001)$ and $\text{MgO}(001)$ surfaces, the effect of surface topology is predominant and significant rehybridization is precluded by the surface topology.

Because defects play an important role in the surface chemistry of mineral and metal oxides, we are developing, in collaboration with researchers at the University of Oregon's Material Science Institute, a real-space density, functional-based method using the Harris-

Foulkes energy functional to compute the atomic and electronic defects in mineral oxides. This method has been used to compute the electronic structure and stability of the perfect (001) , (111) -Mg, and $(-1-1-1)$ -O surfaces of MgO , and use the local density of states at the surface to understand the relative stabilities and differing electronic structures of the surfaces in terms of the different surface chemical bonding. Because the recursion method⁽⁵⁾ is used to solve the Schrödinger equation in real space, the need for supercell networks of periodically repeated defects (used in k -space based methods) is eliminated. This approach allows us to examine isolated defects and, most importantly, charged defects that have a divergent Coulomb energy in supercell network-based methods.

Finally, we have investigated the use of alternative methods [e.g., modified neglect of differential overlap (MNDO)] for the computation of surface reconstructions in complex materials where the usual tight-binding models are inadequate. In addition, the development of the tight-binding bond model⁽⁶⁾ and the Block Recursion Method Library⁽⁷⁾ has continued, and the block recursion method has been used to investigate the properties of disordered materials.⁽⁸⁾

Low-Energy Electron Diffraction and Low-Energy Positron Diffraction from Oxide Surfaces

LEED and low-energy positron diffraction (LEPD) intensity analyses are important tools in the determination of surface atomic structure and the mechanisms of surface relaxation. The application of LEED to the study of insulator surfaces is complicated by surface charging effects. Modulated beam techniques can be used to reduce these effects, and we have carried out a modulated-beam LEED study on $\text{MgO}(001)$. LEED beam intensities were measured from the freshly cleaved (in UHV) $\text{MgO}(001)$ surface using beam energy modulation to avoid electrical charging. A dynamic LEED analysis of the data indicated 1) an interlayer relaxation of $1 \pm 2\%$ and a rumple of the first layer ions of $5 \pm 2.5\%$ of an interlayer spacing, 2) a new attenuation model containing no adjustable parameters gives fits to the data that are comparable to those using a constant V_p , and 3) a weak dependence of the optimum rumpling on the energy range of the data used. The structure determined in this study was in quantitative agreement with other experimental and theoretical structures reported in the literature for this surface.⁽⁹⁾ Work on the (111) and (110) surface of SnO_2 , in collaboration with researchers at the University of Texas at Austin, is currently in progress.

Surface Dynamics

The chemical and physical properties of surfaces are determined not only by the equilibrium (static) surface structure but by the dynamical force fields and surface vibrational structure. The use of quantum-mechanical (QM) molecular dynamics methods to study the dynamical properties of materials has increased greatly over the past few years. One advantage to the use of QM molecular dynamics simulations over conventional QM computations of total-energy derivative properties is the ability of molecular dynamics simulations to investigate temperature effects. During the past year, we have developed a QM molecular dynamic simulation capability based upon the same tight-binding, total-energy models described above. The use of a tight-binding Hamiltonian allows for the accurate computation of atomic forces more efficiently than *ab initio* potentials while retaining the essential quantum-mechanical aspects of interatomic bonding not reproduced by classical potentials. To test the predictive capability of this model QM molecular dynamic simulations were performed to determine both the bulk and (110) surface vibrational spectral densities for GaAs and InP, along with the GaAs(110)-p(1x1)-Sb and InP(110)-p(1x1)-Sb interfaces. These systems were chosen because of the effort that has been spent over the past decade on the development and optimization of their tight-binding Hamiltonians. Consequently, they provide a parameter independent test of the applicability of this approach. Moreover, experimental information on these systems exists, allowing for a direct assessment of the model's predictive capability. Our approach utilizes the tight-binding, total-energy model to compute the Hellman-Feynman forces that are used to drive the molecular dynamics simulation. A method for computing bulk phonon dispersion curves has been developed, and the dependence of both bulk and surface phonon frequencies upon temperature has been explored (Figure 2.1).

The results for the bulk vibrational energies are in semiquantitative agreement with experimental results, displaying approximately the same level of variance as other model computations. The (110) surface vibrational energies are in quantitative agreement with the experimental results, while the computed vibrational energies for the III-V(110)-p(1x1)-Sb(1ML) interface are in semiquantitative agreement with the experimental data. These comparisons give us great confidence that semiempirical models can be developed that are able to quantitatively describe the surface atomic

Figure 2.1. Temperature dependence of the zone center and zone edge spectral density of bulk GaAs as computed by tight-binding QM molecular dynamic simulations. Peak shifts and broadening with increasing temperature are indicative of the anharmonic nature of the potential.

and electronic structure, including surface atomic dynamics, of complex materials.

Physics and Chemistry of Polymers

Chemical sensors are a ubiquitous part of efficient and effective environmental remediation and restoration strategies. They are important for site characterization, for the monitoring of waste processing activities, and for postremediation site monitoring. Thin-film polymers are used as selective layers in many chemical sensor and fuse devices. Unfortunately, little is known about the properties of the polymer films, or their interactions with target species on a molecular level. One key to the improvement of existing selective layers

and the manufacture of new high-performance sensor devices is the development of a detailed understanding of the energetics and mechanisms of target molecule-polymer and polymer-polymer interactions.

Our initial work in this area has concentrated on understanding the polymer-polymer interactions in poly(ethylene terephthalate) (PET). PET has demonstrated a wide range of industrial potential, ranging from applications in the microelectronic and electronic packaging technologies to the medical arena. In our work, the polarized and unpolarized emission spectra for PET were measured with a wavelength resolution greater than that of existing published data, revealing a new phosphorescence peak at 564 nm. More importantly, the CNDO/S3-CI model was used to provide a detailed molecular description of the monomeric absorption and emission, shown schematically in Figure 2.2.⁽¹⁰⁾ There remain, however, several features of PET's emission spectrum (a fluorescence peak at 368 nm and the newly discovered phosphorescence at 564 nm) that are unexplainable in monomeric terms. It has been suggested that these features are attributable to states resulting from the formation of an associated ground-state dimer. To test this hypothesis, molecular dynamics simulations of multiple unit-cell, multiple-strand models of PET were performed at both low (77K) and room (300K) temperatures. These simulations indicated that the polymer strands never deviated significantly from their crystalline orientations. Quantum mechanical (CNDO/S3) CI computations performed on several minimum energy conformations sampled by the molecular dynamics simulation resulted in excited state energy spectra that were essentially identical to those of the monomer chain. This finding indicates a very weak coupling of the electronic structure of neighboring strands. Therefore, our computations indicate that these unexplained emissions must originate in the amorphous region of the film.⁽¹¹⁾

Finally, work on understanding the dual fluorescence arising from TICT in dimethylaminobenzonitrile has continued with *ab initio* computations of the ground-state potential energy surface.

Physics and Chemistry of Minerals

The focus of the mineral chemistry program is to gain a fundamental understanding of the properties of a variety of naturally occurring solid materials using the

Figure 2.2. Schematic indication of the states responsible for the monomeric absorption and emission in poly(ethylene terephthalate). Computed CNDO/S3 energies are shown in electronvolts without parenthesis, while experimentally measured absorptions and emissions are in nanometers without parenthesis. The values in parenthesis provide the unit comparisons.

techniques of *ab initio* solid-state quantum chemistry and molecular dynamics. The efforts include both investigations on specific systems and the development and refinement of the necessary computational methodologies. The current areas of research include 1) the bulk, internal and external surface properties of the clay minerals kaolinite, pyrophyllite, and talc, and 2) the bulk and surface properties of two hydrated nitric acid compounds present in the Antarctic atmosphere.

The computational method currently being used provides SCF solutions to the HF-Roothan equations subject to periodic boundary conditions. This approach is implemented in the program CRYSTAL.⁽¹²⁾ The multistep process⁽¹³⁾ begins with an atomic orbital basis and evaluates the necessary one- and two-electron integrals required to construct the infinite dimension Fock matrix in direct space. The direct space Fock matrix and its associated overlap matrix can then be Fourier transformed into reciprocal space, where they become block diagonal. The solutions to the corresponding matrix equations yield a set of eigenvalues and eigenvectors for each point in

reciprocal space. These results are then used to construct a new direct space density matrix that redefines the direct space Fock matrix. The cycle continues until the total energy converges to a specified tolerance. Some of the physical properties of the solid that can be calculated from the final self-consistent wavefunction include the band structure, total and projected density of states, charge density and density deformation maps, electrostatic potentials, electric fields, and field gradients.

This method utilizes many of the concepts and technologies of molecular quantum chemistry. In particular, the basis sets, integral evaluation routines, and SCF iteration schemes from molecular calculations are directly applicable to condensed-phase systems. The primary advantage of the periodic HF calculations is that they avoid the limitations of treating crystalline systems as finite molecular clusters by exploiting the full crystallographic space group symmetry of the system. The incorporation of translational symmetry into the calculations eliminates "edge" effects associated with truncating finite clusters and enhances the computational feasibility. For materials that contain little or no point symmetry, such as the clays, the calculations are practically limited by the time and size restrictions associated with the storage of certain types of integrals. To alleviate these constraints, a joint project with N. Harrison and V. R. Saunders of the SERC Daresbury Laboratory, United Kingdom, is in progress to develop "direct" methods in which such integrals are reevaluated as needed during the SCF cycles, instead of being permanently stored on disk. This approach will greatly reduce the storage requirements and increase the size of the system (number of atoms in the unit cell) that can be treated.

Electronic Structure of the Clay Minerals

The periodic HF linear combination of atomic orbitals (LCAO) have been used to investigate the properties of the clay minerals kaolinite⁽¹⁴⁾ ($\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$), pyrophyllite⁽¹⁵⁾ ($\text{Al}_2\text{Si}_4\text{O}_{10}(\text{OH})_2$), and talc ($\text{Mg}_3\text{Si}_4\text{O}_{10}(\text{OH})_2$) using STO-3G and modified 6-21G and 6-21G* basis sets. The clay minerals are the abundant natural decomposition products of aluminosilicates in aqueous environments and, hence, affect the chemistry and physics of sedimentary systems. Since these materials are layered structures, they often function as a natural ion-exchange medium for intercollated species. Therefore, the clays play a key role in the subsurface transport of hazardous soil contaminants. The clays are typically formed in small particles

(2 to 40 μm) and often contain substitutional defects, which makes an unambiguous experimental determination of the structure very difficult.⁽¹⁶⁾

One example of this is the ongoing dispute in the literature over the internal hydrogen position in kaolinite.⁽¹⁷⁾ The framework structure of this clay is composed of a sheet of corner sharing SiO_4 tetrahedra linked by common oxygens to a sheet of edge sharing AlO_6 octahedra, which are bound to both inner- and outer-hydroxyl groups. The three degrees of freedom associated with the position of the internal hydroxyl [the OH bond length (r), the Al-O-H angle (θ), and the angle of the OH with respect to the Al-O-Al framework (ϕ)] were optimized in periodic HF calculations. The optimized coordinates are found to be $r = 0.99\text{\AA}$, $\theta = 108^\circ$, and $\phi = 62^\circ$. These optimization calculations were only feasible using the STO-3G basis set and required in excess of 7 Gbytes of disk storage and several CPU hours on a CRAY 2 per point.

Calculations of the physical properties of kaolinite were performed using a modified 6-21G basis set on the optimized structure. These results provide insight into the nature of the chemical bonding in the clays. The computed band structure is indicative of the insulating character of this material. The total and projected valence density of states revealed that the majority of the valence states are composed of oxygen 2s and 2p atomic orbitals with significantly smaller contributions from the overlapping corresponding Si, Al, and H orbitals. The corner sharing SiO_4 tetrahedra in kaolinite form a silicate ring system known as the ditrigonal cavity. It has been proposed that these sites in the crystal are chemically active and function as weak Lewis bases.⁽¹⁸⁾ Crystal charge densities and density deformation maps have been computed in the basal plane of the silicate ring system. The calculations revealed only a slight charge density in the center of the ditrigonal cavity (10^{-4} to 10^{-5} e/bohr³) and an oblate distortion of the oxygen charge density directed towards the center of the ring system. Electrostatic potential maps were computed in the chemically accessible regions of the structure near the ditrigonal cavities. These indicate that the center of the cavity is at a negative potential relative to the nearby silicon framework. Maps of the potential in a plane perpendicular to the layers of the material illustrated the extensive interlayer hydrogen bonding. Similar calculations have been carried out for the clays pyrophyllite (Figure 2.3) and talc. These are 2:1 clays (2 sheets of

Figure 2.3. Structure of 2:1 dioctahedral clay mineral, pyrophyllite ($\text{Al}_2\text{Si}_4\text{O}_{10}(\text{OH})_2$).

" SiO_4 " and one sheet of " AlO_6 " per clay layer). The interlayer bonding in these materials is much weaker than in kaolinite, and hence, they show more extensive intercalation chemistry.

This *ab initio* investigation of the clays demonstrated the feasibility of studying low symmetry solids using periodic HF theory. These techniques provide a powerful means to investigate systems previously judged to be too complex to be treated using *ab initio* methods.

Electronic Structure of Zeolites

Zeolite materials with an aluminosilicate framework containing networks of cavities that permit the transport of a range of molecules and ions that are important in environmental and catalytic processes.⁽¹⁹⁾ Certain zeolites have been used in the extraction of Sr and Cs from nuclear wastes and are involved in a variety of catalytic processes. Although most zeolite structures are very complex with many atoms in a unit cell, the physical properties of these materials can often be understood by studying small prototypical structures.

Sodalite ($\text{Na}_8\text{Al}_6\text{Si}_6\text{O}_{24}\text{Cl}_2$) and its silicon substituted analogue, silica sodalite ($\text{Si}_{12}\text{O}_{24}$), are examples of these. The structure of sodalite-type compounds consists of a network of apex sharing SiO_4 and AlO_4 tetrahedra that form a cubo-octahedron within each unit cell. The naturally occurring compounds of this type often trap a variety of molecular and ionic species inside the caged network. For example, native sodalite incorporates the species $(\text{Na}_4\text{Cl})^{3+}$ in order to charge balance the excess electron density on the framework. Silica sodalite has the same structural framework as sodalite, except all of the Al atoms are replaced with Si atoms. This results in a compound whose framework is naturally charge neutral.

In order to gain a better understanding of the behaviour of species trapped in these cavities, we have investigated the electrostatic properties of clean and substituted sodalite-like compounds using periodic HF theory. An in-depth basis set study has been done on silica sodalite (Figure 2.4). This work tested the agreement of the calculated structural parameters with the experimental values using minimal, Pople split valence, Dunning DZP, and reoptimized split valence basis sets.⁽²⁰⁾ The reoptimized split valence basis with an 8-31G* on Si and an 8-51G* set on O gave the best results.

In order to test the effect of perturbations of the clean lattice due to the presence of trapped species, the electrostatic properties were calculated for silica sodalite compounds containing He, Ne, or Ar in the cage.⁽²¹⁾ These structures contain 18 to 20 atoms per unit cell and the calculations were run with a 6-21G* basis set. The results indicated that the potential in the center of the cage was constant and slightly attractive to a positive test charge. Placing Ne or Ar in the center of the cage had only a minor effect on general features of the potential. However, when He was trapped in the material, the potential calculations revealed extensive polarization of the charge densities centered on the oxygen atoms in the framework.

Current studies are under way to test the effect of trapping larger molecular species in the silica sodalite cage. This work includes calculations on the naturally occurring sodalite-like species (i.e., native sodalite, sodalite perchlorate, and ethylene glycol silica sodalite).

Calculations are also in progress that systematically substitute Al for the Si in the framework. These are

Figure 2.4. Structure of silica sodalite ($\text{Si}_{12}\text{O}_{24}$) (light colored balls are silicons and darker objects are oxygens).

designed to further probe the nature of the bonding in these simple zeolites and investigate the acid/base chemistry associated with such substitutions. In addition, ongoing computational refinements have recently permitted studies to be conducted on larger known zeolites. Current investigations include the study of both siliceous zeolite A and siliceous mordenite, each of which contain over 150 heavy atoms per unit cell.

References

1. M. F. Hochella, Jr., "Mineral-Water Interface Geochemistry," M. F. Hochella, Jr., and A. F. White, eds., *Reviews in Mineralogy* **23** (1990).
2. "Environmental Materials and Interfaces," PNL-SA-19675 (CONF-9009424) Pacific Northwest Laboratory, Richland, Washington, 1991.
3. J. P. LaFemina, and C. B. Duke, *J. Vac. Sci. Tech.* **A9**, 1847 (1991).
4. J. P. LaFemina, *The Handbook of Surface Science, Physical Structures*, W. N. Unertl, ed., Elsevier, Amsterdam.
5. R. Haydock, *Solid State Physics*, H. Ehrenreich, F. Seitz, and D. Turnbull, eds., Vol. 35, Academic Press, New York.
6. A. J. Skinner, and D. G. Pettifor, *J. Phys.: Cond. Matter* **3**, 2029 (1991).
7. T. J. Godin, and R. Haydock, *Comp. Phys. Comm.* **68**, 123 (1991).
8. T. J. Godin, and R. Haydock, *Europhysics Lett.* **14**, 137 (1991).
9. D. L. Blanchard, D. L. Lessor, J. P. LaFemina, D. R. Baer, W. K. Ford, and T. Guo, *J. Vac. Sci. Tech.* **A9**, 1814 (1991).
10. J. P. LaFemina, and G. Arjavalasingam, *J. Phys. Chem.* **95**, 948 (1991).
11. J. P. LaFemina, D. R. Carter, and M. B. Bass, *J. Phys. Chem.* (in press).
12. R. Dovesi, C. Pisani, C. Roetti, and V. R. Saunders, Program Crystal, Quantum Chemistry Program Exchange, Program #577, Indiana University, Department of Chemistry.
13. C. Pisani, R. Dovesi, and C. Roetti, *Hartree-Fock Ab Initio Treatment of Crystalline Systems*, Springer-Verlag, New York, 1988.
14. A. C. Hess, and V. R. Saunders, "Periodic *Ab Initio* Calculations of the Low Symmetry Mineral Kaolinite," *J. Phys. Chem.* (in press).
15. A. C. Hess, and D. R. Fried, "Periodic *Ab Initio* Calculations of the 2:1 Dioctahedral Clay Mineral Pyrophyllite" (in press).
16. D. J. Greenland, and M.H.B. Hayes, eds., *Chemistry of Soil Constituents*, John Wiley & Sons, Inc., New York, 1985.

17. R. F. Giese, "Kaolin Minerals: Structures and Stabilities," *Reviews in Mineralogy: Hydrous Phyllosilicates*, S. W. Bailey, ed., Vol. 9, Mineralogical Society of America, Washington, D.C., 1988.
18. G. Sposito, *The Surface Chemistry of Soils*, Oxford University Press, New York, 1984.
19. J. A. Rabo, ed., *Zeolite Chemistry and Catalysis*, ACS Monograph 171, American Chemical Society, Washington, D.C., 1976.
20. A. C. Hess, and J. C. White, "An *Ab Initio* Hartree-Fock Study of Sodalite and Silica Sodalite" (in press).
21. R. Nada, C.R.A. Catlow, R. Dovesi, and V. R. Saunders, "An *Ab Initio* Hartree-Fock Study of the Ilmenite-Structured MgSiO_3 ," *Proc. Roy. Soc.* (in press).

3. Biomolecular Modeling and Simulation

R. L. Ornstein, G. E. Arnold, M. B. Bass,^(a)
M. D. Paulsen,^(a) et al.^(b)

Biomolecular Modeling and Simulation activities are currently focused on applying existing modeling/simulation methods and developing improved methods for coupled theory-experimental studies in order to uncover structure/function/dynamics relations of key biomacromolecules. The primary focus is to design new and improved catalytic functionality and/or alternative substrate specificity of enzymes/proteins with an emphasis on developing near-, mid- and long-term strategies specific for remediation and restoration

(a) NORCUS postdoctoral fellows at PNL.

(b) F. J. Brockman, Earth and Environmental Science Center, PNL. S. G. Sligar, Departments of Biochemistry and Chemistry, The Beckman Institute, University of Illinois, Urbana-Champaign. D. Filipovic, NORCUS postdoctoral fellow, Departments of Biochemistry and Chemistry, The Beckman Institute, University of Illinois, Urbana-Champaign. R. Rein, Biophysics Department, Roswell Park Memorial Institute, Buffalo, New York, and the Department of Biophysical Sciences, S.U.N.Y. at Buffalo, New York. M. Shibata, Research Scientist, and W. A. Sokalski, Visiting Research Scientist, Roswell Park Memorial Institute. N. Luo, NORCUS postdoctoral fellow, Roswell Park Memorial Institute. J. A. Braatz, B. G. Gladstone, and D. F. Hopkins, U.S. Department of Energy Science and Engineering Research Semester predoctoral interns at PNL. M. P. Hughes, W. A. Jaquysh, and L. A. Thomson, NORCUS predoctoral interns at PNL. S. Sun and J. Lai, NORCUS graduate students, Roswell Park Memorial Institute.

of contaminated DOE sites. Proteins are engineered to change their functionality and/or properties by altering the amino acid sequence. It is impractical to make all possible combinations of amino acid changes for even the smallest protein. Starting with a 3-D x-ray crystal and/or NMR structure as input, computer modeling and simulations can play a valuable, if not essential, and unique role in determining key amino acids for modification.

"Long" Protein Simulations: Motions in Different Time Domains and X-Ray and Solution NMR Structure Differences

"Any attempt to understand the function of proteins requires an investigation of the dynamics of the structural fluctuations and their relation to activity and conformational change."⁽¹⁾ Protein motions range from the rapid local motions of the individual amino acids to slow collective distortions of secondary structure elements or domains. While sidechain rotations at the surface of a protein typically occur in tens to hundreds of picoseconds, torsional rotations of buried groups and relative motion of different domains often have longer characteristic times by one to several orders of magnitude. For practical reasons, most protein molecular dynamics simulations have been performed for less than 100 ps. Occasionally, a protein simulation has been extended beyond 200 ps, but as of 1991, we are not aware of any published protein simulations beyond 400 ps.

Explicitly including only the waters of crystallization in an otherwise implicit solvent model simulation, we have performed a 3,000 ps molecular dynamics simulation on the 76 amino acid protein ubiquitin,⁽²⁾ a 2,000 ps molecular dynamic simulation on the 62 amino acid protein erubatoxin b,⁽³⁾ and a 1,200 ps simulation on the enzyme T4 lysozyme.⁽⁴⁾ These proteins were chosen because of the availability of a wide range of experimental data that can be compared to the simulations (size and other properties were also important considerations). Understanding the limitations and unique information resulting from extended simulations is important in achieving protein/enzyme redesign goals. Some results are discussed for the case of ubiquitin.

The crystal structure of human erythrocytic ubiquitin has been refined at 1.8 Å resolution.⁽⁵⁾ The overall structure of ubiquitin is extremely compact and tightly

hydrogen-bonded and does not contain any disulfide bridges. The solution NMR structure of ubiquitin⁽⁶⁾ has been determined and shown to be virtually identical with the x-ray crystal structure. Our simulated average structure retains much of the character of the starting x-ray crystal structure.

Figure 3.1 shows the main chain root mean square (RMS) deviation of alpha-helix 1 and beta-strand 1.

Figure 3.1. The RMS deviation of all main chain heavy atoms in alpha-helix 1 (thick curve) and beta-strand 1 (dotted curve) plotted versus the time course of the simulation of ubiquitin.

Two points are noteworthy. First, this shows that all secondary structure domains are not equally stable, but rather different domains of a protein may have different ranges of motion. Second, the conformational flexibility of beta-strand 1 was not apparent from the simulation until after 500 ps. Thus, relatively long molecular dynamics simulations that use protein x-ray

crystal coordinates for the starting structure and a computationally efficient solvent representation may be used to gain an understanding of conformational and dynamic differences between the solid-crystal and dilute-solution states. Moreover, such simulations may be able to uniquely characterize conformational mobilities in a wide range of time domains that are essential to understanding structure/function/dynamics relationships.

Structure/Function/Dynamic Relations for the Biodegradative Enzyme Cytochrome P450cam

Only a few important microbial biodegradative enzymes have had their 3-D structures solved by experimental methods. In the absence of such structural information, it is usually not possible to gain an understanding of basic structure/function/dynamic relationships or to use rational engineering methods to redesign an enzyme for enhanced biodegradative functions. High resolution crystal structures are available for camphor-bound cytochrome P450cam, substrate-free and with a variety of substrate analogues and inhibitors.⁽⁷⁾ Cytochrome P450cam, which catalyzes the regiospecific and stereospecific hydroxylation of camphor, contains over 400 amino acids (over 7,000 atoms are involved in our simulations after hydrogens, heme, substrate, and waters of crystallization are added). Initial molecular simulations have focused on obtaining a fundamental molecular understanding of the complementarity and dynamic motions involved in P450cam interactions with its native substrate camphor^(8,9) and the substrate analogues norcamphor^(10,11) and thiocamphor.⁽¹²⁾

While cytochrome P450cam catalyzes the hydroxylation of camphor to 5-*exo*-hydroxycamphor with 100% stereospecificity and 100% efficiency, norcamphor is hydroxylated by this enzyme yielding 45% 5-*exo*-, 47% 6-*exo*-, and 8% 3-*exo*-hydroxynorcamphor with only 12% overall efficiency.¹³ This past year, we performed and compared simulations with camphor-bound or norcamphor-bound P450cam to better understand the underlying factors controlling P450cam regiospecificity and catalytic efficiency.^(9,10) We found that camphor has a significantly reduced level of rotational mobility in the active site in comparison to norcamphor, suggesting that product specificity is, at least partially, controlled by the mobility of the substrate within the

active site. Second, the high mobility of norcamphor in the active site leads to an average increase in separation between the heme iron and the substrate of about 1.0 Å; this increase in separation may well be the cause of the uncoupling of electron transfer from product formation (reduced efficiency) when norcamphor is the substrate. Third, the active site water located in the norcamphor-bound crystal structure possesses mobility that correlates well with the spin-state equilibrium of this enzyme-substrate complex. Therefore, it is not necessary for the substrate to enter this deeply buried active site prealigned for catalytic activity.

The x-ray crystal structure of thiocamphor-bound P450cam⁽¹⁴⁾ gave the most unexpected orientation in the active site of any substrate analogue considered to date. Unlike camphor, carbons 5 and 6 of thiocamphor are positioned away from the heme iron, while carbon 3 is positioned close to the heme iron. Thiocamphor is hydroxylated at carbon 5 (64%), 6 (34%), and 3 (2%) with 98% efficiency.⁽¹⁵⁾ Therefore, the crystallographically observed thiocamphor-P450cam structure is not consistent with the catalytic observation. In an attempt to better understand this puzzling situation, we performed two molecular dynamics simulations on thiocamphor-bound P450cam, starting with the x-ray structure, with and without the presumptive heme iron-oxo reactive intermediate. Figure 3.2 (top) depicts the variation over the simulation for the distance between the heme iron and the 3- and 5-position carbons of thiocamphor for the case without an oxygen ligated to the heme iron; this situation is similar to that in the x-ray crystal structure. In over 95% of the computed structures, carbon 3 is closer to the heme iron; although, there are clearly two such configurations that are also in agreement with the x-ray data. In the second thiocamphor simulation with the presumptive heme iron-oxo intermediate, the distance between the oxygen (of the iron-oxo) and the 3- and 5-position carbons are shown in Figure 3.2 (bottom). Interestingly, during the equilibration the two carbons "switch" such that now the 3-position carbon face is away from the heme, while the 5-position carbon face is close to the heme. Not only do these simulations offer a consistent explanation for the structural and catalytic experiments, they also suggest a unique role for the simulations, since the iron-oxo intermediate is too unstable to be observed.

Redesigning Enzymes to Enhance Environmental Remediation

Contamination in the deep subsurface is commonly dilute, highly dispersed, inaccessible, and exists in large volume. Thus, the nature of deep contamination substantially limits or precludes the application of many remediation strategies that are currently being applied to remediate surface and near-surface contamination. Cost-effective remediation of synthetic substances in deep subsurface environments is, for the most part, limited to novel bioremediation alternatives.⁽¹⁶⁾ The unique advantage of rational redesign is its ability to foster discovery of new enzymic forms that would otherwise have an extremely low probability of evolving biologically under either laboratory-controlled or natural conditions (which is to be expected for an enzyme requiring many simultaneous and specific amino acid changes to efficiently catalyze a significantly different substrate).⁽¹⁷⁾ The objectives of our research are to 1) improve and expand the degradative capacity of biodegradative enzymes in indigenous microorganisms, and 2) design enzymes for extracellular degradation in harsh environments where microorganisms may be unable to grow at adequate levels. Our efforts will be focused on developing systems for remediation of contaminants at DOE sites. Due to the complexity and interdisciplinary nature of our objectives, close collaborations with scientists in the Environmental Sciences Research Center (ESRC) at PNL and several universities have been established. These collaborations play a decisive role in our fundamental and applied enzyme redesign studies.

As a first step in the redesign of the activity of P450cam for a substrate significantly different from camphor, we have chosen to first modify the specificity of P450cam for the substrates camphor and norcamphor. Using the previous structure/function/dynamics relationships of camphor and its analogues bound to P450cam, we are currently attempting to design mutants of P450cam that will 1) have increased efficiency for hydroxylating norcamphor, and 2) preferentially hydroxylate the 3-position of camphor and/or norcamphor. Achieving these intermediate redesign goals will give us considerable confidence and guidance towards the goal of modifying the specificity of P450cam to initiate the degradation pathway for targeted pollutants. Our first prediction involving the former goal is described below.

a) The time course variation for the distances between the heme iron and thioamphor 5- and 3-position carbons are indicated by the dotted and solid curves, respectively.

b) The time course variation for the distances between the oxygen of the heme iron-oxo intermediate and thioamphor 5- and 3-position carbons are indicated by the dotted and solid lines, respectively.

Figure 3.2. Active-site mobility of thioamphor during two molecular dynamics simulations.

We have analyzed the detailed dynamics motion of each amino acid residue that lines the buried active site in camphor-bound⁽⁹⁾ and norcamphor-bound⁽¹⁰⁾ P450cam.

Most of the backbone and sidechain atoms of the active site residues are very stable, but chl-2 of Phe-87 exhibits very significant flexibility. However, when the active site was filled with water but without substrate, simulation showed Phe-87 was dynamically quiet,⁽¹⁸⁾ even though the x-ray crystal structure⁽¹⁹⁾ and time-average simulated structure of substrate-free P450cam are surprisingly similar to those of the camphor-bound enzyme.

On the basis of these results and preliminary modeling and simulations, we proposed replacing Phe-87 by the bulkier aromatic residue Trp. The simulation showed reduced mobility of norcamphor in the active site as compared to the wild-type simulation. The prediction⁽²⁰⁾ (when no experimental evidence was available) suggested that norcamphor would be more tightly coupled to electron transfer, as is indicated by the shorter separation between the substrate center-of-mass and the heme iron for the mutant (Figure 3.3). This shortening in the substrate heme separation is a direct result of the tighter packing of residue 87 in the mutant (Figure 3.4). Per our recommendation, Dr. D. Filipovic in Professor S. G. Sligar's laboratory constructed the mutant and obtained the results on coupling. The theoretical prediction had been tested and confirmed; the mutant had twice the coupling exhibited by the wild enzyme when the substrate was norcamphor. These preliminary results suggest the validity of computational methods in characterizing and predicting the metabolism of small substrates by cytochrome P450cam.

References

1. M. Karplus, and J. A. McCammon, *Ann. Rev. Biochem.* **53**, 263 (1983).
2. R. L. Ornstein, and J. A. Braatz, *J. Biomol. Struct. Dynam.* **8**, a158 (1991); J. A. Braatz, M. D. Paulsen, and R. L. Ornstein, *J. Biomol. Struct. Dynam.* (1992) (in press).

Figure 3.3. Depicts the distance between the center-of-mass of norcamphor and the heme iron for both the Phe-87 → Trp mutant simulation (solid line) and the wild-type simulation (dashed line).

3. B. G. Gladstone, and R. L. Ornstein, Fifth Annual Symposium of the Protein Society, June 1991, Baltimore, Maryland; B. G. Gladstone, and R. L. Ornstein (in preparation).
4. G.E. Arnold and R.L. Ornstein, Fifth Annual Symposium of the Protein Society, June 1991, Baltimore, Maryland.
5. S. Vijay-Kumar, C. E. Bugg, and W. J. Cook, *J. Mol. Biol.* **194**, 531 (1987).
6. P. L. Weber, S. C. Brown, and L. Mueller, *Biochemistry* **26**, 7282 (1987).
7. T. M. Poulos, *Meth. Enzymol.* **205**, in press (1991).
8. M. D. Paulsen, and R. L. Ornstein, *PROTEINS: Struct. Funct. Genetics* **11**, 184 (1991).

Figure 3.4. The distance of the terminal sidechain carbon atom of residue 87 is measured to the heme iron in norcamphor-bound to wild-type or mutant P450cam. The solid line represents the distance to the CH_2 atom of Trp-87 (mutant), while the dashed line represents the distance to the C_γ atom of Phe-87 in the wild-type enzyme.

9. M. D. Paulsen, M. B. Bass, R. L. Ornstein, *J. Biomol. Struct. and Dyn.* **9**, 187 (1991).
10. M. B. Bass, M. D. Paulsen, and R. L. Ornstein, *PROTEINS: Struct. Funct. Genetics* **12** (in press).
11. M. B. Bass, D. F. Hopkins, W. A. Jaquysh, and R. L. Ornstein, *PROTEINS: Structure, Function, and Genetics* **12** (1992) (in press).
12. M. D. Paulsen, M. B. Bass, and R. L. Ornstein, 1991 President's Meeting of the International Society of Quantum Biology and Pharmacology, Abstract #16, Stanford University, Palo Alto, California, September 1991.
13. W. M. Atkins, and S. G. Sligar, *J. Am. Chem. Soc.* **111**, 2715 (1989).
14. R. Raag, and T. L. Poulos, *Biochemistry* **30**, 2674 (1991).
15. W. M. Atkins, and S. G. Sligar, *J. Biol. Chem.* **263**, 18842 (1988).
16. F. J. Brockman, and R. L. Ornstein, Hazardous Materials Control/Superfund '91, in Proceedings of the 12th National Conference, pp. 264-266 (December 3-5, 1991, Washington, D.C.), Hazardous Materials Control Research Institute, Greenbelt, Maryland, 1991.
17. R. L. Ornstein, On Site Bioreclamation: Processes for Xenobiotic and Hydrocarbon Treatment, R. E. Hinchee and R. F. Olfenbuttel, eds., pp. 509-514, Butterworth-Heinemann, Boston, Massachusetts; R. L. Ornstein, in Proceedings of National Research & Development Conference on the Control of Hazardous Materials, pp. 314-319, Hazardous Materials Control Research Institute, Greenbelt, Maryland, 1991.
18. M. D. Paulsen, M. B. Bass, G. E. Arnold, and R. L. Ornstein, 35th Annual Biophysical Society Conference, Abstract W-PM-D3 [*Biophysical J.* **59**, 412a (1991)], San Francisco, California, 1991.
19. T. L. Poulos, B. C. Finzel, and A. J. Howard, *Biochemistry* **25**, 5314 (1986).
20. M. B. Bass, and R. L. Ornstein, Regional ACS Meeting: Molecular Design & Modeling, Indiana University-Purdue University, Abstract #558, Indianapolis, Indiana, 1991; M. B. Bass, and R. L. Ornstein, International Symposium on Cytochrome P450 of Microorganisms, Book of Abstracts, pp. 21-22, Berlin, 1991; M. B. Bass, M. D. Paulsen, and R. L. Ornstein, in Proceedings of the 7th International Conference on Biochemistry and Biophysics of Cytochrome P450: Structure & Function, Biotechnological & Ecological Aspects, A. I. Archakov, ed., Moscow, Russia, 1991 (in press).

4. Molecular Science Software Group

**R. A. Bair, D. F. Feller, M. W. Feyereisen,
R. A. Kendall, M. A. Thompson, M. S. Stave,
and R. J. Littlefield**

Recent advances in computer technology offer opportunities to model the complex molecular processes that

occur in natural systems. The research and development program in Molecular Science Software is focused on the application of these emerging computer technologies and advanced computer architectures to state-of-the-art theoretical and computational techniques in molecular science. This mission includes the development of significant improvements in both the methods and capabilities of molecular science computations, and in the tools for performing these computations and analyzing the results. The current effort addresses areas where significant advances in software technology are needed to fulfill the missions of the MSRC. To address these software needs, a research and development program has been undertaken to

1. develop a new generation of quantum chemistry and molecular dynamics computer codes for large-scale molecular science calculations on cost-effective, massively parallel computer systems
2. develop integrated computer program and data systems with modern graphical user interfaces (GUI) for molecular simulations, with a focus on the program systems described above
3. develop visualization tool kits for analyzing the massive amounts of data obtained from large-scale molecular simulations, with an emphasis on the needs of the computer programs systems described above.

The following sections describe the accomplishments and directions in each of the areas identified above. Of these, the first two areas were actively pursued in FY 1991. In the third area, only limited exploratory work was accomplished with more substantial efforts planned for FY 1992.

Concurrent Supercomputing Consortium

To support the development of molecular simulation codes for massively parallel computers, we are participating in the Concurrent Supercomputer Consortium (CSC), which was organized in 1990 by the California Institute of Technology, Caltech's Jet Propulsion Laboratory (JPL), Argonne National Laboratory (ANL), PNL, Center for Research on Parallel Computation (CRPC), Intel's Scientific Supercomputing Division (SSD), National Aeronautics

Space Administration (NASA), NSF Computational Science and Engineering, Lawrence Livermore National Laboratory (LLNL), Los Alamos National Laboratory (LANL), Purdue University and Sandia National Laboratory (SNL). The mission of the CSC is to advance computational science by applying concurrent computers with unprecedented levels of performance and providing opportunities for collaborations and exchange of information on computational techniques, algorithms, programming methodologies, and software.

In May 1991, the Touchstone Delta System was installed at the California Institute of Technology and acceptance testing began. Researchers in the Molecular Science Software Group, along with R. Littlefield from the MSRC Computing and Information Sciences program, have been active in developing acceptance tests for the Delta system and analyzing results of the test runs. This work has been invaluable to the CSC staff that have been working to accept and integrate the massively parallel Touchstone Delta Computer (with 512 numeric processors).

One of the most difficult aspects of the implementation of quantum chemistry and molecular dynamics codes on massively parallel computer systems is the design of algorithms that perform efficiently on hundreds (or thousands) of processors. The overall experience gained by our group during the acceptance phase has already proven valuable to both the mission of the MSRC and to our collaborators, including the Theoretical and Computational Chemistry Group at Argonne National Laboratory, and the Advanced Research Computing Group at the Science and Research Council (SERC) Daresbury Laboratory in the United Kingdom. As a whole, these efforts lay the technical groundwork necessary to develop many of the quantum mechanical algorithms for moderately parallel systems with a few hundred processors. We are also pursuing the algorithm research necessary to implement quantum chemistry codes for the next tier of massively parallel computers -- those having more than 1,000 processors.

Graphical User Interfaces

Modern GUIs have proven to be both convenient and powerful, substantially reducing the learning time required to run complex application programs, while

providing broad access to the computational power of the computer system. GUIs for the large, complex computational chemistry codes used today would greatly facilitate the use of these codes for molecular simulations, and provide much needed assistance to researchers who are not experts with a particular code or method. GUIs would 1) reduce the time required to prepare the data input, 2) reduce the errors that arise in the preparation of the data input, and 3) facilitate the analysis of the data generated in the simulations. GUIs also provide the foundation for both the graphical analysis of the results from molecular simulations (above) and the integration of discrete computer programs into a comprehensive, problem-oriented package. These computer programs will operate in a distributed computing environment, where steps in a computation are performed on the most appropriate available computing resource, whether local or remote.

In 1991, work continued on the Computational Chemistry Input Assistant (CCIA), an initial prototype of a GUI to support the GAMESS^(a) electronic structure package. The current implementation was done on an Apple Macintosh using SuperCard,^(b) a HyperCard-like development environment that permits rapid assembly of windows and their components, as well as rapid testing of functionality as a result of its ability to run in an interpreted mode.

During the past year, options were added to CCIA to permit easy access to the MSRC basis set library, and a rudimentary help facility was created that can, on request, advise users on the choice of basis sets. Molecules may be built graphically by calling the Chem3D⁽¹⁾ molecular modeling software and the resulting structure will be converted to the necessary cartesian coordinates and analyzed for symmetry elements by CCIA. When the user is satisfied with the input, it can be sent to any computational resource on a network. This may be a desktop workstation or a supercomputer center across the country.

(a) The original version is due to M. Dupuis, D. Spangler, J. J. Wendoloski, National Resource for Computation in Chemistry, University of California, 1980, Program QC01. This version is due to M. W. Schmidt, K. K. Baldrige, J. A. Bontz, J. H. Jensen, S. Koseki, M. S. Gordon, K. A. Nguyen, T. L. Windus and S. T. Elbert, *QCPE Bulletin* 10, 52 (1990).

(b) SUPERCARD V1.5, Silicon Beach Software (1990) 9770 Carroll Center Road, San Diego, California.

Another feature added to CCIA was a database of *ab initio* results. It contains the results of HF and perturbation theory calculations on small molecules using many of the popular basis sets in the chemical literature, as well as the new correlation-consistent⁽²⁾ basis sets. Entries for a particular molecule include the method by which the calculation was performed (e.g., level of perturbation theory, etc.), the total energy, the geometry and a wide assortment of properties. The goal is to provide users of the interface with ready access to information that would be helpful to them at the time they are creating input for the computational chemistry application. In collaboration with Professor David Maier and Ms. Judy Cushing of the Oregon Graduate Institute and Mr. D. M. DeVaney of PNL, we have begun to examine commercially available object-oriented database packages in order to determine their suitability for scientific work.

Parallel Applications for Molecular Computations

The development of accurate, yet computationally tractable, models of the mechanisms, energetics, and dynamics of molecular processes in the environment will require the development of new computational chemistry codes to both reduce the cost and extend the range of the simulations. Because the potential capacity and performance of large-scale parallel computer systems is well in excess of conventional supercomputers, these systems are becoming an increasingly important tool for molecular simulations. Full utilization of the computing capabilities offered by these machines will, in fact, be essential if we are to successfully pursue much of the work proposed in the EMSL (i.e., the modeling and simulation of molecular reactions, molecular spectra, and other molecular processes in solution, on surfaces, and at interfaces). Although a parallel processor system with potentially the same computing power as the fastest uniprocessor system costs substantially less it has proven difficult to deliver this potential savings to the user. Programming environments and computer languages on current parallel platforms are typically variations of serial environments and tools. These did not have parallelism as a design goal and do not afford a straightforward path toward parallel implementation of existing applications. Thus, most current computational chemistry applications do not yet run efficiently on parallel machines. The work to date suggests that, to

make effective use of multiprocessor systems, molecular simulation codes need to be developed to explicitly take advantage of parallel architectures from the beginning, especially in the massively parallel domain where hundreds or thousands of processors are applied to a single simulation.

Parallel Software and Computing Environments

An Intel iPSC/860 (16 node) acquired in 1991 has been put to extensive use in code development and limited production calculations. Code development on this machine includes continuing work on parallel ARGOS, and on implementations of a standard SCF code and the DISCO direct SCF code; development of a molecular dynamics code has also begun. These codes exercise key aspects of a parallel system. The standard SCF code is part of the Utah Mess Kit program system developed at the University of Utah. It is important to develop and test the standard SCF algorithm because this algorithm has features that stress both the computing power of the computational nodes as well as the speed and capacity of the disk input/output (I/O). Another important characteristic of parallel computer systems is the communications bandwidth. Molecular dynamics software stress this component of the machine. These and other codes will form a basis for evaluating competing parallel computer architectures for the MSRC's high-performance computer systems.

Standard SCF implementations have, to date, shown that repetitive read of the two-electron integrals from disk storage devices to form the Fock Matrix is inefficient. Improvements to increase the disk I/O bandwidth include the use of vendor-supplied raw I/O routines that gain a factor of 2 to 4 in performance. Without resorting to specially-written device drivers, this is the maximum disk throughput available from machines like the iPSC/860. However, this throughput is insufficient to keep up with the computations; in addition, it does not scale with the number of computational nodes available. Considering the limited disk I/O capabilities of today's scalable parallel supercomputers, direct techniques (which eliminate disk I/O at the expense of increased numeric operations) are an attractive alternative. In addition, the large primary memories of parallel computers offer opportunities for performance optimization. We have implemented hybrid methods that read as much of the integral list as possible into the aggregate memory of the machine. Then only the remaining portion of the integral list is processed from disk. These hybrid methods can

produce significant performance improvement, and show promise for efficient implementations of quantum mechanical methods where direct techniques are too costly.

Parallelization of the Electronic Structure Code DISCO

We have parallelized the computationally intensive sections of the *ab initio* Direct-SCF program DISCO for the Intel Touchstone Delta computer system. Initial results indicate the performance level of a single processor Cray X-MP can be achieved with about sixteen nodes of the Delta machine. Using all 512 nodes, we have achieved a throughput 250% greater than an eight processor Cray Y-MP/8-128. On 512 nodes, we obtained a speedup of 300 for computing HF energies and 501 for computing the HF atomic forces over the time required on a single node.

Trial calculations were then carried out on the molecule imidazole ($C_3N_2H_4$) using the highly accurate cc-pvdz, aug-cc-pvdz, and cc-pvtz correlation consistent basis sets. Figure 4.1 shows the speedup curves for the three bases, the ideal limit using the master/slave parallel topology (where speedup is equal to the number of slaves), and the speedup based on Amdahl's law:

$$Speedup = \frac{Time(1\text{ node})}{Time(N\text{ nodes})} = \frac{T_s + T_p}{T_s + \frac{T_p}{N}}$$

where T_s is the time required for the serial portion of the routine, and T_p is the time required for the parallelizable portion. Our results indicate that the master's task generation time does not completely overlap with the time spent by the slaves computing integrals. This could be removed by either having the master assign larger tasks, or having a sublevel of masters, each assigning tasks to their own pool of slaves. Currently, overall performance is also limited by bottle necks of serial code, which are computationally insignificant on a single processor machine.

Molecular Dynamics on Massively Parallel Computers

It is clear from the current trends in the computational sciences that, in the future, most gains in processing power will be delivered by massively parallel computing environments. This is particularly evident in the

Figure 4.1. Speedup curves for computing HF energies on the Intel Touchstone Delta.

field of computational chemistry, where computer resources have limited the scope of the problems that can be addressed. The increasing importance of hazardous waste cleanup underscores the need to model large systems important in environmental chemistry. Key among the systems needing study are enzymes important in bioremediation of organic pollutants. For example, Ornstein and coworkers, studying cytochrome P450cam, have demonstrated that rational predictions of specific point mutations near the heme active site can suggest genetic alterations to obtain modified substrate specificity and activity. These model studies have been carried out with traditional molecular mechanics (MM) force field molecular dynamics.

Currently, MM molecular dynamics simulations of biopolymers are typically limited to systems of fewer than 10,000 atoms and simulation times of a few nanoseconds, at most. However, to fully understand the chemistry of the enzyme active site and the effects of

genetic mutations, we need to treat the electrons and nuclei near the active site quantum mechanically. Furthermore, we must examine more realistic MM potentials, especially for the electrostatic interactions. We also need to routinely include solvent molecules and extend simulations from the nanosecond to microsecond time regimes. Clearly, to perform such simulations on systems relevant to bioremediation will require an increase in performance of several orders of magnitude over present day techniques. These requirements suggest the importance of a hybrid quantum mechanical/molecular mechanical (QM/MM)-molecular dynamics method for massively parallel computers.

This project seeks to develop both MM molecular dynamics and QM/MM-molecular dynamics applications to run on massively parallel computers, specifically the newly installed 512 node Intel Touchstone Delta at Caltech. The most computationally demanding portions of an MM-molecular dynamics are the nonbonded interactions, which scale as $O(N^2)$. There are several ways one can map the solution of the non-bonded interactions onto a parallel architecture, such as the Intel Delta. As a test of the computation and communication abilities of the Delta, we completed prototype versions of MM-molecular dynamics kernels, including replicated data, systolic loop, and master-slave topologies. The results of these tests guided the design of the subsequent production MM-molecular dynamics version. The initial version of the production code uses a replicated data method and is currently undergoing its first performance evaluations. Extensions to spatial decomposition methods are planned.

The QM/MM-molecular dynamics method divides the protein into classical and quantum mechanical motifs. The former, representing the bulk of the protein, is treated with traditional MM methods, while the dynamics of the quantum mechanical region is treated with forces derived from a quantum mechanical Hamiltonian. A proper description of the interaction of the quantum mechanical and classical motifs remains a problem of much interest; however, one possible solution has been given.⁽³⁾ We require a semi-empirical model Hamiltonian that is capable of describing transition metal systems, including open shell, as well as electronic excited states. Initially, our work will focus on the intermediate neglect of differential overlap spectroscopic (INDO/S) model.⁽⁴⁾ Progress has been made in improving the ability of this method

to accurately calculate forces, as well as extending it to the neglect of diatomic differential overlap (NDDO) level, where diatomic anisotropies are better described. This work involves a collaboration with Professor D. Edwards at the University of Idaho.

Joint ANL/PNL Parallel Chemical Physics Workshop

The Theoretical Chemistry Group at Argonne National Laboratory and the Theory, Modeling, and Simulation Program in the MSRC jointly organized the "Parallel Computing in Chemical Physics Workshop" held in Willowbrook, Illinois, July 17-19, 1991. The workshop included 29 talks covering a wide range of topics, including parallel electronic structure, parallel molecular dynamics, parallel languages, tools and architectures, and distributed computing. In addition to the papers, discussion groups considered current issues impacting the advancement of the field in each of these areas. The workshop was attended by 41 chemists and computer scientists from six countries. Many of the papers presented at the workshop will be published in an upcoming proceedings issue of *Theoretica Chimica Acta*.

Benchmark Computations

One goal of the computational chemistry user interface project is to provide utilities to assist people in making complex decisions regarding the large number of input parameters that control how the chemical modeling application will work. Because the amount of computer time, high-speed memory, and disk resources vary enormously with the choice of parameters it is vital that scientists have access to detailed information about the performance of different computational models, so they can make informed decisions concerning the efficient use of computer resources while still obtaining results with sufficient accuracy to solve the problem at hand. While part of the database of information needed to document the performance of various computational models can be obtained from the chemical literature, certain important pieces of information are missing and must come from our own benchmark calculations. This will be the case for computational methods that are being developed in the MSRC and for certain high accuracy results on systems that are of interest to the environmental chemistry research being undertaken at PNL.

Methods to Estimate the Full Configuration Interaction Limit

The full configuration interaction (FCI) limit represents the exact solution of the molecular Schrödinger equation for any given one-particle basis set. Since the development of *ab initio* quantum chemistry following World War II, FCI calculations were considered intractable for any but the most trivial system. However, in recent years, much effort has been expended in developing new, more efficient methods for computing FCI wavefunctions. In concert with the rapid improvements in computer hardware, it is now possible to perform FCI's with dimensions in the hundreds of millions, up to even a billion determinants. Nonetheless, the factorial growth in the size of the problem quickly overcomes even the best algorithm and the calculation becomes prohibitive.

Using a systematic sequence of multireference singles and doubles configuration interaction (CI) wavefunctions, it has been shown that sufficient regularity exists in the total energy and selected one electron properties to be able to efficiently extrapolate to the FCI limit. For a number of molecular systems, an exponential function of the form $f(\sum c_i^2) = a + b \cdot \exp[c \cdot (1 - \sum c_i^2)]$, where $\sum c_i^2$ is the sum of the squares of the reference configuration coefficients in the CI wavefunctions, can be used to extrapolate to the FCI energy. For example, this function extrapolates to the FCI energy of N_2 to an accuracy of better than 1 millihartree (Figure 4.2). The actual FCI involved 111 million configuration state functions (CSFs), while the extrapolation procedure used a maximum of 39,000. Tests with the multireference averaged coupled pair functional method and quasidegenerate variational perturbation theory suggest that they too may converge smoothly enough to permit extrapolation.

Ab Initio Studies of Hydrogen Bonding

Hydrogen-bonding is important in the environment. To calibrate the new correlation consistent⁽⁴⁾ basis sets at the SCF and correlated levels of theory, a series of calculations were carried out on the water dimer. The largest set totalled over 400 functions for the dimer (and included up to h-functions on oxygen and g-functions on hydrogen). An analysis of the convergence of the binding energy at the SCF and MP2 level resulted in an estimated complete basis set limit of $\Delta E_{SCF} = -3.55$ kcal/mol and $\Delta E_{MP2} = -4.95$ kcal/mol.

Figure 4.2. Convergence of the total energy as a function of the size of the reference space for N_2 . The FCI corresponds to $\Sigma c_i^2 = 1.0$ and contains 111 million CSFs.

The effects of higher order correlation treatments were examined and found to be only on the order of -0.1 kcal/mol. Thus, the total correlation contribution to ΔE was estimated to be -1.5 kcal/mol. The counterpoise method proposed by Boyes and Bernardi⁽⁵⁾ as a way of estimating the basis set superposition error was found to be effective when used with normal energy-optimized basis sets, but could actually worsen agreement with the estimated limit when used on diffuse function augmented sets at the MP2 level (Figure 4.3). These findings suggest that a modest sized basis set (e.g., the aug-cc-pvdz basis) is a good candidate for use in larger molecules.

Ab Initio Study of the Nonadditive Effects in the Trimer of Ammonia

Intermolecular interactions in atomic and molecular aggregates can depart strongly from pairwise additivity.

Substantial effort has been expended in the last decade to characterize the pairwise interaction. Much less is known about the underlying principles governing the three-body correction to the pair potential. For the rare gases, extensive studies of nonadditive effects have been made in the past, but similar studies concerning the interactions of molecular species are scarce. However, it is clear from previous work that the three-body effect does not originate from a single source. Instead, it is composed of a number of contributions, each with differing physical origin, properties, and behavior relative to variations of the intermolecular geometry.

The goal of this work was to determine the origin and importance of the three-body effect in $(NH_3)_3$. The study was carried out within the framework of intermolecular Møller-Plesset perturbation theory (IMPPT) combined with supermolecular Møller-Plesset perturbation theory (SMPPT). This treatment allows decomposition of the interaction energy into its physically meaningful components. Our work focused on the partitioning of the two-body and the nonadditive three-body effects to components, including electrostatic, exchange, induction (deformation), dispersion, and their respective intrasystem correlation corrections.

The nonadditive interactions in ammonia are significant. In the cyclic trimer, the nonadditive effect is attractive. It is largely due to SCF deformation nonadditivity. However, its anisotropy closely resembles that of the Heitler-London exchange nonadditivity. In the ammonia trimer, this nonadditivity appears to play a more important role than in other polar trimers studied previously. The degree by which the SCF-deformation effect contributes to the entire nonadditivity seems to correlate with proton-donor abilities. In good proton donors (HF and HCl), the nonadditivity is dominated by the SCF deformation. In less efficient proton donors (H_2O and NH_3), the nonadditivity is a mix of two effects dependent on the geometry and strength of the proton donor. In the double donor and double acceptor configurations of the ammonia trimer, the nonadditive effect is repulsive. Since, in solid ammonia, each NH_3 serves as an acceptor and as a donor to three molecules, the overall nonadditivity in the solid is expected to be repulsive.

Figure 4.3. Convergence of the water dimer binding energy as a function of the basis set size. The square symbols represent the original cc-pVxZ basis sets. The circles are the cc-pVxZ results with the counterpoise correction, and the pluses and x's are the corresponding values for the augmented basis sets.

References

1. Chem3D V3.0, Cambridge Scientific Computing, Inc., Cambridge, Massachusetts, 1990.
2. T. H. Dunning, Jr., *J. Chem. Phys.* **90**, 1007 (1989).
3. M. J. Field, P. A. Bash, and M. Karplus, *J. Comp. Chem.* **11**, 700 (1990).
4. J. Ridley, and M. Zerner, *Theo. Chem.* **32**, 111 (1973).
5. S. F. Boys, and F. Bernardi, *Mol. Phys.* **19**, 553 (1970).

Computing and Information Sciences

Computing and Information Sciences

Advances in computer technology offer opportunities to address many problems in the scientific realm and provide advanced computational capabilities to the scientists in the MSRC. However, taking advantage of these opportunities requires research into the computational infrastructure for a scientific computing environment. The Computing and Information Sciences program is designed to perform applied research related to the design and development of a computing environment to support the MSRC scientists and to design and implement the computing infrastructure for the MSRC and the EMSL.

The scientific computing environment requirements involve preconceptual design and applied research in many areas: network design, distributed computing systems, scientific data management, graphics and visualization, high performance computer architectures, programmability and utilization of massively parallel computing, and electronic circuits and computer interfaces for research instruments. The Computing and Information Sciences research programs can be categorized into the three areas.

Scientific Computing Environment Infrastructure. The scientific computing environment involves developing an integrated computing approach for solving scientific problems and performing scientific research. In order to provide an efficient and easy-to-use environment, research and proof of concept developments are necessary to provide

- a network design that 1) meets the high-volume demands of the high performance machines, 2) can be reconfigured easily, and 3) can be managed with minimal support
- an efficient automated approach to the management of heterogeneous distributed computing systems.

The objective is to design a scientific computing environment that will provide capabilities that the scientists need and can operate in an easy and efficient manner. This work also involves collaboration with the Theory, Modeling, and Simulation program to develop the MSCF.

Scientific Computing Software. As massively parallel and distributed computer systems have assumed a preeminent role in high performance computing, more emphasis must be placed on methods and tools to support the development of codes that can effectively utilize the performance offered by these machines. One goal of the Computing and Information Sciences program is to develop techniques for evaluating the performance of a scientific code on a parallel computer. These techniques include modeling proposed codes and computers, as well as analyzing performance data from actual implementations. These techniques can be used both to guide code development and to evaluate computer architectures and configurations proposed for acquisition. A second goal is to develop efficient implementations of kernel algorithms used by computational chemistry applications, using the evaluation techniques to guide the development. In all these areas, our position as a member of the CSC gives PNL and the MSRC an edge in prototyping these tools on the most powerful supercomputer available today, the Touchstone DELTA system. Increased performance on massively parallel machines typically leads to larger problems being solved. Visualization can be a powerful tool to analyze the results of scientific analysis and to monitor the progress of experiments. The goal of the visualization area is to understand the unique visualization needs in the molecular science and environmental science disciplines, and to provide proof of concept software that could serve as a basis for building visualization and GUI toolkits. In the scientific data management realm, the goal is to understand the requirements for the collection and storage of information about the scientific data (meta-data) and determine how to store self-describing information with the data files so that there is an easy transfer of data between applications.

Electronic Circuits and Computer Interfaces. An environment in which there exist a large number of diverse instruments, many of which are unique, provides a challenge for the efficient development of the hardware and software required for real-time control of laboratory instrumentation. Understanding the requirements for the interfaces can eventually lead to the development of software toolkits and standard

electronic modules that can be used by the scientists to develop their interfaces. Another goal of this task is to determine how to integrate the experimental scientific instruments into the overall computing environment without adversely affecting the performance of the instruments during an experiment.

The following sections describe the activities during the past year in each of the areas summarized above. These activities have focused on preconceptual design and proof-of-concept development.

1. Computer Systems Engineering

J. O. Schroeder, J. W. Brothers, D. M. DeVaney, D. R. Jones, R. J. Littlefield, and M. J. Peterson

The Computing and Information Sciences team is working in collaboration with user groups to implement a fully integrated computing environment within the MSRC. This environment will support simulation, experimentation, and analysis within the MSRC scientific community and will prototype the computing environment proposed for the EMSL⁽¹⁾. The goals of the Computer Systems Engineering activity are to

- build a computing environment that meets the technical requirements of staff, visiting scientists, and off-site collaborators and users
- provide efficient and seamless management of data and information from simulations, experiments, technical databases and publications
- provide an integrated environment where a single, multifunctional workstation can be used as the standard user-interactive device to access computing services available in the facility.
- accommodate new requirements and a new computer technology through modular expansion without total systems redesign/replacement
- achieve a reasonable cost of development and operation by using commercially acquired equipment and technologies whenever possible.

To reach these goals, it will be necessary to extend the state of the art in the application of information technology to support scientific research.

An important accomplishment of the computer systems engineering team this past year has been to define the architecture of each element in sufficient detail to develop a baseline cost estimate for its implementation. Cost factors used to develop the estimate were established by reviewing currently available technology, and forecasting industry trends in price and performance.

The key elements that will form the MSRC/EMSL computing environment are identified in Figure 1.1. The integrating element of the computing environment is the network. It will consist of a computer-to-computer network used to couple high-speed systems, many subscriber networks that will provide network attachment points for equipment located in science laboratories and offices, and a backbone network for interconnecting subscriber networks and providing high-speed network services to selected resources. Using network technology available today these networks would be implemented using High-Performance Parallel Interconnect (HiPPI @ 0.8-1.6 Gb/s), Ethernet (@ 10 Mb/s), and Fiber Data Distributed Interface (FDDI @ 100 Mb/s) technologies, respectively.

The computer-to-computer network will service the High-Performance Computer System, the Experimental Computer System, the Data Management System, and the Graphics and Visualization Laboratory. Together, these elements will be the MSCF. High-performance computing will be supplied by one or more advanced architecture (parallel) computers. A network of scientific workstations will also be available as a loosely coupled, parallel high-performance "computer." The Experimental Computer System will be a leading-edge, high-performance system that will provide an opportunity for scientists to evaluate new computer architectures in advance of selecting the next generation production machine. The Data Management System will store technical databases, scientific publications (available in electronic format), and interim and final results from simulations and experiments. This system will have both on-line media that can handle at a minimum

Figure 1.1. MSRC/EMSL Computing Environment.

80-100 GB of data and extensive archival storage capability. Advanced graphics workstations and software in the Graphics and Visualization Laboratory will be used for animation, display, and manipulation of molecular models; analysis and display of experimental or computer generated data; and viewing, in real-time, the progress of simulation and analysis codes running on the high-performance computer.

Additional elements of the computing infrastructure will be the Interface Development Laboratory and the Distributed Computing Systems. The Interface Development Laboratory will support laboratory automation activities by providing the tools and environment necessary to develop electronic interfaces and real-time control and data acquisition software for laboratory instrumentation. A client-server computing model will be adopted for the Distributed Computing Systems element of the architecture with extensive use of server resources to lower the cost of implementing and managing user workstation resources. The servers and workstations

together will provide an appropriate balance of performance for office applications, data visualization, numerical computations, and code development.

The EMSL will include specialized facilities for conferences and user training. Multimedia workstations and projections equipment will be used in these facilities to enhance the education and communication of technical and scientific topics.

We expect that the computer industry will supply many of the required technologies and products, but that a significant challenge will lie in integrating them into a cohesive computing environment. Where our requirements exceed current capabilities (e.g., in the area of high performance computing and scientific data management), we will push as far to the edge of technology as we can and still provide a useable and manageable system. Where development is required, we will work with collaborative partners in industry, academia, and government research organizations.

Our technical approach to implementing the system will couple a traditional engineering approach with prototyping activities, thereby controlling the risk associated with the project. It is important that the elements of the architecture be delivered to users in increments that do not exceed our understanding of the problem. In this manner the implementors of the architecture will have the benefit of seeing their creation as a functioning entity, rather than a paper model, and will be able to alter course as needed to engineer a better solution to users' computing needs.

Reference

1. Computing and Information Sciences Preliminary Engineering Design Study, PNL-7670, April 1990.

2. Distributed Computing Systems

**J. O. Schroeder, W. S. Eggers, and
D. R. McGee**

A major accomplishment in the past year was the development and documentation of the architecture that will be used to extend MSRC distributed computing capabilities. Distributed computing systems include the servers, peripherals, and workstations that bring computing capabilities to the user's office and laboratory work areas. Distributed computing capability has been available in the MSRC since its inception, and is now poised to grow in size and capability as new staff are added and we systematically develop prototypes of distributed computing strategies before transferring them to the EMSL.

Various organizations are working on software to manage and operate distributed computing systems, most notably the Open Software Foundation (OSF) through its Distributed Computing Environment (DCE) and Distributed Management Environment (DME) initiatives^(1,2) and UNIX International (UI) through its efforts to define the Atlas architecture for distributed computing⁽³⁾. These technologies are not yet ready for deployment, and it is not clear at this time which architecture offers the best solution for our needs or which will be delivered in a timely and robust fashion. However, it is clear that industry is attacking the problem and will eventually deliver solutions. The distributed computing architecture

developed for the MSRC is an interim solution that will accommodate production use of computing resources and allow graceful adoption of commercial distributed computing technologies as they become available in 1 to 2 years.

From a hardware perspective, the MSRC computing environment will be divided into sections, each section serving up to approximately 100 clients. A network router device serving each section will attach subscriber networks and local FDDI networks to a backbone network that interconnects all computing resources. This concept is illustrated in Figure 2.1. There will be several network segments in each section. From 6 to 25 computer resources will connect to a subscriber network, depending on the type and utilization of the resources. These resources will form a workgroup, which is defined as a set of equipment that provides computing services to an association of staff sharing a common work interest.

The collective set of work groups served by one router is called a net group. Each net group will have one multifunction server computer that provides a standard set of services to clients. These services will include file and software management, backup, mail and messaging, name resolution, and printing. This approach has advantages with regard to licensing and maintenance of application software, backup of user files, and sharing of data and code. Local disk storage would be used by the operating system, and to store temporary scratch files and the file cache used by the distributed file system.

Additional services provided to all net groups will include authentication, electronic bulletin board access, and system administration. Centralization of critical services will promote more efficient and consistent administration of the computing environment.

In the longer term, we expect that industry standardization efforts will yield a common suite of software for management of heterogeneous workstation equipment. In the near term, we will provide a basic set of software and procedures that allow efficient management of a modest set of workstation equipment. Development of these tools will build on solutions already available and operational within PNL.

Figure 2.1. MSRC Distributed Computing Systems.

Many of the above concepts are being evaluated in the Distributed Computing Systems Testbed. Favorable test results will lead to Beta testing with a small group of users, followed by full deployment to the MSRC computing environment.

References

1. OSF Distributed Computing Environment Rationale, Open Software Foundation, May 1990.
2. OSF Distributed Management Environment Rationale, Open Software Foundation, September 1991.
3. UI-Atlas Distributed Computing Architecture: A Technical Overview, UNIX International, September 1991.

3. Distributed Computing Systems Testbed

T. L. Harper and J. O. Schroeder

The mission of the Distributed Computing Systems (DCS) Testbed is to provide a controlled environment in which distributed hardware and software components can be configured, tested, and evaluated. Specific near term activities of the Testbed are directed toward validating the proposed strategy for enhancing MSRC distributed computing system capabilities in the next 12 to 15 months^(1,2). In the longer term, the testbed will have a major influence on the design of the EMSL distributed computing environment. By using the testbed to evaluate new technologies and explore migration paths from current configurations, new technology can be incorporated while maintaining the high reliability and usability scientists have come to expect from their computational resources.

Capabilities

In the testbed environment, the impact and applicability of new technology can be accurately assessed. Configuration testing can be carried out to determine system permutations that maximize performance. The testbed can also be used to generate performance metrics. This information is used by system designers to evaluate competing vendor products and technologies and to formulate a DCS architecture.

Hardware platforms, including workstations and servers from multiple vendors, have been assembled in the testbed. All of the systems are configured with hardware and software to connect to an isolated Ethernet network. Some of the systems have FDDI network capability, which is the emerging optical networking standard. Network components are available to support AUI, ThinNet, and 10Base-T Ethernet media. In addition, a high-speed network router provides an interconnection between Ethernet and FDDI subnets. Test equipment includes a network protocol analyzer to monitor experiments, induce network traffic, and diagnose and isolate system faults and bottlenecks.

Many of the testbed activities are coordinated with other research institutions. For example, ideas and testing methods are being exchanged with the National Center for Supercomputer Applications (NCSA) and Carnegie Mellon University. The level of cooperation with outside institutions started very modestly but continues to grow. Our willingness to share results obtained through the testbed is the key factor in the development of this cooperation.

The effectiveness of the testbed is greatly enhanced because of a similar project within the Information Systems and Services (IS&S) directorate at PNL. The joint contribution of equipment, software, and labor to the Testbed by IS&S and MSRC increases the productivity of the effort.

Current Activities

The testbed has been operational since August 1991⁽³⁾. Currently, tests are under way to evaluate workstation I/O characteristics and measure Ethernet and FDDI performance. Specific tests include memory-to-memory and disk-to-disk performance between workstations, available network bandwidth as a function of active nodes, and measurement of

network latency. Preliminary results from these tests are available and will be used to guide 1992 procurement strategies and product evaluations. This information is valuable because it allows many low-cost iterations to be made in the design cycle before committing to a implementation strategy.

Benchmarks are being developed to quantify the performance of distributed file systems (DFS). The DFS is an important technology because it allows sharing of data resources in a controllable and manageable fashion for a large organization. A DFS can provide services such as application serving, software license control, load balancing, and centralized management of distributed data. The DFS benchmarking is designed to measure the systems ability to transfer and manage files in a hierarchical directory structure. The emphasis of the testing is on transmission speed and system throughput. The benchmark results are being examined in conjunction with an assessment of features that are not directly measurable, such as ease of use and manageability.

Network configuration alternatives are being explored to determine optimal server location and workstation configurations. These tests provide information on actual observed performance under conditions representative of a production computing environment. Using a router, a combination of Ethernet and FDDI networks, network monitors, and platforms from multiple vendors, realistic situations can be modeled.

Results of these and other tests will be documented and made available to system designers and collaborative partners. A controlled test environment will always be a requirement of the research laboratory for the evaluation of new technology. Technology continues to evolve rapidly and PNL and the MSRC must be prepared to evaluate the potential afforded by currently unforeseen advances in computing.

References

1. EMSL Interim Facility Proposed Computing Environment, June 17, 1991.
2. EMSL Interim Facility Distributed Systems Computing Environment Implementation Plan, June 25, 1991.

3. Task Execution Plan, Evaluation of the Andrew File System, July 22, 1991.

4. Scientific Networking

M. J. Peterson and T. A. Harper

Distributed, high-performance scientific computing places challenging demands on current networking principles. Most current network products and design methodologies are well suited for more predictable business environments of batch processing, electronic mail, file transfers, and printing. The proliferation of large amounts of scientific and theoretical data throughout MSRC and the requirement for moving data among different computing architectures has necessitated a reconsideration of the way in which networks are designed and implemented.

The MSRC networking architecture has been grouped into three general classes of service. The first of these serve the resources that exist at the desktop, and it is currently planned to run 10Mbits/s Ethernet to all but the most demanding desktop computers. These end-user subscriber networks (ESN) are unique in that they will be designed to serve few users, typically between 10 to 30 users per physical subnet. Through partitioning less users per subnet, more bandwidth will be available for each individual user. The flexibility and cost effectiveness of Ethernet make it the viable choice to the desktop for the near term, with newer developing network technologies as candidates for the future.

These ESN subnets will be interconnected with high performance routers to a 100Mbits/s FDDI backbone. This backbone information network (BIN) will interconnect and provide the necessary integration of the subnets (Figure 4.1). Aside from this interconnection service, the BIN technologies will also provide the mechanism to network high performance Workstations at the desktop and shared servers. The BIN may consist of multiple FDDI networks to serve the Ethernet backbone connectivity and the FDDI connected high performance computing needs. Prototyping is currently taking place to help determine the practical limits of utilizing this new high-speed network, and to implement the most optimum configuration.

Certain classes of computation resources require extremely high network bandwidths that are not currently available from Ethernet or FDDI technologies. These high-speed computer-computer networks (C2N) require transfer rates approaching 0.5 to 1.0 Gbits/s. The high-speed parallel processors, graphics/visualization systems, and large data retrieval systems envisioned will utilize this specialized network. Technologies being investigated for this include HIPPI, FiberChannel, and Gbits switching technologies.

All of these networking technologies will require a very tightly integrated network management system to allow effective operation and troubleshooting of the systems. Managing many subnets within a building will require an automated approach. This network management system must include a tool for predicting network traffic bottlenecks and modeling future demands. This integration of current status with future predicted loads will help to expand and reconfigure the MSRC network before problems arise.

The initial prototype installation of ESNs has been designed, and will install 10BASE-T twisted pair Ethernet in the Math building at PNL. The network will be divided into four subnets, with the capability to reconfigure up to eight subnets. These subnets will group users into logical as well as physical networks and will interconnect them with high-speed routers linked by FDDI. The physical plant wiring is designed into a hierarchy of patch panels that allow reconfiguration "on the fly" by simply moving jumper cables. Wiring hub locations are interconnected with a cabling scheme that allows users in the building to be added to any of the physical subnets, depending on their particular requirements. As these requirements evolve and change, users can be regrouped as their needs dictate. This scheme provides a very flexible means of segregating users and also maintains a high level of performance by allowing fewer users to share the Ethernet's limited bandwidth.

Other prototyping efforts involve placement of the network server resources: file storage, printing, applications, network services, routers, and mail. These resources are typically shared by all users, and their placement on the network is critical to maintain a high level of performance. It is preferred to have

Backbone Network

Figure 4.1. Backbone Information Network.

the server resources directly attached to each subnet, but costs may prohibit this configuration. Shared placement on the higher speed BIN FDDI backbone may provide a cost-effective solution with acceptable performance to the users. These different architectures are being prototyped, with performance measurements being conducted to determine acceptable levels of performance. A teamwork approach exists with the other task leaders from the PNL Computer Sciences Department, especially the Distributed Computing Environment that is evaluating computer/network operating systems, file systems, and operations issues dealing with systems management.

The challenges that lie ahead for the MSRC computing network can have a positive impact on the ease in which scientists can apply computational resources to molecular research.

5. Data Storage and Archival

D. M. DeVaney

The data storage and archival approach for the MSRC is based on the requirements of the theoretical and experimental chemists. A database will manage a relatively small volume of information (metadata) about projects, experiments, research, and high-level, summary, and final results. As shown in Figure 5.1, database entries will point to files of detailed scientific data. The volume of these detail files will require massive [an estimated 8 terabytes (8×10^{12} bytes) of data by the late 1990s] and inexpensive storage. The size of some individual files (to tens of gigabytes for some experimental results from the ICR and the gigahertz NMR) and the supercomputer performance level of some of the modeling, analysis, and scientific visualization systems

Figure 5.1. MSRC Database and Archive System Interactions.

requires high-performance storage devices and networks.

We established a 32-gigahertz file from a 3-D experiment on the gigahertz NMR as our benchmark "very large file" for the MSRC. A very large file is characterized as a file that is transmitted over the network (e.g., from an instrument) for long-term storage on the archival system and then is occasionally retransmitted over the network (e.g., to the graphics and visualization laboratory). A HiPPI-class network operating continuously at 100 megabytes per second could transfer a 32-gigabyte file in 5 minutes. However, current transfer rates for peripheral subsystems cannot support a file of that

size. Current striped magnetic disk subsystems optimally support about half this bandwidth, and in practice, a realistic net figure is about one-fourth of that bandwidth, implying a 20 minute net transfer time for this benchmark file over the fastest currently available computer to the computer network. If our benchmark file is staged to unstriped magnetic disk, we expect a sustained rate of transfer of around 4 megabytes/second. This would optimally require about 2.25 hours for the transfer. A real-world transfer probably takes twice that time -- about 4.5 hours. If the file had to be staged from optical disk or narrow (4 mm and 8 mm) cartridge tape, we can expect transfer rates from 4 to 10 times slower than for unstriped magnetic disk, implying 10 to

20 hours optimally for the transfer, with real-world net transfer times probably twice that, (e.g., 20 to 40 hours).

Storage subsystems must be considerably larger to gracefully absorb such a very large file at unscheduled times. For example, using near-term, 11-inch form factor optical technology, this file requires three 10.5-gigabyte platters. Jukeboxes holding up to 50 of these platters are available today. Using near-term, 5.25-inch form factor optical technology, this file requires about 32 1.0-gigabyte platters, or roughly one-third the capacity of a 100 platter jukebox, which is the biggest jukebox currently available. The number of 5.25-inch platters alone presents a problem when the file must be staged from off-line status, as from a platter vault, to physically being mounted on the jukebox. In near-term, 8-mm technology, our benchmark file requires six or seven tape cartridges, which amounts to about one-eighth the capacity of a 50-cartridge jukebox or one-sixteenth the capacity of a 100-cartridge jukebox.

In 1991, we investigated commercial and scientific storage-system products and practices where the total storage requirement is measured in the range of approximately one to several tens of terabytes. These sites are beginning to move from a reliance on large conventional disk farms backed by IBM 3480 tape cartridge technology to newer technologies, including redundant arrays of inexpensive disks (RAID) and 19-mm tape cartridge systems. In these environments, we found UniTREE® software is the defacto standard for managing and staging large files and large collections of files over a hierarchy of peripherals. UniTREE was initially developed as a consolidated file system to satisfy a commercially unmet requirement in the DOE supercomputer community. Our approach is to competitively acquire a commercial UniTREE-like data storage and archival management system.

In the UniTREE model, several levels of hierarchical storage may be defined. Successive layers are further from the user (the computational nodes), are slower, have greater capacity, and use less-expensive media (on a per-byte basis). As shown in Figure 5.2, we have defined four storage levels for MSRC. Level-1 storage is (typically) very-fast-disk storage connected to end-user and computational nodes and is acquired

as part of those nodes. Level-2 storage is up to several hundred gigabytes of network-connected, high-performance (tens to hundreds of megabytes per second transfer rate) striped disk. Level-3 storage is up to several terabytes of medium bandwidth (5 to 50 megabytes per second transfer rate) storage. Level-4 storage is long-shelf-life, very-high-capacity, moderate-speed (e.g., 3 to 10 megabytes per second), off-line (e.g., human tape vault access or human tape-mount required) archival storage.

Optical media is preferred for archival purposes because of its long shelf-life, but transfer rates for optical media of all kinds limit use to Level-4 storage. Media striping techniques could provide almost a tenfold gain in effective transfer rate in the near term (2 to 3 years), and this increase would make optical and tape systems viable for Level-3 storage. Currently, no tape or optical systems use striping.

The IEEE has defined a Mass Storage Reference Model⁽¹⁾ that calls for high-performance, Level-2 peripherals to be direct-connected to high-performance (e.g., HiPPI) networks without a bottleneck server machine as a required intermediary. Instead, a peripheral management and control node is connected with client nodes and the network peripheral through a low-speed (e.g., EtherNet) "backdoor" network. The goal is to reduce costs by eliminating the need for a high-performance server machine without sacrificing performance, indeed while simultaneously enhancing performance. This model is being tested and prototyped at some supercomputer sites (11th IEEE Symposium on Mass Storage Systems, October 1991) and is reportedly proving difficult to implement. Nevertheless, this is the preferred alternative and is the MSRC goal architecture. Competitively available peripheral systems that can support the performance requirements of all four MSRC storage hierarchy levels are emerging.

Reference

1. S. Coleman, and S. Miller, *Mass Storage System Reference Model: Version 4*, developed by the IEEE Technical Committee on Mass Storage Systems and Technology, May 1990.

Figure 5.2. MSRC Data Management System.

6. Database Management Systems

D. M. DeVaney

In 1991, we studied database management systems (DBMS) to determine which, if any, commercial products will best support the needs of the MSRC. Based on our studies thus far, it appears that a new technology, object database management systems (ODBMS), is the best choice for MSRC and have identified at least two acceptable commercial offerings of this technology⁽¹⁾.

During the past three decades, software technology for supporting the development of data-intensive applications has undergone four generations of

evolution, beginning with file systems, hierarchical DBMS, network DBMS, and then relational DBMS.

Conventional (i.e., relational and prerelational) DBMS served the needs of business data processing applications. However, as relational database technology entered the commercial market, a variety of applications began to be identified that are difficult to implement using relational DBMS. These applications include multimedia systems that manage images, graphics, voice, and textual documents; computer-aided design (CAD), engineering (CAE), software engineering (CASE), and manufacturing (CAM) systems; and, of particular interest to the MSRC, scientific modeling and analysis applications.

These applications require facilities for modeling and managing complex nested entities (e.g., engineering and scientific data types and compound data sets); a richer set of data types (e.g., user-defined data types) and long, unstructured data (e.g., 3-D images, audio, and complex textual documents, such as laboratory notebooks); frequently useful semantic concepts (e.g., generalization and aggregation relationships between data); and the concept of temporal evolution of data (i.e., time-dimensioned data and versioned data sets). In addition, scientific modeling and analysis applications are often highly compute-intensive on a large volume of memory-resident data, and impose performance demands that cannot be met by relational and prerelational DBMS. Scientific and modeling applications also require long-duration, interactive, and cooperative transactions.

The need to reduce the cost of developing and operating these applications pointed to the need for a fundamental advancement in database technology. There are two major reasons why the object-oriented paradigm is a sound basis for a new generation of database technology. First, the object-oriented data model can readily subsume the data models of conventional database systems, can represent data and relationships and constraints on the data, and can provide for encapsulation of data and programs that operate on the data and for a uniform framework for the treatment of arbitrary user-defined data types. Second, the object-oriented paradigm (and thus the object-oriented data model), through the notions of encapsulation and inheritance (reuse), satisfies the objective of reducing the difficulty of designing and evolving very large and complex databases.

We established several general requirements that must be met before an ODBMS can be used as the cornerstone in the development of a scientific data management system for the MSRC. The first ODBMS requirement is sufficient performance and functionality. Second, the vendor must offer ODBMS application development tools that support 1) the entire development process, including code generation, testing, and debugging, and 2) graphical aids for designing database schemas. Our third, and final, requirement is that the ODBMS comply with the few existing and proposed information management standards.

We established an ODBMS product evaluation by translating the three general ODBMS requirements into six well-defined product criteria:

1. performance
2. support for standard programming language(s)
3. high productivity development tools
4. complete DBMS functionality
5. portability and heterogeneity
6. compliance with industry standards.

Industry products will be evaluated against these criteria.

During the year, we supported one of Professor David Maier's graduate students (Judy Cushing) at the Oregon Graduate Institute. Dr. Maier is an expert in the field of ODBMS. Ms. Cushing's research included the design of a research chemistry experimental results ODBMS application, implementing the design on four disparate ODBMS platforms, populating the databases with sample MSRC theoretical chemistry results, and evaluating the results of a series of database queries against these databases⁽²⁾. Ms. Cushing's results confirmed and strengthened our evaluation of ODBMS products.

References

1. S. B. Zdonik, and D. Maier. Readings in Object-Oriented Database Systems, Morgan Kaufmann, San Mateo, California, 1990.
2. J. B. Cushing, D. M. DeVaney, D. Feller, and D. Maier. Database Support for Computational Chemistry, in Proceedings of the DARPA Open Object-Oriented Data Base Workshop II, Texas Instruments, Dallas, Texas, September 1991.

7. Self Describing Data

R. S. Barga and D. M. DeVaney

A primary objective of MSRC data management is to develop a system that both manages the scientific data and information produced and "facilitates" scientific inquiry. Hence, it is essential that *relevant* existing data sets can be easily identified, that flexible

tools for searching to obtain appropriate subsets of the data are available, and that processing of the data can be performed independent of the manner in which it was stored.

In addressing these issues, we investigated standard data formats. A standard data format can generalize access to scientific data so methods used for storing and accessing data are independent of applications and computer architectures. In addition, a standard data format would minimize the fraction of the software development effort devoted to dealing with application-dependent data formats.

During our investigation, we performed a desk review of several standard data formats currently in use in a variety of scientific disciplines. For purposes of evaluation, we established a number of key criteria for an MSRC standard data format:

- It must be resource efficient. A data management system that handles large quantities of data can use a standard data format only if it requires a moderate storage overhead at a reasonable cost in central processing unit (CPU) time.
- It must be intuitive. We anticipate that many MSRC users will write their own software to access, manipulate, and generate data. Using the standard data format must be a relatively simple task.
- It must be consistent and unambiguous. Rules and conventions must be kept to a minimum so there can be no question as to the contents of the data file, and the methods for storing data must be consistent across disparate types of data. Indeed, the format should make the underlying data format transparent to the user.
- It must be flexible and powerful. It must represent a wide variety of data generated by computational and experimental chemistry programs (i.e., scalar, vector, tensor, animation, etc.) and store any and all information that can be used to describe and document those data. Furthermore, it must be extensible as the science changes.

- It must be self-describing. It must be readily interpreted by humans and by computers. This will allow researchers and programmers to readily determine the nature of a data set by inspection and react to it.

From this review, one standard data format stood out. The Network Common Data Format, or netCDF, is a software package that standardizes storage and retrieval of scientific data⁽¹⁾. We obtained netCDF and experimented with it using MSRC data sets. We found netCDF includes a library of "C" and FORTRAN programming interfaces that can be used with widely varying scientific data sets by machines of widely varying architectures. We found that the following features distinguish netCDF from other standard data formats:

- *Standardized Data Access.* The library of netCDF functions actually hides the underlying data format.
- *Self-Describing.* Data sets and the parameters that describe their dimensions are stored along with information about the data and the processing history of a data set.
- *C and FORTRAN Interfaces.* The netCDF functions can be invoked from either C or FORTRAN, and data stored using one language may be retrieved in the other.
- *Machine-Independent.* The format underlying netCDF employs an open standard known as XDR (for external Data Representation) that renders netCDF files machine independent.
- *Portable.* The software has been successfully ported to a broad range of computers, from personal computers to supercomputers.
- *Compact.* The storage cost for using netCDF is small.
- *Extensibility.* NetCDF remains an open standard, with proposed extensions and enhancements managed through the Unidata netCDF standards committee.

- *Availability.* NetCDF software is freely available through Unidata, an atmospheric science research program funded by the National Science Foundation and managed by the National Center for Atmospheric Research (NCAR).

While netCDF will provide many of the features required, it does not provide complete and robust support required for a data management system. A collaboration may be established to extend the software to make it more useful for the needs of MSRC.

Reference

1. R. Rew, and G. Davis. *NetCDF: An Interface for Scientific Data Access*. IEEE Computer Graphics and Applications, July 1990.

8. Scientific Laboratory Notebook

D. M. DeVaney, A. Schur, and M. L. Yim

In 1991, we developed a prototype scientific information integration and management tool for the desktop. This system is the Electronic Assistant for Scientific Investigation (EASI)⁽¹⁾.

The individual scientist collects and manipulates large data sets and integrates, synthesizes, generates, and records information. A key tool in this process is the scientist's desktop computer environment. The means to access, interact, and manipulate information by the scientist is a critical determinant of the performance of the system as a whole. The activities of chemists and their interactions with information were abstracted into a common paradigm that meets the critical requirement of easing information access and retrieval. This paradigm was embodied in EASI, a working prototype that provides a structured formal representation of the scientists' working environment.

Engineering the system for usability was critical. To focus attention on users and place them in a primary developer role, four strategies were employed:

- 1) forming a multidisciplinary system development team, 2) conducting Interactive Process Modeling sessions, 3) prototyping, and 4) capturing users evaluations of the prototype⁽²⁾.

A prototype was rapidly developed and incrementally evaluated and extended to test and refine key requirements from the Interactive Process Modeling sessions^(1,3) where user requirements were identified. The laboratory notebook serves as the primary metaphor around which the capabilities necessary to facilitate the scientific process are built. A first time user of EASI accesses laboratory notebooks through their "office." The office provides active graphic aids as bookshelves, desk, table, books, and notebooks. The chemist can point to any book or notebook to open and arrange them anywhere within the office, and can bring new books into the office. Books may also be opened by accessing pop-up menus. Other functions allow books to be renamed, saved, archived (removed from the visual office environment), and closed.

Once a book is opened, many features are available to the scientist for recording and organizing work in the notebook. EASI contains a number of organizing features. A chemist may place bookmarks and section dividers, and add pages sequentially. The system also offers a number of recording options. Most of these options reside outside the notebook. These options are organized around the notebook to give a standard set of capabilities (e.g., hypertext features, annotation, access to software application tools) regardless where the chemist is performing their work (e.g., office, run monitor, scratch pad). Chemists can also annotate any work contained in the system.

A tool was developed to capture, on-line and in real time, user comments during system evaluation.⁽²⁾ This tool, which is analogous to a paper Post-it Note, allows the user to mark and make comments about the system from its behavior to the display format. The comments are placed where the system behavior occurs and captures the context of that event. It was possible to precisely identify where users had difficulty accessing and manipulating information and to find the cause of the problem. Many times the chemists provided the explanation and made it very clear when their expectations were violated. Interestingly, the prototype became the primary tool used to refine its own requirements and an invaluable vehicle to test and articulate the team's ideas.

Many design principles were used to create EASI. None are new but some appeared to be of critical value in providing the chemist with perceived

complete and instant direct control of information access, storage, manipulation, and exploration. These are briefly described within the context of EASI:

- Two forms of hypertext were used to link and record information. Users can create links between information entities where both are within EASI (hyperlink) and where one is outside of EASI (metalink). Metalink provides the ability to record information by reference, to link to such entities as large data sets, literature search results, networks, application tools, visualization packages, and electronic mail.
- The human visual system's ability to form figure/ground relationships was exploited to reduce the need to arrange information hierarchically. The "flattened" presentation method made it possible to present more needed information at one time. Window and pane borders were used to organize the relationships between the figure/foreground items.
- Many people remember where they put items based on their location in space. The graphical interface enabled the use of spatial cues to facilitate the location of items. The visual cues provided to facilitate recall were based on common practice, for example, the office and notebook. In both cases, the chemist may organize information according to his individual visual preference.
- Multiple pathways and entry points provide access to information easily. Information within the notebook can be accessed using bookmarks and section marks, direct page access, and page "turning." Information linked to external sources can be accessed from lists or directly via icon. Any kind of stored entity (e.g., a software tool, a specific file, etc.) could be assigned an application portal. Pointing on a portal gives direct access to the selected entity.
- The ability to move in a transparent fashion from one kind of transaction to another was provided to give literal and perceived instant access to and selection of desired entities of information, application tools, and computational results. The removal of modal operations enabled the chemist to focus their attention on

science rather than aspects of the computer environment in which they were performing their task.

Three major functions to support the scientific process were provided by EASI: 1) a repository for information selected by the scientist; 2) a unifying means to easily manipulate, explore, store, and access the wide variety of computer programs and information necessary for scientific research and other daily activities; and 3) a reference for producing scientific contributions. The chemist must explore, integrate, synthesize, and generate information. EASI appears to facilitate this process.^(1,3)

References

1. A. Schur, D. F. Feller, D. M. DeVaney, J. J. Thomas, and M. L. Yim. EASI: An Electronic Assistant for Scientific Investigation, Vol. I, in Proceedings of the Human Factors Society 35th Annual Meeting, September 1991.
2. S. Hunt, and A. Schur. The User-Oriented Evaluation Process: A Process for Preserving User Needs During iterative System test and Evaluation, Vol II, in Proceedings of the Human Factors Society 35th Annual Meeting, September 1991.
3. A. Schur (attributed in the agenda to D. M. DeVaney). October 1991. "State of the Art in Electronic Notebooks." The 7th Annual Scientific Computing and Automation Conference and Exposition, September 1991. (Also filmed by CNN and aired on Future Watch.)

9. Visual Science Computing

D. R. Jones and J. J. Thomas

Scientific research, in general and computational and experimental chemistry, in particular, require scientists to interpret large volumes of complex data or to execute complex multistep processes requiring access to diverse information. For instance, a molecular dynamics simulation of a single protein can generate 10^5 sets of position and velocity coordinates for each of 10^4 atoms, along with a variety of ancillary information. Graphical display and animation is

essentially the only way in which these kinds of simulation results can be communicated to the researcher. Execution of an electronic structure calculation on a modestly sized molecule may require use of a series of computer codes, each of which can have tens of thousands of arcanelly described (or even undocumented) inputs. The computer graphics technology required for display of data sets (i.e., visualization) is recognized as a key element of the scientific enterprise.⁽¹⁾ The portion of computer codes that communicate directly with the user for problem definition and setup through exchange of text or graphics information (i.e., the user interface) is also recognized as a crucial productivity tool. In the long run, the distinction between these technologies will blur, as the user interface will be the medium through which visualization technology will be provided. The complexity of this situation will surely continue to grow with time.

New technology will have to be developed in user interface and scientific visualization to provide an efficient and highly productive research environment with close ties between research and application. To date, both offer the promise of significantly improving the productivity of the research process. However, many basic questions remain open:

- What software architecture in user interface technology and frameworks allows for the development and integration of complex user interfaces and visualization? This architecture must include definitions for data and control structures appropriate for, and transferable between, scientific applications and the user interface support system.
- What higher order interaction techniques empower the application specialist to define interactions at the problem domain level rather than at the point-and-click level?
- What methods can be used to automatically generate the first order user interface given the application domain control streams?
- What visualization methods will enhance the understandability of complex scientific information and processes in the molecular and environmental sciences?

- How can we embed the intelligence of the application scientists within the user interface?

Lack of answers to these questions limits the effectiveness with which we use computational resources within the scientific process. Both scientists within MSRC⁽²⁾ and throughout the user community⁽³⁾ are calling for improved methods.

Both research and development issues must be addressed before visualization and user interface can achieve their promise of substantial increases in research efficiency and productivity in the chemical sciences. To design effective technology, it is necessary to understand what information a scientist needs to extract from a particular data set, and what form of graphical representation will be effective in communicating that information. To make it practical to incorporate visualization techniques as a routine component of scientific investigation, it is necessary to have a technology that permits facile display and interactive manipulation of the data. Design of an implementation of effective GUIs has similar prerequisites. It is both necessary to understand the information required by the user at any particular stage in a calculation, analysis, or setup of a computer controlled experiment, and to have an appropriate technology framework available that permits facile construction of an interface capturing the problem's information flow.

A real opportunity lies in the combination of user interface and visualization technologies. First, the user interface must incorporate visualization techniques that allow the scientist to create interfaces and communication within the problem domain. Second, the user interface framework is required to provide sharing and discovery of new visualization techniques across application systems. Integration of user interface and visualization technology is a primary goal of this research effort.

Additionally, MSRC scientists require communication of their research to peers and professional groups. In general, scientists and engineers frequently utilize three types (or stages) of visualization:

1. *Personal graphics.* Personal graphics used during the analysis allows the researcher to interact freely with data and observe the results visually on computer graphics monitors. Personal graphics require accurate data representation and

high interactivity rather than image quality. The emphasis is on data understanding and extraction.

2. *Peer graphics.* Peer graphics are used for effective communication between fellow colleagues in their chosen discipline. The image quality for peer graphics should be "better" than that for the personal graphics. The representation of the data should be clear and reproducible with the emphasis on specific information transfer.
3. *Presentation graphics.* Presentation graphics are used when scientists want to visually portray their information to a wider audience, usually for a formal presentation, conferences, or for articles in professional journals. The higher the quality of the material, the more easily understood and effective the presentation. The representation of the data should be clear, concise and accurate, and the image quality should be the best available. Presentation graphics should emphasize both understanding and transfer of information.

The objective of the Visual Science Computing Program is to provide general purpose visualization tools that will help MSRC scientists understand their large, complex data sets.

References

1. ACM, Visualization in Scientific Computing, B. H. McCormich, T. A. DeVanty, and M. D. Brown, eds., ACM SIGGRAPH, New York, 1987.
2. MSCF, Molecular Science Computing Facility Study Group Report, Chairman: Ed Hayes, Rice University, 1990.
3. National Academy Press, Information Technology and the Conduct of Research. The User's View, Report of the Panel on Information Technology and the Conduct at Research, Washington, D.C., 1989.

10. Graphical User Interface Development

D. R. Jones, G. Chin, and J. J. Thomas

Effective exploitation of visualization techniques in general requires a major programming effort, as no comprehensive framework technology for display techniques is generally available. The formation of the new technology called User Interface Management Systems is providing new hope for such a framework, but this technology is still not usable for complex applications. Efforts to develop this technology are under way at both research and commercial institutions.⁽¹⁻⁶⁾

The process of designing user interfaces for computer programs has been active since the first programs were written. Only within the last 10 years has the importance of user interfaces been realized. This has been due to three phenomena:

1. The fraction of user interface code written as compared to methods code for interactive application now ranges from a staggering 50% to 80%.
2. It has become cost effective to include graphics in the interface.
3. Research scientists are stating that the user interface is a limiting factor in utilization of complex computer programs.

Examples like the Macintosh interface give hope that there is a better form of user interface that will greatly improve the usability of computer programs permitting the investigation of more complex phenomena than are now accessible to either theoretical or experimental treatment.

User Interface Development Tool Evaluation

Three commercial systems, DataViews, UIMX, and Application Visualization System (AVS), were

evaluated on their user interface development capabilities. Developed by the V.I. Corporation, DataViews is marketed as a real-time data display tool that allow developers to interactively attach graphical representations to their data. Through DataViews, developers can display their data in the form of graphs, charts or tables, or they can create new 2-D graph representations of their own. UIMX is a user interface development tool initially introduced by AT&T. This tool allows a user interface developer to interactively construct and layout the elements of the user interface. Once the interface has been defined, UIMX automatically generates the source code that may be compiled and linked with the application code. The UIMX copy that we assessed constructed Motif interfaces, but an Open Look implementation is also available. AVS is quickly becoming a *de facto* standard in the area of scientific visualization. In addition to its graphics capabilities, AVS also has some powerful software integration tools and limited user interface development features.

The objective of these evaluations was to assess the current capabilities and limits of commercial user interface technology. In addition, we hoped to find out how applicable this technology is to the needs of the EMSL. After discussion with cognizant MSRC research staff, we came to the conclusion that these systems are missing critical application definition and integration capabilities. In the area of computational chemistry, one of the most prominent problems in developing user interfaces to existing codes is the lack of tools that aid the user in defining the characteristics and logic of his application. Most user interface development tools concentrate on the look and feel aspect of the user interface component, but provide minimal support for the actual definition and integration of the application components. A logical, follow-on approach would be to investigate current user interface research projects that employ methods for extracting application properties directly into the user interface environment. Examples of such systems are UIDE, Stanford's InterViews, and Carnegie Mellon University's Garnet.

Cooperative Research and Development Agreement (CRADA)

In order to successfully complete the computational chemistry goals in developing a GUI, it would be

necessary for other commercial companies to give support to the effort. Following the direction given in the Workshop on Graphical User Interfaces in Computational Chemistry, which was held in September 1990,⁽¹⁾ we initiated the first MSRC CRADA for GUIs. Both MSRC staff and a commercial vendor started to talk about the issues and requirements for an appropriate GUI. This effort is still ongoing, and a final commercial vendor will be selected for continue work in the future.

User Interfaces to Data Acquisition Systems

MSRC experimentalists have markedly different user interface requirements than MSRC theorists and computational chemists. Experimentalists interact directly with physical devices such as pulse lasers and mass spectrometers, and employ computers primarily to capture, display, and analyze resultant data. Computer scientists in the MSRC are working closely with the experimentalists in an effort to understand and support the experimentalists' basic data acquisition and user interface needs. Initially, data acquisition systems with GUIs will be designed and developed for selected pieces of equipment. The user interfaces will provide the experimentalists with a variety of techniques to use in displaying and analyzing their results. Interaction with the experimentalists will provide computer scientists with valuable feedback in the design of GUIs for data acquisition systems.

A subsequent goal in constructing these user interfaces is to develop a basic set or library of data acquisition and user interface modules from which experimentalists may pick and choose. Furthermore, computer scientists wish to develop interactive layout editors from which experimentalists can construct their own data acquisition systems with minimal programming. Such layout editors will allow experimentalists to select data display techniques from tool palettes and spatially organize these tools on the user screen. Currently, most experimentalists construct their own data acquisition systems from scratch. Layout editors and the module library will automate the development of data acquisition systems and remove much of the burden of developing data gathering and user interface code from the experimentalists.

References

1. *Graphical User Interfaces in Computational Chemistry, Workshop Proceedings*, PNL-SA-18847, Pacific Northwest Laboratory, Richland, Washington.
2. G. Pfaff, ed., *User Interface Management Systems*, Springer-Verlag, Berlin, 1985.
3. H. R. Hartson, and D. Hix, *Human-Computer Interface Development: Concepts and Systems*, Vol. 21, Num. 1, ACM Computing Surveys, March 1989.
4. "ACM SIGGRAPH Workshop on Software Tools for User Interface Management" (held at Battelle Seattle) ACM New York. *Computer Graphics* **21**, 2 (Apr.), 71-147 (1986).
5. *ACM SIGGRAPH Symposium on User Interface Software*, (Baniff, Alberta, Canada) ACM New York, 1988.
6. *UIST, User Interface Software and Technology*, ACM Press, Williamsburg, Virginia, 1989.

11. Software Development for Visual Tools

D. R. Jones, E. N. Thornton, and
J. J. Thomas

The goal of visualization is to leverage existing scientific methods by providing new scientific insight through visual methods. Advances in visualization techniques in recent years have been promising.⁽¹⁾ However, several key issues are limiting the broad use of visualization technology. First, the visualization technology developed to date is mostly application dependent since the methods are directly implemented within specific application programs. This means that to apply these methods to other applications requires enormous recoding. Also, we are now just beginning to discover the significance of 3-D visualization. Many significant scientific data sets are of a higher dimensionality.

To support the use of general visualization tools in the MSRC, preliminary studies were made of the research needs and requirements. This initial

investigation established the basis for choosing a general purpose visualization software package. After investigating commercially available systems, it was determined that AVS would be the most likely candidate.

Along with the software, an area of interest was identified for a test program for software development. In collaboration with other MSRC scientists, it was determined that computer graphics work needed to be done in the area of molecular modeling of crystal structures. Scientists have developed computer codes to mathematically model crystals, which could be done with very little input because of the large amount of symmetry in crystals. However, exploring the results of large crystal structures was quite tedious when presented in numerical format. In response to this problem, a software package called Space Crystal was developed in the AVS environment.

Space Crystal is a software package used to graphically build, model, analyze, and visualize crystalline molecular structures. The software uses the original computer code written by MSRC scientists to mathematically build crystals, but enhances the code by graphically linking results directly to the analyst, so that long lists of numbers are replaced by 3-D views of crystals.

As input, the program receives a list of the asymmetric atoms of the crystal (the least set of atoms needed to describe the entire crystal), their location in space, orientation parameters in space, and a space group number (an index to a space group table that contains information needed to build a crystal). Based upon this input, the program can generate the location of all atoms within the unit cell of the crystal. Bonds between atoms are generated in software based upon the covalent radius of atoms and their distance from neighboring atoms. In addition, the unit cell may be replicated in three directions, thus producing a larger crystal. Once built, the crystal may be viewed in various ways, including ball and stick, cylinder, or balls using Van Der Waal's radii. Data, such as location, distance in angstroms, or angle in degrees may be received from the 3-D display of the crystal by picking on individual atoms with the computer's mouse. Atoms are highlighted as they are picked, and data is shown on the display monitor as well as in a separate text window.

Space Crystal not only allows modeling of crystals but also visualization of data produced by hours of computer computation using quantum chemistry codes. Currently, various isosurfaces of the electrostatic potentials of a crystalline molecule may be displayed. If desired, the isosurface could be shown simultaneously with the crystal model or the images could be merged together.

Space Crystal is still currently in alpha development. The first official release, Version 1, is expected in February 1992. Crystal was presented to an International conference in October 1991. Information from this conference will be used to enhance the future versions of Space Crystal.

Reference

1. L. A. Treinish, Tutorial on State of the Art in Data Visualization, ACM SIGGRAPH, 1989.

12. High-Performance Computing

R. J. Littlefield, G. S. Thomas, G. I. Fann, and A. B. Shiflet

The purpose of the Computing and Information Sciences program in high-performance computing is to provide computer science support to the task of exploiting the fastest available computers for computational chemistry. Its primary objectives are to develop analysis and programming techniques that will expand the range of applications for which parallel supercomputers can be used effectively, and to develop evaluation criteria by which computer systems can be designed and selected for use by the MSRC. In particular, the program includes research and development to

- develop techniques for predicting and evaluating the performance of a scientific code on a parallel computer
- evaluate alternate strategies for programming computational chemistry applications
- develop efficient implementations of kernel computations that are important for computational chemistry

- establish which parameters of a parallel computer system are most important to chemistry applications
- improve those parameters in future computer systems.

Approach

The high-performance computing program involves a combination of theoretical and experimental work on large-scale parallel machines. The work focuses on kernel computations for which existing tools and techniques fail to fully exploit the capabilities of parallel machines. The major components of this approach include

- an in-depth study of kernel computations chosen to push the envelope of what parallel machines can do effectively;
 - development of performance models for those kernels
 - improvement of kernel algorithms
- development and evaluation of tools and techniques for
 - performance modeling and visualization
 - programming parallel applications
- active participation in the Concurrent Supercomputing Consortium
- operating system research and vendor collaboration regarding critical parameters of parallel computer systems.

Activities and Results

Work in 1991 focused on two major computational kernels: eigensolution of dense symmetric matrices (needed for electronic structure calculations), and global vector sum (needed for electronic structure calculations), and global vector sum (needed for molecular dynamics using the replicated data

strategy). Both of these kernels are potential bottlenecks in practical applications and have proved difficult to implement efficiently on large-scale parallel machines. In each case, performance modeling techniques have been used to understand the tradeoffs between alternate algorithms and the effect of computer characteristics on performance. New algorithms have been developed, modeled, and studied experimentally on large-scale parallel machines such as Intel iPSC/860 and Intel Touchstone DELTA. This work has resulted in the development of faster algorithms and improved methods for modeling and characterizing parallel machines and programs. This work is described in more detail in a subsequent section, Parallel Computational Kernels.

Staff from the MSRC Computing and Information Sciences program have been active members of the Concurrent Supercomputing Consortium DELTA project team. MSRC provided feedback and direction to Intel regarding development of the DELTA system. Two of the DELTA acceptance test programs were provided by MSRC staff, and the improved global sum algorithms will be incorporated into subsequent releases of the DELTA software.

To support performance modeling for the computational kernels, and effort was begun to study communications system performance of the iPSC and DELTA machines. Early results of this study provided an explanation for previously observed failures of simple performance modeling techniques, and suggested directions for operating system research to improve communications performance. These findings are described in more detail below.

A one-day workshop on parallel eigensolvers was held at PNL in April 1991.

Parallel Computational Kernels

One kernel that has been studied is eigensolution of dense real symmetric matrices. Many electronic structure calculations require eigensolution of matrices having size N of a few hundred. When these calculations are performed on a large scale parallel computer, the number of available processors P may equal or exceed N . Performance in this regime is not well understood. Prior studies of parallel linear algebra, including eigensolvers, have

concentrated on the case $P \ll N$, where it is relatively easy to get good parallel performance by using uniform data distributions to parallelize a conventional sequential algorithm. However, when P is comparable to N (the "large- P regime") the effects of load imbalance and communications costs become more complicated, and significant performance improvements can be obtained by using nonuniform distributions.

A study of the performance of a distributed Jacobi method eigensolver showed that, on average, its performance could be improved by over 30% by using performance modeling to select an appropriate nonuniform distribution [DMCC6]. Subsequent study comparing the Jacobi method against other parallel solvers showed that the Jacobi method is also competitive in absolute speed in the large- P regime [TCA], even though it is relatively slow for small P . These effects are illustrated in Figure 12.1. The study also determined that parallel speedup was limited to modest values (10X) under the conditions that were measured, and that the most important computer system characteristic was communication startup cost.

These studies of eigensolver performance were primarily based on performance modeling, with the models generated manually. This procedure is apparently effective, but experience suggests that it may be too expensive to apply routinely. Subsequent work is planned to develop and evaluate methods of combining performance modeling, visualization, and language annotations such as those of Fortran-D. To this end, performance visualization tools have been obtained from Oak Ridge National Laboratory, Argonne National Laboratory, and the National Center for Supercomputer Applications.

The second kernel that has been studied is global vector sum, which is the task of adding together vectors contained in all processors of a parallel computer system. This operation is a possible bottleneck for molecular dynamics calculations using the replicated data strategy. As with eigensolving, it has been found that different algorithms are appropriate in different regimes (i.e., with different vector lengths and numbers of processors). For the DELTA system, no fewer than three completely different algorithms have been found, each of which is superior by a factor of at least 1.5 in some regimes.

Figure 12.1. Performance of several parallel eigensolvers for a fixed size matrix ($N=200$) and varying numbers of processors. The Jacobi method (BFJ) is relatively slow for small processor counts, but becomes superior with large numbers.

For the long vectors and processor counts that are of most interest in molecular dynamics applications, a new pipelined mesh algorithm has been demonstrated to be 2.5 times faster than the standard unpipelined binary tree method.

Parallel Communication System Performance

To support performance modeling of the computational kernels, a study was performed of the communication system performance of the iPSC/860 and DELTA computers. This study was designed to measure message latency and overhead under less than optimal conditions, such as the slight loss of synchronization that would be caused by imperfect load balance in an actual application. As illustrated by Figure 12.2, such loss of synchronization can cause an increase of up to 405 in the overhead costs of communication. This appears to be attributable to an increase in the number of processor interrupts, and suggests that redesign of the communications kernel of the computer operating system could significantly improve its performance. Research to explore this possibility is being pursued by further

Figure 12.2. Total time to receive and reply to a message, as a function of the timing of the incoming message.

collaborative studies within the framework of the Concurrent Supercomputing Consortium.

13. Data Acquisition System Development

G. A. Anderson

In 1991, the Computer Interface and Electronics team provided support for the development of the Fourier transform ion cyclotron resonance (FTICR) and tandem mass spectrometers (TMS). Both of these systems require high performance data systems to achieve the required instrument performance.

The goal for the FTICR data system is to locate a commercial solution or to influence a promising vendor's development efforts. In 1991, the following vendors and ICR developers were visited:

Rice University
East Cost ICR Meeting
Ohio State University
Extrel
American Cyanamid
Bruker
Harvard University
IonSpec

These visits allowed us to determine the important ICR specifications, define the present instrument limitations, and develop internal goals for the development of a state-of-the-art ICR. A commercial solution, or partial solution, will likely be available in the required time frame.

A high-performance, real-time computer system was purchased and tested to study its feasibility for an ICR (and other high-performance) data system computer. This computer system is a Versa Module Europa (VME)-based multiprocessor platform with a real time UNIX operating system running X windows. A CAMAC crate interface was installed, and software was written to interface to the Rice University CAMAC crate. This system was shipped to Rice and performance tested. These tests included interfacing and data collection from their transient digitizer. This testing allows us to determine the suitability of this platform for rapid prototyping and real-time control and acquisition.

The TMS development efforts are centered on the detector system and image intensifier. Several different systems were discussed and evaluated. A delay line readout system was selected, and development efforts were initiated. Several different delay

lines were built in-house and tested. In addition, a strip-line delay line was designed and then procured through a printed circuit board manufacturer. This device, which was not a full-size, full-resolution unit, was tested and provided promising performance. Additional delay line designs are in progress and research in this area will continue.

A TMS data system conceptual design has been performed. This data system will use commercial components to develop a test system for a full-size, full-resolution delay line. This test system will be developed in 1992 after testing of a full-size delay line if adequate delay line performance is achieved.

14. Electronics and Computer Interfaces

K. D. Steele and G. A. Anderson

The Electronics and Computer Interfaces group provides hardware and software development support to all of the experimentalists within the MSRC. Providing this support from a single group ensures consistency of development methods, simplifies system life-cycle maintenance, avoids duplication of effort, and enables the experimentalists to spend more time on science and less time on engineering.

The Interface Development Laboratory (IDL) will provide the tools and computing environment necessary to allow the Electronics and Computer Interfaces group to accomplish design, development, prototyping, and research activities associated with the data acquisition and instrument control systems employed by EMSL researchers. During FY 1991, initial planning for the IDL was completed.

Design and prototyping work was accomplished on a variety of systems and applications. Applications were developed on Macintosh, IBM PC-AT Industry Standard Architecture (ISA), and VME multiprocessor architectures. In addition to these specific prototyping efforts, we initiated work in the following areas:

- developing a standard system for creating, manipulating, and archiving data files
- developing a standard library of software modules for use with the instruments which will be used in the MSRC Laser Laboratory

- developing a graphical layout editor to be used during the design of future computer interface applications.

Several applications were developed for use with instrument control, data acquisition, and display. These applications support LiF experiments. These applications provide interfaces to boxcar averagers contained within standard network information management (NIM) BIN crates via an IEEE-488 bus, and a tunable dye laser via RS-232C serial line. To provide a comparison with classical software development methods, these applications were produced using a graphical development environment. The graphical development package is a software simulation of a data flow machine. The conclusions reached were that, for several reasons, this environment is not desirable for development and operation of timing-critical applications:

- The data flow paradigm does not support asynchronous interrupts.
- Debugging is complicated since the execution order of computational nodes is difficult to determine.
- The execution speed of developed applications is very slow.

- The graphical development environment does not provide significant advantage, in ease or speed of code development, compared to classical code development methods.

Design was completed and initial development begun for a data acquisition and instrument control system used in Time-Resolved Transient Absorption experiments. The computer system is of DOS-based ISA architecture. The software will control an optical delay line via IEEE-488, and will interface to a data acquisition card that plugs into the computer's internal bus.

The system requirements were established for a computer interface for the Tandem Time-Of-Flight Mass Spectrometry experiments. The application software will provide an interface to instrumentation contained within a CAMAC crate. The physical interface will be implemented via an IEEE-488 bus from an ISA computer to the CAMAC crate. The computer will provide data archiving and display, while the acquisition of the data and synchronization functions will take place within various CAMAC modules.

Initial design was completed for the development of custom electronic circuitry associated with the Transient Absorption of Supersonic Molecular Beam experiments.

Appendixes

Appendix A

MSRC Staff and Associates

Appendix A

MSRC Staff and Associates

MSRC Management

Dr. Michael L. Knotek is Manager of the MSRC and the EMSL, and the Senior Science Director for PNL. From 1985 to 1989, Knotek chaired the National Synchrotron Light Source, a DOE user facility located at Brookhaven National Laboratory. Before joining PNL, Knotek was affiliated with Sandia National Laboratories; the Naval Weapons Centers at Corona and China Lake, California; and Ames Laboratory at Iowa State University. In 1984, Knotek was recognized by the DOE Office of Basic Energy Sciences for outstanding research in solid-state physics for his work on stimulated desorption. In 1985, DOE recognized him for outstanding scientific accomplishments in metallurgy and ceramics for work on stress corrosion cracking in solids. In 1987, he became a Fellow of the American Physical Society.

Knotek earned his Ph.D. in physics at the University of California-Riverside in 1972. He was also a Visiting Scholar at the Cavendish Laboratory, Cambridge University, England. He worked on the Quantum Theory Project at the University of Florida and on transport in organic systems at Oklahoma

State University. Knotek is a member of Sigma Xi, the American Vacuum Society, the American Physical Society, the American Chemical Society, and the American Association for the Advancement of Science.

Knotek was co-chair of the planning study for advanced national synchrotron radiation facilities. He is currently chair of the Proposal Evaluation Board for the Advanced Photon Source at Argonne National Laboratory. He also is a member of the Program Review Panel for the Advanced Light Source at Lawrence Berkeley Laboratory, the Stanford Synchrotron Radiation Laboratory Review Committee, the University of Texas at Austin Science and Technology Advisory Committee, and the American Physical Society Panel on Public Affairs and the Division of Condensed Matter Physics Strategic Planning Committee. He serves on the board of directors of the Idaho Research Foundation, Inc., and the Washington Technology Centers. He has more than 80 publications on topics ranging from theories of charge transport to the effect of radiation on chemical bonding.

Dr. B. Ray Stults is the MSRC Deputy Manager for Programs. Previously, Stults was the manager of PNL's Chemical Sciences Department. Before joining PNL, Stults was an Associate Science Fellow at the Monsanto Company where he supervised research programs in small-molecule, x-ray crystallography; powder diffraction; x-ray adsorption spectroscopy; and surface science.

While with Monsanto, Stults developed an expanded program in surface science, including x-ray photoelectron spectroscopy, secondary-ion mass spectroscopy, and Auger spectroscopy. In addition, he developed a laboratory for protein engineering that included protein x-ray crystallography, interactive

molecular modeling, and macromolecular dynamics simulations.

Stults is a member of the American Chemical Society, the American Crystallographic Association, Sigma Xi, Phi Lambda Upsilon, the American Society for the Advancement of Science, and the Materials Research Society. He received his Ph.D. in organic chemistry from the University of Nebraska and did postdoctoral research with Professor F. A. Cotton at Texas A&M University. He has written more than 50 papers in the areas of crystallography and structure/function analysis for systems ranging from homogeneous catalysts to organic compounds active as agricultural products.

Dr. M. Tom Thomas is the MSRC Deputy Manager for Science Integration and Implementation. He is responsible for ensuring that the EMSL Project technical scope integrates fully with the MSRC and other PNL Research Centers. Also, he is responsible for establishing the appropriate protocols so the EMSL can be used as a collaborative research facility. Previously, he was Manager of the EMSL Project during the planning, conceptual design, and initial start-up phases. Prior to that, he was Manager of Research Operations and Services for the MSRC. Before assuming managerial duties, Thomas was a Staff Scientist and Group Leader responsible for developing an applied surface science group and a surface science analytical facility for PNL. His general research interest is in applying surface science to solve materials problems associated with surface and interface phenomena.

Thomas's research experience includes a study of environmental effects on fracture, grain boundary chemistry, and stress corrosion; material degradation due to corrosion; and wear and tribology. He also is developing standards for surface analytical studies (AES, XPS, SIMS, etc.).

Thomas has conducted research on problems related to industrial surface/interfaces, interfaces in mineral

processing, and silicon/gold interfaces as well as surface problems in fusion technology, including first-wall neutron sputtering and plasma/wall interactions in Tokamak plasma machines.

Thomas is chairman of Committee E42 on Surface Analysis of the American Society for Testing and Materials and is a member of the American Vacuum Society, the American Institute of Mechanical Engineers, the American Physical Society, and the American Association for the Advancement of Science. He received his Ph.D. in solid-state physics/surface physics from Brown University.

Thomas has written more than 50 papers on material science studies in energy technologies involving passivation and corrosion of amorphous materials, effects of grain boundary chemistry on stress corrosion, corrosion of iron alloys, and neutron sputtering from candidate fusion reactor first-wall materials. In addition, he has published review papers on the application of surface science techniques to study problems in tribology, mineral processing, and applications of surface analyses.

EMSL Project Management

Dr. James K. McClusky is Manager of the EMSL Project. McClusky has 28 years of experience in the nuclear and chemical processing fields, with 20 years in project and program management.

Prior to joining Battelle in 1991, McClusky was a Director at EG&G's Rocky Flats Plant. From 1977-1988, he held a number of responsible management positions with Rockwell International at Hanford. In 1983, McClusky led the successful restart of the Purex Chemical Processing Plant for Rockwell.

McClusky received his Ph.D. in nuclear/chemical engineering from Iowa State University in 1971. He was awarded a Post-Doctoral Fellowship at Ames Laboratory during 1972.

McClusky holds several patents in the area of nuclear fuel preparation. He is a member of the American Nuclear Society and the American Institute of Chemical Engineers.

Chemical Structure and Dynamics

Dr. Steven D. Colson became Associate Director for the MSRC Chemical Structure and Dynamics program in 1989. Between 1968 and 1989, Colson progressed from Assistant Professor to Professor of Chemistry at Yale University. Colson has written or contributed to more than 100 papers on photochemistry, photo-physics, and molecular dynamics; electronic structure of molecules, radicals, and molecular ions; processes at the molecule/surface interface; and intermolecular interactions in molecular solids.

In 1984, Colson was chairman for the Gordon Conference on Visible/UV Multiphoton Ionization and Dissociation Processes. He was also chairman of the 21st International Symposium on Free Radicals in 1991. From 1985 to 1987, he was a member of the Petroleum Research Fund Advisory Board. In 1988, he was guest professor at the University of Ulm in West Germany. He has also served as Associate

Editor of the *Journal of Chemical Physics*. Since joining the MSRC, Colson has been appointed an Affiliate Professor in the Chemistry Department at the University of Washington and an Adjunct Professor of Chemistry at Washington State University.

Colson received his Ph.D. in chemistry from California Institute of Technology in 1968 and spent a year as a Postdoctoral Fellow at the National Research Council in Ottawa, Canada before moving to Yale. He is a member of the American Chemical Society. Recently, Colson was awarded a Humboldt Fellowship by the Alexander von Humboldt Foundation in Germany. This award, which was in recognition of his past accomplishments in research and teaching, allowed Colson to study for six months at the Institute for Physical and Theoretical Chemistry in Munich.

Dr. Stephan E. Barlow recently joined the Chemical Structure and Dynamics program as a Senior Research Scientist. He received his Ph.D. in chemical physics from the University of Colorado at Boulder in 1984, working on very-low-temperature ion trapping techniques under the guidance of Professor Gordon H. Dunn of the Joint Institute for Laboratory Astrophysics (JILA) and the National Institute of Standards and Technology (NIST). He was a postdoctoral fellow in the Chemistry Department of the University of Colorado with Professor Charles DePuy and Dr. Veronica Bierbaum. Most recently he was an instructor in the Chemistry Department at the University of Colorado, Denver.

Barlow's work with ion traps led to techniques that allowed gas-phase, ion-molecule reaction kinetics to

be studied at temperatures below 77 K for the first time. His work in Professor DePuy's laboratory led to the development of greatly improved Selected Ion Flow Tube technology. Most recently he has been investigating the nature of clusters and cluster formation, work that will be continuing at the MSRC with new instrumentation.

Barlow's ion-trap work has also led to an interest in the physics of charged-particle storage. He was one of the first to apply the theory of non-neutral plasmas to the behavior of trapped ions. He is using such concepts to develop improved mass spectrometric techniques.

Dr. James P. Cowin is a Senior Research Scientist in the Chemical Structure and Dynamics program. His experimental research program looks at molecular dynamics of molecules adsorbed on surfaces, light-stimulated surface reactions, and charge transfer processes. His work includes making well-characterized models of the water/solid interface, complex

multilayer structures, and single molecule photochemistry with the scanning tunneling microscope. He is the author of more than 30 papers. Before coming to PNL, Cowin was an Assistant Professor of Physical Chemistry at the University of California-Santa Barbara, where he was recognized as one of the pioneers of laser-driven surface reactions. He studied

chlorinated and other halogenated hydrocarbon surface chemistry. He maintains a collaboration and membership there in the Quantized Electronic Structures Group.

Cowin earned his Ph.D. in chemistry at the University of Chicago in 1981 under the direction of Professor Lennard Wharton and built one of the first universal molecular-beam surface machines. His studies included many aspects of surface energy transfer and

structure, most notably the first quantitative use of laser surface heating for kinetic studies and the discovery of the phenomena of rotationally mediated selective adsorption.

Cowin did postdoctoral studies with Dr. John Polanyi at the University of Toronto. He built a beam surface machine and introduced Dr. Polanyi to laser-surface chemistry, now Polanyi's major thrust.

Dr. Donald M. Friedrich, a Senior Research Scientist, is responsible for the experimental study of dynamics at solid/liquid interfaces using the ultrafast laser spectroscopy capabilities in the MSRC. His previous research in PNL's Materials Science Department involved Raman spectroscopic studies of phosphazene polymers and optical thin-film materials.

Prior to joining PNL, Friedrich was Associate Professor of Chemistry at Hope College, where he constructed a laser spectroscopy laboratory dedicated to undergraduate chemistry research. While on sabbatical leave at the University of Minnesota and the University of California-Berkeley, Friedrich used picosecond and resonance Raman methods to study the kinetics and structural changes occurring during excited-state

intramolecular proton transfer in solutions. Friedrich's postdoctoral research at Wayne State University resulted in the first measurement of the polarized two-photon excitation spectrum of benzene. The unexpected nature of the polarization characteristics of this spectrum led to a reassignment of the excited-state vibrations in benzene and a new understanding of vibronic coupling in aromatic molecules.

Friedrich obtained his Ph.D. in chemistry from Cornell University under the direction of Professor Andreas Albrecht. He has written more than 20 publications on the spectroscopy of π -conjugated compounds and free-radicals, inorganic polymers, and thin-film optical materials.

Dr. Wayne P. Hess joined the MSRC's Chemical Structure and Dynamics program as a Senior Research Scientist in 1990. Previously, he was a Postdoctoral Research Associate at the Combustion Research Facility, Sandia National Laboratories, Livermore, California. The thrust of his postdoctoral work involved using cw-LIF to determine precise kinetic rate constants and radical reaction mechanisms. He also studied dissociation dynamics using both LIF and photofragment imaging techniques.

Hess received his Ph.D. in physical chemistry from the University of Colorado in 1988. His graduate studies,

conducted under the guidance of Professor Stephen Leone, involved the application of diode lasers to probe photodissociation quantum yields and chemical dynamics. He is a member of the American Chemical Society and the American Physical Society and has written or contributed to over a dozen publications on topics such as photodissociation dynamics, photofragment imaging, radical kinetics, and Raman spectroscopy.

Hess has presented papers at numerous national and international conferences and meetings.

Dr. Gary R. Holtom joined the MSRC in 1989 as a Senior Research Scientist. He is responsible for laboratory operations in the Chemical Structure and Dynamics program, including safety, organization, equipment specification, and user activities. Previously, Holtom was Head of Laser Operations in the Regional Laser and Biotechnology Laboratories at the University of Pennsylvania, where he maintained fast fluorescence techniques at the state of the art, trained local and visiting researchers, and assisted in interpreting results. He also developed apparatus and analysis methods and created software for data collection and analysis.

Holtom's graduate research involved building lasers and associated data acquisition equipment for two-photon spectroscopic measurements using analysis methods originated by his research director,

Dr. Stephen A. Joyce is a Senior Research Scientist in the Chemical Structure and Dynamics program. He received his Ph.D. in physical chemistry from the Massachusetts Institute of Technology with Professor Jeffrey Steinfield in 1987, studying reactive processes on silicon and silicon dioxide surfaces. He spent two years as a postdoctoral fellow at the National Institute of Standards and Technology, where he studied stimulated desorption of ions and metastable species from surfaces and conducted the first experiments to measure the angular distributions of negative ion desorption. He was most recently a postdoctoral associate at Sandia National Laboratories, where he

Dr. Bruce D. Kay is a Senior Chief Scientist in the Chemical Structure and Dynamics program. His main research interests are in the chemical dynamics and kinetics of gas-surface interactions, which he studies using a combination of molecular beam-surface scattering and laser-based spectroscopic techniques. His current research interests focus on the dissociative chemisorption of halogenated hydrocarbons on model oxide surfaces. Prior to joining the MSRC, Kay was a senior member of the technical staff in the Interfacial Chemistry Division at Sandia National Laboratories, Albuquerque, New Mexico. While at Sandia, Kay initiated and was principal investigator on research programs in the areas of quantum-resolved, gas-surface

Professor Martin McClain. As a Postdoctoral Fellow at Connecticut Wesleyan University under Professor Bryan Kohler, Holtom continued his work with high-resolution cryogenic methods. His primary goal was to relate spectroscopic measurements to photochemistry in octatetraene.

Holtom received his Ph.D. in physical chemistry from the University of California-Berkeley in 1978. He has written or contributed to 20 publications on topics ranging from the design of a birefringent filter for high-power dye lasers to time-resolved spectroscopy of the human brain. In addition, he organized and taught a short course in using lasers in chemical research as well as Chautauqua courses on dye lasers and laser methods in chemical research. The latter courses were intended to refresh high school and college teachers in an area of general interest.

developed the interfacial force microscope to directly measure the interactions of two surfaces at nanometer distances to obtain insight into phenomena such as adhesion and friction.

Joyce is interested in experimental studies of atomically-resolved reactions on solid surfaces using scanning probe techniques such as scanning tunneling and atomic force microscopies. He has over 30 publications on surface chemistry and is a member of the American Physical Society and American Vacuum Society.

scattering; kinetics of hydrogen in and on solids; gas-phase rotational relaxation; and kinetics of sol-gel formation.

Kay received his Ph.D. in chemical physics from the University of Colorado in 1982. He has written or contributed to more than 60 papers and has presented more than 100 lectures at universities and national and international meetings. He is a member of The American Chemical Society and The American Vacuum Society. In 1991 he received the John Yarwood Medal from the British Vacuum Council for his work in the field of state-to-state inelastic and reactive gas-surface scattering.

Dr. Robin S. McDowell joined the Chemical Structure and Dynamics program in 1991 as Senior Chief Scientist and Program Manager. He received his Ph.D. in physical chemistry from the Massachusetts Institute of Technology Spectroscopy Laboratory in 1961. Most of his professional career was spent at Los Alamos National Laboratory, where he was successively staff member, assistant group leader, and Laboratory Fellow. He has also been associated with the Los Alamos Technology Office at Rocky Flats, where he worked on environmental monitoring and remediation.

McDowell's research interests are in studies of molecular structure and dynamics through vibrational infrared and Raman spectroscopy, employing Fourier-transform spectrometers and tunable lasers. He is

Dr. Thomas M. Orlando is a Senior Research Scientist in the Chemical Structure and Dynamics program. His main research interests are in gas-phase cluster ion (nanocrystal) reaction dynamics and in electron, photon, and hyperthermal ion-beam-induced surface chemistry. This work involves the use of a wide variety of low-energy ion-beam techniques such as radiofrequency guided beams (quadrupoles and octopoles), selected ion flow tubes, a 7-tesla ion cyclotron resonance mass spectrometer, and sensitive laser techniques such as laser-induced fluorescence and resonance-enhanced, multiphoton-ionization spectroscopy.

Before joining the MSRC, Orlando did postdoctoral work in the Solid State Sciences Division at Sandia National Laboratories. The first quantum-resolved studies of electron stimulated surface chemical reactions were carried out in collaboration with

Dr. Douglas Ray joined the MSRC's Chemical Structure and Dynamics program as a Senior Research Scientist in 1990. Previously, he was a Postdoctoral Research Associate with W. Carl Lineberger at the Joint Institute for Laboratory Astrophysics in Boulder, Colorado, and a Research Assistant at the University of California-Berkeley.

Dr. Ray received his Ph.D. in physical chemistry in 1985 from the University of California-Berkeley,

also interested in using these techniques for atmospheric sensing and monitoring and in statistical-mechanical calculation of partition functions and thermodynamic properties. In the 1980s he led the successful effort to analyze the complex spectroscopy of SF_6 , UF_6 , and CF_4 , and to apply these results to laser photochemistry, laser isotope separation, and molecular laser development.

McDowell has written 95 papers and several review articles, as well as the entry on "Infrared Spectroscopy" for the *McGraw-Hill Encyclopedia of Science & Technology*. He served as President of the Coblenz Society from 1987 to 1989 and has been active on editorial and conference organizing committees.

Dr. Alan Burns who has pioneered quantum-resolved, electron-stimulated desorption studies. Postdoctoral work in ion spectroscopy and photoinitiated intra-cluster ion-molecule reactions was also done with Professor J. P. Maier at the Institute for Physical Chemistry, University of Basel, Switzerland. Orlando's graduate studies, conducted under the direction of Professor S. L. Anderson at State University of New York, Stony Brook, concentrated on state-resolved studies of polyatomic ion-molecule reaction dynamics.

Orlando has written or contributed to 20 papers that discuss stimulated surface processes, ion-molecule reaction dynamics, and multiphoton spectroscopy. He has also given numerous invited presentations at national and international conferences and is a member of The American Physical Society and The American Vacuum Society.

where his advisor was Richard J. Saykally. He has written or contributed to more than a dozen publications on topics such as ultrafast time-resolved dynamics in mass selected clusters and has been invited to lecture at gatherings such as the 5th International Symposium on Small Particles and Inorganic Clusters in Konstanz, Germany, in 1990.

Dr. Steven W. Sharpe is a Senior Research Scientist in the Chemical Structure and Dynamics program. He received his Ph.D. in physical chemistry from the State University of New York, Stony Brook, in 1986. Following postdoctoral work at the University of Southern California, he joined PNL in 1990. Sharpe's early research dealt with kinetic reaction studies of extraterrestrial atmospheres. His graduate studies, conducted under the guidance of Professor Philip Johnson, included the development of a novel technique for spectroscopic investigations of short-lived

and unstable gas phase chemical species. Postdoctoral studies, conducted under the auspices of Professor Curt Wittig, included the building and implementation of a laser laboratory for the study of weakly bonded, gas-phase molecules.

Sharpe is a member of the American Chemical Society. He has written or contributed to more than a dozen publications, the most recent of which appeared in advances in molecular vibrations and collisional dynamics.

Environmental Dynamics and Simulation

Dr. James E. Amonette is a Research Scientist in PNL's Earth and Environmental Sciences Center and part of the team responsible for the MSRC Environmental Dynamics and Simulation program. Since joining PNL in 1986, he has specialized in the areas of aqueous geochemistry and soil mineralogy. Amonette's primary interest is in using proton-induced, x-ray emission; x-ray diffraction; infrared; and other spectroscopic techniques to characterize solid phases and to predict and monitor reactions associated with the disposal of hazardous wastes in soil

environments. He received his Ph.D. in soil chemistry from Iowa State University in 1988.

Amonette is a member of the Clay Minerals Society, the International Soil Science Society, and the Soil Science Society of America. He has written or contributed to more than two dozen articles and other publications on topics such as oxidative weathering of micas, infrared spectroscopy of iron-chromium hydroxides, and synchrotron-based x-ray scattering studies of soil materials.

Dr. Ben P. Hay is a Senior Research Scientist in PNL's Earth and Environmental Sciences Center and a member of the MSRC Environmental Dynamics and Simulation Group. His current research interests center on the development and application of molecular modeling techniques for the elucidation of structure/activity relationships in coordination chemistry.

Prior to joining PNL, Hay held an appointment as a Postdoctoral Fellow with the Inorganic Structural Group at Los Alamos National Laboratory from 1989 to 1991. There he developed a molecular mechanics model for aqua, nitrate, and β -diketonato complexes of the f-block metal ions. This research was preceded by a two-year appointment as a Postdoctoral Fellow with Professor Athel L. J. Beckwith in the Research School of Chemistry, Australian National University.

Hay elucidated the detailed kinetics and mechanisms for intramolecular rearrangements of alkyl and alkoxy radicals in the liquid phase. This work led to the discovery of two new classes of alkoxy radical precursors.

Under the guidance of Professor Richard G. Finke, Hay received his Ph.D. in Chemistry from the University of Oregon, where his graduate research dealt with kinetic and mechanistic studies of the liquid-phase reactions of organo-cobalt compounds. He is member of the American Chemical Society and has written or contributed to more than a dozen publications on topics such as the thermolysis of coenzyme B₁₂, the reversible cyclization of the 5-pentyloxy radical, synthesis of N-alkoxy-pyridine-2-thiones, and the molecular modeling of lanthanide complexes.

Dr. Charles J. Hostetler is a Senior Research Scientist in PNL's Earth and Environmental Sciences Center and is a member of that center's liaison team to the MSRC. Since joining PNL, he has been involved in applications of geochemistry to contaminant transport. Hostetler's primary contributions have been in the development and application of reactive chemical transport models that include the effects of advection, dispersion, diffusion, aqueous speciation, adsorption, ion exchange, and solubility reactions on the mobility of solutes. His current research interests

include the development of property estimation schemes for organic chemicals and uncertainty analysis for geochemical calculations.

Hostetler received his Ph.D. in planetary sciences at the University of Arizona in 1982. He has written or contributed to more than two dozen publications and reports on topics such as chemical transport models, geohydrochemical codes, and performance assessment analyses of waste disposal facilities.

Macromolecular Structure and Dynamics

Dr. Thomas S. Tenforde is Chief Scientist in PNL's Life Sciences Center and leads the MSRC Macromolecular Structure and Dynamics program. Prior to joining PNL in 1988, he spent 16 years at Lawrence Berkeley Laboratory, most recently as leader of the Environmental Physiology group in the Biology and Medicine Division. Since 1980, Tenforde has served on two dozen national committees and boards and has been a reviewer for federal agencies, private organizations, and more than 20 refereed journals. He is a member of the American Association for the Advancement of Science, the New York Academy of Sciences, the Radiation Research Society, the Biophysical Society, and the Bioelectromagnetics Society (president in 1987-1988). He is currently a member of the National Academy of Sciences Board on Radiation Effects Research, the National Council on Radiation Protection and Measurements (Board of Directors), the American Conference of Governmental Industrial Hygienists, and the IEEE Committee on Man and Radiation.

He chairs the COMAR ELF/VLF subcommittee and the National Research Council committee on the potential health effects of electromagnetic radiation

from the Air Force's Ground Wave Emergency Network.

Tenforde has written or contributed to 185 publications and reports and has presented 80 invited lectures at conferences and symposia during the last 10 years. Recent invited lectures include talks at the 9th International Congress of Radiation Research in Toronto, Canada; the 16th Annual Conference of the Australian Radiation Protection Society; and the International Conference on Advances in Power System Control, Operation, and Management in Hong Kong.

Tenforde attended Harvard University from 1958 to 1962, graduating with an A.B. degree *cum laude*. In 1959, he received the Harvard Dean's Honorary Scholarship. He was a National Institutes of Health (NIH) Predoctoral Fellow at the University of California-Berkeley (UC-B) from 1963 to 1968 and received his Ph.D. in biophysics in 1969. Tenforde was a Bay Area Heart Research Committee Postdoctoral Fellow at UC-B from 1969 to 1971 and an NIH Postdoctoral Fellow from 1971 to 1973.

Dr. Gary P. Drobney, Professor of Chemistry at the University of Washington, was placed under contract during 1991 to head the MSRC NMR program. Ultimately, he will assume the management role for the development of the ultra-high-field (gigahertz) NMR system planned for the EMSL.

Drobney has been affiliated with the University of Washington since 1981, when he accepted a position as a faculty research associate and operations manager of the University's NMR laboratory. He has been a member of the National Science Foundation's Advisory Committee for Biological Facilities Centers since 1987.

Drobney earned his Ph.D. in chemistry at the University of California - Berkeley in 1982. He graduated summa cum laude with a B.S. in chemistry

Dr. Joseph A. Loo is a Senior Research Scientist in PNL's Chemical Methods and Separations group and a member of the MSRC Macromolecular Structure and Dynamics program. Since joining PNL in 1988, his research areas have utilized mass spectrometry for the sensitive detection and structural characterization of large biopolymers. He has an active role in the development of capillary electrophoresis-mass spectrometry for high-resolution separations of complex mixtures, especially electrospray ionization-tandem mass spectrometry for sequencing proteins and oligonucleotides. Dr. Loo is also involved in the development of Fourier transform-ion cyclotron resonance (FTICR) mass spectrometry for ultrahigh resolution detection of biomolecules.

Dr. Alan L. Rockwood is a Senior Research Scientist in the MSRC's macromolecular structure and dynamics group. His research involves investigations of structure and reactivity of molecular and ionic species using a combination of experimental and theoretical methods. He is leading the development team for a new state-of-the-art tandem mass spectrometer for the MSRC and has recently developed a technique for studying fundamental properties of gas-phase ions by thermally induced dissociation. His theoretical work includes *ab initio* potential surface calculations on proton transfer reactions and simple model calculations on the dissociation of highly charged ions.

Prior to joining PNL, Rockwood was an applications chemist at JEOL, USA. He also worked as a mass spectrometrists at the University of Delaware, a

Dr. Richard D. Smith, Senior Staff Scientist, leads PNL's Chemical Methods and Separations group and directs research related to the development of new instrumentation, supercritical fluid solvation phenomena, high resolution separation methods, mass spectrometry, and novel separation processes. For the

from San Francisco State University in 1976. During the past five years, he has published more than 40 articles, reports, and papers in the field of NMR.

Loo received his Ph.D. in analytical chemistry in 1988 from Cornell University. His graduate work, with Professor Fred W. McLafferty, transferred fast ion bombardment, UV-laser desorption, and high energy californium-252 plasma desorption ionization methods to FT-ICR-MS for polymer and biomolecule analysis. He is a member of the American Chemical Society, American Society for Mass Spectrometry, and the American Association for the Advancement of Science. He has written or contributed to more than 30 publications in the area of mass spectrometry.

research scientist at the University of Houston, and a faculty intern at the University of Utah. In addition to his work as a professional scientist, he has carried on an independent research program at home which has resulted in the publication of three papers in *Physical Review* related to electrochemical thermodynamics.

Rockwood earned his Ph.D. in physical chemistry at Utah State University in 1981 under the direction of E. A. McCullough, Jr. He tested and disproved a previously proposed hypothesis for mode-selective chemistry for certain gas-phase isotope exchange reactions. As is characteristic of Rockwood's approach, his thesis research included both experimental (photo chemical) and theoretical (*ab initio* potential surface and kinetic modeling) components aimed at solving a specific scientific problem.

past 5 years, his research has centered on the fundamental aspects and applications of mass spectrometry and the development of new analytical instrumentation, particularly related to large molecule characterization.

Smith has also been the principal investigator for programs studying high temperature kinetics, organized molecular assemblies in fluids, and fundamental fluid phase and heterogeneous phenomena. In 1978, he was awarded the Bituminous Coal Research/R. A. Glenn Award for his work related to the mechanism of fly ash formation during combustion and trace element enrichment on fly ash surfaces.

Smith received R&D 100 Awards for the development of supercritical fluid chromatography-mass spectrometry (1983), the development of the first direct interface combining capillary zone electrophoresis with mass spectrometry (1988), and a process for materials formation from supercritical fluids (1988). In 1988, he patented processes for forming fine powders and thin films from supercritical solutions.

He has presented more than 100 lectures at national and international meetings. He was awarded an Associated Western University-DOE Distinguished Lectureship for the 1990-1991 academic year.

Smith received his Ph.D. in physical chemistry from the University of Utah. He is an Adjunct Professor at Washington State University. He is a member of the American Chemical Society, the American Society for Mass Spectrometry, the American Society for Testing and Materials, and the American Association for the Advancement of Science. Smith has been awarded six patents and has written or contributed to more than 225 papers on mass spectrometry, separation processes, and supercritical fluids and their application to materials processing and separation methods.

Materials and Interfaces

Dr. Jiri (Art) Janata joined the MSRC as Associate Director for the Materials and Interfaces program in 1991. Between 1987 and 1991, he was a professor in the University of Utah's Department of Materials Science and Engineering as well as the Director of the HEDCO Microengineering Facility. While at the University, Janata conducted research programs involving the development of novel chemical sensors, the development of microelectronic probes for medical applications, solid-state chemical sensors for air pollution control, and charge transfer at the liquid/liquid interface. During this period, Janata also was Visiting Professor at the Universität der Bundeswehr München and EPFL Lausanne.

Janata received his Ph.D. (*summa cum laude*) in analytical chemistry from Charles University in Prague, Czechoslovakia, in 1965. He was a Visiting Fellow at Wolfson College, Oxford University from 1986 to 1987 and received the Humboldt Senior Scientist Prize in 1987 and the Heyrovsky Medal in

1990. He currently is a member of the editorial boards for *Sensors and Actuators*, *Biosensors*, *Sensor Technology*, and *Talanta*.

Janata is the author or co-author of more than a dozen book chapters and over 120 refereed publications. His book, *Principles of Chemical Sensors*, was published by Plenum Publishers in 1989 with a second printing in 1990. He also serves as a referee for organizations such as the National Institutes of Health, the National Science Foundation, and the Australian Research Council, and for publications such as *Analytica Chimica Acta*, the *Journal of American Chemical Society*, the *Journal of Inorganic Chemistry*, and *Biosensors and Bioelectronics*.

Janata has received more than a dozen patents. He is a member of the Electrochemical Society, the American Chemical Society, the Society for Electroanalytical Chemistry, and the American Association for the Advancement of Science.

Dr. Donald R. Baer is a Staff Scientist in PNL's Materials Research Section. Since joining PNL in 1976, he has specialized in the use of surface-sensitive techniques to examine effects of surface and interface chemistry on materials properties. He currently is principal investigator and group leader of a team

developing the experimental components of the Materials and Interfaces program for the MSRC.

During 1984 and 1985, Baer was a Visiting Research Fellow in the Department of Materials Science and Engineering at the University of Surrey, Guildford,

England. He is a member of the American Vacuum Society, is secretary of the Applied Surface Science Division, an associate editor of the *Journal of Vacuum Science and Technology*, and an associate editor of *Surface Science Spectra*. He is also a member of the American Physical Society, the Metallurgical Society of the American Institute of Metallurgical Engineers, the American Association for the Advancement of Science, and the American Society for Testing and Materials E42 Committee on Surface Analysis. He

Dr. Bruce C. Bunker, a Senior Research Scientist, is studying the solution chemistry of oxides. His areas of interest include understanding and controlling the formation of oxides and hydroxides from aqueous solutions, oxide dissolution processes, colloid chemistry, and tailoring the surface chemistry of oxides for technological applications in separations, catalysis, sensors, and thin films. Bunker currently is involved in programs to control oxide nucleation and growth in solution and on surfaces via complexation, biomimetic synthesis of oxide films, design and synthesis of inorganic ion exchangers, and synthesis of environmentally relevant materials such as clays.

Prior to joining the MSRC, Bunker was a member of the Electronic Ceramics Division at Sandia National Laboratories, where he supervised research activities in areas such as sol-gel preparation of ferroelectric

Dr. Michael A. Henderson is a Senior Research Scientist in the Materials and Interfaces program. He specializes in the surface chemistry of metal and metal oxide materials under ultrahigh vacuum conditions and examines the structure, dynamics, and reactivity of simple chemical systems at single-crystal surfaces with molecular-level probes such as electron energy loss, secondary ion mass, and Auger electron spectroscopies. The goal of this research is to understand the complex chemistry of environmentally related surface phenomena.

Before joining the MSRC in 1991, Henderson was a postdoctoral researcher at the University of Pittsburgh, where he examined electron- and photon-stimulated processes at metal surfaces. This research

was Subcommittee Chairman for E42.02 (terminology) from 1984 to 1989 and is now serving as membership Secretary.

Baer received his Ph.D. in experimental solid-state physics from Cornell University in 1974 and did postgraduate work in the Department of Physics and Materials Research Laboratory at the University of Illinois, Urbana, between 1974 and 1976. He has written or contributed to more than 85 publications.

thin films and the synthesis and properties of other ceramic materials, including oxide superconductors, high-field varistors, and boron carbides. His earlier work at Sandia involved a wide range of materials chemistry problems, including studies of the mechanisms of dissolution and stress corrosion cracking of glasses, use of NMR and Raman techniques to study glass structure and oxide reactions, solutions techniques for synthesizing oxide superconductors and other ceramics, and development of glasses for use in advanced lithium batteries.

Bunker obtained his Ph.D. in inorganic chemistry from the University of Illinois, Champaign-Urbana, under the direction of Professor Russell Drago. He has written more than 70 publications on various aspects of the structure and chemistry of oxide materials.

yielded interesting results about the anisotropic nature of low-frequency adsorbate vibrations at atomic-high steps. Such vibrations are believed to be the precursors to processes such as surface diffusion and reaction.

Henderson obtained his Ph.D. in physical chemistry in 1988 from the University of Texas, Austin, under the direction of J. Michael White. He has written or contributed to 40 publications on topics including organophosphorus surface chemistry, Fischer-Tropsch catalysis, and surface photochemistry. He is a member of the American Vacuum Society, the American Chemical Society, and the Materials Research Society.

Dr. John P. LaFemina joined the MSRC as a Senior Research Scientist in 1989 and is currently the Program Manager for the Materials and Interfaces program. His research seeks a comprehensive understanding of the atomic and electronic structure, including atomistic dynamics, of insulator and semiconductor surfaces important in areas as diverse as the design of advanced electronic materials, chemical sensors, and subsurface contaminant transport. Between 1985 and 1989, LaFemina was a Captain in the United States Army at the United States Military Academy, serving as a Deputy Director of the Academic Research Division. The Division is a faculty research center consisting of the Science Research Laboratory and the Thomas H. Johnson Photonics Research Center.

In addition to the modeling of surface structure and chemistry, LaFemina's other research interests include

Dr. Gary L. McVay is Manager of PNL's Materials Sciences Department and Deputy Manager of the MSRC. He joined PNL in 1978 as a research scientist and has held positions of increasing responsibility since then. McVay's areas of specialization include ionic mobilities in glass; interactions of ceramic materials with aqueous solutions and the effects of radiation on these interactions; structural water effects on silicate properties; defect mobilities and diffusional properties in semiconductor materials; and neutron irradiation-induced deformation in metals.

Prior to joining PNL, McVay was with the solid-state physics research section at Sandia National

modeling the electronic structure and photophysics of thin film polymers and investigating the morphological dependence of the electronic structure for molecular and periodic systems.

LaFemina earned his Ph.D. at Pennsylvania State University under the direction of Dr. John P. Lowe in 1985. He has written or contributed to more than 60 papers and presentations on topics ranging from the photophysics of polyamide and poly(ethylene terephthalate) to the atomic and electronic structure of compound semiconductor and mineral oxide surfaces. He is a member of the American Chemical Society, the American Physical Society, the American Vacuum Society, and the Materials Research Society.

Laboratories and the EBR-II site at Idaho Falls, Idaho. He received his Ph.D. in ceramic engineering from the University of Missouri in 1970.

McVay is coordinator of the Materials Sciences' Basic Energy Sciences programs at PNL. He is a member of the American Ceramics Society, American Nuclear Society, Materials Research Society, Pacific Northwest Materials Council, Materials Directors' Council, and Advisory Committee of the Washington Technology Center. He currently has 70 open literature publications, has presented numerous invited papers, and was the editor of five proceedings documents.

Theory, Modeling, and Simulation

Dr. Thom H. Dunning, Jr., became Associate Director for the MSRC Theory, Modeling, and Simulation program in 1989. Before joining the MSRC, he was Senior Scientist and Group Leader of the Theoretical and Computational Chemistry Group and Head of the Chemical Dynamics Program at Argonne National Laboratory. Dunning is an Adjunct Professor at Washington State University and an Affiliate Professor at the University of Washington.

Dunning has written or contributed to more than 80 publications on topics ranging from advanced techniques for molecular calculations (including the development of molecular basis sets and generalized valence bond theory) to applications involving the photophysics of atmospheric species and exotic lasermolecules, and chemical reactions involved in combustion. More recently, he has been involved in modeling the structure and energies of ion-water

clusters and the effects of solvation on the energetics and dynamics of chemical reactions.

Dunning is a member of the American Chemical Society, the American Physical Society, and the American Association for the Advancement of Science. Since 1987, he has been the Series Editor for *Advances in Molecular Electronic Structure Theory* published by JAI Press, Inc. Currently, he is a member of the Executive Committee of the Energy Research Supercomputer Users' Group, the National Science Foundation's Chemistry Advisory Committee, and the Policy Board

Dr. Raymond A. Bair joined the MSRC's Molecular Science Software group as a Staff Scientist and Technical Group Leader in 1990. The group's present work is focused on development of electronic structure and molecular dynamics codes for massively parallel computers; graphical user interfaces for computational chemistry codes; graphical visualization and analysis tools; and molecule information databases. The group is also a principal contributor to the design and specification of the Molecular Science Computing Facility.

Bair received his Ph.D. in theoretical chemistry from the California Institute of Technology in 1981. He is a member of the DOE Energy Sciences Network Steering Committee, the American Chemical Society, American Association for the Advancement of

Dr. L. René Corrales is a Senior Research Scientist in the MSRC Theory, Modeling, and Simulation program. His theoretical research program explores the nature of collective properties in complex condensed matter systems and includes the study of collective phenomena associated with atomic and molecular rearrangement such as those observed in amorphous materials, polymeric solutions, and amphiphilic monolayers.

Before coming to PNL, Corrales was a National Science Foundation Postdoctoral Fellow in Chemistry at the University of Texas, Austin, working with Professor Peter J. Rossky on a novel format to model

of the Concurrent Supercomputer Consortium. He also is Vice President of the Advisory Board of the Foundation for Chemical Research (University of Missouri-Rolla) and co-chairman of the Hanford Tank Waste Science Panel.

Dunning was awarded a Woodrow Wilson Fellowship in 1965-66 and a National Science Foundation Fellowship in 1966-69. He received his B.S. in chemistry from the University of Missouri-Rolla in 1965 and his Ph.D. in chemical physics from the California Institute of Technology in 1969.

From 1987 to 1990, Bair was director of new product development with BioDesign Inc., which produced interactive molecular modeling and simulation tools for the chemical industry. Prior to 1987, Bair was an Assistant Scientist and postdoctoral fellow at Argonne National Laboratory, where he studied the energetics and mechanisms of combustion reactions and developed *ab initio* computational methods for vector and parallel computers.

Science, and Sigma Xi. In 1979, he received the Lazlo Zechmeister Fellowship, and from 1976 to 1979 he completed a National Institutes of Health traineeship. He has written or contributed to more than a dozen publications in his field.

covalently bonded materials. This methodology has been applied successfully to the liquid and amorphous states of sulfur and is being extended to study other covalently bonded liquid and amorphous systems.

Corrales earned his Ph.D. in chemical physics at the University of California, San Diego, in 1988 under the direction of Professor John C. Wheeler. His studies included understanding the role of chemical reactions in driving-phase transitions that lead to critical phenomena. In addition, he explored the relationship between lattice polymer solution models and analogous magnetic models in which novel, higher-order critical phenomena were found.

Dr. Liem X. Dang joined the MSRC Theory, Modeling and Simulation program as a Senior Research Scientist in 1991. Previously, he was a Visiting Scientist at the IBM Research Division in San Jose, California and a Postdoctoral fellow at the University of California, San Francisco.

Dang received his Ph.D. in theoretical chemistry in 1985 from the University of California, Irvine, where

his advisor was Professor Max Wolfsberg. Dang's research interest focuses on molecular dynamics simulation of chemical and physical processes. He has written or contributed to more than 20 articles on topics such as protein stability, ionic clusters, and ionic solutions, including the development of nonadditive intermolecular potentials for water-water and ion-water interactions. Dang is a member of the American Chemical Society.

Dr. David F. Feller joined the MSRC as a Senior Research Scientist in 1989. Between 1984 and 1989, he was a Research Associate Scientist and Manager of the Chemistry Computational Facility at Indiana University. His primary areas of research interest are the development and application of techniques for the theoretical investigation of ground-state and excited-state potential energy surfaces as well as the study of molecular properties.

During his academic career, Feller was awarded a Collins Radio Scholarship and the Regents

Competitive Scholarship while at Loras College in Dubuque, Iowa. Subsequently, he received a Proctor and Gamble Scholarship. He earned his Ph.D. in physical chemistry at Iowa State University in 1979.

Feller has written or contributed to more than 60 articles. Most recently, these have included studies of the hyperfine parameters in small radicals and the calculation of momentum distributions for comparison with electron momentum spectroscopy.

Dr. Martin W. Feyereisen recently joined the MSRC as a Research Scientist and is currently developing a program for high performance computing on massively parallel computers and loosely coupled networks. He is conducting research on electron affinities of carbon clusters, chemisorption onto metallic oxides, and basis set accuracy for computing nonlinear optical properties.

Between 1990 and 1991, Feyereisen was a post-doctoral researcher at the University of Utah, where he developed computational methods to determine the structure and property of large systems. He also worked extensively on implementing post Hartree-Fock methods and participated in computational studies of various chemical systems, including stabilities of carbon clusters, proton transfer in solutions, reactivities of substituted purines, and modeling enzymatic behavior.

In 1985, Feyereisen received a fellowship from the Microelectronics and Information Sciences Center of the University of Minnesota. He spent the next five years at the University as a Fellow and Research Assistant, developing methods to accurately compute properties of various chemical systems. Using these methods, he studies systems such as atomic adsorption onto metallic surfaces, electronic spectra and nonlinear optical properties of organic and organometallic molecules, and binding properties of polymers.

Feyereisen earned his Ph.D. in chemistry in 1990 from the University of Minnesota, where his advisor was Dr. Jan Almlöf. He received a 3M Scholarship for Achievement in Chemistry in 1983 and was elected into Delta Epsilon Sigma, a National Honor Society. In 1984, Feyereisen graduated *magna cum laude* from the College of St. Thomas in St. Paul, Minnesota, receiving a B.A. in chemistry.

Dr. Bruce C. Garrett is a Staff Scientist and Technical Group Leader of Molecular Theory and Modeling Group. Before joining the MSRC in 1989, he was a founder and research scientist at Chemical Dynamics Corporation, where his responsibilities included developing and directing basic research in theoretical physical chemistry.

Garrett has been actively engaged in research involving theoretical and computational chemistry, particularly in the field of the dynamics of atomic and molecular collisions. Since 1977, Garrett has been a key participant in the development of practical and reliable methods for predicting rate constants for gas-phase chemical reactions. He has written or contributed to more than 100 publications on a variety of

Dr. Anthony C. Hess became a full-time MSRC Senior Research Scientist in 1991, after spending two years with the organization as a Postdoctoral Fellow. Hess earned his Ph.D in Chemistry at Arizona State University in 1988. His graduate research was a combination of theoretical and experimental studies in the fields of solid state chemistry and geochemistry. During this time, he was vice president of Phi Lambda Upsilon, the Honor Chemical Society. He received his B.S. in Chemistry from North Dakota State University in 1984.

Since joining the MSRC, Hess has initiated a research program that both uses and develops state-of-the-art *ab initio* solid-state quantum chemistry methods for studying complex mineral systems. Hess' ongoing collaboration with the authors of CRYSTAL involves the development and testing of new approaches in

topics including gas-phase reaction dynamics, translational-to-rotational and translational-to-vibrational energy transfer in gas-phase collisions, quenching of electronically excited states in gas-phase and gas/surface collisions, and chemisorption and diffusion of hydrogen on metal surfaces. His interests have recently turned to the study of dynamics in condensed phases.

Garrett received his Ph.D. in physical chemistry from the University of California, Berkeley, in 1977. He is a member of the American Chemical Society and the American Physical Society. He has served as a committee member on the National Research Council Review Panel for the AFOSR High Energy Density Materials program since 1990.

periodic Hartree-Fock theory. Hess' research group is focused on implementing new theoretical methods for studying the physical properties of a range of environmentally and catalytically important materials. These include a diverse class of materials from minerals (e.g., clays, zeolites, and other oxides) to different types of hydrated ices.

Hess is a member of the American Physical Society, the American Chemical Society, the American Geophysical Union, and the American Mineralogical Association. In 1991, Hess has presented invited lectures on "Periodic Hartree-Fock Theory of Low Symmetry Systems: Applications and Perspectives" (Orsay, France), "*Ab Initio* Methods in Geochemistry and Mineralogy" (Arizona State University), and "Experimental and Theoretical Investigations of the Mineral Kaolinite" (American Chemical Society).

Dr. Rick A. Kendall is a Senior Research Scientist in the Molecular Science Software Group interested in developing portable modern *ab initio* quantum chemistry codes on advanced architectures. In FY 1989, he was a Postdoctoral Appointee with the Theoretical Chemistry Group at Argonne National Laboratory's Chemistry Division. In FY 1990, Kendall joined the MSRC as a research scientist in the Theory, Modeling, and Simulation program.

In 1983 Kendall graduated *magna cum laude* from Indiana State University and as a member of Phi Kappa Phi. In 1987 at the University of Utah, he received the Stauffer Graduate Research Prize and

Dr. Delbert L. Lessor, a Staff Scientist in the Applied Physics Center, spends 50 percent of his time with the MSRC Materials Theory Group. His special interests include fundamental surface science and environmental remediation, and he has experience in basic and applied work in both areas. Lessor has participated in deducing structure of molecular adlayers on a number of materials using low energy electron diffraction, with emphasis in these studies shifting from electronic materials to those that are environmentally relevant.

He also models contaminant motion and holdup in soil and groundwater, where the fundamental inter-

Dr. Maureen I. McCarthy joined the MSRC in January 1991 as a Senior Research Scientist. From 1988 to 1991, McCarthy was awarded a fellowship from the Lady Davis Fellowship Trust to pursue postdoctoral research at the Hebrew University in Jerusalem, Israel. She earned her Ph.D. in Chemistry from the University of Colorado in 1988. Her thesis work combined experimental and theoretical techniques in photochemistry. During her graduate studies, she worked at the Fachbereich Chemie Universitat in Frankfurt, Germany, as a NATO Visiting Scientist and was awarded a Doctoral Fellowship by the University (1983-1984). She received her B.S. in Chemistry from Boston College in 1983 (*magna cum laude*, Phi Beta Kappa).

th Graduate Research Fellowship. He earned his Ph.D. from the Chemistry Department at the University of Utah in 1988.

Kendall is a member of the American Chemical Society and the American Physical Society. He has performed and presented research that encompasses vibration-rotation-induced electron detachment of molecular anions, clusters of molecular anions, analytic evaluation of non-Born-Oppenheimer matrix elements, electron affinities of first row atoms, and parallelization of *ab initio* software on both distributed workstations and on traditional parallel supercomputers.

actions at soil particle surfaces are key to remediation schemes such as steam stripping, *in situ* Joule heating, and *in situ* vitrification.

Lessor joined PNL in 1967 after receiving his Ph.D. from Kansas State University. He has developed models and performed computations in diverse areas such as (alpha, neutron) nuclear reactions, electrical phenomena in filtration, phase-contrast microscopy, and multi-electrode multiphase Joule heating.

McCarthy's research interests are in modeling heterogeneous processes that occur at solid surfaces. Her work includes studies on environmentally important systems found in the subsurface and the atmosphere. She is currently involved in an effort that combines methods of *ab initio* solid-state quantum chemistry and classical mechanics to study interfacial dynamics. One goal of this work is to characterize the behavior of contaminants on mineral surfaces.

McCarthy is a member of the American Chemical Society, the American Physical Society, and the American Geophysical Union. She maintains collaborations and has published articles with colleagues in experimental and theoretical chemistry, physics, and geochemistry.

During 1991, McCarthy has given presentations or invited lectures at the West Coast Theoretical Conference, the Oregon Graduate Institute, the

Northwest Regional Meeting of the American Chemical Society, and the Annual Fall Meeting of the American Geophysical Union.

Dr. Rick L. Ornstein joined the MSRC as a Senior Research Scientist in 1988 and was appointed Adjunct Associate Professor in Chemistry at Washington State University in 1990. He was promoted to Staff Scientist at PNL in 1991. From 1985 to 1988, he was a research scientist with Kodak in the Health and Nutrition Department where he set up a peptide/protein modeling and simulation research laboratory. From 1982 to 1985, he was a research staff member/lecturer at Princeton University in the Biochemistry Department.

After receiving his Ph.D. in Biophysical Sciences from the State University of New York, Buffalo, in 1979, he took a postdoctoral fellowship from 1979 to 1982 in the research group of Professor Jacques R. Fresco in the Biochemistry Department at Princeton University. He received his masters degree in Environmental Health Sciences in 1974 from the City University of New York at Hunter College and his B.A. from the City University of New York at Queens College in

Physics in 1972 with minors in biology and chemistry. Ornstein was awarded fellowships during his master's, doctoral, and postdoctoral studies.

Ornstein has published papers in the areas of bio-remediation and structure/function/dynamic relationships of peptides, proteins, nucleic acids, and biodegradative enzymes. He is a member of the American Chemical Society, the Hazardous Materials Control Research Institute, the Protein Society, the Society of Environmental Toxicology and Chemistry, the AAAS, the Richland School District Curriculum Committee, and is listed in Who's Who in Science and Engineering. Ornstein has instructed at York College of C.U.N.Y., Hunter College of C.U.N.Y., Rachel Carson College of S.U.N.Y. at Buffalo, Princeton University, Washington State University, and Rider College. In 1991, he had 11 published papers and 23 abstracts, posters, and presentations.

Dr. Gregory K. Schenter joined the MSRC in 1988 after receiving his Ph.D. in applied physics from Cornell University under the direction of Dr. Richard L. Liboff. At Cornell, his research interests concentrated on classical and quantum transport of charge-carriers in plasmas and semiconductors. Prior to his Ph.D. work he gained experience in both pure and applied physics developing statistical analysis of fission product yield and nuclear decay heat data in addition to the modeling of dye laser physics, reactor physics, and material flows.

His current research interests include the theoretical modeling of chemical dynamics as it occurs in the condensed phase. This work has ranged from the study of fundamental aspects of simplified systems to more realistic systems which may be directly verified by experiment.

He has written over a dozen papers in the areas of nuclear, plasma, semiconductor, and chemical physics.

Dr. Mark A. Thompson recently joined the MSRC as a Senior Research Scientist. Prior to joining the MSRC, he was an Assistant Chemist at Brookhaven National Laboratory. His primary research interests include the theoretical investigation of electron transfer in photosynthetic systems, electronic structure of cofactors and active sites in proteins, and model Hamiltonians useful for studying biological systems. This research includes developing electronic structure and molecular dynamics methods for parallel computing environments.

Thompson received his Ph.D. in physical chemistry in 1990 from the quantum Theory Project at the University of Florida. His research advisor was

Dr. Michael Zerner. He graduated *summa cum laude* from the University of Minnesota with B.S. in biology in 1983. He did his undergraduate research with Dr. Peter Chapman and Dr. Stan Dagley on the catalysis of aromatic compounds by soil bacteria.

While at the University of Minnesota, Thompson was awarded the Alumni Society Scholarship and the Novy Miller Scholarship. In 1984, he received a National Science Foundation Predoctoral Fellowship; and in 1989, a Shell Oil Fellowship. He is a member of the American Chemical Society and Phi Beta Kappa. He has written or contributed to a dozen publications. In 1991, he was an invited speaker at the Sanibel Symposium on Quantum Chemistry and Quantum Biology.

Computing and Information Sciences

Ms. Janice R. Lewis is Acting Manager of the MSRC Computing and Information Sciences program and a Staff Scientist in PNL's Applied Physics Center. Lewis joined PNL in 1977 and has extensive experience in system analysis, system design, and application development. Software development projects for which she has been principle investigator include command and control systems for the U.S. Army as well as emergency response systems for the Hanford site.

Ms. Lewis has managed several software integration projects including the technical management of VISTA, a PNL initiative.

Mr. Gordon A. Anderson is a Senior Research Engineer in PNL's Applied Physics Center and is assigned to the MSRC Computing and Information Science program. He is responsible for the design and development of automated control systems and data acquisition systems, which require extensive use of microprocessor technologies. Some of the systems developed by Anderson as part of Applied Physics Center programs have flown on the NASA space shuttle.

Anderson has analog and digital circuit design experience as well as extensive working knowledge of the Intel 80X86 and Zilog Z80 architectures. He also is experienced in the development of specialized robots and in the application of industrial robots to

Prior to joining PNL, Lewis worked for Washington State University as a software developer and Boeing Computer Services as a systems analyst.

Lewis earned her M.S. in Computer Science from Washington State University and her B.S. in mathematics from the University of Idaho.

specific problems. Anderson has developed a presentation on robotics that has been given to more than 500 students and adults as part of PNL's Sharing Science with Schools Program.

Prior to joining PNL, Anderson worked for Exxon Nuclear on the Laser Isotope Separation Project and at the Fuels Fabrication Plant. During this period, he developed instrumentation and automation equipment.

Anderson received his B.S. in electrical engineering from Washington State University. His publications include a design concept that was featured in EDN Magazine and won a Readers' Choice Award.

Mr. D. Michael DeVaney is a Senior Research Scientist in PNL's Applied Physics Center. Currently, he is assigned to the MSRC Computing and Information Sciences program and is responsible for the Information Management Task for the EMSL Project. This task involves the acquisition, implementation, and integration of a scientific database system and a high-performance hierarchical archival system. The system must be capable of managing up to a few tens-of-terabytes of data with extensive database access and file transfer rates in many tens-to-hundreds of megabytes per second. Object database research support for this task is being conducted at the Oregon Graduate Institute.

DeVaney received his B.S. in mathematics from the University of Washington in 1972. He has since attended Syracuse University and the Massachusetts Institute of Technology. His most recent publications are included in *Proceedings of the DARPA Open*

Dr. Donald R. Jones has been a Staff Engineer in PNL's Applied Physics Center since 1989. Currently, he is assigned to the MSRC Computing and Information Sciences program and is the Visual Engineering Task Leader. His team provides general-purpose computer graphic resources and visual computer tools for all MSRC scientists. His research interests include scientific visualization, graphical user interfaces, data fusion, video technology for the sciences, multimedia for scientific presentations, and scientific animation.

From 1985 to 1989, Jones was the Director of Engineering Animation at Failure Analysis Associates, the largest private U.S. engineering and scientific services consulting firm. While at FaAA, Jones produced more than 30 engineering animations, comprising more than five hours of computer graphics animation. Prior to 1985, Jones was a Research Associate at Brigham Young University's Engineering Computer Graphics Laboratory. He conducted research and development of high-performance graphics devices

Since joining PNL in 1987, DeVaney has specialized in object-oriented and knowledge-based systems, including knowledge engineering and systems analysis, object database, object-oriented programming, rule-based expert systems, and blackboard systems. His research interests also include hypertext/ hypermedia systems and parallel processing. As a member of the Applied Physics Center staff, DeVaney has been a major contributor to the development of a knowledge-based tactical decision aid; a model-based reasoning expert system; a methodology to integrate knowledge-based systems with conventional tactical battle management systems; and an electronic laboratory notebook.

Object-Oriented Database Workshop II, which was held in Dallas, Texas in September; and *Proceedings of the Human Factors Society, 35th Annual Meeting*, which also was held in September.

and state-of-the-art computer graphics software. He is principal author of Dyna-MOVIE.BYU.

Jones received his Ph.D. in engineering at Brigham Young University in 1985. He is a member of the American Society of Mechanical Engineers, the Association of Computing Machines - SIGGRAPH, and the National Computer Graphics Association. Jones also is an associate member of the Institute of Electrical and Electronic Engineers. During 1991, he has been an invited speaker at the DOE Computer Graphics Forum, Boeing Computer Services Workshop on Scientific Visualization, and the 7th Annual Scientific Computing and Automation Conference and Exposition. Currently he is serving as the chairman of the SIG on Scientific Visualization at Hanford. Also, he is a member of the steering committee for the DOE Computer Graphics Forum and a member of SLATEC. He has written or contributed to more than a dozen publications in his field.

Dr. Richard J. Kouzes is a Staff Scientist in the MSRC Computing and Information Sciences program and leader of the Computer Instrumentation and Electronics group, which is conducting experimental research in advanced data acquisition system development and precision atomic mass measurements using FTICR. Kouzes' work includes computer modeling of physical systems to enhance instrument performance so that atomic masses can be measured to higher precision.

Prior to joining PNL, Kouzes was a Senior Research Physicist and lecturer at Princeton University. While at Princeton, he collaborated in two major international solar neutrino experiments, actively pursued research in nuclear techniques for precision atomic mass measurements, and developed several data acquisition systems for nuclear physics applications.

Mr. Richard J. Littlefield leads the Computing and Information Sciences program in High Performance Computing. His current research interests target massively parallel computing for scientific applications and focus on performance modeling and evaluation, algorithm selection, software structure and development tools, and numerical methods. Littlefield's prior work includes the design and construction of several custom parallel computers for image analysis, visualization, and reconstruction of solids scanned with ultrasound using the synthetic aperture focusing technique.

Mr. Marty J. Peterson is a Senior Development Engineer with PNL's Applied Physics Center. Currently he is part of the Systems Engineering Group of the MSRC Computing and Information Sciences program. In this capacity, Peterson is task leader and principal investigator for the design and prototyping of networks to support a large computational-intensive laboratory. Technologies being prototyped include FDDI, with an emphasis on optimization of routed subnetworks and placement of key resources.

Peterson's work is focused on designing a flexible, high-speed hierarchy of networks supporting massively parallel supercomputers, high-speed network file

Kouzes earned his Ph.D. in physics from Princeton University in 1974 under the direction of Professor Rubby Sherr. At Princeton, he built one of the highest-resolution magnetic spectrometers available for nuclear structure research and made a precise measurement of the mass of ^8He . He also developed particle detectors and a data acquisition system for spectroscopic applications.

Kouzes did postdoctoral work at The Indiana University, where he assembled the magnetic spectrometer system and developed the data acquisition system used for the University's Cyclotron Facility. He is a leader in the field of computerized data acquisition and was a founder and past chairman of the IEEE Computer Applications in Nuclear and Plasma Sciences Committee. He is the author of more than 60 papers.

Littlefield has more than 20 years experience in computing, spanning a spectrum from development and support of production software and hardware through research publications. He received his M.S. in computer science and B.S. in mathematics from the University of Washington. Littlefield holds one patent and has written or contributed to more than 20 papers in the areas of parallel computing, analysis of large data sets, computer graphics, and image processing.

server resources, and office workstations. Applications to be supported include computational chemistry codes, large experimental data sets, and graphics and visualization.

Peterson has a strong background in communications design of local area networks, Network Operating Systems, protocols, microwave radio communications, fiber optics technologies, and project management skills. He is a Professional Licensed Electrical Engineer in the state of Washington and is active in the local Institute of Electrical and Electronic Engineers, Sigraph/SigCom, and Professional Engineering societies.

Mr. James Schroeder is a staff engineer with PNL's Applied Physics Center and is currently assigned to the MSRC Computing and Information Sciences program. He leads the Systems Integration and Computing Systems activities for the program. His current technical focus is the integration of diverse computing technologies and resources to meet operational and performance requirements in a timely and cost-effective manner. Schroeder's work experience includes development of real time control and data

collection systems for laboratory, field, and remote sensing applications, data collection and analysis systems for manufacturing industries, system-level software development, and project management. Schroeder earned an M.S. in Electrical Engineering from the University of New Mexico in 1973. His publications are in the areas of distributed data systems and shop floor information systems.

Mr. Kerry D. Steele, a Senior Research Engineer in PNL's Applied Physics Center, is assigned to the MSRC Computing and Information Sciences program. Since joining PNL in 1984, Steele has specialized in the areas of microcomputer data acquisition and control systems, data communications, and information management systems.

In his work with the Applied Physics Center, Steele managed the specification and design of a Command Control Communications and Intelligence system; conceived and led the development of a microprocessor-based interface that links battlefield

reporting systems with information management systems; designed and developed communications software for a multinode integrated network of graphics workstations; and developed software for data acquisition systems for vitrification systems that process contaminated soil. Prior to 1984, Steele served 20 years in the United States Navy, progressing from enlisted technician to commissioned officer. He received his B.S. in electrical engineering from Washington State University in 1984 and has completed graduate-level courses in electrical engineering at WSU.

MSRC Associates

Dr. Gregory E. Arnold

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Biochemistry, Washington State University, 1989 (advisor: Dr. A. Keith Dunker); M.S., Chemistry, Washington State University, 1986; B.S., Chemistry, University of Oregon, 1984.

Dr. Michael B. Bass

Postdoctoral Appointee - Theory, Modeling, and Simulation

Postdoctoral Fellow (Muscular Dystrophy Association Fellowship), Institute of Neuroscience, University of Oregon, 1987-1989 (advisor: Monte Westerfield); Ph.D., Biochemistry, Iowa State University, 1987 (advisor: Herbert J. Fromm); B.S., Biochemistry and Biophysics, Oregon State University, 1982.

Dr. Kenneth M. Beck

Postdoctoral Appointee - Chemical Dynamics and Structure

Postdoctoral Fellow, Chemical Physics, University of Pennsylvania, 1988-1990 (advisor: Marsha I. Lester); Ph.D., Physical Chemistry, University of Illinois-Chicago, 1988 (advisor: Robert J. Gordon); M.S., Chemistry, University of Illinois-Chicago, 1984; B.A., Astronomy/Physics, Northwestern University, 1972.

Dr. David L. Blanchard

Postdoctoral Appointee - Materials and Interfaces

Ph.D., Chemistry, University of Washington, 1989; B.A., Chemistry, Johns Hopkins University, 1983.

Dr. Kirk W. Brown

Postdoctoral Appointee - Chemical Structure and Dynamics

Ph.D., Physical Chemistry, Oregon State, 1991 (advisor: Joseph Nibler); B.S., Chemistry, Idaho State University, 1987.

Dr. Lloyd A. Bumm

Postdoctoral Appointee - Chemical Structure and Dynamics

Postdoctoral Fellow, physics, IBM Zurich Research Laboratory, 1989-1990 (advisor: D. Pohl); Ph.D., Physical Chemistry, Northwestern University, 1991 (advisor: R. P. Van Duyne); M.S., Physical Chemistry, Northwestern University, 1983; B.S., Chemistry, Clarkson University, 1982.

Dr. Ellen L. Chappell

Postdoctoral Appointee - Chemical Structure and Dynamics

Postdoctoral Fellow, Chemical Physics group at SRI International 1990-91 (advisor: David Crosley); Ph.D., Physical Chemistry, University of Oregon, 1990 (advisor: Paul Engelking); B.S., Evergreen State College, 1985.

Professor Walter C. Ermler

Visiting Scientist - Theory, Modeling, and Simulation

Professor of Chemistry and Physics, Stevens Institute of Technology. Ph.D., Chemistry, Ohio State University, 1972; M.S., Chemistry, Ohio State University, 1970; B.S., Chemistry, Northern Illinois University, 1969.

Dr. Thomas J. Godin

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Physics, University of Oregon, 1988; M.S., Physics, University of Oregon, 1984; B.S., Mathematics, University of Rhode Island, 1982.

Dr. Albert T. Hu

Postdoctoral Appointee - Chemical Structure and Dynamics

Postdoctoral Fellow, University of Rochester 1989-91 (advisor: John S. Muentner); Ph.D., Chemical Physics, University of Oregon, 1989.

Dr. Alan G. Joly

NORCUS Postdoctoral Fellow - Chemical Dynamics and Structure

Ph.D., Chemistry, Massachusetts Institute of Technology, 1990 (advisor: Keith Nelson); B.S., Chemistry, University of Rochester, 1985.

Dr. Michael A. Kennedy

Alexander Hollander Postdoctoral Fellow - Macromolecular Structure and Dynamics

Ph.D., Physical Chemistry, University of South Carolina - Columbia, 1989; B.S., Chemistry, Muskingum College, 1984.

Dr. Richard D. Knochenmuss

Senior Research Fellow - Chemical Structure and Dynamics

Staff Scientist, Paul Scherer Institute, Villigen, Switzerland, 1990-91; Postdoctoral Fellow, University of Bern, Switzerland, 1984-89, with Profs. H. Guedel and S. Leutwyler; Ph.D., Washington State University, 1984.

Mlu Kwini

Graduate Student - Chemical Dynamics and Structure

Ph.D. Candidate (1992), Physical Chemistry, University of California-Santa Barbara.

Dr. Maria M. Marino

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Chemistry, Stevens Institute of Technology, 1990; M.S., Chemistry, Stevens Institute of Technology, 1986; B.S., Chemistry, St. Peter's College, 1983.

Dr. Brian L. Maschhoff

Postdoctoral Appointee - Chemical Structure and Dynamics

Postdoctoral Fellow, Rutgers University, 1988-91, with Professor Theodore Madey; Ph.D., Chemistry, University of Arizona, 1988 (Advisor: Professor Neal Armstrong); B.S., Chemistry, University of New Mexico, 1981.

Dr. Robin P. McRae

Postdoctoral Fellow - Theory, Modeling, and Simulation

Ph.D., Chemistry, University of California-Berkeley, 1989 (advisor: A.D.J. Haymet); B.S., Mathematics/Chemistry, University of Puget Sound, 1985.

Dr. Mark D. Paulsen

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Physical Chemistry, University of Wisconsin-Madison, 1989 (advisor: M. Thomas Record); B.S., Chemistry, University of Nebraska-Lincoln, 1982.

Dr. Ronald D. Poshusta

Visiting Scientist - Theory Modeling, and Simulation

Professor of Chemistry, Washington State University. WSU Chemistry faculty since 1967; Ph.D., Chemistry, University of Texas, 1963 (mentor: F. A. Matsen); B.S., Iowa State University, 1957.

Dr. Steven P. Sapers

Postdoctoral Appointee - Chemical Structure and Dynamics

Postdoctoral Fellow, Physical Chemistry, University of Toronto, 1990-1991 (advisor: James Donaldson); Ph.D., Physical Chemistry, University of Colorado, 1989 (advisor: Veronica Vaida); B.A., Chemistry, Brandeis University, 1984.

Dr. Andrew J. Skinner

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Theoretical Solid State Physics, Imperial College, University of London, UK, 1989; B.S., Mathematics, Imperial College, University of London, 1985.

Dr. Scott Smith

Postdoctoral Appointee - Chemical Dynamics and Structure

Ph.D., Chemistry, University of Utah, 1988; B.S., Chemistry, Northern Arizona University, 1983.

Dr. Mark S. Stave

Postdoctoral Appointee - Molecular Science Software

Ph.D., Chemistry, Iowa State University, 1991; B.A., Chemistry, St. Olaf College, 1986.

Dr. Mark O. Trulson

Postdoctoral Appointee - Chemical Dynamics and Structure

National Institutes of Health Fellow, Massachusetts Institute of Technology, 1988-1989 (advisor: Keith Nelson); Ph.D., Chemistry, University of California-Berkeley, 1987 (advisor: Richard Mathies); B.S., Chemistry, Arizona State University, 1981.

Dr. Julia C. White

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Chemistry, Indiana University, 1991 (advisor: Ernest Davidson); B.A., Chemistry, Earlham College, 1986.

Dr. David E. Woon

Postdoctoral Appointee - Theory, Modeling, and Simulation

Postdoctoral Fellow, University of Texas at Arlington, 1990-1991 (advisor: Dennis S. Marynick); Ph.D., Metallurgy, Michigan Technological University, 1989 (advisor: Donald R. Beck); M.S., Physics, MTU, 1987; B.S., Metallurgy, MTU, 1984.

Dr. Sotiris S. Xantheas

Postdoctoral Appointee - Theory, Modeling, and Simulation

Ph.D., Physical Chemistry, Iowa State University, 1990 (advisor: Klaus Ruedenberg); Diploma, Chemical Engineering, National Polytechnic University of Athens, Greece, 1984.

Appendix B

1991 Publications and Presentations

Appendix B

1991 Publications and Presentations

Publications

- G. Atchity, S. S. Xantheas, and K. Ruedenberg, "Potential Energy Surfaces Near Intersections," *J. Chem. Phys.* **95** (3), 1862 (1991).
- K. M. Barkigia, M. Miura, M. A. Thompson, and J. Fajer, "Models of Photosynthetic Chromophores. Molecular Structure of the Bacteriochlorin (2,3,12,13-Tetrahydro-5,10,15,20-Tetraphenyl-porphinato)(pyridine)zinc(II)," *J. Inorg. Chem.* **30**, 2233 (1991).
- S. E. Barlow, T. Dang, and V. M. Bierbaum, "Hydrogen Deuterium Exchange Reactions of Methoxy-Methanol Clusters," *J. Amer. Chem. Soc.* **112**, 6832 (1991).
- A. R. Burns, T. M. Orlando, D. R. Jennison, and E. B. Stechel, "State-Resolved Study of Coadsorption Effects on the Stimulated Dissociation of NO₂ on Pt(111)," *J. Vac. Sci. Technol.* **A9**, 1174 (1991).
- D. W. Chandler, J. W. Thoman, Jr., and W. P. Hess, "Photofragment Imaging: The Photodissociation of bromomethane, bromoethane and bromoethanol," *Inst. Phys. Conf. Ser.* **114**, 355 (1991).
- D. C. Chatfield, R. S. Friedman, D. G. Truhlar, B. C. Garrett, and D. W. Schwenke, "Global Control of Suprathreshold Reactivity by Quantized Transition States," *J. Amer. Chem. Soc.* **113**, 486 (1991).
- D. M. Chipman, I. Carmichael, and D. F. Feller, "Molecular Orbital Studies of Hyperfine Coupling Constants in the H₂CN and H(HO)CN Radicals," *J. Phys. Chem.* **95**, 4702 (1991).
- J. B. Cushing, D. M. DeVaney, D. F. Feller and D. Maiers, "Data Base Support for Computational Chemistry," *Proceedings of the DARPA Open Object-oriented Data Base Workshop II*, Texas Instruments, Dallas, Texas, 1991.
- T. H. Dunning, Jr., S. S. Xantheas, A. C. Hess, D. F. Feller, R. A. Kendall and R. L. Ornstein, "Modeling Molecular Processes in the Environment," in *Proceedings of the First Energy Research Power Supercomputer Users Symposium*, U. S. Department of Energy, Gaithersburg, Maryland, May 21-22, 1991.
- C. G. Edmonds, J. A. Loo, R. R. Ogorzalek-Loo, R. D. Smith, "Techniques in Protein Chemistry II", J. J. Villafranca, ed., p. 487-495, Academic Press, 1991.
- D. F. Feller, and E. R. Davidson, "Theoretical Approaches to ESR Spectroscopy," in Vol. 3 of *Theoretical Models of Chemical Bonding*, Z. B. Maskis, ed., Springer Verlag, 1991.
- J. Fidy, G. R. Holtom, K. G. Paul, and J. M. Vanderkooi, "Binding of Naphthohydroxamic Acid to Horseradish-Peroxidase Monitored by Zinc Mesoporphyrin Fluorescence Line Narrowing," *J. Phys. Chem.* **95**, 4364 (1991).
- B. C. Garrett, and C. F. Melius, "Variational Transition State Theory Calculations of Tunneling Effects on Concerted Hydrogen Motion in Water Clusters and Formaldehyde/Water Clusters," *Theoretical and Computational Models for Organic Chemistry*, S. J. Formosinho et al., ed., p. 35, Kluwer Academic Publishers, Netherlands, 1991.

- B. C. Garrett, and D. G. Truhlar, "Critical Tests of Variational Transition State Theory and Semiclassical Tunneling Methods for Hydrogen and Deuterium Atom Transfer Reactions and Use of the Semiclassical Calculations to Interpret the Overbarrier and Tunneling Dynamics," *J. Phys. Chem.*, **95**, 10374 (1991).
- R. J. Harrison, and R. A. Kendall, "A Parallel Version of ARGOS: A Distributed Memory Model for Shared Memory UNIX Computers," *Theoretica Chimica Acta* **79**, 337 (1991).
- M. Häser, M. W. Feyereisen, and J. Almlöf, "Exploiting Non-Abelian Point-Group Symmetry in Direct Two-Electron Integral Transformations," *Theoret. Chim. Acta* **79**, 115 (1991).
- S. Hunt, and A. Schur, "The User-Oriented Evaluation Process: A Process for Preserving User Needs During Iterative System Test and Evaluation," In *Proceedings of the Human Factors Society 35th Annual Meeting*, Vol. II, September 1991.
- M. A. Kennedy, R. R. Vold, and R. L. Vold, "A Solid State Deuterium NMR Study of Internal Motion in p-Nitroaniline," *J. Magn. Reson.*, **91**, 301-315 (1991).
- M. A. Kennedy, R. L. Vold, and R. R. Vold, "Stepped Frequency NMR Spectroscopy," *J. Magn. Reson.*, **92**, 320-331 (1991).
- J. P. LaFemina, and G. K. Schenter, "Twisted Intramolecular Charge Transfer and the Torsional Potential Function of Dimethylaminobenzonitrile," *J. Chem. Phys.* **94**, 7558 (1991).
- R. J. Littlefield, and K. J. Maschhoff, "Choosing Processor Array Configuration by Performance Modeling for a Highly Parallel Linear Algebra Algorithm," In *Proceedings of the Sixth Distributed Memory Computing Conference*, Portland, Oregon, 1991.
- J. Liu, M. W. Feyereisen, J. Almlöf, C.M. Rohlfing, and S. Saebo, "Ab Initio Studies on Substituted Diamond-like Carbon Clusters," *Chem. Phys. Letters* **183**, 478 (1991).
- J. A. Loo, C. G. Edmonds, and R. D. Smith, "Tandem Mass Spectrometry of Very Large Molecules: Serum Albumin Sequence Information from Multiply Charged Ions formed by Electrospray Ionization," *Anal. Chem.*, **63**, 2488-2499 (1991).
- J. A. Loo, R. R. Ogorzalek-Loo, H. R. Udseth, C. G. Edmonds, and R. D. Smith, "Solvent-Induced Conformational Changes of Polypeptides Probed by Electrospray Ionization Mass Spectrometry," *Rapid Commun. Mass Spectrom.*, **5**, 101-105 (1991).
- G. C. Lynch, R. Steckler, D. W. Schwenke, A.J.C. Varandas, D. G. Truhlar, and B. C. Garrett, "Use of Scaled External Correlation, a Double Many-Body Expansion, and Variational Transition State Theory to Calibrate a Potential Energy Surface for FH_2 ," *J. Chem. Phys.*, **94**, 7136 (1991).
- G. A. Natanson, B. C. Garrett, T. N. Truong, T. Joseph, and D. G. Truhlar, "The Definition of Reaction Coordinate for Reaction-Path Dynamics," *J. Chem. Phys.* **94**, 7875 (1991).
- R. R. Ogorzalek-Loo, H. R. Udseth, and R. D. Smith, *J. Phys. Chem.*, **95** 6412-6415 (1991).
- T. M. Orlando, A. R. Burns, E. B. Stechel, and D. R. Jennison, "Resonance Ionization Detection of Neutral $\text{O}(^3\text{P}_j)$ Produced from Stimulated Surface Processes," *J. Vac. Sci. Technol.* **A9**, 1769 (1991).
- T. M. Orlando, A. R. Burns, E. B. Stechel, and D. R. Jennison, "Quantum-Resolved Studies of Electron-Stimulated Reactions on Adsorbate Covered Pt(111) Surfaces," *Nucl. Instr. Meth. in Phys. Res.* **B58**, 477 (1991).

- R. L. Ornstein, "Rational Redesign of Biodegradative Enzymes for Enhanced Bioremediation: Overview and Status Report for Cytochrome P450," in *Proceedings of National Research and Development Conference on the Control of Hazardous Materials*, pp. 314-319, Hazardous Materials Control Research Institute, Greenbelt, Maryland, 1991.
- J. M. Papanikolas, J. R. Gord, N. E. Levinger, D. Ray, V. Vorsa, and W. C. Lineberger, "Photodissociation and Geminate Recombination Dynamics of I_2^- in Mass-Selected $I_2^-(CO_2)_n$ Cluster Ions," *J. Phys. Chem.* **95**, 8028 (1991).
- R. L. Ornstein, "Why Timely Bioremediation of Synthetics May Require Rational Enzyme Redesign: Preliminary Report on Redesigning Cytochrome P450cam for Trichloroethylene Dehalogenation," *On Site Bioreclamation: Processes for Xenobiotic and Hydrocarbon Treatment*, R. E. Hincee and R. F. Olfenbuttel, eds., pp. 509-514, Butterworth-Heinemann, Boston, Massachusetts, 1991.
- R. L. Ornstein, M. B. Bass, M. D. Paulsen, and G. E. Arnold, "Rational Redesign of the Biodegradative Enzyme Cytochrome P450cam: A Progress Report on Engineering Specificity for Halogenated Hydrocarbon Substrates," in *Proceedings of HAZMACON 91*, T. Bursztynsky, ed., Vol. II, pp. 470-480, JT Litho, Concord, California, 1991.
- V. Parasuk, M. W. Feyereisen, and J. Almlöf, "The Electronic and Molecular Structure of C_{18}^- ," *J. Amer. Chem. Soc.* **113**, 1049 (1991).
- M. D. Paulsen, M. B. Bass, and R. L. Ornstein, "Analysis of Active Site Motions from a 175 psec Molecular Dynamics Simulation of Camphor-Bound Cytochrome P450cam," *J. Biomolecular Structure & Dynamics* **9**, 187-203 (1991).
- M. D. Paulsen, and R. L. Ornstein, "A 175 psec Molecular Dynamics Simulation of Camphor-Bound Cytochrome P450cam," *PROTEINS: Structure, Function, and Genetics* **11**, 184-204 (1991).
- G. Reiser, O. Dopfer, R. Lindner, G. Henri, K. Müller-Dethlefs, and S. D. Colson, "A New Approach to Vibrational Spectroscopy of Ion Clusters: The 'Zero Kinetic Energy (ZEKE)' Photoelectron Spectrum of the Phenol-Water Complex," *Chem. Phys. Lett.* **189**, 1-4 (1991).
- A. L. Rockwood, M. Busman, R. D. Smith "Coulombic Effects in the Dissociation of Large Highly Charged Ions," *Int. J. Mass Spectrom. Ion. Proc.* **111**, 103-129 (1991).
- G. K. Schenter, and C. B. Duke, "Theory of Photoinduced Twisting Dynamics in Polar Solvents: Application to Dimethylaminobenzonitrile in Propanol at Low Temperatures," *Chem. Phys. Lett.* **176**, 563 (1991).
- J. O. Schroeder, E. W. Pearson, J. T. Thomas, and J. W. Brothers, Computing and Information Sciences Preliminary Engineering Design Study, PNL-7670, Pacific Northwest Laboratory, Richland, Washington, 1991.
- A. Schur, D. F. Feller, D. M. DeVancy, J. J. Thomas, and M. L. Yim, "EASI: An Electronic Assistant for Scientific Investigation," *Proceedings of the Human Factors Society 35th Annual Meeting*, Vol. 1, September 1991.
- S.-K. Shin, Y. Chen, S. Nickolaisen, S. W. Sharpe, R. A. Beaudet, and C. Wittig, "Photoinitiated Reactions in Weakly-Bonded Complexes," *Advances in Photochemistry*, Vol. 16, John Wiley & Sons, Inc., 1991.
- R. D. Smith, J. A. Loo, C. G. Edmonds, and H. R. Udseth, *Analytical Applications of Spectroscopy II*, A.M.C. Davies, and C. S. Creaser, eds., pp. 149-164, Royal Society of Chemistry (1991).
- R. D. Smith, H. R. Udseth, C. J. Barinaga, and C. G. Edmonds, "Instrumentation for High-Performance Capillary Electrophoresis-Mass Spectrometry," *J. Chromatogr.*, **559**, 197-208 (1991).
- M. M. Szczesniak, R. A. Kendall, and G. Chalasinski, "Ab Initio Study of the Nonadditive Effects in the Trimer of Ammonia," *Journal of Chemical Physics* **95**, 5169 (1991).

- T. S. Tenforde, "Biological Effects of Static Magnetic Fields," *Int. J. Appl. Electromag. Mat.* **1**, 157-165.
- T. S. Tenforde, "Biological Interactions of Extremely-Low-Frequency Electric and Magnetic Fields," *Bioelectrochem. & Bioenergetics* **25**, 1-17 (1991) [section of *J. Electroanal. Chem.* **320** (1991)].
- Y. Thiel, V. A. Walters, K. B. Wiberg, and S. D. Colson, "High-Resolution Infrared Analysis of the ν_{17} Band of Pyridine," *Int. J. Quant. Chem.* **39**, 423-436 (1991).
- M. A. Thompson, M. C. Zerner, and J. Fajer, "A Theoretical Examination of the Electronic Structure and Excited State of the Bacteriochlorophyll-b Dimer from Rhodospseudomonas viridis," *J. Phys. Chem.* **95**, 5693 (1991).
- P. Valtazanos, S. T. Elbert, S. S. Xantheas, and K. Ruedenberg, "The Ring Opening of Cyclopropylidene to Allene: Global Features of the Reaction Surface," *Theor. Chim. Acta* **78**(5/6), 287 (1991).
- J. M. Van Doren, S. E. Barlow, C. H. DePuy, and V. M. Bierbaum, "Exchange Reactions of $^{18}\text{O}^-$ with Some Di- and Triatomic Molecules," *International J. Mass Spectrom. Ion Processes*, **109**, 305 (1991).
- V. A. Walters, C. M. Hadad, Y. Thiel, S. D. Colson, K. M. Wiberg, P. M. Johnson and J. Foresman, "Assignment of the A State in Bicyclobutane. The Multiphoton Ionization Spectrum, and Calculations of Transition Energies," *J. Amer. Chem. Soc.* **113**, 4782-4791 (1991).
- S. S. Xantheas, G. J. Atchity, S. T. Elbert, and K. Ruedenberg, "Potential Energy Surfaces of Ozone. I," *J. Chem. Phys.* **94**(12), 8054 (1991).
- S. S. Xantheas, S. T. Elbert, and K. Ruedenberg, "The Ring Opening of Cyclopropylidene to Allene: Key Features of the Accurate Reaction Surface," *Theor. Chim. Acta* **78**(5/6), 365 (1991).
- S. S. Xantheas, P. Valtazanos, and K. Ruedenberg, "The Ring Opening of Cyclopropylidene to Allene and the Isomerization of Allene: *Ab Initio* Interpretation of the Electronic Rearrangements in Terms of Quasiatomic Orbitals," *Theor. Chim. Acta* **78** (5/6), 327 (1991).

Manuscripts Not Yet Published

G. E. Arnold, and R. L. Ornstein, "T4 Lysozyme Hinge-Bending from a 2 Nanosecond Molecular Dynamics Simulation Analysis" (in preparation).

R. A. Bair, ed., "Graphical User Interfaces for Computational Chemistry," PNL-SA-18847, CONF-9009303, Pacific Northwest Laboratory, Richland, Washington (in press).

K. M. Barkigia, M. A. Thompson, R. K. Pandey, K. M. Smith, M.G.H. Vicente, and J. Fajer, "Conformational Flexibility of Chlorine Macrocycles. Structural Determinations and Theoretical Calculations of a Benzochlorin, a meso-Dimethyl Chlorine and a Nickel Chlorine," *New J. Chem.* (in press).

M. B. Bass, D. Filipovic, S. G. Sligar, and R. L. Ornstein, "Altering the Coupling Between Cytochrome P450cam: A Theoretical Prediction with Experimental Evidence" (in preparation).

M. B. Bass, D. F. Hopkins, W. A. Jaquysh, and R. L. Ornstein, "A Method for Determining the Positions of Polar Hydrogens added to a Protein Structure that Maximizes Protein Hydrogen Bonding," *PROTEINS: Structure, Function, and Genetics* (in press).

M. B. Bass, M. D. Paulsen, and R. L. Ornstein, "Molecular Dynamics Simulations of Norcamphor-Cytochrome P450cam and Mutations of P450cam Designed to Alter the Product Specificity," In *Proceedings of the 7th International Conference on Biochemistry and Biophysics of Cytochrome P450: Structure & Function, Biotechnological & Ecological Aspects*, A. I. Archakov, ed., Moscow, USSR (in press).

M. B. Bass, M. D. Paulsen, and R. L. Ornstein, "Substrate Mobility in a Deeply Buried Active Site: Analysis of Norcamphor Bound to Cytochrome P450cam as Determined by a 201 psec Molecular Dynamics Simulation," *PROTEINS: Structure, Function, and Genetics* (in press).

M. B. Bass, and R. L. Ornstein, "Comparison of Cytochrome P450cam Dynamics with Bound L- and D-Norcamphor" (in preparation).

J. A. Braatz, M. D. Paulsen, and R. L. Ornstein, "3 nsec Molecular Dynamics Simulation of the Protein Ubiquitin and Comparison with X-ray Crystal and Solution NMR Structure," *J. Biomol. Struct. Dynam.* (in press).

F. J. Brockman, and R. L. Ornstein, "Enhanced Bioremediation of Subsurface Contamination: Enzyme Recruitment and Redesign," In *Proceedings of Hazardous Materials Control '91*, Hazardous Materials Control Research Institute, Greenbelt, Maryland (in press).

A. R. Burns, E. B. Stechel, D. R. Jennison, and T. M. Orlando, "The Effect of Coadsorbed Atomic Oxygen on the Electron-Stimulated Desorption of NO from Pt(111)," *Phys. Rev. B* (in press).

D. E. Buska, and M. J. Peterson, *Study of Network Modeling Tools*, Draft Internal Report, September 1991.

M. Busman, A. L. Rockwood, and R. D. Smith, "Activation Energies for Gas-Phase Dissociations of Multiply Charged Ions from Electrospray Ionization Mass Spectrometry," *J. Phys. Chem.* (in press).

R. J. Cave, S. S. Xantheas, and D. Feller, "Exploiting Regularity in Systematic Sequences of Wavefunctions Which Approach the Full CI Limit," *Theor. Chim. Acta* (in press).

C. G. Edmonds, J. A. Loo, R. R. Ogorzalek-Loo, H. R. Udseth, C. J. Barinaga, and R. D. Smith. *Biochemical Soc. Trans.* (in press).

- D. F. Feller, "Application of Systematic Sequences of Wavefunctions to the Water Dimer" (in preparation).
- M. W. Feyereisen, and R. A. Kendall, "An Efficient Implementation of the Direct-SCF Algorithm on Parallel Computer Architectures," *Theoretica Chimica Acta*. (submitted).
- M. W. Feyereisen, M. Gutowski, J. Simons, and J. Almlöf, "Relative Stabilities of Fullerene, Cumulene, and Polyacetylene Structures for C_n ; $n = 18 - 60$," *J. Chem. Phys.* (in press).
- M. W. Feyereisen, J. Nichols, J. Oddershede, and J. Simons, "Direct Atomic-Orbital Based TDHF Calculation of Frequency-Dependent Polarizabilities," *J. Chem. Phys.* (submitted).
- D. Filipovic, M. B. Bass, R. L. Ornstein, and S. G. Sligar, "A Possible Role for F87 of Cytochrome P450cam in Substrate Access" (in preparation).
- D. Filipovic, M. D. Paulsen, P. J. Loida, S. G. Sligar, and R. L. Ornstein, "Ethylbenzene Hydroxylation by Cytochrome P450cam" (submitted).
- D. Filipovic, M. D. Paulsen, J. D. Ropp, R. L. Ornstein, and S. G. Sgar, "Prediction of the Three-Dimensional Structure of Cytochrome P450lin Using Homology Modeling and Molecular Dynamics" (in preparation).
- B. G. Gladstone, M. B. Bass, and R. L. Ornstein, "Effect of Crystal Packing Forces on the Structure and Mobility of Proteins" (in preparation).
- B. G. Gladstone, and R. L. Ornstein, "2 Nanosecond Molecular Dynamics Simulation of the Protein Erabutoxin b" (in preparation).
- A. C. Hess, and P. F. McMillan, "Ab Initio Methods in Geochemistry and Mineralogy," *Advances in Electronic Theory* 2, T. H. Dunning, Jr., ed., Jai Press (in press).
- A. C. Hess, and V. R. Saunders, "Periodic Ab Initio Hartree-Fock Calculations of the Low Symmetry Mineral Kaolinite," *J. Phys. Chem.* (in press).
- W. P. Hess, D. W. Chandler, and J. W. Thoman, Jr., "Photofragment Imaging: The 205 nm Photodissociation of CH_3Br and CD_3Br ," *Chem. Phys.* (submitted).
- S. A. Kafafi, J. P. LaFemina, and G. K. Schenter, "The Electronic Structure and Conformation of Dimethylaminobenzonitrile," *Structural Chemistry* (in press).
- R. A. Kendall, "Electron Affinities of the First Row Atoms Revisited," *Journal of Chemical Physics* (to be submitted).
- L. B. Knight, Jr., T. Arrington, and D. F. Feller, "An ESR and Ab initio Study of the Methanol Radical Cation" (in preparation).
- L. B. Knight, Jr., and D. F. Feller, "An Experimental and Theoretical Study of the Hyperfine Properties of COH" (in preparation).
- L. B. Knight, Jr., D. F. Feller, and E. R. Davidson, "An Experimental and Theoretical Study of $BC_3(4\Sigma)$ " (in preparation).
- R. Knochenmuss and S. Leutwyler, "Structures and Vibrational Spectra of Water Clusters in the SCF Approximation," *J. Chem. Phys.* (submitted).

- M. Kwini, M. J. Iedema, J. P. Cowin, and T. L. Gilton, "Probing Femtosecond Solvation Dynamics at the Condensed Phase-Vacuum Interface," *J. Phys. Chem.* (submitted).
- J. A. Loo, C. G. Edmonds, H. R. Udseth, R. R. Ogorzalek-Loo, and R. D. Smith, "Electrospray Ionization Mass Spectrometry and Tandem Mass Spectrometry of Large Biomolecules," *Exp. Mass Spectrom.* (in press).
- J. A. Loo, R. R. Ogorzalek-Loo, K. J. Light-Wahl, C. G. Edmonds, and R. D. Smith, "Multiply Charged Negative Ions by Electrospray Ionization from Polypeptides and Proteins," *Anal. Chem.* (in press).
- R. J. Littlefield, "Investigating the Performance of Parallel Eigensolvers for Large Processor Counts, submitted to *Theoretica Chimica Acta* (submitted).
- H. P. Luthi, J. E. Mertz, M. W. Feyereisen, and J. Almlöf, "A Coarse-Grain Parallel Implementation of the Direct SCF Method," *J. Comp. Chem.* (in press).
- M. I. McCarthy, and R. B. Gerber, K. A. Trentelman, P. Strupp, D. H. Fairbrother, P. C. Stair, and E. Weitz, "Photodissociation Dynamics of CH₃I Adsorbed on MgO (100): Theory and Experiment," *J. Chem. Phys.* (in press).
- M. I. McCarthy, and A. C. Hess, "Periodic Hartree-Fock Studies of the Properties of the C₁₂/MgO Interface," *J. Chem. Phys.* (in press).
- V. S. Melissas, D. G. Truhlar, D. G. Truhlar, "Optimized Calculations of Reaction Paths and Reaction-Path Functions for Chemical Reactions," *J. Chem. Phys.* (in press).
- T. M. Orlando, A. R. Burns, D. R. Jennison, and E. B. Stechel, "Electronically-Stimulated Adsorbate Dissociation in the Presence of an Electronegative Coadsorbate: (NO₂ + O) on Pt(111)," *Phys. Rev. B* (in press).
- I. Panas, J. Almlöf, and M. W. Feyereisen, "Ab Initio Methods for Extended Systems," *Intern. J. Quantum Chem.* (in press).
- M. D. Paulsen, and R. L. Ornstein, "Molecular Dynamics Simulation of Substrate-free Cytochrome P450cam and Comparison with Camphor-Bound Simulation" (submitted).
- M. D. Paulsen, and R. L. Ornstein, "Molecular Dynamics Simulations of Camphor-bound Cytochrome P450cam with Active Site Mutation Tyr 96--> Phe" (in preparation).
- M. D. Paulsen, and R. L. Ornstein, "Predicting Product Specificity and Coupling for Catalysis by Cytochrome P450cam" (in preparation).
- M. D. Paulsen, and R. L. Ornstein, "Why X-ray Crystal Structure of Thiocamphor-bound Cytochrome P450cam is Inconsistent with Observed Product Specificity: An Explanation Based on Molecular Dynamics Simulations" (in preparation).
- M. J. Peterson, J. O. Schroeder, W. S. Eggers, D. R. McGee, and J. R. Lewis, *EMSL Interim Facility Proposed Computing Environment*, Draft Internal Report, July 1991.
- A. L. Rockwood, M. Busman, and R. D. Smith, "Thermally Induced Dissociation of Ions in Electrospray Mass Spectrometry," *Rapid Commun. Mass Spectrometry* (in press).
- S. P. Sapers, and W. P. Hess, "Photodissociation of BrCHCH₂OH and ICHCH₂OH: Formation and Characterization of OH (X²Π) Products," *J. Chem. Phys.* (to be submitted).

G. K. Schenter, and J. P. LaFemina, "Semiconductor Surface and Interface Dynamics from Tight-binding Molecular Dynamics Simulations," *J. Vac. Sci. Technol.* (in press).

M. R. Schneider, C. P. Dehnhostel, T. L. Gilton, and J. P. Cowin, "Photoelectron Induced Surface Chemistry: Methyl Chloride on GaAs(100)," *J. Appl. Phys.* (submitted).

J. B. Smith, L. R. Miesbauer, J. Leeds, D. L. Smith, J. A. Loo, R. D. Smith, and C. G. Edmonds, "Mass Spectrometric Analysis of Lens B Crystallines," *Int. J. Mass Spectrom. Ion. Proc.* (in press).

R. D. Smith, J. A. Loo, and C. G. Edmonds, "The Analysis of Biomolecules by Electrospray Ionization-Mass Spectrometry and Tandem Mass Spectrometry," *Clinical Mass Spectrometry*, D. Dm. Desiderio, ed., Plenum Press, New York (in press).

R. D. Smith, J. A. Loo, R. R. Ogorzalek-Loo, M. Busman, and H. R. Udseth, "Principles and Practice of Electrospray Ionization-Mass Spectrometry for Large Polypeptides and Proteins," *Mass Spectrom. Rev.* (submitted).

R. S. Smith, S. D. Colson, T. H. Ballinger, and J. T. Yates, "Catalytic Decomposition of 1,1,1-trichloroethane Over Alumina" (in preparation).

R. S. Smith, S. D. Colson, E. E. Gulcicek, and L. D. Pfefferle, "Methane Oxidative Coupling in a Quenched Reactor: Intermediates and Olefin Products," *J. Chem. Phys.* (submitted).

A. Sokalski, J. Lai, N. Luo, S. Sun, M. Shibata, R. L. Ornstein, and R. Rein, "Ab Initio Study of the Electrostatic Multipole Nature of Torsional Potentials: Hydrogen Peroxide and Methyl Disulfide (CH₃SSCH₃)," *Intern. J. Quant. Chem.: Quant. Biol. Sym.* (in press).

W. A. Sokalski, M. Shibata, R. L. Ornstein, and R. Rein, "Correcting Deficiencies of Atomic Charges Using the Cumulative Atomic Multipole Approach" (submitted).

W. A. Sokalski, M. Shibata, R. L. Ornstein, and R. Rein, "Point Charge Representation of Multicenter Multipole Moments in Calculation of Electrostatic Properties" (submitted).

K. D. Steele, *Comments on the LabVIEW Graphical Development Environment*, Internal Working Report, August 1991.

S. Sun, N. Luo, R. L. Ornstein, and R. Rein "Protein Secondary and Tertiary Structure Prediction By a Reduced Representation Method: Test Case for Seven Small Protein" (in preparation).

S. Sun, N. Luo, R. L. Ornstein, and R. Rein "Protein Structure Prediction: Reduced Representation Approach" (in preparation).

T. S. Tenforde, "Interaction Mechanisms and Biological Effects of Static Magnetic Fields, *Automedica* (in press).

M. A. Thompson, "Theoretical Description of Mutant Photosynthetic Reaction Centers with Altered Photophysical and Electron Transfer Properties" (in preparation).

M. A. Thompson, and J. Fajer, "A Theoretical Model for the Bacteriochlorophyll-g Primary Donor from the Photosynthetic Bacterium *Hellobacterium chlorum*. A Dimeric Structure for P790," *J. Phys. Chem.* (to be submitted).

M. A. Thompson and J. Fajer, "Are Dimers a Fundamental Structure in Photosynthesis?" (in preparation).

- M. A. Thompson, and M. C. Zerner, "A Theoretical Examination of the Electronic Structure and Spectroscopy of the Photosynthetic Reaction Center from *Rhodospseudomonas viridis*," *J. Am. Chem. Soc.* (in press).
- L. A. Thomson, J. A. Braatz, and R. L. Ornstein, "Comparison of Molecular Dynamics Simulations for the Protein Ubiquitin with and without an Explicit 7 Å Shell of Water" (in preparation).
- M. Tomonari, N. Ookubo, T. Takada, M. W. Feyereisen, and J. Almlöf, "Ab Initio Calculations of Nonlinear Optical Properties of Organic Molecules. I: Methods and Applications to Benzene Derivatives" *J. Chem. Phys.* (submitted).
- A. C. Wang, M. A. Kennedy, B. R. Reid, and G. P. Drobny, "A Solid-State Deuterium NMR Study of the Purine Motion in a 12 Base-Pair Oligoribonucleotide," *Biochemistry* (submitted).
- A. C. Wang, M. A. Kennedy, B. R. Reid, and G. P. Drobny, "A Solid-State Deuterium NMR Study of the Relaxation in a 12 Base-Pair Oligoribonucleotide," *J. Am. Chem. Soc.* (submitted).
- X. C. Wang, J. Nichols, M. W. Feyereisen, M. Gutowski, J. Boatz, A. D. Haymet, and J. Simons, "Ab Initio Quantum Chemistry Study of Formamide-Formamidic Acid Tautomerization," *J. Chem. Phys.* (submitted).
- B. E. Winter, K. J. Light-Wahl, and R. D. Smith, "Gas Phase Proton Transfer Reactions Involving Multiply Charged Cytochrome c Ions and Water under Thermal Conditions," *J. Amer. Soc. Mass Spectrom.* (submitted).
- S. S. Xantheas, and T. H. Dunning, Jr., "Solvation of the Hydride Ion: I. Critical Points on the H-+H₂O-+HO-+H₂ Potential Energy Surface" (in preparation).
- S. S. Xantheas, and T. H. Dunning, Jr., "Solvation of the Hydride Ion: II. Optimal Structures of the Clusters H-(H₂O)_n, n=1-4" (in preparation).
- Y. P. Zeng, S. W. Sharpe, C. Wittig, and R. A. Beaudet, "Infrared Spectroscopy of CO₂-D(H)Br and Its Molecular Structure," *J. Chem. Phys.* (submitted).

Presentations

- J. G. Boyle, L. D. Pfefferle, E. E. Gulicek, and S. D. Colson, "Laser Driven Electron Ionization for a VUV Photoionization Time-Of-Flight Mass Spectrometer," *Rev. Sci. Instr.* **62**, 323-333 (1991).
- A. R. Burns, T. M. Orlando, E. B. Stechel, and D. R. Jennison, "State-Resolved Angular Distributions of Neutral Products of Electron-Stimulated Surface Processes," presented at 38th National American Vacuum Society Symposium, Seattle, Washington, November 11-15, 1991.
- A. R. Burns, T. M. Orlando, E. B. Stechel, and D. R. Jennison, "Surface Reactions by Electronic Transitions: NO₂ from O₂ and NO on Pt(111)," presented at New Mexico Chapter of the American Vacuum Society, 27th Symposium, April, 1991.
- S. D. Colson, "ZEKE Photoelectron Spectra of Hydrogen Bonded Clusters," presented at the Chemical Physics Institute Meeting, Coos Bay, Oregon, September 1991.
- S. D. Colson, "Understanding Molecular Processes in Natural & Contaminated Environments," presented at Tulane University, New Orleans, Louisiana, September 1991.
- S. D. Colson, Chairman and Organizer, 21st International Symposium on Free Radicals, Williamstown, Massachusetts, August 1991.
- J. P. Cowin, "Femtosecond Solvation Dynamics at Surfaces," presented at University of Colorado, February 1991.
- J. P. Cowin, "Laser Probes of Surface Dynamics," presented at Oregon Graduate Center, April 1991.
- J. P. Cowin, "Femtosecond Solvation Processes at Surfaces," presented at American Chemical Society Northwest Regional Meeting, La Grande, Oregon, June 1991.
- J. P. Cowin, "Femtosecond Solvation Processes at Surface," presented at 2nd Symposium on Photoinduced Charge, University of Rochester, July 1991.
- J. P. Cowin, "Photodissociation Probes of Femtosecond Solvation Dynamics," presented at Dynamics of Gas-Surface Interactions Symposium, Liverpool, England, July 1991.
- J. P. Cowin, "Femtosecond Solvation Dynamics of CH₃Cl and Cl⁻ at Multilayer Surfaces," presented at 4th Chemical Congress of North America (ACS), New York City, August 1991.
- J. P. Cowin, invited participant at Mini-Workshop on Chemical Dynamics at the Vacuum-Solid Interface, Combustion Dynamics Facility, Berkeley, California, August 1991.
- J. P. Cowin, "Photodissociation Probes of Femtosecond Solvation Dynamics," presented at International Laser Science Symposium, Monterey, California, September 1991.
- B. C. Garrett, "Variational Transition State Theory Calculations of Tunneling Effects on Concerted Hydrogen Motion in Water Clusters and Formaldehyde/Water Clusters," poster at West Coast Theoretical Chemistry Conference, NASA Ames Research Center, Moffett Field, California, March 27-29, 1991.
- B. C. Garrett, "Solvent Effects on Activated Chemical Reactions," poster at Conference on the Dynamics of Molecular Collisions, Lake George, New York, July 7-12, 1991.

W. P. Hess, "Ion Imaging of Photofragments: The Photodissociation of CH_3Br , $\text{CH}_3\text{CH}_2\text{Br}$, and $\text{BrCH}_2\text{CH}_2\text{OH}$," presented at Oregon State University Chemistry Symposium, Corvallis, Oregon, February 1991.

W. P. Hess, "The Photodissociation of $\text{BrCH}_2\text{CH}_2\text{OH}$ by Laser-Induced-Fluorescence and the Photodissociation of Bromomethane, Bromoethane, and Bromoethanol," presented at Telluride Workshop on Reactivity: Effects of Intermolecular Interactions, Telluride, Colorado, July 1991.

D. R. Jennison, A. R. Burns, E. B. Stechel, and T. M. Orlando, "Auger Spectroscopy and Electronically-Stimulated Surface Processes," presented at International Workshop on Auger Spectroscopy, Stockholm, Sweden, September 1991.

R. D. Knochenmuss, "Isomerism and Phase Transitions in Solvent Clusters," presented at XII Conference on Molecular Energy Transfer, Nijmegen, Netherlands, July 1991.

M. Kwini, M. Iedema, and J. P. Cowin, "Femtosecond Solvation Processes at Surfaces," poster session presented at American Physical Society National Meeting, Cincinnati, Ohio, March 1991.

R. S. McDowell, "Modern High-Resolution Infrared Spectroscopy," presented at Washington State University Physics Seminar, Pullman, Washington, September 1991.

T. M. Orlando, "Quantum-Resolved Stimulated Surface Chemistry," presented at the Division of Condensed Matter Physics, Section on Surface Dynamics, American Physical Society Meeting, Cincinnati, Ohio, March 1991.

D. Ray, "Ultrafast Time-Resolved Dynamics in Cluster Ions," presented at Cluster Workshop, Telluride Summer Research Center, Telluride, Colorado, July 1991.

D. Ray, "Ultrafast Time-Resolved Dynamics in Cluster Ions," presented at Tulane University, New Orleans, Louisiana, September 1991.

G. E. Arnold, and R. L. Ornstein, "A Molecular Dynamics Analysis of T4 Lysozyme," Fifth Annual Symposium of the Protein Society, June 1991, Baltimore, Maryland.

G. E. Arnold, and R. L. Ornstein, "Molecular Dynamics Simulations of T4 Lysozyme," *Cellular Biochemistry*, Supplement 15G, UCLA Keystone Conference on Protein Folding, Structure and Function, Keystone, Colorado, April 1991.

G. J. Atchity, S. S. Xantheas, and K. Ruedenberg, "Potential Energy Surfaces near Intersections," poster presentation at the International Symposium *Ab Initio* Methods in Quantum Chemistry, Iowa State University, Ames, Iowa, May 9-11, 1991.

R. A. Bair, "Eigensystems in Quantum Chemistry," presented at the Parallel Eigensolver Workshop, Pacific Northwest Laboratory, Richland, Washington, April 26, 1991.

R. A. Bair, "The Molecular Science Research Center," A segment of the minisymposia: "Intel/DARPA Touchstone Program: Update and Future Directions," presented at the Sixth Distributed Memory Computing Conference, Portland Oregon, April 28-May 1, 1991.

R. A. Bair, "The Molecular Science Software Program and the Molecular Science Computing Facility," presented at the IBM Thomas J. Watson Research Center, Yorktown, New York, July 31, 1991.

M. B. Bass, and R. L. Ornstein, "Molecular Basis of Substrate Specificity for Cytochrome P450cam by Molecular Dynamics Simulations," International Symposium on Cytochrome P450 of Microorganisms, Book of Abstracts, pp. 21-22, Berlin, Germany, July 1991.

M. B. Bass, and R. L. Ornstein, "Mutations Designed to Affect the Product Specificity of Cytochrome P450cam as Studied by Molecular Dynamics Simulations," Abstract #558, Regional ACS Meeting: Molecular Design & Modeling, Indiana University-Purdue University, Indianapolis, Indiana, May 1991.

M. B. Bass, M. D. Paulsen, and R. L. Ornstein, "Molecular Dynamics Simulations of Norcamphor-Cytochrome P450cam and Mutations of P450cam Designed to Alter the Product Specificity," ABSTRACT #6-2, 7th International Conference on Biochemistry and Biophysics of Cytochrome P450: Structure and Function, Biotechnological and Ecological Aspects, Moscow, USSR, July/August 1991.

F. J. Brockman, and R. L. Ornstein, "Enhanced Bioremediation of Subsurface Contamination: Enzyme Recruitment and Redesign," HMCRI's 12th Annual National Conference: Hazardous Materials Control '91, Washington, D.C., December 1991.

D. F. Feller, "Ab Initio Studies of Hyperfine Coupling Constants in the H_2CN and $\text{H}(\text{HO})\text{CN}$ Radicals," 12th Annual West Coast Theoretical Chemistry Conference, NASA Ames, March 27, 1991.

D. F. Feller, and S. S. Xantheas, "Exploiting Regularity in Systematic Sequences of Wavefunctions which Approach the Full CI Limit," Ab Initio Methods in Quantum Chemistry, Ames, Iowa, May 9, 1991.

D. Filipovic, P. J. Loida, S. G. Sligar, M. D. Paulsen, M. B. Bass, and R. L. Ornstein, "Approaches to the Rational Design of the Cytochrome P450cam Active Site," Abstract #9, 1991 President's Meeting of the International Society of Quantum Biology and Pharmacology, Stanford University, Palo Alto, California, September 1991.

B. G. Gladstone, and R. L. Ornstein, "Molecular Dynamics Simulation of Erabutoxin B," Fifth Annual Symposium of the Protein Society, Baltimore, Maryland, June 1991.

R. A. Kendall, "Initial Experiences with Parallel Computing on the Intel Delta, The Good, The Bad, and the Ugly," Workshop on Parallel Computing in Chemistry and Physics, Sponsored by Argonne National Laboratory and Pacific Northwest Laboratory, Chicago, Illinois, July 16-20, 1991.

R. A. Kendall, "Quantum Chemistry on the Intel iPSC/860 and a Network of Workstations," West Coast Theoretical Chemistry Conference, NASA Ames Research Center, March 27-29, 1991.

R. A. Kendall, "Parallel Computing in Theoretical Chemistry and Physics," University of Vienna, September 16-27, 1991.

J. P. LaFemina, and G. K. Schenter, "GaAs Bulk and Surface Phonon Spectra from Tight Binding Molecular Dynamics Simulations," 1991 West Coast Theoretical Chemistry Conference, NASA Ames, Mountain View, California, June 1991.

J. A. Loo, K. J. Light-Wahl, C. Edmonds, and R. D. Smith, "Tandem Mass Spectrometry and MS/MS/MS of Multiply Charged Ions from Large Biomolecules," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

J. A. Loo, K. J. Light-Wahl, R. D. Smith, and R. R. Ogorzalek-Loo, "Protein Structural Effects on Electrospray Ionization Mass Spectra and Tandem Mass Spectra," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

R. P. McRae, "Reaction Dynamics in Solution," Physics Department, University of Puget Sound, February 1991.

R. P. McRae, "Solvent Effects on Activated Chemical Reactions: Zero-Point Energy Constraints and Quantum Mechanical Tunneling," 1991 West Coast Theoretical Chemistry Conference, NASA Ames, Mountain View, California, June 1991.

R. P. McRae, "What is 'Liquid'?" Science Division, University of Puget Sound, February 1991.

R. P. McRae, G. K. Schenter, and B. C. Garrett, "Solvent Effects on Activated Chemical Reactions: Zero-Point Energy Constraints and Quantum Mechanical Tunneling," 1991 West Coast Theoretical Chemistry Conference, NASA Ames, Mountain View, California, June 1991.

R. L. Ornstein, "Enhancing Biodehalogenation by Redesigning Cytochrome P450cam Specificity." HAZMACON '91: Hazardous Materials Technical Conference, Santa Clara, California, April 1991.

R. L. Ornstein, "Environmental Remediation: Current & Future Role of Biological Sciences," HBCU/MI Curriculum Development Workshop, Jackson State University, Jackson, Mississippi, June 1991.

R. L. Ornstein, "Molecular Dynamics Simulations of Proteins: Validation Studies & Cytochrome P450cam Regiospecificity and Active Site Dynamics," Department of Chemistry, Pennsylvania State University, University Park, Pennsylvania, February 1991.

R. L. Ornstein, "Molecular Dynamics Simulations of Proteins: Validation Studies & Cytochrome P450cam Regiospecificity and Active Site Dynamics," Department of Pharmaceutical Chemistry, University of California at San Francisco, San Francisco, California, April 1991.

R. L. Ornstein, "Molecular Modeling and Simulation Studies to Redesign the Substrate Specificity of Biodegradative Enzymes to Enhance Remediation of DOE-site Contaminants," National Research and Development Conference on the Control of Hazardous Materials, Anaheim, California, February 1991.

R. L. Ornstein, M. B. Bass, M. D. Paulsen, and G. E. Arnold, "Rational Redesign of Substrate Specificity for the Biodegradative Enzyme Cytochrome P450cam," In Situ and On-Site Bioreclamation: An International Symposium, San Diego, California, March 1991.

R. L. Ornstein and J. A. Braatz, "3 Nanosecond Molecular Dynamics Simulation of the Protein Ubiquitin," 7th Conversation in the Discipline Biomolecular Stereodynamics, *J. Biomolecular Struct. & Dyn.* 8, a158, S.U.N.Y., Albany, New York, June 1991.

R. L. Ornstein, M. D. Paulsen, and M. B. Bass, "Biodegradation of Halogenated Hydrocarbons by Rational Redesign of Cytochrome P450cam," ACS Division of Biochemical Technology: Biodegradation of Organic Wastes: Molecular Approaches & Field Applications, New York, New York, August 1991.

R. L. Ornstein, M. D. Paulsen, M. B. Bass, G. E. Arnold, J. A. Braatz, and B. G. Gladstone, "Molecular Dynamics Simulations of Proteins: Validation Studies and Cytochrome P450cam Regiospecificity and Active Site Dynamics," 46th Northwest Regional Meeting of the ACS, Eastern Oregon State College, La Grande, Oregon, June 1991.

M. D. Paulsen, M.B. Bass, G. E. Arnold, and R. L. Ornstein, "Active Site Dynamics and Regiospecificity of Cytochrome P450cam as Determined by Molecular Dynamics Simulations," Abstract W-PM-D3 (*Biophysical J.* 59, 412a), 35th Annual Biophysical Society Conference, San Francisco, California, February 1991.

M. D. Paulsen, M. B. Bass, and R. L. Ornstein, "Active Site Dynamics and Regiospecificity of Cytochrome P450cam as Determined by Molecular Dynamics Simulations," Abstract #16, 1991 President's Meeting of the International Society of Quantum Biology and Pharmacology, Stanford University, Palo Alto, California, September 1991.

A. L. Rockwood, "Nomenclature for Mass-to-Charge Ratio," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

A. L. Rockwood, M. Busman, and R. D. Smith, "Are Large Highly Charged Polyatomic Ions Stable?", presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

A. L. Rockwood, M. Busman, H. R. Udseth, and R. D. Smith, "Thermally Induced Dissociation of Large Ions from Electrospray Ionization," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

A. L. Rockwood, J. A. Loo, H. R. Udseth, C. G. Edmonds, and R. D. Smith, "Obtaining Structural Information from Electrospray Ionization-Mass Spectrometry at the Large Molecule Limit," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

K. Ruedenberg, S. S. Xantheas, G. J. Atchity, and S. T. Elbert, "Intersections of Potential Energy Surfaces of Like Symmetry: Exotic or Commonplace? The Case of Ozone," presented at the international symposium "Ab Initio Methods in Quantum Chemistry, Iowa State University, Ames, Iowa, May 9-11, 1991.

G. K. Schenter, and B. C. Garrett, "The Influence of Reaction Path Curvature on Chemical Reaction Dynamics in Solution," 1991 West Coast Theoretical Chemistry Conference, NASA Ames, Mountain View, California, June 1991.

G. K. Schenter, R. P. McRae, and B. C. Garrett, "The Influence of Reaction Path Curvature on Chemical Reaction Dynamics in Solution," Gordon Research Conference on the Chemistry and Physics of Liquids, Holderness School, Plymouth, New Hampshire, August 1991.

R. D. Smith, "Capillary Electrophoresis Electrospray Ionization-Mass Spectrometry of Peptides and Proteins," presented at 11th International Symposium on HPLC of Proteins, Peptides, Polynucleotides, Washington, D.C., October 20-23, 1991.

R. D. Smith, J. A. Loo, C. G. Edmonds, A. L. Rockwood, and H. R. Udseth, "Electrospray Ionization-Tandem Mass Spectrometry for the Characterization of Large Biomolecules," presented at Pittsburgh Conference, Chicago, Illinois, March 4-8, 1991.

R. D. Smith, A. L. Rockwood, J. A. Loo, H. R. Udseth, and C. G. Edmonds, "Obtaining Structural Information from Electrospray Ionization-Mass Spectrometry at the Large Molecule Limit," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

R. D. Smith, A. L. Rockwood, M. Busman, J. A. Loo, R. R. Ogorzalek-Loo, and H. R. Udseth, "Activation and Dissociation of Large Multiply Charged Ions," presented at Ion Activation and Dissociation Symposium, Sanibel Island, Florida, January 28-February 1, 1991.

R. D. Smith, A. L. Rockwood, and H. R. Udseth, "Advanced Methods for Structural Characterization of Large Multiply Charged Ions: Obtaining Charge State Determination and Improved Sensitivity," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

R. D. Smith, H. R. Udseth, J. A. Loo, and C. G. Edmonds, "The Analysis of Peptide and Protein Mixtures by Capillary Electrophoresis-Tandem Mass Spectrometry," presented at Third International Symposium on HPLC, San Diego, California, February 4-6, 1991.

R. D. Smith, H. R. Udseth, A. L. Rockwood, K. J. Light-Wahl, M. Busman, and C. G. Edmonds, "Dissociation of Large Molecular Ions from Electrospray Ionization," presented at Joint Meeting of FACSS XVIII and Pacific Conference, Anaheim, California, October 6-11, 1991.

A. Sokalski, M. Shibata, R. L. Ornstein, and R. Rein, "*Ab Initio* Study of the Electrostatic Multipole Nature of Torsional Potentials: Hydrogen Peroxide and Methyl Disulfide (CH₃SSCH₃)," 35th Sanibel Symposia, St. Augustine, Florida, March 1991.

T. S. Tenforde, "Environmental and Molecular Sciences Laboratory at PNL," presented at the DOE Office of Health and Environmental Research Workshop on Structural Biology, Gaithersburg, Maryland, April 22, 1991.

T. S. Tenforde, "Preliminary Design of the First Gigahertz NMR Spectrometer," presented at the 13th Annual Bioelectromagnetics Society Meeting, Salt Lake City, Utah, June 23-27, 1991.

M. A. Thompson, "Studies of Photosynthetic Systems," invited plenary speaker at the Sanibel Symposium on Quantum Chemistry and Quantum Biology, Florida, March 10, 1991.

M. A. Thompson, "Theoretical Studies of Bacteriochlorophyll Dimers from Photosynthetic Reaction Centers," Invited Seminar, University of Idaho, Moscow, Idaho, February 22, 1991.

H. R. Udseth, J. H. Wahl, C. J. Barinaga, and R. D. Smith, "Interface Contributions to Peak Broadening in CE-ESI-MS," presented at ASMS on Mass Spectrometry and Allied Topics, Nashville, Tennessee, May 19-24, 1991.

A. C. Wang, M. A. Kennedy, B. R. Reid, and G. P. Drobny, "Solid State Deuterium NMR Study of the Purine Motion in an RNA Oligonucleotide." Presented at the 33rd Rocky Mountain Conference, Denver, Colorado, July 27 - August 1, 1991.

S. S. Xantheas and T. H. Dunning, Jr., "Critical Points on the H- + H₂O -> HO- + H₂ Potential Energy Surface," poster presentation at the West Coast Theoretical Chemistry Conference, NASA Ames Research Center, Moffett Field, California, March 27-29, 1991.

S. S. Xantheas, S. T. Elbert, G. J. Atchity, and K. Ruedenberg, "The Lowest Two 1A1 Potential Energy Surfaces of Ozone in the Vicinity of their Intersection Point," presented at the West Coast Theoretical Chemistry Conference, NASA Ames Research Center, Moffett Field, California, March 27-29, 1991.

S. S. Xantheas, D. Feller, and R. J. Cave, "Exploiting Regularity in Systematic Sequences of Wave Functions which Approach the Full CI," poster presentation at the international symposium "*Ab Initio* Methods in Quantum Chemistry," Iowa State University, Ames, Iowa, May 9-11, 1991.

Appendix C

Activities

Appendix C

Activities

Collaborations

Topic: *Ab Initio* Methods for Large Systems
Organization: University of Minnesota
(J. Almlöf)
MSRC Contact: R. A. Bair, R. A. Kendall

Topic: *Ab Initio* Methods in Distributed Computing Environments
Organization: University of Utah
(J. Simons, J. Nichols)
MSRC Contact: R. A. Kendall

Topic: Application Computer Interfaces
Organization: CAChe Inc.
(G. Purvis)
MSRC Contact: M. A. Thompson

Topic: Approximate Path-Integral-Based Methods for Reactions in Solution
Organization: University of Pennsylvania
(G. A. Voth)
MSRC Contact: B. C. Garrett

Topic: Atmospheric Clusters
Organization: University of Washington
(R. Watts)
MSRC Contact: S. W. Sharpe

Topic: Atomic-Scale Models of Heteroepitaxy
Organization: Montana State University
(J. Hermanson)
MSRC Contact: J. P. LaFemina

Topic: Clay Mineralogy
Organization: University of Florida
(C. L. Johnston)
MSRC Contact: A. C. Hess

Topic: Cluster Computing Approaches with the DISCO Program
Organization: IBM, Utah Supercomputer Institute
(J. Nichols)
MSRC Contact: R. A. Kendall

Topic: Comparison Between Fully Numerical and Basis Set Approaches to *Ab Initio* Quantum Chemistry Determinations of Properties
Organization: Utah State University
(E. McCullough)
MSRC Contact: D. F. Feller

Topic: Computer Pattern-Recognition Between Enzyme Substrates and Pollutants
Organization: Pennsylvania State University
(P. Jurs)
MSRC Contact: R. L. Ornstein

Topic: Computer Simulation Program Development
Organization: SUNY at Buffalo/Roswell Park
(R. Rein)
MSRC Contact: R. L. Ornstein

Topic: Concurrent Supercomputer Consortium
Organization: California Institute of Technology (and 12 other institutions)
MSRC Contact: T. H. Dunning

Topic: Defect Structures in Oxides
Organizations: University of Oregon
(R. Haydock)
Pennsylvania State University
(J. Lowe)
MSRC Contact: J. P. LaFemina

Topic: Destruction of Trichloroethylene
Organization: University of Pittsburgh
(J. Yates)
MSRC Contact: S. D. Colson, D. R. Baer

Topic: Direct Participation of Solvent Molecules in
Chemical Reactions
Organizations: Sandia National Laboratory
(C. Mellus)
MSRC Contact: B. C. Garrett

Topic: Excited-State Surface Chemistry/Chemical
Dynamics Facility
Organization: University of California-Berkeley
MSRC Contact: J. P. Cowin

Topic: Experimental and Theoretical Methods for
Treating Molecular Hyperfine Properties
Organization: Furman University
(L. Knight, Jr.)
MSRC Contact: D. F. Feller

Topic: Experimental and Theoretical Investigations of
Hydrated Nitric Acid Ices
Organizations: NOAA
(A. Ravishankara)
Washington State University
(R. Poshusta)
MSRC Contact: M. I. McCarthy

Topic: Experimental and Theoretical Studies of
Environmental Interfaces
Organization: University of Colorado
(S. George)
MSRC Contact: A. C. Hess, M. I. McCarthy

Topic: Experimental and Theoretical Studies of the
Hyperfine Properties of Small Molecules
Organization: Furman University
(L. Knight, Jr.)
MSRC Contact: D. F. Feller

Topic: Fabrication of Solid-State Chemical Sensors
Organization: University of Utah
(D. Petelenz)
MSRC Contact: J. Janata

Topic: Fracto-Emission
Organization: Washington State University
(T. Dickinson)
MSRC Contact: D. R. Baer

Topic: Ground and Excited States in Small Molecules
of Astrophysical Interest
Organization: University of Amsterdam
(G. de Brouckère)
MSRC Contact: D. F. Feller

Topic: High Accuracy Methods for Molecular
Hyperfine Properties
Organization: Notre Dame University
(D. Chipman and I. Carmichael)
MSRC Contact: D. F. Feller

Topic: Hybrid Quantum Chemistry and Empirical
Force Field Methods
Organization: University of Idaho
(W. Edwards)
MSRC Contact: M. A. Thompson

Topic: Hydrogen Bonding - Theory and Experiment
Organization: University of Washington
(R. Watts)
MSRC Contact: S. D. Colson

Topic: Incorporating Continuum Dielectric Solvation
Effects in Variational Transition-State Theory
Organization: University of Minnesota
(D. Truhlar)
MSRC Contact: B. C. Garrett

Topic: Instrument Interface to FTICR Mass
Spectrometer
Organization: Rice University
(R. Smalley)
MSRC Contact: G. A. Anderson

Topic: Intermolecular Interaction Benchmark Calculations

Organization: Oakland University
(M. Szczesniak)

MSRC Contact: R. A. Kendall

Topic: Investigation of the Twisted Intramolecular Charge Transfer Mechanism in Dimethylaminobenzonitrile

Organization: Xerox Webster Research Center
(C. Duke)

MSRC Contact: G. K. Schenter

Topic: Lignin Peroxidase Structure/Function Analysis

Organization: Oregon Graduate Institute
(M. Gold)

MSRC Contact: R. L. Ornstein

Topic: Low-Energy Electron and Positron Diffraction Determinations of Compound Semiconductor and Metal Surfaces

Organizations: Brandeis University
(C. Canter)
Xerox Webster Research Center
(C. Duke)

MSRC Contact: J. P. LaFemina

Topic: Low-Energy Electron Diffraction Determination of Oxide Surface Structures

Organization: University of Texas, Austin
(J. Erskine)

MSRC Contact: J. P. LaFemina

Topic: Massively Parallel Implementation of Full-Potential Linearized Augmented Plane Wave Method

Organization: Oregon State University
(H. Jansen)

MSRC Contact: M. S. Stave

Topic: Methane Conversion

Organization: AMOCO
(M. Barr)

MSRC Contact: S. D. Colson

Topic: Molecular Biology and Biochemistry of Cytochrome P450cam

Organization: University of Illinois, Urbana
(S. Silger)

MSRC Contact: R. L. Ornstein

Topic: Molecular Biology and Biochemistry of Haloalkane Dehalogenase

Organization: University of Gronningen, Netherlands
(D. Janssen)

MSRC Contact: R. L. Ornstein

Topic: Molecular Dynamics Program Development

Organizations: Roswell Park Memorial Institute and
State University of New York-Buffalo
(R. Rein)

MSRC Contact: R. L. Ornstein

Topic: Molecular Dynamics - Theory and Experiment

Organization: University of Washington
(E. Heller)

MSRC Contact: M. O. Trulson

Topic: Molecular Models of Dielectric Polymers

Organizations: Johns Hopkins University
(S. Kafafi)

MSRC Contact: J. P. LaFemina

Topic: Molecule Server Project

Organization: CAChe Scientific Inc.
(G. Purvis),
Oregon Graduate Institute
(D. Meyer, J. Cushing)

MSRC Contact: M. A. Thompson

Topic: Multiconfigurational Approaches in the Computation of Potential Energy Surfaces Governing Chemical Reactions

Organization: Ames Laboratory
(K. Ruedenberg)

MSRC Contact: S. S. Xantheas

Topic: New Methods for Treating the Correlation Problem

Organization: Harvey Mudd College
(R. Cave)

MSRC Contact: D. F. Feller

Topic: NMR of Cytochrome Proteins

Organization: Washington State University
(J. Satterlee)

MSRC Contact: R. L. Ornstein

Topic: NMR of Cytochromes and P450cam

Organization: Washington State University
(J. Satterlee)

MSRC Contact: R. L. Ornstein

Topic: NMR Studies of Nucleic Acid Chemistry

Organization: University of Washington
(G. Drobny, M. Kennedy)

MSRC Contact: T. S. Tenforde

Topic: Object-Oriented Databases for Computational Chemistry

Organization: Oregon Graduate Institute
(D. Maier, J. Cushing)

MSRC Contact: D. M. DeVaney

Topic: Parallel Configuration Interaction Techniques

Organization: Argonne National Laboratory
(R. Harrison)

MSRC Contact: R. A. Kendall

Topic: Parallel Coupled-Cluster Methods

Organization: SERC, Daresbury Laboratory
(A. Rendell, M. Guest)

MSRC Contact: R. A. Kendall

Topic: Parallelization of the COLUMBUS Program System

Organization: University of Vienna
(H. Lischka)

MSRC Contact: R. A. Kendall

Topic: Quantal Calculations of Reaction Rates in Solution

Organization: Northwestern University
(G. Schatz)

MSRC Contact: B. C. Garrett

Topic: Radical Ion Deposition

Organization: University of Colorado
(G. Ellison)

MSRC Contact: J. P. Cowin, S. E. Barlow,
T. M. Orlando

Topic: Reactions on High-Energy Surfaces and Transmission IR Spectroscopy of Oxide Surface Reactions

Organization: University of Pittsburgh
(J. Yates, T. Ballinger)

MSRC Contact: S. D. Colson, D. R. Baer

Topic: Real-Time Measurements of Surface Reaction Dynamics

Organization: University of California-San Diego
(A. Kummel)

MSRC Contact: D. M. Friedrich, J. P. Cowin

Topic: Scalable Algorithms for Grand Challenges

Organization: Argonne National Laboratory
(R. Stevens)

MSRC Contact: R. A. Bair

Topic: Separation Science Project

Organization: University of Idaho
(W. Edwards)

MSRC Contact: M. A. Thompson

Topic: Sequence Attributes Analysis of Protein Structure

Organization: Washington State University
(K. Dunker)

MSRC Contact: R. L. Ornstein

Topic: Site-Directed Mutagenesis and Biochemistry of Cytochrome P450cam

Organization: University of Illinois-Urbana
(S. Sligar)

MSRC Contact: R. L. Ornstein

Topic: Solid State Quantum Chemistry
Organization: University of Torino, Torino, Italy
(C. Pisani, R. Dovesi, C. Roetti)
MSRC Contact: A. C. Hess

Topic: Solid State Quantum Chemistry
Organization: SERC Daresbury Laboratory, England
(V. Saunders, N. Harrison)
MSRC Contact: A. C. Hess

Topic: Surface Atomic and Electronic Structure
of Oxides and Carbonates
Organization: Xerox Webster Research Center
(C. Duke)
Oregon State University
(H. Jansen)
MSRC Contact: J. P. LaFemina

Topic: Synthesis and Design of Artificial Cytochrome
P450s/NMR of Structures
Organization: University of Washington
(T. Sasaki)
MSRC Contact: R. L. Ornstein

Topic: Theoretical and Experimental Studies of
Photodissociation Dynamics on Surfaces
Organizations: Northwestern University
(E. Weltz, P. Stair)
Hebrew University
(B. Gerber)
MSRC Contact: M. I. McCarthy

Topic: Theoretical Geochemistry
Organization: Arizona State University
(P. McMillan)
MSRC Contact: A. C. Hess

Topic: Theoretical Geochemistry
Organization: University of Washington
(S. Ghose)
MSRC Contact: A. C. Hess

Topic: Theoretical Studies of One Electron Properties
Organization: University of Amsterdam
(G. de Brouckère)
MSRC Contact: D. F. Feller

Topic: Theoretical Studies of Heterogeneous Catalysis
Organization: Shell Development Corporation,
Houston
(T. Brownsecombe, M. Skiff)
MSRC Contact: A. C. Hess

Topic: Theoretical Studies of Transition Metal
Compounds Using the Multireference
Averaged Couple Pair Functional Method
Organization: Harvey Mudd College
(R. Cave)
MSRC Contact: D. F. Feller

Topic: Variational Transition State Theory
Organization: University of Minnesota
(D. Truhlar)
MSRC Contact: B. C. Garrett

Topic: X-ray Crystal Structure Determination
of Redesigned P450cam Mutant-Contaminant
Organizations: University of Maryland and CARB
(T. Poulos)
MSRC Contact: R. L. Ornstein

Topic: X-ray Crystal Structure of Haloalkane
Dehalogenase Enzymes
Organization: University of Groningen, Netherlands
(B. Dijkstra)
MSRC Contact: R. L. Ornstein

Topic: X-ray Crystal Structure Determination of
P450cam-Pollutant Complex
Organization: University of California-Irvine
(T. Poulos)
MSRC Contact: R. L. Ornstein

Colloquia Speakers During 1991

- Dr. Dimitris K. Agrafiotis, Harvard University
 Dr. J. Almolf, University of Minnesota
 Dr. Stephan B. H. Bach, Naval Research Laboratory
 Professor Paul Barbara, University of Minnesota
 Dr. Stephan E. Barlow, University of Colorado
 Dr. Bruce Berne, Columbia University
 Dr. Elliot Bernstein, Colorado State University
 Dr. Thomas E. Bitterwolf, University of Idaho
 Dr. C. Brooks, Carnegie Mellon University
 Dr. Gordon E. Brown, Jr., Stanford University
 Dr. Cindy Bruckner-Lea, University of Utah
 Dr. David G. Cahill, IBM T. J. Watson Research Center
 Dr. Roberto Car, University of Geneva, Switzerland
 Dr. Howard Carman, Oak Ridge National Laboratory
 Dr. Richard Cavanaugh, National Institute of Standards and Technology
 Dr. David Chambliss, IBM Almaden Center
 Dr. David Chandler, University of California
 Dr. Christopher E. D. Childsey, AT&T Bell Laboratories
 Dr. Herman M. Cho, California Institute of Technology
 Professor Gerhard Closs, University of Chicago
 Professor Rob D. Coalson, University of Pittsburgh
 Dr. Fred Cohen, University of California-San Francisco
 Dr. Jack Collins, Molecular Research Institute
 Dr. John Cushman, Purdue University
 Professor H. G. K. Daniel, Technical University of Munich, Germany
 Dr. Tom Dickenson, Washington State University
 Dr. Miles Dresser, Washington State University
 Dr. Charles B. Duke, Xerox Corporation
 Dr. Robert A. Eades, Cray Research
 Dr. Jerry Ebner, Monsanto Company
 Professor Gary A. Elcemen, New Mexico State University
 Dr. Paul Ellis, University of South Carolina
 Dr. W. Ermler, Stevens Institute of Technology
 Dr. Charles A. Evans, Callaghan Company
 Dr. D. Farrelly, University of California-Los Angeles
 Dr. David Funk, Los Alamos National Laboratory
 Professor Tom George, Washington State University
 Professor Andy Gewirth, University of Illinois
 Dr. Martyn F. Guest, SERC Daresbury Laboratory, England
 Dr. Audrey Hammerich, Fritz Haber Research Institute, Jerusalem, Israel
 Dr. Stan Harmanek, Czechoslovak Academy of Sciences, Prague, Czechoslovakia
 Dr. P. Jeffrey Hay, Los Alamos National Laboratory
 Dr. Martin P. Head-Gordon, AT&T Bell Laboratories
 Dr. Teresa Head-Gordon, AT&T Bell Laboratories
 Professor D. Michael Heinekey, Yale University
 Dr. Tony Heinz, IBM T. J. Watson Research Center
 Dr. Michael Henderson, University Pittsburgh
 Dr. Vlado Hladý, University of Utah
 Mr. Steven A. Hofstadler, University of Texas-Austin
 Dr. Detlof Hohl, University of Illinois
 Dr. Jan Hrbek, Brookhaven National Laboratory
 Dr. J. T. Hynes, University of Colorado
 Professor Toshiko Ichtye, Washington State University
 Dr. Jiri Janata, University of Utah
 Dr. Henri Jansen, Oregon State University
 Dr. David C. Johnson, University of Oregon
 Professor Phillip Johnson, SUNY at Stony Brook
 Dr. Rowland R. Johnson, Lawrence Livermore National Laboratory
 Professor Harold Jones, University of Ulm, Germany
 Dr. Mira Josowicz, Institute for Physics, Munich, Germany
 Dr. Stephen A. Joyce, Sandia National Laboratory
 Dr. D. Kasik, Electronic Data Systems
 Dr. Bruce Kay, Sandia Laboratories
 Dr. Katsumi Kimura, Institute for Molecular Science, Japan
 Dr. Michael Klein, University of Pennsylvania
 Professor Bruce E. Koel, University of Southern California
 Dr. Bryan E. Kohler, University of California-Riverside
 Dr. Richard Kouzes, Princeton University
 Dr. Randall A. LaViolette, Rockwell International Science Center
 Dr. A. Scott Lea, University of Utah
 Professor Edward C. Lim, University of Akron
 Dr. David H. Live, California Institute of Technology
 Dr. Laurie Locasio-Brown, National Institute of Standards and Technology
 Dr. Gilda Loew, Molecular Research Institute
 Dr. Stephen K. Loh, IBM T. J. Watson Research Center
 Prof. John Loper, University of Cincinnati
 Dr. Horst Lueck, Technical University of Berlin, Germany
 Dr. Yanjun Ma, Brookhaven National Laboratory
 Professor David Maier, Oregon Graduate Institute
 Dr. Allen D. Malony, University of Illinois
 Dr. Massimo Marchi, University of California
 Dr. John L. Markley, University of Wisconsin
 Professor Jeanne McHale, University of Idaho
 Dr. Walter Meier, Cornell University

Professor Roger E. Miller, University of
 North Carolina
 Dr. William H. Miller, University of
 California-Berkeley
 Dr. K. Mueller-Dethlefs, Technical University
 of Munich, Germany
 Dr. Robert W. Odom, Charles Evans & Associates
 Dr. Stanley J. Opella, University of Pennsylvania
 Dr. Thomas M. Orlando, Sandia National Laboratory
 Dr. Steve Otto, Oregon Graduate Institute
 Dr. Russel T. Pack, Los Alamos National Laboratory
 Dr. Robert Parson, Joint Institute for Laboratory
 Astrophysics, University of Colorado
 Dr. Charles H. F. Peden, Sandia National Laboratory
 Dr. Kristin A. Petersen, Los Alamos National
 Laboratory
 Professor David G. Pettifor, Imperial College, London,
 England
 Dr. A. R. Ravishankara, National Oceanic and
 Atmospheric Administration
 Dr. Phillip Reld, University of California-Berkeley
 Professor William Reinhardt, University
 of Washington

Dr. Emil Roduner, Physical and Chemical Institute,
 Zurich, Switzerland
 Dr. Celeste Rohlffing, Sandia National Laboratory
 Dr. Kim Sharp, Columbia University
 Dr. David Sherman, U.S. Geological Survey
 Dr. John D. Simon, University of California
 Prof. Sherwin Singer, Ohio State University
 Dr. Amitabha Sinha, University of Wisconsin
 Dr. David Sonnenfroh, Physical Sciences, Inc.
 Professor Ed Stern, University of Washington
 Dr. Diane M. Szaflarski, University of Colorado - JILA
 Dr. Pat A. Thiel, Iowa State University
 Dr. Russell Tonkyn, Brookhaven National Laboratory
 Dr. Frank Van Swol, University of Illinois
 Dr. Ramabhadran Vasudev, Rutgers University
 Dr. Michael Weaver, Purdue University
 Dr. Klaus Wendt, University of Mainz, Germany
 Dr. Hans Joachim Werner, University of Bielefeld,
 Germany
 Dr. Michael White, Brookhaven National Laboratory
 Prof. Richard V. Williams, University of Idaho
 Professor Curt Wittig, University of Southern
 California

MSRC Technical Advisory Panel

Professor J. Gregory Dash
 Department of Physics
 University of Washington

Professor Mostafa El-Sayed
 Department of Chemistry & Biochemistry
 University of California-Los Angeles

Dr. Robert J. Fletterick
 Biochemistry & Biophysics Department
 University of California-San Francisco

Professor Hans Frauenfelder
 Department of Physics
 University of Illinois

Dr. Jeffrey Hay
 Los Alamos National Laboratory

Professor Yuan T. Lee
 Department of Chemistry
 University of California-Berkeley

Dr. Brian W. Matthews
 Institute of Molecular Biology
 University of Oregon

Dr. Marvin Poutsma
 Oak Ridge National Laboratory

Tours and Visitors During 1991

Universities

Alabama Agricultural and Mechanical University
Central Washington University
Columbia Basin College
Northwest College and University Association of Science
Oregon Graduate Institute
Tulane University
University of Idaho
Washington State University

Industry

Battelle Memorial Institute
Chem-Nuclear Environmental Services, Incorporated
Corporate Communication Conference
General Motors Corporation
Seattle Business Leaders
Tri-Cities Industrial Development Council
Xerox Corporation

International

Japan Research Institute
Sumitomo Heavy Industries
Chubu Electric Power Company

Other National Laboratories

Brookhaven National Laboratory
Oak Ridge National Laboratory
Los Alamos National Laboratory
Sandia National Laboratory

Government

Assistant to the Secretary of Energy
Representatives of Clark County Washington
DOE Deputy Assistant Administrator
DOE Fossil Energy Programs
DOE Office of Congressional Affairs
DOE Office of Basic Energy Sciences
DOE Office of Environmental Restoration and Waste Management
DOE Richland Field Office
General Accounting Office
Governor of South Dakota
Governor of Washington
Mayor of Lacey, Washington
Oregon Secretary of State
U.S. Air Force
Washington Department of Trade and Development

Legislative

Congressional Science Fellow from the Office of Congresswoman Jolene Unsoeld
Legislative Assistant to Senator Slade Gorton
Legislative Assistant to Senator Brock Adams
Representative of U.S. House Interior Appropriations Subcommittee
U.S. House Committee on Energy and Commerce
Washington State Representative R Nehrer
Washington State Senator Jeanette Haynor

Media

Seattle Post-Intelligencer
Tri-City Herald
Washington Post

Appendix D

Acronyms and Abbreviations

Appendix D

Acronyms and Abbreviations

AES - auger electron spectroscopy	cm - centimeter
AFM - atomic force microscopy	CPU - central processing unit
AMP - adenosine 5'-monophosphate	CRADA - Cooperative Research and Development Agreement
amu - atomic mass unit	CRPC - Center for Research on Parallel Computation
ANL - Argonne National Laboratory	CSC - Concurrent Supercomputer Consortium
atm - atmosphere	CSDRF - Cluster Structure and Dynamics Research Facility
AVS - application visualization system	CSF - configuration state function
BIN - backbone information network	CVT - canonical variational theory
C2N - computer-computer networks	DBMS - database management systems
CAD - computer-aided design	DCE - Distributed Computing Environment
CAE - computer-aided engineering	DCS - Distributed Computing Systems
CAM - computer-aided manufacturing	DFS - Distributed File Systems
CASE - computer-aided software engineering	DMABN - dimethylaminobenzonitrile
CASSCF - complete active space self consistent field	DME - distributed management environment
cc - correlation consistent	DNA - deoxyribonucleic acid
CCD - charge-coupled-device	DNP - dynamic nuclear polarization
CCIA - Computational Chemistry Input Assistant	DOE - U.S. Department of Energy
CDF - common data format	EA - electron affinity
CE - capillary electrophoresis	EASI - electronic assistant for scientific investigation
CFS - consolidated file system	ECLS - epitaxial continued layered structure
CI - configurations interaction	EDS - Environmental Dynamics and Simulation (program)
CID - collision-induced dissociation	

EFF - empirical force field	GUI - graphical user interface
EMSL - Environmental and Molecular Sciences Laboratory	GW - gigawatt
EOCS - epitaxial overlapping structure	HF - Hartree-Fock
EOTS - epitaxial on top structure	HiPPI - High-Performance Parallel Interconnect
ER - environmental restoration	Hz - hertz
ESD - electron stimulated desorption	I/O - input/output
ESDIAD - electron-stimulated desorption, ion angular distributions	ICR - ion cluster resonance
ESI - electrospray ionization	IDL - interface development laboratory
ESN - end-user subscriber networks	IEEE - Institute of Electrical and Electronics Engineers
ESR - electron spin resonance	IMPPT - intramolecular Møller-Plesset perturbation theory
ESRC - Environmental Sciences Research Center	INDO/S - intermediate neglect of differential overlap spectroscopic
eV - electron volts	IP - ionization potential
EXAFS - extended x-ray absorption fine-structure	IR - Infrared
FaAA - Failure Analysis Associates	ISA - Industry Standard Architecture
FCI - full configuration interaction	IS&S - Information Systems and Services
FDDI - Fiber Data Distributed Interface	IVR - internal vibrational relaxation
fmol - femtomole	JILA - Joint Institute for Laboratory Astrophysics
fs - femtosecond	JPL - Jet Propulsion Laboratory (Caltech)
FT - Fourier transform	kDa - kilo-Dalton
FTICR - Fourier transform ion cluster resonance	kV - kilovolt
FTIR - Fourier transform infrared	L - liter
FY - fiscal year	LANL - Los Alamos National Laboratory
Gbyte - gigabyte	LAS - laser ablation source
GHz - gigahertz	LCAO - Linear Combination of Atomic Orbitals
GLE - generalized Langevin equation	

LDEI - laser-driven electron impact	MW - molecular weight
LEED - low-energy electron diffraction	NAM - nitric acid monohydrate
LEPD - low-energy positron diffraction	NASA - National Aeronautics and Space Administration
LiF - laser-induced fluorescence	NAT - nitric acid trihydrate
LIMS - laser ionization mass spectrometry	NBO - natural bond orbital (analysis)
LLNL - Lawrence Livermore National Laboratory	NCAR - National Center for Atmospheric Research
LMMS - Large Molecule Mass Spectrometry (program)	NCSA - National Center for Supercomputer Applications
<i>M</i> - molar	NDDO - neglect of diatomic differential overlap
MCSCF - multiconfiguration self-consistent field	netCDF - Network Common Data Format
MD - molecular dynamics	NIH - National Institutes of Health
MHz - megahertz	NIM - network information management
MIMD - multiple instruction-multiple data	NIST - National Institute of Standards and Technology
min - minute	NMR - nuclear magnetic resonance
ML - monolayer	NO - nitric oxide
mm - millimeter	NOE - nuclear Overhauser enhancement
MM - molecular mechanics	NORCUS - Northwest Colleges and Universities Association for Science
MNDO - modified neglect of differential overlap	NPT - constant pressure and constant temperature
MP2 - second-order perturbation theory	ns - nanosecond
MP4 - fourth-order Moeller-Plesset	NSF - National Science Foundation
MPa - megapascal	NVE - number of particles, volume, and energy
MPPT - supermolecular Moeller-Plesset perturbation theory	ODBMS - object database management systems
MR-SDCI - multireference singles and doubles configuration interaction	OSF - Open Software Foundation
MS - mass spectrometry	pdvz - polarized valence double zeta
MSCF - Molecular Science Computing Facility	PEF - potential energy function
MSRC - Molecular Science Research Center	PES - potential energy surfaces

PES - potential energy surface	SNL - Sandia National Laboratory
PET - poly(ethylene terephthalate)	SSD - Scientific Supercomputing Division (Intel)
PNL - Pacific Northwest Laboratory	STM - scanning tunneling microscopy (microscope)
ppb - parts per billion	TBTE - tight-binding total energy
ps - picosecond	TICT - twisted intramolecular charge transfer (twist-induced charge transfer)
QMEFF-MD - quantum mechanical empirical force field molecular dynamics	TMS - tandem mass spectrometers
QMMD - quantum mechanical molecular dynamics	TOF - time of flight
R2PI - resonant two-photon ionization	TST - transition state theory
RAID - redundant arrays of inexpensive disks	UHV - ultrahigh vacuum
REMPI - resonance-enhanced, multiphoton-ionization	UI - UNIX International
RF - radio frequency	UIMS - User Interface Management Systems
RFP - request for proposals	UV - ultraviolet
RGA - residual gas analyzer	V - volt
RHEED - reflection high-energy electron diffraction	VAF - velocity autocorrelation functions
RL - U.S. Department of Energy Richland Field Office	VLSI - very large scale integration
RMS - root mean square	VME - Versa Module Europa
RNA - ribonucleic acid	VTST - variational transition state theory
s - second	VTST - variational transition state theory
S - Siemens	VUV - vacuum ultraviolet
SAC - semiclassical adiabatic tunneling correction	WM - waste management
SAG - semiclassical adiabatic ground-state	XDR - external data representation
SCF - self-consistent field	XPS - x-ray photoelectron spectroscopy
SC - small-curvature	ZEKE - zero kinetic energy
SERC - Science and Research Council	ZGF - Zimmer Gunsul Frasca Associates

Distribution

No. of
Copies

OFFSITE

12 DOE/Office of Scientific and
Technical Information

C. Frank
U.S. Department of Energy
1000 Independence Ave SW
Washington, DC 20585

L. C. Ianniello
Deputy Associate Director
U.S. Department of Energy
Office of Basic Energy Sciences
ER-11, J-304/GTN
Washington, DC 20545

R. S. Marianelli
U.S. Department of Energy
Office of Basic Energy Sciences
ER-14, G-338/GTN
Germantown, MD 20545

W. M. Polansky
U.S. Department of Energy
Office of Basic Energy Sciences
ER-16, G-350/GTN
Germantown, MD 20545

R. Robson
U.S. Department of Energy
Office of Basic Energy Sciences
ER-17, G-358/GTN
Germantown, MD 20545

I. Thomas
U.S. Department of Energy
Office of Basic Energy Sciences
ER-13, J-317/GTN
Germantown, MD 20545

A. Weber
U.S. Department of Energy
Office of Basic Energy Sciences
ER-14, G-338/GTN
Germantown, MD 20545

ONSITE

6 DOE Richland Field Office

W. W. Bixby, A7-50
L. Erickson, A5-10
J. M. Hennig, A5-21
P. W. Kruger, A5-90
J. J. Sutey, A5-90
D. E. Trader, A5-90

43 Pacific Northwest Laboratory

R. C. Adams, K6-52
W. J. Bair, K1-50
J. F. Bagley, K1-64
T. D. Chikalla, P7-75
C. A. Counts, K2-20 (5)
G. H. Cunningham, K1-59
W. V. Discenza, K1-84
D. W. Dragnich, K1-46
J. W. Falco, K6-78
J. S. Hirsch, K1-45
F. C. Hood, K1-60
A. F. Johnston, K1-44
M. L. Knotek, K1-48
W. W. Laity, K2-15
R. C. Liikala, K1-57
L. M. Martucci, P7-35
R. J. Mayer, K1-62
J. L. McElroy, P7-46
G. R. Petersen, K1-40
J.T.A. Roberts, K1-73
A. A. Shepherd, K1-55 (5)
B. D. Shipp, K1-73
G. M. Stokes, K1-74

No. of
Copies

J. L. Straalsund, K1-79
D. B. Van Leuven, B1-40
I. L. White, BWO
W. R. Wiley, A4-25

No. of
Copies

L. D. Williams, K1-41
G. L. Work, K2-51
Publishing Coordination
Technical Report Files (5)

END

DATE
FILMED

6 / 23 / 92