WSRC-RP-91-013

**HYDROSTRATIGRAPHY OF THE GENERAL SEPARATIONS
AREA, SAVANNAH RIVER SITE (SRS), SOUTH CAROLINA
(U)**

by R. K. Aadland

WSRC-RP--91-013

Westinghouse Savannah River Company
Savannah River Site
Aiken, South Carolina 29808

DE93 008499

Other Authors:

M. K. Harris
(WSRC)
C. M. Lewis
(WSRC)
T. F. Gaughan
(WSRC)
T. M. Westbrook
Dames & Moore
Atlanta, GA

This paper was prepared in connection with work done under Contract No. DE-AC09-89SR18035 with the U. S. Department of Energy. By acceptance of this paper, the publisher and/or recipient acknowledges the U. S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper, along with the right to reproduce and to authorize others to reproduce all or part of the copyrighted paper.

oh
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MASTER

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from (615) 576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

DOCUMENT NO. WSRC-RP-91-13
Distribution Category: UC-#

**HYDROSTRATIGRAPHY OF THE GENERAL SEPARATIONS
AREA, SAVANNAH RIVER SITE (SRS), SOUTH CAROLINA (U)**

**Rolf K. Aadland, Mary K. Harris, Cathy M. Lewis,
Thomas F. Gaughan, and Thomas M. Westbrook**

Approved by:

**D.B. Moore, Section Manager
Environmental Sciences Section**

Publication Date: January 1991

**D.B. Moore, Section Manager
Authorized Derivative Classifier**

**Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808**

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY UNDER CONTRACT NO. DE-AC09-89SR18035

CONTENTS

	Page
ABSTRACT	1
ACKNOWLEDGEMENTS	3
I. REGIONAL GEOLOGY	4
A. Physiography	4
B. Geological History	6
C. Coastal Plain Stratigraphy, Lithology, and Structure	11
1) Cape Fear Formation	11
2) Lumbee Group	14
3) Paleocene-Eocene Black Mingo Group	16
4) Middle Eocene Orangeburg Group	18
5) Late Eocene Barnwell Group	20
6) "Upland" Unit	21
7) Quaternary Deposits	22
D. Faulting and Folding of Coastal Plain Sediments	22
1) Soil Fractures	24
II. SURFACE HYDROLOGY OF THE GSA	25
A. Upper Three Runs Creek	25
B. Fourmile Branch	26
III. REGIONAL SUBSURFACE HYDROLOGY OF SRS	28
A. Geologic Setting	28
1) Hydrogeology of the Paleozoic Rocks	28
2) Hydrogeology of the Triassic Rocks	29
3) Hydrogeology of Confining System I	31
4) Hydrogeology of the Coastal Plain Sediments	31
a) Hydrostratigraphic Nomenclature	31
b) Aquifer System I	37
c) Aquifer System II	37
d) Aquifer System I/II	38
IV. HYDROGEOLOGY OF THE GENERAL SEPARATIONS AREA	38
A. General Geology and Hydrogeology	38

CONTENTS (Continued)

	Page
B. Methods and Procedures	44
1) Database.....	44
2) Strip Logs	44
3) Cross Sections	45
4) Lithostratigraphic and Hydrostratigraphic and Interpretation	45
C. Lithostratigraphic Sequence at the GSA	45
1) Confining System I-II.....	47
2) Aquifer System II.....	55
a) Aquifer Unit IIA.....	55
b) Confining Unit IIA-IIB.....	57
c) Aquifer Unit IIB.....	61
• Aquifer Zone IIB ₁	61
• Faulting Associated with Carbonate Buildups.....	67
• Confining Zone IIB ₁ - IIB ₂	69
• Aquifer Zone IIB ₂ (Water Table Aquifer)	76
V. HYDROLOGICAL CHARACTERISTICS OF THE GSA.....	78
A. Identification of the "Uppermost Aquifer"	78
1) Aquifer Zone IIB ₂	84
2) Aquifer Zone IIB ₁	89
3) Aquifer Unit IIA	93
4) Confining Zone IIB ₁ -IIB ₂	95
5) Confining Unit IIA-IIB	96
6) Confining System I-II.....	96
B. Ground Water Flow in Cross Section	97
1) Regional Equipotential Cross Section 1	97
2) Regional Equipotential Cross Section 2.....	98
3) Regional Equipotential Cross Section 3.....	99
4) Regional Equipotential Cross Section 4.....	100
5) Regional Equipotential Cross Section 5A and 5B.....	101
C. Vertical Head Relationships.....	102
1) Head Differences Across Confining Zone IIB ₁ -IIB ₂ (Tan Clay)	102
2) Head Differences Across Confining Unit IIA-IIB (Green Clay).....	104
3) Head Relationships Between the Cretaceous and Tertiary Aquifers	104

CONTENTS (Continued)

	Page
REFERENCES.....	107
APPENDIX I.....	I-1
APPENDIX II.....	II-1

RECEIVED
NOV 1 1963
CST

LIST OF FIGURES

1. Physiographic Subprovinces of the S. C. Coastal Plain.....	5
2. Location of General Separations Area.....	7
3. Surface Water Drainage Map in the Vicinity of the F and H Area Seepage Basins.....	8
4. Tectonic Setting of Eastern U.S.	9
5. Regional Stratigraphic Cross Section	12
6. Coastal Plain Sediments at the SRS	13
7. Surface Drainage Map of SRS	27
8. Comparison of Hydrostratigraphic Units Used at SRS.....	33
9. Hydrostratigraphic Chart	34
10. Schematic Hydrostratigraphic Chart.....	35
11. Regional Hydrostratigraphic Cross Section	36
12. Hydrostratigraphic Nomenclature for the General Separations Area Savannah River Site	40
13. Schematic Map and Cross Section of Fluvial Deposits	41
14. Generalized Marine Depositional Environments	42
15. Overriding - Delta Depositional Environments.....	43
16. GSA Regional Cross Sectional Lines	46
17. Isopach Map of the Lower Carbonate Zone of Aquifer Zone IIB ₂	48
18. Isopach Map of the Upper Carbonate Zone of Aquifer Zone IIB ₁	49
19. Structure Contour Map (Top of Confining System I-II/Ellenton Clays).....	51
20. Geophysical Characteristics of Hydrostratigraphic Units.....	54
21. Isopach Map of Aquifer Unit IIA (Congaree Aquifer)	56
22. Isopach Map of Confining Unit IIA-IIB (Green Clay).....	58
23. Structure Contour Map (Top of Confining Unit IIA-IIB) (Green Clay)	59
24. Isopach Map of Aquifer Zone IIB ₁ (Barnwell/McBean Aquifers).....	65
25. Geophysical Characteristics of Limestone in HSB-122 well.....	68
26. Isopach Map of Confining Zone IIB ₁ -IIB ₂ (Tan Clay)	71
27. Structure Contour Map of Confining Zone IIB ₁ -IIB ₂	72
28. Isopach Map of Aquifer Zone IIB ₂ (Water Table Aquifer)	77
29. Potentiometric Surface Map - Aquifer Unit IIB ₂ (Water Table Aquifer).....	79
30. Potentiometric Surface Map - Aquifer Zone IIB ₁ (Barnwell/McBean Aquifers).....	80
31. Potentiometric Surface Map - Aquifer Unit IIA (Congaree Aquifer)	81
32. Head Difference Map Between Aquifers IIB ₁ and IIB ₂	82
33. Head Difference Map Between Aquifers IIA and IIB ₁	83
34. Head Difference Map Between Aquifer Unit IB of Aquifer System I and Aquifer IIA of Aquifer System II.....	106

LIST OF CROSS SECTIONS

<u>Plate</u>	<u>Regional Cross Sections</u>	<u>In Pocket</u>
1	General Separations Hydrostratigraphic Cross Section 1	
2	General Separations Hydrostratigraphic Cross Section 2	
3	General Separations Hydrostratigraphic Cross Section 3	
4	General Separations Hydrostratigraphic Cross Section 4	
5A	General Separations Hydrostratigraphic Cross Section 5A	
5B	General Separations Hydrostratigraphic Cross Section 5B	

Regional Equipotential Cross Sections

6	Lines of Equipotential Cross Section 1
7	Lines of Equipotential Cross Section 2
8	Lines of Equipotential Cross Section 3
9	Lines of Equipotential Cross Section 4
10A	Lines of Equipotential Cross Section 5A
10B	Lines of Equipotential Cross Section 5B

LIST OF TABLES

1.	Chemical Analyses of Water from Metamorphic and Triassic Rock at the Savannah River Site.....	30
2.	Thickness of the Upper and Lower Carbonate Zones for the General Separations Area.....	50
3.	Elevations of Confining Unit IIA-IIB (Green Clay) and Confining System I-II (Ellenton Clays).....	52
4.	Thickness of Confining Unit IIA-IIB (Green Clay).....	60
5.	Thickness of Aquifer Zone IIB ₁ (Barnwell/McBean Aquifers).....	62
6.	Thickness of Confining Unit IIB ₁ -IIB ₂ (Tan Clay) and the Saturated Portion of Aquifer Zone IIB ₂ (Water Table Aquifer).....	73
7.	Head Difference Between Aquifer Zone IIB ₂ (Water Table Aquifer) and Aquifer Zone IIB ₁ (Barnwell/McBean Aquifers).....	85
8.	Head Difference Between Aquifer Zone IIB ₁ (Barnwell/McBean Aquifers) and Aquifer Unit IIA (Congaree Aquifer)	91

HYDROSTRATIGRAPHY OF THE GENERAL SEPARATIONS AREA, SAVANNAH RIVER SITE (SRS), SOUTH CAROLINA (U)

**By Rolf K. Aadland¹, Mary K. Harris¹, Cathy M. Lewis¹,
Thomas F. Gaughan¹, and Thomas M. Westbrook²**

**¹Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808**

**²Dames and Moore
455 E. Paces Ferry Road
Atlanta, Ga. 30363**

ABSTRACT

Detailed analysis and synthesis of geophysical, core, and hydrologic data from 230 wells were used to delineate the hydrostratigraphy and aquifer characteristics of the General Separations Area at SRS. The study area is hydrologically bounded on the north and northwest by Upper Three Runs Creek (UTRC) and on the south by Fourmile Branch (FB). The Cretaceous-Tertiary sedimentary sequence underlying the study area is divided into two Aquifer Systems; in ascending order, Aquifer Systems I and II. The study concentrated on Aquifer System II, which includes all the Tertiary sediments above the Black Mingo Group (Paleocene) to the water table.

Aquifer System II is further subdivided from base upwards into Aquifers IIA and IIB, which are separated by Confining Unit IIA-IIB (informally termed the "green clay" at SRS). Aquifer IIB is further subdivided from the base upwards into Aquifer zones IIB₁ and IIB₂, separated by Confining zone IIB₁-IIB₂ (informally termed the "tan clay" at SRS). Aquifer IIA is incised by

Upper Three Runs Creek to the northwest and Aquifer zone IIB₁ is incised by Fourmile Branch to the south.

Aquifer zone IIB₂ is variable in thickness and hydraulically discharges groundwater towards the south in Fourmile Branch. Aquifer zone IIB₂ has a downward flow potential into Aquifer zone IIB₁. Aquifer zone IIB₁ (generally 60'-70' thick) has a potentiometric surface above its Confining zone IIB₁-IIB₂, with groundwater flow generally to the south to southeast, also discharging into Fourmile Branch. The zone has downward flow potential into Aquifer IIA.

The waters in Aquifer IIA (generally 40'-50' thick) are artesian and rise 30' to 40' above its Confining Unit IIA-IIB. Groundwater flow is towards UTRC to the north, generally opposite or transverse to the flow direction in the overlying aquifer zones. Carbonate buildups that truncate and/or offset the confining clay beds are interspersed in Confining Unit IIA-IIB. The buildups are potential pathways for the downward movement of water from Aquifer IIB into Aquifer IIA.

In general, the pressure (head) relationships are downward from Aquifer zone IIB₂ to Aquifer zone IIB₁, down to Aquifer Unit IIA. Between Aquifer Unit IIA and Aquifer Unit IB of Aquifer System I, however, an upward pressure gradient is recognized across Confining System I-II throughout the GSA.

A series of lithostratigraphic cross-sections, piezometric gradient profiles, head ratio contour maps, aquifer isopach maps, and potentiometric surface maps illustrate the aquifer characteristics of the study area.

ACKNOWLEDGMENTS

This report is the result of a combined effort by representatives of the Environmental Sciences Section of SRL and the Environmental Restoration Division. The report is a compilation and distillation of the vast amount of data assembled and organized for the Applications for Hazardous Waste Part B Post-Closure Care Permits for the F and H Area Seepage Basins. The study was an equal effort by all the authors.

I. REGIONAL GEOLOGY

A. Physiography

The SRS is located on the Upper Atlantic Coastal Plain about 20 miles southeast of the Fall Line (Figure 1). The Atlantic Coastal Plain is underlain by a seaward-dipping wedge of unconsolidated and consolidated sediments that extend from the contact with the crystalline Piedmont province at the Fall Line to the edge of the continental shelf. Sediment thickness increases from zero at the Fall Line to approximately 1000 feet near the General Separations Area (GSA) to more than 4000 feet at the South Carolina coastline (Colquhoun et al., 1983). The sequence ranges in age from early Late Cretaceous (Cenomanian) to Recent (Hazel, et al., 1977).

Colquhoun and Johnson (1968) divided the South Carolina Coastal Plain into three physiographic belts: Upper, Middle, and Lower Coastal Plain (Figure 1). The Upper Coastal Plain slopes from a maximum elevation of 650 feet msl at the Fall Line to about 250 feet msl on its southeastern boundary. Primary depositional topography of the Upper Coastal Plain has been obliterated by fluvial erosion. The Upper Coastal Plain is separated from the Middle Coastal Plain by the Orangeburg scarp, which has a relief of approximately 100 feet over a distance of a few miles.

Siple (1967) and Cooke (1936) previously divided the Upper Coastal Plain of South Carolina into the Aiken Plateau and Congaree Sand Hills (Figure 1). The Aiken Plateau, where the SRS is located, is bounded by the Savannah and Congaree rivers and extends from the Fall Line to the Orangeburg scarp. Its highly dissected surface is characterized by broad interfluvial areas with narrow, steep-sided valleys. Local relief is as much as 300 feet (Siple, 1967). The plateau is generally well drained, although many poorly drained sinks and depressions exist, especially

Figure 1. Physiographic Subprovinces of the S. C. Coastal Plain

on the topographically high (above 250 feet msl) "Upland" unit. Some depressions represent original depositional topography, whereas others may be related to solution of underlying calcareous sediment (Siple, 1967; U.S. Army Corps of Engineers, 1952).

The General Separations Area (GSA) is an approximately 15-square-mile area near the geographic center of the SRS (Figure 2) and includes the Hazardous Waste Management Facilities (HWMF): the F-Area Seepage Basins, the H-Area Seepage Basins, and the Mixed Waste Management Facility (MWMF). The area is located on the Aiken Plateau within the Coastal Plain province at an elevation of approximately 300 feet above msl. A major surface water divide runs through the GSA between Upper Three Runs Creek (UTRC) and Fourmile Branch (Figure 3). Plant north is used to describe locations in this report.

B. Geological History

The Coastal Plain sedimentary sequence at SRS is underlain by Precambrian to Permian aged metasedimentary and metaigneous rocks in the northern two thirds of the site, and by Late Triassic aged clastic sediments in the southern third of the site. The Paleozoic rocks are predominantly gneisses and schists with lesser amounts of quartzite (Klitgord and Behrendt, 1979) that have been folded, faulted, and intruded by predominantly granitic plutons. This deformation and metamorphism resulted from the accretion of Piedmont exotic terrains to North America. Accretion occurred during the Taconic, Acadian, and Alleghenian orogenies, which culminated about 265 million years ago at the close of the Paleozoic Era (Figure 4). At that time, Africa collided with North America, and the crystalline basement rocks beneath the SRS were emplaced in their present position.

Figure 2. Location of General Separations Area

Figure 3. Surface Water Drainage Map in the Vicinity of the F and H-Area Seepage Basins

Figure 4. Tectonic Setting of Eastern U. S.

Early in the Mesozoic Era (about 240 million years ago), the supercontinent Pangaea, which resulted from Paleozoic collisions, began to break up. The present Atlantic Ocean was created by diachronous rifting between North and South America on the west and Africa and Eurasia on the east, beginning in the north and progressing to the south. Rift basins that accumulated thick (1-3 miles) sequences of sediments in the Late Triassic (approximately 225 million years ago) formed along the east coast of North America .

The Dunbarton basin, which underlies the southeastern one-third of the SRS, is typical of these rift basins (grabens) filled with sediments of Triassic age. The basin is contained within the Paleozoic metamorphic rock beneath about 1200 feet of Coastal Plain sediments in the southeastern part of the SRS. The Triassic sediments consist of poorly sorted, consolidated gravel, sand, silt, and clay. The coarser material is found near the northwest margin where fanglomerates are abundant. By analogy with other East Coast Triassic basins in the Atlantic Coastal Plain, sand, silt, and clay predominate near the center of the basin. The sorting is extremely poor (Hazel, et al., 1977), resulting in extremely low primary permeability in the Triassic rocks. The lithology of the clasts in the sedimentary rock indicates that they were derived from the crystalline metamorphic rock to the northwest of the Dunbarton basin. Many of the sands are arkosic, indicating little transport and/or rapid burial.

Rifting of the continental crust ceased in Early Jurassic time (approximately 190 million years ago), at which time deposition in the basins ceased. Since then both the Paleozoic and Triassic rocks were leveled by erosion and eventually unconformably overlain by the unconsolidated to semiconsolidated Coastal Plain sedimentary sequence. This erosional surface dips approximately 35 feet/mile to the southeast (Klitgord and Behrendt, 1979).

Formation of the Coastal Plain sedimentary sequence at the SRS began with deposition of Late Cretaceous (Austinian/Turonian-Coniacian) aged fluvial and deltaic sediments on the crystalline

basement and Triassic sedimentary wedge. The Coastal Plain sediments began to accumulate at this time, when sea level had risen 650-980 feet above the present level, and the basement complex was being downwarped to the southeast to compensate for sediment loading on the more seaward portions of the continental margin. Deposition of sediments in fluvial and deltaic environments continued into the Early Eocene. During Middle and Late Eocene time, deposition of Coastal Plain sediments at the SRS occurred in shallow marine environments. Deposition was interrupted several times during this period because of drops in sea level.

Sea level fell following the deposition of the Eocene sediments, and except for the deposition of the "Upland" unit, erosion has prevailed on the Coastal Plain since. At present, the "Upland" unit was deposited by streams as they cut down into the underlying strata. At present, the only sedimentation on this portion of the Coastal Plain occurs locally in floodplains.

C. Coastal Plain Stratigraphy, Lithology, and Structure

The sediments of the Atlantic Coastal Plain in South Carolina are stratified sand, clay, limestone, and gravel that dip gently seaward and range in age from Late Cretaceous to Recent. The sedimentary sequence thickens from essentially zero thickness at the Fall Line to approximately 1000 feet at the GSA, to more than 4000 feet at the coast. (Figure 5). Regional dip is to the southeast.

1) Cape Fear Formation

The Cape Fear Formation (Figure 6) at the SRS is composed of poorly sorted, silty-to-clayey quartz sands and interbedded clays. The Cape Fear Formation sediments are more indurated than

Figure 5. Regional Stratigraphic Cross Section

Figure 6. Coastal Plain Sediments at the SRS

the other Cretaceous formations because of their high clay content and abundance of cristobalite in the matrix. The environment of deposition has been interpreted as upper delta plain (Prowell, 1985a). The Cape Fear Formation is about 30 feet thick at the northwestern SRS boundary and thickens to more than 180 feet near the southeastern boundary. Correlation of the basal Cretaceous at the SRS with the type section of the Cape Fear Formation is uncertain. Future correlation studies may reveal that the lowest Cretaceous unit at the SRS should have a different name, and the term is used provisionally in this report.

2) *Lumbee Group*

The Lumbee Group comprises, from oldest to youngest, the Middendorf, the Black Creek, and the Peedee Formations (Figure 6). The time-equivalent units of these three formations are present at the SRS. The lower two units are lithologically similar to the formations at their type localities in northern South Carolina and southern North Carolina. However, the Peedee Formation equivalent at the SRS is dissimilar to the type Peedee Formation. These Cretaceous formations have been referred to collectively as the "Tuscaloosa Formation" in past studies.

The Lumbee Group strata outcrop between the Fall Line and the SRS. The dip of the upper surface of the Lumbee Group is to the southeast at approximately 21 feet/mile across the SRS. Thickness varies across the SRS from 380 feet in the northwestern part of the site to more than 740 feet near the southeastern boundary. In general, Lumbee Group deposits are variably colored sands and clays, but become calcareous downdip from the SRS. At the coast the group is more than 1500 feet thick (Colquhoun, et al., 1983).

The Middendorf Formation, the basal formation of the Lumbee Group, is composed mostly of quartz sand, and unconformably overlies the Cape Fear Formation with a sharp, distinct contact.

The sediments of the Middendorf Formation at the SRS probably were deposited in fluvial and deltaic environments. In the subsurface, the formation is approximately 130 feet thick near the northwestern boundary of the SRS and thickens to more than 180 feet near the southeastern boundary. The Middendorf has been assigned a Late Austinian (Late Santonian) age in the SRS area (Prowell, et al., 1985a).

The Black Creek Formation consists of quartz sands, silts, and clays. It is generally darker and more micaceous and lignitic than the other Cretaceous units. The middle section of the formation is mostly shale, while the lower and upper sections consist of silty micaceous sands in much of the SRS area. In the northwestern part of the SRS, however, the lower portion of the Black Creek Formation is mostly light colored.

Lithofacies suggest a lower deltaic plain environment for the Black Creek Formation, except the light-colored sand in the northwestern part of the SRS. Here the lithofacies suggests an upper deltaic plain environment. The Black Creek is about 110 feet thick at the northwestern boundary (Prowell, et al., 1985a) of the SRS and thickens to more than 250 feet near the southeastern boundary. The Black Creek in South Carolina is assigned a Tayloran and Early Navarroan (Campanian and Early Maastrichtian) age.

The Peedee Formation is composed of light-colored sands and highly colored clays. Because of the difference in lithology between the type Peedee Formation and the laterally equivalent sediments in southwestern South Carolina (Colquhoun, et al., 1983), the deposits in the vicinity of the SRS will be referred to as the "Steel Creek Member" of the Peedee Formation. The lower part of the Steel Creek Member is sandy with a pebble-rich zone at its base, suggesting a basal unconformity. The upper part of the Steel Creek Member is a clay that varies from more than 50 feet to less than 3 feet in thickness at the SRS. Fining upward sands are interbedded with the clay in some areas. Lithofacies suggest an upper delta plain environment in the southwest portion of

SRS and a lower delta plain in the northeast. Dinoflagellates indicate some marine influence in parts of the Steel Creek Member. It is about 110 feet thick at the northwestern SRS boundary and 130 feet thick at the southeastern boundary. The Peedee Formation in South Carolina is Middle and Late Navarroan (Middle and Late Maastrichtian) in age (Prowell, et al., 1985a).

3) *Paleocene-Eocene Black Mingo Group*

In southwestern South Carolina, the Black Mingo Group consists of the Midwayan (Danian-Early Thanetian/Early Paleocene) Rhems Formation, the Early and Middle Sabinian (Middle Thanetian/Late Paleocene) Williamsburg Formation (Colquhoun, et al., 1983) (Figure 6), and the Late Sabinian (Late Thanetian/Early Eocene) Fishburne Formation (Gohn, et al., 1983). These, with the possible exception of the Fishburne, have been traced updip in the subsurface from their type localities in the lower Coastal Plain of South Carolina to the SRS (Figure 1). The upper surface of the group dips approximately 15 feet/mile to the southeast; the group is about 70 feet thick at the northwestern plant boundary, thickens to about 150 feet near the southeastern boundary, and is about 700 feet thick at the coast (Colquhoun, et al., 1983).

The stratigraphic interval that includes the Rhems Formation was named the "Ellenton Formation" and described from well cuttings at the SRS by Siple (1967). At the time, Siple (1967) thought that the age was either Cretaceous or Paleocene. The formation was later dated as Midwayan (Danian to Early Thanetian) and equivalent to the Rhems Formation (Prowell, et al., 1985b). Because it is lithologically different from the type Rhems Formation, the formation is designated as the "Ellenton Member" of the Rhems Formation at SRS. The interval has been informally referred to as the "Ellenton Clays" in previous SRS reports. The Ellenton Member lies unconformably on the Cretaceous sediments and consists mostly of gray, poorly sorted, micaceous, lignitic, silty and clayey quartz sand interbedded with gray clays.

Upper delta plain deposits composed of light-colored, moderately to poorly sorted, micaceous quartz sands are common in the northwestern part of the SRS, with darker, poorly sorted, micaceous, lignitic lower deltaic plain facies becoming dominant in the southeastern part. The Ellenton Member is approximately 40 feet thick at the northwestern boundary of the SRS and thickens to about 100 feet near the southeastern boundary. The Midwayan (Danian to Early Thanetian) section is about 250 feet thick and calcareous at the coast (Colquhoun, et al., 1983).

The deposits near the SRS that are time equivalent to the Sabinian (Mid-Late Thanetian) Williamsburg Formation differ from the type Williamsburg and are designated as the "Snapp Member" of the Williamsburg Formation (Figure 9). The unit is encountered in well P-22 in the southeastern part of the SRS near Snapp Station. The sediments are typically silty, medium- to coarse-grained quartz sand interbedded with clay. Dark, micaceous, lignitic sand also occurs. In the northwest part of the SRS, the Snapp Member is less silty and better sorted, with thinner clay interbeds. The Snapp Member changes from an upper deltaic plain facies in the northeast to a lower deltaic plain facies in the southeastern part of the SRS. The basal contact is probably unconformable. The Snapp Member appears to pinch out in the northwestern portion of SRS and thickens to about 50 feet near the southeastern boundary of the site. The Williamsburg Formation (Colquhoun, et al., 1983) is about 350 feet thick and calcareous near the coast.

Overlying the Williamsburg Formation is the basal Eocene Unit; approximately 30 feet (9 m) thick, mostly tan, yellow orange, and green, moderately sorted sand, with clay beds a few feet thick in the middle and at the top in places. Early Eocene ages, from palynological assemblages, indicate its equivalence to the Fishburne Formation (Gohn et al., 1983), a calcareous unit which occurs down dip near the coast; thus the name Fourmile Member of the Fishburne.

Overlying the Fourmile is 30 feet (9 m) of sands very similar to the Fourmile. Occasional dinoflagellate assemblages indicate that the sands are correlative with the Congaree Formation. Because of the difficulty in distinguishing the Fishburne from the Congaree Formation, the entire 60 foot section has been assigned to the Congaree in previous SRS reports.

4) Middle Eocene Orangeburg Group

The Claibornian (Lutetian-Bartonian) Orangeburg Group consists of the lower Middle Eocene Congaree Formation (Tallahatta equivalent) and the upper Middle Eocene Santee Limestone Formation (Lisbon equivalent) (Figure 6). Here we are not distinguishing the Congaree Formation from the underlying Fishburne Formation. The group is exposed at lower elevations in many places in the vicinity of and at the SRS. The sediments thicken from approximately 100 feet at the northwestern SRS boundary to about 160 feet near the southeastern boundary. Dip of the upper surface is about 12 feet/mile to the southeast across the plant. Downdip at the coast, the Orangeburg Group is about 325 feet thick (Colquhoun, et al., 1983).

The early Middle Eocene Congaree Formation has been traced from the Congaree valley in east-central South Carolina into the SRS area. At the SRS the formation consists of yellow, orange, tan, gray, and greenish gray, well-sorted, fine-to-coarse quartz sands. Thin clay laminae occurs throughout the section, and pebbly layers, clay clasts, and glauconite are present in places. The base of the formation is unconformable.

The Congaree is a shallow marine deposit. The equivalent of the Congaree northwest of the SRS has been mapped as the Huber Formation where it occurs at low elevations (Nystrom and Willoughby, 1982). At these locations it becomes more micaceous and poorly sorted, indicating deposition in fluvial and deltaic environments. The Congaree is about 60 feet thick at the

northwestern boundary of the SRS and about 85 feet thick near the southeastern boundary. The lower Middle Eocene deposits become increasingly calcareous between the SRS and the coast (Colquhoun, et al., 1983).

The late Middle Eocene deposits in the vicinity of the SRS, consisting of carbonates, calcareous quartz sands, glauconitic sands and clays, are designated as the Santee Formation in this report. "McBean Formation" and "Lisbon Formation" have been applied in the past to these sediments. Often the "McBean Formation" of previous SRS reports consists largely of sediments that are now known to be Late Eocene (Dry Branch Formation).

Fine-grained, often glauconitic sands that occur in many places at the SRS at and near the base of the Santee Limestone Formation are referred to as the "Warley Hill Member of the Santee Limestone" in this report, and the green clay beds in the interval are assigned to the "Caw Caw Member of the Santee Limestone". Both the Warley Hill and the Caw Caw lithologies occur at or near the base of the Santee Limestone Formation at the SRS and are the "green clay" beds referred to in many previous SRS reports.

The remainder of the Santee Limestone Formation is assigned to the McBean Member, which is made up of micritic, calcarenitic, and shelly limestone, calcareous quartz sand, and noncalcareous, generally fine-grained quartz sand. The McBean member of the Santee is the dominant member of the formation. Calcareous components of the McBean Member are rare in the northwestern part of the SRS, but are locally more abundant in central portions of the site. To the southeast they are widespread and thick. The McBean Member Carbonates form the lower Carbonate Zone noted in this report.

All three members of the Santee Limestone occur at the surface in the northern areas of SRS, where they are exposed along stream channels. The formation is approximately 40 feet thick at the

northwestern boundary of the SRS and thickens to more than 80 feet near the southeastern boundary.

5) Late Eocene Barnwell Group

Upper Eocene deposits lie unconformably on the Santee Limestone. Huddleston and Hetrick (1985) have recently revised the Upper Eocene stratigraphy of the Georgia Coastal Plain, and their approach has been extended into South Carolina (Nystrom and Willoughby, 1982; Nystrom et al., 1986). These authors elevated the Eocene (Jacksonian-Priabonian) "Barnwell Formation" to the "Barnwell Group", which in Burke County, GA, includes (from oldest to youngest) the Clinchfield, Dry Branch, and Tobacco Road Formations (Figure 6).

The Clinchfield Formation is sandy and contains some clay and siliceous sponge spicules. It can be identified at SRS only when the contrasting carbonates of the Griffins Landing Member of the Dry Branch Formation and the McBean Member of the Santee Limestone are present, with the sand sandwiched between them. It has been identified as being up to 25 feet thick at three localities within the SRS, but cannot be mapped with the available data.

The overlying Dry Branch Formation, which outcrops in many places around and within the SRS, is about 50 feet thick near the northwestern plant boundary and about 80 feet thick near the southeastern boundary. The Dry Branch Formation includes the Twiggs Clay Member, the Irwinton Sand Member, and the Griffins Landing Member, which contains carbonate. The Twiggs Clay Member does not appear to be mappable at SRS, but clay that is lithologically similar to the Twiggs Clay Member is present at various stratigraphic levels within the Dry Branch Formation. This clay lithofacies, which is tan, light gray, and brown, varies up to 12 feet thick in the SRS wells, but is not continuous over long distances. It has been called the "Tan Clay" in

previous SRS reports. The Irwinton Sand Member is composed of tan, yellow, and orange, moderately sorted quartz sand, with interlaminated clays abundant in places. The upper, noncalcareous part of the "McBean Formation" of previous SRS reports is the Irwinton Sand Member.

Previous reports of the SRS geology have assigned all carbonate to the Middle Eocene "McBean Formation." The carbonate of the Griffins Landing Member of the Dry Branch Formation occurs sporadically at the SRS. It is not known to be present northwest of Tinker Creek and Upper Three Runs Creek within the plant boundaries. The Griffins Landing carbonate ranges from 0 to 45 feet thick in the southeastern part of the SRS. The Griffins Landing Member formed in a shallow marine or lagoonal environment. The unit is referred to as the upper carbonate zone in this report. The Tobacco Road Formation, the youngest unit of the Barnwell Group, is probably the typical "Barnwell" of the older literature.

6) *"Upland" Unit*

The "Upland" unit is an informal stratigraphic term applied to fluvial deposits that occur at higher elevations (260 ft above msl at SRS) in many locations in the southwestern South Carolina Coastal Plain (Nystrom and Willoughby, 1982; Nystrom et al., 1985). The sediments are poorly sorted, clayey-to-silty sands, with lenses and layers of conglomerates, pebbly sands, and clays. The lower surface is irregular because of erosion of the underlying deposits that occurred prior to or during deposition of the "Upland". At higher elevations the "Upland" unit occurs at the surface in many places around and within the SRS, but is not present at all higher elevations (Figure 5). The precise age of the "Upland" unit is not known. Pliocene and Miocene ages have been assigned to the unit (Siple, 1967; Doering, 1960; Colquhoun and Steele, 1985; Cooke and MacNeil, 1952; Huddleston and Hetrick, 1979; and McCartan, et al., 1984).

7) Quaternary Deposits

Cooke (1936) identified seven marine terraces in the site region, although the highest three are extensively eroded and are not easily recognized (Siple, 1967). The terrace deposits are a few tens-of-feet thick, and their origin is problematic. Recent alluvium occurs in the main and tributary channels of the Savannah River and other streams in the vicinity of the SRS.

D. Faulting and Folding of Coastal Plain Sediments

The Atlantic continental margin has been tectonically quiescent since late Cretaceous time, and has been characterized by gentle subsidence and relatively uniform sedimentation rates. Faulting has been documented in the sedimentary sequence, however, in both South Carolina and Georgia. Subsurface faulting of sediments as young as Oligocene have been recognized in the Atlantic Coastal Plain province of South Carolina (Higgins, et al., 1978). Isopach and structure-contour maps of Oligocene rocks suggest subsurface faulting in the Atlantic Coastal Plain province of Georgia (Cramer and Arden, 1978). Small-scale faulting and folding through Middle Eocene time have been recognized in shallow sediments of the Atlantic Coastal Plain province near Langley, SC. There, normal faulting is associated with a small graben in the Eocene Barnwell Group (Inden and Zupan, 1975). The Paleozoic Carolina Slate belt argillites and Late Cretaceous sediments of the Middendorf Formation are folded and faulted near Cheraw, SC (Howell and Zupan, 1974).

Reactivation of basement faults associated with the northern flank of the Dunbarton Basin (Pen Branch fault) during the Cretaceous has been documented at SRS (Stieve, Stevenson and Aadland, 1991). Basement related (reactivated) faulting similar to the Pen Branch fault is probably responsible for the offset observed in the Tertiary section between the P-27 well and the HPC-1 well in the GSA (Plate 2). Shallow offsets other than the fault between the P-27 well and the

HPC-1 well are documented in the GSA in this study (Plates 1-10). These offsets are apparently related to slumping of the sediments that overlie and flank carbonate buildups that possibly underwent dissolution and collapse (see p. 66).

Preliminary results from correlation of core and downhole geophysical data from numerous wells used in Phase I of the Pen Branch fault study (Stieve, Stevenson and Aadland, 1991) indicates that the Upland Unit post-dates movement on the Pen Branch fault. Two on going studies: 1) Phase II of the Pen Branch fault study; and 2) the three year multi-disciplined Upland Unit study, are designed in part to precisely date and determine the youngest sediments involved in the most recent movement on the Pen Branch fault, and any other faults documented at depth. In addition, the precise age and most recent movement on the shallow slump features observed in the GSA is not well documented and awaits the results of the Upland Unit study and Phase II of the Pen Branch fault study. Until the results of the two studies are known, the upper or shallow portions of the slump features that underlie the areas at higher elevations in the GSA, which are capped by the Upland Unit, are dashed (Plates 2, 5A, 5B, 7, 10A, 10B). The slump features in the valleys and hill sides are not dashed since they penetrate to the surface due to recent erosion.

The Phase II study of the Pen Branch fault includes high resolution shallow seismic surveys and the addition of numerous boreholes that straddle the Pen Branch fault trace. All are designed to precisely date the most recent movement on the fault by analyzing in detail any and all cross cutting relationships and offsets observed in the shallow strata.

The three year multi-disciplined Upland Unit study is designed to determine the geometry, age, stratigraphic and possible structural relationships between the Upland Unit and underlying units throughout the SRS region. The study involves field mapping, shallow high resolution seismic surveys, paleontological age dating of the Upland Unit and underlying units as well as analysis of sediments from numerous auger holes. The results of the study will determine if and/or where

faulting and/or slump features have involved Upland or younger sediments.

1) Soil Fractures

Road cuts and excavations on interstream areas near the SRS commonly expose a deeply developed soil profile (Newell, et al., 1980). The "A" horizon may be up to 10 feet thick and typically consists of buff to nearly white, structureless, fine- to medium-grained quartz sand. The "B" horizon varies from 2 to 10 feet in thickness and contains iron and aluminum compounds leached from the overlying material. Reticulate mottling of this plinthic horizon is a salient feature.

Weathering effects extend to depths of at least 25 feet. At that depth in the "Upland" unit, feldspar in the originally arkosic unit is weathered completely, and some quartz cobbles are soft enough to crush manually.

In some areas, the intense chemical weathering has produced tensional soil fractures as a result of volume reduction. There has been no vertical displacement along these fractures. They are dominant features in shallow exposures, such as roadside embankments and drainage ditches. At depth they typically terminate as colored bands in the less weathered, parent sedimentary material. These soil fractures have been called "clastic dikes" in earlier reports. This misnomer is unfortunate, because true clastic dikes are caused by episodic earth movements powerful enough to cause liquefaction and upward injection of water-saturated sand through a more brittle overlying stratum. Johnson and Heron (1965) also describe soil fractures resulting from volume changes as "clastic dikes." No true clastic dikes have been observed at the SRS.

II. SURFACE HYDROLOGY OF THE GSA

The SRS is located on the Aiken Plateau (Figure 1). The plateau slopes southeastward approximately 5 feet/mile and is dissected by creeks that drain into the Savannah River. The major tributaries that occur on the SRS are Upper Three Runs Creek, Fourmile Branch, Tinker Creek, Pen Branch, Indian Grave Creek, Steel Creek, and Lower Three Runs Creek (Figure 7). The interstream upland area is flat to gently rolling and characterized by gently dipping units of sand, sandy clay, and clayey sand. The GSA lies between UTRC and Fourmile Branch (Figure 2).

A. Upper Three Runs Creek

The Upper Three Runs Creek watershed drains about 200 mi² of the Upper Coastal Plain northeast of the Savannah River. Significant tributaries to Upper Three Runs Creek are Tinker Creek, a relatively lengthy headwaters branch, and Tims Branch, which receives industrial wastes and wastewater from the Fuel Fabrication facilities (SRS M-Area) and the Savannah River Laboratory (SRL). The drainage basin consists principally of forest land and farmland and has no significant lakes or flow control structures.

The Upper Three Runs Creek stream channel has a low gradient and is meandering, especially in the lower reaches. Its floodplain ranges in width from 0.25 to 1 mile and contains extensive stands (about 98 percent) of bottom land hardwood forest (Faye and Prowell, 1982). Within the SRS the Upper Three Runs Creek valley is asymmetrical, having a steep southeastern side and a gently sloping northwestern side.

B . Fourmile Branch

Fourmile Branch drains about 23 mi² within the SRS, including much of F and H-Areas and C-Area. The creek flows to the southwest into the Savannah River Swamp and then into the Savannah River. The valley is V-shaped, with the sides varying from fairly steep to gently sloping. The floodplain is up to 1000 feet wide. Fourmile Branch receives effluents from F and H-Areas and, until June 1985, received large volumes of cooling water from the production reactor in C-Area. The creek valley has been modified by the cooling water discharge, which has created a delta into the Savannah River Swamp.

Figure 7. Surface Drainage Map of SRS

III. REGIONAL SUBSURFACE HYDROLOGY OF SRS

A. Geologic Setting

Three distinct geologic provinces occur in the subsurface beneath the Savannah River Site: 1) Paleozoic Metamorphic and igneous rocks; 2) lithified mudstones, sandstones, and conglomerates of Triassic age; and 3) the unconsolidated Coastal Plain sediments of Late Cretaceous and Tertiary age.

1) Hydrogeology of the Paleozoic Rocks

The Paleozoic and Triassic rocks comprise the Basement Hydrologic System (Figure 8). The hydrogeology of these deeply buried rocks was studied intensively to assess the safety and feasibility of deep storage of radioactive wastes in these rocks. Water injection and removal tests on packed-off sections of the sequence indicate that two kinds of fractures exist in the Paleozoic bedrock (Siple, 1967). The first consists of minute fractures that pervade the entire rock mass but transmit water slowly. Rocks containing only this kind of fracture are called "virtually impermeable rocks". The second kind of fractures have larger openings that transmit water more quickly. The fractures are generally vertically restricted but can often be traced laterally. Rocks containing these kinds of fractures are called "hydraulically transmissive rocks".

Marine (1979) reported representative hydraulic conductivity values of 3×10^{-4} gal per day per square foot (gpd/ft²) for the virtually impermeable rocks and 0.8 gpd/ft² for the hydraulically transmissive rocks. Analysis of a two-well tracer test indicated a porosity of 0.08 percent in the

hydraulically transmissive fracture zone. Laboratory analyses of cores indicated an average intergranular porosity of 0.13 percent (U.S. Army Corps of Engineers, 1952).

The Paleozoic bedrock is not pumped at the SRS except for testing programs. Immediately south of the Fall Line and to the north of it in the Piedmont province, the Paleozoic metamorphic and igneous rocks are domestic aquifers.

Because of the large amount of fresh ground water in the prolific aquifers of the overlying Coastal Plain sediments, the Paleozoic bedrock will remain an unexploited water source at the SRS. It is also unlikely that the status of the hydrologic regime of the metamorphic and igneous rocks at SRS would change appreciably for many decades. Chemical analyses of ground water from the Paleozoic crystalline rocks are listed in Table 1. The water has a total dissolved solids (TDS) content of about 6000 mg/l; dissolved solids are mainly calcium (500 mg/L), sodium (1300 mg/L), sulfate (2500 mg/L), and chloride (1100 mg/L).

2) Hydrogeology of the Triassic Rocks

Ground water occurs in the primary porosity of the Triassic clastic rocks. However, the hydraulic conductivity is extremely low, and water movement is almost nonexistent. The hydraulic conductivity of the Triassic sequence, as determined from field tests, ranged from 10^{-4} to 10^{-7} gpd/ft² (Cooke, 1936). The average total porosity was 8.0 percent for sandstones and 3.3 percent for mudstones. Average effective porosity was 7.0 percent for sandstones and 0.53 percent for mudstones.

Water is not pumped from the Triassic rocks in the Dunbarton Basin, nor is there likely to be any development in the future, because of the poor water quality and low permeability of the rocks.

Table 1

Table 1 includes chemical analyses of water samples from the Dunbarton Basin. Samples from the deeper wells near the center of the basin have TDS (Total Dissolved Solids) concentrations between 12,000 and 18,000 mg/L, predominantly NaCl (Marine, 1974).

3) Hydrogeology of Confining System I

Overlying the Paleozoic - Triassic Basement Hydrologic System is Confining System I (Figure 8), composed mostly of clayey silty sandstones and gravels of the Cape Fear Formation. Two clay-rich zones at the base of the system form an effective seal that separates the ground water in the Coastal Plain sediments above from the water in the underlying crystalline Paleozoic bedrock and the Triassic Basin. The first zone is a residual clayey material (saprolite) that immediately overlies the bedrock. The second zone is the clay-rich silty sands of the basal Cretaceous-age Cape Fear Formation, which occurs above the saprolite.

4) Hydrogeology of the Coastal Plain Sediments

a) Hydrostratigraphic Nomenclature

The Coastal Plain sedimentary sequence overlies Confining System I and constitutes a multilayer hydrologic system in which retarding beds are interspersed with beds that transmit water more readily. Ground water flow paths and flow velocities for each of these units are governed by the hydraulic properties, the geometry of the individual units, and the distribution of recharge and discharge areas.

Siple (1967) devised the first code for naming hydrostratigraphic units at the SRS. Since that time additional codes have been devised and used in various reports at the SRS (Figure 8). Thus, a generally accepted code has not existed for naming hydrostratigraphic units sitewide.

Consequently, this report will use the new hydrostratigraphic nomenclature system proposed by Aadland and Bledsoe (1990) (Figures 9, 10). The new hydrostratigraphic code provides a systematic classification scheme applicable to the entire SRS (Figure 11).

The classification scheme divides the Coastal Plain sediments at the SRS into two Aquifer systems. Within these aquifer systems are individual aquifer units, which may be further subdivided into zones. Aquifer systems, aquifer units, and zones do not necessarily coincide with stratigraphic boundaries (Figures 9 and 10).

The aquifer units used in this report are defined on the basis of their hydraulic properties relative to the hydraulic properties in the overlying and underlying confining units. The aquifer and confining units are mapped, however, on the basis of the thickness, areal distribution, and continuity of the lithologies of the units (Aadland and Bledsoe, 1990). A summary of the lithologic and hydraulic properties of the new hydrostratigraphic nomenclature is provided in the SRS Baseline Hydrogeologic Investigations Summary Report (Bledsoe, Aadland, and Sargent, 1990).

The proposed nomenclature follows criteria suggested by the South Carolina Hydrostratigraphic Subcommittee for naming aquifers and aquifer systems in South Carolina. Geographic names for the aquifers have not been formally agreed upon by the subcommittee. At present, an interim alpha-numeric hydrostratigraphic nomenclature is used to define and address the regional hydrogeologic characteristics of the SRS aquifers. Geographic names for the aquifers will replace the alpha-numeric system when the subcommittee formally agrees on the nomenclature.

Figure 8. Comparison of Hydrostratigraphic Units Used at SRS

Figure 9. Hydrostratigraphic Chart

Figure 10. Schematic Hydrostratigraphic Chart

Figure 11. Regional Hydrostratigraphic Cross Section

b) Aquifer System I

Aquifer System I overlies Confining System I throughout SRS. Stratigraphically, Aquifer System I includes the majority of the Upper Cretaceous age sediments of the Lumbee Group (Figure 6). In the main body of the SRS, including the GSA, Aquifer System I is divided into Aquifer Units IA and IB. These units are separated by Confining Unit IA-IB (Figures 9 and 10). Aquifer Unit IA includes the sands of the Middendorf Formation and lower third of the Black Creek Formation. Aquifer Unit IB includes the sands in the upper third of the Black Creek Formation and the Peedee Formation. Stratigraphically, Confining Unit IA-IB correlates with confining beds in the middle third of the Black Creek Formation (Aadland and Bledsoe, 1990).

Aquifer System I is overlain by Confining System I-II, which is often referred to as the Ellenton Clays in previous reports. Stratigraphically, the system includes the Paleocene Black Mingo Group and the confining beds of the upper Peedee Formation (Figures 8, 9, 10) (Aadland and Bledsoe, 1990).

In general, Aquifer Unit IB of Aquifer System I is persistent throughout the SRS. However, Confining System I-II to the north of UTRC ceases to be completely effective and Aquifer Unit IB and Aquifer System II are locally in communication (Figure 10). Here Aquifer System I/II is defined, and Aquifer Unit 1B is designated as Aquifer Unit I/IIB of Aquifer System I/II (Aadland and Bledsoe, 1990) (Figure 9).

c) Aquifer System II

Aquifer System II overlies Confining System I-II throughout the lower two-thirds of the SRS, including the GSA. Stratigraphically, it includes the Tertiary sedimentary sequence. The system is

divided into two aquifers, Aquifer Units IIA and IIB, separated by Confining Unit IIA-IIB (Figure 10). Aquifer Unit IIA, usually referred to as the Congaree Aquifer in previous reports, includes the sands of the Congaree Formation, the Four Mile Member of the Fishburne Formation, and locally the Warley Hill Member of the Santee Formation (Aadland and Bledsoe, 1990). Confining Unit IIA-IIB consists of the glauconitic clay beds of the Caw Caw Member located at and near the base of the Santee Formation. The Caw Caw Member is informally termed the "green clay" in previous reports. Aquifer Unit IIB consists of the sandy beds of the Santee Formation and the heterogeneous sedimentary sequences comprising the overlying Barnwell Group.

d) Aquifer System I/II

Aquifer System I/II is not present in the General Separations Area (GSA) at SRS and a discussion of it is not pertinent to this report. For a review of the System, the reader is referred to Aadland and Bledsoe, 1990.

IV. HYDROGEOLOGY OF THE GENERAL SEPARATIONS AREA

A. General Geology and Hydrogeology

The GSA is underlain by unconsolidated sediments occurring with isolated zones of consolidated sediments that range in age from Late Cretaceous to Holocene. This sedimentary sequence is approximately 1000 feet thick in the GSA and is underlain by Paleozoic crystalline basement. The sedimentary sequence strikes northeast and dips to the southeast towards the coast.

Aquifer Systems I and II are present under the GSA (Figure 9). The major ground water producing aquifers are Aquifer Units IA and IB (Middendorf and Black Creek Aquifers of previous reports) and Aquifer Unit IIA (Congaree Aquifer). Major confining beds under the GSA are Confining System I (Cape Fear) and Confining System I-II (Ellenton Clays). Minor confining beds in Aquifer System II are Confining Unit IIA-IIB (Green Clay) and Confining zone IIB₁-IIB₂ (Tan Clay) (Figure 10). This report will be limited to a hydrogeological discussion of the Tertiary-age sediments of Aquifer System II in the GSA.

The Tertiary sediments underlying the GSA are composed of sand, silt, and clay beds, as well as calcareous sediments. The sequence is complex, with abrupt changes in lithology occurring rapidly over short distances. This reflects the alternating nonmarine and marine characteristics typical of near-shore environments. Depositional environments consist of three major types: fluvial, deltaic and marine (Figures 13, 14, and 15).

Depositional environments affect the hydrologic properties of the sediments. Good aquifers consist of the coarsest and/or the most well-sorted sediments. Examples include channel and point bar deposits of both upper and lower delta plains; lower delta plain barrier islands, tidal channels, and beach deposits; and shallow siliclastic shelf sand deposits. Limestones associated with both siliclastic and carbonate shelf deposition can also act as good aquifers. Confining beds occur where the finest grained, least permeable sediments are deposited. Examples are interdistributary-bay and overbank flood deposits on the upper and lower delta plain; swamp and marsh deposits of the lower delta plain; and silt, clay, and marl of the delta front, prodelta, and deep siliclastic shelf deposits.

**Figure 12. Hydrostratigraphic Nomenclature for the General Separations Area
Savannah River Site**

Figure 13. Schematic Map and Cross Section of Fluvial Deposits

Figure 14. Generalized Marine Depositional Environment

Figure 15. Overriding - Delta Depositional Environment

B . Methods and Procedures

1) Database

The subsurface data used in the report comes largely from the F and H-Area Seepage Basins and the Mixed Waste Management Facility (MWMF) and represents the primary source of subsurface information for the GSA hydrostratigraphic interpretations. Subsurface data sources also include interim status wells, deep rock borings, baseline hydrogeological wells (P-wells), and production wells. Data types include geophysical logs (natural gamma, self-potential, resistivity, and caliper), geologist and driller field logs, detailed core logs, well construction data, slug tests, and sieve analyses. A master data list of all well data for the GSA is presented as Appendix I.

2) Strip Logs

For each well with a geophysical gamma log, a hand drawn strip log of the lithologies encountered in the well was constructed as the initial step in the geological interpretation of the data. All the strip logs are documented and available in the F and H Area Seepage Basins Part B Closure Permit Applications. Over 160 strip logs were constructed in this process, all at the vertical scale of 1"= 10'. The core logging descriptive data were given first priority in the lithologic descriptions, with lithologies interpreted from the gamma curves used as a secondary source of comparison. This required the review of the field log and occasional visits to the core repository for more detailed information. The strip logs represented a uniform interpretation of the combined inputs of all of the available subsurface data from the cored and geophysically logged borings. They are maintained in a central repository at SRS.

3) Cross Sections

Six regional cross sections were constructed within the GSA (Figure 16). The six cross sections illustrate the lithostratigraphy and hydrostratigraphy of the region. Detailed descriptions of the lithofacies and thicknesses of the aquifer units observed on the six cross-sections are available in Appendix II. The regional cross sections will provide a database for ongoing and future hydrostratigraphic interpretations within individual facilities in the GSA.

4) Lithostratigraphic and Hydrostratigraphic Maps and Interpretation

A base map of the GSA at a scale of 1"= 1000' was used. At this scale adequate areal coverage of the entire GSA was obtained, while an effective level of detail for the interpretations was maintained. All structure contour maps, isopach maps, potentiometric maps, and head difference maps for the GSA were also constructed at this scale.

C. Lithostratigraphic Sequence at the GSA

The lithostratigraphic sequence at the GSA is composed mostly of terrigenous clastics interspersed with carbonate-rich clastics and limestones. The clastic facies consist of gravel and/or pebble sands, clayey sands, silt, clay, and sandy clay. The calcareous facies consist of calcareous sand, calcareous mud, limestone, sandy limestone, and sandy and muddy limestone. The clastic lithofacies are dominant throughout the GSA and occur within all hydrostratigraphic units. Beds thicken and thin abruptly, pinching out and intertonguing rapidly over short distances.

Figure 16. GSA Regional Cross Sectional Lines

The calcareous lithofacies is discontinuous, but confined to two distinct horizons. The lower horizon (zone) occurs in three separate areas in the GSA along a northeast-southwest trend. The lower horizon is equivalent to the McBean Member of the Santee Formation located near the base of the formation (Figure 10). This horizon thickens and thins rapidly (Table 2 and Figure 17). Slump faulting observed on the various cross-sections typically occurs on the flanks of these carbonate zones, suggesting displacement of overlying sediment due to dissolution and compaction of the limestone horizon. The upper zone is very discontinuous and occurs at two locations in the GSA (Table 2 and Figure 18). This zone occurs within the upper portions of Aquifer zone IIB₁ (Barnwell/McBean Aquifers) and Confining zone IIB₁-IIB₂ (Tan Clay) and is equivalent to the Griffins Landing Member of the Dry Branch Formation (Figure 10).

1) Confining System I-II

Confining System I-II is informally referred to as the "Ellenton Clays" at the SRS. This system attains a thickness of over 100 feet in the GSA. A structure contour map was constructed on the top of the Confining System (Figure 19), with elevations tabulated in Table 3. This system consists predominantly of clays and silts, with interbeds of sandy clay, clayey sands, and sands. Individual layers of clays, silts, and sandy clays range in thickness from 2 to 22 feet. Individual layers of clayey sands and sands range in thickness from 1 to 15 feet.

The clay and silt lenses of Confining System I-II consist of buff and light gray to black clay and sandy clay. The clay is generally very fissile, highly micaceous, and often carbonaceous or lignitic. The interbeds of clayey sands and sands are generally fine to coarse grained and range from moderate to very poorly sorted. Grains are sub-angular to angular. According to Dennehy and others, 1989, the predominant clay minerals forming the matrix of the clays, sandy clays, and clayey sands are kaolinitic. The quartz that is present consists predominantly of cristobalite. The high mica content of the Confining System I-II sediments result in uniformly distinct, natural

Figure 17. Isopach Map of the Lower Carbonate Zone of Aquifer Zone IIB₂

Figure 18. Isopach Map of the Upper Carbonate Zone of Aquifer Zone IIB₁

TABLE 2
THICKNESS OF UPPER AND LOWER CARBONATE ZONES
FOR THE
GENERAL SEPARATIONS AREA

Well	Top Elevation (ft)	Bottom Elevation (ft)	Thickness (ft)***
BGO 6A*	175.8	123.8	52
BGO 8A*	172.3	125.3	47
BGO 10A*	169.1	128.1	41
BGO 14A*	179.2	136.2	43
BGO 16A*	158.8	151.8	6
BGO 18A*	153.9	142.9	11
BGO 25A*	178.7	150.7	28
BGO 26A*	177.1	159.1	18
BGO 27C*	156.9	145.9	11
FSB 100A*	149.8	131.8	18
FSB 106C*	152.0	143.0	11
FSB TA*	160.4	129.4	31
HPT 1A*	152.9	141.9	11
HPT 2A*	154.7	124.7	20
HSB 65A*	160.7	88.7	62
HSB 68A*	160.4	120.4	40
HSB 85A*	160.1	132.1	32
HSB 118A*	145.0	140.0	5
HSB 120A*	101.0	99.0	2
HSB 121A*	147.3	128.3	19
HSB 122A*	149.4	123.4	26
HSB 123A*	154.3	113.3	41
HSB 124A*	138.9	130.9	8
HSB 127C*	157.4	136.4	21
HSB 134C*	144.1	141.1	3
HSB TB*	156.1	108.1	48
P18 TA**	170.7	129.7	41
BGO 26C**	101.1	99.1	2
HSB 126C**	182.3	170.3	12

NOTES:

1. * Lower carbonate zone
2. ** Upper carbonate zone
3. *** Rounded to the nearest whole foot

Figure 19. Structure Contour Map (Top of Confining Systems I-II/Ellenton Clays)

TABLE 3
ELEVATIONS OF CONFINING UNIT IIA-IIB (GREEN CLAY)
AND
CONFINING SYSTEM I-II (ELLENTON CLAYS)

Well	Elevation Bottom of Confining Unit IIA-IIB (Green Clay)	Elevation Top of Confining System I-II (Ellenton Clays)	Thickness of Aquifer Unit IIA (Congaree)
BGO 6A	108.8	--	--
BGO 8A	120.3	--	--
BGO 10A	119.1	--	--
BGO 12A	119.4	--	--
BGO 14A	115.2	--	--
BGO 16A	123.8	--	--
BGO 18A	124.9	--	--
BGO 25A	111.7	--	--
BGO 26A	111.1	--	--
BGO 29A	120.1	--	--
DRB 4	111.0	44.0	67
FSB 76A	115.5	29.5	86
FSB 78A	102.5	--	--
FSB 79A	102.1	20.0	82
FSB 87A	110.6	--	--
FSB 96A	103.7	--	--
FSB 97A	106.8	--	--
FSB 98A	107.7	--	--
FSB 99A	112.3	--	--
FSB 100A	115.8	--	--
FSB 101A	116.9	--	--
FSB TA	116.4	43.5	73
HPT 1A	117.9	52.0	62
HPT 2A	117.7	56.0	62
HPC 1	109.5	46.5	63
HSB 65A	110.7	--	--
HSB 68A	111.4	--	--
HSB 69A	111.1	--	--
HSB 83A	105.9	--	--
HSB 84A	110.4	--	--
HSB 85A	115.1	52.0	63
HSB 86A	110.0	58.0	52
HSB 117A	114.1	--	--
HSB 118A	113.0	--	--
HSB 119A	109.8	--	--

TABLE 3
ELEVATIONS OF CONFINING UNIT IIA-IIB (GREEN CLAY)
AND
CONFINING SYSTEM I-II (ELLENTON CLAYS)
(Continued)

Well	Elevation Bottom of Confining Unit IIA-IIB (Green Clay)	Elevation Top of Confining System I-II (Ellenton Clays)	Thickness of Aquifer Unit IIA (Congaree)
HSB 120A	109.0	--	--
HSB 121A	109.3	--	--
HSB 122A	107.4	--	--
HSB 123A	104.3	--	--
HSB 124A	114.9	--	--
HSB 139A	113.0	--	--
HSB TB	106.1	54.0	52
IDL 3	126.4	28.0	98
P18 TA	113.7	7.0	107
P27 TA	127.1	48.5	79
P28 TA	117.2	59.0	58

NOTES:

1. -- Data not available

Figure 20. Geophysical Characteristics of Hydrostratigraphic Units

gamma ray curve spikes typical of clays (Figure 20). These spikes are very conspicuous in the borings that penetrate Confining System I-II in the GSA.

2) Aquifer System II

Aquifer System II is subdivided into Aquifer Unit IIA (Congaree Aquifer), Confining Unit IIA-IIB (Green Clay), and Aquifer Unit IIB. Aquifer Unit IIB is further subdivided into Aquifer zone IIB₁ (Barnwell/McBean Aquifers), Confining zone IIB₁ - IIB₂ (Tan Clay), and Aquifer zone IIB₂ (Water Table Aquifer) (Figure 12).

a) Aquifer Unit IIA

Aquifer Unit IIA is commonly referred to as the Congaree Aquifer in previous reports at SRS. This unit ranges in thickness from 52 to 107 feet within the GSA (Table 3). An isopach of this unit is presented in Figure 21. The isopach map indicates a thickening of the unit in the western portion of the GSA, a minor thickening to the southeast, and a minor thinning of this aquifer under the MWMF.

Aquifer Unit IIA consists predominantly of sands and clayey sands with thin interbeds of sandy clays, clays, and calcareous sands. Sands and clayey sands range in thickness from 2 to 84 feet. Sandy clays and clays range in thickness from 1 to 9 feet. Calcareous sands range in thickness from 2 to 3 feet. The sands and clayey sands of Aquifer Unit IIA are largely yellow to orange in color and consist of fine to coarse grained, subangular to subrounded quartz. The sands vary from

Figure 21. Isopach Map of Aquifer Unit IIA (Congaree Aquifer)

well to poorly sorted. Relatively small quantities of heavy minerals were noted. The assemblage consists of sillimanite, ilmenite, staurolite, rutile, garnet, and magnetite (Dennehy, et al., 1989).

Aquifer Unit IIA (Congaree Aquifer) typically exhibits a natural gamma ray curve normally associated with clean sands (Figure 20). The contact between the base of Aquifer Unit IIA and the underlying Confining System I-II is readily apparent where the transition occurs from clean sands to clayey/silty sediments.

b) Confining Unit IIA-IIB

Confining Unit IIA-IIB (Caw Caw member of the Santee Formation) is commonly referred to as the "Green Clay" in previous reports at SRS. Previous investigators illustrated this unit as one vertically and horizontally continuous clay zone. The current study has revealed that the "Green Clay" consists of several clay lenses that thicken, thin, and pinch out abruptly.

Confining Unit IIA-IIB ranges in thickness (Table 4) from 2 to 30 feet (Figure 22). The unit is thin to the west and southeast, becoming thicker to the north. A structure contour map of the top of the unit (Figure 23) indicates a southeastern dip with localized lows near the H-Area Seepage Basins and on the northern boundary of the MWMF. These structural lows tend to correspond with the thickest accumulations of the carbonates (Figure 17). In addition, the clays of the confining unit generally thin here or are truncated at the periphery of the carbonate buildups.

Confining Unit IIA-IIB consists predominantly of clays, sandy clays, and clayey sands. Locally, beds of calcareous mud contribute significant thickness to the unit. Minor interbeds of sands are also present. Clay and sandy clay lenses range in thickness from 2 to 12 feet. Clayey sand beds

Figure 22. Isopach Map of Confining Unit IIA-IIB (Green Clay)

**Figure 23. Structure Contour Map (Top of Confining Unit IIA-IIB)
(Green Clay)**

TABLE 4
THICKNESS OF CONFINING UNIT IIA-IIB (GREEN CLAY)

Well	Ground Elev.	Top Elev. IIA-IIB	Bottom Elev. IIA-IIB	Thickness IIA-IIB
BGO 6A	283.8	138.8	108.8	30
BGO 8A	281.3	124.3	120.3	4
BGO 10A	299.1	132.1	119.1	13
BGO 12A	311.4	143.4	119.4	24
BGO 14A	300.2	136.2	115.2	21
BGO 16A	302.8	130.8	123.8	7
BGO 18A	292.9	131.9	124.9	7
BGO 25A	294.7	137.7	111.7	26
BGO 26A	285.1	132.1	111.1	21
BGO 29A	262.1	124.1	120.1	4
DRB 4	250.0	117.0	111.0	6
FSB 76A	291.5	123.5	115.5	8
FSB 78A	270.5	106.5	102.5	4
FSB 79A	216.1	106.1	102.1	4
FSB 87A	285.6	114.6	110.6	4
FSB 96A	277.7	108.7	103.7	5
FSB 97A	283.8	109.8	106.8	3
FSB 98A	280.7	109.7	107.7	2
FSB 99A	285.3	115.3	112.3	3
FSB 100A	283.8	118.8	115.8	3
FSB 101A	282.9	118.9	116.9	2
FSB TA	275.4	118.4	116.4	2
HPT 1A	232.9	121.9	117.9	4
HPT 2A	257.7	121.7	117.7	4
HPC 1	293.5	116.5	109.5	7
HSB 65A	270.7	124.7	110.7	14
HSB 68A	247.4	116.4	111.4	5
HSB 69A	234.1	115.1	111.1	4
HSB 83A	234.9	116.9	105.9	11
HSB 84A	226.4	118.4	110.4	8
HSB 85A	292.1	127.1	115.1	12
HSB 86A	260.0	116.0	110.0	6
HSB 117A	234.1	122.1	114.1	8
HSB 118A	245.0	118.0	113.0	5
HSB 119A	254.8	113.8	109.8	4
HSB 120A	266.0	111.0	109.0	2
HSB 121A	272.3	114.3	109.3	5
HSB 122A	269.4	110.4	107.4	3
HSB 123A	262.3	110.3	104.3	6
HSB 124A	263.9	122.9	114.9	8
HSB 139A	231.5	116.5	113.0	4
HSB TB	267.1	112.1	106.1	6
IDL 3A	257.4	113.4	126.4	5
P18 TA	296.7	122.7	113.7	9
P27 TA	274.1	132.1	127.1	5
P28 TA	285.2	140.2	117.2	23

NOTES: 1. * Rounded to the nearest foot 3. NP Not penetrated
 2. NCU No confining unit present 4. -- Not available

range in thickness from 1 to 14 feet. Calcareous muds range in thickness from 3 to 11 feet and the sand beds range in thickness from 1 to 6 feet.

The clay minerals that constitute the matrix of the units described above consist of illite and smectite according to Dennehy and others, 1989. R. Strom (personal communication) notes that smectite, illite, and kaolinite occur in the unit in that order of abundance. The clays, sandy clays, and clayey sands generally have a distinct dark green color due to the accessory mineral glauconite. The calcareous muds tend to be light green in color. The clayey sands are generally moderately to poorly sorted, with grain sizes ranging from medium to very coarse. They can be consolidated locally within the GSA and are light tan to orange in color, fine to very fine grained, and well sorted. Grains are usually subangular and cemented with silica. The dark green clay facies exhibits a distinct natural gamma ray signature (Figure 20). A large spike on the gamma curve makes this unit easily distinguishable from sediments above and below the horizon.

c) Aquifer Unit IIB

Aquifer Unit IIB is divided from the base upwards into Aquifer zones IIB₁ and IIB₂, separated by Confining zone IIB₁-IIB₂.

- **Aquifer Zone IIB₁** Aquifer zone IIB₁ has been referred to as the "Barnwell/McBean" Aquifer in previous reports. This zone ranges in thickness from 39 to 91 feet (Table 5). The isopach of the unit indicates thinning of the aquifer towards the west near the F-Area Seepage Basins (Figure 24). Thickening of the aquifer occurs to the southeast near the H-Area Seepage Basins.

TABLE 5
THICKNESS OF
AQUIFER ZONE IIB₁
(BARNWELL/McBEAN AQUIFERS)

Well	Elevation Bottom of Confining Zone IIB ₁ -IIB ₂ ^a (Tan Clay)	Elevation Top of Confining Zone IIA-IIB ^b (Green Clay)	Thickness of Aquifer Zone IIB ₁ (Barnwell/ McBean)
BGO 1D	--	--	--
BGO 3D	--	--	--
BGO 5C	215.2	--	--
BGO 6A	194.6	138.8	56
BGO 8A	197.3	124.3	73
BGO 10A	199.1	132.1	67
BGO 12A	215.4	143.4	72
BGO 14A	209.2	136.2	73
BGO 16A	187.8	130.8	57
BGO 18A	198.9	131.9	67
BGO 21D	199.0	--	--
BGO 23D	--	--	--
BGO 25A	204.7	137.7	67
BGO 26A	198.1	132.1	66
BGO 27C	190.9	--	--
BGO 29A	188.1	124.1	64
BGO 30C	190.6	--	--
BGO 31C	185.1	--	--
BGO 33C	193.4	--	--
BGO 35C	194.4	--	--
BGO 37C	192.3	--	--
DRB 4	190.0	117.0	73
FSB 76A	182.5	123.5	59
FSB 78A	145.5	106.5	39
FSB 79A	163.1	106.1	57
FSB 87A	176.6	114.6	62
FSB 88C	180.4	--	--
FSB 89C	179.1	--	--
FSB 90C	178.1	--	--
FSB 91C	162.0	--	--
FSB 92D	142.6	--	--
FSB 93C	155.0	--	--
FSB 94C	151.8	--	--
FSB 95C	156.8	--	--
FSB 96A	160.7	108.7	52
FSB 97A	152.8	109.8	43
FSB 98A	161.7	109.7	52
FSB 99A	175.3	115.3	60

TABLE 5
THICKNESS OF
AQUIFER ZONE IIB₁
(BARNWELL/McBEAN AQUIFERS)
(Continued)

Well	Elevation Bottom of Confining Zone IIB ₁ -IIB ₂ (Tan Clay)	Elevation Top of Confining Zone IIA-IIB (Green Clay)	Thickness of Aquifer Zone IIB ₁ (Barnwell/ McBean)
FSB 100A	181.8	118.8	63
FSB 101A	184.9	118.9	66
FSB 102C	--	--	--
FSB 103C	159.1	--	--
FSB 104C	170.7	--	--
FSB 105C	162.5	--	--
FSB 106C	173.0	--	--
FSB 107C	166.8	--	--
FSB 111C	175.0	--	--
FSB TA	186.4	118.4	68
HC 12A	190.3	--	--
HPT 1A	NCU	121.9	--
HPT 2A	NCU	121.7	--
HPC 1	186.5	116.5	70
HSB 65A	203.7	124.7	79
HSB 68A	189.4	116.4	73
HSB 69A	181.1	115.1	66
HSB 70C	173.9	--	--
HSB 71C	180.4	--	--
HSB 83A	188.9	116.9	72
HSB 84A	182.4	118.4	64
HSB 85A	197.1	127.1	70
HSB 86A	185.4	116.0	69
HSB 100C	193.0	--	--
HSB 101C	189.3	--	--
HSB 102C	180.7	--	--
HSB 103C	181.2	--	--
HSB 104C	183.5	--	--
HSB 105C	176.2	--	--
HSB 106C	177.7	--	--
HSB 107C	182.3	--	--
HSB 108C	202.9	--	--
HSB 109C	189.4	--	--
HSB 110C	188.4	--	--
HSB 111C	171.7	--	--
HSB 112C	180.6	--	--
HSB 113C	173.7	--	--

TABLE 5
THICKNESS OF
AQUIFER ZONE IIB₁
(BARNWELL/McBEAN AQUIFERS)
(Continued)

Well	Elevation Bottom of Confining Zone IIB ₁ -IIB ₂ (Tan Clay)	Elevation Top of Confining Zone IIA-IIB (Green Clay)	Thickness of Aquifer Zone IIB ₁ (Barnwell/ McBean)
HSB 114C	195.5	--	--
HSB 115C	195.8	--	--
HSB 116C	195.3	--	--
HSB 117A	189.1	122.1	67
HSB 118A	182.0	118.0	64
HSB 119A	193.8	113.8	80
HSB 120A	195.0	111.0	84
HSB 121A	191.3	114.3	77
HSB 122A	179.4	110.4	69
HSB 123A	NCU	--	--
HSB 124A	NCU	--	--
HSB 125C	178.6	--	--
HSB 126C	NCU	--	--
HSB 127C	168.4	--	--
HSB 129C	179.8	--	--
HSB 130C	190.9	--	--
HSB 131C	178.5	--	--
HSB 132C	NCU	--	--
HSB 133C	200.4	--	--
HSB 134C	188.1	--	--
HSB 135C	180.0	--	--
HSB 136C	174.9	--	--
HSB 137C	172.8	--	--
HSB 139A	177.5	116.5	61
HSB TB	203.0	112.1	91
IDL 3A	NCU	--	--
P18 TA	168.7	122.7	46
P27 TA	194.1	132.1	62
P28 TA	202.2	140.2	62

NOTE: 1. -- Data not available
2. NCU No confining unit present

Figure 24. Isopach Map of Aquifer Zone IIB₁ (Barnwell/McBean Aquifers)

This aquifer zone consists predominantly of sand and clayey sand sequences with localized zones of abundant calcareous sands, sandy and muddy limestones, and limestones. Minor interbeds of clay and sandy clay are also present. Sands and clayey sands range in thickness from 2 to 60 feet. Calcareous sands range in thickness from 2 to 33 feet. Sandy and muddy limestones and limestones range in thickness from 3 to 30 feet. Figure 18 illustrates the thickness of the two calcareous units in this zone. Figure 24 illustrates the thickness of the entire Aquifer zone IIB₁. Clay and sandy clays range in thickness from 1 to 16 feet. The sand and clayey sand beds are generally yellow to tan, occasionally greenish brown to light brown. The sand is fine to coarse grained, moderately to well sorted, and generally subangular.

The calcareous sands are white to buff in color and contain up to 50 percent calcareous materials. Shell fragments (gastropods, pelecypods, echinoderms, bryozoans, and barnacles) and whole foraminifera are common in the sands. The sand is quartz rich and is generally fine grained, subangular, and well to moderately sorted. Small percentages of glauconite are common.

The sandy and muddy limestones and limestones are white to buff in color and contain greater than 80 percent calcareous material. Glauconite is common in this unit. Many of the limestones are partially to fully consolidated with abundant moldic porosity from fossil molds. Shells are dominantly pelecypods and gastropods. Unconsolidated limestones are generally a "coquina" type shell hash consisting of shell fragments of pelecypods, gastropods, echinoderms, bryozoans, and barnacles. Small silicified layers of this shell hash (<0.5 feet) are occasionally observed in this unit.

The fossiliferous limestones are typical of bioherms observed on the shallow marine shelf. Modern analogues are found in Florida Bay near the Keys (Baars, 1963), where shell mounds are commonly observed surrounded by the silty clayey sands of the shallow shelf. The carbonate

buildups in the GSA are oriented northwest-southeast, paralleling the strike of the shoreline at that time.

The interbeds of clays and sandy clays in this unit are light tan to orange in color. According to Dennehy, et al., (1989) the clay minerals are illite/smectite. The sand is quartz, fine to medium grained, subangular, and well to moderately sorted.

The calcareous sands and limestones contained in this unit are generally not recognizable on natural gamma curves in the GSA. Occasionally, a limestone with good moldic porosity will be distinct on the electric resistivity curve (Figure 25). The clastic lithofacies of Aquifer zone IIB₁ are not easily distinguishable on natural gamma curves alone. Correlation in this zone requires careful examination and interpretation of both core and geophysical data.

- **Faulting Associated With Carbonate Buildups.** A prominent feature on Cross Section #3 (Plate 3) is the offset of beds between FSB-91 and FSB-100. The beds are down thrown approximately 12 feet to the south. The offset between the wells is typical of the shallow "faulting" observed in the GSA. Most often the areas where offset has occurred are related to the flanks of the carbonate buildups found at and near the base of Aquifer zone IIB₁. Geophysical and core data both corroborate the offset in Aquifer zone IIB₁, Confining zone IIB₁ - IIB₂, and Aquifer zone IIB₂, but offset below the carbonate buildups in Aquifer IIA cannot be demonstrated. The precise age and most recent movement on the shallow offsets is not well documented and awaits the results of the Upland Unit study and Phase II of the Pen Branch fault study (see p.23). Until the results of the two studies are known, the upper or shallow portions of the offsets that underlie the areas at higher evaluations in the GSA have been dashed (Plates 2, 5A, 5B, 7, 10A, 10B). The slump features in the valleys and hill sides are not dashed since they penetrate to the surface due to recent erosion.

Figure 25. Geophysical Characteristics of Limestone in HSB-122

Siple (1967) speculated that dissolution of calcareous material in the Santee Limestone has occurred at the SRS and caused subsidence of overlying beds and the formation of depressions. He noted that voids and loosely compacted sediments were encountered during well drilling and mentioned that large amounts of cement grout were used to stabilize the subsurface before construction of heavy structures. Slump features have been recognized in the coastal plain calcareous sediments by Johnson and Heron (1965). In all probability the shallow offsets noted at the GSA are due to slumping of the section above the carbonate buildups due to carbonate dissolution. However, soft sediment deformation such as deltaic slumping is another feasible explanation for the offset (McDowell and Houser, 1983). The localized steepening of dip observed near borings FSB-106, FSB-91, and FSB-100 in the Aquifer zone IIB₁ interval could be caused by the proposed slumping due to carbonate dissolution. Alternately, the overlying clastic sediments may simply be draped over the flanks of the carbonate mounds.

A similar feature on Cross Section #5B (Plate 5B) is the offset beds between FSB-97 and FSB-98. The feature exhibits normal offset, and beds are downthrown to the west with a throw of approximately 9 feet in Aquifer zone IIB₂. Throw decreases to around 3 feet in Confining zone IIB₁-IIB₂ and Aquifer zone IIB₁. The offset apparently does not penetrate Confining Unit IIA-IIB. The offset cuts FSB-97 at 121 feet above msl. Geophysical and core data both corroborate this offset. Again, there is a good possibility of slumping associated with carbonate dissolution as previously discussed. Due to the close proximity of calcareous beds in the nearby boring FSB-100, there is a strong possibility that borings FSB-97 and FSB-98 once contained carbonate beds, and post-depositional carbonate dissolution is a reasonable explanation for the offset observed between FSB-97 and FSB-98.

- **Confining Zone IIB₁ - IIB₂ (Tan Clay).** Confining zone IIB₁-IIB₂ is commonly referred to as the "Tan Clay" in previous reports at SRS. This zone consists of multiple lenses of clay that thicken, thin, and pinch out abruptly and within very short distances. This zone ranges in

thickness from 0 to 33 feet (Table 6 and Figure 26). As indicated, no confining zone was encountered in a group of wells east of the MWMF and in an area west of the F-Area Seepage Basins. At the H-Area Seepage Basins there are three small areas where no confining zone within this interval was identified. Elsewhere within the GSA the zone thickens and thins abruptly. Figure 27 is a structure contour map on top of the confining zone. The structure contours suggest that the thinning of the zone at the H-Area Seepage Basins may be a result of post-depositional erosion.

Confining zone IIB₁ - IIB₂ consists predominantly of clays and sandy clays, with minor to equal interbeds of clayey sands and sands. One occurrence of a 2 foot thick calcareous sand was encountered at wells monitoring the MWMF. Clays and sandy clays range in thickness from 1 to 11 feet. Clayey sands and sands range in thickness from 1 to 8 feet.

Clays and sandy clays are light tan to light green tan in color. According to Dennehy and others (1989), the clay is predominantly composed of kaolinite and tends to be sticky and plastic. The quartz sand is angular to subangular and contains mica and dark heavy minerals. The heavy mineral suite is brown ilmenite, zircon, brown hornblende, staurolite, sillimanite, kyanite, and magnetite. Clay bedding exhibits thin, irregular laminae.

Clayey sands and sands are light tan to light orange in color. The clay matrix in the sand is predominantly kaolinite (Dennehy, et al., 1989). The quartz sands are predominantly medium to coarse grained, with a moderate degree of sorting. Grain size tends to coarsen upward.

The calcareous sand is composed of shell fragments and quartz sand in a clay matrix. The quartz sand is fine grained and subrounded. The thicker clays within the confining zone exhibit a natural gamma spike typical of clays. However, natural gamma spikes may also be related to concentrations of heavy minerals.

Figure 26. Isopach Map for Confining Zone IIB₁ - IIB₂ (Tan Clay)

Figure 27. Structure Contour Map for Confining Zone IIB₁- IIB₂

TABLE 6
DETAILS OF CONFINING UNIT IIB₁-IIB₂ (TAN CLAY)
AND THE SATURATED PORTION OF
AQUIFER ZONE IIB₂ (WATER TABLE AQUIFER)

Well	Ground Elev.	Water Elev. IIB ₂ ^a	Elev. Top Of Confining Zone IIB ₁ -IIB ₂ ^b	Elev. Bottom of Confining Zone IIB ₁ -IIB ₂ ^b	Thickness of Confining Zone IIB ₁ -IIB ₂ ^{b,d}	Saturated Thickness IIB ₂ ^c
BGO 1D	293.0	237.5	NP	NP	--	--
BGO 3D	290.8	235.4	NP	NP	--	--
BGO 5C	294.2	231.3	218.2	215.2	3	13.1
BGO 6A	283.8	231.4	209.6	194.6	15	21.8
BGO 8A	281.3	231.7	213.3	197.3	16	18.4
BGO 10A	299.1	DRY	218.1	199.1	19	--
BGO 12A	311.4	229.4	226.4	215.4	11	3.0
BGO 14A	300.2	DRY	223.2	209.2	14	--
BGO 16A	302.8	229.7	214.8	187.8	27	14.9
BGO 18A	292.9	231.0	204.9	198.9	6	26.1
BGO 21D	283.0	233.9	212.0	199.0	13	21.9
BGO 23D	287.0	235.3	NP	NP	--	--
BGO 25A	294.7	--	214.7	204.7	10	--
BGO 26A	285.1	225.7	214.1	198.1	16	11.6
BGO 27C	273.9	225.5	197.9	190.9	7	27.6
BGO 29A	262.1	--	197.1	188.1	9	--
BGO 30C	272.6	223.6	200.6	190.6	10	23.0
BGO 31C	271.1	224.6	195.1	185.1	10	29.5
BGO 33C	277.4	227.9	201.4	193.4	8	26.5
BGO 35C	271.4	232.2	213.4	194.4	19	18.8
BGO 37C	284.3	235.1	201.3	192.3	9	33.8
DRB 4	250.0	--	204.0	190.0	14	--
FSB 76A	291.5	--	192.5	182.5	20	--
FSB 78A	270.5	207.4	162.5	145.5	17	44.9
FSB 79A	216.1	201.6	173.1	163.1	10	28.5
FSB 87A	285.6	--	175.6	176.6	3	--
FSB 88C	280.4	--	188.4	180.4	8	--
FSB 89C	279.1	213.0	186.1	179.1	7	26.9
FSB 90C	276.1	212.0	185.1	178.1	7	26.9
FSB 91C	277.0	211.2	169.0	162.0	7	42.2
FSB 92D	273.6	210.0	167.6	142.6	25	42.4
FSB 93C	274.0	209.1	169.0	155.0	14	40.1
FSB 94C	278.8	--	173.8	151.8	22	--
FSB 95C	281.8	208.3	172.8	156.8	16	35.5
FSB 96A	277.7	--	169.7	160.7	9	--
FSB 97A	283.8	209.0	165.8	152.8	13	43.2
FSB 98A	280.7	209.9	174.7	161.7	13	35.2
FSB 99A	285.3	210.8	180.3	175.3	5	30.5
FSB 100A	283.8	--	184.8	181.8	3	--
FSB 101A	282.9	--	191.9	184.9	7	--
FSB 102C	198.9	--	NP	NP	--	--
FSB 103C	240.1	--	173.1	159.1	14	--

TABLE 6
THICKNESS OF CONFINING UNIT IIB₁-IIB₂ (TAN CLAY)
AND THE SATURATED PORTION OF
AQUIFER ZONE IIB₂ (WATER TABLE AQUIFER)
(Continued)

Well	Ground Elev.	Water Elev. IIB ₂ ^a	Elev. Top Of Confining Zone IIB ₁ -IIB ₂ ^b	Elev. Bottom of Confining Zone IIB ₁ -IIB ₂ ^b	Thickness of Confining Zone IIB ₁ -IIB ₂ ^{b,d}	Saturated Thickness IIB ₂ ^c
FSB 104C	216.7	203.1	172.7	170.7	2	30.4
FSB 105C	283.5	208.5	173.5	162.5	11	35.0
FSB 106C	233.0	--	179.5	173.0	7	--
FSB 107C	268.8	211.5	180.8	166.8	14	30.7
FSB 111C	274.0	212.9	186.0	175.0	11	26.9
FSB TA	275.4	--	190.4	186.4	4	--
HC 12A	287.3	--	205.3	190.3	15	--
HPT 1A	232.9	--	NCU	NCU	0	--
HPT 2A	257.7	--	NCU	NCU	0	--
HPC 1	293.5	--	192.5	186.5	6	--
HSB 65A	270.7	231.0	208.7	203.7	5	22.3
HSB 68A	247.4	218.6	200.4	189.4	11	18.2
HSB 69A	234.1	219.6	187.1	181.1	6	32.5
HSB 70C	240.9	--	190.9	173.9	17	--
HSB 71C	239.4	225.1	193.4	180.4	13	31.7
HSB 83A	234.9	226.0	196.9	188.9	8	29.1
HSB 84A	226.4	219.0	204.4	182.4	22	14.6
HSB 85A	292.1	237.5	207.1	197.1	10	30.4
HSB 86A	260.0	223.9	211.4	185.4	26	12.5
HSB 100C	258.0	--	199.0	193.0	6	--
HSB 101C	256.3	229.5	195.3	189.3	6	34.2
HSB 102C	256.7	228.0	192.7	180.7	12	35.3
HSB 103C	245.2	225.4	195.2	181.2	14	30.2
HSB 104C	245.5	--	194.5	183.5	11	--
HSB 105C	247.2	225.4	191.2	176.2	15	34.2
HSB 106C	250.7	--	193.7	177.7	16	--
HSB 107C	258.3	--	199.3	182.3	17	--
HSB 108C	263.9	--	212.9	202.9	10	--
HSB 109C	259.4	--	203.4	189.4	14	--
HSB 110C	253.4	--	203.4	188.4	15	--
HSB 111C	253.7	--	186.7	171.7	15	--
HSB 112C	252.6	--	189.6	180.6	9	--
HSB 113C	258.7	--	187.7	173.7	14	--
HSB 114C	261.5	215.0	208.5	195.5	13	15.0
HSB 115C	266.8	--	210.8	195.8	15	--
HSB 116C	255.3	216.4	209.3	195.3	14	16.4
HSB 117A	234.1	--	215.1	189.1	26	--
HSB 118A	245.0	--	214.0	182.0	32	--
HSB 119A	254.8	--	212.8	193.8	18	--
HSB 120A	266.0	--	213.0	195.0	18	--
HSB 121A	272.3	--	197.3	191.3	6	--
HSB 122A	269.4	--	194.4	179.4	15	--

TABLE 6
THICKNESS OF CONFINING UNIT IIB₁-IIB₂ (TAN CLAY)
AND THE SATURATED PORTION OF
AQUIFER ZONE IIB₂ (WATER TABLE AQUIFER)
(Continued)

Well	Ground Elev.	Water Elev. IIB ₂ ^a	Elev. Top Of Confining Zone IIB ₁ -IIB ₂ ^b	Elev. Bottom of Confining Zone IIB ₁ -IIB ₂ ^b	Thickness of Confining Zone IIB ₁ -IIB ₂ ^{b,d}	Saturated Thickness IIB ₂ ^c
HSB 123A	262.3	--	NCU	NCU	0	--
HSB 124A	263.9	--	NCU	NCU	0	--
HSB 125C	229.6	--	193.6	178.6	15	--
HSB 126C	210.3	205.7	NCU	NCU	0	0
HSB 127C	223.4	--	190.4	168.4	22	--
HSB 129C	212.8	208.7	199.8	179.8	20	8.9
HSB 130C	215.9	199.5	215.9	190.9	25	0
HSB 131C	209.5	205.9	193.5	178.5	15	12.4
HSB 132C	238.3	--	NCU	NCU	0	--
HSB 133C	253.4	223.4	204.4	200.4	4	19.0
HSB 134C	236.1	221.9	194.1	188.1	6	27.8
HSB 135C	229.8	--	197.0	180.0	17	--
HSB 136C	225.6	220.8	197.9	174.9	23	22.9
HSB 137C	233.8	222.6	205.8	172.8	33	16.8
HSB 139A	231.5	--	188.5	177.5	11	--
HSB TB	267.1	--	207.1	203.1	4	--
IDL 3A	257.4	--	NCU	NCU	0	--
P18 TA	296.7	218.9	185.7	168.7	17	33.2
P27 TA	274.1	265.9	200.1	194.1	6	65.8
P28 TA	285.2	222.0	215.2	202.2	13	0

NOTES:

1. -- Not applicable
2. a Water Elevation of Aquifer zone IIB₂ (Water Table Aquifer)
3. b Elevation and thickness of Confining Zone IIB₁-IIB₂ (Tan Clay)
4. c Thickness of the saturated portion of Aquifer zone IIB₂ (Water Table Aquifer)
5. d Rounded to the nearest whole foot
6. NCU No confining unit present
7. DRY Dry well
8. NP Not penetrated

- **Aquifer Zone IIB₂** Aquifer zone IIB₂ represents the "water table aquifer". The total thickness of the lithologies of this zone range from 0 feet at Fourmile Branch to 110 feet (Table 6). An isopach of the saturated portion of this zone is presented in Figure 28.

This zone consists predominantly of sands and clayey sands with lesser or equal amounts of clay and sandy clay. Gravel and pebble layers occur locally. Sands and clayey sands range in thickness from 2 to 107 feet. Clays and sandy clays range in thickness from 1 to 18 feet. Gravel and pebble layers range in thickness from 1 to 5 feet. Sands and clayey sands are tan to red to purple and are often highly variegated in color. Sands are fine to very coarse, subangular to angular, and moderately to poorly sorted. Clays and sandy clays are composed predominantly of kaolinite (Dennehy, et al., 1989). Sands are fine to very coarse and subangular to angular. Clay clasts, mica, and dark heavy minerals are also present in the quartz sand. Brown ilmenite and sillimanite are very abundant in the typical heavy mineral assemblage along with zircon, rutile, brown hornblende, and staurolite (Dennehy, et al., 1989).

Clasts within the gravel and pebble beds range in diameter from 0.5 up to 4 inches and are generally well rounded. Gravels are usually found in a matrix of fine to very coarse clayey quartz sand containing abundant mica and white clay balls. The quartz sand is generally coarse and angular to subangular.

The unit is difficult to correlate on geophysical data alone. As in Confining zone IIB₁-IIB₂ (Tan Clay), the natural gamma spikes resembling clays may be attributed to concentrations of potassium-rich heavy minerals. Core and geophysical data were both carefully analyzed in the interpretation and correlations of Aquifer Unit IIB₂.

Figure 28. Isopach Map of Aquifer Zone IIB₂ (Water Table Aquifer)

V. HYDROLOGICAL CHARACTERISTICS OF THE GSA

A. Identification of the "Uppermost Aquifer"

All of the aquifers and confining units above Confining System I-II are included in the "uppermost aquifer" (Figure 12). The RCRA Ground Water Monitoring Technical Enforcement Guidance Document (TEGD) defines the "uppermost aquifer" as "the geologic formation, group of formations, or part of a formation that is the aquifer nearest to the ground surface and is capable of yielding significant amount of ground water to wells or springs" (Chapter 1, Section 1.4, p. 35). The TEGD emphasizes that identification of the lower confining layer (or / boundary) is essential to the proper definition of the "uppermost aquifer". The TEGD also notes that the degree of hydraulic interconnection between the "uppermost aquifer" and lower aquifers must be limited. Using this guidance, the "uppermost aquifer" beneath the GSA includes all the hydrostratigraphic units which lie between the Aquifer zone IIB₂ and the base of Aquifer Unit IIA (Figure 12). Aquifer Unit IIA is the basal hydrostratigraphic unit in the "uppermost aquifer". There are three aquifer units separated by two confining units within the "uppermost aquifer". The Principal Confining Unit which underlies Aquifer Unit IIA is Confining System I-II.

Potentiometric maps were prepared for each of the three aquifer units in the GSA (Figures 29, 30, and 31) at a scale of 1"=1000'. Vertical flow potential across the confining units in the area are illustrated on hydraulic head difference maps in Figures 32 and 33. Water elevation measurements used to produce these maps were taken during the fourth quarter of 1989.

**Figure 29. Potentiometric Surface Map-
Aquifer Unit IIB₂ (Water Table Aquifer)**

**Figure 30. Potentiometric Surface Map -
Aquifer Zone IIB₁ (Barnwell/McBean Aquifers)**

**Figure 31. Potentiometric Surface Map -
Aquifer Unit IIA (Congaree Aquifer)**

Figure 32. Head Difference Between Aquifers IIB₁ and IIB₂

Figure 33. Head Difference Between Aquifers IIA and IIB₁

1) Aquifer Zone IIB₂

Aquifer zone IIB₂ consists of all the saturated strata which lie between the water table and Confining zone IIB₁-IIB₂. This aquifer is included in the "uppermost aquifer" because it is the first hydrostratigraphic unit which lies beneath the GSA. This zone has a general downward potential into the lower aquifer units. A "water table" map is depicted in Figure 29. A ground water divide occurs at the westerly trending topographic high crossing the MWMF (Figure 3). Ground water to the north of the divide flows toward Upper Three Runs Creek. To the south of the divide, ground water flows south toward Fourmile Branch. The F and H Area Seepage Basins are located south of the ground water divide. Horizontal flow in Aquifer Zone IIB₂ at the seepage basins flows toward and outcrops into Fourmile Branch and the wetlands adjacent to the creek. Water table elevations in the GSA range from 199.5 feet near Fourmile Branch to 266 feet above msl at well P-27 (Figure 29). Natural recharge to the water table is from percolation of rainwater. Mounding resulting from excessive local recharge due to water in the seepage basins is evident beneath the basins in both F and H Areas. The steepest hydraulic gradients in the water table are observed between H Area Seepage Basins and Fourmile Branch.

The vertical component for the flow of water in the water table is downward across Confining Zone IIB₁-IIB₂ into Aquifer Zone IIB₁ (Table 7). Confining Zone IIB₁-IIB₂ is a leaky aquitard. The degree of leakage across the confining zone is variable. Leakage depends on the magnitude of the head difference across the confining zone and the local competency of the confining unit. Figure 32 indicates the head differences across the Confining Zone IIB₁-IIB₂. The lateral variability in the sediments that make up the confining zone and offset from faulting which breaches the confining unit at both the F and H Area Seepage Basins, provide the preferential pathways for the vertical migration of water and chemical constituents.

TABLE 7
HEAD DIFFERENCE BETWEEN
AQUIFER ZONE IIB₂ (WATER TABLE AQUIFER)
AND AQUIFER ZONE IIB₁ (BARNWELL/McBEAN AQUIFERS)

Well Cluster	Water Elevation IIB ₂ ^a (Water Table)	Potentiometric Surface IIB ₁ ^b (Barnwell/McBean)	Head Difference*
BGO 1D	237.5	--	--
BGO 3D	235.4	--	--
BGO 5C	231.3	215.8	15.5
BGO 6A	231.4	219.5	11.9
BGO 8A	231.7	222.3	9.4
BGO 10	DRY	--	--
BGO 12	229.4	219.7	9.7
BGO 14	DRY	--	--
BGO 16	229.7	--	--
BGO 18	231.0	--	--
BGO 21	233.9	--	--
BGO 23	235.3	--	--
BGO 25	--	--	--
BGO 26	225.7	--	--
BGO 27	225.5	--	--
BGO 29	--	--	--
BGO 30	223.6	219.1	4.5
BGO 31	224.6	--	--
BGO 33	227.9	223.5	4.4
BGO 35	232.2	227.6	4.6
BGO 37	235.1	228.8	6.3
DRB 4	--	--	--
FSB 76	--	211.1	--
FSB 77	211.0	--	--
FSB 78	207.4	206.3	1.1
FSB 79	201.6	196.4	5.2
FSB 87	--	207.0	--
FSB 88	214.0	210.5	3.5
FSB 89	213.0	209.3	3.7
FSB 90	212.0	208.8	3.2
FSB 91	211.2	208.8	2.4
FSB 92	210.0	--	--
FSB 93	209.1	206.9	2.2
FSB 94	--	--	--
FSB 95	208.3	205.3	3.0
FSB 96	--	--	--
FSB 97	209.0	206.2	2.8
FSB 98	209.9	206.7	3.2
FSB 99	210.8	207.4	3.4
FSB 100	--	--	--
FSB 101	--	--	--
FSB 102	--	195.0	--
FSB 103	--	201.2	--

TABLE 7
HEAD DIFFERENCE BETWEEN
AQUIFER ZONE IIB₂ (WATER TABLE AQUIFER)
AND AQUIFER ZONE IIB₁ (BARNWELL/McBEAN AQUIFERS)
(Continued)

Well Cluster	Water Elevation IIB ₂ ^a (Water Table)	Potentiometric Surface IIB ₁ ^b (Barnwell/McBean)	Head Difference*
FSB 104	203.1	199.5	3.6
FSB 105	208.5	205.3	3.2
FSB 106	--	201.0	--
FSB 107	211.5	208.5	3.0
FSB 108	215.0	--	--
FSB 109	211.3	--	--
FSB 110	204.2	200.3	3.9
FSB 111	212.9	210.1	2.8
FSB TA	--	--	--
HC 12A	--	--	--
HPT 1A	--	--	--
HPT 2A	--	--	--
HPC 1	--	--	--
HSB 65	--	--	--
HSB 66	--	--	--
HSB 67	224.5	--	--
HSB 68	--	217.4	--
HSB 69	219.6	--	--
HSB 70	225.4	224.0	1.4
HSB 71	225.1	223.0	2.1
HSB 83	226.0	224.0	2.0
HSB 84	219.0	213.7	5.3
HSB 85	--	237.5	--
HSB 86	--	--	--
HSB 100	232.2	226.0	6.6
HSB 101	229.5	224.4	5.1
HSB 102	228.0	223.5	4.5
HSB 103	225.4	223.0	2.4
HSB 104	225.5	220.0	5.5
HSB 105	225.4	219.2	6.2
HSB 106	225.4	221.1	4.3
HSB 107	224.2	219.2	5.0
HSB 108	223.1	218.0	5.1
HSB 109	222.4	218.3	4.1
HSB 110	222.0	219.0	3.0
HSB 111	221.9	220.1	1.8
HSB 112	221.8	221.5	0.3
HSB 113	222.7	221.8	0.9
HSB 114	223.5	223.5	--
HSB 115	224.0	224.0	--
HSB 116	225.7	225.2	0.5

TABLE 7

**HEAD DIFFERENCE BETWEEN
AQUIFER ZONE IIB₂ (WATER TABLE AQUIFER)
AND AQUIFER ZONE IIB₁ (BARNWELL/McBEAN AQUIFERS)
(Continued)**

Well Cluster	Water Elevation IIB ₂ ^a (Water Table)	Potentiometric Surface IIB ₁ ^b (Barnwell/McBean)	Head Difference*
HSB 117	225.0	221.7	3.3
HSB 118	--	--	--
HSB 119	--	--	--
HSB 120	--	--	--
HSB 121	--	--	--
HSB 122	--	--	--
HSB 123	--	--	--
HSB 124	--	--	--
HSB 125	--	--	--
HSB 126	205.7	203.6	2.1
HSB 127	218.6	209.7	8.9
HSB 129	208.7	205.5	3.2
HSB 130	200.0	--	--
HSB 131	206.0	203.2	2.8
HSB 132	--	--	--
HSB 133	234.0	229.3	1.7
HSB 134	222.0	220.3	1.7
HSB 135	218.1	206.0	12.1
HSB 136	220.8	217.1	3.7
HSB 137	222.6	220.5	2.1
HSB 138	223.3	--	--
HSB 139	223.3	214.2	9.1
IISB TB	--	--	--
IDL 3A	--	--	--
P18 TA	219.0	--	--
P27 TA	265.9	240.0	25.9
P28 TA	222.0	213.6	8.4

NOTES:

1. DRY Dry well
2. -- Not applicable
3. * Head difference calculated from fourth quarter 1989 data
4. a Elevation of Aquifer zone IIB₂ (Water Table Aquifer)
5. b Potentiometric surface of Aquifer zone IIB₁
(Barnwell/McBean Aquifers)

Hydraulic conductivity estimates for this aquifer zone are based on slug tests, sieve analyses, and pumping tests. Slug tests have been performed on many of the water table wells installed since 1986. The values for the 60 slug tests performed range from a high of 45.40 to a low 0.05 feet per day. The arithmetic means were calculated at each of the three areas: MWMF, F Area Basins, and H Area Basins. The calculated hydraulic conductivities are:

1. MWMF – 2.35 feet/day, n = 17. Range: 0.10 to 15.15 feet/day.
2. F Area Seepage Basins – 2.78 feet/day, n = 15. Range: 0.05 to 23.11 feet/day.
3. H Area Seepage Basins – 10.64 feet/day, n = 28. Range 0.08 to 45.40 feet/day.

The arithmetic mean for the complete dataset (60 values) is 5.26 feet/day.

Two series of single well pumping tests have been performed on Aquifer zone IIB₂. Parizek and Root (1986) conducted thirty-eight single well pump tests in the vicinity of the GSA. Hydraulic conductivity values calculated from these tests were reported to range from 0.30 to 3.6 feet/day with a median value of 0.61 feet/day. A more recent series of tests were performed by Evans (work in progress) which produced values much higher than the highest value of 3.6 feet/day from the Parizek and Root study of 1986.

Sieve analyses are routinely performed on samples of materials obtained from the screened zones of the wells installed for RCRA monitoring. The grain-size analyses have been used to estimate hydraulic conductivity with the Beard and Weyl method, 1973. The values were compiled and arithmetic means were calculated at each of the three areas: MWMF, F Area Basins, and H Area Basins. The means and basis for the calculations are presented by facility as follows:

1. MWMF – 123 feet/day, n = 5. Range: 27 to 315 feet/day.
2. F Area Seepage Basins – 163 feet/day, n = 7. Range: 55 to 479 feet/day.
3. H Area Seepage Basins – 84 feet/day, n = 8. Range 13 to 149 feet/day.

The mean of the complete dataset is 123 feet/day, n = 20, the values range from 13 to 479 feet/day (Thayer, et al., 1990). The method gives elevated estimates of hydraulic conductivity, eg. slug test results for the interval range from .05-45 ft/day, the pump tests .3-3.6 ft/day.

2) Aquifer Zone IIB₁

Aquifer zone IIB₁ consists of all saturated strata which lie between Confining zone IIB₁-IIB₂ and Confining Unit IIA-IIB. This zone is poorly confined and included in the "uppermost aquifer" because it is recharged primarily by downward flow from Aquifer zone IIB₂ above.

The groundwater divide also exists in Aquifer zone IIB₁. Horizontal flow in this aquifer zone is south toward Fourmile Branch in the vicinity of the F and H Area Seepage Basins (Figure 30). Measured water elevations in the GSA range from 195 to 240 feet above mean sea level (msl). The hydraulic gradient in this zone is steepest between the H Area Seepage Basins and Fourmile Branch. The recharge to this unit is from leakage through Confining zone IIB₁-IIB₂.

The vertical component of flow in Aquifer zone IIB₁ is downward across the Confining Unit IIA-IIB into Aquifer Unit IIA. Confining Unit IIA-IIB is more competent than Confining Zone IIB₁-IIB₂, but there is evidence of leakage where the clay beds thin and are discontinuous and where the clays have been breached by slump features. Head differences across Confining Unit IIA-IIB are illustrated on Figure 33 and Table 8.

Hydraulic conductivity estimates for Aquifer zone IIB₁ are based on slug tests, sieve analyses, and pumping tests. The range in the values for the 46 slug tests performed is from a high of 136.0 to a low of 0.13 feet per day (Sirrinc, 1987, 1988; S&ME, 1988 a,b). The arithmetic means were calculated at each of the three areas: MWMF, F Area Seepage Basins, and H Area Seepage Basins. Arithmetic means of the test results were calculated and are presented by facility as follows:

1. MWMF – 1.01 feet/day, n = 3. Range: 0.13 to 1.51 feet/day.
2. F Area Seepage Basins – 3.97 feet/day, n = 16. Range: 0.14 to 24.03 feet/day.
3. H Area Seepage Basins – 12.92 feet/day, n = 27. Range 0.25 to 136.10 feet/day.

Parizek and Root (1986) conducted a series of 51 single-well pump tests from wells screened in Aquifer zone IIB₁ in and around the GSA. Hydraulic conductivity values as reported from the tests range from 0.17 feet/day to 2.6 feet/day with a median value of 0.90 feet/day.

Sieve analyses are routinely performed on samples of materials obtained from the screened zones of the wells installed for RCRA monitoring. The grain-size analyses have been used to estimate hydraulic conductivity using the Beard and Weyl Method, 1973. The values were compiled and arithmetic means were calculated at each of the three areas: MWMF, F Area Seepage Basins, and H Area Seepage Basins. The means and basis for the calculations are presented by facility as follows:

1. MWMF – 110 feet/day, n = 4. Range: 37 to 192 feet/day.
2. F Area Seepage Basins – 199 feet/day, n = 11. Range: 90 to 397 feet/day.
3. H Area Seepage Basins – 271 feet/day, n = 2. Range 185 to 357 feet/day.

TABLE 8
HEAD DIFFERENCE BETWEEN
AQUIFER ZONE IIB₁ (BARNWELL/McBEAN AQUIFERS)
AND AQUIFER UNIT IIA (CONGAREE AQUIFER)

Well Cluster	Potentiometric Surface Aquifer Zone IIB ₁ (Barnwell/McBean)	Potentiometric Surface Aquifer Unit IIA (Congaree)	Head Difference*
BGO 6	219.5	158.1	61.4
BGO 8	222.3	160.5	61.8
BGO 10	219.3	168.5	50.8
BGO 12	219.7	180.0	39.7
BGO 14	221.7	157.3	64.4
BGO 16	--	159.6	--
BGO 18	--	159.7	--
BGO 25	--	158.8	--
BGO 26	--	--	--
BGO 29	--	--	--
DRB 4	--	--	--
FSB 76	214.7	154.3	60.4
FSB 78	208.1	155.1	53.0
FSB 79	196.2	156.4	39.8
FSB 87	209.7	152.9	56.8
FSB 96	--	152.3	--
FSB 97	206.2	151.1	55.1
FSB 98	206.7	150.3	56.4
FSB 99	207.4	149.7	57.7
FSB 100	--	150.6	--
FSB 101	--	150.8	--
FSB TA	--	--	--
HPT 1A	--	--	--
HPT 2A	--	--	--
HPC 1	--	--	--
HSB 65	231.3	169.8	61.5
HSB 68	217.4	169.9	47.5
HSB 69	--	170.3	--
HSB 83	226.0	171.4	54.6
HSB 84	220.8	170.1	50.7
HSB 85	207.6	167.3	40.3
HSB 86	231.6	166.8	64.8
HSB 117	221.7	164.6	57.1
HSB 118	--	166.0	--
HSB 119	--	165.2	--

TABLE 8
HEAD DIFFERENCE BETWEEN
AQUIFER ZONE IIB₁ (BARNWELL/McBEAN AQUIFERS)
AND AQUIFER UNIT IIA (CONGAREE AQUIFER)
(Continued)

Well Cluster	Potentiometric Surface Aquifer Zone IIB ₁ (Barnwell/McBean)	Potentiometric Surface Aquifer Unit IIA (Congaree)	Head Difference*
HSB 120	--	164.7	--
HSB 121	--	170.0	--
HSB 122	--	169.7	--
HSB 123	--	169.8	--
HSB 124	--	192.0	--
HSB 139	214.5	171.7	42.8
HSB TB	--	--	--
IDL 3	--	--	--
P18 TA	--	--	--
P27 TA	--	--	--
P28 TA	--	--	--

NOTES:

1. * Head difference calculated from fourth quarter 1989 data
2. -- Not Applicable

The mean of the overall dataset is 193 feet/day, $n = 17$. The values range from a low of 37 to a high of 397 feet/day (Thayer, et al., 1990).

3) Aquifer Unit IIA

Aquifer Unit IIA consists of all the saturated strata which lie between Confining Unit IIA-IIB and the Principal Confining Unit, Confining System I-II in the GSA (Figure 12). Aquifer Unit IIA is semi-confined and included in the "uppermost aquifer" because a downward potential from Aquifer zone IIB₁ exists, providing recharge and indicating interconnection.

Ground water flow in Aquifer Unit IIA is northwest toward Upper Three Runs Creek as depicted in the potentiometric surface map (Figure 31). Fourmile Branch to the south does not incise Confining Unit IIA-IIB and therefore does not influence flow in the underlying Aquifer IIA . Recharge into Aquifer Unit IIA occurs primarily from leakage through Confining Unit IIA-IIB but also from updip flow.

In places where the confining unit is breached by faulting, this recharge may be locally increased. Evidence of this can be seen south of the H Area Seepage Basins. Perturbations in the equipotential surface (170' contour) of Aquifer Unit IIA (Figure 31) coincide with those in the overlying Aquifer zone IIB₁ (Figure 30) and with the location of slump features, which breach the Confining Unit IIA-IIB. A marked break in the contours of head difference across Confining Unit IIA-IIB in the vicinity of these slump features (Plate 3, Figure 33) is further evidence of localized zones of increased recharge which are controlled by geologic features. Downward flow of water into Aquifer Unit IIA is locally increased.

Discharge is to Upper Three Runs Creek. Any downward component of flow out of this aquifer unit is believed to be minimal. In most of the GSA, an upward flow potential exists between Aquifer Unit IIA and Aquifer System I across the Confining System I-II.

Hydraulic conductivity estimates for this aquifer unit are based on slug tests, sieve analyses, and pump tests. The range in the values for the 8 slug tests performed is from a high of 12.0 to a low of 0.16 feet per day. Arithmetic means of the slug test results were calculated and are presented by facility as follows:

1. MWMF – 4.42 feet/day, n = 3. Range: 0.50 to 11.98 feet/day.
2. F Area Seepage Basins – 0.59 feet/day, n = 2. Range: 0.33 to 0.85 feet/day.
3. H Area Seepage Basins – 6.98 feet/day, n = 3. Range 0.16 to 12.00 feet/day.

The overall mean of the dataset is 4.42 feet/day, n = 8 (Sirrinc, 1987, 1988 a&b; S&ME, 1988).

Aquifer tests have been performed on wells screened in this unit in many locations on the SRS. The aquifer properties discerned from these tests were compiled by Geotrans (1989). Based upon this review, a high degree of variability was noted in the deduced hydraulic conductivity values. Many of the reported ranges of values were taken from transmissivity estimates obtained from small scale pump tests. The variability is therefore more likely attributable to a variable aquifer thickness than in widely ranging permeability differences.

Values (Geotrans, 1989) used for calibrating a GSA area-wide model (38 feet/day) were in the range of the hydraulic conductivity deduced from a recent pumping test performed on well HSB-84A. This test was conducted to gain data relating to aquifer characteristics and ground water chemistry. The test results indicate a range of 30 to 60 feet/day for permeability, and leakage across the Confining System IIA-IIB in this vicinity was suggested (Albenesius, et al., 1990).

A long duration, multiple well test was conducted for the Aquifer Unit IIA in April 1989 (CH2M Hill, 1989). This test employed a fully penetrating pumping well and multiple observation wells screened in the Aquifer Unit IIA and in the overlying Aquifer zone IIB₁. The data indicated that in the vicinity of HPT-2A, Aquifer Unit IIA has a transmissivity of 2290 ft²/day and exhibits a barometric efficiency of 25 percent. The permeability is approximately 37.5 feet/day, utilizing an aquifer thickness of 61 feet. The value agrees with the HSB-84 and HSB-84A pump test. However, a greater permeability is suggested in the HSB-84 vicinity as a result of a thinner aquifer. Transmissivities were very similar. Leakage across the Confining Unit IIA-IIB was suggested in both pumping tests.

Sieve analyses were performed on 17 wells screened in Aquifer Unit IIA within the GSA. Estimates of hydraulic conductivity were made from the sieve data. The overall arithmetic mean for the dataset is 130 feet/day, n = 17. Arithmetic means were also calculated for each of the HWMF facilities.

1. MWMF – 91 feet/day, n = 7. Range: 22 to 205 feet/day.
2. F Area Seepage Basins – 176 feet/day, n = 5. Range: 10 to 479 feet/day.
3. H Area Seepage Basins – 106 feet/day, n = 5. Range 82 to 167 feet/day.

4) Confining Zone IIB₁-IIB₂

Confining zone IIB₁-IIB₂ separates Aquifer zone IIB₂ from Aquifer zone IIB₁. This zone is a leaky confining zone. Laboratory analyses of vertical and horizontal permeability were performed on representative undisturbed samples from this zone. The range in the reported vertical permeability measurements is from a low value of 1.9×10^{-5} feet/day to a high value of 0.12

feet/day (Bledsoe, et al., 1990; Atlanta Testing and Engineering, 1987, 1988). The unit weights of these samples varied from a low of 66.1 pcf (pounds per cubic foot) to a high of 106.6 pcf. The lower unit weights are indicative of a silty and sandy nature.

5) Confining Unit IIA-IIB

Confining Unit IIA-IIB separates Aquifer zone IIB₁ from Aquifer Unit IIA. This unit is more competent than Confining zone IIB₁-IIB₂ but still exhibits leakage. Laboratory analyses of physical characteristics, including horizontal and vertical permeability, were performed on representative undisturbed samples of the clayey portions of Confining Unit IIA-IIB. Nine analyses were performed from wells completed in the GSA. The range in values is from a low of 3.8×10^{-8} feet/day to a high of 2.7×10^{-3} feet/day. The dry unit weights range from 50.9 pcf to 106.3 pcf. In general, the higher unit weights indicate that Confining Unit IIA-IIB is a dense and compact clay. The lower values result from lighter organic material in some samples of the clay.

6) Confining System I-II

Confining System I-II, the principal confining unit in the GSA, separates the basal hydrostratigraphic unit, Aquifer Unit IIA of Aquifer System II, from Aquifer System I (Figures 9 and 12). Confining System I-II was penetrated in 15 borings within the study area. The system attains a thickness of over 100 feet in the GSA. (Figure 19). No streams in the study area incise the system.

Vertical and horizontal hydraulic conductivities of the clay and silt beds are very similar in the GSA. Vertical hydraulic conductivities for Confining System I-II range from 7.7×10^{-5} feet/day to

2.9 x 10⁻⁴ feet/day. Horizontal hydraulic conductivities from 1.0 x 10⁻⁴ feet/day to 1.1 x 10⁻⁴ feet/day (Bledsoe, Aadland, and Sargent, 1990). The consistency of the low conductivity values (vertical and horizontal) support the confining capabilities of this system.

Lithological and geophysical data add additional support to the confining capabilities of this system. Clays and silts are generally very fissile and highly micaceous. The gamma ray curve is very distinct with spikes typical of clays. Resistivity curves support high clay content with a significant drop in resistivity associated with the gamma spikes (Figure 20). The lithological and geophysical characteristics of Confining System I-II are correlative over the entire study area, as illustrated in Cross Sections 1, 2, 3, 4, 5A, and 5B (Plates 1 through 5B).

B. Ground Water Flow in Cross Section

In order to interpret the effect of lithologic variation on ground water flow in the GSA, equipotential lines were plotted on the GSA regional cross sections (Plates 1 through 5B) and are included in this report as Plates 7, 8, 9, 10A, 10B. Water level data are from the fourth quarter 1989. Flow direction on each cross section is described with respect to plant North.

1) Regional Equipotential Cross Section 1

Cross Section 1 (Plate 6) represents the 2.4-mile line of section from north of the MWMF to southeast of the H Area Seepage Basins. The section generally follows the ground water divide noted in the water table surface plan view and Aquifer Zone IIB₁ (Figure 3). The water table surface slopes from 265 to 230 feet above msl along the length of the section. The slope of the equipotential lines indicates a strong tendency for downward movement.

The local change in the slope of the equipotential lines in the vicinity of HSB-85 and BGO-6 is due to the orientation of the line of section; the equipotential lines are more perpendicular to flow at this location. Flow potential is downward with a moderate component of lateral flow. This is in response to the maximum 20 feet of head drop observed along the section in the wells screened just above Confining Unit IIA-IIB. Recharge is indicated all along the line of section with a major influence of the primary recharge exhibited in the area between wells P-27 and HC-12.

Ground water flow in the underlying Aquifer Unit IIA is generally east to west; the total head drop along the section length is approximately 15 feet. The line of section roughly parallels the flow in Aquifer Unit IIA. Wells BGO-8A and BGO-10A were not included in this interpretation. These wells have been identified as having been improperly constructed; they have yielded anomalous water levels since their installation and are currently being replaced.

2) Regional Equipotential Cross Section 2

Cross Section 2 (Plate 7) represents the 3.6-mile line of section from north of the F Area Seepage Basins to east-southeast of the H Area Seepage Basins. This is essentially along the regional dip of the GSA and subparallel to the ground water divide through the MWMF. This section line is generally downgradient from the ground water divide evident in the shallow system. Recharge is indicated as occurring primarily from the east to the west with a strong downward potential all along the length of the section. Lateral flow is illustrated, however, in the shallower portions of Aquifer zone IIB₂ and the equipotential lines "open" somewhat between HSB-117 and DRB-4, possibly in response to some additional recharge. The ground water mound depicted in the thickness of the saturated water table (Figure 29) apparently corresponds to the recharge. Overall,

the Aquifer zone IIB₁ and IIB₂ reveal strong recharge from the vicinity of P-27, with a predominant downward flow potential and an indication of a prevalent lateral tendency for flow further to the west.

Underlying the shallow Aquifer zone IIB₁ and IIB₂ is Aquifer Unit IIA. It reveals lateral flow from east to west with a total drop in head from 175 feet to 155 feet above msl. Toward the west beneath the F Area Seepage Basins the lines of equipotential in Aquifer Unit IIA bend in response to an upward potential for flow observed between the upper and lower portions of the aquifer. This tendency has been observed in the historical water level record in limited areas within the GSA, generally in close proximity to Upper Three Runs Creek. The upward tendency may be attributed to the head or pressure relief in UTRC or to localized permeability differences.

3) Regional Equipotential Cross Section 3

Cross Section 3 (Plate 8) represents the 2.7-mile line of section from northeast of the F Area Seepage Basins to south-southwest of the basins. The section roughly follows the regional strike. Recharge is indicated all along the line of section. Ground water flow is indicated to have a stronger lateral component for Aquifer zone IIB₂ than in the underlying Aquifer zone IIB₁. The subtle mound can be discerned around the 210-foot equipotential line that is likely attributable to some residual recharge from the inactive seepage basins. The equipotential lines become more closely spaced from FSB-91 to the south, as Fourmile Branch (the discharge point for the Water Table Aquifer) is approached. No apparent effects of the offset drawn between the wells FSB-100 and FSB-91 is perceived in the equipotential lines in this orientation and scale.

Beneath Confining Unit IIA-IIB, Aquifer Unit IIA is flowing generally updip in a northwesterly direction towards UTRC. The stronger component of flow in Aquifer Unit IIA illustrated in this

section, is perpendicular to the regional strike of the section. The equipotential line, 155 feet above msl, is contorted in response to the upward potential for flow existing between the upper and lower portions of Aquifer Unit IIA in this portion of the GSA.

4) Regional Equipotential Cross Section 4

Cross Section 4 (Plate 9) represents the 1.3-mile line of section from northeast of the MWMF to the south-southeast of the H Area Seepage Basins. The section roughly follows the regional strike. Similar to the orientation of Cross Section 3, recharge is occurring all along the line of section with potential for ground water flow from the north to south. Between BGO-6 and BGO-35, the line of cross section parallels the equipotential lines and is therefore perpendicular to the direction of ground water flow. In this part of the section a dominant tendency for vertical flow can be discerned. From BGO-35 to the south, the cross section line crosses the equipotential lines (in plan view) and becomes parallel to the ground water flow direction. Here, the equipotential lines indicate a stronger tendency for lateral flow. In the vicinity of HSB-116, HSB-115, and HSB-86, each of the wells screened above and below the Confining zone IIB₁-IIB₂ have essentially identical pressures. No head difference is present and lateral flow tendencies predominate in this area. Moving south towards Fourmile Branch, the shallower water table zone equipotentials are affected by proximity to the creek, causing pressure relief and flow towards the creek. The underlying Aquifer zone IIB₁ appears only slightly affected by the proximity of the creek, and the possibility of some localized underflow is indicated within the floodplain.

Beneath Confining Unit IIA-IIB, water in Aquifer Unit IIA is flowing generally transverse to this line of the section west to northwest towards UTRC. However, the pressure drop is exhibited as lateral flow occurring generally in a north to northwest direction (Figure 31).

5) Regional Equipotential Cross Sections 5A and 5B

Cross Sections 5A and 5B (Plates 10A and 10B) represent the 2.3-mile long east-west line of section from east of the MWMF to the western edge of the F Area Seepage Basins. The discussion of the equipotential lines for these sections will be treated as one single line of section. Recharging conditions are indicated all along the 5A/5B line of cross section. The 5A portion of the sectional line is located just to the north of the ground water divide through the GSA and therefore downgradient of the divide on the north side (Figure 3). Little control is available for the construction of equipotentials further to the east of BGO-5 along Cross Section 5A, but there appears to be a change in the pressure relationship associated with the line of offset depicted near BGO-5. The offset is a single fault essentially parallel to the line of cross section (Figure 16). The bend in the line of Cross Section 5A causes the line of section to intersect the fault twice. Generally the equipotentials in this area indicate recharging conditions. Although the potential for lateral flow is evident, the major component of flow is perpendicular to the line of section, i.e., to the north and northwest.

BGO-14 and BGO-25 represent the midpoint of the line of cross section (Figure 16). The orientation of the equipotentials in this area remains fairly constant indicating recharging conditions with a prevalent vertical potential. As the transition is made from the high ground in the MWMF (along the northeast side of Cross Section 5B) the equipotentials curve more toward the vertical, indicating a stronger component of lateral flow. At the southwest end of Cross Section 5B, the 210-foot line of equipotential bends in response to an area of increased recharge. This recharge is apparently caused by the F Area Seepage Basins. Along the length of Cross Section 5B the drop in head in a westerly to southwesterly direction is slightly more than 15 feet.

In the underlying Aquifer Unit IIA the initial portions of the Cross Section 5A proceeding from east to west are actually found to fall between the contoured piezometric levels of 155 to 160 feet

above msl. Additionally, three of the wells screened in this zone (BGO-8A, BGO-10A, and BGO-12A) produce anomalous water levels and were not incorporated into the dataset. As a result, no equipotential lines are produced for the 5A portion of Cross Section 5 for Aquifer Unit IIA.

Near the matchline between Cross Sections 5A and 5B, the line of section turns to the south. The resulting water levels, in profile, are subparallel to the potentiometric lines, and the flow is therefore perpendicular to the cross sectional line or to the north towards UTRC. Approaching the F Area Seepage Basins to the west, equipotentials indicate lateral flow towards the west and directly beneath the Seepage Basin F-3. Near FSB-99 the equipotential lines strongly converge, indicating lateral flow to the north-northeast towards UTRC.

C. Vertical Head Relationships

The uppermost aquifers beneath the GSA are comprised of three waterbearing units separated by two leaky aquitards. The shallowest waterbearing zone, Aquifer zone IIB₂, overlies the poorly confined Aquifer zone IIB₁, which in turn overlies Aquifer Unit IIA. The pressure relationships in general are downward from Aquifer zone IIB₂ to Aquifer zone IIB₁ down into Aquifer Unit IIA. Maps of the pressure differences were prepared on a scale where 1"=1000 feet for the GSA. These differences were then contoured in order to evaluate the pressure relationships throughout the GSA.

1) Head Differences Across Confining Zone IIB₁-IIB₂

The differences in the elevations of the water table surface and the potentiometric surface of Aquifer zone IIB₁ were plotted spatially and contoured utilizing a 5-foot contour interval (Figure 32). Confining zone IIB₁-IIB₂ supports an average head difference of 5 to 10 feet. The overall

study area contains localized areas of more than 25 feet of head difference to areas of no head difference. The head difference indicates that Aquifer Zones IIB₁ and IIB₂ are not in complete hydraulic communication. However, the zones are considered generally connected and the head difference is downward from Aquifer Zone IIB₂ to Aquifer Zone IIB₁.

The areas contoured with average head differences (5 to 10 feet) attest to the general low permeability of the confining materials and the evidence of the response to recharge to the water table. The map illustrates two large areas of no head difference. These areas are located just east of the MWMF and northwest of the F Area Seepage Basins. These areas are associated with an absence of Confining zone IIB₁-IIB₂ as previously discussed. Additionally, the area with no observed head difference near IDL-3 is also associated with a large roadcut and valley slope down to Upper Three Runs Creek.

No zones of "0" head difference are evident around the F Area Seepage Basins. However, in the vicinity of H Area Seepage Basins, four localized areas of no head difference are observed. Of these four areas, only the one found near HSB-124 and HSB-123 appears to be related to the absence of the underlying confining unit. Just south (downgradient) of the Basin H-4 a small area of high head difference which may be related to the offset in the sediments observed is noted.

In the subsurface database, 46 points were available for the determination of the difference in head across the Confining Zone IIB₁-IIB₂. At these locations the thickness of the confining unit is also known. The head difference divided by the thickness of the unit produces a gradient in feet of head difference per foot of clay, which can be used to further evaluate the head differentials. The arithmetic mean of the gradient calculation is 0.5 feet per foot. The coefficient of variation of these values is high, approaching 120 percent, indicating an extreme range in the values. This occurrence is primarily due to the range of the head differences and the variable thickness of the confining unit.

2) Head Differences Across Confining Zone IIA-IIB

A map of the differences in head observed from the wells in the GSA completed above and below Confining Unit IIA -IIB was prepared in a fashion similar to the head difference map for Confining Zone IIB₁-IIB₂ (Figure 33). The map illustrates a ridge of maximum head difference across the GSA trending in an east-west direction. The largest head difference (66.2 feet) is noted at the northwest boundary of the study area at cluster FC-1. The lowest head difference was 39.8 feet occurring at cluster FSB-79. The lines of equal head difference, as contoured in the vicinity of the H Basins, indicate an increase beneath the basins tending in a general north-south direction. This area corresponds to the largest calcareous body situated in the lower calcareous zone. The response may be related to the localized high permeability suspected for these materials based on the drilling notes and slug tests values.

The differences in head at each cluster location were divided by the associated thickness of the Confining Unit IIA-IIB. The resulting value is a gradient calculated in feet per foot of the loss in pressure across the clay materials comprising the confining unit. This value is a reasonable estimate of the confining capability of the unit expressed in a relative sense throughout the GSA. The subsurface database contains 20 values available for analysis of the relative head loss across this clay boundary. The mean of the data is 9.4 feet per foot with a coefficient of variation of 71 percent, indicating that the data are very consistent throughout the GSA.

3) Head Relationships Between the Cretaceous and Tertiary Aquifers

The contact of the Confining System I-II with the underlying unit (Aquifer System I) is marked by an erosional unconformity representing Cretaceous-age sediments overlain with Tertiary-age sediments. In the GSA the ground water in the Cretaceous sediments is under higher pressure than

in Aquifer Unit IIA that overlies Confining System I-II. This was first discovered from preliminary subsurface reconnaissance work undertaken by the Corps of Engineers (1952).

The upward pressure gradient across the Tertiary/Cretaceous boundary has been recognized in the GSA since the 1960s with the maps prepared by Reichart (1969). Later, Marine (1983) plotted water elevations of the aquifers above and below Confining System I-II and concluded that the underlying upper Cretaceous Aquifer (Aquifer Unit IB) in this report (Figure 15) has a potentiometric surface some 10 feet above the potentiometric surface of the Tertiary-age Aquifer Unit IIA within some portions of the SRS.

Since 1981 ground water development on the SRS has increased as many facilities converted from surface water to ground water and additional facilities have been constructed. Ground water usage has almost doubled (Stephenson and Steele, 1988). These demands, coupled with increased offtake from downdip municipal pumpage, has produced a measurable effect of the head relationships between Aquifer System I and Aquifer Unit IIA on site. Although the upward pressure gradient has been observed to decline and the unity line between the aquifers has wandered somewhat with time, the upward pressure gradient has been found to be persistent beneath the F and H Area Seepage Basins.

Within the GSA the upward pressure gradient ranges from 0 to 20+ feet. An updated version of the pressure gradient map between Aquifer System I and Aquifer System II has been prepared from fourth quarter 1989 data and is presented as Figure 34 (Bledsoe, personal communication, 1990). Historical data indicate that the upward pressure gradient has remained essentially constant since 1988.

Figure 34. Head Difference Between Aquifer Unit IB of Aquifer System I and Aquifer II of Aquifer System II

REFERENCES

Aadland, R. K., and H. W. Bledsoe, "Classification of Hydrostratigraphic Units at Savannah River Site, South Carolina." Savannah River Laboratory, WSRC-RP-90-987, Westinghouse Savannah River Company, Aiken, S.C. (1990).

Albenesius, P. W., M. B. Amidon, and T. F. Gaughan, "HSB 84A Pumping Test." Waste Management Technology, WSRC-TR-90-313, Westinghouse Savannah River Company, Aiken, S. C. (1990).

Baars, D. L., 1963, Petrology of Carbonate Rocks: in. Bass, R. O. ed., Shelf Carbonates of the Paradox Basin; A Symposium: Fourth Field Conference: Four Corners Geological Society, pp. 101-129.

Beard, D. C., and P. K. Weyl. "Influence of Texture on Porosity and Permeability of Unconsolidated Sand." American Association of Petroleum Geologists; Vol. 57 No. 2. (1973).

Bledsoe, H. W., R. K. Aadland, and K. S. Sargent, "SRS Baseline Hydrogeologic Investigations Summary Report." Savannah River Site, Savannah River Laboratory, WSRC-RP-90-1010-Westinghouse Savannah River Company, Aiken, S. C. (1990).

Bledsoe, H. W., Personal Communication, (1990).

CH₂M Hill, "Congaree Aquifer Test." Savannah River Site, (1989).

Colquhoun, D. J., I. D. Woollen, D. S. Van Nieuwenhuise, G. G. Padgett, R. W. Oldham, D. C. Boylan, J. W. Bishop, and P. D. Howell. "Surface and Subsurface Stratigraphy, Structure and

Aquifers of the South Carolina Coastal Plain." Report for Department of Health and Environmental Control, Ground Water Protection Division, published through the Office of the Governor, State of South Carolina, Columbia (1983).

Christensen, E. J. and D. E. Gordon, "Technical Summary of Groundwater Quality Protection Program at Savannah River Plant." Vol. I, Site Geohydrology and Solid and Hazardous Wastes; DPST-83-929. E. I. du Pont de Nemours & Company, Savannah River Laboratory, Aiken, S. C. (1983).

Colquhoun, D. J., and H. S. Johnson, Jr. "Tertiary Sea-Level Fluctuation in South Carolina." *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 5, pp. 105-126 (1968).

Colquhoun, D. J., and K. B. Steele, "Chronostratigraphy and hydrostratigraphy of the northwestern South Carolina Coastal Plain": Project No: G868-05, Annual Cooperative Grant Agreement Nol 13040 R-83-591, Interim Technical Report to Water Resources Research Institute, Clemson University, Clemson, South Carolina, 15 p (1985).

Cooke, C. W. "Geology of the Coastal Plain of South Carolina." U. S. Geological Survey Bulletin 867 (1936).

Cooke, C. W., and F. S. MacNeil, "Tertiary Stratigraphy of South Carolina": U.S. Geological Survey Professional Paper 243-B, p. 19-29 (1952).

Cramer, H. R., and D. D. Arden, Jr. "Faults in Oligocene Rocks of the Georgia Coastal Plain." *Geological Society of America Abstracts with Programs, Southeast Section*, v. 10, p. 166 (1978).

Dennehy, K. F., D. C. Prowell, and P. B. McMahon "Reconnaissance Hydrogeologic Investigation of the Defense Waste Processing Facility and Vicinity." Savannah River Plant South Carolina, U. S. Geological Survey, Water Resources Investigations Report 88-4221, 74 pp (1989).

Doering, J. A. "Quaternary Surface Formations of Southern Part of Atlantic Coastal Plain." Journal of Geology, v. 68, pp. 182-202 (1960).

Faye, R. E., and D. C. Prowell. "Effects of Late Cretaceous and Cenozoic Faulting on the Geology and Hydrology of the Coastal Plain Near the Savannah River, Georgia, and South Carolina." U. S. Geological Survey Open File Report 82-156 (1982).

Geotrans, Inc. "Characterization of Groundwater Flow and Transport in General Separations Area. Savannah River Plant, Effluents of Closure Options on Groundwater Flow at H-Area Seepage Basins (1989).

Geotrans, Inc. "Characterization of Ground Water Flow and Transport in the General Separations Area, Savannah River Plant, Effects of Closure Options on Groundwater Flow at the F-Area Seepage Basins" (1988).

Gohn, F. S., J. E. Hazel, L. M. Bybell, and L. E. Edwards. "The Fishburne Formation (Lower Eocene), a Newly Defined Subsurface Unit in the South Carolina Coastal Plain." U. S. Geological Survey Bulletin 1537-C, pp. 1-16 (1983).

Hazel, J. E., L. M. Bybell, R. A. Christopher, N. O. Frederiksen, F. E. May, D. M. McLean, R. Z. Poore, C. C. Smith, N. F. Sohl, P. C. Valentine, and R. J. Witmer. "Biostratigraphy of the Deep Corehole (Clubhouse Crossroads Corehole 1) near Charleston, South Carolina," Studies

Related to the Charleston, South Carolina, Earthquake of 1886 -- A Preliminary report, ed., D. W. Rankin, U. S. Geological Survey Professional Paper 1028, pp 71-89 (1977).

Higgins, B. B., G. S. Gohn, and L. M. Bybell. "Subsurface Geologic Evidence for Normal Faults in the South Carolina Coastal Plain Near Charleston." Geological Society of America Abstracts with Programs, Southeast Section, v. 10, p. 171 (1978).

Howell, D. E., and A. W. Zupan. "Evidence for Post-Cretaceous Tectonic Activity in the Westfield Creek Area, North of Cheraw, South Carolina." South Carolina State Development Board, Division of Geology, Geologic Notes, v. 18, pp. 98-105 (1974).

Huddlestun, P. F., and J. H. Hetrick. "Upper Eocene Stratigraphy of Central and Eastern Georgia." Georgia Geologic Survey Bulletin 95, p. 78 (1985).

Inden, R. F., and A. W. Zupan. "Normal Faulting of Upper Coastal Plain Sediments, Ideal Kaolin Mine, Langley, South Carolina." South Carolina State Development Board, Division of Geology, Geologic Notes, v. 19, pp. 159-165 (1975).

Johnson, H. S., Jr., and S. D. Heron, Jr. "Slump Features in the McBean Formation and Younger Beds, Riley Cut, Calhoun County, South Carolina." South Carolina State Development Board, Division of Geology, Geologic Notes, v. 9, pp. 37-44 (1965).

Klitgord, K. D., and J. C. Behrendt. "Basin Structure of the U. S. Atlantic Margin." Geological and Geophysical Investigations of Continental Margins. American Association of Petroleum Geologists Memoir 29, eds., J. S. Watkins, L. Montadert, and P. W. Dickerson, pp. 85-112 (1979).

Marine, I. W., and G. E. Siple. "Buried Triassic Basin in the Central Savannah River Area, South Carolina and Georgia." *Geological Society of America Bulletin*, v. 85, pp. 311-320 (1974).

Marine, I. W., "Hydrology of Buried Crystalline Roch at Savannah River Plant Near Aiken, South Carolina" DOE/SR-WM-79-2. 220 pp., 9(1979).

Marine, I. W., "Geohydrology of a Buried Triassic Basin at Savannah River Plant, South Carolina." *American Association of Petroleum Geologists Bulletin*, v. 58, pp. 1825-1837 (1974).

McCartan, Lucy, E. M. Lemon, Jr., and R. E. Weems, "Geologic Map of the Area Between Charleston and Orangeburg, South Carolina": U.S. Geological Survey Miscellaneous Investigations Series Map I-1472, scale 1:250,000.

McDowell, R. C. and B. B. Houser. "Distribution of Small - Scale Deformation Structures Upper Coastal Plain, South Carolina and Georgia." U. S. Geological Survey Map MF-1538 (1983).

Newell, W. L., M. H. Pavich, D. C. Prowell, and H. W. Markewich. "Surficial Deposits, Weathering Processes, and Evolution of an Inner Coastal Plain Landscape, Augusta, Georgia." *Excursions in Southeastern Geology; Guidebook*, ed., R. W. Frey, Geological Society of America 1980 Annual Meeting, Atlanta, Georgia, American Geological Institute, Guidebook, v. 2, pp. 527-544 (1980).

Nystrom, P. G., Jr., R. H. Willoughby, and L. E. Kite. "Cretaceous-Tertiary Stratigraphy of the Upper Edge of the Coastal Plain Between North Augusta and Lexington, South Carolina." *Carolina Geological Society Field Trip Guidebook*, South Carolina Geological Survey (1986).

Nystrom, P. G., Jr., and R. H. Willoughby. "Geological Investigations Related to the Stratigraphy in the Kaolin Mining District, Aiken County, South Carolina." Carolina Geological Society Field Trip Guidebook, South Carolina Geological Survey, p. 183 (1982).

Parizek, R. R., and R. W. Root, "Development of a Ground Water Velocity Model for the Radioactive Waste Management Facility, Savannah River Plant, Aiken, South Carolina." DPST-86-658, E. I. du Pont de Nemours & Company, Savannah River Laboratory, Aiken, South Carolina (1986).

Prowell, D. C., R. A. Christopher, L. E. Edwards, L. M. Bybell, and H. E. Gill. "Geologic Section of the Undip Coastal Plain from Central Georgia to Western South Carolina." U. S. Geological Survey, Text with Map MF-1737, p. 11 (1985a).

Prowell, D. C., Edwards, L. E., and Frederiksen, N. O.; "The Ellenton Formation in South Carolina—A revised age designation from Cretaceous to Paleocene: U. S. Geological Survey Bulletin 1605-A. pp. A63-A69, (1985b).

Reichart, S.O., "Flow of Groundwater in Tertiary Sediments at the Savannah River Plant, Near Aiken, South Carolina." Draft Report, Savannah River Plant, Aiken, S. C. (1969).

S&ME Inc. "Borehole Permeability Analysis F-Area Assessment Wells at the Savannah River Plant, Aiken, SC." (1988B).

S&ME Inc. "Borehole Permeability Analysis for Burial Ground Parameter Wells at the Savannah River Plant, Aiken, SC." (1988A).

Siple, G. E. "Geology and Ground Water of the Savannah River Plant and Vicinity, South Carolina." U. S. Geological Survey Water-Supply Paper 1841 (1967).

Sirrine Environmental Consultants. "Field Permeability Tests H-Area Seepage Basins Savannah River Plant." (1988A).

Sirrine Environmental Consultants. "Field Permeability Tests H-Area Seepage Basin Plume Assessment Well Savannah River Plant." (1988B).

Sirrine Environmental Consultants. "F-Area Compliance Monitoring Program Field Permeability Tests." (1987).

Stephenson, D. E., and J. E. Steele, "New Production Wells and Head Reversal in the Separations Areas of the Savannah River Plant (SRP)." Savannah River Laboratory, DPST-88-473, E. I. du Pont de Nemours & Company, Aiken, S. C. (1988).

Stieve, A. L., Stephenson, D. E. and R. K. Aadland; "Pen Branch Fault Program: Consolidated Report on the Seismic Reflection Surveys and the Shallow Drilling (U)"; U.S. DOE Report WSRC-TR-91-87; Westinghouse Savannah River Co, Aiken S.C. 26 pp. (1991).

Strom, R. N., 1991, Personal Communication.

Thayer et al, 1990, Personal Communication

U. S. Army Corps of Engineers, Charleston District. Geologic-Engineering Investigations, Savannah River Plant. DPWR-52, E. I. du Pont de Nemours and Company, Savannah River Plant, Aiken, SC (1952).

Veatch, Otto, and L. W. Stephenson, "Preliminary Report on the Geology of the Coastal Plain of Georgia": Georgia Geologic Survey Bulletin 23, pp. 237-296 (1911).

APPENDIX I

MASTER LIST OF WELLS
GENERAL SEPARATIONS AREA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note: * = Elevation below sea level; NA = Not available)
 ^Source is unedited HP data
 Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
^ BG 52	75910.4	55524.3	NA	289.8	243.8-223.8	NA	7/06/76	NA	NA
^ BG 53	76157.3	55073.9	NA	285.7	234.7-214.7	NA	7/07/76	NA	NA
^ BG 54	75837.9	54830.3	NA	277.2	235.2-215.2	NA	7/07/76	NA	NA
^ BG 55	75525.3	54590.5	NA	276.9	234.9-214.9	NA	7/08/76	NA	NA
^ BG 56	75206.5	54481.9	NA	274.9	230.9-210.9	NA	7/20/76	NA	NA
^ BG 57	75000.4	54820.0	NA	272.6	234.6-214.6	NA	7/19/76	NA	NA
^ BG 58	74790.9	55162.3	NA	278.2	238.2-218.2	NA	7/16/76	NA	NA
^ BG 59	74593.4	55508.3	NA	282.7	237.7-217.7	NA	7/09/76	NA	NA
^ BG 60	74386.3	55850.3	NA	275.5	235.5-215.5	NA	7/12/76	NA	NA
^ BG 61	74075.4	56360.8	NA	275.0	245.0-225.0	NA	7/12/76	NA	NA
^ BG 62	73971.6	56530.9	NA	272.5	242.5-222.5	NA	7/12/76	NA	NA
^ BG 63	73754.5	56870.5	NA	274.2	244.2-224.2	NA	7/13/76	NA	NA
^ BG 64	73547.0	57212.4	NA	285.3	247.3-227.3	NA	7/13/76	NA	NA
^ BG 65	73340.6	57552.7	NA	290.9	250.9-230.9	NA	7/14/76	NA	NA
^ BG 66	73585.0	57805.0	NA	296.0	251.0-231.0	NA	7/14/76	NA	NA
^ BG 67	73954.1	57902.6	NA	294.7	244.7-224.7	NA	7/15/76	NA	NA
EGO 1D	73737.9	58779.3	293.0	295.1	245.0-225.0	267.0-216.0	9/10/87	(30-72)	(0-71)
EGO 2D	74552.9	58809.7	294.9	296.9	238.9-218.9	248.9-209.9	8/12/87	NA	NA
EGO 3D	75351.3	58809.2	290.8	292.7	247.6-227.6	259.8-215.8	9/10/87	(33-68)	0-68)
EGO 4D	76150.1	58803.7	295.6	297.5	240.6-220.6	252.2-213.1	8/13/87	NA	NA
EGO 5C	76476.9	58794.5	294.2	296.1	193.2-183.2	198.2-166.2	9/04/87	(33-127)	(0-128)
EGO 5D	76477.5	58784.8	294.2	296.3	241.8-221.8	248.9-214.2	9/04/87	NA	NA
EGO 6A	76487.2	58316.8	283.8	285.6	117.5-107.5	135.2-97.8	10/08/87	(0-181)	(0-179)
EGO 6C	76487.1	58307.0	283.5	285.6	168.0-158.0	179.5-149.5	9/02/87	NA	NA
EGO 6D	76487.3	58297.1	283.2	285.5	237.2-217.2	248.2-208.2	8/31/87	NA	NA
EGO 7D	76494.5	57917.2	280.5	282.4	235.5-215.5	249.5-207.5	8/15/87	NA	NA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

^Source is unedited HP data

Update6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
BGO 8A	76569.0	57618.3	281.3	283.2	115.3-105.3	124.8-91.3	10/07/87	(0-181)	(0-182)
BGO 8C	76596.0	57619.0	281.3	283.1	179.8-169.8	188.3-161.3	8/28/87	NA	NA
BGO 8D	76588.0	57617.0	281.1	283.2	236.1-216.1	248.1-206.1	8/28/87	NA	NA
BGO 9D	76811.6	57478.9	283.2	285.1	229.2-209.2	239.7-199.2	8/20/87	NA	NA
BGO 10A	76805.2	57050.9	299.1	300.9	121.1-111.1	132.1-101.1	10/05/87	(0-192)	(0-188)
BGO 10C	76805.0	57041.0	299.3	300.0	167.3-157.3	181.3-144.3	9/15/87	NA	NA
BGO 10D	76805.0	57030.0	299.5	300.0	250.5-230.5	262.5-219.5	9/16/87	NA	NA
BGO 11D	76805.1	56651.3	303.3	305.3	236.3-216.3	249.8-203.3	8/24/87	NA	NA
BGO 12A	76804.6	56250.7	311.4	313.4	116.4-106.4	136.4-93.4	10/02/87	(0-211)	(0-211)
BGO 12C	76805.2	56241.1	311.6	313.6	163.6-153.6	179.6-146.6	10/01/87	NA	NA
BGO 12D	76805.2	56231.1	311.8	313.7	237.8-217.8	252.8-210.8	9/29/87	NA	NA
BGO 13D	76805.3	55840.0	316.5	318.5	248.5-228.5	268.7-218.5	10/12/87	NA	NA
BGO 14A	76377.5	55838.3	300.2	301.9	119.6-109.6	132.6-100.2	10/19/87	(0-196)	(0-196)
BGO 14C	76367.7	55839.0	300.1	302.0	202.1-192.1	214.4-185.1	10/16/87	NA	NA
BGO 14D	76357.5	55839.6	300.0	302.0	249.6-229.6	266.3-220.0	10/15/87	NA	NA
BGO 15D	75973.5	55859.1	296.7	298.7	238.7-218.7	249.4-196.7	8/25/87	NA	NA
BGO 16A	75757.0	56194.2	302.8	305.0	112.5-102.5	123.3-92.8	12/08/87	(2-211)	(0-175)
BGO 16D	75751.4	56202.1	302.3	304.6	237.3-217.3	245.9-209.3	8/24/87	NA	NA
BGO 17D	75599.6	56399.4	296.0	298.3	224.0-204.0	233.5-198.0	7/08/87	NA	NA
BGO 18A	75599.9	56699.7	292.9	295.2	109.5-99.5	121.3-89.0	10/30/87	(0-201)	(0-166)
BGO 18D	75600.0	56711.2	292.6	294.9	239.6-219.6	250.1-201.0	11/04/87	NA	NA
BGO 19D	75350.0	56997.3	284.9	287.2	213.9-193.9	223.4-186.9	7/15/87	NA	NA
BGO 20D	74962.2	57113.8	281.3	283.7	236.3-216.3	248.3-206.8	3/02/88	NA	NA
BGO 21D	74688.5	57470.7	283.0	285.4	237.7-217.7	247.2-211.0	11/11/87	(16-72)	NA
BGO 22D	74482.2	57817.3	284.2	286.5	214.2-194.2	220.7-180.2	7/23/87	NA	NA
BGO 23D	74238.1	58133.0	287.0	289.2	242.0-222.0	249.2-210.6	8/06/87	(0-76)	(0-73)

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note: * = Elevation below sea level; NA = Not available)

~Source is unedited HP data

Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
BGO 24D	74012.4	58438.8	291.0	293.2	241.0-221.0	248.2-192.0	7/29/87	NA	NA
BGO 25A	76158.5	55668.1	294.7	296.5	114.1-104.1	136.7-94.7	10/19/87	(31-187)	(0-187)
BGO 26A	76144.6	55014.2	285.1	287.2	91.0-81.0	109.3-63.1	3/08/89	(11-210)	(0-212)
BGO 26D	76128.0	55015.2	283.5	285.5	233.5-213.5	256.4-203.5	12/01/88	NA	NA
BGO 27C	75666.3	54671.4	273.9	276.0	164.9-154.9	177.4-141.9	3/14/89	(14-130)	(0-127)
BGO 27D	75677.3	54680.2	274.3	276.3	229.3-209.3	250.5-199.3	12/02/88	NA	NA
BGO 28D	75348.3	54457.9	275.1	277.4	230.1-210.1	248.5-200.1	12/16/88	NA	NA
BGO 29A	75560.0	54103.5	262.1	264.2	112.5-102.5	125.8-83.2	2/05/89	(11-195)	(0-196)
BGO 30C	75181.0	54512.3	272.6	274.5	177.9-167.9	197.1-142.1	4/01/89	NA	(0-117)
BGO 30D	75187.7	54499.2	272.8	274.8	227.8-207.8	249.8-197.8	2/04/89	NA	NA
BGO 31C	74978.0	54816.2	271.1	273.1	186.4-176.4	197.5-168.1	4/03/89	(11-115)	(0-115)
BGO 31D	74985.3	54841.7	271.6	273.7	231.1-211.1	254.0-201.6	2/11/89	NA	NA
BGO 32D	74727.0	55250.2	279.5	281.7	234.5-214.5	255.7-204.5	12/15/88	NA	NA
BGO 33C	74479.7	55681.4	277.4	279.4	187.8-177.8	198.3-157.4	2/05/88	(12-120)	(0-118)
BGO 33D	74468.7	55695.4	278.1	280.3	233.1-213.1	249.3-203.1	12/05/88	NA	NA
BGO 34D	74228.8	56082.6	272.7	274.9	232.7-212.7	245.5-202.7	12/07/88	NA	NA
BGO 35C	73953.9	56545.7	271.4	273.4	171.9-161.9	186.5-151.4	3/21/89	(11-120)	(0-118)
BGO 35D	73946.0	56556.5	271.4	273.5	239.4-219.4	252.6-209.4	12/07/88	NA	NA
BGO 36D	73743.8	56888.1	273.3	275.4	243.3-223.3	259.9-213.3	12/08/88	NA	NA
BGO 37C	73498.2	57279.2	284.3	286.3	178.8-168.8	203.1-161.3	3/22/89	(11-115)	(0-115)
BGO 37D	73490.8	57292.9	285.1	287.3	246.1-226.1	263.8-220.1	12/09/88	NA	NA
BGO 38D	73329.3	57557.5	289.3	291.6	242.3-222.3	259.3-212.3	12/13/88	NA	NA
BGO 39D	73583.5	57831.0	293.7	295.7	244.7-224.7	262.2-214.7	12/14/88	NA	NA
DRB 2	72600.0	56900.0	281.6	284.6	NA	NA	NA	NA	(0-2000)
DRB 3	75500.0	56900.0	285.3	285.2	75.8-65.8	NA	3/13/70	NA	(0-1900)
DRB 4	73900.0	53400.0	250.0	252.4	107.1-99.1	NA	1/20/76	NA	(0-1950)

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

Source is unedited HP data

Updated 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
DRB 6	73958.0	60005.0	269.1	272.1	NA	NA	NA	NA	(0-1900)
DRB 7	72800.0	60000.0	277.8	280.8	NA	NA	NA	NA	(0-2000)
F 1	75604.7	51302.8	281.4	281.9	256.9-251.9	NA	3/06/56	NA	NA
F 2	75677.4	51484.3	280.2	281.4	217.0-207.0	219.2-207.0	1/26/56	NA	NA
F 9	75015.1	50651.3	270.0	269.3	213.1-203.1	NA	1/09/56	NA	NA
F 10	75155.3	50444.3	277.0	280.1	276.5-266.5	276.0-266.0	3/05/56	NA	NA
F 14	74807.1	50010.9	NA	275.7	NA	NA	3/01/62	NA	NA
F 15	74716.0	50738.3	NA	253.5	NA	NA	3/01/62	NA	NA
F 16	74597.1	50029.0	NA	255.9	NA	NA	3/01/62	NA	NA
F 17	74303.4	50466.4	NA	239.8	NA	NA	3/01/62	NA	NA
F 18A	74170.2	50108.0	231.4	233.4	204.4-194.4	212.4-194.4	1/16/85	NA	NA
F 19	74245.9	49672.9	NA	243.9	NA	NA	3/01/62	NA	NA
F 24	76069.0	51322.9	NA	291.0	NA	NA	3/01/62	NA	NA
F 25	75083.8	51322.1	NA	260.9	NA	NA	3/01/62	NA	NA
FC 1A	79664.5	53114.7	316.7	318.6	103.6-98.6	NA	7/23/76	(2-219)	(0-213)
FCB 2	76679.7	55046.7	305.2	307.3	235.2-205.2	NA	9/03/81	NA	NA
FCB 3	76427.8	54874.4	300.3	302.3	225.3-195.3	NA	10/15/81	NA	NA
FCB 4	76780.4	54605.9	304.5	306.6	234.5-204.5	NA	11/04/81	NA	NA
FCB 5	76492.6	54773.0	301.7	303.9	237.1-217.1	246.9-204.7	7/15/88	NA	NA
FCB 6	76582.1	54733.4	308.4	310.5	235.1-215.1	243.3-205.4	7/13/88	NA	NA
FCB 7	76913.9	54957.1	313.1	315.4	238.3-218.3	246.0-203.1	7/07/88	NA	NA
FET 1D	76165.6	53299.9	268.0	270.0	226.9-206.9	234.0-198.0	9/26/88	NA	NA
FET 2D	76045.8	52981.2	267.9	270.0	229.5-209.5	236.3-202.9	8/05/88	NA	NA
FET 3D	75961.0	53025.7	283.2	285.2	223.0-203.0	230.8-195.2	8/04/88	NA	NA
FET 4D	75959.3	53149.0	284.7	286.9	225.1-205.1	230.4-193.7	8/05/88	NA	NA
FSB 76	76141.6	51388.8	291.7	294.2	226.7-196.7	230.7-196.7	10/01/81	NA	NA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

^Source is unedited HP data

Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
FSB 76A	76131.9	51391.6	291.5	293.9	48.9-38.5	54.6-33.6	10/24/84	(2-250)	(0-242)
FSB 76B	76122.4	51394.0	291.4	293.8	111.4-100.9	121.1-96.1	10/31/84	NA	NA
FSB 76C	76112.4	51396.4	291.3	293.6	166.2-155.7	175.4-150.2	11/05/84	NA	NA
FSB 77	75129.4	50713.1	271.0	273.3	216.0-186.0	220.0-186.0	10/07/81	NA	NA
FSB 78	74764.0	50164.7	270.1	272.6	215.1-185.1	220.1-185.1	10/08/81	NA	NA
FSB 78A	74757.7	50172.8	270.5	272.6	38.7-28.3	48.7-19.7	11/20/84	(0-248)	(0-250)
FSB 78B	74765.9	50178.8	270.8	272.8	93.7-83.3	102.7-76.2	11/28/84	NA	NA
FSB 78C	74772.5	50170.2	271.4	273.5	152.0-142.2	155.0-135.0	11/30/84	NA	NA
FSB 79	73663.1	50139.7	215.3	217.8	203.3-173.3	208.3-173.3	10/09/81	NA	NA
FSB 79A	73664.5	50149.6	216.1	218.1	35.9-25.4	42.5-20.5	11/13/84	(1-202)	(0-202)
FSB 79B	73666.1	50159.2	216.2	218.2	92.4-81.9	98.9-76.4	11/15/84	NA	NA
FSB 79C	73668.0	50171.3	216.1	218.4	161.2-151.4	171.2-145.2	2/27/84	NA	NA
FSB 87A	75601.7	50115.8	285.6	287.8	45.0-34.5	62.0-27.0	10/22/84	(2-261)	(0-261)
FSB 87B	75597.0	50104.9	285.5	287.5	101.7-91.2	112.7-83.0	11/01/84	NA	NA
FSB 87C	75591.9	50093.4	285.4	287.5	160.7-150.2	170.7-143.2	11/07/84	NA	NA
FSB 87D	75586.3	50081.1	285.3	287.3	218.2-188.8	226.7-182.7	11/15/84	NA	NA
FSB 88C	75619.4	51518.0	280.4	283.0	168.4-158.4	175.9-145.4	10/07/87	NA	(0-134)
FSB 88D	75621.8	51527.0	280.1	282.4	221.1-202.1	230.3-192.1	10/07/87	NA	NA
FSB 89C	75553.2	51345.2	279.1	281.3	166.1-156.1	178.6-145.9	10/07/87	(14-127)	(0-123)
FSB 89D	75548.3	51335.8	278.9	281.2	221.9-201.9	231.3-191.9	10/07/87	NA	NA
FSB 90C	75382.9	51148.6	276.1	278.4	168.1-158.1	181.3-145.1	10/07/87	NA	(0-128)
FSB 90D	75376.9	51140.7	276.1	278.6	225.1-205.1	234.6-196.1	10/07/87	NA	NA
FSB 91C	75213.3	50953.5	277.0	279.3	159.1-149.1	175.9-141.0	10/07/87	(14-131)	(0-134)
FSB 91D	75207.6	50946.6	276.9	279.2	220.9-200.9	230.8-189.9	10/07/87	NA	NA
FSB 92C	75053.2	50564.0	273.6	275.7	157.6-147.6	164.6-131.6	10/07/87	NA	(0-114)
FSB 92D	75045.8	50557.6	273.7	275.9	221.7-201.7	230.5-191.7	10/07/87	NA	NA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)
 ~Source is unedited HP data
 Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
FSB 93C	74897.3	50458.3	274.0	276.2	152.0-142.0	162.2-133.5	10/07/87	(33-135)	(0-136)
FSB 93D	74888.5	50452.4	273.9	276.1	217.9-197.9	230.5-187.9	10/07/87	NA	NA
FSB 94C	74869.0	50180.0	278.8	281.1	149.8-139.8	159.2-128.8	10/07/87	NA	(0-144)
FSB 94D	74863.4	50188.6	278.8	281.1	221.8-201.8	231.1-188.8	10/07/87	NA	NA
FSB 95C	74971.7	50016.7	281.8	284.0	155.8-145.8	164.6-137.8	10/07/87	(31-138)	(0-135)
FSB 95D	74977.5	50008.9	281.8	284.3	227.8-207.8	236.5-196.8	10/07/87	NA	NA
FSB 96A	74882.2	49778.7	277.7	279.8	95.7-85.7	103.4-77.7	10/07/87	(16-199)	(0-196)
FSB 97A	75171.2	49965.7	283.8	286.1	95.8-85.8	106.1-77.8	10/07/87	(15-198)	(3-196)
FSB 97C	75179.6	49970.6	283.8	286.1	153.8-143.8	159.8-133.8	10/07/87	NA	NA
FSB 97D	75188.9	49975.5	283.9	286.0	216.9-196.9	227.4-186.9	10/07/87	NA	NA
FSB 98A	75389.8	50121.6	280.7	283.0	94.7-84.7	103.7-76.7	10/07/87	(15-195)	(2-170)
FSB 98C	75381.2	50116.5	280.8	283.1	156.8-146.8	170.8-116.8	10/07/87	NA	NA
FSB 98D	75371.9	50111.6	280.8	283.1	218.8-198.8	229.8-188.8	10/07/87	NA	NA
FSB 99A	75675.6	50314.8	285.3	287.6	102.5-92.5	111.3-85.3	10/07/87	(0-194)	(3-196)
FSB 99C	75683.7	50320.6	285.2	287.7	167.2-157.2	175.6-147.2	10/07/87	NA	NA
FSB 99D	75691.7	50326.9	285.1	287.6	218.1-198.1	230.0-188.1	10/07/87	NA	NA
FSB 100A	75534.4	50958.4	283.8	286.0	105.8-95.8	113.8-85.8	10/07/87	(0-193)	(0-199)
FSB 101A	75719.0	51191.3	282.9	285.2	102.9-92.9	114.0-82.9	10/07/87	(15-189)	(4-164)
FSB 102C	73582.9	50834.8	198.9	201.1	155.9-145.9	168.1-128.9	2/01/88	NA	(0-70)
FSB 103C	74210.0	49651.3	240.1	242.4	157.1-147.1	172.5-135.1	2/01/88	NA	(0-103)
FSB 104C	73872.7	49248.6	216.7	219.1	160.7-150.7	174.2-136.7	1/20/88	NA	(0-80)
FSB 104D	73865.1	49255.4	216.9	219.2	210.4-190.4	215.4-181.9	1/20/88	NA	NA
FSB 105C	75234.7	49828.0	283.5	285.8	151.5-141.5	168.2-136.0	1/18/88	NA	(0-167)
FSB 105D	75244.3	49833.3	283.7	285.8	223.7-203.7	241.1-193.7	1/18/88	NA	NA
FSB 106C	74190.1	50651.3	233.0	235.1	166.0-156.0	177.0-143.0	2/05/88	NA	(6-83)
FSB 107C	75184.0	51158.1	268.8	270.9	160.8-150.8	174.4-133.8	1/15/88	NA	(8-130)

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

^Source is unedited HP data

Updated 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
FSB 107D	75177.2	51149.8	268.7	271.0	220.9-200.9	230.9-192.7	1/23/88	NA	NA
FSB 108D	76260.7	51142.3	295.8	298.0	223.8-203.8	230.8-193.8	10/07/87	NA	NA
FSB 109D	75855.9	50488.6	290.8	293.1	225.8-205.8	238.9-191.8	10/07/87	NA	NA
FSB 110C	74190.7	50150.6	232.2	234.0	154.2-144.2	153.2-125.7	9/09/87	NA	NA
FSB 110D	74193.3	50141.6	232.6	234.5	211.1-191.1	224.6-182.6	1/26/88	NA	NA
FSB 111C	75383.3	51526.3	274.0	276.3	169.0-159.0	181.2-149.0	2/05/88	NA	(0-133)
FSB 111D	75382.4	51515.9	274.4	276.6	221.7-201.7	236.4-194.4	2/05/88	NA	NA
FSB TA	75649.1	51658.3	275.4	277.3	*235.4-*199.1	*282.0-171.8	11/12/88	(0-558)	(0-555)
~ FTF 1	77413.3	53179.8	281.2	283.1	241.2-221.2	NA	6/02/72	NA	NA
~ FTF 2	77336.0	53275.1	279.4	281.3	239.4-219.4	NA	6/15/72	NA	NA
~ FTF 3	77235.3	53244.8	278.2	280.0	221.2-218.2	NA	9/16/72	NA	NA
~ FTF 4	77132.9	53268.2	276.6	278.5	236.6-216.6	NA	6/08/72	NA	NA
~ FTF 5	77035.6	53168.3	275.3	277.3	235.3-215.3	NA	6/10/72	NA	NA
~ FTF 6	77151.4	53062.0	276.9	278.0	236.9-216.9	NA	6/09/72	NA	NA
~ FTF 7	77235.9	53089.7	278.1	280.0	226.1-222.1	NA	9/08/73	NA	NA
~ FTF 8	77336.2	53059.9	279.6	281.5	239.6-219.6	NA	6/13/72	NA	NA
~ FTF 9	77482.8	52769.5	270.4	271.9	236.4-216.4	NA	6/18/73	NA	NA
~ FTF 10	77336.0	52905.0	268.6	271.1	235.1-215.1	NA	7/03/72	NA	NA
~ FTF 11	77180.7	52748.8	269.8	271.2	235.8-215.8	NA	7/05/72	NA	NA
~ FTF 12	77321.4	52648.5	269.0	271.7	235.0-215.0	NA	11/13/73	NA	NA
~ FTF 13	76637.8	53098.4	284.1	285.8	236.1-216.1	NA	11/17/73	NA	NA
~ FTF 14	76188.8	52107.7	268.6	270.7	238.6-218.6	NA	9/10/73	NA	NA
~ FTF 15	76732.0	53230.0	284.5	286.5	227.5-197.5	247.5-197.5	8/14/84	NA	NA
~ FTF 16	76758.6	52879.8	286.8	288.6	233.8-203.8	252.8-202.8	10/22/84	NA	NA
~ FTF 17	76872.0	52884.0	287.6	289.6	230.6-200.6	247.6-200.6	8/17/84	NA	NA
~ FTF 18	76955.8	52879.2	286.3	288.0	232.3-202.3	236.3-202.3	10/15/84	NA	NA

GENERAL SEPARATION'S AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

^Source is unedited HP data

Updated 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
~ FTF 19	77139.1	52670.4	285.3	287.2	228.3-198.3	245.3-198.3	8/18/84	NA	NA
~ FTF 20	77015.0	52500.0	285.3	287.1	228.3-198.3	240.3-198.3	8/21/84	NA	NA
~ FTF 21	76866.7	52498.6	285.7	287.5	228.7-198.7	240.7-198.2	10/25/84	NA	NA
~ FTF 22	76751.3	52494.7	284.6	286.8	242.6-212.6	249.6-212.6	8/29/84	NA	NA
~ FTF 23	76611.8	52660.3	284.2	286.0	231.2-201.2	244.2-201.2	8/07/84	NA	NA
~ FTF 24A	77256.6	52780.8	268.4	270.3	232.7-212.7	250.4-212.6	5/02/84	NA	NA
~ FTF 25A	77308.4	52868.7	269.1	271.2	232.8-212.8	NA	4/30/84	NA	NA
~ FTF 26	77250.0	52875.4	268.8	270.9	226.3-206.3	238.8-188.8	5/09/84	NA	NA
~ FTF 27	77227.2	52823.5	268.5	270.5	243.5-213.5	265.5-215.5	4/20/84	NA	NA
~ H 2	72180.5	58786.3	262.8	262.9	244.5-234.5	244.8-229.8	3/08/56	NA	NA
~ H 4	72071.4	58741.3	258.7	258.8	245.2-235.2	240.7-229.7	3/08/56	NA	NA
~ H 6	72009.1	58335.4	259.4	260.1	235.2-225.2	244.4-225.4	4/26/56	NA	NA
~ H 7	71949.2	58336.1	257.3	257.8	234.9-224.9	240.3-225.3	4/25/56	NA	NA
~ H 8	71615.4	58233.9	257.1	257.2	228.4-218.4	NA	4/17/56	NA	NA
~ H 9	71572.6	58275.3	248.6	248.9	217.4-207.4	NA	4/19/56	NA	NA
~ H 10	71607.2	57822.8	254.5	257.3	232.5-222.5	237.5-222.5	5/01/56	NA	NA
~ H 11	71565.9	57779.4	246.0	249.3	222.0-212.0	226.0-211.0	5/02/56	NA	NA
~ H 12	71529.9	58490.3	NA	234.4	NA	NA	12/31/57	NA	NA
~ H 13	71327.4	58231.7	NA	244.4	NA	NA	2/20/62	NA	NA
~ H 14	71213.1	58071.9	NA	242.3	NA	NA	2/20/62	NA	NA
~ H 15	71231.1	57812.9	NA	240.1	NA	NA	2/20/62	NA	NA
~ H 16	71339.8	57631.9	NA	241.5	NA	NA	2/20/62	NA	NA
~ H 17	71479.4	57582.6	NA	243.3	NA	NA	2/20/62	NA	NA
~ H 18A	71339.6	57337.7	237.5	239.5	227.5-217.5	231.5-217.5	1/09/85	NA	NA
~ H 19	71434.2	57041.7	NA	246.6	NA	NA	2/20/62	NA	NA
~ HC 8A	77481.8	60058.5	NA	262.7	NA	NA	5/15/67	(1-240)	(0-250)

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note: * = Elevation below sea level; NA = Not available)

~Source is unedited HP data

Updated 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
^ HC 12A	73187.0	59504.0	287.3	NA	NA	NA	NA	(1-135)	(0-130)
^ HC 12B	73186.9	59488.4	287.3	290.1	NA	NA	6/23/67	NA	(0-112)
^ HC 35D	71918.0	58548.0	NA	NA	NA	NA	NA	(100-164)	(0-162)
HPT 1A	74774.4	60587.0	232.9	235.4	109.9-59.9	121.3-47.9	10/17/88	(17-185)	(0-174)
HPT 1B	74857.4	60590.0	233.1	235.6	133.1-123.1	143.6-121.1	10/12/88	NA	NA
HPT 2A	75061.8	60200.5	257.7	260.2	112.5-62.5	122.4-42.7	10/12/88	(17-215)	(2-177)
HPC-1	70395.4	62493.6	293.5	-	BORING	-	-	(11-998)	NA
^ HR8 11	71945.7	59559.8	257.4	259.2	237.6-207.9	252.4-207.4	10/19/84	NA	NA
^ HR8 12	71780.1	59330.1	255.7	257.5	235.9-206.3	250.7-204.7	10/22/84	NA	NA
^ HR8 13	71559.6	59300.2	251.2	253.1	231.4-201.7	246.2-201.2	10/22/84	NA	NA
^ HR8 14	71431.4	59612.1	251.8	253.6	231.9-202.3	246.8-200.8	10/23/84	NA	NA
HSB 65	72425.6	58432.0	270.0	272.0	242.0-212.0	246.0-212.0	9/24/81	NA	NA
HSB 65A	72436.2	58436.0	270.7	273.6	73.2-62.5	75.5-57.5	8/01/84	(0-241)	(0-193)
HSB 65B	72445.6	58439.4	270.9	273.7	133.3-123.3	142.3-110.4	8/01/84	NA	NA
HSB 65C	72439.6	58447.1	270.5	273.6	218.6-207.8	227.6-200.5	8/01/84	NA	NA
HSB 66	72429.2	56928.3	278.2	280.2	223.2-193.2	228.5-193.2	10/14/81	NA	NA
HSB 67	71505.0	58424.3	235.8	237.8	230.8-200.8	234.8-200.8	9/24/81	NA	NA
HSB 68	71528.0	56901.0	248.3	250.1	243.3-213.3	246.3-213.3	9/28/81	NA	NA
HSB 68A	71526.9	56892.1	247.4	249.4	58.0-47.5	60.4-42.3	8/01/84	(0-231)	(0-230)
HSB 68B	71525.5	56882.1	247.2	250.0	134.5-123.5	143.6-116.6	7/01/85	NA	NA
HSB 68C	71524.1	56872.7	246.7	250.1	179.5-168.4	187.5-161.5	7/01/85	NA	NA
HSB 69	71546.9	56475.1	234.0	236.0	229.0-199.0	232.0-199.0	9/25/81	NA	NA
HSB 69A	71549.4	56465.1	234.1	236.6	93.1-83.1	101.9-72.1	3/07/88	(7-156)	(0-155)
HSB 70	72606.9	55758.9	240.8	242.8	235.8-205.8	238.8-205.8	9/25/81	NA	NA
HSB 70C	72597.3	55757.1	240.9	243.1	174.9-164.9	186.9-150.9	3/22/88	NA	(0-90)
HSB 71	72875.9	55279.2	239.4	241.4	234.4-204.4	237.4-204.4	9/29/81	NA	NA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

~Source is unedited HP data

Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
HSB 71C	72866.6	55281.5	239.4	241.6	181.9-171.9	193.9-139.4	3/24/88	NA	(0-100)
HSB 83A	71648.6	58606.1	234.9	237.3	76.0-65.2	79.7-60.3	8/01/84	(0-224)	(0-143)
HSB 83B	71639.6	58594.9	234.8	237.0	132.0-121.1	141.2-113.2	7/01/85	NA	NA
HSB 83C	71636.9	58614.8	234.4	237.1	171.2-160.2	177.8-153.3	7/01/85	NA	NA
HSB 83D	71628.1	58601.7	234.4	237.0	228.7-198.7	232.7-190.7	7/01/85	NA	NA
HSB 84A	71586.2	56359.1	226.4	228.7	75.6-64.4	87.2-57.2	7/01/85	(2-211)	(0-173)
HSB 84B	71603.3	56352.4	226.8	228.9	132.7-121.6	140.4-113.8	7/01/85	NA	NA
HSB 84C	71597.1	56360.1	227.1	229.1	181.8-170.9	189.1-164.8	7/01/85	NA	NA
HSB 84D	71583.9	56349.9	225.8	228.8	219.6-199.6	223.0-194.5	7/01/85	NA	NA
HSB 85A	73791.9	58943.4	292.1	294.4	71.1-61.1	71.4-56.1	7/01/84	(2-273)	(0-213)
HSB 85B	73789.3	58953.3	292.0	294.5	143.2-133.2	162.7-128.3	7/01/84	NA	NA
HSB 85C	73802.3	58947.4	292.0	294.1	224.2-214.2	232.2-207.2	8/01/84	NA	NA
HSB 86A	72520.2	55985.9	260.0	262.4	73.9-63.1	76.2-58.2	8/01/84	(2-229)	(0-182)
HSB 86B	72519.0	55976.9	259.6	261.9	124.0-113.8	134.8-106.3	8/01/84	NA	NA
HSB 86C	72529.7	55984.6	260.5	262.9	199.4-189.4	209.0-181.4	8/01/84	NA	NA
HSB 86D	72522.1	55996.5	260.7	263.0	236.6-206.6	245.6-201.6	8/01/84	NA	NA
HSB 100C	72077.2	58806.5	258.0	260.2	163.0-153.0	173.6-143.0	11/12/87	NA	(0-115)
HSB 100D	72073.8	58796.9	257.9	260.1	236.9-216.9	251.3-202.9	10/28/87	NA	NA
HSB 101C	72001.9	58604.4	256.3	258.5	176.3-166.3	184.5-156.3	11/05/87	(26-94)	(0-100)
HSB 101D	71997.5	58594.8	256.4	258.7	236.1-216.1	246.8-206.1	11/11/87	NA	NA
HSB 102C	71960.1	58399.7	256.7	259.0	176.7-166.7	187.7-159.2	11/18/87	NA	(0-96)
HSB 102D	71962.4	58393.4	256.3	258.6	236.3-216.3	245.3-208.6	11/19/87	NA	NA
HSB 103C	71593.9	58323.6	245.2	247.4	169.2-159.2	180.2-150.2	10/30/87	NA	(0-86)
HSB 103D	71588.1	58315.6	245.4	247.6	233.7-213.7	241.6-205.4	11/02/87	NA	NA
HSB 104C	71376.8	58082.6	245.5	247.9	173.5-163.5	183.1-150.5	11/03/87	NA	(0-95)
HSB 104D	71370.2	58075.8	245.6	247.8	230.6-210.6	240.0-203.6	10/29/87	NA	NA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

~Source is unedited HP data

Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
HSB 105C	71447.3	57883.8	247.2	249.5	162.2-152.2	174.2-142.2	11/05/87	(31-105)	(0-97)
HSB 105D	71454.8	57877.4	247.2	249.5	231.8-211.8	243.3-202.2	11/06/87	NA	NA
HSB 106C	71720.9	57651.5	250.7	252.9	168.7-158.7	178.5-150.7	10/26/87	NA	(4-92)
HSB 106D	71727.8	57644.8	250.7	252.9	230.7-210.7	239.0-202.7	10/26/87	NA	NA
HSB 107C	71698.5	57432.0	258.3	261.6	168.3-158.3	179.8-146.1	10/26/87	(31-100)	(0-100)
HSB 107D	71696.6	57412.2	260.0	262.3	235.1-215.1	245.2-205.0	11/12/87	NA	NA
HSB 108C	71688.7	57155.5	263.9	266.2	196.1-186.1	206.4-174.9	10/14/87	NA	(0-85)
HSB 108D	71688.0	57145.6	264.0	266.3	232.0-212.0	241.9-204.0	10/26/87	NA	NA
HSB 109C	71684.8	56895.6	259.4	261.6	178.4-168.4	198.2-157.9	11/11/87	(30-100)	(0-98)
HSB 109D	71685.6	56885.5	259.0	261.2	233.0-213.0	248.3-204.0	11/10/87	NA	NA
HSB 110C	71779.3	56680.7	253.4	255.7	181.4-171.4	192.2-163.4	10/12/87	NA	(0-78)
HSB 110D	71785.2	56672.1	253.4	255.6	231.4-211.4	239.4-203.4	10/28/87	NA	NA
HSB 111C	71919.4	56501.9	253.7	256.0	150.7-140.7	161.2-130.7	11/17/87	(29-116)	(0-100)
HSB 111D	71926.2	56494.5	253.7	256.0	195.7-185.7	203.9-179.2	9/21/87	NA	NA
HSB 111E	71932.8	56487.2	253.7	255.9	231.7-211.7	242.7-203.7	10/23/87	NA	NA
HSB 112C	72156.4	56417.4	252.6	254.9	150.6-140.6	161.0-132.6	11/20/87	NA	(0-85)
HSB 112D	72161.6	56408.1	252.8	255.1	198.3-188.3	206.0-167.8	10/08/87	NA	NA
HSB 112E	72166.6	56399.5	252.7	255.1	231.7-211.7	243.2-202.7	10/27/87	NA	NA
HSB 113C	72312.3	56160.4	258.7	261.0	161.7-151.7	176.9-143.7	11/24/87	(35-106)	(0-107)
HSB 113D	72302.7	56164.3	258.7	260.9	235.7-215.7	248.4-208.7	11/20/87	NA	NA
HSB 114C	72464.6	56107.0	261.5	263.8	195.6-185.6	205.5-174.5	10/16/87	NA	(0-86)
HSB 114D	72474.2	56104.6	261.8	264.0	232.8-212.8	242.5-205.8	10/21/87	NA	NA
HSB 115C	72653.2	56043.2	266.8	269.3	192.8-182.8	206.4-173.8	10/07/87	(0-85)	(0-84)
HSB 115D	72662.3	56039.8	266.9	269.1	233.9-213.9	243.3-206.9	10/20/87	NA	NA
HSB 116C	72888.1	55989.1	255.3	257.5	190.5-180.5	197.3-170.3	10/20/87	NA	(0-82)
HSB 116D	72898.1	55988.2	254.5	256.8	234.5-214.5	243.0-208.0	9/02/87	NA	NA

GENERAL SEPARATIONS AREA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

~Source is unedited HP data

Update 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
HSB 117A	72733.6	55170.1	234.1	236.3	94.1-84.1	104.1-74.1	1/25/88	(22-160)	(0-159)
HSB 117C	72740.7	55162.9	234.0	236.3	174.0-164.0	182.5-154.0	1/27/88	NA	NA
HSB 117D	72747.6	55155.6	234.1	236.3	219.1-199.1	225.1-190.1	1/27/88	NA	NA
HSB 118A	72696.4	55775.6	245.0	247.3	101.0-91.0	111.0-81.0	11/25/87	(31-165)	(0-155)
HSB 119A	73082.5	56100.2	254.8	257.1	103.3-93.3	113.3-84.8	10/05/87	(33-167)	(0-165)
HSB 120A	73395.1	56431.9	266.0	268.2	101.0-91.0	111.0-81.0	11/13/87	(33-185)	(0-182)
HSB 121A	72024.8	57389.6	272.3	274.6	98.3-88.3	108.1-78.3	11/02/87	(33-191)	(0-186)
HSB 122A	72195.9	57747.4	269.4	271.6	95.4-85.4	105.2-77.4	10/16/87	(31-190)	(0-186)
HSB 123A	72189.8	58124.8	262.3	264.5	102.3-92.3	109.7-78.3	10/08/87	(31-183)	(0-180)
HSB 124A	72199.6	58514.6	263.9	266.2	103.1-93.1	111.4-82.9	9/16/87	(33-179)	(0-174)
HSB 125C	71503.6	58592.8	229.6	231.9	155.6-145.6	164.6-133.6	3/17/88	NA	(0-96)
HSB 125D	71498.2	58584.1	229.4	231.7	219.4-199.4	223.4-187.4	3/17/88	NA	NA
HSB 126C	70627.7	57178.2	210.3	212.6	181.3-176.3	190.6-129.8	2/03/88	NA	(0-76)
HSB 126D	70633.4	57169.6	210.5	212.7	200.5-190.5	206.7-185.0	2/05/88	NA	NA
HSB 127C	71210.1	56792.1	223.4	225.7	158.4-148.4	169.1-128.4	3/02/88	NA	(0-92)
HSB 127D	71218.9	56788.0	223.8	226.1	217.8-197.8	221.8-188.3	3/02/88	NA	NA
HSB 129C	71830.4	55110.0	212.8	215.1	157.8-147.8	168.8-122.8	3/18/88	NA	(0-86)
HSB 129D	71837.1	55103.4	212.7	214.7	205.2-185.2	210.2-176.7	3/17/88	NA	NA
HSB 130C	70762.4	54643.6	215.9	218.3	169.9-159.9	178.9-133.9	3/23/88	NA	(0-77)
HSB 130D	70757.2	54651.7	216.1	218.6	202.1-182.1	209.6-174.1	3/23/88	NA	NA
HSB 131C	70374.7	56894.9	209.5	211.7	158.5-148.5	169.2-134.5	2/26/88	NA	(0-69)
HSB 131D	70365.0	56891.1	209.8	212.1	205.7-195.7	208.6-172.8	2/26/88	NA	NA
HSB 132C	71472.4	58787.7	238.3	240.5	178.6-168.6	188.8-123.3	2/22/88	NA	(0-111)
HSB 132D	71469.5	58799.3	238.5	240.7	226.5-206.5	235.6-198.3	2/22/88	NA	NA
HSB 133C	71949.5	59110.3	253.4	255.6	183.3-173.3	193.8-152.9	1/25/88	NA	(0-99)
HSB 133D	71943.5	59102.3	253.1	255.3	228.5-208.5	236.1-200.1	1/26/88	NA	NA

GENERAL SEPARATIONS ~ REA MASTER WELL LIST

Well Data (Note- * = Elevation below sea level; NA = Not available)

~Source is unedited HP data

Updated 6/27/90

Well	North	East	Ground Elev. (ft)	TOC Elev. (ft)	Screen Elev. (ft)	Sand Pack Elev. (ft)	Date Installed	Cored Interval (depth in ft)	Geophysics (depth in ft)
HSB 134C	71210.3	58289.9	236.1	238.4	159.1-149.1	170.1-140.1	2/22/88	NA	(0-93)
HSB 134D	71217.3	58296.5	235.9	238.1	225.8-205.8	232.4-199.9	2/15/88	NA	NA
HSB 135C	71390.2	56560.8	229.8	232.0	157.3-147.3	169.8-134.8	3/24/88	NA	(0-95)
HSB 135D	71396.7	56552.8	229.9	232.3	219.9-199.9	224.9-187.9	3/11/88	NA	NA
HSB 136C	71900.3	55949.6	225.6	227.9	170.5-160.5	181.1-135.6	3/08/88	NA	(0-90)
HSB 136D	71906.0	55941.7	225.7	228.0	220.2-200.2	223.7-188.7	3/07/88	NA	NA
HSB 137C	72269.9	55700.2	233.8	236.0	173.8-163.8	184.3-143.8	3/10/88	NA	(0-90)
HSB 137D	72278.9	55696.1	234.3	236.6	225.3-205.3	231.3-193.3	3/14/88	NA	NA
HSB 138D	73160.2	55260.7	250.1	252.4	228.1-208.1	236.6-200.1	3/24/88	NA	NA
HSB 139A	71127.4	57365.4	231.5	233.7	97.6-87.6	105.9-77.5	2/12/88	(11-154)	(0-151)
HSB 139C	71129.8	57374.5	231.5	233.8	158.5-148.5	169.2-117.5	3/16/88	(91-114)	(0-153)
HSB 139D	71133.2	57384.4	231.7	233.8	226.7-206.7	231.2-196.7	2/16/88	NA	NA
HSB TB	72394.0	58696.1	267.1	269.6	*252.9-*295.1	*234.7-*462	12/07/88	(11-730)	(0-730)
IDL 3A	78210.0	50240.0	257.4	260.4	NA	NA	NA	NA	(0-300)
P 18TA	67576.5	47652.8	296.7	298.7	*531.9-*558.3	*516.3-*263.3	6/13/85	(0-952)	(0-950)
P 27TA	70382.0	64022.9	274.1	275.8	*532.7-*542.9	*515.9-*560.9	10/07/86	(0-899)	(0-899)
P 28TA	79284.3	55441.1	285.2	287.7	*424.8-*504.8	*462.0-*494.6	7/08/86	(0-842)	(0-838)

APPENDIX II

DETAILED DESCRIPTIONS OF THE LITHOFACIES OF THE HYDROLOGIC UNITS ON REGIONAL CROSS SECTIONS

Cross Section #1

Cross Section #1 (Plate 1) represents a 2.4-mile line of section from north of the MWMF to southeast of H-Area Seepage Basins. The section follows the southeast regional dip of the GSA. Topography is relatively flat with elevations ranging from 270 to 292 feet above msl. The depression between borings P-28 and BGO-10 is an unnamed stream that drains into UTRC. The bedding is essentially flat.

Cross Section #1 illustrates all of the hydrostratigraphic units previously described. Confining System I-II was penetrated in borings P-28, HSB-85, and P-27, attaining a thickness of over 100 feet in P-28 and P-27. Little data are available in the vicinity of HSB-65. The dashed line represents where the top of the confining system would be located based on stratigraphic position.

Aquifer Unit IIA was not fully penetrated in borings BGO-10, BGO-8 and BGO-6. Where the unit was fully penetrated it consists of approximately 55 to 75 feet of clean sand with minor interbeds of clayey sand. Confining Unit IIA-IIB is interpreted to thicken and thin rapidly on Cross Section #1. The unit is approximately 20 feet thick in P-28 to the northeast. At the MWMF the unit thickens to 30 feet in BGO-6 and thins to 5 feet to the southeast at P-27. The units between HSB-85 and P-27 consist of two separate lenses of clay. The dashed line represents where the lenses would meet based on available data. The lithofacies of Confining Unit IIA-IIB consists predominantly of clays and sandy clays with minor interbeds of calcareous muds.

The thickness of Aquifer zone IIB₁ varies from 60 to 75 feet. The calcareous sands, mud, and limestone beds in the zone attain a thickness of 45 feet to the northeast, but thin to approximately 25 feet to the southeast. Clastic interbeds observed within the calcareous facies near BGO-6 are not uncommon. Generally, lithofacies vary quickly and abruptly within short distances. This could be indicative of local changes in the depositional environment such as water depth. Clastic lithologies may have been calcareous at the time of deposition with post-depositional ground water movement possibly removing the calcareous material within localized areas.

Confining zone IIB₁-IIB₂ is a series of thin clay beds, sandy clay beds, and clayey sand beds attaining of thickness of 20 feet to the northwest. To the southeast it thins to 5 feet and becomes a clay of intermediate thickness between HC-12 and P-27.

Aquifer zone IIB₂ has an average thickness ranging from 70 to 80 feet. The zone becomes slightly thicker to the southeast attaining a thickness of 90 feet. To the northeast this zone becomes very clay-rich in borings P-28, HC-12 and P-27. At other locations this zone is predominantly clayey sand with few beds of clean sand. The dip of the beds steepens between HSB-85 and HC-12 within Confining zone IIB₁-IIB₂ and Aquifer zone IIB₂. This is interpreted to be a local facies change or a localized post-depositional erosional surface.

2) Cross Section #2

Cross Section #2 (Plate 2) represents a 3.6-mile line of section from north of F-Area Seepage Basins to east-southeast of H-Area Seepage Basins. This is essentially along the regional dip. Topography is undulating with elevations ranging from 304 to 208 feet above msl. An unnamed

tributary of Fourmile Branch exists between HSB-TB, HSB-133 and HPC-1. The incised area between DRB-4 and HSB-117 is an effluent stream for the F-Area facility to the north. The bedding is essentially flat lying.

A prominent structural feature on Cross Section #2 is the normal fault between HPC-1 and P-27. This fault is downthrown to the east with a throw of approximately 20 feet. Data accumulated in the Pen Branch Fault investigation suggests the offset may actually be an anticlinal flexure overlying and controlled by deeper seated faulting associated with reactivation of basement faults. Thus this feature is considered to be structurally controlled.

All of the hydrostratigraphic units are illustrated on this cross section. Confining System I-II attains a thickness of approximately 100 feet in DRB-4. The geophysical and core data provide good correlation for this system. At the H-Area Seepage Basins the system was not fully penetrated. The dashed line represents where the top of Confining System I-II would be located based on stratigraphic position. Confining System I-II consists predominantly of clay/silts, sandy clay/silts, and clayey sands of equal thicknesses. Few clean sands are present.

Aquifer Unit IIA was not fully penetrated in HSB-117, HSB-119, HSB-65, or HSB-123. Where the unit was fully penetrated it consists predominantly of sands and clayey sands. The unit becomes clayey at FSB-76 and FSB-TA. HSB-65 contains a 20-foot thick calcareous sand bed. Both the clay and calcareous sand beds are attributed to local facies change.

Aquifer Unit IIA thins considerably to the east-southeast. In the northeast the unit attains a thickness of around 100 feet thinning to 50 to 65 feet in the east-southeast area. The thickening to the northeast may be related to the absence of carbonate sediments that tend to dominate in the vicinity of the H-Area Seepage Basins. To the northwest the clay beds of Confining Unit IIA-IIB tend to be stratigraphically higher than in the areas of carbonate buildup. The clay beds of the

confining unit underlie the carbonates and are stratigraphically lower in the section. Aquifer Unit IIA is thinnest in these areas.

Confining Unit IIA-IIB is relatively thin throughout Cross Section #2, attaining a maximum thickness of 15 feet in HSB-65. Elsewhere the unit is approximately 5 feet thick. Cross Section #2 illustrates the separate lenses of green clays that thicken and thin, and pinch out abruptly. The clay lens in IDL-3 pinches out before meeting a lower clay lens in FSB-76. This upper clay lens appears downdip at HPC-1 and truncates abruptly into a limestone lithofacies in Aquifer zone IIB₁. No lower clay lenses are present at HPC-1.

Aquifer zone IIB₁ thins to around 50 feet in the north of the F-Area Seepage Basins, thickens to about 90 feet in the H-Area Seepage Basins and thins again to around 65 feet east-southeast of the basins. This fluctuation in thickness is attributed to the thick carbonate sequence present near the H-Area Seepage Basins. The isolated carbonate section in FSB-TA was interpreted to be carbonate, based on the daily detailed drilling report. Lost circulation and cuttings data all indicate the presence of a calcareous zone.

Aquifer zone IIB₁ consists predominantly of clayey sand beds. Several clean sand beds are scattered throughout this zone. The calcareous facies of Aquifer zone IIB₁ consists mainly of limestones and sandy limestones beds. A few thin clastic beds are intertonguing with the calcareous beds. This is attributed to a local facies change or carbonate dissolution within the clastic beds, reducing the percentage of carbonate.

Confining zone IIB₁-IIB₂ thickens and thins very abruptly on Cross Section #2, ranging in thickness from 0 to 25 feet. No confining zone was encountered in the vicinity of IDL-3 and HSB-123. Between HSB-133 and HPC-1 the clays of the confining zone are interpreted to be discontinuous and, based on available data, may or may not be interconnected.

Aquifer zone IIB₂ thickens and thins in response to the topography along the cross section. This zone becomes thin in the vicinity of HSB-117 with a thickness of approximately 40 feet and increases to the north and east-southeast to a thickness of 90 to 100 feet. Clayey sediments dominate this zone with the exception of FSB-TA and IDL-3. FSB-TA may possibly represent a sand infilled channel.

3. Cross Section #3

Cross Section #3 (Plate 3) represents a 2.7-mile line of section from northeast of the F-Area Seepage Basins to south-southwest of the F-Area Basins. The section roughly follows the regional strike. Topography is undulating with elevations ranging from 180 feet above msl at Fourmile Branch to 297 feet at P-18. The beds are essentially flat lying with the exception of the northeast area between P-28 and FSB-76 where Confining zone IIB₁-IIB₂ and Aquifer zone IIB₂ show a slight steepening of the beds to the southwest. Within this area Confining zone IIB₁-IIB₂ could be a post-depositional erosional surface resulting in a change of dip.

A prominent feature on Cross Section #3 is the offset of beds between FSB-91 and FSB-100. Beds are downthrown approximately 12 feet to the south. This feature is possibly structurally controlled or sedimentologically controlled. Often areas of offset are related to carbonate buildups. There is the strong possibility of slumping associated with carbonate dissolution. Slump features were recognized in the coastal plain calcareous sediments by Johnson and Heron (1965). Features of this type are also discussed by McDowell and Houser (1983). The offset has been projected to the top of Confining System I-II however, this interpretation is based on the limited data available. Geophysical and core data both corroborate the offset in Aquifer zone IIB₁, Confining zone IIB₁-IIB₂, and Aquifer zone IIB₂.

A localized steepening of dip is observed in borings FSB-106, FSB-91, and FSB-100 within Aquifer zone IIB₁, in association with the area of offset. The steepening could be caused by the proposed slumping due to carbonate dissolution. Another possibility is clastic sediments draped over the flanks of the carbonate mounds.

Confining System I-II is represented by four borings on Cross Section #3 (Plate 3). In P-28 the system attains a thickness of approximately 100 feet; in P-18, a thickness of around 50 feet; and in FSB-79 and FSB-76, only the very top (approximately 10 feet) was penetrated. The dashed line represents where the top of the confining system would be located based on stratigraphic position for borings FSB-91, FSB-100, and FSB-101.

At the northeast end of the section at P-28 Aquifer Unit IIA reaches a thickness of approximately 55 feet, thickening to the south-southwest to 105 feet at P-18. The thickening and thinning of this unit and its relation to the top of Confining System I-II is well illustrated in Figures 19 and 21. The structure contours on Confining System I-II reveal a regional dip to the south-southeast while the isopach of Aquifer Unit IIA illustrates a thickening in the same direction. This supports the interpretation of channel erosion into the Ellenton Clays of Confining System I-II infilled by sands of Aquifer Unit IIA (Congaree Aquifer) deposited on the erosional unconformity. Aquifer Unit IIA is composed predominantly of sands and clayey sands. Minor interbeds of clay and sandy clay are present. A thin stringer of calcareous sand is present in P-18.

Confining Unit IIA-IIB consists of three separate lenses in the southwest between P-18 and FSB-79. Only the thinner lower lens continues through the F-Area Seepage Basins. The unit thickens around FSB-76, and continues to thicken, and becomes two distinct clay lenses to the northeast. Generally, the unit consists of clay and sandy clay beds with minor clayey sand and sand beds.

Aquifer zone IIB₁ is relatively uniform in thickness varying from 45 to 60 feet throughout Cross Section #3. The zone is predominantly calcareous to the southwest near P-18, becoming less calcareous at the F-Area Seepage Basins. To the northeast only clastic lithofacies are found. The zone is primarily clayey sand beds with a few thick, clean sand beds.

Confining zone IIB₁-IIB₂ is about 17 feet thick to the southwest at P-18. The zone thins to 2 to 3 feet thick in the vicinity of FSB-91 and FSB-100 and thickens again to about 12 feet at P-28 to the northeast. The lithofacies are primarily clay with minor interbeds of clayey sand.

Aquifer zone IIB₂ becomes very thin around Fourmile Branch. Thicknesses range to over 100 feet to the northeast and southwest. The lithofacies are quite clayey to the northeast at P-28 and FSB-76. Around the F-Area Seepage Basins sands and clayey sands dominate the lithofacies. To the southwest at P-18 the section includes equally thick beds of sand, clays, clayey sands, and sandy clay.

4. Cross Section #4

Cross Section #4 (Plate 4) represents a 1.3 mile line of section from northeast of the HWMF to the south-southeast of H-Area Seepage Basins. The section roughly follows the regional strike. This section intersects Cross Section #1, Cross Section #2 and Cross Section #5A. Topography is relatively flat with the steepness increasing near Fourmile Branch. Elevations range from 284 feet at BGO-6 to 192 feet at Fourmile Branch. The bedding is essentially flat lying.

Confining System I-II was only penetrated at HSB-86 on Cross Section #4. Approximately 30 feet of sandy clay, silt, and clayey sand beds of equal thickness are present at this location. The

dashed line indicates where the top of the confining system would be located based on its stratigraphic position at HSB-129 and HSB-130.

Aquifer Unit IIA was fully penetrated at HSB-86 and partially penetrated at HSB-119 and HSB-120. The unit attains a thickness of 50 feet at HSB-86. The unit is composed of thick sand and clayey sand beds. One calcareous sand stringer was encountered at HSB-120. The dashed line indicates where the top of Aquifer Unit IIA would be located based on stratigraphic position.

Confining Unit IIA-IIB consists of a relatively thin, sandy clay to the south at HSB-86. The unit becomes progressively thinner at HSB-120, having a thickness of only 2 feet. To the northeast the unit thickens and is composed of four separate lenses of clay and calcareous mud with minor interbeds of clayey sand. The confining unit attains a thickness of 30 feet at BGO-6.

Aquifer zone IIB₁ is variable in thickness. To the south-southeast the zone is approximately 60 feet thick. In the vicinity of the H-Area Seepage Basins the zone thickens to around 80 feet. To the northeast at BGO-6 the zone thins to approximately 60 feet. The zone consists predominantly of sands and clayey sands with minor interbeds of thin clays. One muddy limestone is present at BGO-6.

Confining zone IIB₁-IIB₂ is very variable in thickness on Cross Section #4. The zone attains a maximum thickness of 32 feet to the south-southeast at HSB-137 and a minimum thickness of 9 feet to the south where it is incised by Fourmile Branch. The confining zone consists predominantly of clay and sandy clay beds 1 to 13 feet thick interbedded with clay sand and sand beds 2 to 8 feet thick.

Aquifer zone IIB₂ is truncated by Fourmile Branch. The zone attains a maximum thickness of approximately 75 feet to the northeast between BGO-21 and BGO-6. The lithofacies are very

clayey throughout the zone. Interbeds of clayey sands and sands intertongue with the clayey sediments throughout Cross Section #4. Occasional pebble and gravel beds are present to the northeast.

5) Cross Sections #5A and #5B

Cross Sections #5A and #5B (Plates #5A and #5B) represent an east-west 2.3-mile line of section from east of the MWMF to the western edge of the F-Area Seepage Basins. There are 18 borings represented on these two sections. This section was split into "A" and "B" for the ease of handling. A match line is provided for correlation. Topography is undulating with elevations ranging from 311 above msl at BGO-12 to 233 feet above msl at HPT-1. HPT-1 and HPT-2 are near the flanks of an unnamed tributary of UTRC. Beds are essentially flat lying on the outer edges of the cross sections. Around the MWMF the dip steepens in Aquifer zone IIB₁, Confining zone IIB₁-IIB₂, and Aquifer zone IIB₂.

The dip could be the result of clastic deposits draped over the flanks of carbonate mounds. Localized post-depositional erosional surfaces could also change the dip. An example of this would be the beds between BGO-12 and BGO-10. Here dissolution of the carbonate beds in Aquifer zone IIB₁ could result in slumping of the overlying sediments.

A prominent feature on Cross Section #5B is the offset beds between FSB-97 and FSB-98. The feature exhibits normal offset, and beds are downthrown to the west with a throw of approximately 9 feet in Aquifer zone IIB₂. Throw decreases to around 3 feet in Confining zone IIB₁-IIB₂ and Aquifer zone IIB₁. The offset apparently does not penetrate Confining Unit IIA-IIB. The offset cuts FSB-97 at 121 feet above msl. Geophysical and core data both corroborate this offset. This feature is either structurally or sedimentologically controlled. There is a good possibility of slumping associated with carbonate dissolution as previously discussed. Due to the close

proximity of calcareous beds in the nearby boring FSB-100, there is a strong possibility that borings FSB-97 and FSB-98 once contained carbonate beds and post-depositional carbonate dissolution is a reasonable explanation for the offset observed between FSB-97 and FSB-98. Slump features were recognized in the calcareous sediments described by Johnson and Heron (1965). Soft sediment deformation such as deltaic slumping is another feasible explanation for the offset. Features of this type are discussed in detail by McDowell and Houser (1983).

An offset of beds is also on Cross Section #5B. A normal down to the south feature is represented on the Cross Section between borings BGO-6 and BGO-5 and HPT-2. The line of section intersects the offset so that the offset is displayed twice. Well BGO-5 is on the upthrown side of the offset and wells BGO-6 and HPT-2 are on the downthrown side. The offset has a throw of 5 to 7 feet in Aquifer IIB₂ and Confining zone IIB₁-IIB₂. The throw increases to approximately 10 feet in Aquifer zone IIB₁. The offset possibly dies out in Aquifer zone IIB₁, but data is limited in this area. Again the offset may be due to slumping of the younger sediments overlying the area where carbonate dissolution had occurred.

Confining System I-II was only penetrated at FSB-TA to the west and at HPT-1 and HPT-2 to the east. The dashed line indicates the confining system's stratigraphic position. Where the system is observed it consists of clays/silts and sandy clays with interbeds of clayey sands and sands.

Aquifer Unit IIA was only fully penetrated at FSB-TA to the west and at HPT-1 and HPT-2 to the east. Total thickness ranges from 60 to 70 feet at these locations. The unit is composed predominantly of clean sand beds with minor interbeds of clayey sand and sand beds. A thinning of the unit is inferred beneath BGO-6 and BGO-5 due to the thickening of the Confining Unit IIA-IIB that overlies Aquifer Unit IIA.

Confining Unit IIA-IIB is thin from the western edge of the F-Area Seepage Basins and becomes very thin on the southwestern edge of the MWMF. Thicknesses range from 3 to 5 feet at these locations. On the southern edge of the MWMF the unit thickens to 25 feet and remains thick until borings HPT-1 and HPT-2 are reached to the west. Here the unit is 4 to 5 feet thick. The confining unit is predominantly a clay and sandy clay until it reaches the southern edge of the MWMF at BGO-25, BGO-14, and BGO-12. The unit thickens and contains clays and clayey sands with interbedded sands. The clayey sand and sand lithofacies were included in the confining zone because this lithofacies is predominantly consolidated indurated sandstone in this area, where it is a very fine to fine sand sized and tightly cemented with silica cement.

West of BGO-12 the unit contains calcareous muds, clay, and clayey sands thinning back to a clay at BGO-8. At BGO-6 the unit thickens to 30 feet. The clays separate into four separate lenses and become calcareous at the top of the unit. Interbeds of clayey sand are included in this confining unit (Confining Unit IIA-IIB). Due to limited data this unit is dashed with question marks between BGO-8 and HPT-2. To the east the unit becomes a clay and sandy clay thinning to 5 feet.

Aquifer zone IIB₁ is variable in thickness. At the western edge of the F-Area Seepage Basins the thickness is approximately 45 feet. To the west and at the MWMF the zone thickens to 95 feet. From the F-Area Seepage Basins the zone is predominantly composed of clastic lithofacies with calcareous lithofacies present on the eastern edge of the basins near FSB-100 and FSBTA. Clastic lithofacies are composed of relatively thick sands (up to 20 feet thick) with minor interbeds of clayey sands, clays, and sandy clays. Calcareous lithofacies consist of calcareous sands and muddy and sandy limestones. The limestone in FSB-TA is based on daily detail drilling reports compiled by the drilling and oversight personnel. Lost circulation and cuttings data all indicate the presence of a calcareous zone.

Aquifer zone IIB₁ is predominantly a calcareous lithofacies under the MWMF with the exception of BGO-12 and BGO-6. Well BGO-12 contains no calcareous sediments. Based on the isopach of the lower carbonate zone (see Figure 17) it is possible that the carbonate deposition was absent at this location. However, another possibility is the previously discussed carbonate dissolution of the sediments. The overlying steeply dipping clastic beds in BGO-12 could be a result of slumping from carbonate dissolution of calcareous beds.

The calcareous lithofacies under the MWMF are composed predominantly of calcareous sands, muddy limestone, and sandy limestones with minor interbeds of calcareous muds. The clastic lithofacies in this area includes interbeds of sands and clayey sands with minor interbeds of clay and sandy clay of equal thicknesses.

To the east Aquifer zone IIB₁ in HPT-1 and HPT-2 consists predominantly of clastic lithofacies with minor interbeds of calcareous sands in the lower portion. The clastic lithofacies include interbeds of equal thickness of sands and clayey sands. Minor clays and sandy clays are also present.

Confining zone IIB₁-IIB₂ varies in thickness throughout Cross Sections #5A and #5B. Thicknesses range from 5 to 20 feet. Lithofacies are predominantly clastic. One thin stringer of calcareous sand is present at BGO-25 at the southwestern edge of the MWMF. Clastic lithofacies are predominantly clay and sandy clay with minor interbeds of clayey sand and sand from the western edge of the F-Area Seepage Basins to the southwestern edge of the MWMF. At this location the clay beds become a series of thin lenses with interbeds of clayey sand and sands. At the very eastern edge of Cross Section #5A Confining zone IIB₁-IIB₂ was not encountered at HPT-1 and HPT-2. This is illustrated in the isopach map of Confining zone IIB₁-IIB₂ (Tan Clay) (see Figure 26).

Confining zone IIB₁-IIB₂ is variable in thickness due to topographic undulations. Lithological thickness is approximately 100 feet at FSB-TA and 23 feet at HPT-1. The zone is composed entirely of clastic lithofacies; predominantly sands and clayey sands. This zone becomes very sandy at the F Area Seepage Basins and in the vicinity of BGO-8. Elsewhere abundant lenses of clay are present.

PHYSIOGRAPHIC SUBPROVINCES OF THE SOUTH CAROLINA COASTAL PLAIN

FIGURE 1

LOCATION OF GENERAL SEPARATIONS AREA

FIGURE 2

LOCATION OF GENERAL SEPARATIONS AREA

FIGURE 2

NOTE: Arrows Denote Approximate Drainage Pattern

SOURCE: 1:48,000 Savannah River Plant Topographic Map

Figure 3. Surface Water Drainage Map in the Vicinity of the F and H Area Seepage Basins

TECTONIC SETTING OF EASTERN UNITED STATES

LATE PRECAMBRIAN EXTENSION

LATE PRECAMBRIAN SPREADING

LATE PRECAMBRIAN - EARLY CAMBRIAN

MIDDLE ORDOVICIAN - SILURIAN (TACONIC OROGENY)

DEVONIAN - EARLY CARBONIFEROUS (ACADIAN OROGENY)

LATE CARBONIFEROUS - PERMIAN (ALLEGHENIAN OROGENY)

REGIONAL STRATIGRAPHY

PREPARED BY: ROLF K. AAO
DRAWN BY: ROD A. I
MODIFIED FROM D. COLQUHOUN

CROSS-SECTION

1989
T
ETAL. 1983

SOUTHEAST

DWG. NO. R00016

COASTAL PLAIN SEDIMENTS AT THE SRS

Groups	Formations	Period/Epoch	N. American Stage	European Stage
	"Upland Unit"	Age Unknown	Age Unknown	Age Unknown
Barnwell Group	Tobacco Road Sand	Tertiary	Jacksonian	Priabonian
	Dry Branch Formation			
	Irwinton Sand Mbr Griffins Landing Mbr			
Orangeburg Group	Clinchfield Formation		Clalbornian	Bartonian
	Santee Limestone Formation			Lutetian
	McBean Member Cow Cow Member Warley Hill Member			Ypresian
Black Mingo Group	Congaree Formation		Sabinian	Thanetian
	Snapp Mbr Williamsburg Fm		Midwayan	Danian
	Ellenton Member, Rhems Formation			
Lumbee Group	Steel Creek Member, Peedee Formation	Late Cretaceous	Navarroan	Maestrichtian
	Black Creek Formation		Tayloran	Campanian
	Middendorf Formation		Austinian	Santonian
	Cape Fear Formation			— ? — Coniacian? Turonian?

Crystalline Triassic Basement
Newark Supergroup

Triassic or Paleozoic or Precambrian

SURFACE DRAINAGE MAP OF THE SAVANNAH RIVER SITE

FIGURE 7

SOURCE: URANIUM SOLIDIFICATION FACILITY SITE CHARACTERISTICS,
APPENDICES 3A, 3B, 3C: OCTOBER, 1989.

Geologic Age	COMPARISON OF H				
	Siple (1967)	SRP Baseline Hydrogeologic study		GeoTrans (1989)	Auc
Tertiary	Hawthorn Aquifer	Upland Unit		Aquifer 4	Tert A
	Barnwell Aquifer	Barnwell Group	Tobacco Rd FM		
			Dry Branch FM		
	McBean Aquitard Green Clay	McBean Formation Green Clay			
	Congaree Aquifer	Congaree		Aquifer 3	
	Ellenton Aquitard	Williamsburg Formation		Aquitard 2	Confi
		Ellenton Formation			
Cretaceous	Upper Tuscaloosa Aquifer	Pee Dee Formation		Aquifer 2	Blk A
	Mid Tuscaloosa Clay Aquitard	Black Creek Formation		Aquitard 1	Coni
	Lower Tuscaloosa Aquifer	Middendorf Formation		Aquifer 1	MI A
	Basal Clay Aquitard	Cape Fear Formation		Base of Model	Confi
Triassic or Paleozoic Basement					

NOTE:

- (1) Included within the different units can be individual zones of relatively higher or lower permeability.
- (2) Hydrostratigraphic boundaries do not necessarily coincide with stratigraphic boundaries.

SAVAN

Add
Prep

HYDROSTRATIGRAPHIC UNITS USED AT SRS

Price (1988)		Proposed Nomenclature Aadland & Bledsoe (1990)				
dry Sand lifer	Zone 8		Aquifer System 1/11	Aquifer unit 1/11C	Aquifer Unit 11B ⁽¹⁾	Aquifer System 11
	Zone 7	7a
		7b				
		7c				
	Zone 6					
Zone 5	5b	Aquifer Unit 11A				
	5a					
ing Unit	Zone 4		Confining Unit 1/11B-1/11C		Confining System 1-11	
Creek lifer	Zone 3	3b	Aquifer Unit 1/11B	Aquifer Unit 11B ⁽²⁾	Aquifer System 1	
		3a				
ing Unit	Zone 2	2c	Confining Unit 1/11A-1/11B	Confining Unit 1A-B		
endorf lifer		2b	Aquifer unit 1/11A	Aquifer Unit 1 A		
		2a				
ing Unit	Zone 1		Confining System I			Confining System I
			Paleozoic - Triassic Basement Hydrologic System			

SAH RIVER SITE

and Bledsoe, 1990

red By: JA Pivovar

Paleozoic - Triassic
Basement
Hydrologic System

FIGURE 8

HYDROS

Sat

Hydrostratigraphic Units

Paleozoic - Triassic
Basement
Hydrologic System

RATIGRAPHIC CHART

nnah River Site

Rolf K. Aadland
SS-SRL 10/26/89

Prepared by: JA Pivovar

FIGURE 9

N. American Stage	European Stage	Groups	Formations (Price et al., 1988)	
Age Unknown	Age Unknown		Upland Unit	
Jacksonian	Priabonian	Barnwell Group	Tobacco Road Sand	
			Dry Branch Formation	Irwinton Sand Mbr Twigg Clay Mbr Griffith Landing Mbr
			Clinchfield Formation	
Clabornian	Bartonian	Orangeburg Group	Santee Limestone	McBean Member Cox Cove Member Warley Hill Member
	Lutetian			Congaree Formation
Sabinian	Ypresian	Black Mingo Group	4 Mile Mbr Fishburne Fm	
Midwayan	Thanetian		Snapp Mbr Williamsburg Fm	
	Danian		Ellenton Member, Rhems Formation	
Navarroan	Maestrichtian	Lumbee Group	Steel Creek Member, Peedee Formation	
Tayloran	Campanian		Black Creek Formation	
Austinian	Santonian		Middendorf Formation	
	— ? — Coniacian? Turonian?		Cape Fear Formation	

Aquifer System 1/11

Paleozoic Crystalline Basement
Triassic Newark Supergroup

SCHEMATIC HYDROS
OF THE SAVANNAH
Aadland & F

STRATIGRAPHIC CHART

H RIVER SITE

Bedsoe, 1990

Prepared by: JA Pivovar

FIGURE 10

REGIONAL HYDROSTRATIGRAPHY

PREPARED BY: ROLF K. AADLAI
DRAWN BY: ROD A. BOL
MODIFIED FROM D. COLQUHOUN E

HIC CROSS-SECTION

1989

. 1983

FIGURE 11

DWG. NO.
R00017

Geologic Age	Lithostratigraphic Units		Hydrostratigraphic Units		
Tertiary		Upland Unit	Aquifer Unit IIB	Aquifer Zone IIB ₂	Aquifer System II
	Barnwell Group	Tobacco Rd FM		Confining Zone IIB ₁ -IIB ₂	
		Dry Branch FM		Aquifer Zone IIB ₁	
	McBean Formation Green Clay			Confining Unit IIA-IIB	
	Congaree		Aquifer Unit IIA		
	Williamsburg Formation		Confining System I-II		
	Ellenton Formation				

**HYDROSTRATIGRAPHIC NOMENCLATURE
FOR GENERAL SEPARATIONS AREA
SAVANNAH RIVER SITE**

(Aadland, 1990)

T.L. Mickley

FIGURE 12

SCHEMATIC MAP AND CROSS SECTION OF FLUVIAL-ENVIRONMENT DEPOSITS

GENERALIZATION OF MARINE DEPOSITIONAL ENVIRONMENT

OVERRIDING-DELTA DEPOSITIONAL ENVIRONMENT

UPPER
DELTA

LOWER
DELTA

SEE SECTION ON CORRECTION PROBLEMS FOR DISCUSSION OF WELLS.

SOURCE: WATER RESOURCE COMMISSION REPORT, NUMBER 155, 1989.

FIGURE 15

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	EMR NO.	PROJECT NO.
TITLE		
GENERAL SEPARATIONS AREA REGIONAL CROSS-SECTIONAL LINES		
DRAFTED BY TRACY W. MICKLEY	DATE	
GEOLOGY BY R. AADLAND M. K. HARRIS	DATE	
GEOLOGY BY T. WESTBROOK	DATE	
DRAWING NO. S6416116	DATE

FC 1

P 28

IDL 3

DRB 4 (8)

P 18 (3100')

F

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	ENR NO.	PROJECT NO.
TITLE		
ISOPACH MAP LOWER CARBONATE ZONE AQUIFER ZONE IIB ₁		
DRAFTED BY	DATE	
SHANNON F. VICK		
GEOLOGY BY	DATE	
AADLAND, HARRIS		
GEOLOGY BY	DATE	
AND WESTBROOK		
DRAWING NO.		
S6416109

PLANT TRUE
NORTH NORTH

NOTE:

1. CONTOUR INTERVAL = 25'

HPC 1

P 27
(8)

FEET

FC 1

P 28

IDL 3

DRB 4

FOURMILE BRANCH

P 18
(3180')

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	ENV. NO.	PROJECT NO.	
TITLE ISOPACH MAP UPPER CARBONATE ZONE AQUIFER ZONE IIB₁		DRAFTED BY TRACY W. MICKLEY	DATE
		GEOLOGY BY R. AADLAND M.K. HARRIS	DATE
		GEOLOGY BY T. WESTBROOK	DATE
		DRAWING NO. S6416112	DATE

NOTES:

CONTOUR INTERVAL = 5'

FIGURE 18

FC 1

P 28
(59)

30
IDL 3
(28)

40

50

60

DRB 4
(44)

The map shows the study area with sampling stations marked by dots and labeled with HSB numbers. The stations are distributed along the coast and in the river. The map includes a coastline, a river, and a grid of latitude and longitude coordinates.

FOURMILÉ BRANCH

FOURMILE BRANCH

P 18
• (3100')

UNIT ED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA

EWR NO.

PROJECT NO.

TITLE

STRUCTURE CONTOUR MAP
CONFINING SYSTEM
I - II

DRAFTED BY

SHANNON F. VICK

DATE

GEOLOGY BY

ADDLAND, HARRIS

DATE

GEOLOGY BY

AND WESTBROOK

DATE

DRAWING NO.

S6416107

PLANT TRUE
NORTH NORTH

LEGEND:

- TRACE OF OFFSET
D - DOWNTROW

NOTE:

1. CONTOUR INTERVAL = 10'
2. (52) ELEVATION OF PICK IN FEET ABOVE (MSL)

FIGURE 19

GEOPHYSICAL CHARACTERISTICS OF HYDROSTRATIGRAPHIC UNITS

UNITED STATES DEPARTMENT OF ENERGY	
SAVANNAH RIVER SITE	
DATE	BY
GEOPHYSICAL CHARACTERISTICS OF HYDROSTRATIGRAPHIC UNITS WELL P-27	1. TOP OF CONFINING SYSTEM I/II 2. AQUIFER UNIT IIA 3. CONFINING UNIT IIA - IIB 4. AQUIFER ZONE IIB ₁ 5. CONFINING ZONE IIB ₁ - IIB ₂ 6. AQUIFER ZONE IIB ₂

DUI 333' -0" +
 TD 899' -0"

FIGURE 20

VERTICAL SCALE
 0 20

FC 1

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	EMR NO.	PROJECT NO.
TITLE		
ISOPACH MAP AQUIFER UNIT IIA		
DRAFTED BY SHANNON F. VICK	DATE	
GEOLOGY BY ADDLAND, HARRIS	DATE	
GEOLOGY BY AND WESTBROOK	DATE	
DRAWING NO. S6416108

FIGURE 21

FC 1

P 28
(23)

IDL 3
(5)

10

20

FSB 108
FSB 109
FSB 99
FSB 87
FSB 98
FSB 105
FSB 97
FSB 96
FSB 95
FSB 94
FSB 78
FSB 103
FSB 110
FSB 79
FSB 106
FSB 101
FSB 100
FSB 88
FSB 89
FSB 111
FSB 92
FSB 91
FSB 107
FSB 90
FSB 77
FSB 114
FSB 113
FSB 104
FSB 102

DRB 4
(6)

BGO 13
BGO 26
BGO 25
BGO 14
BGO 27
BGO 28
BGO 30
BGO 31
BGO 32
BGO 33
BGO 34
BGO 15
BGO 16
BGO 17
BGO 21
BGO 22
BGO 23
BGO 24

FOURMILE BRANCH

HSB 138
HSB 71
HSB 117
HSB 119
HSB 116
HSB 118
HSB 70
HSB 86
HSB 137
HSB 114
HSB 129
HSB 136
HSB 115
HSB 113
HSB 112
HSB 111
HSB 110
HSB 109
HSB 108
HSB 107
HSB 106
HSB 105
HSB 104
HSB 103
HSB 102
HSB 101
HSB 100
HSB 99
HSB 98
HSB 97
HSB 96
HSB 95
HSB 94
HSB 93
HSB 92
HSB 91
HSB 90
HSB 89
HSB 88
HSB 87
HSB 86
HSB 85
HSB 84
HSB 83
HSB 82
HSB 81
HSB 80
HSB 79
HSB 78
HSB 77
HSB 76
HSB 75
HSB 74
HSB 73
HSB 72
HSB 71
HSB 70
HSB 69
HSB 68
HSB 67
HSB 66
HSB 65
HSB 64
HSB 63
HSB 62
HSB 61
HSB 60
HSB 59
HSB 58
HSB 57
HSB 56
HSB 55
HSB 54
HSB 53
HSB 52
HSB 51
HSB 50
HSB 49
HSB 48
HSB 47
HSB 46
HSB 45
HSB 44
HSB 43
HSB 42
HSB 41
HSB 40
HSB 39
HSB 38
HSB 37
HSB 36
HSB 35
HSB 34
HSB 33
HSB 32
HSB 31
HSB 30
HSB 29
HSB 28
HSB 27
HSB 26
HSB 25
HSB 24
HSB 23
HSB 22
HSB 21
HSB 20
HSB 19
HSB 18
HSB 17
HSB 16
HSB 15
HSB 14
HSB 13
HSB 12
HSB 11
HSB 10
HSB 9
HSB 8
HSB 7
HSB 6
HSB 5
HSB 4
HSB 3
HSB 2
HSB 1

P 18
(9)

UNIT ED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	EMR NO.	PROJECT NO.
TITLE		
ISOPACH MAP CONFINING UNIT IIA - IIB		DRAFTED BY TRACY W. MICKLEY
		GEOLOGY BY R. AADLAND M.K. HARRIS
		GEOLOGY BY T. WESTBROOK
		DRAWING NO. S6416106

NOTES:

CONTOUR INTERVAL = 10'

HPC 1
(17)P 27
(15)

FEET

0 500 1000

FC 1

140

P 28
(140)

130

IDL 3
(131)

120

110

FSB 108
FSB 76 (124)
FSB 109
FSB 101 (119)
FSB 100
FSB 88
FSB 89
FSB 85
FSB 91
FSB 107
FSB 90
FSB 80
FSB 86
FSB 87 (115)
FSB 99 (115)
FSB 105 (115)
FSB 98
FSB 97 (110)
FSB 96 (109)
FSB 95
FSB 94
FSB 78 (107)
FSB 103
FSB 104
FSB 79 (106)
FSB 110
FSB 106
FSB 102

F-2
F-3
U-111
D

BGO 13
BGO 14 (136)
BGO 25 (132)
BGO 26 (132)
BGO 27
BGO 28 (124)
BGO 29
BGO 30
BGO 31
BGO 32
BGO 33
BGO 34

DRS 4
(117)

FOURMILE BRANCH

HSB 119
HSB 114
HSB 116
HSB 117 (122)
HSB 71
HSB 118 (118)
HSB 70
HSB 98
HSB 137
HSB 114
HSB 129
HSB 136
HSB 130

FOURMILE BRANCH

P 18
(123)
(3100')

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	EMR NO.	PROJECT NO.
TITLE		
STRUCTURE CONTOUR MAP CONFINING UNIT IIA - IIB		
DRAFTED BY	SHANNON F. VICK	DATE
GEOLOGY BY	AADLAND, HARRIS	DATE
GEOLOGY BY	AND WESTBROOK	DATE
DRAWING NO.	S6416102

FIGURE 23

FC 1

P 28
(62)

IDL 3

DRB 4

P 18
(3100')
(46)

SAVANNAH RIVER SITE

PROJECT NO.

ISOPACH MAP
AQUIFER ZONE
IIB₁

TASTE

DATE _____

T. WESTBROOK

DATE _____

DRAWING NO.

NO.
S6416105

(62) = PLOT VALUE IN FEET

HPC 1
(78)

P 27
(62)

A horizontal scale bar labeled "FEET" in bold capital letters. The bar has three major tick marks labeled "0", "500", and "1000". There are four smaller tick marks between each major tick mark, dividing each 500-foot segment into five 100-foot segments.

FIGURE 24

GEOPHYSICAL CHARACTERISTICS OF LIMESTONE IN HSB 122

FIGURE 25

FC 1

10

P 28

P 18
(3100')

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA

ENV. NO.

PROJECT NO.

TITLE

ISOPACH MAP
CONFINING ZONE
IIB₁ - IIB₂

DRAFTED BY

TRACY W. MICKLEY

DATE

GEOLOGY BY

R. AADLAND
M.K. HARRIS

DATE

GEOLOGY BY

T. WESTBROOK

DATE

DRAWING NO.

S6416104

LEGEND:

- CONFINING ZONE
NOT PRESENT

NOTES:

CONTOUR INTERVAL = 10'

HPC 1

P 27

FC 1

220

P 28

P 18
(3100')

FIGURE 27

FC 1

P 28

IDL 3

P 18
(3100')

UNIT ED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	ENR NO.	PROJECT NO.	
TITLE ISOPACH MAP AQUIFER ZONE IIB₂ (SATURATED WATER TABLE ZONE)		DRAFTED BY TRACY W. MICKLEY	DATE
		GEOLOGY BY R. AADLAND M. K. HARRIS	DATE
		GEOLOGY BY T. M. WESTBROOK	DATE
		DRAWING NO. S6416103

NOTES:
CONTOUR INTERVAL = 10'

FIGURE 29

P 28

IDL 3

20:

P 18
6 (3100')

210

PLANT TRUE
NORTH NORTH

UNIT ED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	ENR NO.	PROJECT NO.
TITLE		
POTENTIOMETRIC SURFACE OF THE AQUIFER ZONE IIB ₁		DRAFTED BY T MICKLEY/S VICK
		GEOLOGY BY R. AADLAND M.K. HARRIS
		GEOLOGY BY T. WESTBROOK
		DRAWING NO. S6416114
		DATE

NOTES:

1. CONTOUR INTERVAL = 5'
2. DATA FROM THE FOURTH QUARTER 1989

HPC 1

P 27 •
(240)

FIGURE 30

FC 1
(142.5)

P 28
(146.4)

IDL 3

145

150

112
155

160

165

170

P 18
(3188')
(187.7)

FSB 108
FSB 99
FSB 109
FSB 76 (150.7)
FSB 101
FSB 88 (150.8)
FSB 7A
FSB 111
FSB 90
FSB 107
FSB 91
FSB 89
FSB 92
FSB 77
FSB 93
FSB 94
FSB 95
FSB 96
FSB 97
FSB 98
FSB 100
FSB 103
FSB 104
FSB 106
FSB 110
FSB 113
FSB 78
FSB 79 (156.6)
FSB 102

BGO 29
(158.5)

BGO 26
(159.1)

BGO 25
(158.8)

BGO 14
(157.3)

BGO 15
(159.6)

BGO 16
(159.7)

DRB 4

HSB 138
HSB 71
HSB 117
(164.6)
HSB 118
(168)
HSB 70
(166.8)
HSB 137
HSB 114
HSB 129
HSB 136
HSB 130

(165.7) HSB 119
HSB 115
HSB 116
HSB 117
HSB 118
HSB 119
HSB 120
HSB 121
HSB 122
HSB 123
HSB 124
HSB 125
HSB 126
HSB 127
HSB 128
HSB 129
HSB 130
HSB 131
HSB 132
HSB 133
HSB 134
HSB 135
HSB 136
HSB 137
HSB 138
HSB 139
HSB 140
HSB 141
HSB 142
HSB 143
HSB 144
HSB 145
HSB 146
HSB 147
HSB 148
HSB 149
HSB 150
HSB 151
HSB 152
HSB 153
HSB 154
HSB 155
HSB 156
HSB 157
HSB 158
HSB 159
HSB 160
HSB 161
HSB 162
HSB 163
HSB 164
HSB 165
HSB 166
HSB 167
HSB 168
HSB 169
HSB 170
HSB 171
HSB 172
HSB 173
HSB 174
HSB 175
HSB 176
HSB 177
HSB 178
HSB 179
HSB 180
HSB 181
HSB 182
HSB 183
HSB 184
HSB 185
HSB 186
HSB 187
HSB 188
HSB 189
HSB 190
HSB 191
HSB 192
HSB 193
HSB 194
HSB 195
HSB 196
HSB 197
HSB 198
HSB 199
HSB 200

(164)

(176)

FOURMILE BRANCH

FOURMILE BRANCH

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA

EWR NO.

PROJECT NO.

TITLE

POTENTIOMETRIC SURFACE
AQUIFER UNIT
IIA

DRAFTED BY

TRACY W. MICKLEY

DATE

GEOLOGY BY

R. AADLAND
M.K. HARRIS

DATE

GEOLOGY BY

T. WESTBROOK

DATE

DRAWING NO.

S6416115

DATE

NOTES:

CONTOUR INTERVAL = 5'

PLANT TRUE
NORTH NORTH

FIGURE 31

FC 1

P 28

DRB 4

FOURMILE BRANCH

P 18 (3188')

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	ENR NO.	PROJECT NO.
TITLE		
HEAD DIFFERENCE BETWEEN AQUIFERS IIB ₁ AND IIB ₂		
DRAFTED BY	TRACY W. MICKLEY	DATE
GEOLOGY BY	R. AADLAND M.K. HARRIS	DATE
GEOLOGY BY	T. WESTBROOK	DATE
DRAWING NO.		S6416110

NOTE:
CONTOUR INTERVAL - 5

LEGEND:
- CONFINING ZONE NOT PRESENT

FC 1
(66.2)

65

60

P 28
(68.3)

IDL 3

65

60

55

50

45

40

DRB 4

FOURMILE BRANCH

HSB 138

HSB 119

HSB 117
(57.1)

HSB 118

HSB 78

HSB 137

HSB 129

HSB 136

HSB 111

HSB 84
(40.3)

HSB 130

FOURMILE BRANCH

P 18
(3100')

BGO 13

BGO 12
(39)

BGO 14
(64.4)

BGO 26

BGO 25

BGO 15

BGO 18

BGO 17

BGO 27

BGO 28

BGO 30

BGO 31

BGO 32

BGO 33

BGO 34

FSB 108

FSB 76

FSB 101

FSB 88

FSB TA

FSB 111

FSB 69

FSB 107

FSB 90

FSB 109

FSB 99

FSB 87
(56.8)

FSB 98

FSB 105

FSB 97
(55.1)

FSB 96

FSB 95

FSB 94

FSB 93

FSB 78
(53.0)

FSB 103

FSB 110

FSB 106

FSB 113

FSB 79
(39.8)

FSB 102

FSB 104

FSB 100

FSB 101

FSB 102

FSB 103

FSB 104

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	ENR NO.	PROJECT NO.
TITLE		DRAFTED BY
HEAD DIFFERENCE BETWEEN AQUIFERS IIA - IIB ₁		TRACY W. MICKLEY
		DATE
		GEOLGY BY
		R. AADLAND
		M.K. HARRIS
		DATE
		GEOLGY BY
		T. WESTBROOK
		DATE
		DRAWING NO.
		S6416111

NOTES:

CONTOUR INTERVAL = 5'
 (61.4) = HEAD DIFFERENCE (IN FEET)
 *WELLS BGO-12 & BGO-10 LEVELS
 NOT USED IN CONTOURING DUE TO
 SUSPECT WELL CONSTRUCTION

FIGURE 33

GENERAL SEPARATIONS

C

AREA REGIONAL HYDROSTRATIGRAPHIC CROSS-SECTION 1

2770' -0"

405' -0"

THE SCALE IN THIS
AREA IS EXAGGERATED
FOR CLARITY

HC895

HC12

ELEV 292' -0"

ELEV 288' -0"

10-135' -0"

?

A

?

5230' -0"

P27

ELEV 274' -0"

300

290

280

270

260

250

240

230

220

210

200

190

180

170

160

150

140

130

120

110

100

90

80

70

60

50

40

D

C

B

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY
- SAND PACK INTERVAL
- SCREENED INTERVAL

DUI = DEPTH USED FOR INTERPRETATION
 TD = TOTAL DEPTH
 GSA = GENERAL SEPARATIONS AREA
 NR = NO RECOVERY
 GEOPHYSICAL CURVE = NATURAL GAMMA
 — = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

SCALE

HORIZONTAL 0 200 400

VERTICAL 0 5 10 15 20

PLATE 1

DOC. NO. 800002

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	GSA	SRS NO.	PROJECT NO.	F & H RCRA PERMIT
TITLE	GENERAL SEPARATIONS HYDROSTRATIGRAPHIC CROSS - SECTION #1			
PREPARED BY	W. A. BOLT			
REVIEW BY	J. A. LARLAND, M. K. HARRIS			
DATE	JAN 1981			
DATE	JAN 1981			
DATE	JAN 1981			
DATE	JAN 1981			

TD 899' - 0"

G

GENERAL SEPARATIONS AREA REGIONAL H

CROSS-SECTION # 2

HYDROSTRATIGRAPHIC

2

2430' -0"

480' -0"

750' -0"

THE SCALE IN THIS AREA
IS EXAGGERATED FOR CLARITY

THE SCALE IN THIS
IS EXAGGERATED FOR C

HSB65
ELEV 278' -0"

HSB123
ELEV 262' -0"

TD 183' -0"

50'-0" 600'-0" 3750'-0"

SCALE IN THIS AREA
GENERATED FOR CLARITY

HS8TB
ELEV 267'-0"

HS8133
ELEV 253'-0"

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY

- SAND PACK INTERVAL
- SCREENED INTERVAL

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

DUI = DEPTH USED FOR INTERPRETATION

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

SCALE
HORIZONTAL 0 200 400
VERTICAL 0 10 20

AQI

140

130

120

110

100

90

80

70

60

50

40

30

20

10

0

-10

-20

-30

-40

-50

-60

CONFINING UNIT IIA - IIB

AQUIFER
UNIT
IIA

TOP OF CONFINING
SYSTEM I/II

TD 300' -0"

DUI-273'-0"±
TD 558'-8"

DWI 386' -8"
TD 1938' -8"

TO 98'-0"

?

OUI 277' -0"+
TO 730' -0"

PLATE 2

FIG. 15
980004

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

GSA

DRY NO.

PROJECT NO.
F & H RCRA PERMIT

TITLE

GENERAL SEPARATIONS
HYDROSTRATIGRAPHIC
CROSS-SECTION #2

DESIGNED BY
TED A. BOLT
DRAWN BY
R. ADLAND, M. HARRIS
CHECKED BY
CH. LEVY, TF. OHLERMAN
IN CHARGE
J. H. WESTBROOK

DATE	REVISION	BY
6-24-81	REVISED FAULT LINE AND AFFECTED AREA	9FV

GENERAL SEPARATIONS A

CH

6550' - 8'

IS AREA REGIONAL HYDROSTRATIGRAPHIC CROSS-SECTION # 3

5100' - 0'

P2
ELEV 21

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY
- SAND PACK INTERVAL
- SCREENED INTERVAL

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

DUI = DEPTH USED FOR INTERPRETATION

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

SRS

150
140
130
120
110
100
90
80
70
60
50
40
30
20
10
0
-10
-20
-30
-40

AQUIFER
ZONE
IIB,

CONFINING UNIT IIA-IIB

AQUIFER
UNIT
IIA

TOP OF CONFINING
SYSTEM I/II

DUI 330'-0"
TD 952'-0"

TO 2

DUI 325' -0" +
TO 428' -0"

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

SCALE

HORIZONTAL 0 200 400

VERTICAL 0 5 10 15 20

SRS

PLATE 3

D.K. NO. 000005

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

AREA	GSA	ENV. NO.	F & H RCRA PERMIT
TITLE	GENERAL SEPARATIONS HYDROSTRATIGRAPHIC CROSS SECTION #3		
DESIGNED BY	R. BOLT / S. VICK	DATE	
DRAWN BY	R. AARLAND, M. HARRIS	DATE	
CHECKED BY	M. LEWIS, T. GAUGHAN	DATE	
IN CHARGE	R. WESTBROOK	DATE

GENERAL SEPARATIONS AREA REGIONAL F

CROSS-SECTION # 4

LEGEND:

TD = TOTAL DEPTH

OSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

PLANT NORTH
TRUE NORTH

DIATE 4

000000

TD 90' -0"

110
100
90
80
70
60
50
40
30
20
10
0

CONFINING UNIT IIA-IIB

AQUIFER
UNIT
IIA

TOP OF CONFINING
SYSTEM I/II

?

?

?

?

?

PLATE 4

UNITED STATES DEPARTMENT OF ENERGY	
SAVANNAH RIVER SITE	
GSA	F & H RCRA PERMIT
GENERAL SEPARATIONS HYDROSTRATIGRAPHIC CROSS SECTION #4	
REVIEWED BY	DATE
ROD A. BOLT	
REVIEWED BY	DATE
R. AASLAND, MK. HARRIS	
CM. LEVITS, TY. GAUGHAN	
TM. WESTBROOK	

GENER

GENERAL SEPARATIONS AREA REGIONAL

CROSS-SECTION # 5A

ONAL HYDROSTRATIGRAPHIC

5A

490' -0"

1920' -0"

480' -0"

SCALE IS EXAGGERATED
IN THIS AREA
FOR CLARITY

SCALE IS EXAGGERATED
IN THIS AREA
FOR CLARITY

BG05
ELEV 294' -0"

HPT2
ELEV 258' -0"

HPT1
ELEV 233' -0"

TD 128' -0"

TD 215' -0"

TD 185' -0"

290

280

270

260

250

240

230

220

210

200

190

180

170

160

150

140

130

120

110

100

90

80

70

60

50

40

30

TD
GSA
NR
GEO
DUI

PLANT
NORTH

GRAPHIC

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY
- SAND PACK INTERVAL
- SCREENED INTERVAL

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES.

GEOPHYSICAL CURVE = NATURAL GAMMA

DUI = DEPTH USED FOR INTERPRETATION

SCALE

HORIZONTAL 0 200 400

VERTICAL 0 5 10 15 20

PLATE 5A

ME	REVISION	NP0
8-24-91	REVISED FAULT LINES AND AFFECTED AREA	SFV

UNITED STATES DEPARTMENT OF THE INTERIOR		
SAVANNAH RIVER		
AREA	SECTION	PROJECT
GSA		F & E
TITLE		
GENERAL SEPARATIONS HYDROSTRATIGRAPHIC CROSS - SECTION #5/

PLATE 5A

SS0007

UNITED STATES DEPARTMENT OF ENERGY		
SAVANNAH RIVER SITE		
AREA	PROJECT NO.	PROJECT NO.
GSA	F & H RCRA PERMIT	
TITLE	REVIEWED BY	DATE
GENERAL SEPARATIONS HYDROSTRATIGRAPHIC CROSS - SECTION #5A	ROD A. BOLT	DATE
	R. AADLAND, MK. HARRIS	DATE
	CH. LEWIS, TF. GAUGHAN	DATE
	TM. WESTBROOK	DATE
REVISION	DATE	DATE
6-24-91	REVISED FAULT LINES AND AFFECTED AREA	SFV

GENERAL

AL SEPARATIONS AREA REGIONAL HYDROST

CROSS - SECTION # 5B

PROSTRATIGRAPHIC

LEGEND:

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES.

GEOPHYSICAL CURVE = NATURAL GAMMA

DUI = DEPTH USED FOR INTERPRETATION

TD 198'-0"

TD 198'-0"

TD 196'-0"

DUI 273'-0"±
TD 558'-0"

PLANT
NORTH
N

TD 188'-0"

TD 196'-0"

TD 212'-0"

MATCH LINE

SECT5B

SECT5A

DATE	REVISION	APP
8-24-91	REVISED FAULT LINE AND AFFECTED AREA	GFV

PI

TITLE

GE

H

CR

PLATE 5B

DATE		REVISION		PROJECT NO.	
8-24-91		REVISED FAULT LINE AND AFFECTED AREA		880000	
DATE		REVISION		PROJECT NO.	
				880000	
DATE		REVISION		PROJECT NO.	
				880000	
DATE		REVISION		PROJECT NO.	
				880000	
DATE		REVISION		PROJECT NO.	
				880000	

UNITED STATES DEPARTMENT OF ENERGY			
SAVANNAH RIVER SITE			
AREA	GSA	DATE	PROJECT NO.
TITLE	F & H RCRA PERMIT		
GENERAL SEPARATIONS			
HYDROSTRATIGRAPHIC			
CROSS - SECTION #5B			
PREPARED BY	ROD A. BOLT	DATE	
REVIEWED BY	R. AADLAND, MK. HARRIS	DATE	
CH. LEVIS, TF. CAIKHAN		DATE	
TM. WESTBROOK		DATE	

300
290
280
270
260
250
240
230
220
210
200
190
180
170
160
150
140
130
120
110
100
90
80
70
60
50
40
30

AQUIFER SYSTEM II

AQUIFER
UNIT
IIB

AQUIFER
ZONE
IIB₂

CONFINING
ZONE
IIB₁ - IIB₂

AQUIFER
ZONE
IIB₁

CONFINING
UNIT
IIA - IIB

AQUIFER
UNIT
IIA

P28
ELEV 285' -0"

GENERAL SEPARA

2950' -0"

600' -0"

683' -4"

THE SCALE IN THIS
AREA IS EXAGGERATED FOR CLARITY

THE SCALE IN THIS
AREA IS EXAGGERATED FOR CLARITY

BG010

ELEV 299' -0"

BG08

ELEV 281' -0"

BG06

ELEV 284' -0"

* well currently
on rehab/replacement
list - water levels
are not included in contouring.

CROSS-SECTION # 1

OSTRATIGRAPHIC

5230' -0"

0"

- GROUND WATER DIVIDE -

ELEV

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY
- SAND PACK INTERVAL
- SCREENED INTERVAL

DUI = DEPTH USED FOR INTERPRETATION

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

SCALE

HORIZONTAL 0 200 400

VERTICAL 0 5 10 15 20

120
110
100
90
80
70
60
50
40
30
20
10
0
-10
-20
-30
-40
-50
-60

AQUIFER
UNIT
IIA

TOP OF CONFINING
SYSTEM I/II

DUI 345' -0" ±

TD 428' -0"

?

* well currently
on rehab/replacement
list - water levels
are not included in contouring.

QUI 333' -0"±
TD 899' -0"

— CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

PLATE 6

SRN 980671

UNITED STATES DEPARTMENT OF ENERGY			
SAVANNAH RIVER SITE			
AREA	ORG NO.	PROJECT NO.	RCRA PERMIT
GSA		F & H	
TITLE		PREPARED BY	DATE
GENERAL SEPERATIONS HYDROSTRATIGRAPHIC CROSS-SECTION #1 WITH LINES OF EQUIPOTENTIAL FOR FOURTH QUARTER 1989		ROD A, BOLT	DATE
		REVIEW BY H. AARLAND, MK. HARRIS	DATE
		CM LEWIS, TF GAUGHAN	DATE
		TM WESTBROOK	DATE

GENERAL SEPARATIONS AREA REGIONAL CROSS-SECTION

ONAL HYDROSTRATIGRAPHIC

! # 2

750' -0'

600' -0'

3750' -0'

THE SCALE IN THIS AREA
IS EXAGGERATED FOR CLARITY

HSB7B
ELEV 267' -0"

HSB133
ELEV 253' -0"

235

▽ 240

245

250

234.0

229.3

TO 99' -0"

230

169.8

225

170

175

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY

- SAND PACK INTERVAL
- SCREENED INTERVAL

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

DUI = DEPTH USED FOR INTERPRETATION

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

130

CONFINING UNIT IIA - IIB

120

110

100

90

80

70

60

50

40

30

20

10

0

-10

-20

-30

-40

-50

-60

AQUIFER
UNIT
IIATOP OF CONFINING
SYSTEM I/II

TD 300' -0"

223.4

B

169.8

A

170

TD 183'-0"

169.8

NR

A

TD 210'-0"

?

?

DUI
TD

HORIZONTAL SCALE AND
DISTORTION IMPOSED
ON THE
PRESENTED AS A
QUANTITATIVE
GENERALIZED
RNS

DATE	REVISION	APP'D
6-24-81	REVISED FAULT LINE AND AFFECTED AREA	SFV

PLATE 7

UNITED STATES DEPARTMENT OF ENERGY			
SAVANNAH RIVER SITE			
AREA	GSA	PROJECT NO.	F & H RCRA PERMIT
TITLE		PREPARED BY	DATE
GENERAL SEPERATIONS HYDROSTRATIGRAPHIC CROSS-SECTION #2 WITH LINES OF EQUIPOTENTIAL FOR FOURTH QUARTER 1989		ROD A. BOLT	DATE
		REVIEWED BY	DATE
		R. AAZLAND, MK. HARRIS	DATE
		CM LEWIS, TF GAUGHAN	DATE
		TM WESTBROOK	DATE

GENERAL SEPARATIONS A CI

6550' -0"

ONS AREA REGIONAL HYDROSTRATIGRAPHIC CROSS-SECTION # 3

APHIC

8"

5100' - 8"

GENERATED IN
A FOR
ITY

FS876

ELEV 291' - 6"

215
(est.)

211.1

215

210

205

150.7

8

LEGEND:

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

DUI = DEPTH USED FOR INTERPRETATION

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

SRS

150.7

B

154.3

155

-0"

DUE TO VARIANCES IN HORIZONTAL SCALE AND
WITH RESPECT TO THE DISTORTION IMPOSED
BY THE VERTICAL EXAGGERATION THE
EQUIPOTENTIAL LINES ARE PRESENTED AS A
QUALITATIVE RATHER THAN QUANTITATIVE
REPRESENTATION OF THE GENERALIZED
GROUND WATER FLOW PATTERNS

OUT
TO

PLATE 8

SWR NO. 888873

UNITED STATES DEPARTMENT OF ENERGY			
SAVANNAH RIVER SITE			
AGE	GSA	SWR NO.	F & H RCRA PERMIT
TITLE		PREPARED BY	DATE
GENERAL SEPERATIONS HYDROSTRATIGRAPHIC CROSS-SECTION #3 WITH LINES OF EQUIPOTENTIAL FOR FOURTH QUARTER 1989		ROD A. BOLT	DATE
		REVIEW BY	DATE
		R AADLAND, MC HARRIS	DATE
		CM LEWIS, TF GAUGHAN	DATE
		TH WESTERBOOK	DATE

GENERAL SEPARATIONS AREA REGIONAL HYDROLOGIC CROSS-SECTION # 4

HYDROSTRATIGRAPHIC

LEGEND:

TD = TOTAL DEPTH

GSA = GENERAL SEPARATIONS AREA

NR = NO RECOVERY

GEOPHYSICAL CURVE = NATURAL GAMMA

— = CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES

PLANT NORTH
TRUE NORTH

160
150
140
130
120
110
100
90
80
70
60
50
40
30
20
10
0

ZONE
IIB₁

CONFINING UNIT IIA-IIB

AQUIFER
UNIT
IIA

TOP OF CONFINING
SYSTEM I/II

DUE TO VARIANCES IN HORIZONTAL
WITH RESPECT TO THE DISTOR
BY THE VERTICAL EXAGGERATIO
EQUIPOTENTIAL LINES ARE PRE
QUALITATIVE RATHER THAN QU
REPRESENTATION OF THE GENE
GROUND WATER FLOW PATTERNS

PLATE 9		JOB NO. 890874	
UNITED STATES DEPARTMENT OF ENERGY			
SAVANNAH RIVER SITE			
CDR GSA	DRG NO.	PROJECT NO. F & H RCRA PERMIT	
TITLE		PREPARED BY	DATE
GENERAL SEPERATIONS HYDROSTRATIGRAPHIC CROSS-SECTION #4 WITH LINES OF EQUIPOTENTIAL FOR FOURTH QUARTER 1989		ROD A. BOLT	
		REVIEW BY	DATE
		R AADLAND, YK HARRIS	
		CM LEVITS, TJ GAUGHAN	DATE
		TH WESTBROOK	DATE

ERAL SEPARATIONS AREA REGIONAL HYDROS

CROSS-SECTION # 5A

[] DATA IN BRACKETS
NOT USED IN
INTERPRETATION
(WELLS SCHEDULED FOR
REPLACEMENT)

YDROSTRATIGRAPHIC

APHIC

LEGEND:

- CALCAREOUS SAND
- CALCAREOUS MUD
- CLAYEY SAND
- SAND
- GRAVEL/PEBBLES
- SILT/CLAY
- SANDY LIMESTONE
- LIMESTONE
- SANDY MUDDY LIMESTONE
- MUDDY LIMESTONE
- SANDY CLAY
- SAND PACK INTERVAL
- SCREENED INTERVAL

TD = TOTAL DEPTH
GSA = GENERAL SEPARATIONS AREA
NR = NO RECOVERY
--- CONFINING ZONE, UNIT, AND SYSTEM BOUNDARIES.
GEOPHYSICAL CURVE = NATURAL GAMMA
DUI = DEPTH USED FOR INTERPRETATION

[] DATA IN BRACKETS
NOT USED IN
INTERPRETATION
(WELLS SCHEDULED FOR
REPLACEMENT)

01E
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

PLANT NORTH
TRUE NORTH

SECT 5B SECT 5A

80000
45000
65000
45000

FSBTA
FSB101A-BG01A
FSB100A-BG025
FSB99A-BG026A
FSB98
FSB97
BG029A
BG027A

SE

PLATE 10A

UNITED STATES DEPARTMENT

SAVANNAH RIVER

DUE TO VARIANCES IN HORIZONTAL SCALE AND WITH RESPECT TO THE DISTORTION IMPOSED BY THE VERTICAL EXAGGERATION THE EQUIPOTENTIAL LINES ARE PRESENTED AS A QUALITATIVE RATHER THAN QUANTITATIVE REPRESENTATION OF THE GENERALIZED GROUND WATER FLOW PATTERNS

DATE	REVISION	APP'D
6-24-91	REVISED FAULT LINES AND AFFECTED AREA	SFV

AREA: GSA
SUB NO.:
PROJECT NO.: F & H
TITLE:
GENERAL SEPERATIONS
HYDROSTRATIGRAPHIC
CROSS-SECTION #5A WITH
LINES OF EQUIPOTENTIAL
FOR FOURTH QUARTER 1989

PLATE 10A

DWG NO. 888875

VERTICAL SCALE AND
HORIZONTAL SCALE
IMPOSED
ON THE
SENTED AS A
INITIATIVE
CALIZED

DATE	REVISION	BY
6-24-91	REVISED FAULT LINES AND AFFECTED AREA	SFV

UNITED STATES DEPARTMENT OF ENERGY			
SAVANNAH RIVER SITE			
AREA	DWG NO.	PROJECT NO.	
GSA		F & H RCRA PERMIT	
TITLE		PREPARED BY	DATE
GENERAL SEPERATIONS HYDROSTRATIGRAPHIC CROSS-SECTION #5A WITH LINES OF EQUIPOTENTIAL FOR FOURTH QUARTER 1989		ROD A. BOLT	
		REVIEWED BY	DATE
		R AADLAND, MK HARRIS	
		DATE	
		CM LEWIS, TF GAUGHAN	DATE
		DATE	
		TH WESTBROOK

ERAL SEPARATIONS AREA REGIONAL HYDROS

CROSS - SECTION # 5B

OSTRATIGRAPHIC

49.7

150.6

150.8

155

150

TD 198'-0"

TD 198'-0"

TD 19

CULAR
IE PAGE

DUI 273'-0"±
TD 558'-0"

DUE TO VARIANCES IN HORIZONTAL SCALE AND WITH RESPECT TO THE DISTORTION IMPOSED BY THE VERTICAL EXAGGERATION THE EQUIPOTENTIAL LINES ARE PRESENTED AS A QUALITATIVE RATHER THAN QUANTITATIVE REPRESENTATION OF THE GENERALIZED GROUND WATER FLOW PATTERNS

ME	
6-24-81	REVISED FAULT LIT

159.1

TO 196'-0"

MATCH LINE

SECT5B SECT5A

PLANT NORTH
TRUE NORTH

SECT5B SECT5A

MATCH LINE

MATCH LINE

SECT5B SECT5A

FSBTA
FSB101A
FSB100A
FSB99A
FSB98
FSB97
BG012
BG010
BG08
BG06
BG05C
HPT2A
HPT1A
BG025
BG014
BG029A
BG027A
BG026A

PLATE 10B

SS0076

UNITED STATES DEPARTMENT OF ENERGY

SAVANNAH RIVER SITE

DATE	REVISION	BY
6-24-91	REVISED FAULT LINE AND AFFECTED AREA	SFV

AREA	GSA	DATE		PROJECT NO.	F & H RCRA PERMIT
TITLE	GENERAL SEPERATIONS HYDROSTRATIGRAPHIC CROSS-SECTION #58 WITH LINES OF EQUIPOTENTIAL FOR FOURTH QUARTER 1989				
PREPARED BY	R.A. BOLT/S.F. VICK	DATE			
REVIEWED BY	R. AADLAND, MK HARRIS	DATE			
CH. LEVITS, TF GAUGHAN		DATE			
TH WESTBROOK		DATE			

AL SCALE AND
ON IMPOSED
THE
ENTED AS A
TITATIVE
LIZED

END

**DATE
FILMED**

5 / 18 / 93