DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

STUDIES OF YRAST AND CONTINUUM

STATES IN A - 100-200 NUCLEI

Progress Report for 1992

Patrick J. Daly

Principal Investigator

Purdue University, West Lafayette, IN 47907

January, 1993

Prepared for

THE U.S. DEPARTMENT OF ENERGY

UNDER GRANT NO. DE-FG02-87ER40346

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Abstract

This report summarizes progress in nuclear structure research for the year 1992. Highlights include spectroscopic results for hard-to-reach neutron-rich and proton-rich products of heavy ion reactions.

I. Research Activities

This has been a active, productive year for our research group and good general progress has been made. The heart of our research program continues to be a lively and fruitful experimental collaboration with the Khoo-Janssens Argonne team in investigating superdeformation and related structural problems in high-spin physics, as well as the spectroscopy of otherwise hard-to-reach neutron excessive nuclei produced in deep inelastic heavy ion reactions. Most of our experiments are performed using heavy ion beams from the ATLAS accelerator and on-line γ -ray measurements with modern detector arrays. The bulk of our data analysis is performed on our relatively new VAX system at Purdue, which has proven to be an important practical step forward, boosting the effectiveness and morale of our group.

In the operation of the collaboration, running the experiments is a joint undertaking, but the individual research projects thereafter naturally subdivide into two distinct categories. The emphasis in this report is on those projects for which the leadership came from the Purdue contingent, notably, the spectroscopy of neutron-rich nuclei, the yields and mechanisms of deep inelastic heavy ion reactions and the first studies of μ s isomers using the Argonne FMA. The Argonne scientists provided direction and drive for the superdeformation and high-spin physics studies; this work is represented in the publications and talks lists, but only those particular studies where Purdue participants made outstanding contributions are detailed further in this report. Visitor Rafal Broda (Cracow) has again this year contributed substantially to several of our projects, and Jan Blomqvist (Stockholm) continues to be an invaluable source of theoretical advice and inspiration.

Spectroscopy of Neutron-Rich Products of Heavy Ion Collisions

In earlier work¹⁾ we found that $\gamma\gamma$ coincidence measurements could be used to identify inelastic and transfer reaction products from collisions of 255 MeV

^{60}Ni ions with ^{92}Mo . Typically, mutual excitation of light and heavy reaction products was observed, providing clearcut identification of the reaction involved. Population of moderately high-spin (up to $I=12$) non-collective states was seen, along with large energy transfer, which sometimes gave rise to neutron evaporation. It seemed to us that $\gamma\gamma$ techniques might have potential for examining at high resolution some aspects of inelastic/transfer processes, and for exploring the yrast spectroscopy of certain neutron excessive nuclei that are inaccessible by fusion-evaporation reactions.

Neutron-Rich sdf Shell Nuclei

Our first spectroscopic applications of these techniques were to neutron-rich sdf shell nuclei, produced in reactions of ^{34}S , ^{36}S and ^{37}Cl beams on ^{160}Gd targets just above the Coulomb barrier. High quality $\gamma\gamma$ data came to μs as byproducts of our ATLAS studies of superdeformed (SD) bands in the $A = 190$ region, and spectroscopic information about deep inelastic reaction products could be extracted by emphasizing lower multiplicity subsets of the coincidence events. Gates placed on known γ -rays in the $A \sim 160$ products selected individual reaction channels and identified coincident γ -rays in light product partners. Transfer of protons from projectile to target, with or without exchange of neutrons, was found to be generally favored, leading to excited neutron-rich light nuclei, some of them difficult to reach by other means. Notable results include the observation of yrast cascades up to the highest known yrast states in four $N = 20$ isotones ^{34}Si , ^{35}P , ^{36}S and ^{37}Cl . In two $N = 19$ isotones ^{33}Si and ^{34}P , and two $N = 22$ isotones ^{38}S and ^{39}Cl , previously unknown yrast states have been identified. These results test the predictions of up-to-date shell model calculations in a region where such calculations are thought to be rather accurate. A paper based on this work has been planned in detail and will soon be submitted for publication.

Neutron-Rich Tin Region

The most complete and fruitful data analyses have been performed for the systems $^{122,124}\text{Sn} + 325 \text{ MeV } ^{76}\text{Ge}$ and $^{122,124}\text{Sn} + 344 \text{ MeV } ^{80}\text{Se}$. These particular studies were undertaken to explore the yrast spectroscopy of heavy tin nuclei – not accessible by fusion-evaporation – and the in-beam and off-beam γ -ray data obtained were found to include new information about more than 40 nuclei around $A=124$ and $A=80$. These are products of deep inelastic reactions involving transfer²⁾ of many nucleons between target and projectile, and population of excited states with moderately high spins. The data analysis yielded many interesting things including:

- a) Long-lived $(\nu h_{11/2})^n$ seniority $\nu = 2$ 10^+ isomers in ^{122}Sn and ^{124}Sn . Their $B(E2)$ values pinpointed half-filling of the $\nu h_{11/2}$ subshell close to $N=73$, and provided³⁾ a revealing comparison with less complete results for $N=82$ isotones and $Z=82$ isotopes;
- b) Microsecond M2 isomers in ^{119}Sn , ^{121}Sn and ^{123}Sn , which have been characterized and interpreted⁴⁾. In ^{119}Sn and ^{121}Sn , higher-lying nanosecond E2 isomers of $(\nu h_{11/2})^n$ $\nu = 3$ character have also been identified – their decays amplify what we know about the $\nu h_{11/2}$ subshell filling.
- c) Yrast cascades have been identified in ^{121}Sb , ^{123}Sb , ^{125}Sb and in ^{115}In , ^{117}In and ^{119}In , nuclei that from the point of view of yrast spectroscopy are untouched up to now.
- d) Useful spectroscopic findings have also been made for neutron-rich products around $A=80$, including $^{72-74}\text{Zn}$, $^{76-78}\text{Ge}$ and $^{80-84}\text{Se}$.

Making isotopic assignments for previously unknown γ -ray cascades proved to be one of the biggest problems. The assignments for the Sb and In isotopes were based in part on some overlap with radioactivity data, and in part on observed γ - γ cross-coincidences between light and heavy reaction partners. However, the primary reaction products often have sufficient excitation energy for subsequent evaporation of neutrons, so the interpretation of the cross coincidences is not

always quite straightforward. The problems involved are similar to those encountered in fission product spectroscopy studies.

The reactions studied to date populated yrast states in tin region nuclei lying left of ^{124}Sn . In an upcoming ATLAS study using a ^{136}Xe beam we hope to be able to explore nuclei that are more neutron-rich by up to 4 neutrons (see below).

$^{130}\text{Te} + ^{64}\text{Ni}$ System

We were also able to examine thick target $\gamma\gamma$ data for the system $^{130}\text{Te} + 262$ MeV ^{64}Ni , which were originally taken as part of the ATLAS program of super-deformed band studies in Hg nuclei. Analysis of lower multiplicity subsets of the $\gamma\gamma$ data identified yrast cascades in such nuclei as ^{126}Te , ^{128}Te , ^{130}Te and the semi-magic ^{64}Ni and ^{66}Ni . Although the ^{64}Ni beam energy chosen for the SD band studies was not favorable for spectroscopy of deep inelastic products, the results of these analyses were most encouraging, and we plan follow-up experiments at a more favorable 275 MeV ^{64}Ni beam energy using the GA.SP detector system.

Population Patterns in Deep Inelastic Heavy Ion Reactions

For the reactions of $^{122,124}\text{Sn}$ targets with ^{76}Ge and ^{80}Se ions, 10-15% above the Coulomb barrier, we have performed extensive product yield determinations by in-beam, off-beam and radioactivity measurements. The results include the yields - on the same scale - of:

- 1) 10^+ and $19/2^+$ μs isomers in $^{118-124}\text{Sn}$
- 2) the ^{123}Sn and ^{125}Sn $3/2^+$ "ground states" - from radioactivity
- 3) eight high-spin β -decay isomers in $^{116}\text{In}-^{123}\text{In}$.
- 4) several fusion-evaporation products

One firm and striking result is that the 10^+ and $19/2^+$ isomers in the seven nuclei from ^{118}Sn to ^{124}Sn are populated with comparable cross sections from the ^{124}Sn target.

The overall distribution of yields with A and Z for the $^{124}\text{Sn} + ^{80}\text{Se}$ system is illustrated for both projectile-like and target-like even-even products in Fig. 1, which shows relative $\gamma\gamma$ coincidence intensities (between the $2^+ \rightarrow 0^+$ transition and all γ -rays feeding the 2^+ state directly in each case) for the different nuclei. Analysis of these distributions gave an estimate of the average number of neutrons evaporated from the excited reaction products, which was then used to extract the pattern of primary products. This pattern may be broadly understood in terms of N/Z equilibration in the dinuclear system formed in the H.I. collisions. Calculated minimum potential energy surfaces can be used to explain the gross features of the observed yield pattern as well as to predict the outcome of future experiments using other target-projectile combinations. For example, yrast states in tin nuclei with $A > 124$ were not detectably populated in the $^{124}\text{Sn} + ^{80}\text{Se}$ reaction, but considerations of N/Z equilibration indicate that ^{125}Sn , ^{126}Sn and ^{127}Sn should all be seen in an upcoming $^{124}\text{Sn} + ^{136}\text{Xe}$ study at ATLAS.

Yrast Isomers in ^{151}Yb Studied Using the Argonne FMA

Product recoils from the reaction $^{96}\text{Ru} + 255 \text{ MeV } ^{58}\text{Ni} \rightarrow ^{154}\text{Hf}^*$ were analyzed using the Argonne Fragment Mass Analyzer (FMA), and transported to a collector foil behind the focal plane, where decay of μs isomers could be studied under low background conditions. The recoils were dispersed in A/q as they traversed the position sensitive focal plane detector, and recoil- γ -ray coincidences could thus be used to assign masses for specific γ -ray cascades (Fig. 2). By use of these methods, yrast isomers with half-lives of $2.6(7) \mu\text{s}$ and $20(1) \mu\text{s}$ have been firmly assigned to the exotic $N=81$ nucleus ^{151}Yb , and the main features of the isomeric

Fig. 1: Relative yields determined for even-even products of the reaction $^{124}\text{Sn} + 344 \text{ MeV } ^{80}\text{Se}$. Target and projectile nuclei are labelled. The larger yields are well correlated with the minimum nuclear potential energy contours calculated for this dinuclear system.

decays have been established. A short paper describing this work has been submitted to Phys. Rev. C.

These results came from the first test experiment of this kind with the Argonne FMA, and much higher detector sensitivity should be achievable in future experiments. A particularly important improvement will be the installation of Si detectors around the catcher foil for measuring conversion electrons from stopped product nuclei.

Feeding of Superdeformed Bands

This work, guided by T. L. Khoo with major inputs from T. Lauritsen and Purdue post-doc P. Benet, explored how the compound nucleus formed by fusion-evaporation is sometimes trapped into a secondary minimum in the nuclear potential energy surface. Experimental measurements for the $^{160}\text{Gd}(^{36}\text{S},4n)^{192}\text{Hg}$ reaction included a) SD band intensity and its variation with spin; b) (I,E) entry distributions c) quasicontinuum spectra, and d) E_γ - E_γ correlation matrices. A model based on Monte Carlo simulations was developed to follow the history of γ -ray cascades associated with the feeding of SD bands. These simulations correctly described not only the SD entry distribution but also the SD band intensity and its variation with spin, with well defined ranges of the SD band energy and well depth. The experimental and model results suggest that the SD band lies 3.3-4.3 MeV above the normal yrast line when it decays around spin 10, and that the SD well depth is about 4 MeV around spin 40.

A Letter⁵⁾ based on this investigation was recently published.

Spectroscopy of A = 188-192 Hg Nuclei

For the past two years, our student Ian Bearden has been resident at Argonne under the day-to-day supervision of Robert Janssens. During this time, he has gained wide experience, participating in many different kinds of experiments but with particular emphasis on SD band investigations in Hg nuclei. He has played

Fig. 2: Mass spectra gated on a) all transmitted $A = 151$ and 152 ions b) known γ -rays in ^{151}Tm c) known γ -rays in ^{152}Yb d) the 203, 259 and 600 keV γ -rays. The conclusion is that these three γ -rays occur in an $A=151$ nucleus.

a notable role in the characterization of a ^{189}Hg superdeformed band⁶⁾, and in demonstrating that there is no SD band of appreciable intensity in ^{188}Hg .

For his Ph.D. thesis he has been extracting from the $\gamma\gamma$ data acquired in the SD band experiments information about "normal" states in these Hg nuclei. He has, of course, considerably extended the known level schemes and at this stage the experimental portion of his thesis work is more-or-less complete. He is now concentrating on interpretative aspects, particularly the cranked shell model approach.

References

1. R. Broda et al., Phys. Lett. B251, 245 (1990).
2. H. Freiesleben and J. V. Kratz, Phys. Rep. 106, 1 (1984).
3. R. Broda et al., Phys. Rev. Lett. 68, 1671 (1992).
4. R. H. Mayer et al., Z. Phys. A342, 247 (1992).
5. T. Lauritsen et al., Phys. Rev. Lett. 69, 1222 (1992).
6. I. G. Bearden et al., Z. Phys. A341, 491 (1992).

II. Personnel

Senior Staff

Patrick J. Daly, Professor of Chemistry, Principal Investigator
Zbigniew W. Grabowski, Professor of Physics
Rafal Broda, Visiting Professor
Bogdan Fornal, Research Associate

Graduate Students

Ian G. Bearden
Ralf H. Mayer
David Nisius
Laura Richter
Michele Sferrazza

1992 Publications and Submitted Papers

1. "Characterization of the Superdeformed Band in ^{189}Hg ", I. G. Bearden, R. V. F. Janssens, M. P. Carpenter, E. F. Moore, I. Ahmad, A. M. Baxter, Ph. Benet, P. J. Daly, M. W. Drigert, P. B. Fernandez, U. Garg, Z. W. Grabowski, T. L. Khoo, T. Lauritsen, W. Reviol, D. Ye, Z. Phys. A341, 491 (1992).
2. "New $19/2^+$ Isomers in ^{119}Sn , ^{121}Sn and ^{123}Sn ", R. H. Mayer, B. Fornal, R. Broda, I. G. Bearden, Z. W. Grabowski, S. Lunardi, P. J. Daly, T. Lauritsen, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, Y. Liang, Z. Phys. A342, 247 (1992).
3. "Yrast Isomers in Tin Nuclei from Heavy Ion Collisions and Filling of the $h_{11/2}$ Subshell", R. Broda, R. H. Mayer, I. G. Bearden, Ph. Benet, P. J. Daly, Z. W. Grabowski, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, T. Lauritsen, E. F. Moore, S. Lunardi, J. Blomqvist, Phys. Rev. Lett. 68, 1671 (1992).
4. "Shape Driving Effects in ^{193}Tl from the Spectroscopy of Yrast and Near-Yrast States", W. Reviol, M. P. Carpenter, U. Garg, R. V. F. Janssens, I. Ahmad, I. G. Bearden, Ph. Benet, P. J. Daly, M. W. Drigert, P. B. Fernandez, T. L. Khoo, E. F. Moore, S. Pilotte, D. Ye, Nucl. Phys. A548, 331 (1992).
5. "Isomeric Decay Studies Using a Recoil Mass Separator: Exotic N = 82 and 83 Nuclei ^{153}Yb , ^{153}Lu , ^{154}Lu and ^{154}Hf " J. H. McNeill, A. A. Chishti, P. J. Daly, W. Gelletly, M. A. C. Hotchkis, M. Piiparinen, B. J. Varley, P. J. Woods, J. Blomqvist, Z. Phys. A (in press), to appear Jan.-Feb. 1993.
6. "Feeding of Superdeformed Bands: the Mechanism and Constraints on Band Energies and the Well Depth", T. Lauritsen, Ph. Benet, T. L. Khoo, K. Beard, I. Ahmad, M. P. Carpenter, P. J. Daly, M. W. Drigert, U. Garg, P. B. Fernandez, R. V. F. Janssens, E. F. Moore, F. L. H. Wolfs and D. Ye, Phys. Rev. Lett. 69, 1222 (1992).
7. "Yrast Isomers in Exotic N=81 Nucleus ^{151}Yb Studied Using the Argonne Fragment Mass Analyzer", D. Nisius, B. Fornal, I. G. Bearden, R. Broda, R. H. Mayer, Z. W. Grabowski, P. J. Daly, C. N. Davids, I. Ahmad, B. B. Back, K. Bindra, M. P. Carpenter, W. Chung, D. Henderson, R. G. Henry, R. V. F. Janssens, T. L. Khoo, T. Lauritsen, Y. Liang, F. Soramel and A. V. Ramayya, submitted to Phys. Rev. C, November 1992.
8. "A Pocketful of Tin Isomers From Heavy Ion Collisions", P. J. Daly, R. Broda, B. Fornal, R. H. Mayer, D. Nisius, I. Bearden, Ph. Benet, Z. Grabowski, T. Lauritsen, M. Carpenter, R. V. F. Janssens, T. L. Khoo, Y. Liang, S. Lunardi and J. Blomqvist, in The Building Blocks of Nuclear Structure, ed. Aldo Covello (World Scientific, Singapore 1992).
9. "Collective Oblate Bands in ^{196}Pb ", J. R. Hughes, Y. Liang, A. Kuhnert, R. V. F. Janssens, I. Ahmad, I. G. Bearden, J. A. Becker, M. J. Brinkman, J. Burde, M. P. Carpenter, P. J. Daly, J. E. Draper, M. A. Deleplanque, R. M. Diamond, C. Duyar, B. Fornal, U. Garg, Z. W. Grabowski, E. A. Henry, R. G. Henry, W. Hesselink, N. Kalantar-Nayestanaki, W. H. Kelly, T. L. Khoo, T. Lauritsen, R. H. Mayer, D. Nisius, A. J. M. Plomper, T. R. B. Oliveira, W. Reviol, E. Rubel, F. Soramel, F. S. Stephens, M. A. Stoyer, D. Vo, and T. F. Wang, submitted to Phys. Rev. C as Rapid Communication, Dec. 1992.

1992 Conference Contributions and Other Talks

1. "Studies of Nuclear Structure with Heavy Ions", Graduate Research Seminar, Purdue Physics Department, January 1992 - Z. W. Grabowski and P. J. Daly.
2. "Population Patterns in $^{124}\text{Sn} + 325 \text{ MeV } ^{76}\text{Ge}$ Collisions" Heavy Ion Discussion Group, Argonne NL, January 1992 - B. Fornal.
3. "Yrast Isomers in $Z = 49, 50$ and 51 Nuclei from $^{124}\text{Sn} + 340 \text{ MeV } ^{80}\text{Se}$ Reactions", Nuclear Chemistry Seminar, Lawrence Livermore Laboratory, March 1992 - P. J. Daly.
4. "Spectroscopy of Neutron-Rich Nuclei following Deep Inelastic Collisions", Nuclear Structure Seminar, Lawrence Berkeley Laboratory, March 1992 - P. J. Daly.
5. "New $19/2^+$ Isomers in ^{119}Sn , ^{121}Sn and ^{123}Sn ", R. H. Mayer, B. Fornal, R. Broda, I. G. Bearden, Z. W. Grabowski, S. Lunardi, P. J. Daly, T. Lauritsen, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, Y. Liang, Washington DC Meeting, April 1992, abstract in Bull. Am. Phys. Soc., 37 (1992).
6. "Yrast Cascades in $^{121,123,125}\text{Sb}$ from Heavy Ion Reactions on $^{122,124}\text{Sn}$ ", D. Nisius, B. Fornal, R. H. Mayer, R. Broda, I. G. Bearden, Z. W. Grabowski, P. J. Daly, T. Lauritsen, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, Y. Liang, Washington DC Meeting, April 1992, abstract in Bull. Am. Phys. Soc., 37 (1992).
7. "Isotopic Yields for Heavy Ion Reactions on ^{124}Sn from γ -ray Measurements", B. Fornal, R. H. Mayer, R. Broda, I. G. Bearden, Z. W. Grabowski, P. J. Daly, T. Lauritsen, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, Y. Liang, Washington DC Meeting, April 1992, abstract in Bull. Am. Phys. Soc., 37 (1992).
8. "High Spin State Studies in the Nucleus ^{189}Hg ", I. G. Bearden, A. M. Baxter, R. V. F. Janssens, M. P. Carpenter, P. J. Daly, Ph. Benet, M. W. Drigert, P. B. Fernandez, U. Garg, Z. W. Grabowski, E. F. Moore, T. L. Khoo, W. Reviol, F. L. H. Wolfs, D. Ye, Washington DC Meeting, April 1992, abstract in Bull. Am. Phys. Soc. 37 (1992).
9. "The Feeding of Superdeformed Band in ^{192}Hg ", T. Lauritsen, Ph. Benet, T. L. Khoo, I. Ahmad, I. G. Bearden, M. P. Carpenter, P. J. Daly, M. W. Drigert, R. V. F. Janssens, Y. Liang, E. F. Moore, P. B. Fernandez, Washington DC Meeting, April 1992, abstract in Bull. Am. Phys. Soc. 37 (1992).
10. "A Pocketful of Tin Isomers from Heavy Ion Collisions", 4th International Spring Seminar on Nuclear Physics, Amalfi, Italy, May 1992 - P. J. Daly.
11. "Shape Coexistence at High Spin in ^{188}Hg ", I. G. Bearden, M. P. Carpenter, A. M. Baxter, R. V. F. Janssens, I. Ahmad, Ph. Benet, P. J. Daly, M. W. Drigert, P. B. Fernandez, B. Fornal, U. Garg, Z. W. Grabowski, T. L. Khoo, R. H. Mayer, E. F. Moore, W. Reviol, D. Ye, Int. Conf. on Nucl. Structure at High Angular Momentum, Ottawa, May 1992; AECL - 10613, p. 18 (1992).

12. "Shape Driving Effects in ^{193}Tl from the Spectroscopy of Yrast and Near-Yrast States", W. Reviol, M. P. Carpenter, U. Garg, R. V. F. Janssens, I. Ahmad, I. G. Bearden, Ph. Benet, P. J. Daly, M. W. Drigert, P. B. Fernandez, T. L. Khoo, E. F. Moore, S. Pilotte, D. Ye, Int. Conf. on Nucl. Structure at High Angular Momentum, Ottawa, May 1992; AECL - 10613, p. 24 (1992).
13. "Entrance Channel Dependence in the Population of the Superdeformed Bands in ^{191}Hg ", F. Soramel, T. L. Khoo, R. V. F. Janssens, I. Ahmad, M. P. Carpenter, T. Lauritsen, Y. Liang, B. Fornal, I. Bearden, Ph. Benet, P. J. Daly, Z. W. Grabowski, R. Mayer, D. Ye, U. Garg, W. Reviol, M. W. Drigert, Int. Conf. on Nucl. Structure at High Angular Momentum, Ottawa, May 1992; AECL - 10613, p. 52 (1992).
14. "Calculations of the Decay of Superdeformed Bands and Search for γ -Rays Connecting SD and Normal States", T. Lauritsen, T. L. Khoo, E. F. Moore, M. P. Carpenter, P. Fernandez, R. V. F. Janssens, Y. Liang, M. Freer, A. Wuosmaa, Ph. Benet, P. J. Daly, I. Bearden, P. J. Daly, B. Fornal, D. Ye, U. Garg, M. W. Drigert, Int. Conf. on Nucl. Structure at High Angular Momentum, Ottawa, May 1992; AECL - 10613, p. 53 (1992).
15. "Feeding of Superdeformed Band in ^{192}Hg : The Mechanism and Constraints on the SD Band Energies and Well Depth", Ph. Benet, T. Lauritsen, T. L. Khoo, I. Ahmad, K. Beard, I. G. Bearden, M. P. Carpenter, P. J. Daly, M. W. Drigert, P. B. Fernandez, U. Garg, R. V. F. Janssens, Y. Liang, E. F. Moore, W. Reviol, D. Ye, Int. Conf. on Nucl. Structure at High Angular Momentum, Ottawa, May 1992; AECL - 10613, p. 54 (1992).
16. "Higher Superdeformed Band Members in ^{190}Hg : Evidence for a Band Interaction?", I. G. Bearden, R. V. F. Janssens, M. P. Carpenter, I. Ahmad, P. J. Daly, M. W. Drigert, U. Garg, T. L. Khoo, T. Lauritsen, W. Reviol, R. Wyss, Int. Conf. on Nucl. Structure at High Angular Momentum, Ottawa, May 1992; AECL - 10613, p. 55 (1992).
17. "A Pocketful of Tin Isomers from Heavy Ion Collisions", Nuclear Seminar, Legnaro National Lab., Italy, May 1992 - P. J. Daly.
18. "Gamma-ray Spectroscopy of Multiple Nucleon Transfer Reactions in Tin Nuclei", 2nd Int. Symp. on Nuclear Excited States, Lodz, Poland, June 1992 - Z. W. Grabowski.
19. "Gamma-ray Spectroscopy of Massive Transfer Products from Heavy Ion Reactions", ACS Symp. on Nuclear Shapes, Washington DC, Aug. 1992 - B. Fornal.
20. "Zakopane School Enters Television Age: Conference Summary Talk", XXVII International School on Nuclear Physics, Zakopane, Poland, Sept. 1992 - P. J. Daly.
21. " γ -Ray Studies of Neutron Rich Nuclei Around A=80 Produced in Deep-Inelastic Collisions", B. Fornal, I. G. Bearden, R. Broda, R. H. Mayer, D. Nisius, Z. W. Grabowski, P. J. Daly, M. P. Carpenter, R. Henry, R. V. F. Janssens, T. L. Khoo, T. Lauritsen, Y. Liang, Santa Fe APS Meeting, Oct. 1992, abstract in Bull. Am. Phys. Soc. 37, 1279 (1992).

22. "New $27/2^-$ Isomers in ^{119}Sn and ^{121}Sn ", R. H. Mayer, B. Fornal, R. Broda, D. Nisius, I. G. Bearden, Z. W. Grabowski, P. J. Daly, T. Lauritsen, Y. Liang, R. Henry, M. P. Carpenter, R. Henry, R. V. F. Janssens, T. L. Khoo, Santa Fe APS Meeting, Oct. 1992, abstract in Bull. Am. Phys. Soc. 37, 1280 (1992).
23. "Evidence for a Band Interaction in the ^{190}Hg SD Band", I. G. Bearden, R. V. F. Janssens, M. P. Carpenter, I. Ahmad, P. J. Daly, M. W. Drigert, U. Garg, Z. W. Grabowski, T. L. Khoo, T. Lauritsen, W. Reviol, R. Wyss, Santa Fe APS Meeting, Oct. 1992, abstract in Bull. Am. Phys. Soc. 37, 1285 (1992).
24. "Identification of Yrast Isomers in ^{151}Yb : Results from the Argonne Fragment Mass Analyzer", D. Nisius, B. Fornal, I. G. Bearden, R. Broda, R. H. Mayer, Z. W. Grabowski, P. J. Daly, C. N. Davids, I. Ahmad, F. Soramel, M. P. Carpenter, R. Henry, R. V. F. Janssens, T. L. Khoo, T. Lauritsen, Y. Liang, K. Bindra, W. Chung, B. B. Back, D. Henderson, A. V. Ramayya, Santa Fe APS Meeting, Oct. 1992, abstract in Bull. Am. Phys. Soc. 37, 1307 (1992).
25. "Nuclear Structure Studies by Deep Inelastic Heavy Ion Reactions", Nuclear Physics Seminar, University of Milan, Italy, Nov. 1992 - P. J. Daly.
26. "FMA Study of Microsecond Isomers in ^{151}Yb ", Fragment Mass Analyzer Workshop, Argonne NL, Nov. 1992 - D. T. Nisius.
27. "Feeding Yrast Isomers in Heavy Ion Collisions: Fusion and Deep Inelastic Products" Nuclear Physics Seminar, University of Valencia, Spain, Dec. 1992 - P. J. Daly.

END

DATE
FILMED
5128193