Precision Measurement of the Direct CP Violation Parameter ϵ'/ϵ
via the four $K \rightarrow 2\pi$ decay modes And
A High Sensitivity Search for CP Violating Rare K_L Decays

Progress Report
for period May 1, 1991 - April 30, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Yau W. Wah
The University of Chicago
5801 S. Ellis Ave
Chicago, IL 60637

April 1992
Prepared for
The U.S. Department of Energy
Award no. DOE DE F602 290ER40560 Task J

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

PROGRESS OF THE CP GROUP DURING THE LAST GRANT YEAR

Much has happened in the past year. The progress has been in three areas: (A) results from the complete E731 data set which was collected in the 1987/88 fixed target run; preparations for and the taking of the data for (B) E773 (CPT symmetry test) and (C) E799 (rare decay study); and finally (D) R&D for a new detector to further study ϵ'/ϵ and rare decays.

(A) Results from the complete E731 data set :

Last summer we announced results from the entire data set. Earlier, we had published a result based upon 20% of the data set which was $\epsilon'/\epsilon = (-4 \pm 14(\text{stat.}) \pm 6(\text{syst.})) \times 10^{-4}$ (E731, 20% sample), in agreement with a class of super-weak models. Since the first publication, much better understanding of the acceptance, energy resolution, and of the backgrounds have allowed major reductions in the systematic error. This was accomplished by detailed studies of a sample of about 10^8 Ke_3 decays recorded with little bias. This analysis was performed upon the ACP system at the University of Chicago. First, the 20% sample was reanalyzed and the central value became -1×10^{-4} : the result moved by only 1/2 of the old systematic error even though nearly every aspect of the analysis was improved. The preliminary result for the FULL E731 data set, given at Fermilab in mid July, is $\epsilon'/\epsilon = (6.0 \pm 5.8(\text{stat.}) \pm 3.2(\text{syst.}) \pm 1.8(\text{monte-carlo})) \times 10^{-4}$ which combines to:

$$\epsilon'/\epsilon = (6.0 \pm 6.9) \times 10^{-4} \quad (\text{E731, full sample}).$$

The result is still consistent with zero.

TABLE I: RESULTS ON ϵ'/ϵ [10^{-4}]

Data Set	ϵ'/ϵ	Statistical Error	Systematic Error
NA31 '86	33	7	8++
E731 20%	-4	14	6
NA31 '88	17	7	7++
E731 20% reanalysis ⁺	-1	14	3.2
E731 Full⁺	6	6.1	3.2
NA31 '89 ⁺	21	5	6.5++
NA31 Full⁺	23	3.4	6.5

⁺Preliminary

⁺⁺Correlated errors

NA31 also announced a new result this summer and a listing of all of the recent results is given in Table I as shown above. It is seen that there is still a disagreement (at the 2σ level) and that while NA31 is dominated by (common) systematic uncertainties, E731 is statistically dominated.

Our group has also made very precise determinations of other parameters of the neutral kaon system with the same 2π data samples, using exactly the same techniques of background subtraction and acceptance corrections. The reported results, where both statistical and systematic errors are included, are:

$$\tau_s = (0.8912 \pm 0.0013) \times 10^{-10} \text{ sec};$$

$$\Delta m = (0.5339 \pm 0.0020) \times 10^{10} \text{ sec}^{-1};$$

$$\Phi_{+-} = (43.2 \pm 1.6)^0;$$

$$\Delta\Phi = \Phi_{+-} - \Phi_{00} = (-0.6 \pm 1.6)^0.$$

These results are either comparable to or exceed in precision the best previous determinations. In particular, the value for Φ_{+-} is in good agreement (as expected from CPT symmetry) with the expected "superweak phase", $43.5^0 \pm 0.14^0$, unlike the PDG value of $46.0^0 \pm 1.2^0$, which is largely based upon an NA31 result. The result for τ_s is three times better than the best previous determination, and the value for $\Delta\Phi$ is a factor of two better than earlier results of both E731 and NA31. This work will be the thesis of one Chicago PhD student.

We have as well made other measurements with this data set, some of which are given in the following Table:

PROCESS	RESULT	COMMENT (see publication list)
$K_L \rightarrow \pi^0 e e$	$< 7.5 \times 10^{-9}$ (90%)	published
$K_L \rightarrow \pi^0 \gamma \gamma$	$(1.86 \pm 0.85) \times 10^{-6}$	published
$K_L \rightarrow \pi^+ e^- \pi^0 \nu$	$(5.1 \pm 0.2 \pm 0.5) \times 10^{-5}$	720 events; form factor analysis underway; one Chicago PhD thesis student
$K_L \rightarrow \pi^+ \pi^- \gamma$ (IB)	$(1.40 \pm 0.05) \times 10^{-5}$	$E_{\gamma^*} > 20 \text{ MeV}$, to be published
$K_S \rightarrow \pi^+ \pi^- \gamma$ (IB)	$(4.59 \pm 0.14) \times 10^{-3}$	$E_{\gamma^*} > 20 \text{ MeV}$, to be published
$K_L \rightarrow \pi^+ \pi^- \gamma$ (DE)	$(3.04 \pm 0.14) \times 10^{-5}$	to be published
$\eta \rightarrow \gamma$	$(2.6 \pm 0.5 \pm 0.2) \times 10^{-3}$	First observation, to be published
$\Phi_{+-\gamma}$	$(41 \pm 28 \pm 11)^0$	First observation, to be published

In addition, from other data samples of E731, two physics results have been submitted to Phys. Rev. Lett. for publication. They are,

a) the first limit of the decay mode $K_L \rightarrow \pi^0 \nu \nu$: This decay is dominated by short distance effects and is almost entirely direct CP-violating within the Standard Model. Cuts were developed to reject the background processes $\Lambda \rightarrow n \pi^0$ and $K_L \rightarrow \pi^+ e^- \gamma \nu$. No candidate events were seen after these cuts were applied. We find $\text{BR}(K_L \rightarrow \pi^0 \nu \nu) < 2.2 \times 10^{-4}$ at the 90% confidence level. We have submitted this result to Phys. Rev. Lett. on 20th April, 1992.

The decay $K_L \rightarrow \pi^0 \nu \nu$ has been considered theoretically, but because of the formidable experimental difficulties, no search for this mode has previously been performed. The major attraction of this mode is its potential for the study of direct CP violation in the Standard Model. There is no long-distance contribution to $K_L \rightarrow \pi^0 \nu \nu$ and, based on existing upper limits for the $K^+ \rightarrow \pi^+ \nu \nu$ mode, the amplitude from indirect CP violation is negligible compared to direct CP-violating effects. The dominant, direct CP-violating, contribution to the decay is proportional to the imaginary part of V_{td} . In the Wolfenstein parametrization of the CKM matrix, $V_{td} = A \lambda^3 (1 - \rho - i \eta)$ and the Standard Model predicts

$$\text{BR}(K_L \rightarrow \pi^0 \nu \nu) = 1.5 \times 10^{-10} (M_t / 100 \text{ GeV})^2 A^4 \eta^2,$$

where M_t is the mass of the top quark. In the Standard Model, the magnitude of η can be related to the known value of ϵ : it is estimated that η lies somewhere between about 0.1 and 0.6. Based on the value $A = 0.85 \pm 0.09$, and the mass of the top quark determined indirectly from LEP data to be $157 \pm 40 \text{ GeV}/c^2$, the branching fraction for $K_L \rightarrow \pi^0 \nu \nu$ could be as large as 2×10^{-10} .

(b) the first measurement of the quadratic slope parameter in the $K_L \rightarrow 3\pi^0$ decay Dalitz plot: from a sample of 5.1 million $3\pi^0$ sample, we measure the quadratic slope parameter h in the Dalitz plot and it is $-(3.3 \pm 1.1 \pm 0.7) \times 10^{-3}$. This is the first measurement of the $3\pi^0$ slope parameter, and also the most sensitive measurement to date of any of the quadratic slope parameters for the charged and neutral kaons.

In order to check the validity of the $\Delta I = 1/2$ rule in the quadratic term, we combine our result with the linear and quadratic slope parameter values from the most sensitive charged K experiment according to the isospin formalism. We use the values from one experiment instead of the Particle Data Group averages since the latter are calculated separately for the K^+ and K^- decays, whereas the experimental values are calculated for the combined data set taking into account the significant correlations that exist. We obtain a value of $(31 \pm 10)\%$ for the $\Delta I = 3/2$ and $\Delta I = 1/2$ amplitude ratio ζ_3/ζ_1 , where the

contribution to the error from the charged K data is a little larger than that from our result.. This atypically large value of ζ_3/ζ_1 suggests a violation of the $\Delta I = 1/2$ rule in the quadratic term, and may provide a clue to the dynamical origin of the $\Delta I = 1/2$ rule. However, this conclusion is subject to the theoretical uncertainties in the Coulomb corrections applied to the charged K data. Further improvement in verifying the status of the $\Delta I = 1/2$ rule in the quadratic term will require simultaneous improvements in the slope parameter measurements in both the neutral as well as the charged K sector, and the reduction of systematic uncertainties in the charged K sector.

(B) E773

This experiment is a dedicated study of CPT symmetry, through the 2π decays of the neutral kaon. The same detector as for E731 is used with important upgrades. First, we (Task B) built a totally active regenerator, consisting in 24 small scintillators each with two phototubes. This is the first time such a regenerator has been built and we have shown that the inelastic background can be reduced substantially.

Data taking was completed on September 30th and enough statistics was collected to give a result on $\Delta\Phi$ to better than 0.5° . Two Chicago PhDs should result from this data and preliminary results are expected within the next fiscal year.

(C) E799 Phase I

This experiment focuses upon rare decays. We finished an 6 weeks data taking run on 8th of January 1992. Sensitivities approaching 10^{-10} for a variety of rare decays should be achieved. Our group (Task J) has provided 3 large transition radiation detectors for off-line electron identification. The modes to be studied are $K_L \rightarrow \pi^0 e e$, $\pi^0 \mu \mu$, $\mu \mu \gamma$, $\pi^0 \mu e$, $e e \gamma \gamma$. There are several '4-tracks' decay modes that we are sensitive to, namely $K_L \rightarrow e e e e$, $\mu \mu e e$, and $\pi^+ \pi^- e e$. In addition, we should be able to cleanly measure the $\pi^0 \rightarrow e e$ branching ratio, using the very clean $3\pi^0$ decays. Two more Chicago PhDs should result from this data. We are currently in the midst of reducing collected data (~1000 8mm cassettes). Some preliminary results may come as early as this summer.

(D) E799 Phase II / E832

We have, during this grant year, submitted a proposal to Fermilab to make another measurement of ϵ'/ϵ . Our goal is to achieve a precision of 0.0001 or a factor of about 6

better than with E731. We will be using the same technique with a substantially improved detector and beamline. The most important detector upgrade will be the electromagnetic calorimeter. Our Chicago NSF supported colleagues have been studying both BaF2 and pure CsI and have selected the CsI option. The same detector will be used for rare decay studies (E799 Phase II), reaching a sensitivity for 4-body decays of 10^{-11} . This program is nicknamed 'KTeV' (by Doug Jensen of Fermilab).

A comprehensive design report on the KTeV program was produced outlining the physics, the beam, the siting, and the detector with much detail. This KTeV report also describes the full detail of the calorimeter material tests and thus its justification for various parameters. The Fermilab Physics Advisory Committee has approved the program in February 1992 and we plan to run in 1994.

SUMMARY OF THE PROGRESS REPORT OF TASK J

Much happens during the last grant year as described above. This task (TASK J) specifically is responsible for the following:

- 1) as the scientific spokesman for E799 ;
- 2) built the first generation TRD for E799 Phase I. Data was taken during the three months running period. The performance of this detector will be comprehensively investigated and with the learned knowledge, we will proceed to design and build the second generation TRD for E799 Phase II ;
- 3) as advisor for two PhD graduate students, D. Harris who holds an Department of Education Fellowship and K. MacFarland who holds an National Science Foundation Fellowship. Both of them will have their PhD thesis based on data of E799 Phase I. Both of their supports (three years graduate fellowship limit of both Department of Education and National Science Foundation) will also be terminated by late 1992. Further progress on the issue, although not imminent, do need guidance from the Department of Energy ;
- 4) the lately submitted paper on the search of $K_L \rightarrow \pi^0 \nu \nu$ is the outcome of a senior thesis of an Chicago undergraduate student, Greg Graham with Y. Wah as advisor. He was supported by this task as undergraduate summer research student. Another Chicago senior (Kent Konkol) is working on the $K_L \rightarrow \pi^0 \pi^+ \pi^- \gamma$ decay mode which has never been seen before. He will submit this work as his senior thesis with Y. Wah as advisor. We expect to publish the result at the end of this summer.

Publication for past three years :

Journal publication:

1. 'A Cluster Finding Trigger Processor For a High Energy Physics Experiment Searching For New Source of CP Violation' IEEE Trans.Nucl. Sci. **36**,358(1989).
2. 'Search for $K_L \rightarrow \pi^0 \gamma \gamma$ ' Phys. Rev. Lett. **63**, 28 (1989).
3. 'A Cluster Finding Trigger Processor for a High Energy Experiment ' NIM A**291**, 577 (1990).
4. 'New Limit on $K_L \rightarrow \pi^0 e^+ e^-$ ' Phys. Rev. D**41**, 3546 (1990).
5. 'A Determination of ϵ'/ϵ by the Simultaneous Detection of the Four $K_L \rightarrow 2\pi$ Decay Modes' Phys. Rev. Lett. **64**, 1491 (1990).
6. 'A Measurement of the Phase Difference between η_{00} and η_{+-} in $K_L \rightarrow 2\pi$ Decays' Phys. Rev. Lett. **64**, 2976 (1990).
7. 'Measurement of the branching ratio of the decay $K_L \rightarrow \pi^0 \gamma \gamma$ ' Phys. Rev. D **44**, R573 (1991).

Submitted to journal:

1. 'A Measurement of the quadratic slope parameter in the $K_L \rightarrow 3\pi^0$ decay dalitz plot' Accepted publication by Phys. Rev. Lett. .
2. 'Search for the decay $K_L \rightarrow \pi^0 \nu \nu$ ' Submitted to Phys. Rev. Lett., 20 April 1992.

Conference Contributions :

1. 'A Progress Report of Fermilab Experiment E-731: A Precision Experiment to Measure ϵ'/ϵ and Rare Decay modes of the Neutral Kaon' The Second International Symposium on the Fourth Family of Quarks and Leptons, Annals of New York Acad. of Science, Vol.578, 162 (1989).
2. 'Precision Experiments to Determine $\text{Re}(\epsilon'/\epsilon)$ by And to Study Neutal Kaon Rare Decay', Proceeding of the 1989 Theoretical Advanced Study Institute in Elementary Particle Physics (TASI), World Scientific,725 (1989).
3. 'Background Estimate for $K_L \rightarrow \pi^0 \mu \mu$ with a Main Injector Fixed Target Beam and Detector', Proceedings of the Workshop on Physics at the Main Injector; Fermilab, Edited by S. Holmes and B. Winstein, 237 (1989).
3. 'New Results on rare neutral kaon decay from Fermilab Experiment E731', Proceeding of the 25th International Conference on High Energy Physics, Vol.II, 1221 (1990).

4. 1992 Aspen Winter Conference on Elementary Particle Physics, Jan 1992; Edited by T. Appelquist, P. Ramond, and S. Meshkov.

Reports :

1. Conceptual Design Report : Kaons At the Main Injector. Fermilab FN-568 (1991).
2. KTeV Design Report. Fermilab FN-580 (1992).

END

**DATE
FILMED**

8 / 11 / 92