STUDIES IN THEORETICAL HIGH ENERGY PARTICLE PHYSICS

Technical Progress Report

Henrik Aratyn, Lee Brekke, Wai-Yee Keung, and Uday Sukhatme

Department of Physics
University of Illinois at Chicago
801 W. Taylor Street, Rm #2236
Chicago, Illinois 60687-7059

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

January 1993

Prepared for
THE U.S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-84ER40173

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ABSTRACT

This is a technical progress report for grant no. DE-FG02-84ER40173 describing research carried out in 1991-1992.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

TECHNICAL PROGRESS REPORT (Grant No. DE-FG02-84ER40173)

In this report, we describe technical progress achieved in various areas of research in 1991-1992. A detailed account of our work can be found in the following articles, which have either been published or circulated in preprint form. The publications are labeled by the names of physicists in our group [A = Aratyn, B = Brekke, K = Keung, S = Sukhatme].

- A1. Symplectic Geometry and Infinite-Dimensional Symmetry Groups
By H. Aratyn, E. Nissimov and S. Pacheva
Talk at the XXth International Conference on Differential Geometric Methods in Theoretical Physics, June 1991, New York, in Differential Methods in Theoretical Physics, World Scientific (1992) 234-243.
- A2. Kac-Moody Construction of Toda Type Field Theories
By H. Aratyn, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
Phys. Lett. **254B**, 372-380 (1991).
- A3. Infinite-Dimensional Noether Symmetry Groups and Quantum Effective Actions from Geometry
By H. Aratyn, E. Nissimov and S. Pacheva
Phys. Lett. **255B**, 359-366 (1991).
- A4. Conformal Theories, Integrable Models and Coadjoint Orbits
By H. Aratyn, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
In Proceedings of the VI J.A. Swieca Summer School, Particles and Fields, Campos do Jordão, Brazil, 14-26, January, (1991), World Scientific (1992) 560-575.
- A5. Higher Spin Symmetries and w_∞ algebra of the Conformal Affine Toda Model
By H. Aratyn, C.P. Constantinidis, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
Phys. Lett. **281B**, 245-253 (1992).
- A6. Classical r -matrices and Poisson Bracket Structures on Infinite Dimensional Groups
By H. Aratyn, E. Nissimov and S. Pacheva
Phys. Lett. **284B**, 273-282 (1992).
- A7. A New Deformation of W -Infinity and Applications to the Two-loop WZNW and Conformal Affine Toda Models
By H. Aratyn, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
Phys. Lett. **293B**, 67-71 (1992).
- A8. On Two-Current Realization of KP Hierarchy
By H. Aratyn, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
28 pp. July 1992, to appear in Nucl. Phys. B.
- A9. R-Matrix Formulation of KP Hierarchies and Their Gauge Equivalence
By H. Aratyn, E. Nissimov and S. Pacheva and I. Vaysburd
Phys. Lett. **294B**, 167-176 (1992).

- A10. Hirota's Solitons in the Affine and the Conformal Affine Toda Models
By H. Aratyn, C.P. Constantinidis, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
November 1992, 36 pp. Submitted for publication.
- A11. Basics of W Infinity Algebras and Some Applications in Physics
By H. Aratyn, L.A. Ferreira, J.F. Gomes and A.H. Zimerman
Talk at the XIII-th meeting of the Particles and Fields of the Brazilian Physics Society, Caxambu, September 1992.
- B1. Anyons from Bosons
By L. Brekke, A.F. Falk, S. Hughes and T.D. Imbo
Phys. Lett. B271, 73 (1991).
- B2. Alice Strings, Magnetic Monopoles and Charge Quantization
By L. Brekke, W. Fischler and T.D. Imbo
Phys. Rev. Lett. 67, 3643 (1991).
- B3. Statistics of 2D Solitons
By L. Brekke and T.D. Imbo
Int. J. Mod. Phys. A7, 2589 (1992).
- B4. Knots and Links of Nonabelian String
By L. Brekke, H. Dykstra, S.J. Hughes and T.D. Imbo
Phys. Lett. B288, 273 (1992).
- B5. Novel Spin and Statistical Properties of Nonabelian Vortices
By L. Brekke, H. Dykstra, A.F. Falk and T.D. Imbo
University of Illinois at Chicago preprint UICHEP-TH/92-15 (Submitted to Phys. Lett. B).
- B6. Avoiding the Dirac Quantization Condition on a Torus
By L. Brekke and T.D. Imbo
University of Illinois at Chicago preprint UICHEP-TH/92-2 (1992).
- B7. Spinning Particles, Braid Groups and Solitons
By L. Brekke, M.J. Dugan and T.D. Imbo
University of Illinois at Chicago preprint UICHEP-TH/92-17 (1992).
- B8. Rings and Balls
By L. Brekke, S.J. Hughes and T.D. Imbo
University of Illinois at Chicago preprint UICHEP-TH/92-19 (1992).
- K1. The Chromoelectric Dipole Moment of the Heavy Quark and Purely Gluonic CP Violating Operators
By W.-Y. Keung, D. Chang, T.W. Kephart and T.C. Yuan
Phys. Rev. Lett. 68, 439-443 (1992).

- K2. Anomalous Dimensions for All Dimension - Eight CP Odd Operators with Only Photons and Gluons
By W.-Y. Keung, D. Chang, I. Phillips and T.C. Yuan
Phys. Rev. D46, 2270-2271 (1992).
- K3. An Effective Field Theory for the Neutron Electric Dipole Moment
By W.-Y. Keung, D. Chang, T.W. Kephart and T.C. Yuan
Nucl. Phys. B384, 147-167 (1992).
- K4. Neutron Electric Dipole Moment due to Higgs-Boson Exchange in Left-Right-Symmetric Models
By W.-Y. Keung, D. Chang, X.-G. He, B. McKellar and D. Wyler
Phys. Rev. D46, 3876-3883 (1992).
- K5. Simple Mechanism for Large Neutrino Magnetic Moments
By W.-Y. Keung, K.S. Babu, D. Chang and I. Phillips
Phys. Rev. D46, 2268-2269 (1992).
- K6. A Class of Models with Large Neutrino Magnetic Moment
By W.-Y. Keung, D. Chang and I. Phillips
FERMILAB-PUB-92-143-T, submitted to Nucl. Phys. B.
- K7. CP Violation in the Decay of Neutral Higgs Boson into $t\bar{t}$ and W^+W^-
By W.-Y. Keung and D. Chang
Fermilab preprint, FERMILAB-PUB-92-172-T (1992).
- K8. CP Violation in Top Pair Production at an e^+e^- Collider
By W.-Y. Keung, D. Chang and I. Phillips
CERN preprint CERN-TH-6658/92 (1992).
- S1. Mapping of Shape Invariant Potentials Under Point Canonical Transformations
By R. De, R. Dutt and U. Sukhatme
Jour. Phys. A25, L843-L850 (1992).
- S2. Bound States in the Continuum from Supersymmetric Quantum Mechanics
By J. Pappademos, U. Sukhatme and A. Pagnamenta
University of Illinois at Chicago preprint UICHP-TH/92-9 (1992).
- S3. Singular Superpotentials in Supersymmetric Quantum Mechanics
By P. Panigrahi and U. Sukhatme
University of Illinois at Chicago preprint UICHP-TH/92-7 (1992).
- S4. Path Integral Solutions for Shape Invariant Potentials Using Point Canonical Transformations
By R. De, R. Dutt and U. Sukhatme
Phys. Rev. A46, 6869 (1992).
- S5. Supersymmetry and Tunneling in an Asymmetric Double Well
By A. Gangopadhyaya, P. Panigrahi and U. Sukhatme
Phys. Rev. A47 (1993).

- S6. Semiclassical Approach to Quantum Mechanical Problems with Broken Supersymmetry
By R. Dutt, A. Gangopadhyaya, A. Khare, A. Pagnamenta and U. Sukhatme
University of Illinois at Chicago preprint UICHEP-TH/92-14 (1992).
- S7. Gluon Structure Functions in Nuclei from the Quark Cluster Model
By U. Sukhatme, G. Wilk and K. Lassila
Zeit. Phys. C53, 439 (1992).
- S8. Evidence for Six-Quark Clusters in QCD Processes
By A. Petridis, C. Carlson, K. Lassila and U. Sukhatme
Proc. Amer. Phys. Soc. Meeting, Vancouver, Canada (1991).
- S9. Shadowing of the Gluon in Fixed Target Experiments
By K. Lassila, A. Petridis, U. Sukhatme and G. Wilk
Phys. Lett. B297, 191 (1992).
- S10. Lognormal Multiplicity Distributions and the Dual Parton Model
By U. Sukhatme
Proc. XXII International Symposium on Multiparticle Dynamics, Santiago de Compostela, Spain (1992).
- S11. Dual Parton Model
By A. Capella, U. Sukhatme, C.I. Tan and J. Tran Thanh Van
University of Paris, Orsay preprint (1992); to be published in Physics Reports.

We give below a brief description of the main areas in which new physics results were obtained.

I. H. Aratyn

A. Symmetry Structure of Conformal Affine Toda Model and KP Hierarchy (A2,A5,A7,A8)

Building on results previously obtained by us in [A2], showing that the conformal affine Toda models can be obtained via Hamiltonian reduction from a two-loop Kac Moody algebra, we proposed in [A5] a systematic procedure to analyze the higher spin symmetries of the conformal affine Toda models. Our method was based on an explicit construction of infinite towers of extended conformal symmetry generators. Two fundamental building blocks of this construction were special spin-one and -two primary fields characterizing the conformal structure of these models. We have obtained as a special limit the algebra of area preserving diffeomorphisms on a 2-manifold (w_∞ algebra). We therefore suspected that there exists formulation in which the extended conformal symmetry appears as a special deformation of the algebra w_∞ of area preserving diffeomorphisms of a 2-manifold and constitutes therefore a closed algebra. To obtain such a formulation we constructed in [A7] first a centerless W -infinity type of algebra in terms of a generator of a centerless Virasoro algebra and an abelian spin-1 current, which conventionally emerges in the study of pseudo-differential operators on a circle or alternatively within KP hierarchy with Watanabe's bracket. We then designed in this framework a new deformation technique producing the extended conformal symmetry as a special deformation of the algebra w_∞ . Now the generators of the extended conformal symmetry spanned a closed algebra known in literature as W_∞ , which is a quantum counterpart of the algebra w_∞ of area preserving diffeomorphisms of a 2-manifold. This technique applied both to the two-loop WZNW and conformal affine Toda models, establishing henceforth W_∞ invariance of these models.

Since W_∞ algebra describes a first Poisson bracket structure of the KP hierarchy characterizing a large class of integrable models we turned in [A8] our attention towards KP hierarchy. We have formulated a simple description of the KP hierarchy and its multi-hamiltonian structure in terms of two Bose currents. This turned out to be a natural setting to study our deformation scheme connecting various W -infinity algebras and relations between two fundamental nonlinear structures. Applications of our method were given for the Conformal Affine Toda model, WZNW models and discrete KP approach to Toda lattice chain.

B. Solitons in the Affine Toda and Conformal Affine Toda Models (A10)

Most recently in [A10] we used Hirota's tau-function method formulated as a recursive scheme to construct complete set of soliton solutions for the affine Toda

field theory based on an arbitrary Lie algebra. Our solutions included a new class of solitons connected with two different type of degeneracies encountered in the Hirota's perturbation approach.

We also derived an universal mass formula for all Hirota's solutions to the Affine Toda model valid for all underlying Lie groups. Embedding of the Affine Toda model in the Conformal Affine Toda model played a crucial role in this analysis.

C. Classical r -matrices and Poisson Bracket Structures on Infinite-Dimensional Groups (A6)

Starting with a canonical symplectic structure defined on the cotangent bundle T^*G we derived in [A6], via Dirac Hamiltonian reduction, Poisson brackets (PB's) on arbitrary infinite-dimensional group G (admitting central extension). We found that the PB structures were given in terms of an r -operator kernel related to the two-cocycle of the underlying Lie algebra and satisfying a differential classical Yang-Baxter equation. The explicit expressions of the PB's among the group variables for the $(N,0)$ for $N = 0,1,\dots,4$ (super-) Virasoro groups and the group of area-preserving diffeomorphisms on torus were also obtained.

D. R-Matrix Formulation of KP Hierarchies and Their Gauge Equivalence (A9)

The Adler-Kostant-Symes R-bracket scheme was applied in [A9] to the algebra of pseudo-differential operators to relate the three integrable hierarchies: KP and its two modifications, known as non-standard integrable models. All three hierarchies were shown to be equivalent and a connection was established in the form of a symplectic gauge transformation. This construction resulted in a new representation of the W -infinity algebras in terms of four boson fields.

II. L. Brekke

A. Statistics of Particles and Solitons (B1, B3,B5,B7,B8)

We have developed topological techniques for determining the allowed statistics for particles in quantum mechanics and solitons (and other topological objects) in field theories. We have also extended our topological methods to include particles with spin moving on very general manifolds and the relationship between spin and statistics. In particular, we demonstrated that certain field theories living on a spatial circle can have various exotic forms of statistics including ambistatistics, where the superselection rule between bosons and fermions is effectively violated. This exotic form of statistics was also seen in the behavior of nonabelian vortices in 2+1 dimensions. These vortices can arise when nonabelian gauge theories are spontaneously broken down to discrete nonabelian subgroups. In the general case, fractional statistics, para-statistics and

their nonabelian generalizations were found to occur as well as the usual Bose and Fermi statistics.

We also investigated the behavior of composite systems of two or more particle types in quantum mechanics in two spatial dimensions. The allowed linking of the world lines of the particles in space-time leads to possible additional phases in the quantum theory. In particular, by choosing these phases appropriately, two bosons can form a composite which has an arbitrary statistical angle. This behavior was also seen for two distinct soliton types in particular 2+1 dimensional nonlinear sigma models. We explicitly constructed a topological term which generates the additional phase in a limit where the solitons are forbidden to overlap. Similarly, we explicitly constructed topological terms corresponding to the threading of ball-like or ring-like solitons through the center of other ring-like solitons (of distinct type) as a function of time in 3+1 dimensions. The methods we used should be more generally useful in examining linking phenomena in field theories.

B. Charge Quantization in the Presence of an Alice String (B2)

Alice strings are a class of cosmological strings in which electric charge changes sign after circling the string. We showed that charge must be quantized in the presence of such a string. Field theories possessing Alice strings were shown to also contain magnetic monopoles. The charge quantization coming from the presence of Alice strings is consistent with the Dirac quantization condition from the magnetic monopoles.

C. Knotting and Linking of Nonabelian Flux (B4,B6)

When a 3+1 dimensional nonabelian gauge theory is spontaneously broken down to a nonabelian discrete subgroup the string-like configurations which carry nonabelian fluxes can occur. The presence of the nonabelian flux constrains the way in which the strings can knot or link with each other. We analyzed these constraints using a general topological framework and identified some particular configurations which we call "essentially knotted". These configurations cannot be deformed into the vacuum without overlapping the cores of two segments of string with noncommuting flux elements. We also showed how the usual Dirac quantization condition can be modified for a charged particle on a toroidal surface by threading nonabelian flux through the holes in the torus.

III. W.-Y. Keung

A. Electric Dipole Moments (K1,K4)

Following the recent studies of Weinberg and Barr and Zee on the electric dipole moments of the neutron and the electron, we have set up the correct QCD evolution

equation for the relevant operators, in particular, the chromoelectric dipole moments of quarks which play an important role in the neutron electric dipole moment. We have shown the importance of the dimension 8 operator due to the QCD enhancement. Complete anomalous dimensions of dimension 8 operators have been worked out as well as the relevant matching conditions when these dimension 8 operators are induced by the chromoelectric dipole moment of the heavy quarks. We have also performed detailed phenomenological studies on the left-right-symmetric model for its contribution to the neutron electric dipole moment.

B. Neutrino Physics in Gauge Theories (K5,K6)

We have studied a mechanism which gives a small mass but a large magnetic moment for the neutrino such that the solar neutrino deficit problem can be explained. We have shown that the simple Zee Model of neutrino mass already incorporates an automatic mechanism to provide a large transitional neutrino magnetic moments. Other scenarios, with a vector quark or a triplet Higgs, beyond the standard model have also been studied.

C. CP Violation in the High Energy Colliders (K7,K8)

We have shown that the CP nonconservation can manifest itself in the asymmetry in the energy spectra between the lepton and the anti-lepton from the decays of heavy particles, like W^+W^- or $t\bar{t}$.

IV. U. Sukhatme

A. Supersymmetric Quantum Mechanics (S1-S6)

During the past year, we have continued our study of various aspects of supersymmetric quantum mechanics. Using point canonical transformations, it has been possible to map all known shape invariant potentials into two classes. The eigenfunctions in these two classes are given by hypergeometric and confluent hypergeometric functions respectively [S1]. We have also explicitly evaluated the Feynman kernels for all shape invariant potentials [S4].

In the past, our group has constructed the most general family of potentials, all of which have the same bound state energies and scattering amplitudes. If this formalism of generating isospectral families is applied to a state in the classical continuum, one gets the unusual occurrence of bound states in the continuum, generated in a systematic fashion [S2]. This purely quantum mechanical phenomenon was first discussed by von Neumann and Wigner shortly after the birth of quantum mechanics, and was generally regarded as a mathematical curiosity. However, it has very recently been experimentally observed in semiconductor heterostructures grown by molecular beam epitaxy.

We have carefully examined the role of singular superpotentials and their relation to explicit breaking of supersymmetry, occurrence of negative energy eigenstates and breakdown of the degeneracy theorem [S3].

Finally, our group has started to examine quantum mechanical problems with broken supersymmetry [S6]. A number of exactly solvable examples have been found, all of which can be mapped to shape invariant potentials with unbroken supersymmetry. A semiclassical formalism for broken supersymmetry problems has been developed. It has the desirable feature of giving exact answers for analytically solvable problems.

B. Parton Structure Functions in Nuclei (S7-S9)

Multiquark clusters in nuclei have been shown to provide a phenomenological description of the EMC effect, the Drell-Yan process and the backscattering of hadrons from nuclear targets. We have also looked at J/ψ production and ways to study the gluon EMC effect [S8,S9] which is predicted in the multiquark cluster framework.

C. Dual Parton Model (S10-S11)

The multiplicity distributions for proton-proton collisions have been computed using the dual parton model and a lognormal function to describe single chain e^+e^- multiplicity distributions [S10]. The agreement with pp multiplicity data was found to be very good provided that the valence quark momentum distribution functions in a proton at small values of x are taken to be of the form $x^{-\gamma}$ with $\gamma \simeq 0.7$, somewhat steeper than the canonical $x^{-1/2}$ behavior used in previous work. We have also written a detailed review of the dual parton model and submitted it to Physics Reports for publication [S11].

THEORETICAL HIGH ENERGY PHYSICS AT THE UNIVERSITY OF ILLINOIS AT CHICAGO

At present, there are five faculty members actively involved in theoretical high energy physics research at UIC. They are Professors Henrik Aratyn, Wai-Yee Keung, Tom Imbo, Antonio Pagnamenta and Uday Sukhatme. Dr. David Barclay joined our group as a postdoctoral fellow in September 1992. Prior to that our postdoctoral fellow was Dr. Lee Brekke (1990-1992). [Dr. Brekke will continue to be a postdoctoral fellow through August 1993, supported by university funds]. We have been fortunate in obtaining extensive support from the university in the form of partial funding for numerous long term visitors. During the past four years, our visitors have been Prof. Ranabir Dutt, Visva Bharati University, India; Prof. K. V. L. Sarma, Tata Institute of Fundamental Research, India; Prof. J. Jayaraman, Univ. Federal da Paraiba, Brazil; Dr. A. Khare, Institute of Physics, Bhubaneswar, India; Mr. Q. M. Wang, Anhui Normal University, China. Also, Dr. L. A. Ferreira, Univ. Paulista do Estado de Sao Paulo, Brazil was a visitor for two months in 1992 and Prof. A. H. Zimmerman, Brazil visited for two months each in 1990 and 1991. The theoretical physics effort benefits substantially from interaction with the experimental high energy physics group involved in performing or planning experiments at Fermilab, RHIC and the SSC. In August 1992, Dr. Tom Imbo [formerly a Junior Fellow of the Harvard University Society of Fellows] joined our department as a tenure-track assistant professor. His addition to our group will further strengthen the research effort at UIC.

THEORETICAL HIGH ENERGY PHYSICS AT THE UNIVERSITY OF ILLINOIS AT CHICAGO

Proposed Technical Program (7/1/93-6/30/94)

Henrik Aratyn, David Barclay, Wai-Yee Keung, Uday Sukhatme

1. Aratyn

A1. Integrable field theoretical models via reduction

We intend to elaborate, in a specific field-theoretic setting, on the general problem: what precise assumptions and restrictions are to be imposed on a particular physical model in order to reduce it to the form of a completely integrable lower dimensional system? Some remarkable progress has already been made in this direction, but much work still remains to be done. To mention just one open problem: it is still not known which integrable hierarchy describes the matrix model representation of subcritical superstrings, though attempts have been made to identify it. We hope that the concept of gauge equivalence introduced by us into the setting of KP hierarchy will shed some light on this problem.

A2. Symplectic-geometric approach to R-matrix Hamiltonian structures

We plan to continue developing our symplectic-geometric approach to R-matrix Hamiltonian structures on infinite-dimensional groups and carry out its natural geometric generalization, through the Lie-Poisson analysis, suited to describe the semiclassical limit of quantum groups in the infinite-dimensional case. We expect that:

- this will open the avenue to find new representations for W_{∞} algebra which, was recently found to be the symmetry algebra in a large variety of important physical problems;
- this will produce methods of hamiltonian reduction to obtain W_{∞} from some affinization of the algebra of linear transformations in an infinite-dimensional linear space, which will also provide constructive approach to the representation theory of the former.
- this will shed light on the meaning of the group of dressing transformations which is defined for each integrable system. Specifically, we shall study the KP integrable hierarchy and its modifications. We shall aim to answer questions like: what is the algebra of Poisson brackets of the dressing transformations; what is the relation between the infinite number of local conserved charges (the Hamiltonians), pertinent to each integrable field theory, and the generators of the dressing transformations which correspond to non-local symmetries.

A3. Quantization of models with infinite-dimensional symmetry groups.

We shall investigate various aspects of quantization of models with infinite-dimensional symmetry groups. In particular:

- We would like to study quantization of KP hierarchy. The solution to this problem presumes finding a deformation of the universal enveloping algebra of the algebra of the fundamental Poisson brackets, i.e., the universal quantum R-matrix and the corresponding quantum group structure, as a second step.
- We plan to contribute to the very interesting program, initiated by Zamolodchikov, to construct new quantum integrable models, based on integrable perturbations of conformal field theories. We shall be aimed at developing a systematic Lagrangian field-theoretic approach for construction integrable deformations. The main idea here stems from the fact that adding the perturbing composite fields does not change the Hamiltonian structure (the fundamental Poisson brackets) of the original conformal model. Thus, deriving the infinite set of independent integrals of motion in involution of the full (deformed) integrable field theory amounts to the construction of the set of all functionals of the fundamental conformal fields whose mutual Poisson brackets vanish.

2. Barclay

B1. WKB-Type Series

We intend to continue to explore the properties of both the WKB and SWKB series, including the broken-SUSY WKB series recently introduced by the UIC group. Much work has been done elsewhere on case-by-case comparisons of the two approximations, but a general understanding of their relative performances is still a completely open question. As a first step towards a solution, our priority is resolving the question of in how many cases the approximations are exact.

B2. QCD Perturbation Series

Our current interest in QCD centers on the flavor dependence of the perturbative coefficients. Apart from the question of how this may force modifications in our existing jet-fraction project, this topic relates to the large-order behavior and renormalons via the large- N expansion and to the scheme-dependence problem via the BLM scale-fixing method. In collaboration with Dr. C.J. Maxwell and Matthew Reader (Durham) we are pursuing a "pole approximation" in the flavor dependence which will yield estimates of uncalculated coefficients in QCD series and which should thus be of use phenomenologically. This approach will further increase the utility of the Effective Charge Formalism in the analysis of QCD data as previously advocated. We are also, engaged in trying to understand the significance of the new scheme invariants (again related to the flavor dependence) recently identified by Grunberg and we will continue in trying to extend partial results relating to these already obtained.

3. Keung

K1. Collider Physics

We will continue our efforts in looking for the CP violating effect in the high energy colliders. There may be large CP violation in the Higgs interaction. The effect could appear in the energy asymmetry between the lepton and the anti-lepton from the decays of heavy particles. Another measurable quantity of CP nonconservation is the up-down asymmetry with respect to the reaction plane. Detailed phenomenological analysis will be pursued.

K2. $g-2$

Based on our previous experience in the two-loop diagrams, we will perform a complete two-loop calculation for the Higgs contribution to the $g-2$ electron anomalous magnetic moment in the Standard Model. Non-decoupling behavior in the limit of a heavy Higgs boson will be discussed. Our calculation will be relevant to the test of the Standard Model as the experimental precision on $g-2$ is improving.

K3. Neutrino Physics

Different mechanisms for giving a large magnetic moment to the neutrino have been studied as a solution to the puzzle that the neutrino flux from the Sun is lower than that of the canonical prediction in the standard solar model. We will confront such scenario with data from those current experiments dedicated to the detection of the solar neutrinos.

K4. Beyond Static Limit in the Quarkonium Decays

Based on our previous work on the bound states of a heavy quark pair, we wish to develop a systematic program to calculate the annihilation rate beyond the static limit. This may provide a better understanding of the strong interaction.

4. Sukhatme

S1. Supersymmetric Quantum Mechanics

Quantum mechanical potentials satisfying the property of shape invariance are well known to be algebraically solvable. Shape invariance implies that the supersymmetric partner potentials $V_+(x)$ and $V_-(x)$ have the same x -dependence except for a change of parameters $a_1 = f(a_0)$. In all known shape invariant potentials, a_1 and a_0 are related by a translation $a_1 = a_0 + \alpha$. We have recently investigated the scaling ansatz $a_1 = qa_0$, and found new types of shape invariant potentials. A special case of our results are the self-similar potentials recently discussed by Shabat and Spiridonov. Our potentials seem to be q -deformations of familiar potentials like the harmonic oscillator and Pöschl-Teller potential.

q-deformations of potentials corresponding to multi-soliton systems are currently being investigated.

S2. Multichain Interactions in Soft Hadron Physics

In the dual parton model, independent multi-chain diagrams correspond to unitarity corrections to the leading two chain, single Pomeron exchange diagram. Recently, the consequences of multi-chain interactions have been considered by Pajares, Braun and others. We plan to study the approach in more detail using lognormal distributions for single chain inputs, since these have been shown to give a very economical description of proton-proton multiplicity data. The consequences of such a study should be of special importance at the energies of LHC and SSC.

END

DATE
FILMED

4 129/93