Capillary Effects in Drainage in Heterogeneous Porous
Media: Continuum Modeling, Experiments and Pore
Network Simulations

Topical Report

By
Baomin Xu
Yanis C. Yortsos

DOE/BC/14600--43

DE93 000134

April 1993

Work Performed Under Contract No. DE-FG22-90BC14600

Prepared for
U.S. Department of Energy
Assistant Secretary for Fossil Energy

Thomas B. Reid, Project Manager
Bartlesville Project Office
P.O. Box 1398
Bartlesville, OK 74005

Prepared by
University of Southern California
Petroleum Engineering Program
Department of Chemical Engineering
Los Angeles, CA 90089-1211

MASTER

Contents

1	INTRODUCTION	2
2	THEORY	4
3	EXPERIMENTS	8
4	PORE NETWORK SIMULATIONS	12

List of Figures

1	Schematic saturation response to heterogeneity in permeability	7
2	Permeability map of experimental set up	9
3	Experimental saturation profiles at various times (a) $\varepsilon = 35.3$, (b) $\varepsilon = 3.53$	10
4	Saturation jump vs. ε	11
5	Numerical simulations for the conditions of Figure 3	13
6	Typical lattices used in pore network simulations	15
7	Saturation profile and displacement patterns for permeability increase (1:2) and for $Ca = 1.5 \times 10^{-5}$	17
8	Saturation profile and displacement patterns for permeability increase (1:2) and for $Ca = 1.5 \times 10^{-4}$	18
9	Saturation profile and displacement patterns for permeability increase (1:2) in a ramp of 20 lattice spacings for $Ca = 1.5 \times 10^{-4}$	19
10	Saturation profile and displacement patterns for uniform permeability and $Ca =$ 1.5×10^{-4}	20
11	Saturation profile and displacement patterns for permeability decrease (2:1) and $Ca = 1.5 \times 10^{-5}$	21
12	Saturation profile and displacement patterns for sinusoidal variation and $Ca =$ 1.5×10^{-4}	22
13	Saturation profile and displacement patterns for the conditions of Figure 8 and $M = 10$	24
14	Saturation profile and displacement patterns for the condition of Figure 8 and $M = 0.1$	25
15	Invasion percolation with an abrupt decrease (2:1) in permeability at various stages	27
16	Invasion percolation with an abrupt increases (1:2) in permeability and at various stages	28

ABSTRACT

We investigate effects of capillary heterogeneity induced by variations in permeability in the direction of displacement in heterogeneous porous media under drainage conditions. The investigation is three-pronged and uses macroscopic simulation, based on the standard continuum equations, experiments with the use of an acoustic technique and pore network numerical models. It is found that heterogeneity affects significantly the saturation profiles, the effect being stronger at lower rates. A good agreement is found between the continuum model predictions and the experimental results based on which it can be concluded that capillary heterogeneity effects in the direction of displacement act much like a body force (e.g. gravity). A qualitative agreement is also found between the continuum approach and the pore network numerical models, which is expected to improve when finite size effects in the pore network simulations diminish. The results are interpreted with the use of invasion percolation concepts.

1 INTRODUCTION

Displacement processes in porous media, such as oil reservoirs, soils and water aquifers, find important engineering applications in oil recovery and in subsurface remediation and cleanup. Of particular significance to such processes is the inherent heterogeneity of medium properties, key among which are permeability and porosity. While recognized in early studies, a systematic investigation of the effects of heterogeneity has been undertaken mostly during the last decade [1]. In immiscible displacement, pore-scale studies led to the development of novel statistical tools to characterize displacement patterns in random media, such as Invasion Percolation (IP) and Diffusion-Limited-Aggregation (DLA) [2]. In miscible displacement, large-scale investigations on effects of randomness and correlation have led to classifications of flow regimes in terms of correlation length, degree of heterogeneity and mobility ratio [3]. For example, for a fixed correlation length, an increase in the heterogeneity index leads successively to a transition from a viscous fingering (VF) to a dispersive and to a channeling regime.

The preponderance of large-scale heterogeneity investigations on miscible displacement (for an exception, see [5]) reflects mainly the premise that on a large scale, differences between immiscible and miscible displacements are minimal. However, this hypothesis is not unconditionally nor uniformly valid, and depends on both geometric and flow properties. At sufficiently low capillary numbers (more precise estimate for which are developed later in the text), for example, permeability gradients have effects similar to gravity, leading to gradient percolation patterns (both stable and unstable) [6]; sharp changes in permeability in the direction of flow significantly change steady-state saturation profiles [7]; random distribution of permeabilities attribute large-scale percolation features (such as large-scale trapping) to displacements [8]; while correlated fields drastically alter the percolation aspects of the displacement [9]. All these are capillary heterogeneity effects driven by permeability heterogeneity.

A simple, but convenient, first-step to explore capillary heterogeneity effects is by using the continuum equations that are presumed to model immiscible displacement. This was undertaken in recent studies, where we investigated the solutions of these equations in 1-D flows (where capillary heterogeneity effects are maximized) under both steady-state (concurrent or counter-current) and transient conditions [10]. In such displacements, capillary and permeability (and wettability)

heterogeneity are interrelated via the Leverett relationship $P_c = \frac{\gamma}{\sqrt{k}} J(S)$, where γ is the interfacial tension and J a dimensionless function of the saturation S . It was found that during drainage in a medium where the permeability k suddenly changes, the response of the non-wetting saturation is a well-pronounced kink in the saturation profile, pointing downwards for a permeability increase and upwards in the opposite case. The size of the kink was shown to decrease with an increase in the capillary number, Ca . At steady-state flow, the continuum predictions were tested with experiments and found to be in excellent agreement [7]. Good experimental support was also found for transient drainage in experiments independently performed, and with no intention to test the theory, in eolian sandstones which involve spatial permeability oscillations [10].

The preliminary agreement between continuum theory and experiments is somewhat surprising. Capillary heterogeneity predominates at low Ca , where capillary effects lead to fractal percolation patterns, at least over the length scale determined by the viscous correlation length. In these regimes the continuum approach, and particularly the Leverett J-function, is least likely to apply. More generally, the validity of the continuum equations for drainage in heterogeneous media has not been investigated. To resolve these questions and to provide a rigorous test of the existing theory we conducted a series of experiments involving a well-characterized heterogeneity and we developed alternative models based on a pore-network simulator. Experimental and pore-network results were then compared to the continuum theory. In addition, a pore-level analysis was employed to interpret experimental, pore-network model and continuum results.

Under the conditions tested, the experimental results were found to be in good agreement with the continuum equations. Pore network simulations also showed the qualitative trends expected from the continuum theory, although a quantitative agreement was not attempted due to the strong finite size effects and fluctuations in the pore network model. The paper is organized as follows: We first proceed with a brief review of the continuum theory and its implications. The experimental procedures and results are subsequently presented and compared to the theory. Then, a pore network simulator is developed and used to simulate these processes in various heterogeneous media. The final section focuses on interpretation of the various findings based on percolation theory and on attempts to unify the results obtained.

2 THEORY

We summarize briefly the results obtained from the continuum models. Consider the constant rate, immiscible, two-phase displacement (drainage) in an 1-D system with properties that vary in the direction of displacement. We assume that the non-wetting phase (“oil”) displaces the wetting phase (“water”) and that the usual Darcy’s law applies to yield the equation

$$\phi \frac{\partial S_o}{\partial T} + q \frac{\partial}{\partial X} \left[f_o(S_o) - \frac{(\rho_w - \rho_o)}{\mu_w q} g_x k_{rw}(S_o) f_o(S_o) - \frac{k k_{rw}(S_o)}{\mu_w q} f_o(S_o) \frac{\partial P_c}{\partial X} \right] = 0 \quad (1)$$

where standard notation has been employed [10], gravity effects were included and the permeability may also be variable in space. Capillary heterogeneity is contained in the capillary term P_c and can be a result of spatial variations in wettability, pore space configuration, or pore size. All these variations are understood in a macroscopic sense, so that the continuum formulation is relevant. Although wettability heterogeneity may directly be included, here we concentrate on effects associated with pore size alone or equivalently on effects of permeability heterogeneity.

The connection between permeability and capillarity is well known. It is implicitly contained in the Laplace-Young equation ($P_c = 2\gamma \cos\theta/r$), a particularly useful form of which is the Leverett relationship

$$P_c = \frac{\gamma \cos\theta}{\sqrt{k/\phi}} J(S_w) \quad (2)$$

where θ is the contact angle, γ the interfacial tension and r a characteristic pore size. A derivation of (2) using percolation theory on random pore networks in the absence of gradients (such as induced by gravity or viscous effects) can be obtained, e.g. by following approaches such as that of Heiba et al. [11]. Additional support of the relation between permeability and capillarity can be found in [12]. Although pore space configuration and geometry may exert additional effects, they are secondary so that dimensionless quantities, such as porosity, or the functions of relative permeabilities k_{ro} , k_{rw} and capillary pressure J may be taken independent of the permeability change. We must stress, however, that the ensuing theory remains qualitatively the same even if a weak permeability dependence is ascribed to these functions. Taking a reference characteristic value k^* , the dimensionless function $\tau = \sqrt{k/k^*}$, which roughly follows the variation of an average pore size, is the relevant dimensionless variable that measures capillary heterogeneity in our problem.

In the usual dimensionless notation, $x = /L$, $t = \frac{q}{\phi L(1-S_{wr})}T$, where L is a macroscopic characteristic length (e.g. core length), denoting $u = \frac{S_o}{1-S_{wr}}$, where $M = \frac{\mu_w}{\mu_o}$ is the viscosity ratio, $D(u) = k_{rw}f_o$, $G(u) = k_{rw}Jf_o$, $H(u) = k_{rw}f_oJ'$, $\varepsilon = \frac{\gamma \cos \theta \sqrt{k^* \phi}}{L \mu_w q}$, $\varepsilon_g = \frac{k^*(\rho_w - \rho_o)g_x}{\mu_w q}$ and $\tau'(x) = d\tau(x)/dx$, equation (1) is rewritten as follows:

$$\frac{\partial u}{\partial t} + \frac{\partial}{\partial x} \left[F(u) - \varepsilon_g D(u) \tau^2(x) + \varepsilon G(u) \tau'(x) \right] = \varepsilon \frac{\partial}{\partial x} \left[\tau(x) H(u) \frac{\partial u}{\partial x} \right] \quad (3)$$

In the above, important variables are the inverse macroscopic capillary number ε , the gravity number ε_g , and the permeability heterogeneity Bond number $\tau'(x)$, the latter obtained from the heterogeneity function $\tau(x)$.

While permeability heterogeneity enters in both sides of (3), its main effect to the saturation profile is through the term $\varepsilon G(u) \tau'(x)$ on the left hand side. This novel, position-dependent addition to the fractional flow function F carries the main thrust of heterogeneity in the present context. The modification of the fractional flow is akin to the effect of gravity in homogeneous systems (compare with $\varepsilon D(u)$ in (3)), with which it mathematically coincides when τ is a linear function of position. This similarity was exploited in [6], where a modification of gradient percolation was proposed to describe slow drainage. The combination $\varepsilon \tau'(x)$ is the ratio of two length scales, the mean-field viscous correlation length, $l_c \sim \sqrt{k} C a^{-1}$, over the permeability heterogeneity length, $l_k \sim \sqrt{k} / \left(\frac{d\sqrt{k}}{dx} \right)$. Capillary heterogeneity effects are expected to be important as long as this ratio is not small. It is also worth noting that the role of capillarity (ε) in the present context is a subtle one. Theoretically, as long as sharp, step-like discontinuities exist in permeability (τ), the smallest value of ε should be sufficient to trigger a response. However, in our simulations, and certainly in natural rocks, the permeability has a bounded variation, thus the response amplitude would be expected to diminish as ε decreases.

Before we proceed with the solution of the initial-boundary value problem (3) we shall briefly discuss the steady-state solution for a linear change in τ , which is physically instructive. Consider the heterogeneity model (Figure 1a)

$$\tau = \begin{cases} \tau_- & x < x_- \\ a(x - x_-) + \tau_- & x_- < x < x_+ \\ \tau_+ & x > x_+ \end{cases} \quad (4)$$

where $a = (\tau_+ - \tau_-)/\Delta x$, $\Delta x = x_+ - x_-$, and subscripts $+$ and $-$ refer to downstream and upstream regions, respectively. Then, the steady state version of (3) is

$$F(u) + \lambda G(u) - \varepsilon \tau(x) H(u) \frac{du}{dx} = F_i \quad (5)$$

where we defined $\lambda = a\varepsilon$, the gravity term has been omitted for simplicity and F_i is the flow fraction of the non-wetting phase in the vicinity of the permeability jump (where F_i could be different from its inlet value). Equation (5) was solved in a self-consistent manner in [7] and [10]. It was found that for decreasing permeability ($a < 0$) the saturation builds up from its upstream value u_i ($F(u_i) = F_i$) in the high permeability region, $x < x_-$, to a maximum value u_- (at $x = x_-$), before it declines continuously inside the region of permeability variation ($x_- < x < x_+$) to its asymptotic value u_i (Figure 1b). For $x \leq x_-$ the saturation remains constant at u_i until another heterogeneous region is encountered. For increasing permeability ($a > 0$), a similar analysis leads to a “mirror image” of the previous (rightmost part of Figure 1b). (Yortsos and Chang, 1990). The physical interpretation is that for the non-wetting fluid to flow at steady conditions in a progressively lower permeability medium its saturation must decrease in accordance with capillary continuity. Since, however, in the continuum theory both asymptotic saturations must be equal, this can only be accomplished if the non-wetting fluid builds up a high enough saturation in the high permeability part. One should note the difference with the picture dictated from the static capillary continuity criterion, which requires a step-like change in saturation across the jump from high permeability to low permeability. Both the amplitude $\Delta u = |u_- - u_i|$ of the saturation peak and its width in the $x < x_-$ zone are increasing functions of λ , which can be rearranged to read

$$\lambda = \frac{\gamma \sqrt{\phi}}{\mu_o q} \left(\frac{\sqrt{k_+} - \sqrt{k_-}}{2\Delta X} \right) \quad (6)$$

Similar results are also obtained when the injected fluid is wetting but only under conditions of primary imbibition, which involves meniscus displacement. Secondary imbibition proceeds by the additional mechanisms of film flow and may not be amenable to the simple continuum description above. Finally, we should note that for the case of drainage, where $F_i = 1$, Δu should vanish at a decreasing permeability jump, contrary to an increasing jump where Δu should attain a maximum value. The steady-state predictions were tested experimentally in [7] for the case of drainage and found to be in good agreement.

Figure 1: Schematic saturation response to heterogeneity in permeability

To test the continuum theory under the general unsteady-state conditions, we performed displacement experiments on model porous media, which consisted of slices of packed glass beads of different sizes, thus different permeabilities, and we simulated the process using pore network simulation. In this paper we shall focus on drainage processes only. Due to hyperdiffusion phenomena [13] secondary imbibition displays qualitatively different features and must be analyzed separately.

3 EXPERIMENTS

An acoustic technique [14] was used to determine the saturation profile inside the porous medium. Saturation measurements were derived from the velocity variations of a sound wave along the sample (of length $L = 30 \text{ cm}$ and cross-sectional area $A = 4 \times 2 \text{ cm}^2$). From the calibration curve [15] and the accuracy of the relative velocity measurements (10^{-4}), the overall resolution in saturation is estimated at 1%. The acoustic device has been improved so that we can obtain saturation profiles along the main core direction at different times t with a spatial resolution of 2 mm (150 measurements along the length of the sample) in order to track precisely the saturation variations.

The porous media consists of packs of glass beads of different diameter d . The sample was patiently filled with slices of various diameters of glass beads, the close packing ensured by shaking after each filling. We describe experiments on six-slice samples of alternating permeability (large/small) $k_l/k_s/k_l/k_s/k_l/k_s$, each slice of 5 cm thickness. To impart the desired heterogeneity, two bead sizes of diameters $d_s = 50 \text{ }\mu\text{m}$ and $d_l = 180 \text{ }\mu\text{m}$, respectively, were used. Two other diameter pairs ($80, 280 \text{ }\mu\text{m}$) and ($50, 90 \text{ }\mu\text{m}$) were also used. The measured permeability was found in agreement with the Kozeny-Carman value $k = \frac{\phi^3 d^2}{180(1-\phi)^2}$ for the porosity value $\phi \simeq 0.4$. The permeability map of the sample (corresponding to simple jumps) is sketched in Figure 2. The fluids used were water as a wetting fluid ($\mu_w = 1 \times 10^{-3} \text{ Pa.s}$, $\rho_w = 10^3 \text{ kgm}^{-3}$) and nonane as the non-wetting fluid ($\mu_0 = 0.7 \times 10^{-3} \text{ Pa.s}$, $\rho_0 = 780 \text{ kgm}^{-3}$), with interfacial tension $\gamma = 25 \text{ mJm}^{-2}$ and $\cos\theta = 1$. Experiments corresponded to primary drainage processes. The sample was initially saturated with water and was subject to injection of the lighter fluid (oil) in the direction from top to bottom. The flow rate q varied from 0.3 cm/h to 150 cm/h corresponding to a wide variation in

Figure 2: Permeability map of experimental set up

the capillary number values, from 3×10^{-8} to 1.5×10^{-4} . We investigated six different flow rates (0.3, 6, 15, 30, 60 and 150 cm/h) corresponding to the following ϵ values (700, 35, 14, 7, 3.5, and 1.4). The data are shown as oil saturation versus reduced sample length at different times, expressed in terms of the pore volume fraction t .

Figure 3 shows two such profiles for various values of time. First, we shall comment on the qualitative features of the experimental results. As the displacement progresses, it is affected substantially by the heterogeneities encountered. The profiles rather rapidly assume a steady-state to which they become locked, suggesting that the steady-state interpretation should be of relevance. The experiments show that a large amount of the wetting-phase remains trapped behind regions of permeability increase. Although the degree of trapping and the profiles themselves are affected by the capillary number, several features are common at steady-state: Sharp saturation jumps (within the experimental resolution of 2 mm) occur at the positions of permeability changes (corresponding to $\xi = 0.17$, 0.55 and 0.88 for a permeability decrease and to $\xi = 0.35$, 0.72 and 1.00 for a permeability increase, respectively, in the schematic of Figure 2). In the regions of high permeability the saturation profile becomes approximately flat and assumes a high value, without displaying any remarkable features. By contrast, significant dips in the profile occur in the

Figure 3: Experimental saturation profiles at various times (a) $\varepsilon = 35.3$, (b) $\varepsilon = 3.53$

Figure 4: Saturation jump vs. ϵ

low permeability regions, with the saturation decreasing continuously before the region of sharp permeability increase is encountered. The size of the dip ΔS varies with the flow rate, decreasing with an increase in the latter. Heterogeneity effects are mitigated as the flow rate increases, although a weak imprint was still left at the highest numbers tested.

The features are in qualitative agreement with the previous steady-state theory, which predicts sharp changes in saturation when regions of permeability changes are traversed, a significant response of the saturation profile as a permeability change region is approached (with a downwards kink in the oil saturation before a permeability increase, such as at $\xi = 0.35$ and 0.72), an insignificant response in the high permeability region for the pure drainage case ($F_i = 1$) and the dependence of all these effects on the capillary number. For a quantitative comparison, however a numerical simulation is needed. We first simulated the steady-state results by solving (3). A particularly useful measure of the saturation response is the variation ΔS across a region of permeability increase. Plotted in Figure 4 are theoretical and experimental results for ΔS as a function of λ . Because the latter contains the unknown parameter ΔX , which is difficult to estimate a priori, this parameter is adjustable. All other functions were evaluated using the simple expressions of [10]. The theoretical predictions agree reasonably well with the experiments at all flow rates for a width

Δx of the permeability jump region equal to 0.015 (corresponding to $\Delta X = 4 \text{ mm}$). This estimate suggests that during the packing large and small sizes interpenetrated in a small region extending over a few millimeters. When the other bead size pairs were used (50, 90 μm and 180, 280 μm) the value of λ obtained was the same for constant ε , implying that the slope of the permeability jump ($a \sim (\sqrt{k_l} - \sqrt{k_s}) / \Delta X$) can be considered as constant. Since $\sqrt{k} \sim d$, ΔX must be proportional to the bead size ($\Delta X \sim d$) in order for this to be valid, in which case we can postulate $a \sim (d_+ - d_-) / (d_+ + d_-)$. This assumption was used in all our subsequent calculations.

Having determined ΔX we proceeded to simulate the transient profiles. Figure 5 shows numerical results corresponding to the experiments of Figure 3. Despite the simplicity of the model, in particular as it regards the relative permeability and capillary pressure expressions, which were not independently evaluated, quite a reasonable agreement between simulation and experiment is found both for steady-state and transient profiles. At this point it should be remarked that as long as the relative permeability and capillary pressure functions were approximated by expressions qualitatively consistent with percolation trends the sensitivity to the particular shape of the functions was weak. The simulations reproduced all large-scale features of the experiments at all capillary numbers tested. Certain small scale features were not successfully reproduced, either because of device artifacts due to local porosity or because of wettability variations, which also affect capillarity. However, such fine-tuning can be performed, if desired. In particular, it should be possible to relate saturation to permeability fluctuations, and thus to extract statistical characteristics of the permeability field, particularly as it regards fractal statistics and correlations. This possibility is currently under investigation.

4 PORE NETWORK SIMULATIONS

The good agreement between experiments and the simple continuum model is somewhat unexpected. Capillary heterogeneity effects are significant at low values of the capillary number (large values of λ). During transient displacement, the advancing front is dominated by capillary effects leading to fractal structures, where continuum models are least likely to apply. Since the ultimate saturation distribution is affected by the transient process, it is interesting to inquire whether the

Figure 5: Numerical simulations for the conditions of Figure 3

continuum predictions can also be supported by a pore-network level analysis, where the advance of the interfaces is followed from pore to pore. To accomplish this task, a drainage pore network simulator was developed.

The basic structure of the simulator parallels that of Blunt et al. [16] and it is based on the following premises and rules. Interfaces between the two phases reside only in pore bodies (sites), which can be partially or fully occupied. In the former case the capillary pressure across the interface is set to zero. After it fully occupies a site, the interface can move to an adjacent site provided that the capillary pressure threshold of the connecting pore throat (bond) is exceeded. If many sites on the front are partially occupied, the one to be filled next is that with the minimum time required for pore filling. Pressure drops during fluid flow are only associated with pore throats, where a Poiseuille-type law is used to relate pressure drop to flow rates. An *SOR* technique was used to calculate pressure fields in the two fluids. The main difference between the present model and that of [16] is that the simulator solved displacement at constant flow rate, rather than constant pressure drop, a process which required additional iteration of the flow field. Often, and particularly at low capillary number values, the interfaces of the front would lock at all places and convergence could not be reached. In such cases, we followed Invasion Percolation rules and advanced the interface neighboring the largest pore. Simulations were carried in 2-D square lattices of size 100×21 and 100×60 , for capillary number values ranging between 1.5×10^{-6} and 1.5×10^{-3} , three different values of the viscosity ratio ($M = 0.1, 1$ and 10) and for various forms of heterogeneity. Because of the connection between permeability and average pore size, the simulation of permeability heterogeneity can be accomplished by a variation of the pore size. The latter included a sudden increase or decrease in the average pore size in the direction of displacement by a factor of 2 or 4, a smooth variation from one zone to another, as well as a sinusoidal variation. Pore sizes were randomly and uniformly distributed around their mean value. Typical schematics of the lattices simulated are shown in Figure 6, where the pore sizes are exaggerated. It should be pointed out that because of the small size of the network (100×21), strong finite size effects in the form of strong fluctuations in the saturation profile exist in our simulations. These effects persist for the case of a homogeneous sample as well, where the average pore size is constant, but tend to diminish as the size or the capillary number increase.

We shall concentrate mostly on simulations with equal viscosities ($M = 1$). Simulations results

(a)

(b)

(c)

(d)

Figure 6: Typical lattices used in pore network simulations

for a pore size ramp increase of five lattice spaces are shown in Figures 7 and 8 for two different values of the capillary number. A response similar to that predicted by the continuum model and observed in the experiments, namely the reduction of the non-wetting phase saturation as the permeability increase region is approached, followed by a sharp increase, is evident. The downwards kink diminishes as the capillary number increases, although not as smoothly as in the experiments, mostly we suspect as a result of finite size effects. Also, because of the overlap with the boundary end effect at the outlet of the lattice, the saturation level does not rise, following the jump, to the high level observed in either experiments or continuum simulations. A better agreement was found in the simulations involving a larger lattice, however. In all cases, the response diminished substantially as the capillary number increased to levels higher than 10^{-3} . To illustrate the effect of heterogeneity (λ) on the profile, we simulated a heterogeneous system where the ramp extent is 20 lattice spacings (Figure 9). It is apparent that the effect of heterogeneity here is all but lost. Certain differences with the “homogeneous” case (Figure 10) do exist, however, as a result of the differences in viscous effects because of the different permeabilities in the two regions. These viscous effects exist in our previous simulations as well, and can partly explain the not so satisfactory match of the saturation response with the continuum theory or the experiments downstream of the heterogeneity. We expect that this difference would diminish and eventually disappear as the lattice size is increased.

Simulations involving a permeability decrease were also conducted. Figure 11 shows results obtained for a decreasing ramp of 5 lattice spacings. The response is qualitatively similar to the theoretically expected. The non-wetting saturation rises as the permeability decline region is approached and subsequently decreases substantially as the low permeability region is entered. The upwards kink in the saturation profile predicted from the theory is thus observed. As in the previous case, the effect of increasing capillary numbers is to smooth out the heterogeneity. The combined effects of alternating regions of increasing and decreasing permeability can be quite dramatic. Figure 12 shows pore network simulations with a sinusoidal variation in the average pore size of a wavelength equal to $1/5$ of the system length. According to the continuum theory [10], the saturation response should also oscillate with a slight phase lag depending on the capillary number. Both predictions are clearly observed in the pore network simulations. We must point out that similar effects were recently observed in a related simulation on multifractal lattices [17].

Figure 7: Saturation profile and displacement patterns for permeability increase (1:2) and for $Ca = 1.5 \times 10^{-5}$

Figure 8: Saturation profile and displacement patterns for permeability increase (1:2) and for $Ca = 1.5 \times 10^{-4}$

Figure 9: Saturation profile and displacement patterns for permeability increase (1:2) in a ramp of 20 lattice spacings for $C'a = 1.5 \times 10^{-4}$

Figure 10: Saturation profile and displacement patterns for uniform permeability and $Ca = 1.5 \times 10^{-4}$

Figure 11: Saturation profile and displacement patterns for permeability decrease 1.5×10^{-5}

Figure 12: Saturation profile and displacement patterns for sinusoidal

We conclude by presenting two illustrative examples of the effect of the viscosity ratio. Figure 13 shows saturation profiles corresponding to a sudden permeability increase in the middle of the sample at a favorable viscosity ratio, $M = 0.1$. Because viscous effects are here favorable, the homogeneous displacement is compact in all regions, leading to a high efficiency and high saturation values. As the region of permeability increase is encountered, the corresponding saturation kink is very small and the capillary effect is insignificant. In contrast, for unfavorable viscosity ratio values ($M = 10$), the displacement is much less efficient and heterogeneity effects are quite noticeable in the saturation profile (Figure 14). Both effects of M were anticipated in the continuum study of Chang and Yortsos [10].

To interpret the pore network simulation results, it is necessary to consider the application of percolation theory. Let us first discuss invasion percolation (IP), namely displacement at constant rate at low Ca , in a medium involving a sudden decrease in permeability at the boundary of two regions 1 and 2. According to the IP rules, the front advances by occupying the largest pore available to it. Before the front reaches the boundary, the pattern is given by percolation corrected by finite size effects (Figure 15a). In the upstream region 1 of high permeability, pores of size greater than r_d can be penetrated, where r_d is another measure of the capillary pressure level, $P_c = \frac{2\gamma}{r_d}$. The corresponding percolation probability p_1 is related to the pore distribution in this region, $\alpha_1(r)$, by

$$p_1 = \int_{r_d}^{\infty} \alpha_1(r) dr \quad (7)$$

Given the width of the lattice W , p_1 can be obtained from finite size scaling which involves the percolation threshold of the lattice p_c

$$|p_1 - p_c|^{-\nu} \sim W \quad (8)$$

The extent of the fractal regime is quenched by the width of the lattice, which sets the characteristic length of the system. The corresponding saturation in region 1 satisfies the percolation scaling

$$S_1 \sim (p_1 - p_c)^\beta \quad (9)$$

Upon encountering the boundary between the two regions, the front would penetrate only a small

Figure 13: Saturation profile and displacement patterns for the conditions of Figure 8 and $M = 10$

Figure 14: Saturation profile and displacement patterns for the condition of Figure 8 and $M = 0.1$

part of the low permeability region 2 (Figure 15b). The extent is determined from the corresponding percolation probability

$$p_2 = \int_{r_d}^{\infty} \alpha_2(r) dr \quad (10)$$

which is related to p_1 via (7). Clearly $p_2 < p_1$ for all r_d . The extent of frontal advance in region 2 satisfies the scaling law

$$l_2 \sim |p_2 - p_c|^\nu \quad (11)$$

As long as $l_2 < W$, penetration of region 2 is limited to a short tail. As the displacement proceeds, the probability p_1 and the saturation of region 1 steadily increase with only small changes in S_2 and l_2 (Figure 15c-15d). Percolation of region 2 is reached (Figure 15e) when $l_2 \sim W$, which occurs at the critical value p_2^* , where

$$|p_2^* - p_c|^{-\nu} \sim W \quad (12)$$

Upon percolation, subsequent increase in the saturation of either region requires that either another region of lower permeability is encountered (for example if the outlet end is a semi-permeable membrane) or that viscous effects are significant to provide the necessary pressure increase (see also below). The relationship between the saturation values S_1 and S_2 at any time requires knowledge of the size distribution functions α_1 and α_2

$$\frac{S_2}{S_1} = \left(\frac{p_2 - p_c}{p_1 - p_c} \right)^\beta \quad (13)$$

Because the capillary pressure is also related to r_d , the above is another representation of the Leverett J-function. It is apparent that in this case the Leverett J-function formalism remains valid and that the continuum approach should be applicable as long as W is large enough. The events described above can also be interpreted in terms of (stable) gradient percolation but at an infinitely large Bond number, where front widths (or tails) are of infinitesimal extent.

Consider next IP in a medium involving a sharp increase in permeability (Figure 16). The percolation of region 2 begins immediately when the front contacts the heterogeneity, at which time p_1 and S_1 are still given by (7) and (9). However, S_2 will not follow (13). Instead, as long

(a)

(b)

(c)

(d)

(e)

(f)

Figure 15: Invasion percolation with an abrupt decrease (2:1) in permeability at various stages

(a)

(b)

(c)

(d)

(e)

(f)

Figure 16: Invasion percolation with an abrupt increases (1:2) in permeability and at various stages

as there is no subsequent region of lower permeability downstream or viscous pressure drops are not accounted for, the probability percolation in region 2 remains fixed at the value p_2^* given by (12), implying that the two saturations S_1 and S_2 are momentarily the same. Capillary pressure continuity can only be enforced in region 1 by a re-adjustment and partial withdrawal of its menisci towards pore bodies, to bring about a drop in the capillary pressure and a possible (although small) drop in the saturation S_1 . During this transient, which however depends strongly on W , we expect the Leverett J-function formalism to fail. It is only after contact with a lower permeability region or if some viscous pressure drops develop, and only after the capillary pressure has increased to the original level to satisfy (8), that further penetration of region 1 takes place. This problem is somewhat analogous to capillary fingering in unstable gradient percolation, where percolation fractal regimes exist throughout the process. Figure 16 shows the sequence of events in this case.

The preceding suggests the possibility of hysteresis in the capillary pressure-saturation diagram, depending on the direction of displacement, namely whether it is from high-to-low permeability or inversely, and the possibility that the Leverett formalism is only conditionally valid. Fortunately, this would be the case only if finite size effects are strong (small values of W). Indeed, while the displacement pattern follows the schematics of Figures 15 and 16, the transients where continuum approaches would fail correspond to fractal regimes where saturation values are expected to be negligibly small for large W . For such cases, the saturation in the low permeability region 1 would remain infinitesimally small before its capillary pressure threshold is reached, and would increase thereafter following the typical continuum approach. We conclude that hysteresis, if it exists, is infinitesimal for large enough widths W , thus the Leverett J-function formalism applies locally in each permeability region regardless of the position of the permeability block or the direction of displacement (as also assumed in our recent study of large scale percolation of drainage [8]).

The previous analysis provided a description of the displacement under static conditions. Consider, next, displacement in the presence of viscous effects, which introduce a viscous correlation length ξ_ν . In a displacement in a homogeneous region, ξ_ν measures the extent upstream of the front where fractal concepts apply, and beyond which continuum descriptions are applicable. Two different expressions for the dependence of ξ_ν on Ca were developed by Wilkinson [18], and by Blunt et al. [19], independent of the viscosity ratio M . In a separate publication we shall consider the dependence of ξ_ν on Ca and M following Lenormand's work [20], where it will be shown

that the two different cases $M \ll 1$ and $M = O(1)$ or higher need to be considered. We find $\left(\frac{\xi_\nu}{r^*}\right) \sim Ca^{-\frac{\nu}{1+t+\nu}}$ for $M \ll 1$, and $\left(\frac{\xi_\nu}{r^*}\right) \sim Ca^{-\frac{\nu}{1+\nu}}$ for $M = O(1)$ or higher, where t and ν are conductivity and correlation length percolation exponents and r^* the average pore size. These expressions differ from those in either [18] or [19].

Consider, first, the problem involving a sudden permeability decrease. Here, an analysis similar to the previous applies, with the important exception that the variable r_d is now space-dependent. For convenience, we shall use the boundary position as the origin of the space coordinate, which increases in the direction of displacement. Clearly, r_d is an increasing function of x . When the front reaches the boundary, the extent of the frontal advance into region 2 is still given by (11), where p_2 is evaluated at $r_d(0)$. As previously, penetration would be insignificant, until condition (12) is reached. Before this, the displacement is confined in the region 1 of high permeability, and the heterogeneity boundary behaves as a no-flow boundary for the non-wetting phase. Because of the insignificant frontal advance, the only possibility for further displacement is by lateral advancement of the menisci in region 1. The result is that the non-wetting pressure becomes almost uniform at least in the (percolation) region near the boundary

$$p_o(-\xi_\nu, t) \sim p_o(0, t) \quad (14)$$

while, at the same time, the wetting phase continues to flow

$$p_w(-\xi_\nu, t) - p_w(0, t) \sim q\xi_\nu\mu_w \quad (15)$$

A capillary pressure gradient rapidly develops

$$P_c(0, t) - P_c(-\xi_\nu, t) \sim q\xi_\nu\mu_w \quad (16)$$

The existence of a capillary pressure gradient in the direction opposite to the viscous pressure gradient and flow causes the non-wetting saturation to increase preferentially in the region closest to the heterogeneity boundary, thus the initially concave saturation profile progressively changes curvature and approaches the shape of an upwards kink. This process continues until percolation of region 2 is reached, namely when condition (12) is satisfied. Because the capillary pressure gradient is in a direction opposite to viscous pressure drop, the regime ceases to be fractal the

moment equation (16) becomes in effect, which roughly corresponds to the time the front reaches the boundary for the first time. It follows that soon after the boundary is encountered the macroscopic equations can reasonably well predict the subsequent saturation evolution. Thus, the extent of the region of the saturation response to the heterogeneity would scale with the mean field result Ca^{-1} rather than with ξ_ν . The effect of M can be similarly interpreted. For favorable ratio, $M \gg 1$, the saturation pattern following the fractal regime is much more compact resembling a piston-like displacement, thus any saturation response to the heterogeneity has no room to develop and the profile acquires, no discernible features. By contrast, when $M \ll 1$, the pattern is much sparser, thus the saturation response is strong and lasts for a considerable time.

The situation is qualitatively different in the case of a sudden decrease in permeability. Upon encounter of the heterogeneity boundary, penetration of region 2 commences immediately, much like in the case where viscous effects are absent. However, because of the viscous pressure drops, saturation profiles are now concave and not flat. As a result, the upstream saturation in region 2 (and particularly the region on the downstream side of the heterogeneity boundary) begins to rise following the rules of a regular displacement. Until such time that the capillary pressure at the boundary, $P_c(0, t)$, has reached the percolation level for region 1 (equation (8) with p_1 evaluated at $r_d(0)$), no substantial change in the saturation in region 1 takes place. The familiar downwards kink of the saturation response thus emerges. During the period of a frozen saturation in region 1, a re-adjustment of the menisci may be necessary to balance the lower capillary pressure experienced during the invasion of the high permeability region 2. Neither the continuum nor the pore network model capture this feature. However, made by both models by excluding the menisci movement are not significant. In particular, the two models do predict the downwards kink in the saturation profile. Upon the subsequent onset of penetration of region 1 at the point near the boundary, the fractal regime ceases to exist and a continuum description should be a reasonably good alternative. As before, the spatial extent of the dip in the profile should scale inversely proportional to Ca . If the permeability contrast is so high that the percolation threshold for region 1 cannot be reached even when region 2 becomes fully occupied, the spatial extent of the dip should also scale as Ca^{-1} , provided that region 1 is long enough to accomodate it. Steady-state profiles at variable rates across a discontinuity were used in the experimental study of [7]. The effect of M is similar to the previous case.

CONCLUSIONS

In this paper, experiments and pore network simulations were used to study primary drainage in a heterogeneous porous medium involving changes in the permeability in the direction of displacement and to test the validity of macroscopic, continuum models. Both experiments and pore level model confirmed the validity of the macroscopic predictions, at least as far as the basic features of the saturation response is concerned. An analysis of the results based on percolation arguments provided additional support to the adequacy of continuum models in describing the response of the saturation to the heterogeneity. It should be pointed out that these results apply only for drainage and not for imbibition, which is qualitatively different due to the different flow mechanisms. In particular, for secondary imbibition, the standard continuum models do not reproduce well preliminary experiments suggesting that improved continuum models are needed. This subject is currently under study. The sensitivity of the saturation to heterogeneity at low displacement rates suggests the possibility of heterogeneity identification from saturation maps. This idea, originally suggested in [10], is also currently under investigation.

ACKNOWLEDGEMENTS

This work was partly supported by NATO travel grand CRG 90 L053. MC, NR and DS would like to acknowledge support by GDR CNRS *Physique des Milieux Heterogenes Complex*. BX and YCY were supported by DOE Contract DE-FG22-90BC14600. All these sources of support are gratefully acknowledged.

REFERENCES

1. Lake, L.W. and Carroll, H.B. jr. (Editors), *Reservoir Characterization* (Academic Press, Orlando, FL.) 1986; Feder, J., *Fractals* (Plenum Press, New York, N.Y.) 1988.
2. Lenormand, R., *J. Phys.: Condens. Matter* **2** (1990) SA79.
3. Waggoner, J.R., Castillo, J.L. and Lake, L.W., paper SPE 21237 presented at the 11th SPE Symposium on Reservoir Simulation, Anaheim, CA Feb. 17-20 (1991).
4. Sorbie, K.S., Feghi, F., Pickup, G.E., Ringrose, P.S. and Jensen, J.L., proc. ECMOR III, Delft, Holland, 17-19 June (1992).

5. Van Batenburg, D.W., Bruining, J., Bakken, H. and Palmgren, L.T.S., paper SPE 22588 presented at the 1991 SPE Annual Fall Meeting, Dallas, TX Oct 6-9 (1991).
6. Chaouche, M., Rakotomalala, N., Salin, D., Xu, B. and Yortsos, Y.C., *Phys. Rev. Lett.*, submitted (1993).
7. Chaouche, M., Rakotomalala, N., Salin, D. and Yortsos, Y.C., *Europhys. Lett.*, **21**, (1993) 19.
8. Yortsos, Y.C., Satik, C., Bacri, J.-C. and Salin, D., *Trans. Porous Media* in press (1993).
9. Satik, C. and Yortsos, Y.C., paper presented at the 1991 AIChE Annual Meeting, Los Angeles, CA (1991).
10. Chang, J. and Yortsos, Y.C., *Trans. Porous Media*, **5** (1990); Stubos, A.K., Satik, C. and Yortsos, Y.C., *Int. J. Heat Transf.*, in press (1993); Chang, J. and Yortsos, Y.C., *SPE Res. Eng.*, (1992) 285.
11. Heiba, A.A., Sahimi, M., Scriven, L.E. and Davis, H.T., paper SPE 11015 presented at the 57th Annual Technical Conference and Exhibition of the Society of Petroleum Engineers, New Orleans, LA, Sept., 26-29 (1982).
12. Katz, A.J. and Thompson, A.H., *Phys. Rev. B* **34** (1987) 8175.
13. Bacri, J.-C., Leygnac, C. and Salin, D., *J. Physique Lett.* **46** (1985) L467; Bacri, J.-C., Rosen, M. and Salin, D., *Europhys Lett.*, **11** (1990) 127.
14. Bacri, J.-C., Hoyos, M., Lenormand, R., Rakotomalala, N.R., Soucemariadin, A. and Salin, D., *J. Phys. III*, **1** (1991) 1455.
15. Bacri, J.-C. and Salin, D., *Geophys. Res. Lett.*, **13** (1986) 326.
16. Blunt, M. and King, P.R., *Transp. Porous Media*, **6** (1991) 407.
17. Lenormand, R., Kalaydjian, F., Bieber, M.T. and Lombard, J.M., paper SPE 20475 presented at the 1990 SPE Annual Fall Meeting, New Orleans, LA, Sept (1990).
18. Wilkinson, D., *Phys. Rev. A* **30** (1984) 520.

19. Blunt, M., King, M.J. and Sher, H., preprint (1991).
20. Lenormand, R., Proc. Roy. Soc. London, Ser. A, **423** (1989) 159.

END

**DATE
FILMED**

6 / 17 / 93