015027

1 OF 1

PROGRESS REPORT

DE92 015027

"Regulation of Cell Division in Higher Plants"**Grant No. DE-FG02-90ER20008**

Two reprints and one manuscript-in-preparation are attached. These are briefly summarized below, followed by a description of more recent work. Some of the results reported in the first two manuscripts were obtained while the application for the present grant was pending.

Feiler HS and TW Jacobs (1990) Cell division in higher plants - a *cdc2* gene, its 34-kDa product, and histone-H1 kinase activity in pea. Proc Nat Acad Sci (USA) 87:5397-5401.

In this paper, we described the existence of a high molecular weight histone H1 kinase activity from etiolated pea epicotyl hooks, as well as immunological evidence for the existence of a p34^{cdc2} kinase in a variety of higher plants. The pea p34^{cdc2} seemed to consist of a multiplicity of forms around 34 kDa. While we suggested at the time that this polymorphism was consistent with multiple phosphorylated forms of pea p34, more recent work in animal systems leads us to include now the possibility that these may represent the products of multiple *cdc2*-like genes (see below). Work in progress is aimed at resolving this uncertainty. We also reported the amino acid sequence of our PCR-cloned pea *cdc2* gene product. This paper represented the first biochemical evidence for an MPF-like complex in a higher plant.

Feiler H, Jacobs T (1991) Cloning of the pea *cdc2* homologue by efficient immunological screening of PCR products. Plant Mol Biol 17:321-333

This paper describes the details and results of a novel method we developed for positively identifying *cdc2* PCR products from pea. The method relied on a p34^{cdc2}-specific monoclonal antibody (from David Beach at Cold Spring Harbor Laboratory) for screening expressed PCR fragments produced from degenerate *cdc2* primers and plant cDNA templates. The method was 100% efficient, permitting us to distinguish between what appears to be an authentic pea *cdc2* gene and a *cdc2*-like gene from this species. Reports from a number of plant and animal laboratories have since confirmed that higher eukaryotic genomes carry multiple *cdc2*-like genes. The current nomenclature identifies the products of these *cdc2*-like genes as "CDK2's", for "Cyclin-Dependent Kinase 2" (the authentic *cdc2* being CDK1). Further structural and functional analyses are needed in order to determine whether either our pea *cdc2* or *cdc2*-like genes actually encode cyclin-dependent protein kinases (see below). We also showed in this paper that, at relatively high stringency, our partial pea *cdc2* clone appeared to represent a maximum of two genes in the pea genome.

Prewett ME, HS Feiler H and TW Jacobs (1992) Developmental expression of MPF in pea (*in preparation for PLANT CELL*)

We have begun to ask some developmental questions regarding the distribution of *cdc2* transcripts and other aspects of the MPF complex in peas. In this manuscript, for which a final increment of data (see below) and round(s) of revision are still lacking, we describe a developmental survey of pea *cdc2* and cyclin transcript levels as well as a parallel characterization of the levels of histone H1 kinase activity. In order to accurately quantify pea *cdc2* transcript levels, we used the PCR-generated partial cDNA clone in RNase protection experiments. We surveyed a variety of

tissues and found that levels of *cdc2* mRNA per microgram total RNA closely reflect the mitotic state of the tissues: etiolated stem and root tips have more, roots and stems have less, leaves and tendrils hardly any at all, and seeds, none detectable (Figure 2). It is likely that, were we to dissect out germinating embryos from imbibed seeds, that we might see a signal above the massive null background of the cotyledon. This finding presents an interesting cellular/developmental challenge: What is the mechanism by which *cdc2* transcription is turned off during seed (esp. cotyledon) maturation, since one can assume that it is probably *on* when cotyledonary cells are dividing during seed maturation?

We used PCR to obtain a partial pea cyclin clone. Primary structural characterization of this gene and its product await recovery of a full length clone (see below). Once again, we found that mitotic tissues display more cyclin transcript, on a per microgram total RNA basis, than their less-mitotic counterparts (Figure 3).

We used precipitation with p13-Sepharose as a means of affinity purifying the pea "MPF" complex from a variety of tissues. This method has become the industry standard for p34^{cdc2} kinase assays and is based on the natural affinity of p13, the fission yeast cell cycle regulator, for the p34-cyclin complex. When this matrix was used to precipitate histone H1 kinase activity from a variety of pea tissues, a result consistent with that described above was obtained: activity was concentrated in mitotic tissues, when activity was expressed on a per microgram protein basis (Figure 4).

In order to refine this analysis and eliminate developmental differences from one tissue to the next, we selected the 10 cm pea root tip as a developmental gradient along which to investigate pea MPF activity and its components. Five successive 2 mm segments from the pea root tip are collected and pooled from a population of pea roots. The meristem proper is located in the second segment (2-4 mm from the tip). We were surprised to find that the levels of *cdc2* transcript were relatively constant along the root gradient (Figure 5), on a per cell basis, whereas the pea cyclin message was much more abundant, on a per cell basis in the meristematic zone of the root tip. The cyclin profile was closely paralleled by the p13-precipitable histone H1 kinase activity along the root tip gradient (Figure 7). These results suggest that, within the pea root tip developmental profile, cyclin and p34^{cdc2} kinase activity reflect -- and possible control -- mitotic activity of cells, whereas the level of *cdc2* transcription is relatively unregulated on a per cell basis. This may reflect the potential for pea root cells to re-enter the proliferative state upon stimulation by *Rhizobium* or other developmental programs such as lateral root emergence or cambial growth.

Up to now, we have characterized *cdc2* and cyclin message levels in addition to p34^{cdc2} kinase enzyme activity. Data on levels of p34^{cdc2} protein are still lacking and very much needed. The original antibody used to identify this clone (see above) is no longer available and only recently have we generated antisera which can now be used for this assay (see below). It must be noted, however, that any conclusions drawn from this experiment's data, as well as those described above, must be qualified. Recent reports from animal and yeast cell cycle systems suggests that multiple *cdc2* and cyclin genes within an organism are the rule, and that our experiments to date are but a vignette of much larger, more complex picture. Until we can verify the function of our clones by yeast complementation tests, we cannot make any assumptions about the *in vivo* role of their gene products. And until we have full-length cDNA clones, we cannot perform such tests. Nonetheless, preliminary quantitative immunoblotting experiments suggest that the level of p34^{cdc2} protein in our system tends to follow the pattern of its enzyme activity, both along the root tip gradient and in the tissue/organ survey.

Recent Work

Functional characterization of clones. The accepted diagnostic for the authenticity of a *cdc2* or cyclin clone is a complementation test of the cognate yeast mutant. We have tried unsuccessfully to

complement a temperature sensitive fission yeast *cdc2* mutant a number of times, under a variety of temperature conditions, with the chimeric yeast/pea *cdc2* gene which was the product of our *cdc2* cloning/verification scheme (described in the second manuscript, above). We can only conclude from this negative result that this chimeric construct does not function in yeast. However, until we have recovered a full-length version of this pea cDNA, of which our PCR clone constitutes about 60%, we can draw no firm conclusions about this sequence's role in pea, despite its convincing primary sequence character. It should also be noted that not all heterologous *cdc2* genes, when introduced into mutant fission yeast, are capable of complementation. The past year's revelations of a large family of *cdc2*-like genes in essentially all higher eukaryotes surveyed suggests that this protein kinase motif has proliferated to fill a number of cellular roles, only some of which may involve the regulation of the cell cycle.

Pea cyclins and cdc25. As mentioned above, we obtained a ~200 bp PCR clone which appears to represent part of a pea cyclin cDNA. The original PCR products from this experiment included a diversity of fragment sizes, and Houqi Chen, the postdoctoral associate on this project, has pursued these signals further. Recently, Dr. Chen has obtained two more pea cyclin PCR clones, bringing our total to three. Thus, we now have a handle on at least some of peas's diversity of these important regulatory subunits. We await longer clones for further characterization and publication of these genes and their products. Both my graduate student Heidi Feiler and Dr. Chen have labored heroically to obtain a PCR clone of a pea *cdc25* homologue, but to no avail. Heidi, in particular, sequenced upwards of 30 PCR clones, none of which encoded any homology to published *cdc25* proteins. It may well be that this step in the regulatory network (the phosphatase which dephosphorylates p34^{cdc2}, thus triggering M-phase) is a point of evolutionary departure among yeasts, animals and plants. However, some broadly crossreactive antibodies against evolutionarily distant *cdc25* phosphatases were recently reported and we will try to obtain these for screening our (awaited, see below) pea cDNA expression library, should a pea root tip immunoblot look promising with the antisera.

Obtaining full-length clones. Published reports of large families of both *cdc2*-like genes as well as cyclins in higher eukaryotes restrict the breadth of conclusions which can be drawn from the data presented in the manuscript in preparation, above. In these studies, we have assayed the levels of but one *cdc2* transcript and one cyclin transcript, among the many which are likely to be present in these tissues. In addition, we have used very generic reagents for assaying p34^{cdc2} kinase protein and enzyme activity levels (an anti-chimeric pea/yeast p34^{cdc2} antibody and p13-Sepharose, respectively, both of which probably detect/precipitate multiple forms). We have therefore recently placed a high priority on obtaining gene- and protein-specific probes for all of the *cdc2* and cyclin PCR clones we have obtained. These would be, respectively, DNA sequences representing the 3'-untranslated portions of cDNAs and anti-peptide antibodies directed against, most probably, the carboxy terminus of each putative gene product.

To this end, we purchased two off-the-shelf pea cDNA libraries (from Clontech) and have constructed one from root tip cDNA. The commercial libraries were screened intensively, only to yield a single short clone which constituted a subset of our existing pea *cdc2* PCR clone. Our homemade libraries have been too small to merit screening for these very low abundance cDNAs. A technical blunder at the company with whom we had contracted aborted the first of our subsequent ventures into custom-made pea root tip cDNA (expression) libraries. A second such attempt, through a more established company, is currently in progress and should be available for our screening in about three weeks. When such a library is available to us, we will have five excellent pea cDNA probes with which to screen it: a *cdc2*, a *cdc2*-like gene and three cyclins.

Immunological tools and experiments. Gene product-specific antisera are absolutely essential for progress in the kinds of experiments we wish to perform. Given the assumption now that peas probably have multiple *cdc2* and cyclin gene products, the reagent/technique of choice is immunoprecipitation with anti-peptide antibodies directed against the carboxy terminus of each putative protein. Without 3' terminal cDNA sequence information, we have not been able to raise such antisera. Instead, we have focused on raising whatever antibodies we can, given the tools available. Heidi Feiler successfully raised some very useful antisera against the chimeric pea/yeast p34^{cdc2} protein which was the gene product of the clone obtained from our PCR product screening strategy. As mentioned above, we are now using this antiserum to evaluate p34^{cdc2} levels in a pea developmental survey and along the root tip gradient.

Since p34^{cdc2} is regulated by phosphorylation on a tyrosine in animal cells and fission yeast (but not budding yeast, according to two articles in last week's Nature!), our access to this gene should provide us with an opportunity to determine whether such a regulatory strategy also occurs in plants. Such a demonstration has been elusive until now in the plant kingdom. We obtained four different anti-phosphotyrosine (P-tyr) antisera and tested each of them for ability to detect a phosphorylated tyrosine in p13-Sepharose precipitates from pea root tip extracts. No positive signals were seen. At least three interpretations can be offered for this negative result. First, the amount of phosphotyrosine present in the precipitates may have been too small to be detectable, either because of rogue phosphatases in the extract or because the extracts simply contained much more non-phosphorylated than phosphorylated p34^{cdc2}. Second, anti-phosphotyrosine antibodies are notoriously selective for specific P-tyr contexts and perhaps the pea P-tyr's in p34^{cdc2} are not appropriate for any of four employed. Third, perhaps pea p34^{cdc2} is simply not regulated by tyrosine phosphorylation. This issue will only be resolved when we can generate some high-affinity, p34^{cdc2} antibodies with which the protein can be immunoprecipitated quantitatively, for western and phosphoaminoacid analysis.

Cell Cycle Synchrony Experiments. We are pleased to report that we have been able to establish a highly reproducible pea root tip synchrony procedure, with which we have begun to observe facets of *cdc2* activity in the higher plant cell cycle. Recall that a synchronous cell system is absolutely essential for sorting out the roles of these regulators through the various stages of the eukaryotic cell cycle, G1, S, G2 and M. We were fortunate to receive a protocol from the notebooks of Jack Van't hof (Brookhaven N.L.), which he had successfully applied to pea root tips many years ago. The procedure involves arresting hydroponically-grown pea root tip meristems in S phase of cell cycle by incubating them for 12 hours in 5-aminouracil (5AU), a thymidine analog. When the drug is washed out, the root tip meristems proceed out of S-phase and through G2, M and G1 in upwards of 60% synchrony. Heidi Feiler has performed such an experiment approximately a dozen times now, collecting root tip samples at one hour time points following release from 5AU blockage, assaying their cells' DNA content by flow cytometry. M-phase appears to come at approximately 8 hours following release, and is, as always, the shortest phase. Thus, we can now obtain material, from intact plants -- as opposed to cultured plant cells -- highly enriched for any stage of the cell cycle, for biochemical and molecular analyses.

Using the the 5AU synchrony system, we have examined histone H1 kinase (= p34^{cdc2} kinase) activity in p13-Sepharose precipitates, throughout the pea cell cycle. Surprisingly at first, we observed high activity in S-phase, very low activity in G2 and G1 and moderately high activity in M. Whereas we had expected the activity to be confined to M-phase, following the animal and yeast cell paradigms, we now suspect that our generic reagent, p13-Sepharose, was probably precipitating an S-phase histone H1 kinase which may well be composed of some of the other p34^{cdc2} and cyclin family members described above. The relatively low levels in M phase cells probably reflect the brevity of this phase of the cycle and the consequent stochastic difficulty in catching cells within it,

regardless of the efficiency of the synchronization method employed. Indeed, we are examining at present another cell cycle blocker which will reportedly arrest cells in M-phase, providing an enrichment for cells at this critical juncture of the cycle.

We have also examined total cellular proteins across the pea root meristem cell cycle time course, as well as total cellular protein kinase phosphorylation products at hourly time points in synchronous populations. We were surprised and delighted to be able to detect about 7 proteins on one-dimensional protein gels, whose levels appear to be elevated in either S-phase and/or in M-phase. These are excellent candidates for cyclins and, in fact, their behavior formally classifies them as cyclins, by the original definition of the term. Of equal interest was our finding of a variety of proteins which show a very distinctive and cell cycle stage-specific pattern of phosphorylation. None of these corresponds to a known cell cycle regulatory protein. Indeed, some may be cytoskeletal elements. Nonetheless, their clarity of visualization via one-dimensional electrophoresis and autoradiography invites questions as to their identity, and no less, the identities of the protein kinases responsible for their periodic phosphorylation. Heidi Feiler is currently writing up her doctoral dissertation and is, in so doing, preparing a manuscript describing in greater detail the results obtained from her synchrony experiments.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Research Plans – 3rd Year

Our overall objectives remain unchanged since the inception of this project. Some of the details have evolved due to the many advances in studies of yeast and animal cell division controls over the past few years. Specifically, the realization that plants probably have several *cdc2* and cyclin genes and gene products now requires that we search carefully for all of these family members that we can find expressed from the pea genome and use the clones obtained to design gene-specific and protein-specific probes for the experiments originally described. The p13-Sepharose affinity matrix and generic anti-p34^{*cdc2*} kinase antibodies turn out to be less revealing reagents than we had originally anticipated.

Therefore, considerable resources are being devoted to obtaining a complete set of full-length cDNA clones for *cdc2* and cyclin genes from pea root tips. As stated above, a custom-made cDNA library is on order and five probes are available for screening it. Dr. Chen will also be performing further *cdc2* PCRs on pea cDNA in order to determine whether more expressed *cdc2* genes are present.

We have already used PCR (the "R.A.C.E." procedure) to obtain the 3' ends of our first pea *cdc2* cDNA. Once the sequence of this clone is completed, we will use the derived information for the same two purposes to be served when similar data are obtained from the rest of our clones. That is, we will have a carboxy terminal peptide synthesized, conjugate it to a carrier protein and raise antibodies against the complex. These will be affinity-purified and used to determine where in the plant, and when in the cell cycle, this protein is present. We will also use immunoprecipitation to assay the protein's phosphorylation state in tissues in which we initially find the protein to be present by immunoblotting. Finally, we will design 3'-untranslated region-specific probes in order to determine transcript levels for this and each of the other cell cycle regulators for which we obtain analogous sequence information.

In summary, we have modified and refined our plan of attack, based on the current model of cell cycle regulation and its component parts, as revealed in model yeast and animal cell systems since the inception of this project.

1992 Project Summary

Regulation of Cell Division In Higher Plants

T. Jacobs, Department of Plant Biology

Cell division is arguably the most fundamental of all developmental processes. In higher plants, mitotic activity is largely confined to foci of patterned cell divisions called meristems. From these perpetually embryonic tissues arise the plant's essential organs of light capture, support, protection and reproduction. Once an adequate understanding of plant cell mitotic regulation is attained, unprecedented opportunities will ensue for analyzing and genetically controlling diverse aspects of development, including plant architecture, leaf shape, plant height, and root depth. The mitotic cycle in a variety of model eukaryotic systems is under the control of a regulatory network of striking evolutionary conservation. Homologues of the yeast *cdc2* gene, its catalytic product, p34, and the cyclin regulatory subunits of the MPF complex have emerged as ubiquitous mitotic regulators. We have cloned *cdc2*, *cdc2*-like and cyclin genes from pea. As in other eukaryotic model systems, p34 of *Pisum sativum* is a subunit of a high molecular weight complex which binds the fission yeast p13 protein and displays histone H1 kinase activity *in vitro*. Our primary objective in this study is to gain baseline information about the regulation of this higher plant cell division control complex in non-dividing, differentiated cells as well as in synchronous and asynchronous mitotic cells. We are investigating *cdc2* and cyclin expression at the levels of protein abundance, protein phosphorylation and quaternary associations.

**DATE
FILMED
8 10 71 92**