AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

**DOE/ER/60631-11
(DE93014482)**

**RADIATION PHYSICS, BIOPHYSICS, AND RADIATION BIOLOGY
Progress Report for the Period December 1, 1992—November 30, 1993**

**By
E. J. Hall
M. Zaider**

May 1993

Work Performed Under Contract No. FG02-88ER60631

**College of Physicians & Surgeons of Columbia University
New York, New York**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from (615)576-8401.

Available to the public from the National Technical Information Service, U. S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

RADIATION PHYSICS, BIOPHYSICS, AND RADIATION BIOLOGY

Progress Report
for Period December 1, 1992 - November 30, 1993

E. J. Hall, Principal Investigator
M. Zaider, Co-Principal Investigator

*Center for Radiological Research
Department of Radiation Oncology
College of Physicians and Surgeons of Columbia University*

May 1993

Prepared for
THE U.S. DEPARTMENT OF ENERGY
UNDER GRANT NO. DE-FG02-88ER60631

TABLE OF CONTENTS

	Page
Introduction	2
Staff News	4
The Columbia Colloquium and Laboratory Seminars	5
Center for Radiological Research Staff	6
Teaching	8
RADIATION PHYSICS AND MICRODOSIMETRY	
Measurement of Fundamental Physical Parameters of a Wall-less Ultraminiature Proportional Counter P. Kliauga	10
Microdosimetry of Low Energy Photon Sources P. Kliauga	17
A Study of Excited States in Trans-Polycetylene in the Hartree-Fock, Tamm-Dancoff, and Random-Phase Approximation M.G. Vracko & M. Zaider	21
Electron Transport in Condensed Water M. Zaider, M.G. Vracko & A.Y.C. Fung	25
BIOPHYSICAL MODELLING	
A Mathematical Description of Sublethal Damage Repair and Interaction During Continuous Low Dose-Rate Irradiation M. Zaider & C.S. Wu	33
A Chromosomal "Fingerprint" for Past Exposure to Densely-Ionizing Radiation D.J. Brenner & R.K. Sachs	40
Correlations between α/β and $T_{1/2}$: Implications for Clinical Biological Modelling D.J. Brenner	44

CELLULAR AND MOLECULAR BIOLOGY

Cell-Cycle Dependent Radiation-Induced Transformation of C3H 10T _{1/2} Cells R.C. Miller, C.R. Geard, M.J. Geard & E.J. Hall	51
The Inverse Dose-Rate Effect for Oncogenic Transformation by Charged Particles R.C. Miller, G. Randers-Pehrson, L. Hieber, S.A. Marino, M. Richards & E.J. Hall	55
Sister Chromatid Exchange Induction as a Function of Charged Particle Linear Energy Transfer C.R. Geard	58
Human Chromosome Specific Changes in a Human-Hamster Hybrid Cell Line (AL) C.R. Geard & G. Jenkins	62
SSCP Analysis of <i>ras</i> Oncogenes in Neutron Transformed Mouse Cell Lines G.A. Freyer, R.C. Miller & Y. Yu	64

INTRODUCTION
STAFF NEWS
COLLOQUIA
STAFF LIST
TEACHING

INTRODUCTION

Research at the Center for Radiological Research is a blend of physics, chemistry and biology and epitomizes the multidisciplinary approach towards understanding the mechanisms involved in the health problems resulting from human exposure to ionizing radiations. To an increasing extent, the focus of attention is on biochemistry and the application of the techniques of molecular biology to the problems of radiation biology.

The following are some research highlights from the past year.

- Construction of a new wall-less ultra miniature counter is nearing completion. This instrument will make possible measurements of fundamental physical parameters on this type of counter as well as practical monodosimetric measurements of neutrons and charged particles.
- A mathematical model describing the production of single-strand and double-strand breaks in DNA as a function radiation quality has been completed.
- For the first time Monte Carlo techniques have been used to obtain directly the spatial distribution of DNA moieties altered by radiation. This information - which currently is not obtainable from experiment - was obtained by including in the transport codes a realistic description of the electronic structure of DNA.
- A collaborative effort with Dr. M.N. Varma of DOE has established that hit-size effectiveness functions in 1- μ m cellular volumes accurately predict available radiobiological data. This makes proportional counters available at the present time applicable to radiation protection
- Major components of our research activities involve oncogenic transformation assays. In a collaboration with the Medical Research Council in the U.K., we have investigated structure activity relationships for the potential oncogenicity of a new generation of bioreductive drugs that function as hypoxic cytotoxins.
- Experimental and theoretical investigation of the "inverse dose rate effect", whereby medium LET radiations actually produce an enhanced effect when the dose is protracted, is now at a point where the basic mechanisms are reasonably understood - and the complex interplay between dose, dose rate and radiation quality which is necessary for the effect to be present can now be predicted - at least in vitro.
- In terms of early radiobiological damage, a quantitative link has been established between basic energy deposition and "locally multiply damaged sites", the radiochemical precursor of DNA double strand breaks; specifically, the spatial and energy deposition requirements necessary to form LMDs have been evaluated.
- For the first time, a mechanically understood "biological fingerprint" of high-LET radiation has been established. Specifically measurement of the ratio of inter-to intra-chromosomal aberrations produces a unique signature from alpha-particles or neutrons.
- The Pulsed Brachytherapy concept, in which low dose rate brachytherapy is replaced with pulses of radiation, typically ten minutes in every hour, has now been taken up by a

commercial manufacturer and machines are in use in 9 centers around the U.S. and Europe. Over 100 patients have been treated. The limits of the technique are being validated in experimental systems.

- For the first time, cells of human origin have been transformed by a single dose of α -particles (10 cGy) which produce tumors in immune-suppressed mice. SSCP analysis failed to reveal any changes in p53. However, transfection studies are consistent with the measure of an activated dominant oncogene in these cells. We are currently working on isolating this gene. We have studied a large collection of (40) neutron transformed C3H 10T1/2 cells for damages in the ras oncogenes. Using the technique of SSCP analysis, no altered ras gene were detected in this large survey.

The productivity of the Laboratory continues at a high level, as evidenced by the steady stream of papers to peer-reviewed journals during the year, and invitations to members of the staff to participate in national and international meetings. Senior faculty are frequently called upon as consultants, reviewers and site visitors by government and private bodies.

Dr. Hall continues to serve on the BRER (Biological Research on the effects of Radiation) Committee and the BEIR VI Committee, of the National Academy of Sciences as well as Committee 1 of NCRP while Dr. Geard serves on the American Cancer Society Study Section in Prevention Diagnosis and Therapy.

Dr. Zaider has been invited to serve as a member of the "Expert Group on Hit Size Effectiveness Approach; Implications for Radiation Protection". He has also become a member of the Scientific Committee 88 of NCRP concerning the subject "Fluence as a Basis of a System of Radiation Protection for Astronauts".

Dr. Kiauga has been elected as a full member of the European Radiation Dosimetry Working Group 10, preparing a report on low pressure proportional counters in microdosimetry.

Funding levels have been steady during the past year. Drs. Hei and Brenner and Lieberman successfully competed for awards from the National Institute of Environmental Health, National Cancer Institute and the American Cancer Society.

STAFF NEWS

Dr. Eric Hall received the RSNA Gold Medal, the highest honor of The Radiological Society of North America. In early 1993 the Trustees of Columbia University appointed Dr. Hall to a named chair as the fourth Higgins Professor of Radiation Biophysics. The Society of Therapeutic Radiology and Oncology elected Dr. Hall Secretary for 1992-1993.

Dr. Charles Geard and Dr. David Brenner were elected to serve on the Council of the Radiation Research Society.

Dr. Marco Zaider has been invited to join the editorial board of Radiation and Environmental Biophysics.

Dr. Kliauga has been selected to be a Lead Investigator on the newly-constituted Radiation Physics and Biology Program of the Cancer Center.

Dr. Brenner was promoted to Associate Professor of Radiation Oncology with tenure.

Dr. C-Z Chen joined the Laboratory as Associate Research Scientist from the University of California at San Francisco.

Dr. Richard Miller rejoined our staff in January 1993.

Dr. Young Yu will be leaving us at the end of June to pursue a career in medicine.

Albert Fung has defended successfully his doctoral work at the physics department. His project "Electron Scattering off DNA Molecules", was performed under the supervision of Prof. Zaider.

Mrs. Jeanne Kramer retires this year after 33 years in the Center as a senior secretary. Her skills and sense of humor will be missed very much.

Mrs. Marie Burchett who was the secretary for our group at RARAF resigned in December.

THE COLUMBIA COLLOQUIUM AND LABORATORY SEMINARS

At intervals of approximately one month during the academic year, a regular colloquium has been held to discuss ongoing research. Dr. Gerhard Randers-Pehrson organized them and scheduled the speakers. These have been attended by the professional staff, graduate students, and senior technical staff of this Laboratory and RARAF, as well as scientists from other departments of the College of Physicians & Surgeons interested in collaborative research. Attention has focused on recent findings and future plans, with special emphasis on the on the inter-disciplinary nature of our research effort.

During the year, we have been pleased to welcome a number of visitors who have presented formal seminars and/or spent time discussing ongoing research with various members of the Laboratory. These have included Dr. Debbie Fisher, Rutgers University, New Jersey, Prof. B. L. Cohen, University of Pittsburgh, Dennis Heller, Carleton University, Ontario, Canada, Dr. Frank Ellis, The Churchill Hospital, Oxford, U.K., and Dr. Robert Ullrich, University of Texas, Galveston, Texas.

CENTER FOR RADIOLOGICAL RESEARCH

RESEARCH STAFF

- * **ERIC J. HALL**, D. Phil, D.Sc, FACR, Professor of Radiology and Radiation Oncology (Director)
- * **CHARLES R. GEARD**, Ph.D., Professor of Clinical Radiation Oncology (Associate Director)

- * **DAVID J. BRENNER**, Ph.D., Associate Professor of Radiation Oncology (in Public Health Environmental Science)
CHUN-ZHANG CHEN, Ph.D., Associate Research Scientist
- * **GREG A. FREYER**, Ph.D., Assistant Professor of Radiation Oncology
ZHU-YING HE, M.D., Staff Associate
TOM K. HEI, Ph.D., Assistant Professor of Radiation Oncology
KEVIN HOPKINS, M.S., Staff Associate
- * **PAUL J. KLIAUGA**, Ph.D., Assistant Professor of Radiation Oncology
HOWARD B. LIEBERMAN, Ph.D., Assistant Professor of Radiation Oncology
BRADFORD LOUCAS, Ph.D., Associate Research Scientist
- * **STEPHEN A. MARINO**, M.S., Senior Staff Associate
STEWART G. MARTIN, Ph.D., Associate Research Scientist
RICHARD C. MILLER, Ph.D., Assistant Professor of Clinical Radiation Oncology
CHANG-QING PIAO, M.D., Senior Staff Associate
- * **GERHARD RANDERS-PEHRSON**, Ph.D., Associate Research Scientist
- * **MARCO ZAIDER**, Ph.D., Associate Professor of Radiation Oncology and Public Health
YUQI ZHAO, Ph.D., Associate Research Scientist

POST-DOCTORAL FELLOWS

- * **MARJAN VRACKO-GROBELSEK**, Ph.D., Post-doctoral Research Scientist
KHALIFAH SIDIK, Ph.D., Post-doctoral Research Scientist
YONG YU, M.D., Staff Associate
LI XUN ZHU, M. D., Post-doctoral Research Scientist

DESIGN AND INSTRUMENT SHOP

- * **GARY W. JOHNSON, A.A.S., Senior Staff Associate (Director)**
- * **NICHOLAS LUDAS, Senior Research Worker**
DAVE CASSIMY, A.A.S., Instrument Maker

TECHNICAL STAFF

- JULIE FEINSTEIN, M.A., Senior Technician**
- JOSE GARCIA, Laboratory Assistant**
- MARCUS GEARD, B.A., Technician B**
- LAKSHMI GORIPARTHI, B.S., Research Worker**
- GLORIA JENKINS, B.A., Research Worker**
- ZENIN KURUNTHOTTICAL, B.A., Technician B**
- * **JESUS PEREZ, Accelerator Technician**
- * **MARCIA RICHARDS, B.S., Research Worker**

ADMINISTRATIVE AND SECRETARIAL STAFF

- * **MONIQUE REY, B.A., Center Administrator**
- * **MARIE BURCHETT, Senior Secretary (until December 1992)**
YVETTE CLAUSELL, A.A., Executive Secretary
- * **MARY COADY, Administrative Assistant**
- * **KAREN EVANS, B.A., Junior Accountant**
C. JEANNE KRAMER, Senior Secretary

- * **Partially or wholly supported by the Department of Energy**

TEACHING

Dr. Eric Hall taught in the Oncology Section of the course on "Abnormal Human Biology" for second-year medical students. In addition, a laboratory elective for medical students in "Cellular and Radiation Biology" was offered by Drs. Hall and Geard.

Drs. Brenner and Lieberman taught a course, "Radiation and Life" for non-science undergraduates, under the auspices of the Department of Biological Sciences. Dr. Brenner has taught a new course "Radon, Risk and Remedy" in the School of Public Health. He is supervising the Ph.D. work of Augustinus, in the School of Public Health.

Dr. Zaider is the Director of the Graduate Program in Medical Physics/Health Physics, offered jointly by the School of Public Health and the College of Physicians & Surgeons. Dr. Zaider is also supervising the Ph.D. work of Albert Fung of the Physics Department, Columbia University. He has also joined the Medical Physics program at the School of Engineering, Columbia University. He was invited to participate in several Thesis Examination Committees, at the Department of Physics, at the Department of Nuclear engineering, and at the School of Public Health.

During the Fall, a didactic course in Radiobiology was given by Drs. Hall, Geard, Brenner, Freyer, Hei, Miller, and Lieberman, for Columbia Presbyterian residents in Radiation Oncology and other institutions in the New York area. Drs. Hall and Geard also taught a course in Radiobiology for residents in Diagnostic Radiology. A similar course was taught at Montefiore Hospitals, as well as a one-day refresher course. Drs. Geard and Miller lectured to Radiological Technologists at United Hospitals, Port Chester, NY.

Dr. Kliauga taught an experimental course on "Nuclear Instrumentation", for M.S. students. Drs. Zaider and Brenner are lecturing on radiation effects in the courses entitled "Industrial Toxicology" and "Environmental Sciences," while Dr. Geard is lecturing in the course "Environmental and Molecular Toxicology." Additionally, Dr. Zaider is conducting a laboratory on the detection, measurement, and risk evaluation of radon in housing environments.

Dr. Greg Freyer teaches in the Medical Genetic and Cell Biology and Anatomy Programs at Columbia University. He is also a member of the Summer High School Science Teacher Program, and hosts a high school teacher, in his laboratory.

RADIATION PHYSICS AND MICRODOSIMETRY

MEASUREMENT OF FUNDAMENTAL PHYSICAL PARAMETERS OF A WALL-LESS ULTRAMINIATURE PROPORTIONAL COUNTER

Paul Kliauga

Much recent biological research supports the view that close range interactions, of the order of nanometers, can be a dominant factor in the production of biological damage from ionizing radiation. Until now, energy distributions at this level could be calculated and inferred, but could not be measured experimentally. A new instrument, designed and developed in this Center displays the capability of achieving this goal. This newly designed wall-less proportional counter is in process of construction. The design is based on a previously produced conventional walled counter called the UMC (ultra-miniature counter) which has been shown to be successful in the measurement of single-event energy deposition distributions at simulated tissue equivalent sizes of about 10 nm, far lower than any previously achieved. The new wall-less design will provide much more accurate spectra for those high LET radiations whose spectra are most distorted by the so-called wall effect, such as neutrons. In addition, this counter, unlike the UMC, will allow a probe by microbeams of charged particles and ultraviolet radiation. This will permit for the first time quantitative studies on relevant physical parameters which can be related to the fundamental physics of proportional counter operation and also allow comparison with modeling studies of avalanche processes in regions of large E/p .

The relative success of the ultraminiature proportional counter (UMC) in measuring single event ionization spectra at low pressures simulating about 10 nm in unit density tissue raises questions as to why the counter is able to have adequate resolution and gain, whereas conventional size counters have failed in this regard. We propose to investigate the physics of proportional counters at high E/p , making measurements of those parameters of proportional counter operation which should help to shed light on the limitations as well as the capabilities of this instrument for measuring "nanodosimetric" distributions of ionizations. Specifically, it is proposed to do a series of physical measurements on the following parameters:

1. Absolute gas gain as a function of E/p and wire diameter.
2. Single ion counter response as a function of E/p .
3. Avalanche radius as function of anode potential and wire diameter.
4. It is important to be able to compare the measured spectra with theoretical predictions. In this way the reliability as well as the limitations of the counter for simulating nanometer site sizes can be assessed. To this end, we will form a collaboration with Dr. David Brenner to develop Monte Carlo techniques to simulate spectra in nanometer size sites. Such a collaboration is very valuable in that it provides feedback for both the theoretical code and evidence for "proper" operation of the counter.

These measurements will provide physical parameters for testing models of proportional counter avalanche production, such as the recently proposed Monte Carlo model of Pierre

Segur and coworkers. In order to accomplish these measurements, a wall-less type ultra miniature counter must be used. The counter which was proposed for development at the last contract renewal is near completion, and such tests will be begun soon.

The new wall-less UMC design

The primary instrument of this proposed research plan is a new wall-less counter now under development at our laboratory. It utilizes elements of the UMC and the "Q" counter as prototypes in its design, but most closely resembles the "Q" counter. The Q counter has been in use for several years, and has been shown to possess possibly the best combination of gain and resolution of all the counters built in this laboratory, which is a group of considerable size and variety. The reasons for a particular counter excelling are not always amenable to scientific evaluation, because the actual construction of such a counter is still as much "art" as science. The basic principles are known, of course, but the finished product is the result of craftsmanship which can radically alter the characteristics of a counter. In this regard, the experience gained in our machine shop and laboratory over the years arguably make this laboratory the best (if not perhaps the only) site for development of such a counter. Among the many variables of construction which can affect performance are: precision alignment of field tubes along the axis, perfect circular concentricity of the field tubes to maintain true cylindrical geometry (a non-trivial problem), center wire size and smoothness, and pitch and size of the helix wires. There are many others, such as providing leak paths to reduce probability of electrical breakdown under application of high voltages to the center wire, but this gives an idea of the magnitude of the problem.

These problems are severe enough in the construction of "normal" size proportional counters, of the order of centimeter dimensions, but their difficulty is at least an order of magnitude greater when the outer dimension of the counter's collecting volume is of the order of a millimeter or less! The new wall-less UMC (WUMC) is shown in Fig. 1 below. The collecting volume is defined by a 12.5 μm stainless steel helix 1.0 mm in diameter. A good deal of trial and error experimentation was required to be learn how to construct such a helix which would have adequate pitch (turns/cm), have enough "spring" to maintain its shape, and not become distorted due to bowing or stretching in the center. The helix design was chosen over parallel wire designs when it was learned that in spite of its own problems, the helix design was still simpler to produce reliably and reproducibly.

Fig. 1. The new wall-less UMC, front and side views, showing the yoke which holds the entire assembly together.

The size of the counter's collecting volume was chosen as 1 mm as a compromise. Although the UMC had a 0.5 mm diameter, it was felt that counting rate at some of the RARAF lower energy neutron beams may be too low without an unacceptably high beam intensity, thereby reducing target lifetime. The 1 mm size should still be small enough to maintain acceptable "wall-lessness" in a reasonably-sized container. Previous experience with both the UMC and the Q counter indicate that such a design should prove capable of measurements of microdosimetric spectra down to 10 to 20 nm. The exact lower level is difficult to predict, being dependent upon some of those intangibles mentioned above.

Once constructed, it is intended to use this counter to perform the measurements mentioned above. This will be accomplished by mounting the counter on a precision slides with reproducible positioning better than 10 μm in directions transverse to the beam. The beam will enter into the chamber through a collimated aperture of a few micrometers in diameter, and a solid state detector mounted behind the counter with a similar aperture will insure coincidence between a counter signal and the desired trajectory of the incoming particle.

Single electron spectra can be obtained by using an ultraviolet source, such as a UV laser, directed at the inside of one of the helix wires. The single electron will be released at the

boundary of the collecting volume and the resulting signal can be measured to determine accurately the single electron response function of the counter as a function of E/p .

Calibration will be achieved by an intense uncollimated alpha source using the alpha "edge", i.e., the maximum energy an alpha particle can deposit. This has a number of advantages compared to the conventional use of a collimated source and a "peak", which requires precise knowledge of both alpha energy and chord length. Neither is relevant in the edge technique. Furthermore, at very small site sizes, the peak may be so broad that accurate determination is difficult. The edge method was tried with the Q counter and found to give quite satisfactory results.

Figure 2 shows the end of the beam pipe mounted onto the outer chamber of the WUMC. The beam pipe protrudes some distance into the chamber to allow a close approach of the counter, to a distance of about 3 mm. In experiments examining the extent of the avalanche region, this minimizes the delta ray contribution, since the particle entrance will be so close to the sensitive volume, there will be negligible contribution of delta rays entering from outside the collecting volume.

Fig. 3a. Center Wire=100V; Field Tube=0V; Cathode=-500V.

Fig 3b. Center Wire=300V; Field Tube=0V; Cathode=-300V

Fig. 3c. Center Wire=500V; Field Tube=0V; Cathode=-100V.

It is important in counter design to insure that nearby insulators do not appreciably perturb the field in the sensitive volume, and to know the proper ratios of electrode potentials to obtain a uniform cylindrical field. Figures 3a-3c illustrate equipotential distributions obtained by solving the two dimensional Poisson equation with cylindrical symmetry for different configurations of center wire and cathode potentials. The cathode is assumed solid for simplicity, and the field tube is at ground potential. the difference between center wire and cathode is constant at 600 Volts. Only the region near the ends of the field tubes are shown. The best symmetry is obtained for center wire and cathode at equal potentials with opposite polarity.

Figure 4 is a solution for the equipotential surfaces using a realistic representation of the helix wires, drawn to the proper scale, and using the potentials found in Fig. 3b. It is evident that the field region within the sensitive volume is still a good approximation to a cylindrically symmetric field. The major difference is the regions of weak potential at the midpoints between helix wires, where diffusion may be a problem for electrons formed there.

The counter is mounted on precision slides allowing adjustment in three mutually orthogonal directions. The axes of two of those directions can be chosen independently by rotating the counter yoke in its mount on the "z" slide. Thus the counter can be scanned relative to the beam axis either laterally or longitudinally. The slide along the beam axis (the "x" axis) has a long range travel allowing motion of about 25 cm in the beam direction. The other motions are fine and allow about 2 mm travel with a precision of 10 μm . Figure 5 is a frontal view along the beam axis showing the x and y slides and the cables for their motion adjustment.

Fig. 5. Front view along the beam axis, showing counter mounted in the chamber and slides for motion adjustment in x and y directions.

MICRODOSIMETRY OF LOW ENERGY PHOTON SOURCES

Paul Kliauga

Systems for quantifying the biological effectiveness of different radiations for purposes of radiation protection currently rely on the assignment of a numerical value called the quality factor, which multiplies the absorbed dose, D , to provide a quantity called dose equivalent, H :

$$H = Q \cdot D.$$

Q is given as a function of LET, and the resulting quantity H for a given LET is supposed to represent the dose of the radiation of interest which produces the same biological effect as an absorbed dose D of x-rays or ^{60}Co .

The limitations of LET as a parameter for defining radiation quality are well known, and alternative approaches based on microdosimetric principles have been suggested, using specific quality functions $q(y)$, as the microdosimetric equivalent of the quality factor $Q(\text{LET})$. Project 3 by M. Zaider attempts to calculate this quantity for photons from the initial slopes of dose response curves for a given biological endpoint:

$$\alpha_i = \int_0^\infty q(y) f_i(y) dy$$

where the α_i are the initial slopes and the $f_i(y)$ are the single event microdosimetric spectra for radiation i .

The solution for $q(y)$ is obtained from a set of integral equations of this form which obviously depend on the biological endpoint chosen, implicit in the input of the α_i . From the point of view of the physics, the only input is the set of single event microdosimetric spectra $f_i(y)$ which should cover a wide range of photon energies and site sizes.

A spherical wall-less proportional counter with no metallic components was developed in this laboratory years ago, (Kliauga, P. and Rossi, H.H., 1982) but was not used for a systematic measurement of photon microdosimetry. A few spectra for ^{125}I and ^{241}Am were measured, but this was with a view to show that in comparison with a conventional wall-less counter containing metal electrodes, there was a substantial difference in the spectra. This is due to the fact that at lower photon energies, in the range up to about 100 keV, the interaction cross section of photons in low-Z tissue equivalent materials undergoes a transition from primarily photoelectric to Compton. Metallic materials are primarily photoelectric throughout this energy range. Thus, the presence of metals distorts the secondary electron spectrum with an overabundance of more energetic photoelectrons.

The original non-metallic counter used a graphite coated human hair as the "center wire", or anode. While this provided a satisfactory tissue-like element, it was difficult to work with, often non-uniform in diameter and relatively large (90 μm), not easy to obtain, and subject to breakage because of poor tensile strength. The availability of carbon impregnated nylon fibers as solved this problem, and the counter has been rebuilt using such a fiber as the anode, with a diameter of 2.5 μm . A cut-away view of the counter is shown in Fig. 1.

As preliminary experiments preparatory to a large scale systematic investigation of a number of low energy photon sources using this counter, we have measured the single event spectra for ^{125}I and ^{103}Pd , with mean photon energies of 28 and 21 keV, respectively. Brachytherapy type seeds of these isotopes were mounted in a slit milled into the inside surface of a tissue equivalent plastic plug which was inserted through the outer wall of the counter. In this

geometry, the seed is flush with the inner wall of the TE shell. Different depths can be simulated by attaching material to the plug on top of the seed. Spectra are shown in Fig. 2 for ^{125}I and ^{103}Pd at depths of 1 cm of A150 tissue equivalent plastic. At 1 mm depth, the spectra are virtually identical to Fig. 1. Comparison of 1mm and 1 cm spectra for the same isotope, shown in Fig. 3, reveal very minor differences, which is reflected in the shift of y_D by only a few percent. Values of y_D for these isotopes are given in Table I.

Fig. 1. The non-metallic proportional counter. The "hair" has been replaced by a carbon-impregnated nylon fiber. Two holes are drilled through the outer wall, one to insert the photon source "seeds", the other for a calibration source.

Fig. 2. Iodine and Palladium at 1 cm depth of tissue equivalent plastic.

Fig. 3. Comparison of Palladium spectra at 1 mm and 1 cm depths of A150 tissue equivalent plastic and 1 μm site size. The 1 cm spectrum is the one slightly higher in the peak area.

Table I. Comparison of ^{125}I and ^{103}Pd at two depths of A150 tissue equivalent plastic. Values of y are in $\text{keV}/\mu\text{m}$.

	Iodine		Palladium	
	1 mm depth	1 cm depth	1 mm depth	1 cm depth
y_F	1.46	1.52	1.77	1.54
y_D	3.59	3.68	3.90	4.00

References

Kliauga, P. and rossi, H.H. (1982). Studies with encapsulated sources of ^{125}I . III. Microdosimetry using a non-metallic wall-less proportional counter. Int. J. Radit. Onc. Biol. Phys. 1963-1968.

A STUDY OF EXCITED STATES IN TRANS-POLYACETYLENE IN THE HARTREE-FOCK, TAMM-DANCOFF, AND RANDOM-PHASE APPROXIMATION

M. G. Vracko and M. Zaidar

1. Introduction

Trans-polyacetylene (C_2H_2)_n has been the topic of numerous theoretical and experimental studies. The elementary gap is about 1.4 to 1.8 eV and in some circumstances (as doped polymer) it displays high electric conductivity. Calculations of the electronic structure have served as a base for different models treating the elementary excitations in one-dimensional semiconductors. Furthermore, they have been used to describe transport phenomena. The character of these excitations, on the other hand, has been a matter of controversy.

The elementary cell of polyacetylene consists of two hydrocarbon units and thus can be treated efficiently using quantum chemical calculations. Moreover, the modest amount of output data (seven doubly occupied bands in the case of two CH units per elementary cell) provides a clear picture for theoretical discussions. Another advantage is the relative abundance of experimental measurements performed on polyacetylene crystalline films.

In the present work we treat the correlation effects in the exciton approximation, which can be understood within the particle-hole propagator formalism (1). The energy bands in polyacetylene can be separated in σ and π bands, and we report the excitonic transitions for different bands. Furthermore, we define several levels of approximation by taking either one or two-time ordering in the two-particle propagator or the band-band interaction into account (1-3).

2. Theoretical Methods and Computational Details

The brief description of the exciton formalism is given in Refs. (1-3).

The calculations were performed with the STO-6G basis. The two CH units in the elementary cell had the following parameters: C-C distance is 1.377 Å, C-H 1.08 Å, and the C-C-C angle is 120°. The polymer is oriented along the z direction.

3. Results and Discussion

The band structure obtained is shown in Fig. (1). Bands 7 and 8 have π and π^* character, i.e. the Hartree-Fock eigenfunctions are combinations of p_y functions (oriented perpendicular to the polymer plane). On the other hand, the eigenfunctions of the bands 3 to 6 and 9 to 12 are located in the polymer plane with no p_y contributions (σ and σ^* bands). The bands 6 and 7 are crossing near the Γ point.

Table 1 shows the three lowest exciton energies calculated at different levels of approximation. The first three rows show the exciton energies calculated with occupied and unoccupied bands of different symmetries. By comparing these results with those in the last row it is obvious that the σ - π coupling plays a minor (if any) role in the exciton calculation. The difference between the TD and RP approximation is of the order of 0.8 eV. Compared to the HF picture, both approximations (TD and RP) lead to a lowering of the exciton energies (by about 1 eV for π - π^* and 5 eV for σ - σ^* excitons). In Table 1B we report the excitons at

$K=0.25 \text{ \AA}^{-1}$. From Fig. (1) it is obvious that the exciton energies become larger for $\pi-\pi^*$ excitons. The gap, which is included in Eq. (1) via Eq. (2), becomes larger for the non-zero momentum-transfer excitations. The opposite is true for $\sigma-\pi^*$ and $\pi-\sigma^*$ excitons.

The electron-loss-spectra show a sharp peak at 4.9 eV and also a wide, strong peak at about 22.5 eV. The lowest calculated energies for the $\pi-\pi^*$ and $\sigma-\sigma^*$ singlet excitons are 4.225 eV and 21.536 eV. The spectra also display small overlapping peaks at 9 eV, 12.5 eV and 17 eV, which may belong to the $\sigma-\pi^*$ and $\pi-\sigma^*$ excitons.

Figs. 2 (A,B,C) show the exciton densities of states within the approximations described above.

4. Conclusions

This work belongs to the class of *ab initio* calculations that include correlation effects in the treatment of trans-polyacetylene. Several calculations have been performed in the framework of the single-particle Green's function formalism. This means that they treat the transitions from the ground state to the quasi-particle bands as band-to-band transitions. The exciton picture, on the other hand, includes two-particle states. In comparison with previous work, we have introduced both time orderings in the particle-hole propagator. Additionally, we have investigated the excitons between bands of different symmetry.

The energy-loss function, which is a measurable quantity, for instance with electron energy loss spectroscopy, may serve as a significant test for calculations of the type shown here. This quantity is very sensitive not only to the correct positioning of the excitation bands, but also to the wave functions used to calculate the transition probabilities. It is also of interest to establish whether the large peak observed at about 21 eV is of collective nature (e.g. plasmonic excitation) or whether it continues to represent single-particle transitions. Such calculations are currently underway in our laboratory.

References:

1. M. G. Vracko and M. Zaider, Int. J. Quantum Chem. **43**, 321 (1992).
2. M. G. Vracko and M. Zaider, Int. J. Quantum Chem. (accepted)
3. M. G. Vracko and M. Zaider, CRR Ann. Rep. 1992, 11; *ibid.* 23.

Figure 1: The Hartree-Fock band structure of trans-polyacetylene.

Figure 2 A: The exciton density of states (DOS) in the Hartree-Fock approximation

Figure 2 B: The exciton density of states (DOS) in the Tamm-Dancoff approximation

Figure 2 C: The exciton density of states (DOS) in the Random-Phase approximation

Table 1: The three lowest exciton energies. The three levels of approximation (HF, TD and RP) are defined in the text (energies in eV).

K=0					
occ. - unocc.	HF	TD		RP	
		singlet	triplet	singlet	triplet
$\pi-\pi^*$	5.144	4.075	/	3.225	2.467
		5.582	5.205	4.024	4.948
		7.323	5.586	4.192	5.586
$\sigma-\pi^*$	17.265	10.733	10.204	10.732	10.203
		11.929	11.607	11.065	10.534
		14.472	14.298	11.927	11.602
$\pi-\sigma^*$	22.115	16.739	16.036	16.557	15.835
		17.253	16.565	16.784	16.130
		18.628	18.159	17.131	16.462
$\sigma-\sigma^*$	26.587	22.297	16.579	21.536	15.460
		23.192	18.125	22.361	17.104
		23.739	21.859	22.610	17.487
$\sigma, \pi \quad - \quad \sigma^*, \pi^*$		4.022	/	3.167	2.358
		5.586	5.189	3.963	4.881
		7.276	5.586	4.105	5.586
K=0.25Π/a					
occ. - unocc.	HF	TD		RP	
		singlet	triplet	singlet	triplet
$\pi-\pi^*$	9.283	5.070	/	4.165	/
		8.183	6.903	5.152	5.276
		8.668	7.470	6.938	7.141
$\sigma-\pi^*$	21.171	10.337	9.892	10.335	9.891
		12.465	12.048	10.479	10.032
		13.863	13.556	12.463	12.044
$\pi-\sigma^*$	21.973	16.359	15.735	16.190	15.578
		16.785	16.216	16.403	15.799
		18.651	18.161	16.708	16.144
$\sigma-\sigma^*$	28.359	22.092	16.574	21.699	15.902
		23.126	18.073	22.165	16.858
		23.727	21.717	22.796	17.567
$\sigma, \pi \quad - \quad \sigma^*, \pi^*$		5.020	/	4.061	/
		8.181	6.888	5.072	5.087
		8.655	7.170	6.878	6.858

ELECTRON TRANSPORT IN CONDENSED WATER

M. Zaider, M.G. Vracko and A.Y.C. Fung

Introduction

Electron, and more generally charged-particle transport in biomatter represents an important step in describing the biological effects of ionizing radiation. Transport calculations are expected to provide a description of the spatial and temporal pattern of those radiation-induced molecular alterations deemed responsible for the biological effect. Traditionally, this task has been accomplished with the aid of Monte Carlo simulation codes where charged particles interact with a "generic" biological material, most often water vapor. Cellular density is accounted for by scaling distances to the density of liquid water. The physics of charged particle interaction with water vapor is reasonably well understood, and it is quite abundantly supported with good quality experimental cross sectional data. This is the primary reason for the widespread usage of vapor-based calculations.

More recently efforts have been made to treat explicitly interactions in **condensed** matter¹. The implementation in an event-by-event Monte Carlo transport code of our previously-reported interaction cross sections for crystalline water (cubic ice) is reported here.

Electron (and charged particle) transport in a condensed medium is usually treated in the so-called **semi-classical** approximation². In between energy loss events primary and secondary electrons are assumed to travel along rectilinear trajectories (quantum mechanically, wave packets are introduced at this point). The quantum mechanical system consisting of the primary electron and the crystal is taken to be separable in the sense that:

$$|e, \text{crystal}\rangle = |e\rangle |\text{crystal}\rangle. \quad (1)$$

One may then apply the first Born approximation and use **stationary** (time-independent) properties of the crystal to obtain interaction probabilities.

Within this approximation, the double-differential cross section for electron scattering resulting in energy loss ω and momentum transfer q is given by:

$$\frac{d^2\sigma}{d\omega dq} = \frac{2}{\pi N v^2} \frac{1}{q} \text{Im} \left[-\frac{1}{\epsilon(q, \omega)} \right]. \quad (2)$$

N is the total electron density in the crystal, v is the velocity of the incident electron, m is the electron mass, and $\epsilon(q, \omega)$ is the frequency (ω) and wave-vector (q) dependent dielectric response function of the crystal. The cross section, σ , is per crystal electron. In Eq(2) it is assumed that σ does not depend on the direction of q . k_i and k_f are the initial (final) momenta of the projectile, and θ is the polar scattering angle. ω and q are not independent quantities:

$$\begin{aligned} q &= k_i - k_f, \\ \omega &= \frac{1}{2m} (k_i^2 - k_f^2). \end{aligned} \quad (3)$$

An expression equivalent to Eq(2) is

$$\frac{d^2(\Lambda^{-1})}{d\omega d\Omega} = \frac{1}{(\epsilon\pi a_0)^2} \left(\frac{k_i}{k_f}\right) \frac{1}{q^2} \text{Im}\left[-\frac{1}{\epsilon(q, \omega)}\right]. \quad (4)$$

Here, momentum transfer is replaced with the scattering solid angle, $d\Omega$; $\Lambda=1/(N\sigma)$ is the mean free path of the electron and a_0 is the Bohr radius. The quantity $\text{Im}[-1/\epsilon]$ is termed the **energy-loss function** of the crystal; it is related to the more familiar quantity, **differential oscillator strength**, as follows:

$$\frac{df(\omega, q)}{d\omega} = \frac{2\omega}{\pi\omega_p^2} \text{Im}\left[-\frac{1}{\epsilon(q, \omega)}\right] \quad (5)$$

where ω_p is the plasma frequency for an electron gas ($\omega_p=4\pi N$). The energy-loss function satisfies the Thomas-Reiche-Kuhn sum rule:

$$\int_0^{\omega_p} \frac{df(q, \omega)}{d\omega} d\omega = 1. \quad (6)$$

Other quantities useful in electron transport result from Eq(2) in an obvious manner. Thus, the single-collision spectrum is given by:

$$\frac{d(\Lambda^{-1})}{d\omega} = \frac{2}{\pi} \frac{1}{k_i^2} \int_{q_{\min}}^{q_{\max}} \frac{dq}{q} \text{Im}\left[-\frac{1}{\epsilon(q, \omega)}\right], \quad (7)$$

where:

$$q_{\min, \max} = k_i \pm \sqrt{(k_i^2 - 2\omega)}. \quad (8)$$

Within the approximation stated above, the electron transport problem requires the evaluation of the dielectric response function. This is discussed below.

The evaluation of $\epsilon(q, \omega)$

Experimental information on $\epsilon(q, \omega)$ for biomaterials (including water) is rather limited; it is typically restricted to the $q=0$ region of the (q, ω) plane, as obtained from measurements of the refraction index or electron scattering. Since electron transport requires cross sections at non-zero momentum transfer, it is necessary to evaluate $\epsilon(q, \omega)$ theoretically. The basis for this evaluation is:

$$\epsilon_{KK'}(q, \omega) = \delta_{KK'} + \frac{4\pi e^2}{|q+K||q+K'|} \sum_{k, n, m} \frac{F_{(q+K)nm}(k) F_{(q+K')nm}(k)}{E_m(k+q) - E_n(k) - \omega - i\delta}. \quad (9)$$

In this expression the crystal is described in terms of one-electron wave functions $|nk\rangle$ of band index n , crystal wave vector k and energy $E_n(k)$. K and K' are reciprocal lattice vectors, Ω is the crystal volume and δ^+ is an infinitesimal positive number (the limit $\delta^+ \rightarrow 0$ is taken at the end). The summation in Eq(9) is over all occupied states, n , and unoccupied (conduction) states, m . Also:

$$F_{qnm}(k) = \langle nk | e^{-iqr} | m, k+q \rangle. \quad (10)$$

A direct evaluation of Eq(9) requires a complete quantum mechanical description of the crystal. For all but the simplest systems this is not possible, primarily because of computing limitations. In a previous paper we have described calculations of the dielectric response function for cubic ice; the results were obtained within the tight-binding approximation. We have evaluated directly Eq(9) (but with local field corrections ignored, that is $K=0$, $K'=0$), fitted available optical data ($q=0$), and extended then the result in the (ω, q) plane. An example of calculated $\epsilon(q, \omega)$ is shown in Fig. 1. These results may be used directly in the Monte Carlo transport code.

Figure 1: The energy-loss function, $\text{Im}[-1/\epsilon(q, \omega)]$, calculated as a function of frequency and wave vector.

The transport code for condensed water

The code described here (symbolically termed DELTAI) is a modified version of our water-vapor based code, PROTON⁴. Features common to both codes are not being further discussed here. This includes, for instance, the treatment of electron elastic scattering for which no data exist on condensed water other than at subexcitation energies. These latter are treated separately as described below. The treatment of vibrational excitation is also similar to that used in PROTON⁴; the excitation energies for these processes

were taken, however, from data on ice⁵.

The evaluation of $\epsilon(q, \omega)$ of Eq(9) requires computationally-intensive calculations of the single-particle band structure and wave functions. The dielectric response function was evaluated in the domain $0 < \omega < 3.6$ a.u., $0 < |q| < 9$ a.u. Although separate calculations have been performed along q_x , q_y , q_z (and combinations thereof) the results used in the code DELTAI represent spherically-symmetric averages of these quantities.

The problem of analytically continuing the energy loss function, $\text{Im}[-1/\epsilon(q, \omega)]$ at large q values (at a fixed ω) has been discussed by Dillon et al⁶ for the case of secondary electrons resulting from the ionization of molecules by charged particles. We have used an expression similar to that suggested in the cited paper, namely:

$$\text{Im}\left[-\frac{1}{\epsilon(q, \omega)}\right] = \sum_i \frac{A_i(\omega) q^{2i}}{[(q^2 - 2 \times \omega)^2 + 8Iq^2]^M} \quad (11)$$

Although in the molecular case parameters such as I and M have well defined interpretations¹¹ we have used them here simply as adjustable constants. Fig.2 shows an example of a spectrum, $d\sigma(\omega, q)/dq$, as a function of q for 1-keV electrons. Here $\omega = 21.8$ eV.

Figure 2. The spectrum, $d\sigma(\omega, q)/dq$, as a function of q for 1-keV electrons.

The single-collision (secondary electron) spectrum, $d\sigma(\omega, T)/d\omega$, for primary electrons with kinetic energy T can be obtained by integrating the expressions, Eqs(2) and (11), over all allowed momentum transfers. To extrapolate this spectrum at energies $\omega > 100$ eV we have used the analytic form proposed by Green and Sawada⁷:

$$\frac{d\sigma_{GS}(\omega, T)}{d\omega} = \frac{A(T)}{[\omega - B(T)]^2 + \Gamma^2(T)}, \quad (12)$$

where $\omega \in [0, T/2]$ and, again, as suggested⁷ and Γ are parameterized as follows:

$$B(T) = a_1 - \frac{1}{T + a_2}, \quad (13)$$

$$\Gamma(T) = \frac{a_3 T}{T + a_4}. \quad (14)$$

Fig.3 shows the single collision spectrum, $d\Lambda^{-1}(\omega)/d\omega$, as a function of ω for 1-keV electrons. The total cross section, σ , as a function of the kinetic energy of the electron, T , is displayed in Fig.4. Also shown in this figure is the total cross section for water vapor as used in the code DELTA⁴. Over the range of electron energies shown σ for condensed water is larger than its counterpart for the vapor phase. Total cross sections are obtained either by direct integration of $d\sigma/d\omega$ or - at larger energies - through the device of the Fano plot.

Figure 3. The single collision spectrum, $d\Lambda^{-1}(\omega)/d\omega$, for 1-keV electrons.

Figure 4: The total cross section, σ , as a function of the kinetic energy, T , of the electron. The results for ice (solid) and vapor water (dash) are shown.

The treatment of excitonic states in condensed water is somewhat problematic. As already anticipated, a significant reduction in the probability of excitonic transitions occurs as water vapor condenses. Spectroscopic analyses⁸ reveal two transitions, at 8.7 and 10.4 eV, which

appear in both cubic and amorphous ice, and could therefore be associated with molecular excitons. Based on this we have retained in the code the molecular Rydberg states (9.998 and 11.057 eV) and the dissociative continua at 7.44 eV and 9.85 eV. As opposed to molecular excitations, excitonic transitions are induced with momentum transfer. Improved calculations need to take this into account.

The transport of subexcitation electrons

A recurrent problem in electron transport is the treatment of subexcitation electrons (typically of energy below several eV). The quantity of interest here is the thermalization range of these electrons and its statistical distribution (range straggling). In water vapor subexcitation electrons interact with molecular, vibrational and rotational modes. Cross sections for these processes are available⁴. In condensed water, however, the situation is quite different: a) interaction cross sections as a function of energy and momentum transfer are largely unknown [an exception are the recent results of Michaud and Sanche⁹ that give total cross sections for electrons above 2.7 eV interacting with films of amorphous ice]; and b) quantum mechanical considerations make it difficult to continue to use Monte Carlo schemes based on random walk of electrons with **well defined** momentum along **well defined** straight-line trajectories when the wave length spans many target molecules. To put it differently, the assumptions behind Eq(1) are no longer valid. Monte Carlo studies of the transport of subexcitation electrons in solid water have been nevertheless undertaken¹⁰.

Another possibility is to use experimental data on electron photoinjection from metals into electrolyte solutions (PIMES). The details of this technique have been recently reviewed by Konovalov et al¹¹. We have used this information to obtain an explicit analytic expression for the range distribution of subexcitation electrons. This is:

$$h_E(l) = \frac{1}{2} f_1(l, E) \left[1 + \frac{l[x - m(E)]}{\sigma^2} \right], \quad (15)$$

where $h_E(l)$ is the probability distribution function for the range, l , of an electron of energy E and $f_1(l, E)$ is a normal distribution [mean $m(E)$ and variance σ^2]. Fig.5 shows several distribution functions for the range of subexcitation electrons. The simulation of these distributions has been implemented in the code DELTAI.

Fig 5: The range distribution of subexcitation electrons at the energies indicated.

Microdosimetric results

In this section we present several results obtained from electron tracks in condensed water. In Fig.6 we compare our results for the stopping power of water for electrons (solid line) with results obtained in other calculations. There appears to be fair agreement between our calculations and the stopping power evaluations of Ashley¹² and Kutcher and Green¹³. However, the earlier results of Ritchie et al²³ are significantly larger. For high-energy electrons we obtain for the energy per ion pair (w) a value of 21.2 eV, in

line with some experimental determinations. At the same electron energy the water vapor result is $w=29$ eV. For comparison purposes, Turner et al¹ report $w=24.6$ eV for liquid water and $w=30$ eV for vapor, while Kutcher and Green¹³ obtain $w=28.5$ eV and $w=30.7$ eV, respectively.

The question of whether there are significant spatial differences between the distribution of inelastic energy deposition events in vapor and condensed phase may be addressed by calculating microdosimetric distributions. While - as already indicated - it is not clear that in the condensed phase the location of the energy loss event is identical with the location of the energy absorption event we have assumed such differences (if real) to be much less than micrometer dimensions. In Fig.7 we compare microdosimetric spectra for 30-keV electrons in 1- μ m diameter spheres containing water vapor (solid curve) or condensed water (dashed curve). One notes a larger number of events at high linear

energy (y) values in the condensed phase. This is reflected in increased values of the frequency average (y_F) and the dose average (y_D) of y . To the extent that y_D is an index of biological effectiveness¹⁴ the results shown in Fig.7 suggest the potential importance of solid state microdosimetry in radiobiological applications.

It is a pleasure to acknowledge illuminating discussions with Prof. R. Ritchie and Dr. M. Inokuti. The possibility of obtaining the range of subexcitation electrons from the PIMES data was suggested to us by Dr. D. Brenner.

References

1. Turner, J.E., Paretzke, H.G., Hamm, R.N., Wright, H.A. and Ritchie, R.H. Comparative study of electron energy deposition and yields in water in the liquid and vapor phases. *Radiat. Res.* **92**, 47-60 (1982).
2. Ashcroft, N.W. and Mermin, N.D. *Solid State Physics*, Saunders College, Philadelphia (1976). Chapter 12.
3. Zaider, M., Fry, J.L. and Orr, D.E. Towards an *ab initio* evaluation of the wave-vector and frequency-dependent dielectric response function for crystalline water. *Radiat. Prot. Dosim.* **31**, 23-28 (1990).
4. Zaider, M., Brenner, D.J. and Wilson, W.E. The application of track calculations to radiobiology I. Monte Carlo simulation of proton tracks. *Radiat. Res.* **95**, 231-247 (1983).
5. Bertie, J.E., Labbe, H.J. and Whalley, E. *J. Chem. Phys.* **50**, 4501 (1969).
6. Dillon, M.A., Inokuti, M. and Wang, Z. Analytic representation of the generalized oscillator strength for ionization. *Radiat. Res.* **102**, 151-164 (1985).
7. Green, A.E.S. and Sawada, T. Ionization cross sections and secondary electron distributions. *J. Atmos. Terr. Phys.* **34**, 1719-1728 (1972).
8. Shibagaki, T., Onuki, H. and Onaka, R. Electronic structure of water and ice. *J. Phys. Soc. Jpn.* **42**, 152-158 (1977).
9. Michaud, M. and Sanche, L. Absolute vibrational cross sections for slow electrons (1-18 eV) scattering in solid H₂O. *Phys. Rev.* **A36**, 4684-4699 (1987).
10. Goulet, T. and Jay-Gerin, J.P. Thermalization distances and times for subexcitation electrons in solid water. *J. Phys. Chem.* **92**, 6871-6874 (1988).
11. Konovalov, V.V., Raitsmiring, A.M. and Tsvetkov, Y.D. Thermalization lengths of "subexcitation electrons" in water determined by photoinjection from metals into electrolyte solutions. *Radiat. Phys. Chem.* **32**, 623-632 (1988).
12. Ashley, J.C. Stopping power of liquid water for low-energy electrons. *Radiat. Res.* **89**, 25-31 (1982).
13. Kutcher, G.J. and Green, A.E.S. A model for energy deposition in liquid water. *Radiat. Res.* **67**, 408-425 (1976).
14. Kellerer, A.M. and Rossi, H.H. A generalized formulation of dual radiation action. *Radiat. Res.* **75**, 471-488 (1978).

BIOPHYSICAL MODELLING

A MATHEMATICAL DESCRIPTION OF SUBLETHAL DAMAGE REPAIR AND INTERACTION DURING CONTINUOUS LOW DOSE-RATE IRRADIATION

M. Zalder (with C. S. Wu)

Introduction

The clinical use of low dose-rate brachytherapy has a long history. With this type of treatment, a high radiation dose can be delivered to a tumor locally with a rapid dose fall-off in the surrounding normal tissue. This therapeutic advantage of nonuniform dose distribution is achieved by means of its geometry. The effect of dose-rate on the therapeutic gain is however harder to determine.

Cells irradiated at low dose-rate are subject to three processes⁽¹⁾: a) sublethal damage repair, b) repopulation, and c) redistribution of cells in the cell cycle. A typical implant (such as the treatment of cervix carcinoma) consists of irradiating the tumor at a dose rate of 50 cGy/hr for 40 hrs. At this dose rate, cell repopulation is unimportant because the irradiation slows the cell progression through the G2 block. It is widely believed that the survival properties of populations of cells exposed to protracted doses of radiation are dominated by repair processes, the effect of reassortment of cells within the cell cycle, and the difference in radiosensitivity in each phase of the mitotic cycle. It is desirable to increase the cell damage by taking advantage of the fact that cells progress into a more sensitive phase of the cycle. The influence of cell-cycle redistribution on cell killing is more dramatic for certain cell lines (such as HeLa cells) than for others⁽²⁾. Survival properties due to variation in the phase-dependent radiosensitivity has generally been regarded as minimal. There has been no theoretical analysis on the effect of changes in radiosensitivity through cell cycle on the overall survival properties.

We describe a mathematical model to calculate the survival property of cells that are progressing through the cell cycle during irradiation. Our model includes, a) phase-specific parameters, α and β , b) different recovery rates in each phase of the cell cycle, and c) the interaction of sublesions produced in different phases of the cycle. The formalism presented here is conceptually (and, to a certain extent, mathematically) equivalent to our previous treatment of cellular survival following exposure to mixed-fields of radiation⁽³⁾.

Mathematical Model

The linear-quadratic formalism^(4,5) for dose-effect curve is expressed as

$$\epsilon(D) = \alpha D + \beta D^2 \quad (1)$$

or, for survival probability

$$S(D) = e^{-\alpha D - \beta D^2} \quad (2)$$

The survival expression, Eq(2), obtains if one assumes a Poisson distribution of the number of lesions, and also if lesions produce the effect (cell killing) independently. The yield of sublesions is a measure of the sublethal damage in the cell. Furthermore, cellular repair mechanisms may remove some of the sublethal damage.

Sublethal damage recovery will reduce the quadratic term of Eq (2) by a factor, q :

$$q(x) = 2x - x^2 [1 - e^{-\frac{1}{x}}] \quad (3)$$

where $x = t_0/T$, T is the irradiation time and t_0 is a recovery constant. The expression, Eq(3), has been calculated from:

$$q(t) = \int_0^t \tau(t') h(t') dt' \quad (4)$$

$h(t)$ is the distribution of time intervals between pairs of sublesions and $\tau(t) = \exp(-t/t_0)$ quantifies the amount of sublesion left unrepaired during the time interval, t . The expression, Eq(2), has been extended to the situation involving sequential exposures to two different radiations⁽⁶⁾, 1 and 2. Thus

$$-\ln S(D) = \sum_{i=1}^2 [\alpha_i D_i + \beta_i q(\frac{t_0}{t_i}) D_i^2] + 2q_{12}(\frac{t_0}{t_1}, \frac{t_0}{t_2}, \frac{t_0}{T}) (\beta_1 \beta_2)^{\frac{1}{2}} D_1 D_2 \quad (5)$$

where

$$q_{12}(x, y, z) = xye^{-1/z} (1 - e^{-1/x})(1 - e^{-1/y}), \quad (6)$$

t_1 and t_2 are the respective irradiation time from radiation 1 and 2, and T is the time interval between irradiations. The meaning of these expressions is obvious: the first two terms represent effects produced by each radiation individually. the third term represent interactions between sublesions produced by the first radiation (and left unrepaired) and sublesions produced by the second radiation.

The situation of cells being exposed to radiation while moving through the mitotic cycle [and changing their radiosensitivity parameters (α, β, t_{0i})] is analogous to the sequential exposure scheme described above. To make the derivation below more manageable and easier to follow, we divide the mitotic cell cycle in 4 intervals (G1, S, G2, M), assign to each stage parameters $\alpha_i, \beta_i, t_{0i}$, assume that the cell spends the time t_i ($i=1, 2, 3, 4$) in each interval, and do not consider the effects of cell division. This latter assumption may be removed, if necessary, but it appears easier to treat it non-analytically (i.e. with Monte Carlo techniques). The time a cell spends in each stage of the cell cycle depends on dose, and a mathematical formalism has been proposed recently to account for this dependency⁽⁷⁾. Given the absorbed dose, D , we shall assume t_i known. The main difference between Eq(6) and the q expression necessary for cycling cells is that the recovery constants, t_{0i} , change with cell cycle stage, i . The new expression is derived below:

To be more specific we shall be concerned for the moment with the factor- q_{14} describing the repair and interaction of two sublesions, one produced in G1 ($i=1$) and the other in M ($i=4$), see Fig. 1. We apply Eq.(4). The distribution of time intervals, $h(t)$, is given by

$$h(t) = \int f(u) f(u+t) du \quad (7)$$

where $f(u)$ is the dose rate (assumed constant), see Figure 1. Note that $u \in [0, t_1]$ and $t \in [t_M - (u + t_4), t_M - u]$, where $t_M = t_1 + t_2 + t_3 + t_4$. From Eq.(4):

Figure 1: Graphical representation of the parameters characterizing the radiosensitivity of the cells in different stages of the mitotic cycle.

$$q_{14} = \int_0^{t_1} du \int_{t_M - u - t_4}^{t_M - u} \exp\left(-\frac{t_1 - u}{t_{01}} - \frac{t_2}{t_{02}} - \frac{t_3}{t_{03}} - \frac{t + u + t_1 - t_M}{t_{04}}\right) dt \quad (8)$$

This can be readily integrated. The result is:

$$q_{14} = \frac{t_{01} t_{04}}{t_1 t_4} \left[1 - e^{-\frac{t_1}{t_{01}}}\right] \left[1 - e^{-\frac{t_4}{t_{04}}}\right] e^{-\sum_{i=2}^3 \frac{t_i}{t_{0i}}} \quad (9)$$

The general expression is:

$$q_{ik} = \frac{t_{0i} t_{0k}}{t_i t_k} e^{-\sum_{l=i+1}^{k-1} \frac{t_l}{t_{0l}}} \left[1 - e^{-\frac{t_i}{t_{0i}}}\right] \left[1 - e^{-\frac{t_k}{t_{0k}}}\right] \quad (10)$$

Eq.(5) is replaced with:

$$-\ln S(D) = \sum_i \left[\alpha_i D_i + \beta_i q\left(\frac{t_{i0}}{t_i}\right) D_i^2 \right] + 2 \sum_{\substack{i,k \\ i < k}} q_{ik} (\beta_i \beta_k)^{1/2} D_i D_k \quad (11)$$

We have chosen the HeLa cells for this analysis because it is a human cervix cell line and because its cell kinetics is known. Terasima and Tolmach⁽⁸⁾ were the first to report on the variation in radiosensitivity of cells exposed to radiation during the cell cycle. Mitchell et. al.⁽⁹⁾ have measured survival curves for HeLa cell cultures synchronized initially in mitosis and irradiated at dose rates of 0.37, 0.74 and 1.54 Gy/hr. From the data of Mitchell et. al.⁽⁹⁾, we

have obtained the phase-specific radiosensitivity parameters, α and β , and also the reduction factor q . The repair time constant t_0 was taken as 1.43 hr⁽⁹⁾. Table 1 shows these parameters for each phase of the cell cycle.

Table I Phase-specific radiosensitivity parameters used to calculate the survival curves for HeLa cells.

phase	α (Gy ⁻¹)	β (Gy ⁻²)
G1	0.22	0.290
S	0.21	0.050
G2	0.20	0.048
M	0.60	0.005

Griffith and Tolmach⁽¹⁴⁾ have examined the response of HeLa cells before and after the arrest point in G2 phase. These authors have found that HeLa cells undergo an abrupt increase in sensitivity at or near the arrest point, and the radiosensitivity of the cells beyond this point is close to that of mitotic cells. Based on these results, we have assigned the radiosensitivity of mitotic cells to G2 cells that have passed the arrest point, i.e. about 2 hr after the S/G2 boundary. Under continuous low dose-rate irradiation a block at G2+M phase has been experimentally demonstrated for HeLa cells by Mitchell et. al.⁽⁹⁾ In this study, a linear relation of 3.0 hr per Gy/hr between G2 block duration and dose rate was used for V79 cells.

In a separate analysis⁽⁷⁾ we have developed a mathematical model for cell-cycle progression under continuous low dose-rate irradiation, and applied this model to the data of Mitchell et. al.⁽⁹⁾ Phase duration obtained from this analysis are given in Table 2.

Table II Phase durations (hr) for HeLa cells at different dose rates.

phase	0.37 Gy/hr	0.74 Gy/hr	1.54 Gy/hr
G1	10.7	8.8	8.3
S	9.2	8.9	10.2
G2	45.2	39.3	64.0
M	8.9	12.9	10.8

The choice of repair times, t_0 , is difficult. Analyses of human cell lines survival give inconsistent results⁽¹⁵⁾. For instance, for human cervix carcinoma cells, repair time constants ranging from 0.5 to 8.0 hr. have been reported⁽¹⁶⁾. A variation of t_0 with the phase of the cell cycle has been observed for V79 cells⁽¹⁷⁾. Similar information on human cells does not appear to be available. To demonstrate the effect of repair time on cellular inactivation, we have used repair time constants ranging from 0.1 to 7 hrs in these calculations.

Results and Discussion

Fig. 2 Survival probability for synchronous cells irradiated at a dose rate of 0.74 Gy/hr. The result from our model (solid line) is compared with the result obtained if synergistic term is neglected (dashed line). The parameters used in this calculation are indicated in Table 1. The same repair time constant was used for all stages of cell cycle: $t_0 = 3$ h.

Figure 2 shows calculated survival curves for synchronous HeLa cells irradiated at low dose-rate (0.74 Gy/hr) with repair time constant $t_0 = 3$ hr. The parameters used for this calculation are listed in Table 1 and 2. Our model (solid line) shows a lower survival probability than the result obtained in a calculation (dashed line) where the inter-phase interaction between sublesions is ignored. The calculation was performed for synchronous cells in order to demonstrate more closely these effects. The sudden changes of slope show the positions on the survival curves where cells change phase during exposures. As the dose-rate increases, the probability of sublesion repair decreases, resulting in an increased inter-phase interaction among sublesions, and therefore a larger difference in survival response between the two models.

Fig. 3 The points represent surviving fraction of HeLa cells from Mitchell et. al.⁽⁹⁾ for various dose rates. The lines represent fits to the data obtained from our model.

In Fig. 3 we compare our model with the data of Mitchell et. al.⁽⁹⁾ At each of the 3 dose rates employed (0.37, 0.74 and 1.54 Gy/h) the effect of radiosensitivity change is apparent. We note the inverse dose-rate effect at $D > 12$ Gy for 0.37 Gy/h and 0.74 Gy/h obtained both experimentally and theoretically. The quality of the data are such that we cannot further assess the contribution to this inverse dose-rate effect of changes in radiosensitivity versus changes in repair constants, during the mitotic cycle.

Fig. 4 shows the effect of repair time on the surviving probability with and without the inter-phase interactions between sublesions. As the recovery constant, t_0 , increases, the difference in survival response between the models increases, as expected. At 0.74 Gy/hr, there is no difference in survival between the two models for $t_0 = 0.1$ hr. For $t_0 = 7$ hr, however, the difference may be as much as two orders of magnitude (e.g. at 15 Gy). This difference will further increase as dose rate increases.

In conclusion:

1. A consistent application of the dual radiation action mechanism (under whatever label the investigator may want to use it, e.g. the "incomplete repair model⁽¹⁸⁾") requires an expression in which sublesion interaction and repair is explicitly taken into account for all the phases of the mitotic cycle. We have provided a mathematical description of this statement.
2. For a variety of practical situations (dose rate, cell lines, dose) the neglect of a) the variation of radiosensitivity through the mitotic cycle, b) variation of repair ability through the cycle, and c) inter-phase sublesion interaction may lead to

significant errors.

3. This study has underlined several cellular parameters that need further experimental investigation before a more precise calculation can be performed. For example, the phase-specific radiosensitivity parameters (α and β) and repair time constant (t_0) need to be measured for various cell lines. The latter is very difficult to measure, and currently no reliable information exists.

The authors would like to thank Drs. W.K. Sinclair, R. Nath, C.S. Lange and K.T. Wheeler for helpful discussions.

References

1. Hall, E. J. Radiobiology for the Radiologist, 3rd ed. Lippincott, Philadelphia (1988).
2. Mitchell, J.B. Bedford J. S. and Bailey S. M. Dose-rate effects in mammalian cells in culture III. Comparison of cell killing and cell proliferation during continuous irradiation of six different lines. Radiat. Res. **79**, 520-536 (1979).
3. Zaider, M. and Wu, C. S. The biological effects of mixed-radiation fields on cellular system of variable radiosensitivity (OER and cell-cycle stage). In proceedings of the international workshop on macro and microdosimetry and treatment planning for neutron capture therapy, 1991.
4. Lea, D. E. and Catcheside, D. G. The mechanism of the induction by radiation of chromosome aberration in tradescantia. J. genet. **44**, 216-245 (1942).
5. Kellerer, A. M. and Rossi, H. H. The theory of dual radiation action. Curr. Top. Radiat. Res. **8**, 85-158 (1972).

6. Zaider, M. and Rossi, H. H. The synergistic effects of different radiations. *Radiat. Res.* **83**, 732-739 (1980).
7. Zaider, M and Minerbo, G. N. A mathematical model for cell cycle progression under continuous low dose-rate irradiation. *Radiat. Res.* (in press), (1992).
8. Terasima, T. and Tolmach, L. J. Variations in several responses of HeLa cells to x-irradiation during the division cycle. *Biophys. J.* **3**, 11-33 (1963).
9. Mitchell, J. B., Bedford, J. S. and Bailey, S. M. Dose-rate effects on the cell cycle and survival of S3 HeLa and V79 cells. *Radiat. Res.* **79**, 520-536 (1979).
10. Painter, R. B. and Roberson, J. S. Effect of irradiation and theory of role of mitotic delay on the time course of labelling of HeLa S3 cells with tritiated thymidine. *Radiat. Res.* **11**, 206-217 (1959).
11. Weiss, B. G. Perturbations in precursor incorporation into DNA of x-irradiated HeLa S3 cells, *Radiat. Res.* **20**, 600-618 (1963).
12. King, C. R. , Nath, R. and Rockwell, S. Effects of continuous low dose-rate irradiation: Computer simulation, cell tissue kinet. **21**, 339-351 (1988).
13. Dillehay, L. E. A model of cell killing by low-dose-rate radiation including repair of sublethal damage, G2 block, and cell division. **124**, 201-207, (1990).
14. Griffith, T. D. and Tolmach, L. J. Lethal response of HeLa cells to x-irradiation in the latter part of the generation cycle. *Biophys. J.* **16**, 303-318 (1976).
15. Steel, G. G. , Deacon, J. M. , Duchesne, G. M., Horwich, A., Kelland, L. R. and Peacock, J. H. The dose-rate effect in human tumor cells. *Radiother. Oncol.* **9**, 299-310 (1987).
16. Kelland, L. R. and Steel, G. G. Differences in radiation response among human cervix carcinoma cell line. *Radiother. Oncol.* **13**, 225-232 (1988).
17. Reddy, N. M. S., Mayer, P. J. and Lang, C. S. The saturated repair kinetics of Chinese hamster V79 cells suggests a damage accumulation -interaction model of cell killing. *Radiat. Res.* **121**, 304-311 (1990).
18. Thames, H. D. An "incomplete-repair" model for survival after fractionated and continuous irradiation. *Int. J. Radiat. Biol.* **47**, 319-339 (1985).
19. Segal, L. The infinite and the infinitesimal in models for natural phenomena. *Rev. Mod. Phys.* **63**, 225- 238 (1991).

A CHROMOSOMAL "FINGERPRINT" FOR PAST EXPOSURE TO DENSELY-IONIZING RADIATION

D. J. Brenner, R. K. Sachs (University of California, Berkeley)

We suggest, and give evidence that densely-ionizing radiations such as alpha particles from radon daughters, or fission neutrons, leave a distinctive chromosomal "fingerprint" which may be detected long after radiation exposure. Specifically, they produce an anomalously low ratio of inter-chromosome to intra-chromosome inter-arm exchange-type chromosome aberrations, in comparison with either x rays or chemical carcinogens. Such a distinctive and measurable marker would have significant scientific, legal, and societal implications.

Introduction

Several groups of people, such as nuclear power workers, inhabitants of high-level radon areas, uranium miners, and persons exposed to A-bomb tests, have potentially been exposed to significant doses of densely-ionizing radiation, such as alpha particles or neutrons.¹ Consequently, determination of whether individuals, particularly those who later developed cancer, did in fact receive significant radiation doses, is an important societal and legal issue. Thus there has been considerable interest in finding a detectable biological "fingerprint" preferentially produced by densely-ionizing radiation. Several possible candidates have been suggested²⁻⁴ but no corresponding mechanistic basis has as yet been outlined.

Nature of the chromosomal "fingerprint"

On mechanistic grounds, the ratio of inter-chromosome to intra-chromosome exchange-type chromosome aberrations should be, and is, significantly different after exposure to densely-ionizing radiations, compared with sparsely-ionizing radiations such as x rays, or other environmental carcinogens.

Exchange-type chromosomal aberrations are formed by interaction between pairs of DNA double-strand breaks (DSB), either on different chromosomes, or on different arms of the same chromosome, or on a single chromosome arm (see Table 1). The non-lethal chromosomal aberrations are stable and can be measured⁵⁻⁷ in irradiated cells and their progeny many decades after exposure.

Table 1. Relevant chromosomal aberration types. The F value is the ratio of the second column yield to the third.

	<i>Inter-chromosomal</i>	<i>Intra-chromosomal Inter-arm</i>
<i>Stable (non lethal)</i>	Translocations	Pericentric Inversions
<i>Unstable (lethal)</i>	Dicentrics	Centric Rings

If all DSB produced in a cell nucleus were equally likely to interact with one another, the ratio (F) of inter-chromosomal to intra-chromosomal, inter-arm aberrations would be approximately the number of chromosomal arms (92 in humans). However it has long been

known⁸ that pairs of DSB on different chromosomes that are distant from each other within the nucleus are less likely to interact, resulting in a decrease in the F value. For example, x-ray exposure, which produces DSB roughly uniformly, yields F values around 15 (see below).

Densely-ionizing radiations, however, exhibit a unique property that reduces the F value even further. These radiations produce spatially inhomogeneous energy depositions, and thus DSB, that are much closer together than those produced by sparsely-ionizing radiations such as x rays, or chemical carcinogens. Consequently it is expected that intra-chromosomal aberrations would be further increased relative to inter-chromosomal aberrations, the resulting smaller F value representing a "fingerprint" of densely-ionizing radiation.

Experimental data

There is much data in the literature, both *in vivo* and *in vitro*, on yields of inter- and intra-chromosomal aberrations. Often, F values are not well determined, because of the smaller number of intra-chromosomal aberrations produced. Where adequate data are available, as we discuss, suggest that densely-ionizing radiation does produce characteristically low F values.

Some of the data refer to stable aberrations, whilst others refer to lethal aberrations. In terms of a detectable chromosomal "fingerprint," only the long-lasting stable aberrations are of interest. However, it would be expected^{6,7} and there is evidence (see below) that F values for unstable aberrations follow the same pattern as those for stable aberrations, and thus may be used to augment the stable-aberration results.

In vivo, there are many measurements of F values after accidental or radiotherapeutic exposures to sparsely-ionizing radiation. For Chernobyl-region inhabitants, a ratio of 37 ± 19 was measured;⁹ for individuals exposed to γ rays in the Goiania accident, the measured ratio was 20 ± 3 ;¹⁰ for radiotherapy patients, where data are adequate for analysis, the mean F value is 18 ± 9 ;¹¹ these data deriving from both stable and unstable aberrations. For densely-ionizing radiation, the most extensive *in-vivo* data are for individuals injected with the alpha-particle-emitting contrast agent Thorotrast: the largest study yields $F = 5.0 \pm 0.3$,¹² and smaller studies report similar numbers. Measurements on workers exposed to plutonium (emitting alpha particles) yield $F = 5.6 \pm 3.0$ for stable aberrations, and $F = 4.5 \pm 2.0$ for unstable aberrations.¹³ Finally, measurements¹⁴ on workers accidentally exposed to radiation consisting of ~88% (by dose equivalent) densely-ionizing neutrons yielded $F = 5.7 \pm 3.5$ for stable aberrations, and 5.0 ± 2.4 for unstable aberrations.

In vitro, several very large-scale studies have been conducted. The largest study for sparsely-ionizing radiation yielded¹⁵ $F = 16.7 \pm 0.9$, in agreement with the value of 16 ± 5 obtained from a literature survey.¹⁶ By contrast, analysis of human lymphocyte cells exposed to densely-ionizing neutrons¹⁷ gave a significantly smaller value of $F = 5.6 \pm 0.5$.

Discussion

In summary, F values *in-vitro* are consistent with those measured after irradiation *in vivo*, both showing significantly smaller values for densely-ionizing radiations compared with x or γ rays. The data suggest that an F value of around 5 is characteristic of densely-ionizing radiation, in contrast to significantly larger values for x or γ rays.

Although chemical carcinogens often produce damage at specific sites within the genome, such damage is unlikely to be preferentially located on pairs of sites on opposite arms of one chromosome.¹⁸ Chemical carcinogens would therefore not be expected to produce anomalously low F values; although few adequate data sets are available, a typical value is 30 ± 5 for bleomycin-induced damage *in vitro*.¹⁹

Using F values from stable aberrations, possibly measured long after exposure, presupposes no differential loss over time of translocations relative to pericentric inversions. No data are available on the relative loss rates of these aberrations. However, the structural similarities between these aberration types suggest that differential loss would be unlikely.

F values for stable chromosome aberrations in individuals can be measured using Giemsa staining techniques,⁵⁻⁷ and faster measurement techniques can be developed using fluorescent in-situ hybridization.⁷ The results should provide stronger evidence than currently possible as to whether an individual was exposed to a significant dose of densely-ionizing radiation.

This chromosomal "fingerprint" would facilitate epidemiological studies investigating the relationship between radiation exposure and cancer. "Molecular epidemiology" is a powerful tool for chemical risk assessment,²⁰ and the same techniques could be applied to epidemiological studies of, for example, radon effects.

References

1. Ionizing Radiation: Sources and Biological Effects (United Nations, New York, NY, 1982)
2. Vahakangas KH, et al. Mutation of p53 and ras genes in radon-associated lung cancer from uranium miners. *Lancet* 1992; **339**: 576-580.
3. Kadhim KA, et al. Transmission of chromosomal instability after plutonium α -particle irradiation. *Nature* 1992; **355**: 738-740.
4. Aghamohammadi SZ, Goodhead DT, Savage JRK, Induction of sister chromatid exchanges (SCE) in G₀ lymphocytes by plutonium-238 α -particles. *Int J Radiat Biol* 1988; **53**: 909-915.
5. Evans HJ, Buckton KE, Sumner A, Cytological mapping of human chromosomes. *Chromosoma* 1971; **35**: 310-325.
6. Savage JRK, Papworth DG, Frequency and distribution studies of asymmetrical versus symmetrical chromosome aberrations. *Mutat Res* 1982; **95**: 7-18.
7. Lucas JN, et al. Rapid translocation frequency analysis in humans decades after exposure to ionizing radiation. *Int J Radiat Biol* 1992; **62**: 53-63.
8. Sax K, An analysis of X-ray induced chromosomal aberrations in Tradescantia. *Genetics* 1940; **25**: 41-68.
9. Stephan G, Oestreicher U, An increased frequency of structural chromosome aberrations in persons present in the vicinity of Chernobyl during and after the reactor accident. *Mutat Res* 1989; **223**: 7-12.
10. Ramalho AT, Nascimento ACH, The fate of chromosomal aberrations in ¹³⁷Cs-exposed individuals in the Goiania radiation accident. *Health Phys* 1991; **60**: 67-70.
11. Zaslav A-L, Stamberg J, Shende A, Persistent chromosome damage induced by localized radiotherapy for lymphoma. *Cancer Genet Cytogenet* 1988; **30**: 245-251. Stephan G, Diener A,

- Vogl T, Chromosome aberrations in peripheral lymphocytes of patients irradiated with 15-MeV photons for Morbus Hodgkin. *Strahlenther Onkol* 1990; **166**: 460-465. Martin RH, et al. A comparison of chromosomal aberrations induced by in vivo radiotherapy in human sperm and lymphocytes. *Mutat Res* 1989; **226**: 21-30.
12. Sasaki MS, et al. Chromosome aberration frequency and radiation dose to lymphocytes by alpha-particles from internal deposit of Thorotrast. *Radiat Environ Biophys* 1987; **26**: 227-238.
 13. Tawn EJ, Hall JW, Schofield GB, Chromosome studies in plutonium workers. *Int J Radiat Biol* 1985; **47**: 599-610.
 14. Bender MA, Gooch PC, Persistent chromosome aberrations in irradiated human subjects. *Radiat Res* 1962; **16**: 44-53. 1963; **18**: 389-396.
 15. Lloyd DC, Edwards AA, Prosser JS, Chromosome aberrations induced in human lymphocytes by *in vitro* acute x and gamma radiation. *Radiat Protec Dosim* 1986; **15**: 83-88.
 16. Hlatky L, Sachs RK, Hahnfeldt P, The ratio of dicentrics to centric rings produced in human lymphocytes by acute low-LET radiation. *Radiat Res* 1992; **129**: 304-308.
 17. Pohl-Ruling J, et al. Chromosomal damage induced in human lymphocytes by low doses of D-T neutrons. *Mutat Res* 1986; **173**: 267-272.
 18. Kihlman BA. Actions of chemicals on dividing cells. New Jersey: Prentice-Hall, 1966:143-157.
 19. Dresch J, Schmid E, Bauchinger M, The cytogenetic effect of bleomycin on human peripheral lymphocytes in vitro and in vivo. *Mutat Res* 1978; **56**: 341-353.
 20. Perera FP, Santella RM, Brenner D, Young TL, Weinstein IB. Applications of biological markers to the study of lung cancer causation and prevention. *IARC Sci Publ* 1988; **89**: 451-459.

CORRELATIONS BETWEEN α/β AND $T_{1/2}$: IMPLICATIONS FOR CLINICAL BIOLOGICAL MODELLING

D. J. Brenner

There seems little question that future directions in radiotherapy will involve altered or customized fractionation schedules (Cooper, 1990). Recent interest in accelerated and hyper-fractionation, high dose-rate brachytherapy, and the search for predictive assays for tumor proliferation, all exemplify this trend.

The linear-quadratic model, the formalism for which was first described by Lea and Catcheside (1942), is now the modelling tool of choice (e.g. Fowler, 1989) for investigating alternative fractionation schedules. In this model the biological effect, E , of an acute dose of radiation, D , is related to a combination of dose and dose-squared terms:

$$E/\beta = (\alpha/\beta) D + D^2 \quad (1)$$

where α/β is an empirically-determined parameter. When the dose is protracted (with fractionation or low dose rate), there is time for sublethal damage repair to act on the dose-squared component, but the linear dose term will be unchanged. Thus the dose-squared term is reduced and the effect is now related to

$$E/\beta = (\alpha/\beta) D + G D^2 \quad (2)$$

where the quantity G (≤ 1) can be calculated from the particular fractionation / LDR scheme used, and a parameter to represent the speed of sublethal damage repair. For repair governed by first-order kinetics, this parameter can be represented by one (or more) half times for sublethal damage repair, $T_{1/2}$.

Thus, in principle, knowing α/β and $T_{1/2}$ (and/or their distributions), it is possible to investigate the equivalence of any fractionation scheme to any other, either for early or for late effects. There are, in fact, some situations where α/β but not $T_{1/2}$ need be known - in particular, a fractionation schedule with many hours between treatments, where time for full sublethal damage repair is allowed; for brachytherapy, or accelerated or hyper-fractionated regimes, however, both parameters need to be known.

Note that, in general, we also need to consider proliferation during the overall times of the treatments; here, for illustrative purposes, we avoid this by assuming that the various treatments take the same overall time.

Available data for α/β and $T_{1/2}$ derived from results in the clinic have been summarized by Thames et al (1990), for normal tissues in animals by Thames and Hendry (1987), and for animal tumors by Williams et al (1985) and Guttenberger et al (1990). *In-vitro* data for $[\alpha, \beta, T_{1/2}]$ from cells of human origin have been reviewed by Brenner and Hall (1991). Figure 1 indicates that, for the parameter α/β , the human, animal, and *in-vitro* data appear consistent with each other, all showing a very broad range of values.

Figure 1: Comparison of α/β values derived from cells of human origin in vitro (shaded histogram; Brenner & Hall, 1991), from clinical data (full curve: log-normal distribution fitted to data for human tumors from review by Thames et al (1990)), and from animal tumor data (dashed curve: log-normal distribution fitted to data for clamped tumors from review by Williams et al (1985)).

Figure 2: Summary of α/β and $T_{1/2}$ data for normal tissues in animals, where these parameters were obtained together. Data are from Thames et al, 1984, Dale et al, 1988, Dewit et al, 1987, Ang et al, 1985, Vegesna et al, 1985, Vegesna et al, 1989, Travis et al, 1987, Steel et al, 1989, Ang et al, 1987, Thames et al, 1988, Scalliet et al, 1989, Stuben et al, 1991, Rojas et al, 1989, and Travis et al, 1985. The p value refers to the probability, estimated using Kendall's τ test (Kendall, 1938), that α/β and $T_{1/2}$ are independent.

only one published report in which both α/β and $T_{1/2}$ were obtained from the same clinical results; this was from Turesson and Thames (1989) for post-mastectomy irradiation. For normal tissues in animals, there are at least 16 data sets in which α/β and $T_{1/2}$ were obtained together (Thames et al, 1984; Dale et al, 1988; Dewit et al, 1987; Ang et al, 1985; Vegesna et al, 1985; Vegesna et al, 1989; Travis et al, 1987; Steel et al, 1989; Ang et al, 1987; Thames et al, 1988; Scalliet et al, 1989; Stuben et al, 1991; Rojas et al, 1989; Travis et al, 1985). The data are shown in Fig. 2. Only two such data sets are known for animal tumors, from Guttenberger et al (1990) and Rojas et al (1989). *In vitro*, where experiments may be performed rather more easily and with better statistics, 36 paired data sets for $[\alpha/\beta, T_{1/2}]$ from cells of human origin were reviewed by Brenner and Hall (1991).

The normal-tissue animal data in Fig. 2 show an *inverse correlation* - as α/β increases, $T_{1/2}$ decreases. Appropriate non-parametric (i.e. model independent) tests for correlation are the Spearman rank test (Spearman 1904) or the Kendall τ test (Kendall 1938). For these normal-tissue animal data, both tests indicate that the probability of α/β and $T_{1/2}$ being independent is less than 0.1%. On the other hand, the animal-tumor data of Guttenberger et al (1990) and Rojas et al (1989), though limited in number for correlation analysis, do not show

this inverse correlation, and suggest a possible *positive correlation* (significant at 80% confidence level).

In light of the highly-significant correlation found in the normal-tissue animal data, we have investigated possible relationships between α/β and $T_{1/2}$ in *in-vitro* data for cells of human origin. The data for cells of human origin are shown in Figs. 3 and 4. Figure 3 shows these *in-vitro* data for *normal* cells of human origin and, using the non-parametric tests described above, we can say, at the 90% confidence level, that the data are *inversely correlated* (the probability that the parameters are independent is 9.2%). Figure 4 shows the corresponding *in-vitro* data for *tumor* cells of human origin. In this case we can say with 90% (or 95%) confidence that the parameters α/β and $T_{1/2}$ have a *positive correlation* (the probability that they are independent is 4.6%).

Thus both the *in-vivo* and the *in-vitro* data appear to suggest that α/β and $T_{1/2}$ are correlated - an inverse correlation for normal cells, and a positive correlation for tumor cells. As pointed out by Ang et al (1987), if all other factors were kept constant (the effect, E , and the quadratic term, β), then a fluctuation (say upward) in the regression estimate of α/β would necessitate a compensating downward fluctuation in the estimate of G (see Equation (2)), and hence in $T_{1/2}$, thus making for an inverse correlation. However, there is no reason to suppose that the estimate of β would not also fluctuate in such estimates, which would eliminate this "curve-fitting" explanation of the observed correlations. It should be noted that maximum-likelihood methods were used to obtain most of the data sets discussed here (Thames et al, 1986, Brenner & Hall, 1991), and it is thus unlikely that these correlations are spuriously produced by the data fitting procedures used.

If these correlations are real, they may well represent the influence of a third "hidden" variable. For example, cellular proliferation - and thus the fraction of cells in different parts of

the cell cycle - is often considered to be a determinant of the α/β ratio (Hall, 1988), and it is quite plausible to suppose that repair rates will also be different in different parts of the cell cycle, due to such factors as chromatin structure and enzymatic availability.

Apart from the mechanistic questions raised, there are important methodological implications for the use of the linear-quadratic model in designing alternative fractionation regimes. From Fig. 1, it is clear that the use of single "representative" values of α/β is unwise; this is also true for $T_{1/2}$, which has an even broader range of measured values (Brenner & Hall, 1991). Prima facie, a reasonable approach would be to consider an appropriate range of α/β values and $T_{1/2}$ values, and repeat the calculations for parameter values covering these ranges; this is implicitly done, for example, by Fowler (1989; 1990), Dale (1986) and Thames and Hendry (1987); at that point the results might be averaged, or the most conservative accepted, or the results examined to see which are in best accord with clinical data.

Such an approach, however, will not work if the α/β and $T_{1/2}$ really are correlated. If this is the case, as the discussions above seem to suggest, then only paired sets of parameters, i.e. [$\alpha/\beta, T_{1/2}$] sets obtained from analysis of the same data set, should be used. This approach will certainly decrease the number of parameter sets that can be used for such calculations, but should lead to more reliable design of new fractionated and brachytherapy regimens.

References

- Ang, K.K., Thames, H.D., van der Kogel, A.J. & van der Schueren, E., 1987. Is the rate of repair of radiation-induced sublethal damage in rat spinal chord dependent on the size of dose per fraction? *International Journal of Radiation Oncology, Biology and Physics*, 13, 557-562.
- Ang, K.K., Xu, F.-X., Landuyt, W., & van der Schueren, E., 1985. The kinetics and capacity for repair of sublethal damage in mouse lip mucosa during fractionated irradiations. *International Journal of Radiation Oncology, Biology and Physics*, 11, 1977-1983.
- Brenner, D.J. & Hall, E.J., 1991. Conditions for the equivalence of continuous to pulsed low dose rate brachytherapy. *International Journal of Radiation Oncology, Biology and Physics*, 20, 181-190.
- Cooper, J.S., 1990. Will altered fractionation schemes alter the future? *International Journal of Radiation Oncology, Biology and Physics*, 19, 1621-1622.
- Dale, R.G., 1986. The application of the linear-quadratic models to fractionated radiotherapy when there is incomplete normal tissue recovery between fractions, and possible implications for treatments involving multiple fractions per day. *British Journal of Radiology*, 59, 919-927.
- Dale, R.G., Trott, K.R. & Huczkowski, J., 1988. Possible dose rate dependence of recovery kinetics as deduced from a preliminary analysis of the effects of fractionated irradiations at varying dose rates. *British Journal of Radiology*, 61, 153-15.
- Dewit, L., Oussoren, Y. & Hart, G., 1987. Modification by cis-diamminedichloroplatinum (II) of the radiation dose response for intestinal crypt cells in mice. *Radiation Research*, 111, 445-456.

- Fowler, J.F., 1989. The linear-quadratic formula and progress in fractionated radiotherapy. *British Journal of Radiology*, 62, 679-694.
- Fowler, J.F., 1990. How worthwhile are short schedules in radiotherapy?: A series of exploratory calculations. *Radiotherapy and Oncology*, 18, 165-181.
- Guttenberger, R., Kummermehr, J. & Chmelevsky, D., 1990. Kinetics of recovery from sublethal damage in four murine tumors. *Radiotherapy and Oncology*, 18, 79-88.
- Hall, E.J., 1988. *Radiobiology for the Radiologist* (J. P. Lippincott, Philadelphia).
- Kendall, M.G., 1938. A new measure of rank correlation. *Biometrika*, 30, 81-93.
- Lea, D.E. & Catcheside, D.G., 1942. The mechanism of the induction by radiation of chromosome aberrations in *Tradescantia*. *Journal of Genetics*, 44, 216-245.
- McNally, N.J., 1989. Can cell kinetic parameters predict the response of tumours to radiotherapy? *International Journal of Radiation Biology*, 56, 777-786.
- Rojas, A., Joiner, M.C., & Johns, H., 1989. Recovery kinetics in mouse skin and CaNT tumours. *Radiotherapy and Oncology*, 16, 211-220.
- Scalliet, P., Landuyt, W. & van der Schueren, E., 1989. Repair kinetics as a determining factor for late tolerance of central nervous system to low dose rate irradiation. *Radiotherapy and Oncology*, 14, 345-353.
- Spearman, C., 1904. The proof and measurement of association between two things. *American Journal of Psychology*, 15, 72-101.
- Steel, G.G., Kelland, L.R. & Peacock, J.H., 1989. The Radiobiological basis for low dose rate. In *Brachytherapy 2*, ed. by R.F. Mould (Nucletron, The Netherlands).
- Stuben, G., Landuyt, W., van der Schueren, E., van der Kogel, A. & Reijnders, A., 1991. Estimation of repair parameters in mouse lip mucosa during continuous and fractionated low dose-rate irradiation. *Radiotherapy and Oncology*, 20, 38-45.
- Thames, H.D., Ang, K.K., Stewart, F.A. & van der Schueren, E., 1988. Does incomplete repair explain the apparent failure of the basic LQ model to predict spinal cord and kidney responses to low doses per fraction? *International Journal of Radiation Biology*, 54, 13-19.
- Thames, H.D., Bentzen, S.M., Turesson, I., Overgaard, M. & van den Bogaert, W., 1990. Time-dose factors in radiotherapy: a review of the human data. *Radiotherapy and Oncology*, 19, 219-235.
- Thames, H.D. & Hendry, J.H. *Fractionation in Radiotherapy*. London: Taylor and Francis; 1987.
- Thames, H.D., Rozell, M.E., Tucker, S.L., Ang, K.K. Fisher, D.R. & Travis, E.L., 1986. Direct analysis of quantal response data. *International Journal of Radiation Biology*, 49, 999-1009.

- Thames, H.D., Withers, H.R. & Peters, L.J., 1984. Tissue repair capacity and repair kinetics deduced from multifractionated or continuous irradiation regimens with incomplete repair. *British Journal of Cancer* 49, Suppl. VI, 263-269.
- Travis, E.L., Peters, L.J., McNeill, J., Thames, H.D. & Karolis, C., 1985. Effect of dose-rate on total body irradiation: Lethality and pathologic findings. *Radiotherapy and Oncology*, 4, 341-351.
- Travis, E.L., Thames, H.D., Watkins, T.L. & Kiss, I., 1987. The kinetics of repair in mouse lung after fractionated irradiation. *International Journal of Radiation Biology*, 52, 903-919.
- Turesson, I. & Thames, H.D., 1989. Repair capacity and kinetics of human skin during fractionated radiotherapy, erythema, desquamation, and telangiectasia after 3 and 5 year's follow-up. *Radiotherapy and Oncology*, 15, 169-188.
- Vegesna, V., Withers, H.R., Thames, H.D. & Mason, K.A., 1985. Multifraction radiation response of mouse lung. *International Journal of Radiation Biology*, 47, 413-422.
- Vegesna, V., Withers, H.R. & Taylor, J.M., 1989. Repair kinetics of mouse lung. *Radiotherapy and Oncology*, 15, 115-123.
- Williams, M.V., Denenkamp, J. & Fowler, J.F., 1985. A review of α/β ratios for experimental tumors: implications for clinical studies of altered fractionation. *International Journal of Radiation Oncology, Biology and Physics*, 11, 87-96.

CELLULAR AND MOLECULAR BIOLOGY

CELL-CYCLE DEPENDENT RADIATION-INDUCED TRANSFORMATION OF C3H 10T1/2 CELLS

Richard C. Miller, Charles R. Geard, Marcus J. Geard and Eric J. Hall

Introduction

Changes in radiosensitivity of cells exposed to X rays as a function of position in the cell cycle are well documented; with cell lethality as an endpoint, cells are most sensitive to X-rays in G₂/M and most resistant in late S. For cell lines with a long G₁ phase, there is often a second period of resistance in G₁. For exposure to neutrons the pattern of radiosensitivity as a function of cell age is qualitatively similar but the magnitude of the changes is smaller.

The importance of a variation of radiosensitivity for oncogenic transformation with phase of the cell cycle in C3H 10T1/2 cells is of considerable interest because of biophysical models that have been proposed recently to explain the so-called inverse dose/rate effect for neutrons. Various experimenters, including ourselves, have shown that low doses and low dose-rates of certain radiations of intermediate and high LET result in a higher incidence of oncogenic transformation than the same dose delivered in an acute exposure. To account for this anomalous effect, a biophysical model was proposed by Rossi and Kellerer (1), which postulated the existence of a narrow "window" of sensitivity. The present series of experiments were designed to test this hypothesis.

Materials and Methods

Cell culture conditions

C3H 10T1/2 clone 8 cells between passages 8-14 were grown in Eagle's basal medium (BME) and 10% calf serum supplemented with iron plus gentamycin to minimize bacterial contamination. Calf serum with iron supplement was selected after tests showed it capable of supporting uniform clonal growth and transformation induction of cells exposed to 6 Gy of X rays.

Cell synchrony techniques

Cells were made synchronous by the mitotic shake-off method. Briefly, one day prior to the shake-off procedure, stock C3H 10T1/2 cells were plated into 150 cm² culture flasks at 2 x 10⁶ cells/flask. Then culture flasks were struck firmly to dislodge poorly attached cells and the relatively loose and rounded cells held in M phase. This collection was discarded. Medium with serum was then added back to the flasks and returned to the incubator for 30 mins. Flasks were again removed from the incubator and firmly struck to dislodge mitotic cells and the supernatant was collected and placed in an ice bath. Complete medium was added back to the cell culture flasks which were returned to the incubator for an additional 30 mins. and this cycle was repeated on three more occasions.

Synchrony and cell age distribution evaluation

At the time of mitotic shake off a small aliquot of cells was placed on a microscope slide and examined to assess mitotic purity. Additionally, aliquots of cells were placed into single chamber Nunc slide flasks and, after 10 mins. at 37°C, bromodeoxyuridine (Brd Urd) was added. At each time interval at which cells were irradiated, one flask was removed from the incubator, medium gently discarded and ice cold 3 parts methanol:1 part acetic acid carefully added to fix cells *in situ*. Cells were stained in 0.3% Giemsa stain in 0.3M Na₂HPO₄ at pH 10.3 for up to 60 mins. This staining procedure allows for differentiation between cells that have incorporated Brd Urd into their DNA versus cells without such incorporation. Cells from each irradiation time as well as additional intermediate times, were examined for the fraction of cells in mitosis, the number of cells in different mitotic stages, the proportion of paired early G₁ cells, and the fraction of cells showing Brd Urd incorporation. From these observations the mitotic index and the cumulative fraction of S phase cells was determined allowing for assessments of actual frequencies of synchronized mitoses and of the different stages of the cell cycle.

Irradiation procedures

Cells in 100 mm dishes were irradiated with 3 Gy of X rays at various times from 0 to 18 hrs post mitotic collection. In the case of the initial (0 time) irradiation, cells were in suspension. Cells were attached at the later irradiation times. Irradiation was carried out using a Westinghouse 250 kVp X ray machine at room temperature at a dose rate of 0.82 Gy/min.

Clonogenic and oncogenic transformation assay

Within 2 mins. of the completion of irradiation, cells were removed from the dishes by trypsinization and replated into 100 mm-diameter culture dishes at cell concentrations estimated to result in either 300 clonogenic cells per dish for the 6 week transformation assay, or 30 clonogens for the 2 week cell-survival assay. At the end of either incubation period, cells were fixed in formalin and stained with Giemsa. Morphologically transformed foci types II and III were identified according to the criteria of Reznikoff et al. (2,3).

Results and Discussion

Cell synchrony post mitotic shake-off

The procedure used to collect non-perturbed mitotic cells results in relatively low yields with a mitotic purity of approximately 80%. At the time of shake-off it was noted that the greater part of the mitoses were paired attached telophase cells. Cells at anaphase and telophase are more rounded with fewer cellular attachments to substrate than cells in prophase or metaphase. It should be noted that this post metaphase category of cells is not seen when spindle inhibitors are used. When cells showing incorporation of Brd Urd were examined it was noted that the first mitotic cells showing incorporation of Brd Urd were present at 6 hours. That is, these cells were in late S phase at the time of shake-off and then incorporated Brd Urd before moving through G₂ into mitosis. Therefore, it is probable that the G₂ period is 5 to 6 hrs. When paired G₁ cells were specifically observed, it was found that a significant fraction of such cells were labelled at 6 hrs. This, therefore, suggests that the G₁ period is about 5-6 hrs. Overall, it is estimated that the average cell cycle time of cycling cells is about 17-18 hrs, which is similar to that previously found for asynchronous populations of C3H

10T1/2 cells using both tritiated thymidine and Brd Urd labelling procedures (unpublished observations).

These estimates of the duration of the various phases of the cell cycle are transferred to Figure 1 where data are shown for cell survival (upper panel) and transformation incidence (lower panel) following a dose of 3 Gy of X rays administered at various times following the harvesting of mitotic cells.

Figure 1 Upper panel: Percentage of cells surviving a dose of 3 Gy as a function of time after mitotic shake-off. Lower panel: Transformants per surviving cell as a function of time after mitotic shake-off. In both cases the error bars are one standard error from the mean. Data were pooled from replicate experiments.

The age response function for survival (upper panel in Figure 1) shows cells to be sensitive in G_2 and M while there are resistant peaks in both G_1 and late S. This is both qualitatively and quantitatively similar to previously published data for cell lines, such as HeLa, that have a significant G_1 period. By contrast the age response function for oncogenic transformation shows a single and radiosensitive period at 14 to 18 hours post shake-off. While only approximate, the half-width of this sensitive period is about 2 hours.

It is not possible to be very precise in delineating the exact time in the cycle at which this sensitivity is expressed, because populations of C3H 10T1/2 cells have a range of cell cycles and synchrony tends to be quickly lost. However, it is clear that cells exposed at time zero, immediately after shake-off, i.e. cells in mitosis or very early G_1 do not show this increased sensitivity. It appears to be cells passing through G_2 , but before they round up for mitosis, that represent the very sensitive cohort.

Overall, for this dose of X rays (3 Gy) there is a ten-fold variation in sensitivity to oncogenic transformation between the most resistant S phase cells and the most sensitive G_2 cells. This compares with a 2-1/2 fold variation in cell

survival between the most sensitive and most resistant phases of the cell cycle.

It is interesting to note from the lower panel of Figure 1 that, when a synchronous population of C3H 10T1/2 cells is irradiated, the vast majority of the transformed foci would appear to derive from cells in a relatively narrow time period of the cell cycle in G_2 . The experimental observation of a window of sensitivity supports the hypothesis of Kellerer and Rossi (1) upon which their model to explain the inverse dose-rate effect was based.

References

1. H.H. Rossi and A.M. Kellerer, The dose rate dependence of oncogenic transformation by neutrons may be due to variation of response during the cell cycle. *Int. J. Radiat. Biol.* 50, 353-361 (1986).
2. C.A. Reznikoff, D.W. Brankow, and C. Heidelberger, Establishment and characterization of a cloned line of C3H mouse embryo cells sensitive to postconfluence inhibition of division. *Cancer Res.* 33:3231-3238 (1973).
3. C.A. Reznikoff, J.S. Bertram, D.W. Brankow, and D. Heidelberger, Quantitative and qualitative studies of chemical transformation of cloned C3H embryo cells sensitive to postconfluence inhibition of cell division. *Cancer Res.* 33, 3239-3249 (1973).

THE INVERSE DOSE-RATE EFFECT FOR ONCOGENIC TRANSFORMATION BY CHARGED PARTICLES

Richard C. Miller, Gerhard Randers-Pehrson, Ludwig Hleber,

Stephen A. Marino, Marcia Richards, and Eric J. Hall

Introduction

Several groups of individuals are exposed over an extended period of time to low doses of ionizing radiation that includes a substantial proportion of intermediate to high LET (Linear Energy Transfer) radiation. Such individuals include radiation workers in certain areas of the nuclear power industry, shuttle astronauts in low earth orbit, as well as the crews of commercial jet airliners.

For most cellular endpoints, lowering the dose-rate, or protracting the exposure over a longer time period by fractionation, leads to a reduction or sparing effect, at least for X or γ rays.

An exception to this generalization is the anomalous "inverse dose-rate effect" observed for oncogenic transformation induced by neutrons. The incidence of transformation induced by a given dose of neutrons was enhanced if the dose was given continuously at low dose rate, or delivered in a series of dose fractions over 5 days compared with a single acute exposure (1). Miller et al (2) suggested that the enhancement of transformation by fractionation was neutron-energy dependent. Experiments with neutrons suffer from the disadvantage that the secondary charged particles produced when the neutrons are absorbed cover a wide range of LET values. Therefore, the present experiments were designed to investigate the LET dependence of the inverse dose-rate effect using charged particles of defined LET in the track segment mode.

Material and Methods

Exponentially growing C₃H 10T $\frac{1}{2}$ clone 8 cells were plated onto specially fabricated culture dishes (30mm diameter) at 2×10^5 cells per dish between 44 and 50 hrs prior to irradiation. The special dishes had thin mylar bottoms (6 μ m) to allow penetration of the charged particles. Immediately after the final irradiation, cells were trypsinized and replated into 100-mm-diameter tissue culture petri dishes at cell concentrations estimated to result in either 300 clonogenic cells per dish for the transformation assay or 30 clonogenic cells for the cell-survival assay. Cells were grown in Eagle's basal medium with 10% heat-inactivated calf serum supplemented with iron (CBS-iron) from Hyclone Labs. Inc., and 25 μ g/ml gentamycin. CBS-iron was found to uniformly support C3H10T1/2 cell growth; plating efficiencies varied from 15 to 48% while all radiation-induced transformation phenotypes displayed patterns of occurrences similar to that observed when using FBS. At the end of the incubation period, cells were fixed in formalin and stained with Giemsa. Cell survival was determined by the colony assay method while transformed foci of types II and III were identified according to criteria described by Reznikoff et al. (3, 4).

Irradiations with charged particles were performed using the track segment facility. The track segment irradiation fixture consisted of an enclosed wheel that allowed irradiations to occur

under physiological conditions (37°C temperature with humidified 95% air and 5% CO₂). Monolayer cell cultures growing in thin-bottomed (6-μm mylar) culture dishes were irradiated from below. Cells were exposed to 25-, 40-, 75-, 90-, 120-, 150- and 200- keV/μm charged particles. Total doses were 0.6 Gy for 25 keV/μm exposures, 0.3 Gy for 40 keV/μm and 0.2 Gy for the remainder of the high LET particles and were selected to elicit similar levels of cell killing (54-85% survival). For each radiation quality, the doses were delivered in three equal fractions at intervals of 0.3, 1.5, 15, 45, and 150 min.

Results and Discussion

Cells were irradiated with seven different LETs, each delivered in 3 dose fractions, with various time intervals between the fractions. The doses used were 0.6Gy for 25 keV/μm, 0.3Gy for 40keV/μm deuterons and 0.2Gy for 75, 90, 120, 150 and 200 keV/μm and were selected to be equitoxic at 70% survival. As can be seen in Fig. 1, enhancement of transformation incidence is seen only for LET values between 40 and 120 keV/μm, and the maximum enhancement of transformation incidence is evident when the interfraction interval is about 150 minutes.

The data of Figure 1 are summarized in Figure 2 in which the maximum change in transformation incidence caused by fractionation is plotted as a function of LET. Plotted in this way, the data clearly demonstrate that the inverse dose-rate effect i.e., the enhancement of transformation by fractionation, occurs only over a limited range of LET values. No enhancement is seen for radiations of low or very high LET.

Figure 2 Change in transformation frequency for C₃H 10T $\frac{1}{2}$ cells irradiated for the minimum (0.3 min.) and maximum (150 min.) fractionation intervals as a function of LET. Error bars represent ± 1 SD.

The experiments reported in this paper, using charged particles of defined LET, demonstrate convincingly that the "inverse dose-rate effect," i.e., the enhancement of transformation incidence when a dose of radiation is protracted by fractionation is LET dependent. Only LETs in the range between 40 and 120 keV/μm produce an enhanced effect due to fractionation; radiation of either lower or higher LET does not. It is further supported by the results of Bettega et al. (5) who showed an enhanced effect of fractionation by 4.3 MeV α particles (LET 101=keV/μm), and also by the results of Hieber et al. (6) who

demonstrated enhancement by fractionation with α particles having an LET of 100 keV/μm but not with α particles having an LET of 147 keV/μm (7).

References

1. C.K. Hill, A.Han, and M.M. Elkind, Fission-spectrum neutrons at low dose rate enhance neoplastic transformation in the linear, low dose region (0-10 cGy). *Int. J. Radiat. Biol.* **46**, 11-14 (1984).
2. R.C. Miller, D.J. Brenner, C.R. Geard, K. Komatsu, S.A. Marino, and E.J. Hall, Oncogenic transformation by fractionated doses of neutrons. *Radiat. Res.* **114**, 589-598 (1988).
3. C.A. Reznikoff, D.W. Brankow, and C. Heidelberger, Establishment and characterization of a cloned line of C3H mouse embryo cells sensitive to postconfluence inhibition of division. *Cancer Res.* **33**, 3231-3238 (1973).
4. C.A. Reznikoff, J.S. Bertram, D.W. Brankow, and C. Heidelberger, Quantitative and qualitative studies of chemical transformation of cloned C3H mouse embryo cells sensitive to postconfluence of cell division. *Cancer Res.* **33**, 3239-3249 (1973).
5. D. Bettega, P. Calzolari, G. Noris Chiorda, and L. Tallone-Lombardi. Transformation of C3H 10T $\frac{1}{2}$ cells with 4.3 MeV α particles at low doses: single and fractionated dose effects. *Radiat. Res.* **131**, 66-71 (1992).
6. L. Hieber and A.M. Kellerer, Transformation of C3H 10T $\frac{1}{2}$ cells by acute and fractionated doses of 2.7 MeV α particles. *Radiation Research: A Twentieth-Century Perspective*, Chapman, J.D., Dewey, W.C. and Whitmore, G.F., eds. Academic Press, New York Vol. 1, 346. (abstract) (1991).
7. L. Hieber, G. Ponsel, H. Roos, S. Fenn, E. Fromke and A.M. Kellerer, Absence of a dose-rate effect. *Int. J. Radiat. Biol.* **52** 859-869, (1987).

SISTER CHROMATID EXCHANGE INDUCTION AS A FUNCTION OF CHARGED PARTICLE LINEAR ENERGY TRANSFER

Charles R. Geard

In this study cycling Chinese hamster ovary cells were exposed to well characterized charged particles of defined linear energy transfer (LET) in order to establish the LET dependent inducibility of S.C.E.'s as a biological endpoint with a capacity for evaluating potential mutagen-carcinogens. The LET dependence for a number of biological endpoints as expressed by R.B.E.'s greater than 1 has been instructive in determining appropriate quality factors for diverse radiations. In contrast to the majority of biological endpoints S.C.E. induction was determined to decline and not increase on a dose dependent basis as charged particle LET changed from 10 to 120 keV per micrometer.

Chinese hamster ovary cells (CHO-AA8, generously provided by Dr. Larry Thompson LLNL) were grown on 6 μm thick mylar foil attached to stainless steel rings. Cell dishes containing firmly attached late log phase cells were placed in a wheel such that at the time of irradiation each dish was passed over a wedge shaped slit at a rate dependent on the dose to be delivered. Cells were irradiated at the Radiological Research Accelerator Facility (RARAF) based in Irvington, NY where a 4 MeV van de Graaff accelerator was used to irradiate cells with protons depositing energy at 10 keV/ μm (intracellular), deuterons at 40 keV/ μm , helium-3 particles at 80 keV/ μm and helium-4 particles at 120 keV/ μm . Absorbed doses ranged from 0.125 to 4 Gy at dose rates ranging from 4 to 14 Gy/min. Irradiating 20 dishes per experiment in a wheel typically took 10 minutes or less. Following irradiation within 2-3 mins. BrdUrd [5 μM] was added to the cells which were returned to the incubator and maintained under dim lighting conditions until the conclusion of each experiment (to restrict photosensitization). At 12 hrs post-irradiation colcemid (1 μM) was added to each cell dish and cells harvested at 18 hrs (6 hrs accumulation of mitotic cells). Chromosomal spreads were prepared and cells stained with 3% Giemsa in 0.3 M Na_2HPO_4 at pH 10.3. A 10-20 min. staining period using this "reverse" Giemsa technique results in chromosomes at the 2nd mitosis post incorporation of BrdUrd being differentially stained such that interchanges between chromatids (S.C.E.'s) are readily observed. From each dose in each experiment with a total of 75 dose-LET points, 500 or 1000 chromosomes i.e. ~25 or ~50 cells, were evaluated for the frequencies of S.C.E.'s using a 60x planapochromatic oil immersion objective.

Sister Chromatid Exchanges and Absorbed Dose

The design of this study whereby BrdUrd was nearly always added to cells immediately post-radiation precludes directly induced interchanges between sister chromatids and allows for evaluation of those S.C.E.'s produced as a consequence of the processing of DNA damage. The frequencies of S.C.E.'s per chromosome for the 40 and 120 keV/μm particles are shown in Tables 1 and 2 respectively and all data for induced S.C.E.'s per chromosome versus absorbed dose are shown in Figure 1. The level of S.C.E.'s in unirradiated cells which were similarly handled was 0.26 and 0.24 per chromosome for the 10 keV/μm experiments and 0.21 and 0.28 per chromosome in the 80 keV/μm experiments. These levels are much closer to

those shown in Table 1 than those in Table 2, which series of experiments was carried out over a four month period distinct from all others. The aspects of cell handling, media and growth control which could contribute to this difference were not determinable and over all 15 experiments control S.C.E. levels varied by a factor of about 2.5. The lowest control incidence was for the one experiment (with 40 keV/μm deuterons) when BrdUrd was present and partially incorporated into DNA at the time of irradiation. Since this experiment showed no evidence of a radiosensitized divergence in S.C.E. incidences results were included with the other five experiments. Results are shown in detail in Tables 1 and 2 to illustrate variation between experiments but particularly to examine the effects of low doses of alpha particles where there is interest as to their effectiveness as put forward in other studies and because of the possible role of radon daughter alpha particles in initiating human health problems. The within and between experiment variability was greater with the alpha particles than with the other charged particles with one experiment having no dose point with an S.C.E. incidence above the control level. This inefficiency in S.C.E. induction could possibly be attributed to the high basal levels, however absolute S.C.E. levels equal or greater were often observed in other experiments with a lower basal level. When all data for induced S.C.E.'s per chromosome are considered (Figure 1) protons at 10 keV/μm are the most effective at inducing S.C.E.'s with an apparent decline in effectiveness as LET increases. That is, this response is the opposite to that which has been observed for chromosomal aberrations and other endpoints where biological effectiveness increases from 10 to about 100 keV/μm. At both 0.5 (or 0.6) and 1.0 (or 1.2) Gy the order of effectiveness for S.C.E. induction is 10 > 40 > 80 > 120 keV/μm, with differences in frequencies being two fold or greater. The nature of the dose responses is unclear however initial linear responses followed by plateauing (sub-linear) responses at higher doses for each LET are probable. At the lower doses of the 80 and 120 keV/μm alpha particles (0.125 - 0.3 Gy) differences only represent near 10% increases over control and even the most effective dose (4 Gy of 10 keV/μm protons) only produces a 170% increase over control indicating that ionizing radiations over a range of LET's are

inefficient producers of S.C.E.'s relative to many chemicals which produce orders of magnitude effects. This system convincingly demonstrates a response to low LET (10 keV/ μ m) radiation and a lesser response to high LET alpha particles (80 or 120 keV/ μ m). The results obtained here with 120 keV/ μ m alpha particles were rather variable (Table 2) with the low dose responses in general being insignificantly different from control, however at these dose levels the number of particles per nucleus is low perhaps contributing to the variability.

Table 1
Sister chromatid exchanges per chromosome induced by 40 keV/ μ m deuterons^a

Absorbed dose [Gy]	<u>Experiment</u>						Mean
	1	2	3	4	5	6	
0	0.22 ^b	0.20	0.22	0.23	0.23	0.16	0.207
0.25	0.31 ^c	0.28 ^c	0.33 ^c	0.28 ^c	0.24	0.18	0.270 ^c
0.5	0.30 ^c	0.28 ^c	0.39 ^c	-----	0.35 ^c	0.25 ^c	0.314 ^c
1.0	0.36 ^c	0.37 ^c	0.44 ^c	-----	0.34 ^c	0.48 ^c	0.398 ^c
2.0	0.41 ^c	0.35 ^c	0.49 ^c	-----	0.38 ^c	-----	0.407 ^c

a. CHO-AA8 cells attached to 6 μ m thick Mylar in specially constructed cell dishes were irradiated in track segment mode. Bromodeoxyuridine (10^{-5} M) was added to cells 2-3 mins post-irradiation except in Expt. 6 when it was 30 mins. before irradiation and 2nd division mitotic cells were colcemid accumulated from 12-18 hrs. post-irradiation. Dashes represent inadvertent mishaps.

b. Mean S.C.E.'s per chromosome from samples of 500 (25 cells) or 1000 (50 cells) chromosomes; within experiment standard errors were derived from actual distributions (individual errors not shown). Distributions were tested against Poisson expectations (variance : mean ratios = 1) and found to vary inconsistently, with a trend to overdispersion throughout (not shown).

c. Significantly different from control at $P < 0.05$

Table 2

Sister chromatid exchanges per chromosome exchanges per chromosome induced by 120 keV/ μ m alpha particles^a

Absorbed dose [Gy]	<u>Experiment</u>					Mean
	1	2	3	4	5	
0	0.36 ^b	0.38	0.39	0.42	0.36	0.382
0.125	0.40	0.41	0.39	0.40	0.47 ^c	0.414
0.25	0.43 ^c	-----	0.41	0.36	0.43 ^c	0.408
0.5	0.44 ^c	0.50 ^c	0.46 ^c	0.38	0.50 ^c	0.456 ^c
1.0	0.44 ^c	0.46 ^c	0.51 ^c	0.39	0.45	0.450 ^c

a. footnote as for Table 1

b. footnote as for Table 1

c. Significantly different from control at $P < 0.05$.

HUMAN CHROMOSOME SPECIFIC CHANGES IN A HUMAN-HAMSTER HYBRID CELL LINE (A_L)

Charles R. Geard and Gloria Jenkins

Ionizing radiations are relatively efficient inducers of chromosomal changes and it is apparent that such changes play a pivotal role in assessing the consequences of radiation exposures. Chromosomal changes largely contribute to cell killing, to mutation and to oncogenic change. Chromosomal aberrations have been determined to be readily quantifiable with predictive value in cases of unknown radiation exposure and with clear dose and LET dependent effectiveness. However, the visualization of chromosomal changes by standard light microscopy only results in a fraction, albeit a significant fraction, of chromosome changes being recorded. In particular, interchanges between chromosomes, chromosomal translocations, which are essentially non-lethal and specific instances of which have been determined to be characteristic of particular human cancers are frequently missed. The complementary aberration to a translocation, a dicentric chromosome, is readily visualized, however translocations require analysis of chromosomal banding patterns or some means of recognizing interchanges between different chromosomes. This can be achieved by hybridizing chromosome specific DNA to mitotic spreads where the enquirer DNA is tagged directly or indirectly with a fluorochrome. In hybrid cell lines whole species DNA can be used to highlight a chromosome or chromosomes of interest. In this way using fluorescent *in situ* hybridization (FISH) all changes capable of fluorescence microscope magnification i.e. down to a fraction of a micrometer, can be visualized and quantified.

This approach has been used to quantify changes in human chromosome 11 in the human-hamster hybrid cell line A_L. The human chromosome 11 is the only human chromosome in this cell line, which has been used with some success to evaluate mutational changes at specific loci on the human chromosome.

Mitotic cells from the human-hamster hybrid cell line (A_L) which contains human chromosome 11 and which has been utilized in mutational studies were collected by mitotic dislodgement. Harvests routinely exceeded 95% mitoses. Cells were allowed to attach for 1.5 hrs. then irradiated with 0.5 to 10 Gy of Cs-137 γ -rays at a dose rate of 1.1 Gy/min. Following irradiation of the synchronized G1 cells, mitoses were collected 16-20 hrs later. Chromosomal spreads were prepared, denatured and hybridized with a fluorescein tagged probe against total human DNA. Counter staining was achieved with propidium iodide. Spreads were examined by fluorescence microscopy and mitoses from each dose point in 3 separate experiments examined for ploidy level, the number of human chromosomes per cell and all categories of aberrations involving the human chromosome. For the 3096 cells examined 81.3% were diploid cells, of which 88.5% contained 1 human chromosome 11, 5.8% contained 2 copies and 5.7% 0 copies. Of the remaining 18.7% of cells, 85.2% were tetraploid cells with 2 copies of human chromosome 11. There was a linear relationship between chromosome 11 changes and absorbed dose with a yield of 0.1 aberrations per Gy. The yield of dicentrics, translocations and terminal deletions which involve one lesion on the human chromosome was linear while the yield of interstitial deletions which arise from 2 interacting lesions on the human chromosome was curvilinear. The frequencies of dicentrics

and translocations was approximately equal, while there was a high (40-60%) incidence of incomplete exchanges between human and hamster chromosomes. These observations along with the high level of induced terminal deletions (failed interactions) indicate that interactions between lesions in hamster and human DNA lack the fidelity of human-human or hamster-hamster DNA lesion interactions. Assumptions made about the universality of response for this system should be treated with caution.

SSCP ANALYSIS OF *RAS* ONCOGENES IN NEUTRON TRANSFORMED MOUSE CELL LINES

Greg A. Freyer, Richard C. Miller and Yong Yu

While transformation induced by neutron radiation has been extensively studied, little is known about the molecular basis of this process. In fact, little is known about the molecular basis of transformation induced by any type of radiation. There are *in vivo* studies which reported that the *ras* oncogene was found to be activated in some radiation transformed mouse cell lines. In one of these studies, mice were irradiated with 0.44 Me V neutrons. Of 24 tumors studied, only 4 had mutations in *ras*, all in *Kras*. Other studies using x-rays or γ -rays found some cells contained activated *Ki-ras*. *In vitro* studies have not yielded the same results. Two studies with mouse and hamster cell lines transformed with x-rays concluded that none of several oncogenes tested were activated, including any of the *ras* oncogenes. These results leave several unanswered questions. Why this apparent difference between *in vivo* and *in vitro* studies? Possibly it is due to the fact that cell lines are already partially transformed in that they are immortalized. When *ras* is activated, is this a primary or secondary event? Predictably ionizing radiation, particularly neutrons, would not be expected to produce point mutations such as the activating event found with *ras*. What oncogenes were activated in the cell lines in which *ras* was not found?

The aim of this study was to make a careful survey of a large collection of cell lines transformed by neutrons to determine whether activated *ras* could be identified. Previous studies had relied on detecting *ras* by transfection and RFLP analysis or differential oligonucleotide hybridization. Our study employed the method of single-strand conformational polymorphism (SSCP) analysis which is a rapid method for detecting point mutations allowing one to detect any change in a particular sequence (outlined in Fig 1). SSCP analysis is carried out as follows: A set of oligonucleotides (20 bases) are synthesized which flank a 100 to 300 bp long sequence of interest. The oligonucleotides are used to amplify the sequence from DNA or cells (as few as 1000), in a polymerase chain reaction (PCR) containing α [^{32}P]dNTPs. The amplified sequence is denatured and analyzed by electrophoresis on a neutral sequencing

Figure 1. Scheme for SSCP analysis.

sized gel. Under these conditions, DNA migration is based on sequence composition. Thus, the two strands migrate as two distinct bands on the gel. A single base change in the sequence will cause a detectable shift in the normal migration of the two bands. The PCR products can then be sequenced directly to confirm the SSCP result.

There are two positions in *ras* where point mutations are found which activate the oncogenic phenotype. These correspond to codons 12 or 13 in exon 1 or 59 or 61 of exon 2. This is true for all three *ras* genes. Therefore, oligonucleotides bracketing these positions were synthesized for exon 1 and 2 of both *Kras* and *Nras* (see Table). The sequence for mouse *Hras* is not available.

TABLE OF SSCP OLIGONUCLEOTIDE PRIMERS

Gene	Name	Primer Sequence	Length of PCR Product (bp)
<i>Kras</i> exon 1	K1F	CTTGTGTGAGACATGTTCTA	171
	K1R	TATCGTAGGGTCGTAATCAT	
<i>Kras</i> exon 2	K2F	CTCCTACAGGAAACAAGTAG	128
	K2R	CACAAAGAAAGCCCTCCCCA	
<i>Nras</i> exon 1	N1F	GCTGGTGTGAAATGACTGAG	117
	N1R	TGGTGGGATCATATTCATCC	
<i>Nras</i> exon 2	N2F	GGTGAGACCTGCCTGCTGGA	103
	N2R	ATACACAGAGGAACCCTTCG	

C3H 10T1/2 cells were transformed by 6Mev neutrons using a single dose of 0.6 cGy. SSCP analysis was carried out on thirty five foci. Positive controls were used which contained mutations at codons 12/13 or 59/61 in *Nras* or *Kras*. The studies failed to detect any *Nras* or *Kras* mutations. These results would argue that neutrons do not lead to activation of *ras* oncogenes and future studies will need to focus on other oncogenes or a strategy which can screen for any activated oncogene. Examples of two SSCP gels are shown in Fig 2.

PUBLICATIONS

- Brenner, D.J. Radon - Current challenges in cellular radiobiology. *Int. J. Radiat. Biol.* 61:3-13, 1992
- Brenner, D.J. and Hall, E.J. Radiation-induced oncogenic transformation: the interplay between dose, dose protraction, and radiation quality. *Advan. Radiat. Biol.* 16: 167-179, 1992.
- Brenner, D.J. and Ward, J.F. Constraints on energy deposition and target size of multiply-damaged sites associated with DNA double-strand breaks. *International Journal of Radiation Biology* 61: 737-748, 1992.
- Brenner, D.J. Correlations between α/β and $T_{1/2}$: Implications for clinical biological modelling. *Brit. J. Radiol.* 65: 1051-1054, 1992.
- Brenner, D.J., Miller, R.C., Marino, S.A., Geard, C.R., Randers-Pehrson, G., and Hall, E.J. Inverse dose rate effects for neutrons: General features and biophysical consequences. *Radiat. Protec. Dosim.* 44: 45-48, 1992.
- Brenner, D.J., Medvedovsky, C., Huang, Y., and Worgul, B.V. Accelerated heavy particles and the lens VIII. Comparison between the effects of iron ions (190 keV/ μ m) and argon ions (88 keV/ μ m). *Radiation Research* 133: 198-203, 1993.
- Brenner, D.J. Dose, volume and tumor-control predictions in radiotherapy. *Int. J. Radiat. Oncol. Biol. Phys.* 26: 171-179, 1993.
- Brenner, D.J., Hall, E.J., Randers-Pehrson, G. and Miller, R.C. Model considerations on the dose-rate/LET dependence of oncogenic transformation by charged particles. *Radiation Research* 133: 365-369, 1993.
- Brenner, D.J., and Sachs, R.K., Generalized Microdosimetric Calculations of Cell-to-Cell Variance. *Radiat. Protec. Dosim.*, In Press (1993).
- Brenner, D.J. The Influence of Cell Killing on Radiation Oncogenesis: Possible Implications for High-LET Risk Assessment at Medium Doses. *Health Physics*, In Press (1993).
- Fung, A.Y.C. and Zaider, M. Energy-loss function for biological material: polycytosine. *Radiat. Protec. Dos.*, In Press (1993).
- Geard, C.R. Cytogenetic assays for genotoxic agents. *Lens and Eye Toxicity Research* 9 (3&4): 413-428, 1992
- Hahnfeldt, P., Hearst, J.E., Brenner, D.J., Sachs, R.K., and Hlatky, L.R. Polymer models for interphase chromosomes. *Proc. Nat. Acad. Sci.*, In Press (1993)
- Hall, E. J. Astor, M., and Brenner, D. J., Biological intercomparison of neutron beams used for radiotherapy generated by p^+ -Be in hospital-based cyclotrons. *British Journal of Radiology* 65: 66-71, 1992.
- Hall, E. J. and Brenner, D. J., The dose rate effect in interstitial brachytherapy - A controversy resolved. *British Journal of Radiology* 65: 242-247, 1992.

- Hall, E.J. and Brenner, D.J., Needles, Wires and Chips - Advances in brachytherapy. Clin. Oncol. 4: 249-256, 1992.
- Hall, E.J. and Brenner, D.J. The biological effectiveness of neutrons: Implications for radiation protection. Radiat. Protec. Dosim. 44: 1-9, 1992.
- Hall, E.J. and Brenner, D. J., The radiobiology of radiosurgery: Rationale for different treatment regimes for AVM's and malignancies. Int. J. Radiat. Oncol. Biol. Phys. 25: 381-385, 1993.
- Hall, E.J. and Brenner, D.J. The 1991 George Edelstyn memorial lecture; Needles, Wires and Chips - Advances in Brachytherapy. Clinical Oncology 4: 249-256, 1992.
- Hall, E. J. Review article: The gene as theme in the paradigm of cancer. The British Journal of Radiology 66: 1-11, 1993.
- Hall, E. J. and Freyer, G.A. The molecular biology of radiation carcinogenesis. In: Physical and Chemical Mechanisms in Molecular Radiation Biology, Plenum Press, 1991
- Hall, E.J., Geard, D. R. and Brenner, D.J. Risk of breast cancer in ataxia telangiectasia, Correspondence, New England Journal of Medicine 326: 1358-1359, 1992
- Kliauga, P., Nanodosimetry of heavy ions using a miniature cylindrical counter of wall-less design, Radiat. Prot. Dosim., In Press (1993)
- Kliauga, P. and Amols, H. Measurement of the spectrum and dose contribution of photoneutrons from a 20 MV clinical acclerator using a miniature proportional counter, Submitted, Phys. Med. Biol
- Medvedovsky, C., Worgul, B.V., Huang, Y., Brenner, D.J., Tao, F., Miller, J., Zeitlin, C., and Ainsworth, E.J. The Influence of dose, dose rate, and particle fragmentation on cataract induction by energetic iron ions. Advan. Space. Sci., In Press (1993).
- Miller, R. C., Geard, C. R., Geard, M.J. and Hall, E.J. Cell-cycle-dependent radiation-induced oncogenic transformation of C3H 10T-1/2 cells. Radiat. Res. 130: 129-133, 1992
- Rossi, H.H. and Zaider, M. Compound Dual Radiation Action. I. General Aspects. Radiat Res. 132: 178-183, 1992
- Sidik, K., Lieberman, H.B. and Freyer, G.A. Repair of DNA damaged by UV and ionizing radiation by cell-free extracts prepared from *schizosaccharomyces pombe*. Proc. Natl. Acad. Sci. 89: 12112-12116, 1992
- Tishler, R.B. and Geard, C.R. Correlation of sensitizer enhancement ratio with bromodeoxyuridine concentration and exposure time in human cervical carcinoma cells treated with low dose rate irradiation. Int. J. Radiation Oncology Biol. Phys. 22: 495-498, 1992
- Tishler, R.B., Geard C.R., Hall, E.J. and Schiff, P.B. Taxol sensitizes human astrocytoma cells to radiation. Cancer Res. 52: 3495-3497, 1992
- Tishler, R.B., Schiff, P.B., Geard, C.R. and Hall, E.J. Taxol: A novel radiation sensitizer. Int. J. Radiation Oncology Biol. Phys. 22: 613-617, 1992

Varma, M.N. and Zaider, M. A non-parametric, microdosimetric-based approach to the evaluation of the biological effects of low doses of ionizing radiation. In: Biophysical Modelling of Radiation Effects, IOP Publishing, Adams Hilger, p.145, 1992

Varma, M.N. and Zaider M. Hit size effectiveness in relation to the microdosimetric site size. Radiat. Prot. Dos., In Press (1993)

Vracko, M.G., Zaider, M. A calculation of the exciton energies in periodic systems with helical symmetry: Application to a hydrogen fluoride chain. Int. J. Quantum Chem. 43: 321-326, 1992

Vracko, M.G. and Zaider, M. A study of excited states in transpolyacetylene in the Hartree-Fock, Tamm-Dancoff and Random-Phase approximation. Int. J. Quantum. Chem. In Press, (1993)

Worgul, B.V., Brenner, D.J., Medvedovsky, C., Merriam, G.R. Jr., and Huang, Y. Accelerated heavy particles and the lens: VII The cataractogenic potential of 450 MeV/amu ^{56}Fe ions. Invest. Opthal. Vis. Sci. 34: 184-193, 1993.

Wuu, C.S. and Zaider, M. A mathematical description of sublethal damage repair and interaction during continuous low dose-rate irradiation. Radiat Prot. Dos., In Press (1993)

Zaider, M. and Rossi, H.H. On the question of RBE reversal at high doses. Radiat. Res. 130: 117-120, 1992

Zaider, M. and Minerbo, G.N. The response of cell populations to protracted doses of radiation: cell cycle progression. Radiat. Res. 133: 20-26, 1993

Zaider, M. The application of the principle of "dual radiation action" in biophysical modeling. In: Biophysical Modelling of Radiation Effects, IOP Publishing, Adam Hilger, p. 47, 1992

Zaider, M. A mathematical formalism describing the yield of radiation-induced single- and double-strand DNA breaks, and its dependency on radiation quality. Radiat Res. 134: 1-8, 1993

Zaider, M., Vracko, M.G., Fung, A.Y.C. and Fry, J.L. Electron transport in condensed water. Radiat. Prot. Dos., In Press (1993)

**DATE
FILMED**

7 / 29 / 93

END