WAVE DYNAMICS ON FALLING FILMS
AND ITS EFFECTS ON HEAT/MASS TRANSFER

Progress Report
for Period June 15, 1992 - June 14, 1993

Hsueh-Chia Chang

University of Notre Dame
Notre Dame, IN 46556

December 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Prepared for
The US DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-92ER14269

MASTER

Progress Report:

Wave evolution on a falling film is important in many industrial processes such as optical fiber coating, paper coating, pollutant scrubbers for industrial gas effluents etc. since it is responsible for both non-uniform coating and enhanced heat/mass transfer rates. It is also a physically intriguing spatio-temporal pattern formation process that has attracted much academic attention. If one follows the waves on a vertical plane downstream from its inception near the inlet, one can typically observe four wave regions. In the inception region, monochromatic waves of a very specific wavelength and speed (twice the interfacial velocity) are formed. These waves grow downstream and excite its overtone through a weakly nonlinear mechanism. This mechanism arrests the linear exponential growth of the waves. However, the front slopes of the waves continue to steepen until a highly asymmetric wave with a steep front and a gentle back slope appears in the third region. Such "solitary" waves retain their speed and shape for a long distance before transverse variation of their crests develops and begins to cause pinching and coalescence of adjacent crests. Such irregular and non-stationary behavior constitutes the fourth and final wave region which persists indefinitely downstream. The waves in the inception region have been well-understood from linear stability theory (see review by S. P. Lin, "Film waves on fluid interfaces", Editor R. E. Meyer, pg 261, Academic Press, 1983) developed in the twenty years after the classical experiments of Kapitza and Kapitza in 1949. Some weakly nonlinear theories for the second regime have also been developed by Lin (JFM, 63, 417 (1974)) and by us more recently (H.-C. Chang, Phys, Fluids, A1, 1314 (1989)) to explain the excitation of the linearly stable overtone. However, the more important third and fourth wave regions on a vertical plane remained unexplored until the present project. If the film flows down a cylindrical fiber, as in fiber coating processes, the wave transitions are distinctly different from those on a vertical plane. As Quere's recent experiment (Europhys. Lett, 13, 721 (1990)) indicates, saturated waves are not observed here unless the film thickness h is below a critical value h_c . If h is in excess of h_c , the waves grow

very rapidly to form drops with characteristic dimension close to the capillary length scale $H = (\sigma/\rho g)^{1/2}$. In the first year of this DOE project, we have resolved some important open problems in these two wave formation processes as detailed below:

1) Wave evolution on vertical plane

From our 1989 study, it was found that one expects two families of solitary waves in the third wave region, one faster than twice the interfacial velocity and one slower. However, waves in these regimes have relatively large amplitudes that are beyond the description of the classical weakly nonlinear theory. Moreover, most of the data for this region are on thick films that escape the lubrication approximation, thin-film waves being difficult to measure. As a result, one has to study the full Navier-Stokes equation with non-negligible dispersion instead of some of its limiting approximations. In collaboration with Professor E. A. Demekhin of the Applied Mathematics department of the Krasnodar University in Russia, we have been able to overcome this major obstacle for the first time and we are able to quantitatively compare our constructed solitary waves to measured waves in region III (see Figure 1). We indeed obtained two families of waves whose small-amplitude members can be described by weakly nonlinear bifurcation theories. We also carried out a stability analysis that predicts that a slow wave will be selected first before a fast wave is approached. This is also consistent with experimental data. The interaction between the fast solitary waves leads eventually to the non-stationary fourth wave region. We have developed a coherent structure theory that describes well the inception of this final region and provides a promising theory for the fully-developed "interfacial turbulence" further downstream. Our results will appear in the following journals:

- (a) "Laminarizing effects of dispersions in an active-dissipative nonlinear medium", accepted by Physica D.
- (b) "Construction of stationary waves on a falling film", accepted by Comp. Fluid Dynamics.

- (c) "Nonlinear Evolution of waves on falling film", accepted by Journal of Fluid Mechanics.
- (d) "Interfacial waves on a falling film", tentative title for an invited review for Annual Review of Fluid Mechanics (1993).

All four articles acknowledge financial support by this grant. We are also in the final stage of building our experiment which will permit laser resolution of the wave profiles and wave dynamics for comparison to our existing and further results. Our theoretical direction will be to use coherent structure theory to describe non-stationary dynamics in the fourth regime with a low-dimensional dynamics system which is constructed from a weak-interaction theory of the solitary waves. We are also ready now to apply our knowledge of interfacial dynamics to study its effect on heat and mass transfer. A preliminary theoretical study is underway.

2. Drop formation during coating of vertical fibers

We have associated the drop formation phenomenon with a finite-time blow-up solution of the evolution equation describing the coating process. It is analytically shown that when the film thickness is small, strongly nonlinear coupling between mean-flow and axial curvature produces another self-similar solution, a large solitary wave solution. Although the solitary wave solution is unstable, it generates many stable periodic solutions via the Silnikov bifurcation mechanism. These new attractors reduce the domain of attraction of the blow-up solution to a negligible neighborhood of the blow-up solution. As a result, drops will form (the blow-up is attracting) only if the solitary wave solution does not exist. Using matched asymptotic analysis, we are able to determine the critical film thickness beyond which solitary wave solutions do not exist. The analytical result yields

$$h_c = 1.68 R^3 H^{-2}$$

where R is the fiber radius and H is the capillary length. This is favorably compared to the experimental data for drop formation (see Fig. 2). We have hence used dynamical

systems tools to obtain an analytical estimate of an industrially important process. The following paper

(a) "Drop formation during coating of vertical fibers"

has been submitted to the Journal of Fluid Mechanics. We are currently working with Professor Bei Hu of the Mathematics department at Notre Dame on mathematical estimates of the "blow up" and "wave breaking" (gradient becoming infinity) phenomena of general evolution equations and their relationship to the actual processes. We are also using our techniques to estimate apparent dynamic contact angles of large capillary drops on a falling film. These dynamic contact angles play an important role in interfacial dynamics. We intend to eventually extend our study to predict the onset of air entrainment, an industrially important hydrodynamic instability in paper coating. A collaboration with Professor Cyrus Aidun of the Institute of paper Science and Technology at Georgia Tech has begun. His group has begun to carry out some pertinent experimental studies.

THEORY

EXPERIMENT

$$\langle R \rangle = 18.8 \text{ (water)}$$

Figure 1

Figure 2

END

DATE
FILMED
6/11/93