Final Report on Grant DE-FG06-88ER13890 (expired 5/22/91)**D-Erythroascorbic acid: Its Preparations, Chemistry, and Metabolism (Fungi and Plants).**

**Frank A. Loewus, Institute of Biological Chemistry, Washington State University, Pullman,
Washington 99164-6340**

and

**Paul A. Seib, Department of Grain Science and Industry, Kansas State University,
Manhattan, Kansas 66506**

Introduction

Oxalic acid is widely distributed in nature where it commonly occurs as its insoluble mineral salts (1). Plants produce characteristic deposits of calcium oxalate which are usually imbedded in unique crystal-forming cells referred to as crystal idioblasts (2, 3, 4). The origin of oxalate in plants has received considerable attention and glycolate metabolism has been generally regarded as a prime precursor candidate (5) although studies on the metabolism of L-ascorbic acid single out that plant constituent as well (6, 7). In the case of L-ascorbic acid metabolism in geranium leaf, it was shown that cleavage of the carbon chain at the C2-C3 position led to oxalic acid and L-(+)-tartaric acid as final products (8, 9). Extension of these experiments to oxalate-accumulating plants that contain little or no tartaric acid revealed the presence of a comparable L-ascorbic acid metabolism with the exception that the cleavage products were oxalic acid and L-threonic acid or products of L-threonic acid metabolism (10, 11). It was the subsequent oxidation of L-threonic acid that led to L-tartaric acid accumulation in geranium leaf (**Figure 1**).

Mild alkaline oxidation of L-ascorbic acid to form oxalic acid and L-threonic acid was first used in studies on the structure of L-ascorbic acid (12). Later, Isbell and Frush reported that excess alkaline hydrogen peroxide afforded oxalic acid and L-threonic acid in high yield (13). A reasonable mechanism for cleavage of L-ascorbic acid at the endiolic bond is found in studies on the photooxygenation of L-ascorbic acid (14) (**Figure 2**). Presumably, analogs of L-ascorbic acid that differ only in the substituent at C4 also form a hydroperoxide in the presence of alkaline hydrogen peroxide and subsequently yield oxalic acid and the corresponding aldonic acid or its lactone.

We became interested in such a possibility when we discovered that L-ascorbic acid was rare or absent in certain yeasts and fungi whereas a L-ascorbic acid analog, D-*glycero*-pent-2-enono-1,4-lactone (trivial nomenclature: D-erythroascorbic acid), was present (15, 16, 17). It has long been known that oxalate occurs in yeasts and fungi (1) and its production plays a role in plant pathogenesis (18, 19, 20). As to the biosynthetic origin of fungal oxalic acid there is little information although it is generally assumed that oxaloacetate or possibly, glycolate, might be that precursor (21).

In anticipation of the possibility that D-*glycero*-pent-2-enono-1,4-lactone might function as an oxalic acid precursor, Dr. Paul Seib and his group at Kansas State University devised three practical chemical syntheses for D-erythroascorbic acid to supply us with material for peroxygenative degradation studies of oxalic acid production. The first was a four step procedure beginning with D-glucose (25). More recently, Gan and Seib (26) shortened the synthesis to three steps starting from commercially available 3,4-O-benzylidene-D-ribo-1,5-lactone. They have also devised a synthetic route from D-xylose (27). Details of their studies are found in a report prepared by Dr. Seib that is included in the Materials and Methods section.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

FEB 10 1993

12
MASTER

Our initial studies which were motivated by our search for a culturable, single-cell, L-ascorbic acid-producing organism amenable to studies on L-ascorbic acid biosynthesis had led us to test *Lypomyces starkeyi*, the yeast reportedly containing a significant amount of L-ascorbic acid (22, 23). Use of highly sensitive, specific, and selective HPLC analytics involving amperometric (electrochemical) detection (24) revealed that the purported L-ascorbic acid was, in fact, the 5-carbon analog noted above, D-glycero-pent-2-enono-1,4-lactone = trivially named D-erythroascorbic acid (16). Structures of L-ascorbic acid and D-erythroascorbic acid are compared in **Figure 3**. Subsequent studies during this project revealed that D-erythroascorbic acid was present in several species of edible fungi. It may well be that this analog is as common in fungi as L-ascorbic acid is in higher plants.

As Liang et al. reported (25), D-erythroascorbic acid is cleaved at the C2/C3 double bond to form oxalic acid and D-glyceric acid, in analogy to the peroxygenative pathway for L-ascorbic acid (14). We have confirmed this in a kinetic study with alkaline hydrogen peroxide that identified optimal conditions for stoichiometric oxalic acid production from L-ascorbic acid and D-erythroascorbic acid (**Figure 4**).

To study the biosynthesis and metabolism of D-erythroascorbic acid in fungi, we selected *Sclerotinia sclerotiorum* (Lib.) de Bary, ATCC 18687, an oxalate-producing fungus that is a serious pathogen on numerous crops, notably legumes, leafy vegetables, small grains, and oil seeds such as sunflower (28). D-Erythroascorbic acid was detected amperometrically at 0.7 volts following HPLC on a Bio-Rad organic acid column. Oxalic acid fails to produce a strong signal below 1.0 volt in this system so this product was measured separately by HPLC using a Waters IonPak A column and ion conductivity detection. Submerged culture with shaking, produced mycelia in which very little D-erythroascorbic acid accumulated (<25 µg/g dry weight of mycelia after 15 days on Difco Y-M broth). After many months of experiments involving submerged cultures, we devised a new approach involving microporous rafts floating on growth media. This procedure led to greatly enhanced recovery of D-erythroascorbic acid and permitted us to test selected carbohydrates as carbon sources. Murakawa et al. (29) reported that *Candida* species grown on D-arabinose, D-arabinonic acid or D-arabinono-1,4-lactone as compared to D-glucose contained higher levels of D-erythroascorbic acid. As seen in **Figure 5**, D-erythroascorbic acid production by *S. sclerotiorum* was greater when D-arabinose was the carbon source in a 4 day culture. It also revealed an unknown electrochemically detectable product $R_{AA} = 0.87$ (where R_{AA} = retention time of L-ascorbic acid) as compared to D-erythroascorbic acid ($R_{AA} = 1.14$) (**Figure 6**). The unknown had a uv spectrum that closely resembled D-erythroascorbic acid (which is virtually indistinguishable from L-ascorbic acid). Its structure is still unknown but we think it might possibly be an open-chain form of D-erythroascorbic acid. A comparison of raft-suspended cultures of *S. sclerotiorum* grown on D-glucose, D-arabinose, and/or D-arabinono-1,4-lactone for 4 days revealed that those cultures with D-arabinose had higher levels of D-erythroascorbic acid, unknown and oxalic acid than those provided with D-glucose or a combination of D-glucose and D-arabinose (**Figure 7**). There was no synergism when D-arabinose was coupled with D-arabinono-1,4-lactone. When D-arabinose was supplied singly, the 4 day culture produced nearly 2 mg of D-erythroascorbic acid and 600 mg of oxalic acid/g dry weight of mycelia. It should be emphasized here that the enantiomeric form of the pentose precursor used in these studies was D-arabinose whose configuration at C-4 is compatible with D-erythroascorbic acid. Whether other D-pentoses also function similarly needs to be examined.

Oxalic acid was mostly secreted and accumulated in the broth (Figure 8). Extraction of the hyphal mat from a 10 day culture with buffer (to obtain free oxalic acid) or acid (to recover calcium oxalate) led to only minor recoveries. Virtually none of the oxalate was present in sclerotia, the dense dark aggregates of hyphae that form in cultures held for 4 days or more.

The availability of D-[5-³H]arabinose permitted an experiment in which *S. sclerotiorum*, grown for 4 days on 0.1% D-[5-³H]arabinose. The mycelial extract was analyzed for ³H in D-erythroascorbic acid, the $R_{AA} = 0.87$ unknown, and oxalic acid by anion exchange chromatography (Figure 9). Three major peaks of ³H accounted for most of the extracted radioactivity. The first, a peak at fraction 21, was the electrochemically-reactive unknown; the second, a shoulder at fraction 27 was glyceric acid; and the third at fraction 42-45 was D-erythroascorbic acid. Oxalic acid was unlabeled as expected. Although the experiment needs to be repeated with appropriate ¹⁴C-labeled precursors that will tag the 2-carbon fragment, this preliminary study strongly supports the view that D-erythroascorbic acid is cleaved at the C2/C3 position to yield oxalic acid and D-glyceric acid.

The stereomeric assignment of D-erythroascorbic acid (which is identical to L-ascorbic acid apart from the absence of an asymmetric carbon on the sidechain) was strengthened by two experimental observations. First, ascorbic acid oxidase utilizes both L-ascorbic acid and D-erythroascorbic acid, albeit the latter at a rate about one-third the former (Figure 10). Second, D-erythroascorbic acid can replace L-ascorbic acid during growth of Vitamin C-requiring *Manduca sexta* larvae although it fails to support ecdysis to pupae and adults (27).

Host-pathogen studies on *Sclerotinia*, *Sclerotium*, and related genera extend backward many decades and the observation that oxalic acid formation and polygalacturonase production have key roles during the infective process is well documented (30, 31, 32 and references therein). Advances in cellular and molecular biology have opened new approaches. These include consideration of signals associated with development of infectious structures (33), cloning of *S. sclerotiorum* genes encoding cell wall degrading enzymes (34), cytoplasmic processes in *Aspergillus* associated with oxalate biosynthesis (35) and selective inhibition of plant enzymes by oxalate (36). Our research on oxalic acid biosynthesis in *S. sclerotiorum* will contribute to a better understanding of these related studies, thus further advancing our understanding of the infective process.

Although a major portion of our study dealt with biosynthesis and metabolism of erythroascorbic acid in *S. sclerotiorum* due to the experimental practicality of this fungus, to its agronomic importance and to its close relationship to oxalic acid biosynthesis, the significance of the project in terms of renewable resources should also be mentioned. We now know that D-erythroascorbic acid is a relatively common constituent of yeasts and fungi. Use of these organisms in conversion of carbohydrate biomass to combustible fuels involves production of D-erythroascorbic acid. It would be useful to know if this compound is an essential component microorganismal growth. It would also be useful to know if this ascorbyl analog has practical uses in medicine, nutrition, industrial applications and the like. Since the organisms we have used can produce up to 2% of dry weight as this compound, its extraction and recovery from spent cells following the fermentation process may have economic value.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Materials and Methods

Chemicals

Reagent grade chemicals were used. Ascorbic acid was recrystallized from water (1g/mL) with a 10-fold excess of glacial acetic acid. D-Erythroascorbic acid was prepared by a 4-step synthesis that involved oxidation of D-glucose to K D-arabinonate, lactonization, conversion to the 2-keto methyl ester, and tautermization. The final crystalline product had a melting point of 160-162°C and $[\alpha]_D^{25} = -11 \pm 1$ ($c = 5$, H₂O) (25). Stock solutions of ascorbic acid and erythroascorbic acid (2mg/mL) in 5 mM dithiothreitol remained stable and storable at -20°C for extended periods. Working standards (200 ng/mL) were prepared daily. A stock solution of oxalic acid (20 mg/mL) was prepared from freshly sublimed crystals and stored at -20°C. A working standard (20-200 µg/mL) was prepared daily.

Synthesis of D-erythroascorbic acid from D-glucose or D-xylose

In the course of this project, two schemes which were more practical than the one first used, were devised to chemically synthesize D-erythroascorbic acid starting from D-glucose. In the two schemes, either C-1 or C-6 of D-glucose was lost, so that a variety of isotopically-labeled derivatives of D-erythroascorbic acid can be synthesized from labeled D-glucose. In scheme II, labeled D-xylose may also be used as starting material.

Scheme I began with O₂-oxidation of D-glucose to give calcium D-arabinonate in 74% yield. Epimerization of the latter in hot aqueous calcium hydroxide under pressure gave a 3:7 mixture of D-ribonate and D-arabinonate in 22% yield. Unreacted D-arabinonate, due to its low solubility in water, was readily recovered from the reaction mixture in 50-60% yield, and recycled in the scheme. Three cycles of the epimerization reaction, starting with 105 g of calcium D-arabinonate, gave crude D-ribonate in 80% yield. Following acidification of the technical-grade product, high-performance liquid chromatography showed a 4:1 syrupy mixture of D-ribono- and D-arabinono lactones. Reaction of this mixture with benzaldehyde and concentrated hydrochloric acid gave crystalline 3,4-*O*-benzylidene-D-ribono-1,5-lactone in 59% yields. The acetal of this lactone was oxidized with magnesium oxide in acetone to its 2-keto derivative in 76% yield. Acid-catalyzed methanolysis gave a syrupy mixture of products, which upon tautomerization in hot methanolic sodium acetate followed by removal of sodium ions gave 78% D-erythroascorbic acid. This five-step synthesis yielded 20% of D-erythroascorbic acid from D-glucose (26).

Scheme II started with D-xylose, which can be prepared in 60% yield from D-glucose using procedures that are described in the literature. D-Xylose was then isomerized to a mixture of D-xylulose:D-xylose (8:2, w/w) using xylose isomerase in the presence of borate. After removal of borate, the syrupy mixture of sugars was acetonated to give mostly 2,3-*O*-isopropylidene-D-xylulose plus some 1,2:3,5-di-*O*-isopropylidene-D-xylose. The mono-isopropylidene, which was isolated by solvent extraction in 70% crystalline yield, was catalytically oxidized in aqueous solution at pH 8 to give 2,3-*O*-isopropylidene-β-D-*threo*-pentulosonic acid in 85% yield. Finally, 2,3-*O*-isopropylidene-β-D-*threo*-pentulosonic acid was converted to D-erythroascorbic acid in methanolic hydrogen chloride in 64% yield. The overall isolated yield from D-xylose was 31%, and from D-glucose was 18% (27).

D-Erythroascorbic acid was slightly less readily oxidized at pH 4-8 than L-ascorbic acid by electrochemical oxidation or by O₂ in water.

Sources of organisms used in study

Whetzelinia sclerotiorum (previously named *Sclerotinia sclerotiorum*), ATCC #18687, *Sclerotium cepivorum* strain Sherren ATCC #56041, *Lipomyces starkeyi* ATCC #12659, and *Penicillium griseorosium* (previously named *Penicillium cyaneo-fulvum*) ATCC #10431 were obtained from the American Type Culture Collection, Rockville, MD. Dikaryotic stocks of *Lentinus*, *Pleurotus* and *Flammulina* species (37) were generously provided by Professor Philip G. Miles, State University of New York at Buffalo.

Cultural conditions

L. starkeyi was maintained on 3% Sabouraud's glucose broth and 2% agar, pH 6.9, as recommended by ATCC recipe #28. Cell suspensions were prepared by transfer from solid to liquid media and growth for 7-9 days in dark at 25°C. Fungal cultures were maintained on Bacto YM agar (Difco #0712-01-8) and transferred to Bacto YM broth (Difco #0711-01) for growth of experimental tissue. Mycelial mats were grown on agar in petri dishes for 1 week. Inoculation of broth for submerged cultures or floating rafts of aerobic growth was done with 7 mm disks cut from the mats with a sterile cork borer. Submerged cultures were grown at 25°C in 250-mL erlynmeyer or 1-L low form flasks on an orbital or reciprocating shaker at 130 r/min or 70 strokes/min, respectively. Raft cultures were grown on microporous polypropylene membranes supported by plastic frames (#M3027, Sigma Chem Co, St Louis MO) in Magenta GA-7-3 plastic vessels (#V8380, Sigma Chem Co, St Louis, MO). Rafts were floated on 100 mL of media except in experiments involving radioactive substrates in which case the plastic legs of the rafts were removed and the rafts floated on 25 mL of media.

Harvest and extraction

Submerged cultures were recovered by draining the mycelia on 20 µm mesh nylon bolting cloth followed by vacuum filtration on a glass filter (Whatman GF/C) until a damp mat was formed. This mat was thoroughly wetted for 2-3 min with 2 mM dithiothreitol and again drawn down until all filtrable liquid was extracted. The mycelia was weighed and sliced into small cubes. An aliquot was taken for dry weight determination and the remaining material stored at -70°C.

Frozen samples of mycelia were pulverized in liquid nitrogen and taken up in 2 mM dithiothreitol (approx. 3 mL/g). After extraction for 30 min at 25°C, samples were centrifuged at 10,000g for 10 min. Supernatant (1 mL) was applied to a column of 2 mL of Dowex 50 H⁺, 200-400 mesh resin and followed with sufficient distilled water to recover 5 mL of eluate. The diluted eluate was filter-centrifuged (Bio-Analytical Co., West Lafayette IN) through a 0.45 µm membrane and analyzed for erythroascorbic acid and/or oxalic acid by HPLC.

Assay of ascorbic acid and erythroascorbic acid

Ascorbic acid and erythroascorbic acid were determined by HPLC with electrochemical detection. In most instances the column of choice was an organic acid column, HPX-87, with an ion exclusion pre-cartridge (Bio-Rad Labs, Richmond, CA) that provided baseline resolution of ascorbic acid and erythroascorbic acid peaks (16, 24). Assays were automated through the use of a Spectro-Physics Model 8875 autosampler and Model 4270 computing integrator with data acquisition module. When detection of oxalic acid was included in the ascorbic acid or erythroascorbic acid assays, a polymeric reverse phase column, 25 cm x 4.6 mm PLRP-S

(Polymer Labs Inc, Amherst, MA), was employed (38; 39). The mobile phase of choice when using the polymeric column was 1.8% tetrahydrofuran and 0.3% metaphosphoric acid in water as recommended (39).

Assay for oxalic acid

Oxalic acid was readily assayed by HPLC in conjunction with conductivity detection (41). A 0.46 x 5 cm IC Pak Anion column with precolumn (Waters, Milford MA) was used in conjunction with a M430 ion conductivity detector (Waters, Milford MA) to measure conductivity changes in column eluates. Preliminary studies led to the decision to operate the column isocratically with "B3" eluent (12.65 mM boric acid, 1.70 mM gluconic acid, 4.72 mM lithium hydroxide, 0.75 mM glycerin and 12% acetonitrile). Choice of this system provided rapid elution of oxalic acid (approx. 6 min at a pump flow rate of 1.2 mL/min) and minimum interference from inorganic anions that might be present in the extract.

References

1. Hodgkinson A 1977 Oxalic Acid in Biology and Medicine. Academic Press, London 325 p.
2. Libert B, Franceschi VR 1987 Oxalate in crop plants. J Agric Food Chem 35:926-938.
3. Franceschi VR 1989 Calcium oxalate formation is a rapid and reversible process in *Lemna minor* L. Protoplasma 148:130-137.
4. Franceschi VR 1987 Oxalic acid metabolism and calcium oxalate formation in *Lemna minor* L. Plant Cell Environ 10:397-406.
5. Raven JA, Griffiths H, Glidewell SM, Preston T 1982 The mechanism of oxalate biosynthesis in higher plants: investigations with the stable isotopes ^{18}O and ^{13}C . Proc Royal Soc London B216:87-101.
6. Loewus FA 1988 Ascorbic acid and its metabolic products. In The Biochemistry of Plants, Volume 14, Carbohydrates, J Preiss, ed, Academic Press, San Diego, CA pp 85-107.
7. Li X, Franceschi VR 1990 Distribution of peroxisomes and glycolate metabolism in relation to calcium oxalate formation in *Lemna minor* L. Eur J Cell Biol 51:9-16.
8. Wagner G, Loewus FA 1973 The biosynthesis of (+)-tartaric acid in *Pelargonium crispum*. Plant Physiol 52:651-654.
9. Williams M, Saito K, Loewus FA 1979 Ascorbic acid metabolism in geranium and grape. Phytochemistry 18:953-956.
10. Nuss RF, Loewus FA 1978 Further studies on oxalic acid biosynthesis in oxalate-accumulating plants. Plant Physiol 61:590-592.
11. Helsper JP, Loewus FA 1982 Metabolism of L-threonic acid in *Rumex x acutus* L. and *Pelargonium crispum* (L.) L'Hér. Plant Physiol 69:1365-1368.
12. Herbert RW, Hirst EL, Percival GV, Reynolds RJW, Smith F 1933 The constitution of ascorbic acid. J Chem Soc 1270-1289.
13. Isbell HS, Frush HL 1979 Oxidation of L-ascorbic acid by hydrogen peroxide: preparation of L-threonic acid. Carbohydr Res 72:301-304.
14. Kwon B-M, Foote CS, Khan SI 1989 Photooxygenation of ascorbic acid derivatives and model compounds. J Am Chem Soc 111:1854-1860.

15. Leung CT, Loewus FA 1985 Concerning the presence and formation of ascorbic acid in yeasts. *Plant Sci* 38:65-69.
16. Nick JA, Leung CT, Loewus FA 1986 Isolation and identification of erythroascorbic acid in *Saccharomyces cerevisiae* and *Lipomyces starkeyi*. *Plant Sci* 46:181-187.
17. Dumbrava VA, Pall ML 1987 Control of nucleotide and erythrosacorbic acid pools by cyclic AMP in *Neurospora crassa*. *Biochim Biophys Acta* 926:331-338.
18. Bateman DF, Beer SV 1965 Simultaneous production and synergistic action of oxalic acid and polygalacturonase during pathogenesis by *Sclerotium rolfsii*. *Phytopathology* 55:204-211.
19. Maxwell DP, Lumsden RD 1970 Oxalic acid production by *Sclerotinia sclerotiorum* in infected bean and in culture. *Phytopathology* 60:1395-1398.
20. Lumsden RD 1979 Histology and physiology of pathogenesis in plant diseases caused by *Sclerotinia* species. *Phytopathology* 69:890-896.
21. Maxwell DP 1973 Oxalate formation in *Whetzelinia sclerotiorum* by oxaloacetate acetylhydrolase. *Physiol Plant Pathol* 3:279-288.
22. Heick HMC, Atewart HB, Graff GLA, Humpers JEC 1969 Occurrence of ascorbic acid in the yeast *Lipomyces starkeyi*. *Can J Biochem* 47:752-753.
23. Heick HMC, Graff GLA, Humpers JEC 1972 The occurrence of ascorbic acid among the yeasts. *Can J Microbiol* 18:597-600.
24. Grün M, Loewus FA 1983 Determination of ascorbic acid in algae by high-performance liquid chromatography on strong cation-exchange resin with electrochemical detection. *Anal Biochem* 130:191-198.
25. Liang YTS, Liu XS, Seib PA 1990 Synthesis of D-erythroascorbic acid. *Carbohydr Chem* 9:75-84.
26. Gan L-X, Seib PA 1991 Synthesis of D-erythroascorbic acid from D-glucose.
27. Shao Y-Y, Seib PA, Kramer KJ, Van Galen DA 1993 Synthesis and properties of D-erythroascorbic acid and its Vitamin C activity in the Tobacco Hornworm, *Manduca sexta*. *J Agric Food Chem*, in press.
28. Willetts HJ, Wong JA-L 1980 The biology of *Sclerotinia sclerotiorum*, *S. trifoliorum*, and *S. minor* with emphasis on specific nomenclature. *Bot Rev* 46:101-165.
29. Murakawa S, Sano S, Yamashita H, Takahashi T 1977 Biosynthesis of D-erythroascorbic acid by *Candida*. *Agric Biol Chem* 41:1799-1800.
30. Willetts HJ, Wong JA-L 1980 The biology of *Sclerotinia sclerotiorum*, *S. trifoliorum*, and *S. minor* with emphasis on specific nomenclature. *Bot Rev* 46:101-165.
30. Lumsden RD 1979 Histology and physiology of pathogenesis in plant diseases caused by *Sclerotinia* species. *Phytopathology* 69:890-896.
31. Purdy LH 1979 *Sclerotinia sclerotiorum*: diseases and symptomatology, host range, geographic distribution, and impact. *Phytopathology* 69:875-880.
32. Smith VL, Punja ZK, Jenkins SF 1986 A histological study of infection of host tissue by *Sclerotium rolfsii*. *Phytopathology* 76:755-759.
33. Hoch HC, Staples RC 1991 Signaling for infection structure formation in fungi. In: *The Fungal Spore and Disease Initiation in Plants and Animals*, Cole GT, Hoch HC, eds Plenum Press pp 25-46.
34. Waksman G 1988 Molecular cloning of genes expressed specifically during induction of cell wall degrading enzymes from *Sclerotinia sclerotiorum* and preliminary identification

- of a fungal beta-galactosidase encoding gene by expression in *Escherichia coli*. *Curr Genet* 14:91-93.
35. Kubicek CP, Schreierl-Kunar G, Wohrer W, Rohr M 1988 Evidence for a cytoplasmic pathway of oxalate biosynthesis in *Aspergillus niger*. *Appl Environ Microbiol* 54:633-637.
 36. Kleczkowski LA, Randall DD, Edwards GE 1991 Oxalate as a potent and selective inhibitor of spinach (*Spinacia oleracea*) leaf NADPH-dependent hydroxypyruvate reductase. *Biochem J* 276:125-127.
 37. Chang S-T, Miles PG 1989 Edible mushrooms and their cultivation. CRC Press Inc, Boca Raton FL 337 p.
 38. Lloyd LL, McConville JA, Warner FP, Kennedy JF, White CA 1987 HPLC analysis of L-ascorbic acid (vitamin C) using a polymeric reversed-phase adsorbant. *LC-GC* 5:338-341.
 39. Bushway RJ, King JM, Perkins B, Krishnan M 1988 High-performance liquid chromatographic determination of ascorbic acid in fruits, vegetables and juices. *J Liq Chromatogr* 11:3415-3423.
 40. Jones WR, Jandik P, Heckenberg AL 1988 Gradient elution of anions in single column ion chromatography. *Anal Chem* 60:1977-1979.

Figure 1

Kwon, Foote & Khan, 1989

Figure 2

Figure 3

Peroxygenation of L-AA and D-EAA Oxalate Production

Figure 4

Products from Glc and/or Ara *Sclerotinia sclerotiorum*, 4 days growth

Expt #1

Figure 5

Figure 6

Products from Selected Carbon Sources

Sclerotinia sclerotiorum, 4 days growth

Figure 7

Oxalate Production from Glc and/or Ara

Sclerotinia sclerotiorum, 10 days growth

Figure 8

Extract from D-[5-³H]Ara-labeled S.s. Anion-exchange, 0 - 0.1 N Formic Acid

Figure 9

Oxidation of EAA or AA with AA Oxidase

Figure 10

END

DATE
FILMED

4 / 7 / 93