THE SINGLE ELECTRON CHEMISTRY OF COALS

Report for the period
July 1 1990 to December 30 1990

John W. Larsen
and Parveen Kaushal

Department of Chemistry
Lehigh University
Bethlehem, PA 18015

January 18, 1991

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE UNITED STATES DEPARTMENT OF ENERGY

MASTER

OBJECTIVES AND SCOPE OF WORK

Objectives

To investigate and characterize the single electron reactions of alkyl and alkoxy aromatic compounds in order to determine the role these reactions play in the chemistry of coal.

Scope of work

1. Attempts will be made to demonstrate that the radicals from inertinite maceral group will initiate the polymerization of 4-vinylpyridine.
2. The molecule, N,N-diphenyl-phenylenediamine, will be deposited in coals to characterize their native free radicals.
3. Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) will be used to characterize the numbers and strengths of single electron donors in coals.

INTRODUCTION

Our work on single electron transfer in coals led us to the knowledge that the energetics of bond cleavage in radical cations is 20-40 kcal/mole lower than the corresponding homolytic bond cleavage energies.¹⁻³ Having made excellent progress in the other areas covered by this proposal, we are extending our studies to the investigation of the formation and cleavage reaction of radical cations in coals.

The formation of a radical cation requires the transfer of an electron from a neutral molecule to an appropriate electron acceptor (oxidant). As a first step, we seek oxidants which will form radical cations from functional groups typical of those in coals. We must also study the decomposition behavior of bonds typical of those found in coals. Alkyl and alkoxy aromatic compounds were chosen as the electron donors because of their common occurrence in coals.⁴

RESULTS AND DISCUSSION

When benzyl phenyl ether (BPE) was refluxed in acetonitrile (82°C) in the presence of tris-(1,10-phenanthroline)-iron(III) perchlorate (for details of procedure see experimental section) the products identified were phenol (P), benzyl acetamide (BAC) and 2-(phenylmethyl)phenol, the rearranged product (RP), shown in equation (1). Benzyl phenyl ether disappeared in the first few

minutes of refluxing (Table 1, for product yield see Table 2) while only 55% reaction took place in two hrs. at room temperature. The reaction products are consistent with the decomposition of the benzyl phenyl ether radical cation. Presumably iron(III)phenanthroline oxidized BPE via an outer-sphere electron transfer (ET) s^{-1} to give the radical cation [equation(2)] which undergoes C-O bond cleavage [equation (3)]. The radical and cation produced from it react to give the products shown [equations (4)-(6)].

In order to study whether the rearranged product (Eq. 6) is formed by an inter or intra molecular reaction [eq. (7)-(10)], an equimolar mixture of BPE (0.025g, 1.36×10^{-4} mol, 0.03 mol dm^{-3}) and p-bromobenzyl-p-bromophenylether (BPE Br) (0.047g, 1.36×10^{-4} mol, 0.03 mol dm^{-3}) was refluxed in acetonitrile in the presence of tris-(1,10-phenanthroline) iron(III) perchlorate. Almost all of the BPE reacted in the first 0.5 hr. of the reflux time while only 49% of BPE Br reacted. The products were phenol (48 mole %), benzylacetamide (37 mole %), the rearranged product (RP) (15 mole %) and two more having retention times of 5.4 min. and 13.4 min. These last two are products of the direct reaction between bromosubstituted BPE and the iron(III) complex. The bromo-substituted compound may not have been oxidized to form a radical cation.

If Intra molecular

If Inter molecular

When BPE was refluxed (0.05 mol dm^{-3}) in acetonitrile for 20 hrs., in the absence of iron(III) complex, no reaction was observed.

Other iron(III) oxidizing reagents were studied to see if they can oxidize BPE efficiently. Iron powder was used (0.1 mol dm^{-3}) with a slow stream of air in refluxing acetonitrile for 20 hrs. No oxidation was observed. When iron (III) chloride (0.08 mol dm^{-3}) was refluxed with BPE (0.05 mol dm^{-3}) in acetonitrile, $45 \pm 5\%$ of the BPE reacted in 1 hr. and $55 \pm 5\%$ in 18.5 hrs. See Table 2 for the products and mass balance.

Other single electron transfer oxidizing reagents were also tried with BPE (see Table 3). Cerium(IV) ammonium nitrate is considered to be a typical single electron outer sphere oxidant even though it does have some inner-sphere oxidizing properties. It reacted with BPE, yielding PhOH, ArCH₂ONO₂ and an unidentified

product having the general structure ArCH_2X where X contains N and/or O. Other single electron oxidants listed in Table 3 did not react with BPE.

Ideally, we would like to use O_2 as the oxidant and run the reaction using a catalytic amount of a metal based single electron oxidant. There is a possibility that iron(II)(1,10-phenanthroline) perchlorate could be oxidized by O_2 to the iron(III) complex. To test this idea, iron(III)(1,10-phenanthroline) perchlorate ($0.025 \text{ mol dm}^{-3}$) was used in a catalytic amount and refluxed with BPE ($0.137 \text{ mol dm}^{-3}$, 6-fold excess) in acetonitrile with a stream of air passing through the solution.

Diagram to show the recycling of iron complex.

$84 \pm 10\%$ conversion of BPE took place in 23 hrs. while almost all

(96 $\pm$ 10%) of the substrate reacted in 47 hrs. The products were P, BAC, RP [2-(phenylmethyl)phenol] and 4-(phenylmethyl)phenol. See Table 2 for product yield. A control experiment duplicating these reactions but omitting the iron(III) complex was run. No reaction took place and all the substrate was recovered at the end.

This work was extended to other substrates, for example, neopentylbenzene [$\text{PhCH}_2\text{C}(\text{CH}_3)_3$], bibenzyl ($\text{PhCH}_2\text{CH}_2\text{Ph}$), 1-methoxynaphthalene (PhPhOCH_3), and 1-methylnaphthalene (PhPhCH_3). Each substrate was refluxed with an oxidant in a solvent. For results, see Table 4. Neopentylbenzene reacted with the iron (III) complex but it was not a clean system and it was very difficult to identify the products. 1-Methoxynaphthalene and 1-methylnaphthalene were not expected to react because they would give very unstable methyl carbonion ions and H^+ . They did not react. The absence of dimer formation is interesting and worth further study.

EXPERIMENTAL

Materials

The solvents acetonitrile and benzonitrile were dried by refluxing them over calcium hydride for a minimum of 48 hrs followed by distillation. The oxidizing reagents tris(4-bromophenyl)aminium hexachloroantimonate, ammonium cerium(IV) nitrate, copper(II) acetate, copper(II) chloride, cupric acetylacetonate, ammonium hexachloroiridate(IV), iron(III) chloride and the iron powder were purchased from Aldrich while the iron(III)(1,10-phenanthroline)₃ complexes were prepared as described below and their purity was checked by u.v. spectroscopy.

The u.v. spectra were obtained on a Perkin-Elmer Lambda 5 UV/VIS spectrometer. Spectra were recorded at wavelength ranging between 200 nm to 800 nm using 2 nm slit width. Solutions containing the complex were examined in 1cm quartz cells. Pure acetonitrile was used as a reference sample.

The organic substrates which were not commercially available were also synthesized. The compounds were characterized by n.m.r., melting points, and gas chromatography (gc).

Iron(III)(1,10-phenanthroline)₃ Complexes

Tris-(1,10-phenanthroline)-iron(III) perchlorate was prepared by the literature method.⁸ 1,10-Phenanthroline (4.0 g, 0.02 mol) and iron(II) sulphate heptahydrate (1.84 g, 6.6×10^{-3} mol) were added to 40 cm³ water. The mixture was warmed and stirred until almost all the solid had dissolved and the solution was filtered into a flask which was immersed in an ice bath. 10 cm³ dilute sulfuric acid (ca. 1 M; 7 cm³ conc. H₂SO₄ in 120 cm³ distilled water) was added to this solution and after further cooling excess lead(IV) oxide (ca. four fold) was also added. The mixture was stirred until all the iron(II) (red) was converted to the iron(III) (blue) complex. Excess lead(IV) oxide was filtered off using a sintered funnel (could also use centrifuge) and sodium perchlorate (2.4 g, 0.02 mol) was added to the filtrate. A blue precipitate appeared in the solution which was filtered and recrystallized from dilute perchloric acid (1:1 v/v). The iron(III) complex was dried under vacuum (0.4 Torr) overnight and it was characterized by u.v. spectroscopy (λ_{max} (acetonitrile) for iron(III) complex was 594 nm, lit⁹ 592 nm). The synthesis of the complex was finished in one day because the complex decomposes in aqueous solutions.

An attempt was made to synthesize tris-(1,10-phenanthroline)-iron(III) tetrafluoroborate and hexafluorophosphate by the literature method¹⁰ but we could not oxidize the solution with lead(IV) oxide.

Benzyl Phenyl Ether^{11,12}

Benzyl phenyl ether was prepared¹¹ by treating equal molar quantities of potassium phenoxide (freshly prepared from equal molar quantities of phenol and potassium hydroxide) with benzyl bromide in a 1:1 mixture of ethanol and tetrahydrofuran (THF with stabilizer or Inhibitor BHT). Temperature of the refluxing mixture was kept as low as possible but the final yield was low.

High yield was obtained when it was synthesized by the literature method⁵ from benzyl chloride, phenol and potassium carbonate in acetone.

In both cases product was recrystallized twice from ethanol and dried under vacuum (0.4 Torr) overnight. It was white solid, mp 38-39° (lit.¹² 39°) and ¹H n.m.r. (CDCl₃,) 5.1(2H, s), 6.8-7.6 (10H, m). Only one peak was observed except solvent(CH₃CN) in gc chromatograph.

p-Bromobenzyl-p-bromophenyl ether¹²

A mixture of 4-bromobenzyl bromide (10 g, 0.04 mol), 4-bromophenol (7.6 g, 0.044 mol), potassium carbonate (5.5 g, 0.04 mol) and acetone (8 cm³) was refluxed overnight. After cooling the mixture was diluted with 8 cm³ of water and the product was extracted with ether (3 x 10 cm³). The ether solution was washed with aqueous alkali (sodium bicarbonate) and dried over magnesium sulphate. After removing the ether the product was recrystallized from

ethanol and dried over night under vacuum (0.4 Torr).

It is a white solid, yield 4.0 g (30%), and mp 111-111.5° (lit.¹² 111°).

Reactions

The apparatus was dried overnight in an oven at 110°C and assembled while it was hot. Benzyl phenyl ether (0.0503 g, 2.74×10^{-4} mol) and tris-(10-phenanthroline)-iron(III) perchlorate (16% molar excess, 0.2839 g, 3.17×10^{-4} mol) were placed in a two necked round bottom flask equipped with magnetic stirrer and water condenser under nitrogen in the dark. To this was added 5 cm³ freshly dried acetonitrile through a rubber septum using syringe and the flask was lowered into a preheated (110 °C) oil bath. During reaction the color of the solution changed from blue to red.

To follow the reaction, 1 cm³ of the reaction mixture was withdrawn through the septum at known intervals and worked up as described below.

The usual method of analysis involved quenching the reaction (1 cm³) with about an equal volume (1 cm³) of saturated aqueous sodium bicarbonate solution (pH=8). The organic materials were extracted with chloroform (3 x 1 cm³ each time), excess sodium chloride salt, dried over magnesium sulphate and analysed by gas chromatography (0.4 ul for each injection) on a column using naphthalene as an internal standard.

Product Analysis

The identification of the products was based on gc and gc/ms data. The gas chromatograph has a SPB1/SUPELCO 30 m long column with a flow rate of 2 cm³/min and a split ratio of 60/1. The temperature programming was initially for 6 min. at 110°C, program rate 110-200°C, 10°C min⁻¹, and finally for 5 min. at 200°C. Injector and detector temperature were both at 220°C.

The gc/ms was model 59/95 with data system RTE VII. The gas chromatograph was SPB5, 30 m/0.523 mm ID capillary column packed with fused silica. The temperature programming was as for the gc except that initial temperature was 70° instead of 110°.

The final concentrations of reactant and products were calculated from the areas relative to the standard (naphthalene unless otherwise stated) after corrections for the relative response of the species. Hence the mass balance was calculated for the whole reaction.

Table I. Recovery of Starting Material from Reaction 1 in Refluxing Acetonitrile.

Time (hr.)	PhOCH ₂ Ph recovered
0.25	5
0.5	3
1.0	3
1.5	0

Table 2. Product yields (weight %)^a for benzylphenylether (BPE) reacted in acetonitrile at 82°C.

Oxidizing reagent	P	BAC	RP sub ^a at 2	% Conv
Iron(III) (1,10-phen) perchlorate	52	31	13	100
Iron(III) chloride	16	24	18	58
Iron(III) (1,10-phen) perchlorate (catalytic amount)/ air (O ₂)	28	45	22a	96

a) Reported yields are accurate to $\pm$ 10%.

b) 17% 2 substituted and 5% 4 substituted.

Table 3. Benzlphenylether Conversions in Refluxing Acetonitrile with a variety of Oxidizing Agents.

Oxidizing reagent	Outer sphere(OS)/	% conv ($\pm$ 10%)
	Inner sphere(IS) mechanism ^a of BPE	
Tris(4-bromophenyl)	IS	35 (22 hrs., 25 °C)
ammonium hexachloro		68 (1/2 hr., 82 °C)
antimonate ^a		
Cerium(IV)ammonium	mainly OS/IS	51 (92 hrs., 82°C)
nitrate		
Ammonium hexachloro	OS/IS	0 (74 hrs., 82°C)
iridate(IV)		
Copper(II)chloride	very weak OS	0 (91 hrs., 82°C)
Copper(II)acetate	very weak OS	0 (116 hrs., 82°C)
Cupric acetylacetonate	OS	0 (22 hrs., 82°C)

^a Product analysis and mass balance was difficult because a lot of side reactions of the radical and cation produced from the substrate took place with the oxidizing reagent. Bromine substituted products were identified, it was not a clean system.

Table 4. Oxidations of Hydrocarbions with Fe(III)

(1, 10-phenanthroline)₃ Perchlorate.

Substrate	Oxidant	Solvent	Temp/°C	Time/hrs.	Reaction took place
PhCH ₂ C(CH ₃) ₃	A	PhCN	188	44	Yes (83 % Conv)
PhCH ₂ C(CH ₃) ₃	B	CH ₃ CN	82	72	No
PhCH ₂ C(CH ₃) ₃	B	PhCN	188	43	No
PhCH ₂ CH ₂ Ph	B	PhCN	188	43	No
PhPhOCH ₃	A	CH ₃ CN	82	72	No
PhPhOCH ₃	B	CH ₃ CN	82	44	No
PhPhCH ₃	A	CH ₃ CN	82	72	No

A Iron(III)(1,10-phenanthroline) perchlorate

B Tris(4-bromophenyl)aminium hexachloroantimonate

REFERENCES

1. A) L. W. Reichel, G. W. Griffin, A. J. Muller, P. K. Das, and S. N. Ege, Can. J. Chem., 1984, 62, 424; H. F. Davis, P. K. Das, L. W. Reichel, G. W. Griffin, J. Am. Chem. Soc., 1984, 106, 6968.
2. A. Okamoto, D. R. Arnold, Can. J. Chem., 1985, 63, 2340.
3. D. M. Camaioni, J. A. Franz, J. Org. Chem., 1984, 49, 1607.
4. W. Wiser, DOE Symposium Series on Scientific Problems Relevant to Coal Utilisation, Conf. No. 770509 (West Virginia University) Oak Ridge Laboratories, TN, Nov., 1977.
5. S. Fukuzumi, C. L. Wong, J. K. Kochi, J. Am. Chem. Soc., 1980, 102, 2928.
6. C. J. Schlessener, C. Amatore, J. K. Kochi, J. Am. Chem. Soc., 1984, 106, 3567; J. Phys. Chem., 1986, 90, 3747.
7. C. J. Schlessener, J. K. Kochi, J. Org. Chem., 1984, 49, 3142.
8. (a) N. Sutin and B. M. Gordon, J. Am. Chem. Soc., 1961, 83, 70. (b) M. H. Ford-Smith and N. Sutin, J. Am. Chem. Soc., 1961, 83, 1830.
9. C. L. Wong and J. K. Kochi, J. Am. Chem. Soc., 1979, 101, 5593.
10. Richard E. Desimone and Russell S. Drago, J. Am. Chem. Soc., 1970, 92, 2343.
11. G. A. Russell and Roger C. Williamson, Jr., J. Am. Chem. Soc., 1964, 86, 2357.
12. S. G. Powell and Rogger Adams, J. Am. Chem. Soc., 1920, 42, 646.

END

**DATE
FILMED**

3 / 26 / 93