**Thermodynamic Properties for Polycyclic Systems
By Non-Calorimetric Methods**

**Progress Report
for Period August 1, 1992 - September 30, 1993**

W. V. Steele, R. D. Chirico, and T. D. Klots

**IIT RESEARCH INSTITUTE
NATIONAL INSTITUTE FOR PETROLEUM AND ENERGY RESEARCH
P.O. BOX 2128
BARTLESVILLE, OKLAHOMA 74005**

March 1993

Prepared for

**THE U.S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-92ER14288**

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. References herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ABSTRACT

This report describes the progress made in several directions (Research Paths) on the project Thermodynamic Properties of Polycyclic Systems by Non Calorimetric Methods. A detailed vibrational spectroscopic study of furan, pyrrole, and thiophene has been completed. These compounds form part of the base of five-membered ring systems on which the rest of the research program will be built. Several methyl-substituted derivatives were also studied. The results will be used to confirm the model for alkyl-substitution in the ring systems. Gas-phase spectra and fundamental-frequency assignments were completed for 2,3- and 2,5-dihydrofuran. Those compounds initiate work on ring-puckering within the research program. A paper describing the need for third virial estimation, when using the virial equation of state to derive thermodynamic properties at pressures greater than 1 bar was completed.

I. OBJECTIVES

The main objective of this research project is the development of the thermodynamic properties for polycyclic molecules containing five-membered ring(s) via assigned spectra and statistical mechanics. A goal is the attainment of consistent spectroscopic and calorimetrically derived properties over a wide temperature range. Specific emphasis will be placed on thermodynamic property values for reduced temperatures greater than 0.7 (approaching the critical region). The fulfillment of these objectives will facilitate the development of a consistent theoretical interpretation of the fundamental vibrational modes for ring systems with as many as four rings. Such an interpretation will provide a more fundamental understanding of the foundation of estimation techniques such as group-additivity and molecular-mechanics approaches. Success in the goals would aid Chemical & Material Science researchers in designing, via computer simulation, optimized molecules for specific applications.

II. INTRODUCTION

This report summarizes progress during the first year: August 1, 1992 through July 31, 1993 on the research project, "THERMODYNAMIC PROPERTIES OF POLYCYCLIC SYSTEMS BY NON-CALORIMETRIC METHODS," which is supported by DOE contract DE-FG02-92ER14288. This research program provided a valuable complement to the experimental programs currently in progress here at NIPER for the Advanced Exploration & Process Technology (AEPT) division of the Department of Energy. A portion of those experimental programs focuses on the calorimetric determination of thermodynamic properties of key polynuclear heteroatom-containing aromatic molecules. This project for the Office of Energy Research focuses on the non-calorimetric determination of thermodynamic properties through the extension of existing correlation methodologies and through molecular spectroscopy with statistical mechanics.

This Progress Report outlines the research accomplished to date and planned for completion during the remainder of the first year. Progress in each of the three Research Paths detailed in the original proposal to OER is discussed below. In the first six months significant progress was made in the research and the Milestones detailed in the original proposal remain "on track". Highlights of the research accomplished include:

- A detailed vibrational spectroscopic study of furan, pyrrole, and thiophene was completed. For the first time, the fundamental vapor-phase vibrational frequencies were completely defined for all three molecules. (A paper describing the research was submitted to *Spectrochimica Acta*.)
 - Improved ideal-gas thermodynamic functions were calculated for each compound using the now complete vapor-phase fundamental frequencies.
 - Effects due to anharmonicity were shown to be less than the uncertainty interval in the calorimetrically derived ideal-gas entropies.
- Gas-phase vibrational spectra were obtained for 2-methylfuran, 2- and 3-methylthiophene. (A paper describing the research is in progress.)
 - Results showed good agreement between calorimetrically-derived ideal-gas entropies and values derived using statistical mechanics in a narrow region where extrapolation of the experimentally-derived calorimetric values was a minimum.
 - Details of the affects due to anharmonicity await extension of the calorimetric data in the coming quarter.
- Complete gas-phase fundamental frequency assignments were made for 2,3-dihydrofuran and 2,5-dihydrofuran. (A paper describing the research is in progress.)
 - The assignments are sufficient to allow calculation of ideal-gas entropies for either molecule with an uncertainty interval of $\pm 0.05 R$ at 400 K.
- A paper, "Reconciliation of calorimetrically and spectroscopically derived thermodynamic properties at pressures greater than 0.1 MPa for benzene and methylbenzene; the importance of the third virial coefficient," was completed.
 - Research accomplished delineated the significance of the third virial coefficient even at 1 bar pressure.
 - A literature correlation for estimation of third virials was shown to be applicable to monoaromatics.

III. DETAILS OF PROGRESS

III.1 Research Path 1: Assignment of vibrational spectra for polycyclic molecules containing 5-membered rings

[Application and extension of the scaled force constant method;⁽¹⁾ based on the method of Pulay et al.^(2,3)]

Figure 1 shows some of the interrelationships between the ring systems being studied in this research. The early work in this project was directed at establishing clearly the frequency assignments for the monocyclic building blocks. A review of previous vibrational studies for several of these single-ring compounds was made, including furan, oxazole, pyrrole, imidazole, thiophene, thiazole and cyclopentadiene. The search revealed that, despite many past investigations, several ambiguities remained in the vibrational assignment for some of these molecules, including for furan and pyrrole. Common problems included the lack of a measured vapor phase frequencies; or disagreement between the various assignments. Without a firm basis for the assignments of these single-ring compounds, the progress of the vibrational assignments for the bicyclic compounds would be impeded. Also, the single-ring results can be used to assess anharmonic contributions to molecular thermodynamics, and that requires a complete set of vapor frequencies.

A detailed vibrational spectroscopic study of furan, pyrrole, and thiophene was completed. Special emphasis was placed on obtaining high-quality Raman vapor spectra, since only for furan has the Raman vapor spectrum previously been observed. The vapor phase vibrational frequencies for these molecules are now completely known for the first time. This will benefit other areas of molecular research, including high-level *ab initio* calculations of the vibrational frequencies for these important test molecules. A full description of the work is found in the paper, "COMPLETE VAPOR PHASE ASSIGNMENT FOR THE FUNDAMENTAL VIBRATIONS OF FURAN, PYRROLE AND THIOPHENE," which is being submitted to *Spectrochimica Acta* for peer review (copy enclosed with this Progress Report; Exhibit 1).

Figure 1. Two-ring compounds to be studied within research path 1 of this proposed project, and their monocyclic analogs. The highlighted two-ring compounds were studied in the previous project for OER.

Related work for some methyl derivatives of these compounds has also been completed. Spectroscopic studies have been made on 2-methylfuran, and 2- and 3-methylthiophene. Much of the basis for their assignments comes from previous work by Scott^(4,5) who analyzed *liquid-phase* spectra for these molecules and evaluated the force fields of these compounds, i.e. furan and thiophene and their deuterium and methyl derivatives. Calorimetrically derived values for the ideal-gas entropies of these compounds have been reported in the literature.⁽⁶⁻¹⁰⁾ By studying them, the results provide insight into the spectroscopic and calorimetric effects of adding a methyl group to a ring system. The same kind of agreement between the spectroscopic and calorimetric results for the parent compounds is sought for the methyl derivatives.

Methyl-substitution often lowers the melting point of a parent organic molecule. Lowering of the melting point can facilitate the collection of the vibrational spectra. It is anticipated that methyl derivatives will be used in the study of some of the two and three-ringed molecules planned in this research.

The gas-phase spectra determinations for 2-methylfuran, and 2- and 3-methylthiophene resulted in several assignments being changed for each molecule, principally because of the new information provided by the IR vapor bandshapes and Raman vapor spectra. The methyl torsion and moments of inertia have been previously studied for each molecule using microwave spectroscopy.⁽¹¹⁻¹³⁾ With the proposed set of vibrational frequencies, the ideal-gas thermodynamic properties were calculated using statistical mechanics. Figure 2 shows the difference in ideal-gas entropy between the spectroscopic and calorimetric determinations for 2-methylfuran. The vapor pressures of 2-methylfuran have been measured from 250 - 490 K,⁽⁶⁻⁸⁾ while its liquid heat-capacities have been determined from 190 - 310 K.⁽⁶⁾ In the approximate region of overlap, the agreement in ideal-gas entropy is good. These results suggest that the assignment and methyl torsion are likely modeled correctly. However, in order to assess the effects of anharmonic contributions to the ideal-gas entropy, the liquid-phase heat-capacity range will have to be extended to higher temperatures (see below).

A paper describing the results of the research on the methyl-substituted compounds is in preparation.

Figure 2. A plot of the difference in ideal-gas entropy for 2-methylfuran: $S(\text{spectroscopy}) - S(\text{calorimetry})$. The thick bar indicates the temperature range where the calorimetric data is most reliable.

III.11 Research Path 2: Ring puckering and its ramifications.

Topics of study will include, 1) assignment of vibrational spectra for mono- and polycyclic molecules with ring puckering, 2) the effect of the ring puckering on derived thermodynamic properties, and 3) ramifications for group-contribution methods for property predictions.

Spectroscopic studies of 2,3-dihydrofuran and 2,5-dihydrofuran are nearing completion. These molecules contain a low-frequency ring-puckering motion, which has been extensively studied in the past.⁽¹⁴⁻¹⁹⁾ The thermodynamic implications of such a ring-puckering vibration has for the most part been worked out.^(20,21) However, without knowledge of the other $3n-6$ molecular vibrations, the total molecular statistical entropy cannot be computed.

Other than its ring-puckering transitions, the vibrational spectrum of 2,3-dihydrofuran has not been previously analyzed. There has been one previous attempt, that by Fortunato,⁽²²⁾ to make the assignment for 2,5-dihydrofuran. The low-frequency ring-puckering mode complicates the IR and Raman vapor spectra for both of these molecules, particularly for 2,3-dihydrofuran because of its lower symmetry and because of its non-planarity in the lowest two puckering levels (the ground and first excited states).

The extensive work by Villarreal *et al.*^(23,24) on the spectroscopy of cyclopentene has been used as a template. Scale factors for cyclopentene were determined by adjusting the calculated AM1 force field to best fit the assigned vibrational frequencies for the parent compound and three of its deuterium derivatives. These scale factors were then used to make a prediction of the vibrational frequencies for 2,5-dihydrofuran. An assignment for 2,5-dihydrofuran was then made by considering the IR liquid, IR vapor, Raman liquid, and Raman vapor spectra and the calculated frequencies. The new assignment, given in Table I, differs mainly from the Fortunato⁽²²⁾ assignment in the choice of one B1 vibration.

The scale factors for 2,5-dihydrofuran were adjusted to fit the observed frequencies with those calculated from the scaled AM1 force field. Scale factors for cyclopentene and 2,5-dihydrofuran were combined to predict frequencies for 2,3-dihydrofuran. An assignment was made based on its predicted frequencies and observed spectra. This assignment is given in Table I. The vibrational frequencies were in very good agreement with the predicted values (less than 20 cm^{-1}). However, two A'' vibrations are not identifiable from the recorded spectra, presumably because they are too weak. These two A'' vibrations, ν_{22} and ν_{24} , are calculated at ~ 1080 and 818 cm^{-1}

and have estimated windows of uncertainty of 20 and 25 cm^{-1} respectively. A recording of the solid state IR spectrum of 2,3-dihydrofuran is planned in an attempt to observe these vibrations. Sharper bands in the solid may enable observation of a weaker band that is hidden in the liquid or vapor spectra. Full details of the spectra will be published in a paper in progress.

Table I. Fundamental vibrational frequencies for 2,5-dihydrofuran and 2,3-dihydrofuran in the vapor and liquid phases (in cm^{-1}).

2,5-dihydrofuran			2,3-dihydrofuran		
	vapor	liquid		vapor	liquid
<i>A1 species</i>			<i>A' species</i>		
1	3096	3088	1	3122.0	3110
2	2865	2852	3	(2935) ^a	2923
3	1625.5	1618.5	5	1624.8	1618.5
4	1489.5	1483	7	1459	1453
5	1363	1360	8	1378	1375
6	1097.0	1097	11	1141.0	1136.5
7	982.5	982	13	995.5	994
8	901.5	890	15	920	913
9	740	737	17	674	668
<i>B1 species</i>			<i>A' species</i>		
15	(3092) ^a	3083	2	3110.5	3099
16	(2865) ^c	(2852) ^c	4	(2875) ^a	2863
17	1480	1475	6	1482	1476
18	1350	1346	9	1305	1299
19	1307	1304	10	1278	1276
20	1092	1080	12	1066	1057
21	925	923	14	929.5	927
22	801	799	16	830	826
<i>A2 species</i>			<i>A'' species</i>		
10	2888	~2880	18	(2980) ^a	2965
11	1195	1200	20	1220	1225
12	1039	1035	22	1070 $\pm$ 20 ^b	
13	890 $\pm$ 20 ^b		23	885.2	887
14	394	400	26	447.5	450
<i>B2 species</i>			<i>A'' species</i>		
23	2883	(2875) ^a	19	(2905) ^a	2892
24	1167	1174	21	1192.0	1187
25	1011	1012	24	825 $\pm$ 30 ^b	
26	661	669	25	705.0	709
27	RP	NA	27	RP	NA

a) Values in parenthesis have estimated phase shift corrections applied.

b) These values are best estimates based on the scaled force field fit.

c) Placed at same frequency as corresponding A1 vibration (ν_2).

Figure 3. Candidate compounds for study within research path 2. Compounds with ring-puckering or large-amplitude out-of-plane deformations.

III.III Research Path 3: Application of Path 1 and Path 2 results to test physical-property correlations

The research in this area will use the results obtained in Research Paths 1 and 2, in combination with available calorimetric results already obtained within separate projects at NIPER, to test the validity of several physical-property correlations. The primary areas of research will be, 1) to test the efficacy of widely used vapor-pressure equations (Antoine,⁽²⁵⁾ Cox,⁽²⁶⁾ Wagner,⁽²⁷⁾ etc.) in the extrapolation of vapor-pressure data beyond the range of the experimental measurements; and 2) to test corresponding-states type virial-coefficient correlations^(28,29) in their application to polycyclic aromatics. The correlation for second virial coefficients is well established.⁽²⁸⁾ The primary concentration will be on the third virial coefficient correlation.⁽²⁹⁾

This area of research is due to start in the second year of this project when the spectroscopy is sufficiently advanced. As a prelude to the research in this area, the paper, "Reconciliation of calorimetrically and spectroscopically derived thermodynamic properties at pressures greater than 0.1 MPa for benzene and methylbenzene; the importance of the third virial coefficient," was completed. That paper is a "carry-over" from the research completed in the previous project for OER. A copy of the manuscript is enclosed (Exhibit 2)

IV. OUTLINE OF RESEARCH FOR REMAINDER OF FIRST YEAR

IV.1 Research Path 1: Assignment of vibrational spectra for polycyclic molecules containing 5-membered rings

[Application and extension of the scaled force constant method;⁽¹⁾ based on the method of Pulay et al.^(2,3)]

In this Research Area refined scale factors for furan, pyrrole, and thiophene will be determined. Studies leading to gas-phase fundamental vibrational frequencies for thiazole, oxazole, and isoxazole will be completed. Application of these results to the spectroscopy of two ring compounds will also begin. Two ring compounds that will be studied are indole, benzofuran, and indene. Some work has already begun on these compounds.

IV.11 Research Path 2: Ring puckering and its ramifications.

Topics of study will include, 1) assignment of vibrational spectra for mono- and polycyclic molecules with ring puckering, 2) the effect of the ring puckering on derived thermodynamic properties, and 3) ramifications for group-contribution methods for property predictions.

Results and methodology derived in the dihydrofuran studies will be extended into studies leading to gas-phase fundamental vibrational frequencies for phthalan and 4,5-dihydro-2,3-benzofuran. Results obtained from the spectroscopy will be compared with ideal-gas values derived previously for both compounds under DOE Fossil Energy Programs at NIPER.

V. REFERENCES

1. Collier, W. B. *J. Chem. Phys.* **1988**, *88*, 7295.
2. Pulay, P.; Fogarasi, G.; Pang, F.; Boggs, J. E. *J. Am. Chem. Soc.* **1979**, *101*, 2550.
3. Pulay, P.; Fogarasi, G.; Boggs, J. E. *J. Chem. Phys.* **1980**, *74*, 3999.
4. Scott, D. W. *J. Mol. Spectrosc.* **1971**, *37*, 77 .
5. Scott, D. W. *J. Mol. Spectrosc.* **1969**, *31*, 451.
6. Kobe, K. A., Ravicz, A. E., Vohra, S. P. *J. Chem. Eng. Data* **1956**, *1*, 50.
7. Carlson, H. G., Westrum, E. F. Jr. *J. Chem. Eng. Data* **1965**, *10*, 134 .
8. Eon, C., Pommier, C., Guiochon, G. *J. Chem. Eng. Data* **1971**, *6*, 408.
9. Pennington, R. E., Finke, H. L., Hubbard, W. N., Messerly, J. F., Frow, F. R., Hossenlopp, I. A., Waddington, G. *J. Am. Chem. Soc.* **1956**, *78*, 2055.
10. McCullough, J. P., Sunner, S., Finke, H. L., Hubbard, W. N., Gross, M. E., Pennington, R. E., Messerly, J. F., Good, W. D., Waddington, G. *J. Am. Chem. Soc.* **1953**, *75*, 5075.
11. Norris, W. G., Krisher, L. C. *J. Chem. Phys.* **1969**, *51*, 403.
12. Pozdeev, N. M., Gunderova, L. N., Shapkin, A. A.. *Opt. Spektrosk.* **1970**, *28*, 254.
13. Ogata, T., Kozima, K. *J. Mol. Spectrosc.* **1972**, *42*, 38 .
14. Ueda, T., Shimanouchi, T. *J. Chem. Phys.* **1967**, *47*, 4042.
15. Ueda, T., Shimanouchi, T. *J. Chem. Phys.* **1967**, *47*, 5018 .
16. Carreira, L. A., Lord, R. C. *J. Chem. Phys.* **1969**, *51*, 3225 .
17. Green, W. H. *J. Chem. Phys.* **1969**, *50*, 1619.
18. Durig, J. R., Carreira, L. A. *J. Chem. Phys.* **1972**, *56*, 4966.
19. Durig, J. R., Carter, R. O., Carreira, L. A. *J. Chem. Phys.* **1973**, *59*, 2249.
20. Dorofeeva, O. V., Gurvich, L. V., Jorish, V. S. *J. Phys. Chem. Ref. Data* **1986**, *15*, 437.
21. Dorofeeva, O. V. *Thermochemica Acta* **1992**, *194*, 9 .
22. Fortunato, B. *Gazz. Chim. Ital.* **1976**, *106*, 799.
23. Baumann, L. E., Killough, P. M., Cooke, J. M., Villarreal, J. R., Laane, J. *J. Phys. Chem.* **1982**, *86*, 2000.
24. Villarreal, J. R., Laane, J., Bush, S. F., Harris, W. C. *Spectrochimica Acta* **1979**, *35A*, 331.
25. Antoine, C. *C. R. Acad. Sci.* **1888**, *107*, 681.
26. Cox, E. R. *Ind. Eng. Chem.* **1936**, *28*, 613.
27. Wagner, W. *Cryogenics* **1973**, *13*, 470.
28. Pitzer, K. S.; Curl, R. F. Jr. *J. Am. Chem. Soc.* **1957**, *79*, 2369.
29. Orbey, H.; Vera, J. H. *AIChE Journal*, **1983**, *29*, 107.

Publications from this research

1. Klots, T. D.; Chirico, R. D.; Steele, W. V. *Complete Vapor Phase Assignment for the Fundamental Vibrations of Furan, Pyrrole and Thiophene*. Submitted to *Spectrochimica Acta*, March 1993.
2. Klots, T. D.; Collier, W. B. *Fundamental Vibrational Frequency Assignments for 2,3-Dihydrofuran and 2,5-Dihydrofuran*. In preparation for submission to *Spectrochimica Acta* in April 1993
3. Klots, T. D.; Chirico, R. D.; Steele, W. V. *Complete Vapor Phase Assignment for the Fundamental Vibrations of 2-Methylfuran, 2-, and 3-Methylthiophene*. In preparation for submission to *Spectrochimica Acta* in April 1993
4. Steele, W. V.; Chirico, R. D. *Reconciliation of Calorimetrically and Spectroscopically Derived Thermodynamic Properties at Pressures Greater Than 1 Bar for Benzene and Methylbenzene; The Importance of the Third Virial Coefficient*. To be submitted to *Ind. Eng. Chem. Res.*, April 1993.

Presentations

1. Klots T. D. *Vibrational Assignments for Bicyclic Aromatic Compounds using a Scaled Quantum-Mechanical Method.* presented at the 47th. Ohio State International Symposium on Molecular Spectroscopy, June 15-19, 1992.
2. Klots T. D. *IR Overtone Region for the Out-of-plane Wags of Bicyclic Aromatic Compounds: Clues to Hidden Frequencies.* presented at the 47th. Ohio State International Symposium on Molecular Spectroscopy, June 15-19, 1992.

END

DATE
FILMED
5/13/93