NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

Endothermic Photo-Catalytic Reactions

Final Report

NREL/TP--253-4277

DE91 002142

H.W. Prengle, Jr., W.E. Wentworth, K.C.
Polonczyk, M. Saghafi, J.A. Wilking, and
K.S. Kramer
University of Houston
Houston, Texas

NREL Technical Monitor: R. Gerald Nix

National Renewable Energy Laboratory
(formerly the Solar Energy Research Institute)
1617 Cole Boulevard
Golden, Colorado 80401-3393
A Division of Midwest Research Institute
Operated for the U.S. Department of Energy
under Contract No. DE-AC02-83CH10093

Prepared under Subcontract Nos. XX-7-07028-1
and XX-6-06006-1

April 1992

MASTER

This report describes subcontracted research. The report is unreviewed and expresses only the opinions of the author[s]. It has been prepared for reproduction from the best available copy.

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or any agency thereof.

Printed in the United States of America
Available from:
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

Price: Microfiche A01
Printed Copy A05

Codes are used for pricing all publications. The code is determined by the number of pages in the publication. Information pertaining to the pricing codes can be found in the current issue of the following publications which are generally available in most libraries: *Energy Research Abstracts (ERA)*; *Government Reports Announcements and Index (GRA and I)*; *Scientific and Technical Abstract Reports (STAR)*; and publication NTIS-PR-360 available from NTIS at the above address.

. EXECUTIVE SUMMARY .

The overall objective of this report is to present the results of an investigation to provide guidelines for future experimental work, on solar energy driven endothermic photo-catalytic reactions, and primarily to select candidate synthesis reactions which lead to high \$-value products. Since the initial report a group of exothermic aromatic and olefin oxidation reactions have been included also, as possible night-time reactions.

An intensive literature search was conducted to find properties, market demand, and prices of pertinent chemicals; meeting four criteria: 1) -the reaction must be endothermic and favorable; 2) -the reaction must be catalytic; 3) -the product must be produced from low cost feedstocks; and 4) -the product must have a sales price >\$1.00/lb. A price histogram of some 200 chemicals was generated indicating a bimodal distribution - a low price (< \$0.85/lb) category, and a higher price (> \$1.00/lb) category. An attempt was made to correlate price as a function of molecular weight and structural (complexity) factor; the latter was 4-5 times more important.

Initial examination of low cost feedstocks to high value products lead to consideration of n-paraffins to aromatics and substituted aromatics. Fifteen candidate endothermic synthesis reactions, meeting the above criteria, are suggested. The ratio of product price by reactant cost indicates ~ 5-8 for the best possibilities; all can be visualized as starting with low cost paraffin and methanol feedstocks.

The revised report contains an extensive section on reaction thermodynamics of the synthesis reactions, aromatic oxidation reactions, and olefin oxidation reactions, giving the pertinent thermodynamic reaction quantities as a function of temperature over the range 298-1000 K.

Presently operating commercial processes for the candidate and related reactions are discussed; including the initial reaction steps, as well as synthesis of the final products.

The investigation leads to six primary conclusions concerning, price distribution of chemicals, importance of conversion of paraffins to aromatics, alkylation-hydroxylation of the aromatic ring, reaction selection by the product price to reactant cost ratio, catalyst for optimum yield of aromatics, and the importance of olefin conversion to oxygenated products.

. TABLE OF CONTENTS .

	page
Executive Summary	iii
List of Tables and Figures	v
Introduction and Background	1
Literature Search	2
Candidate Reactions	13
Reaction Thermodynamics	17
Synthesis Reactions & Oxidation Reactions	
Commercial Processes	38
Conclusions and Recommendations	42
Selected References	43
APPENDIX	
A. Estimation of Thermodynamic Properties	a-1
B. Recommended Catalysts for Endothermic Cyclization- Addition Reactions	b-1
C. Chemical Prices- CHEMICAL MARKETING REPORTER (9 June 1986)	c-1
D. Top 50 Chemicals Production Turned Back Up in 1987 (4-11-88)	d-1
E. World Chemical Outlook: USA & Foreign Trade (12-14-87)	e-1
F. HPI's Role in Chemical's Future (Feb 1988)	f-1
G. Catalyst Production & Use	g-1
(5-17-88)	

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Pertinent Primary Feedstocks Data	6
2	Pertinent Intermediate Chemical Data	7
3	Pertinent Product Data	8
4	Pertinent Specialty Chemical Data	9
4a	Chemical Structure Factors in Ascending Order	12
5	Favorable Synthesis Reactions	14
6	Price/Cost Ratios and Annual Production	15
7-12	Synthesis Reactions - thermodynamic calculations and temperature dependent functions	18
13-16	Oxidation Reactions of Aromatics - thermodynamic calculations and temperature dependent functions	24
17-25	Oxidation Reactions of Olefins - thermodynamic calculations and temperature dependent functions	28
26	Price/Cost Ratios and Annual Production, Oxidation Products	37
A-1	Benson Group Contributions to Ideal Gas Properties	a-2
B-1	Recommended Catalysts for the Endothermic Cyclization-Addition Reactions	b-1
C-1	Chemical Prices	c-1

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	Price Histogram of Chemicals	3
2	Price vs. MW Scatter Plot	4
3	Price as a Function of Mol. Wt. & SF	4
4	Aromatics from Coal Gas and Tar Light Oil	39
5	Aromatics from Petroleum by Reforming	39

INTRODUCTION & BACKGROUND

Although the objective of this study was to recommend candidate reactions for experimentation of photo-catalytic endothermic reactions, for continuity and completeness a brief summary of past and present related work will be discussed. Previously, Wentworth and coworkers [1] had determined an appropriate reactor configuration which would allow the direct use of high flux solar radiation to carry out endothermic chemical reaction. By implementing a Xenon lamp to simulate the full solar spectrum over a shallow 1 cm diameter fluidized bed reactor (relaxed bed height of 0.5 cm), Wentworth was able to compare the effects of high flux radiation with those produced by thermal input alone. From these results it was observed that photo-catalysis produced higher yields and increased reaction rate constants in numerous cases than those achieved by conventional industrial processes. In lieu of these findings, it became obvious that an appropriate set of criteria should be defined and implemented as a guideline for future work.

The pertinent criteria for selecting candidate reactions are:

1. the reaction must be endothermic;
2. the reaction must be conventionally catalytic;
3. the product can be synthesized from low cost feedstocks;
- and 4. the product must be of high \$-value (>\$1.00). These criterion then lead to favorable economic advantages. Due to the availability of heat induced by the direct use of solar radiation, it is advantageous to choose reactions which are endothermic in nature. High market prices of specialty chemicals can be traced to two obvious effects: either the product is presently synthesized

from expensive feedstocks or the common industrial synthesis is complicated by process constraints (e.g. catalyst deactivation). Utilizing photo-catalysis could produce a high value product from an alternate cheap feedstock and may possibly overcome the inherent problems associated with standard industrial methods.

LITERATURE SEARCH

The search began by considering overall market demands in the current chemical industry [2]. The top 50 chemicals produced were reviewed and are attached in the Appendix [3]. A tabulation of chemicals with high production rates was formed. Early in the project it was expected that oxygen and nitrogen containing compounds would be of high value.

A detailed listing of price quotations of common chemicals is attached in the Appendix and was used in this project to assign a \$-value to the chemicals of interest [4]. A price histogram is presented in Figure 1. It is interesting to note that it is bi-modal, indicating a distinct separation between low and high value products at \$1.00. The first mode has a mean value of \$0.4 and the second has a mean value of \$1.42. This analysis based on current market value lead to the separation of the list as follows primary feedstocks, intermediate chemicals, products, and specialty chemicals.

As presented in Figure 2, by generating a scatter plot of chemical prices versus molecular weight, it became apparent that a relationship existed. It was further suggested by Prengle [5], that this relationship was also dependent upon the complexity of

FIGURE 1 - PRICE HISTOGRAM - 1ST DIST. MEAN = 0.4127, S.D. = 2.158E-02
 2ND DIST. MEAN = 1.422, S.D. = 3.122E-02

FIGURE 2 Price vs. MW Scatter Plot

FIGURE 3 Price as a Function of Mol. Wt. & SF

structure. The chemicals were arranged in order of complexity and assigned a structure factor. After analyzing fundamental groups, the following guidelines were chosen to assign values to more complex compounds.

1. Straight chain molecules were given a value of 0.10 for C_1-C_6 , and 0.20 for C_7 to each carbon in the chain. example: pentane, $CH_3CH_2CH_2CH_2CH_3$ $SF = 5(0.1) = 0.50$.

2. Groups that were arranged in the iso- and tert-configurations were multiplied by 1.15 and 1.25, respectively. example: Isobutane, $CH_3CH(CH_3)_2$ $SF = 0.2 + 1.15(0.20) = 0.43$.

3. Double bonds were accounted for by doubling the contribution. example: n-Butene, $CH_3CH_2CH=CH_2$ $SF = 2(0.20) + 0.20 = 0.60$.

4. Halide substitutions were given a contribution of 0.10. example: vinyl chloride, $CH_2=CHCl$ $SF = 2(0.20) + 0.10 = 0.50$.

5. An aromatic ring was given a value of 0.35. Substituents on an ortho and para directed ring were multiplied by 1.1 for ortho substituents, 1.3 for para substituents, and 1.6 for meta substituents. Meta directed rings were similarly multiplied by 1.1 for meta, 1.3 for ortho, and 1.6 for para. example: Dinitrophenol $HOC_6H_3-2,4(NO_2)_2$ $SF = (1.6)(1.1)(0.35 + 0.10 + 0.50) = 1.67$.

6. Oxygen containing groups were given the following contributions: alcohols, $-OH = 0.10$; aldehydes, $-CHO = 0.30$; carboxylic acids, $-COOH = 0.35$; ether, $-O- = 0.425$; ketones, $-CO- = 0.45$; and esters, $-COO- = 0.60$. example: Acetone, CH_3COCH_3 $SF = 0.45 + 0.20 = 0.65$.

New structure factors should be assigned according to the guidelines above and compared to similar structures found in Tables

Table 1 Pertinent Primary Feedstocks Data

Chemical	Symbol	Price (\$/lb) (6-6-86)	MW	SF	Production (billion lbs/yr) (1975)	Hf (298K) (kcal/mol)	Sf (298K) (kcal/mol)
1 Xylenes	(pet.)				*5.250		
m-Xylene	1,3-(CH ₃) ₂ C ₆ H ₄	0.360	106.17	0.88		4.12	28.1
p-Xylene	1,4-(CH ₃) ₂ C ₆ H ₄	0.195	106.17	0.72	*4.720	4.29	28.1
o-Xylene	1,2-(CH ₃) ₂ C ₆ H ₄	0.125	106.17	0.61	0.703	4.54	29.1
2 Bromide	Br ₂	0.330	159.80	0.20	0.408	0.00	0.0
3 Isobutylenes	(CH ₃) ₂ C=CH ₂	0.320	56.10	0.63	0.550	-1.67	15.1
4 Phosphoric Acid	H ₃ PO ₄	0.290	98.00	0.40	21.889		
5 n-Butenes	CH ₃ CH ₂ CH=CH ₂	0.260	56.12	0.60	1.480	-1.67	15.1
6 Ethyl Alcohol	CH ₃ CH ₂ OH	0.194	46.07	0.30	1.429	-56.12	-40.1
7 Toluene	C ₆ H ₅ CH ₃	0.190	92.13	0.45	*4.990	11.95	29.1
8 Ethylene	CH ₂ =CH ₂	0.185	28.05	0.40	*30.550	12.50	16.1
9 Ethylene Dibromide	BrCH=CHBr	0.170	185.85	0.60	0.275		
10 Ethylene Dichloride	ClCH=CHCl	0.170	96.94	0.60	*13.630	0.30	5.1
11 Propylene	CH ₂ =CHCH ₃	0.160	42.08	0.50	*14.93	4.83	14.1
12 C ₆ -C ₇ Paraffins		0.160			*4.41		
n-Hexane	CH ₃ (CH ₂) ₄ CH ₃		86.17	0.52		-39.96	0.1
heptane	CH ₃ (CH ₂) ₅ CH ₃		100.17	0.57		-44.86	1.1
13 Cyclohexane	C ₆ H ₁₂	0.156	84.16	0.60	*1.740	-29.43	7.1
14 Cumene	(CH ₃) ₂ CHC ₆ H ₅	0.145	120.20	0.55	*3.330	0.94	32.1
15 Benzene	C ₆ H ₆	0.123	78.11	0.35	*9.730	19.82	30.1
16 Urea	H ₂ NCONH ₂	0.107	60.06	0.40	*13.110	-58.70	-36.1
17 Formaldehyde	HCHO	0.106	30.05	0.30	*5.700	-27.70	-26.1
18 Normal Paraffins	C _n H _{2n+2}	0.09-0.10			0.615		
Methane	CH ₄		16.04	0.10		-17.88	-12.1
Ethane	CH ₃ CH ₃		30.07	0.20		-20.24	-7.1
Propane	CH ₃ CH ₂ CH ₃		44.09	0.30		-24.82	-5.1
n-Butane	CH ₃ (CH ₂) ₂ CH ₃		58.12	0.40		-29.81	-3.1
n-Pentane	CH ₃ (CH ₂) ₃ CH ₃		72.15	0.50		-35.00	-1.1
19 LPG		0.100			220.664		
20 Chlorine	Cl ₂	0.098	70.91	0.20	20.963	0.00	0.0
21 Borax	Na ₂ B ₄ O ₇ ·10H ₂ O	0.096			1.247		
22 Ammonia	NH ₃	0.085	17.05	0.15	32.400	-10.92	-3.0
23 SR gasoline		0.061			190.875		
23 Methanol	CH ₃ OH	0.079	32.04	0.20	5.980	-45.08	-35.1
24 Kerosene		0.056			2.356		
25 Heating Oil (No. 2)		0.049			5.623		
26 Residual Fuel Oil (No. 6)		0.041			1.735		
27 Natural Gas		0.029-0.047					
28 Sulfuric Acid	H ₂ SO ₄	0.038	98.05	0.30	79.218	-194.45	-164.1
29 Coal		0.014-0.028					

NOTE: * indicates current figure (6-6-86)

() indicated estimated figure

Table 2 Pertinent Intermediate Chemical Data

Chemical	Symbol	Price	MW	SF	Production	Hf(298K) (kcal/mol)	Gf(298K) (kcal/mol)
		(\$/lb) (6-6-86)			(billion lbs/yr) (6-6-86)		
1 Nonylphenol, Ethoxylated		0.490			0.092		
2 Triethanolamine	C3H9N	0.480	59.11	0.60	0.089	-5.70	23.64
3 Isopropylene Oxide	C3H6O	0.475	58.08	0.73	*1.880	-22.17	-6.16
4 Triethylene Glycol	(CH2OCH2CH2O)2	0.470	150.18	1.65	0.088	-191.40 (1)	
5 Monoethanolamine (2-Aminoethanol)	H2NCH2CH2OH	0.460	61.09	0.40	0.083	-65.60 (1)	
6 Acrylonitrile	C3H3N	0.455	53.06	0.75	*2.350	44.20	46.68
7 Linear Alkybenzenes		0.450			0.495		
8 Acetic Anhydride	C4H6O3	0.435	102.09	0.65	1.458	-137.60	-113.93
9 Dinitrotoluene	C7H6N2O4	0.430	182.13	0.60	0.273	-10.20	-
10 Aniline	C6H7N	0.420	93.12	0.50	0.407	20.76	35.84
10 Ethylene Glycol Monobutyl Ether	HOCH2CH2OOC4H9	0.415	118.18	1.13	0.131		
11 Diethanolamine	(C2H5O)2NH	0.410	105.14	0.90	0.086	-17.30	17.23
11 Ethyl Acetate	CH3COOC2H5	0.410	88.12	0.90	0.171	-105.86	-76.25
12 Methyl Chloroform (1,1,1-Trichloroethane)	CH3CCl3	0.405	133.41	0.55	0.455	-33.10	-18.52
13 Dodecyl Phthalate		0.395			0.110		
14 Vinyl Acetate	C4H6O2	0.390	86.09	0.95	*2.110	86.86	-
15 Trichloroethylene	C1Cl=CCl2	0.385	131.29	0.73	0.293	-1.40	4.75
16 Stearic Acid	CH3(CH2)16COOH	0.370	284.50	0.95	0.094	-211.74 (1)	
17 Acetaldehyde	C2H4O	0.370	44.05	0.40	0.900	-35.76	-31.86
18 Carbon Tetra- chloride	CCl4	0.360	153.84	0.58	0.906	-24.00	-13.92
19 Ethylenediamine Tetraacetic Acid (EDTA)		0.360			0.050		
20 Methylene Chloride (Dichloromethane)	CH2Cl2	0.350	84.93	0.33	0.457	-22.80	-16.46
21 Methyl Isobutyl Ketone (4-Methyl-2-Pentanone)	(CH3)2CHCH2COCH3	0.350	100.16	0.98	0.150	-52.50 (1)	
22 Ethylene Oxide	C2H4O	0.350	44.05	0.60	*5.930	-12.58	-3.13
23 Chloroform	CHCl3	0.345	119.39	0.48	0.262	-24.20	-16.38
24 n-Butyl Alcohol	C4H9OH	0.340	74.12	0.50	0.490	-65.59	-36.01
25 Ethylene Glycol Dimethyl Ether	HOCH2CH2OCH3	0.340	76.11	0.83	0.090		
26 Perchloroethylene (Tetrachloroethylene)		0.310			0.679		
27 Ethylene Glycol	HOCH2CH2OH	0.310	62.07	0.40	*4.680	-93.05	-72.77
28 Sorbitol	C6H14O6	0.300	182.17	0.90	0.134	-320.00	-226.00
29 n-Butyraldehyde	C4H8O	0.295	72.10	0.70	1.066	-49.00	-27.43
30 Phenol	C6H6O	0.290	94.11	0.45	*2.780	-23.03	-7.86
31 Vinyl Chloride	H2C=CHCl	0.280	62.50	0.50	*2.780	6.40	12.31
32 Acetone	CH3COCH3	0.270	58.08	0.65	*1.800	-52.00	-36.59
33 Phthalic Anhydride	C8H4O3	0.270	146.12	0.50	0.702	-110.10	-75.12
34 Ethyl Chloride	C2H5Cl	0.265	64.52	0.40	0.575	-26.70	-14.34
35 Methyl Chloride	CH3Cl	0.260	50.49	0.20	0.366	-19.32	-13.72
36 Acetic Acid	CH3COOH	0.250	60.05	0.45	*2.680	-103.93	-90.03
37 Lignosulfonates		0.250			0.945		
38 Methyl Ethyl Ketone (2-Butanone)	CH3CH2COCH3	0.235	72.12	0.75	0.425	-56.97	-34.91
39 Ethylbenzene	C6H5CH2CH3	0.230	106.17	0.55	*8.550	7.12	31.21
40 Styrene	C6H5CH=CH2	0.210	104.16	0.75	*7.450	35.02	51.10
41 Carbon Disulfide	CS2	0.210	76.13	0.30	0.479	27.96	15.99
42 Isopropyl Alcohol	(CH3)2CHOH	0.206	60.11	0.40	*1.120	-65.15	-41.15

NOTE: * indicates current figure (6-6-86)

(1) indicated estimated figure

Table 3 Pertinent Product Data

Chemical	Symbol	Price	MW	SF	Production	Hf(298K) (kcal/mol)	Gf (kcal/mol)
		(\$/lb) (6-6-86)			(billion lbs/yr) (1975)		
1 Acrylamide	$\text{H}_2\text{C}=\text{CHCONH}_2$	1.000	70.07	1.20	0.060	(-32.65)	(=
2 Acetylene	$\text{CH}\equiv\text{CH}$	0.960	26.04	0.60	0.462	54.19	
3 Trichlorotrifluoroethane (Fluorocarbon 113)	Cl_3CCF_3	0.935	187.38	0.95	0.063	-178.20	
4 Glycerin		0.890			0.272		
5 O-Cresol	$2\text{-CH}_3\text{C}_6\text{H}_4\text{OH}$	0.870	108.13	0.61		-30.74	
6 Caprolactam	$\text{C}_6\text{H}_{11}\text{NO}$	0.860	101.15	0.80	0.713		
7 Epichlorohydrin	$\text{C}_3\text{H}_5\text{OCl}$	0.860	92.53	0.95	0.241	92.53	
8 Tetrahydrofuran	$\text{C}_4\text{H}_8\text{O}$	0.860	74.10	0.95	0.088	-44.03	
9 Butadiene	C_4H_6	0.800	54.09	0.80	*2.530	38.77	
10 Monosodium Glutamate		0.800			0.038		
11 Naphthalene	C_{10}H_8	0.770	128.16	0.80	0.380	36.08	
12 Furfural	$\text{C}_5\text{H}_4\text{O}_2$	0.750	96.08	0.95	0.165	-46.60 (1)	
13 Dichlorodifluoromethane (Fluorocarbon 12)	Cl_2CF_2	0.740	120.91	0.56	0.393	-115.00	
14 Pentaerythritol	$\text{C}_5\text{H}_{12}\text{O}_2$	0.720	104.15	0.90	0.101	-219.70	
15 sec-Butyl Acrylate		0.710	128.17		0.380		
16 Butyl Acrylate	$\text{C}_7\text{H}_{10}\text{O}_2$	0.690	126.15	0.75	0.179	-80.10 (1)	
17 Bisphenol A	$\text{C}_{15}\text{H}_{16}\text{O}_2$	0.670	228.28	1.00	0.293	-88.20 (s)	
18 Phosgene	CCl_2O	0.670	98.92	0.90	0.796	-52.80	
19 Acrylic Acid	$\text{C}_3\text{H}_4\text{O}_2$	0.670	72.06	0.90	0.215	-80.36	
20 Ethyl Acrylate	$\text{C}_5\text{H}_8\text{O}_2$	0.660	100.11	1.00	0.240	-88.70 (1)	
21 Allyl Chloride	$\text{C}_3\text{H}_5\text{Cl}$	0.650	76.53	0.55	0.240	-0.15	
22 Diethylene Glycol	$\text{C}_2\text{H}_6\text{O}_2$	0.640	78.13	0.60	0.323	-93.05	
23 Trichlorofluoromethane (Fluorocarbon 11)	Cl_3CF	0.640	137.37	0.58	0.270	-68.00	
24 Dimethylamine	$\text{C}_2\text{H}_7\text{N}$	0.635	45.08	0.50	0.117	-4.50	
25 Hexamethylenetetramine	$\text{C}_6\text{H}_{12}\text{N}_4$	0.630	146.24	0.70	0.062		
26 Methyl Methacrylate	$\text{C}_5\text{H}_8\text{O}_2$	0.620	100.11	0.90	0.546	-88.70 (1)	
27 Adipic Acid	$\text{C}_6\text{H}_{10}\text{O}_4$	0.590	146.14	0.80	*1.300	-236.50 (s)	
28 Cresylic Acid	$\text{CH}_3\text{C}_6\text{H}_4\text{OH}$	0.580	108.13	0.60	0.045	-30.74	
29 Melamine	$\text{C}_3\text{H}_6\text{N}_6$	0.580	128.14	0.80	0.083		
30 Linear Alcohols		0.570			0.506		
31 Cyclohexanone	$\text{C}_6\text{H}_{10}\text{O}$	0.565	98.14	0.80	0.554	-55.00	
32 Chloroacetic Acid	$\text{C}_2\text{H}_3\text{O}_2\text{Cl}$	0.560	94.50	0.80	0.065	-122.60	
33 Ethylene Glycol Monoethyl Ether Acetate		0.555			0.093		
34 Maleic Anhydride	$\text{C}_4\text{H}_2\text{O}_3$	0.530	98.06	0.85	0.216	-112.43	
35 Diacetone Alcohol	$\text{C}_6\text{H}_{12}\text{O}_2$	0.520	106.08	0.80	0.297		
36 Ethylene Glycol Monoethyl Ether	$\text{HOCH}_2\text{CH}_2\text{OCC}_2\text{H}_5$	0.510	90.12	0.90	0.178	-46.10	

NOTE: * indicates current figure (6-6-86)

() indicated estimated figure

Table 4 Pertinent Specialty Chemical Data

Chemical	Symbol	Price (\$/lb)	BB	SF	Production (billion lbs/yr)	Hf(298K) (kcal/mol)	Gf(298K) (kcal/mol)
1 Vitamin E		12.428			0.002		
2 Benzyl Formate	BCOOCB2C6H5	10.500	136.16	1.30		-81.4 (1)	
3 Diacetyl	CB3COCOCB3	9.250	86.09	1.60		-55.4	
4 Dimethyl Benzyl Carbinyl Acetate		6.950					
5 Allyl Bromide	CB2=CBCH2Br	5.500	120.98	1.00		11.8	19.11
6 Cyclamen Aldehyde	C3B7C6H4CB2CBCH3CHO	4.850	190.29	2.50			
7 Ascorbic Acid (Vitamin C)	C6B8O6	4.763	176.14	2.40	0.024	-321.30 (s)	
8 Vanillin	1CB0C6H3-3OCB3-4OH	4.750	152.16	2.81		-127.2 (1)	
9 o-Dianisidine		4.250					
10 Capric Aldehyde	CB3(CB2)8CHO	3.950	156.27	2.20		-89.09	-15.9
11 Allyl Caproate		3.900					
12 Benzophenone	C6B5COC6H5	3.500	182.21	2.00		-4.3 (1)	
13 Benzyl Propionate		3.350					
14 Acetoacet-m-Xylidide		3.330					
15 2,6-Dichloro-4-Nitro- aniline	2,6Cl2-4NO2-C6B2NH2	3.300	207.02	1.70			
16 N,N-Diethyl-m-Toluidine	C6B4-1N(C2H5)2-2CH3	3.180	163.26	1.90			
17 Benzylidene Acetone		2.950					
18 Acetoacet-o-Chloroanilide		2.850					
19 Acetyl Salicylic Acid	C6B4-1COOH-2COCB3	2.800	168.16	2.87	0.025	-190.8 (s)	
20 Diethyl Toluamide	CB3C6H4-3CON(CB2CB3)2	2.750	191.27	2.38			
21 Guaiacol	C6B4-1OH-2OCB3	2.700	124.15	1.90		(14.78)	(21.5)
22 Acetoacet-o-Anisidide		2.700					
23 N-Acetyl-p-Amino Phenol		2.540					
24 o-Anisidine	C6B4-1NH2-2OCB3	2.270	123.16	1.90		-41.1 (s)	
25 Aminoacetic Acid	C6B1ONHCB2CO2C2H5	2.180	179.22	1.50		-158.8 (s)	
26 Hydroxyethyl cellulose		2.070			0.030		
27 Acetyltriethyl citrate		2.060					
28 Dicapryl Phthalate		2.000					
29 Dinitrophenol	C6B3-1OH-2,4(NO2)2	1.950	184.11	1.67		-55.5 (s)	
30 Hydroquinone	1,4-(OH)2C6H4	1.950	110.11	1.20		-66.00	(-33.85)
31 n-Butyl Ether	C4B9OC4B9	1.850	130.23	1.60		-78.8	-21.16
32 N,N-Diethylaniline	C6B5N(C2H5)2	1.830	149.24	1.30		9.6	62.72
33 Camphor	C8B12O	1.800	152.24	0.40			
34 Diethyl Oxalate	C2H5OCCOOC2H5	1.800	146.14	2.20		-189.9 (1)	
35 Resorcinol	1,3-(OH)2C6H4	1.794	110.11	1.20	0.019	(-64.19)	-
36 Cellulose Acetate Butyrate		1.750			0.065		
37 Catechol	1,2-(OH)2C6H4	1.686	110.11	1.10		(-62.39)	-
38 Salicylic Acid	C6B4-1OH-2COOH	1.680	138.12	1.30	0.031	(-99.50)	(-72.82)
39 m-Cresol	C6B4-1OH-3CH3	1.650	108.13	1.10		-31.63	-9.69
40 Methyl Parathion		1.650			0.054		
41 Benzyl Benzoate	C6B5COOCB2C6H5	1.650	212.25	1.60		-73.7 (s)	
42 Diethylenetriamine	HN(CB2CB2NH2)2	1.600	103.17	1.20			
43 Acetoacet-o-Toluidide		1.580					
44 Ethylaniline	C6B4-1NH2-3CH2CB3	1.580	121.18	1.12		13.4	49.58
45 n-Butyl Lactate	CB3CB(OH)COOC4B9	1.580	146.19	1.50			
46 Tricresyl Phosphate, Tritotyl Phosphate	C21B21O4P	1.550	368.37	1.90	0.051		
47 Crotonic Acid	C4B6O2	1.500	89.09			-103.1 (s)	
48 Tetraethylene Glycol Diacrylate		1.500					

Table 4 Pertinent Specialty Chemical Data

Chemical	Symbol	Price (\$/lb)	EW	SF	Production (billion lbs/yr)	Hf(298K) (kcal/mol)	Gf(298K) (kcal/mol)
49 2,4-Xylydine	C6H3-1NH2-2,4(C6H3)2	1.500	121.18	1.30		-19.2 (1)	
50 Itaconic Acid	CH2=C(COOH)CH2COOH	1.450	130.10	1.50		-201.05 (s)	
51 Quinoline	C5H4C4H3N	1.430	129.16	1.35		35.7 (1)	
52 Hexahydrophthalic Anhydride		1.420					
53 Methyl Anthranilate		1.410					
54 Isatoic Anhydride	CH2C4H2O3	1.400	112.09	1.40		-125.1 (s)	
55 Diethyl Carbonate		1.400					
56 Nitroaniline	C6H4-1NH2-2NO2	1.370	138.13	1.10		-3 (s)	
57 o-Phenylphenol	C6H4-1OH-2C6H5	1.350	170.21	1.10			
58 Ethyl Benzoate	C6H5-COOC2H5	1.350	150.18	1.25			
59 Dicyclohexylamine	C6H11-NH2-C6H11	1.350	181.33	0.90			
60 Ethyl Butyrate		1.350					
61 Terpinyl Acetate		1.350					
62 Triethylamine	(CH3CH2)3N	1.330	101.19	1.20		-23.8	26.36
63 Tributylamine	(CH3CH2CH2CH2)3N	1.330	185.36	1.25		-75.6 (1)	
64 Benzyl Alcohol	C6H5CH2OH	1.320	124.15	1.00		(-23.99)	
65 Ethylenediamine	NH2CH2CH2NH2	1.300	60.10	0.90	0.069	-17.30	17.23
66 Cellulose Acetate	[C6H7O2(OCOCH3)3]x	1.300			0.726		
67 2,4-D Butyl Ester		1.300					
68 Acetoacetanilide	C6H5NHCOCH3	1.290	135.17	1.40		-50.30 (s)	
69 Acetyltributyl Citrate		1.280					
70 Benzaldehyde	C6H5CHO	1.250	106.13	1.25		-19.00 (1)	
71 Dicyclohexyl Phthalate	C6H4-1,2(COOC6H11)2	1.250	330.43	1.90			
72 Carboxymethyl Cellulose		1.250			0.063		
73 2,4-Dichlorophenoxy Acetic Acid	C6H3-2,4Cl2-1OCH2COOH	1.250	221.02	1.70	0.080		
74 2,4-Dinitroaniline	C6H3-2,4(NO2)2-1NH2	1.220	183.13	0.98		-22.40 (s)	
75 Benzyl Acetate		1.200					
76 Butyrolactone	C4H6O2	1.200	86.09	1.60		-99.20 (1)	
77 Citric Acid	HOC(CH2COOH)2COOH	1.190	192.14	1.80	0.204	-369.00 (s)	
78 Diethyl Ethanolamine		1.180					
79 Di-Isopropylamine	(1C3H7)2NH	1.170	133.19	1.10		-23.80	26.36
80 tert-Butylamine	NH2C(CH3)3	1.170	73.14	1.40			
81 p-Cresol	p-C7H8O	1.150	108.13	1.70		-32.97	-7.34
82 Dimethyl Ethanol-amine		1.150					
83 Chlorodifluoromethane (Fluorocarbon 22)	ClCHF2	1.140	86.40	1.10	0.132	-119.90	-112.47
84 Dibutylamine	(C4H9)2NH	1.120	129.25	0.95		-41.80	
85 2,4-D acid		1.100					
86 Di-tert-Amylphenol		1.040					
87 N,N-Dimethylaniline	C6H5N(CH3)2	1.030	121.18	0.85		20.20	55.26
88 Butyl Cyclohexyl Phthalate		1.010					
89 Toluene Diisocyanate	C6H3-2,4(N=C=O)2-1CH3	1.010	174.16	1.50	0.479		
90 Acrylamide	CH2=CHCONH2	1.000	71.08	1.80		(-32.85)	(-10.06)
91 Allene	H2C=C=CH2		40.00	1.00		(70.74)	(53.98)

NOTE: * indicates current figure (6-6-86)

() indicated estimated figure

1-4. Examples in ascending order, covering the entire SF-range are shown in Table 4a.

Price can be described as follows

$$P(\$/lb) = a_0 + a_1 \cdot MW + a_2 \cdot SF \quad (1)$$

After rearranging equation (1) can be written as

$$P(\$/lb) = a_0 + a_1 \left(MW + \frac{a_2}{a_1} SF \right) = a_0 + a_1 \cdot X \quad (2)$$

Figure 3 represents price correlated as a function of molecular weight and structure factor revealing a linear relationship with a coefficient of regression of 0.901. This correlation is useful in estimating market value of compounds not included in this report.

It was necessary to obtain thermodynamic data to calculate the heat of formation and ground-state free energy values at desired temperatures. Data available in the literature was first obtained and then the method of Benson (described in the Appendix) was chosen to estimate unknown properties [6-8]. This data was needed to calculate heat of reactions and ground state free energies of candidate reactions found throughout the literature [9-12]. Tables 1-4 present a summary of all pertinent chemical data obtained in the search.

Table 4a Chemical Structure Factors in Ascending Order

Compound	SF	Molecular Structure Comment
1) Methane	0.10	CH_4 , simplest carbon molecule
2) Ammonia	0.15	NH_3 , simplest amine group
3) Chlorine	0.20	Cl-Cl , diatomic Halogen
4) Formaldehyde	0.30	HCHO , simple aldehyde
5) Benzene	0.35	C_6H_6 , simple aromatic
6) Ethylene	0.40	$\text{H}_2\text{C}=\text{CH}_2$, straight chain olefin
7) Acetic Acid	0.45	CH_3COOH , carboxylic acid attached to a methyl group
8) Propylene	0.50	$\text{H}_2\text{C}=\text{CHCH}_3$, straight chain olefin
9) Cumene	0.55	$\text{C}_6\text{H}_5-\text{CH}(\text{CH}_3)_2$, benzene ring with iso-propyl substituent
10) Methyl Chloroform	0.60	$\text{CH}_3\text{C}(\text{Cl})_3$, tetrachlorine configuration on ethane
11) Trichloro-ethylene	0.73	$\text{ClCH}=\text{CCl}_2$, straight chain olefin with 3 halogens, one iso-configuration
12) Ethylene Glycol Monoethyl Ether	0.93	$\text{HOCH}_2\text{CH}_2\text{OC}_2\text{H}_5$, ether group
13) Butadiene	0.80	$\text{CH}_2=\text{CHCH}=\text{CH}_2$, straight chain diene
14) Ethylaniline	1.12	$\text{C}_6\text{H}_4-1\text{NH}_2-3\text{CH}_2\text{CH}_3$, aniline with ethyl group in the meta position
15) Ethyl Benzoate	1.25	$\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$, benzoic ester with ethyl group
16) N,N-Diethyl-aniline	1.31	$\text{C}_6\text{H}_5\text{N}(\text{C}_2\text{H}_5)_2$, aniline with 2 ethyl groups attached to Nitrogen
17) Diacetyl	1.60	$\text{CH}_3\text{COCOCH}_3$, diketone with 2 methyl groups
18) Dinitrophenol	1.67	$\text{C}_6\text{H}_3-1\text{OH}-2,4(\text{NO}_2)_2$, phenol with 2 nitr groups in the ortho and para positions
19) 2,6-Dichloro-4-Nitroaniline	1.70	$2,6\text{Cl}_2-4\text{NO}_2-\text{C}_6\text{H}_2-\text{NH}_2$, aniline with NO_2 group in the para position and 2 Cl groups both in the ortho position
20) Vanillin	2.81	$4\text{CHOC}_6\text{H}_3-2\text{OCOC}_2\text{H}_5-1\text{OH}$, phenol with an aldehyde and an ester group attached in the ortho and para positions

CANDIDATE REACTIONS

Fifteen (15) possible candidate endothermic chemical reactions, listed in Table 5, were chosen and examined thermodynamically, using "ground state" enthalpies and free energies of reaction. More detailed summaries of the thermodynamic values are presented by Tables 7-16 incl. in the section on Reaction Thermodynamics.

The reactions utilize the feedstock compounds of n-hexane, benzene, toluene, methanol, ammonia, water, and hydrogen bromide. The product compounds include benzene, toluene, phenol, aniline, resorcinol, cresols, salicylic acid, acetanilide, benzaldehyde, benzyl alcohol, and others. Cyclization-addition reactions of n-paraffins to aromatics, and n-paraffins plus methanol to alkylated and hydroxylated aromatics, look feasible and produce valuable products. Possible catalysts for these reactions, which are photo-activated, are the transition metal oxides of titanium, vanadium, chromium, and molybdenum, especially V_2O_5 . Other possible catalysts are mentioned in Appendix B.

As a discriminator of the product cost advantage, the product price to reactant(s) cost ratio is proposed, and values are presented in Table 6. In general reactions were sought that would give ratios of 5 to 10. It will be noted that cresols, resorcinol, benzaldehyde, benzyl alcohol, guaiacol, and ethyl aniline fall in this range of values.

As experimental reaction rate data are obtained, this list of favorable reactions (high conversion and selectivity) will be shortened and process possibilities will come into sharper focus.

TABLE 5 - Favorable Synthesis Reactions

Reaction	Products	$\Delta H_R^\circ(298)$	$\Delta H_R^\circ(500)$	Product Value (\$/lb)
1) $n\text{-C}_6\text{H}_{14}$	benzene	59.78	62.02	0.123
2) $n\text{-C}_6\text{H}_{14} + \text{CH}_3\text{OH}$	toluene	42.17	44.95	0.190
3) $n\text{-C}_6\text{H}_{14} + \text{CH}_3\text{OH}$	phenol	47.12	49.98	0.290
4) $n\text{-C}_6\text{H}_{14} + \text{NH}_3$	aniline	71.65	74.90	0.420
5) $n\text{-C}_6\text{H}_{14} + 2\text{CH}_3\text{OH}$	Resorcinol	31.24	36.35	p 1.950 m 1.790 o 1.686
6) $\text{C}_6\text{H}_5\text{CH}_3 + \text{H}_2\text{O}$	o-cresol	15.88	16.67	p 1.150 m 1.650 o 0.870
7) $n\text{C}_6\text{H}_{14} + \text{CH}_3\text{OH}$	m-cresol	56.41	59.97	"
8) $\text{C}_6\text{H}_5\text{OH} + \text{CO}_2$	salicylic acid	17.58	16.04	1.680
9) $\text{C}_6\text{H}_5\text{NH}_2 + \text{CH}_3\text{CHO}$	CH_3CONH - acetanilide	2.72	4.30	1.29
10) $\text{C}_6\text{H}_5\text{CH}_3 + \text{H}_2\text{O}$	benzaldehyde	27.85	29.08	1.25
11) $\text{C}_6\text{H}_6 + \text{CH}_3\text{OH}$	benzyl alcohol	4.27	6.87	1.32
12) $\text{C}_6\text{H}_5\text{CH}_3 + \text{NH}_3$	2,4-xylylidine	19.37	21.18	1.50
13) $\text{C}_6\text{H}_5\text{OH} + \text{CH}_3\text{OH}$	guaiacol	11.91	11.78	2.70
14) $\text{CH}_2=\text{CHCH}_3 + \text{HBr}$	$\text{CH}_2=\text{CHCH}_2\text{Br}$ allyl bromide	15.58	16.29	5.50
15) $n\text{-C}_8\text{H}_{18} + \text{NH}_3$	ethylaniline	74.14	76.51	1.58

TABLE 6 - Price/Cost Ratios and Annual Production

REACTION	REACTANT COST (\$/lb)	PRODUCT	PRODUCT VALUE (\$/lb)	RATIO ⁺	ANNUAL ⁺⁺ PRODUCTION
1a) nC_6H_{14}	0.089	benzene	0.123	1.54	16,960
* 1b) nC_6H_{18}	0.089	ethylbenzene	0.230	2.58	9302
2) $nC_6H_{14} + CH_3OH$	0.168	toluene	0.190	1.13	12,490
3) $nC_6H_{14} + CH_3OH$	0.168	phenol	0.290	1.73	3398
4) $nC_6H_{14} + NH_3$	0.174	aniline	0.420	2.41	1242
5) $nC_6H_{14} + 2CH_3OH$	0.247	resorcinol	1.95 (p) 1.79 (m) 1.69 (o)	7.89 7.25 6.83	31
6) $C_6H_5CH_3 + H_2O$	0.195	o-cresol	1.15 (p) 1.65 (m) 0.87 (o)	5.90 8.46 4.46	350
7) $nC_6H_{14} + CH_3OH$	0.168	m-cresol	1.15 (p) 1.65 (m) 0.87 (o)	6.85 9.82 5.18	
8) $C_6H_5OH + CO_2$	0.340	salicylic acid	1.68	4.94	55
9) $C_6H_5NH_2 + CH_3CHO$	0.790	acetanilide	1.29	1.63	
10) $C_6H_5CH_3 + H_2O$	0.195	benzaldehyde	1.25	6.41	14.5
11) $C_6H_6 + CH_3OH$	0.202	benzyl alcohol	1.32	6.53	7.46
12) $C_6H_5CH_3 + NH_3$	0.275	2,4-xylidine	1.50	5.45	
13) $C_6H_5OH + CH_3OH$	0.369	guaiacol	2.70	7.32	
14) $CH_2=CHCH_3 + HBr$	0.545	$CH_2=CHCH_2Br$ allyl bromide	5.50	10.1	
15) $nC_6H_{18} + NH_3$	0.174	ethylaniline	1.58	9.08	

* added to table

+ Product Price/Reactant(s) Cost

++ 10^6 lbs/year

Notes to Accompany Table 6

REACTANTS	COST
nC_6H_{14}	$\left(\frac{\$21.63}{\text{bbl}}\right)^{(1)} \left(\frac{\text{bbl}}{42\text{gal}}\right) \left(\frac{\text{gal}}{19.28\text{ft}^3}\right)^{(2)} \left(\frac{3.322\text{ft}^3}{\text{lb}}\right)^{(2)} = \$1.089/\text{lb}$
CH_3OH	\$0.079/lb
NH_3	\$0.085/lb
H_2O	\$0.005/lb
$C_6H_5CH_3$	\$0.190/lb
C_6H_5OH	\$0.290/lb
$C_6H_5NH_2$	\$0.420/lb
CH_3CHO	\$0.370/lb
C_6H_6	\$0.123/lb
$CH_2=CHCH_3$	\$0.160/lb
HBr	\$0.385/lb
nC_8H_{18}	\$0.089/lb
CO_2	\$0.050/lb ⁽³⁾

(1) "Statistics", Oil and Gas Journal, October 19, 1987.

(2) "Engineering Data Book", 9th ed., Gas Processors Suppliers Association, pp. 16-3 (1972).

(3) Price quoted from Liquid Carbonics CO₂ Corporation, Houston, TX.

--- All other prices were obtained from "Chemical Prices", Chemical Marketing Reporter, Schnell Publishing Company, Inc., New York, pp. 32-39, June 9, 1986.

REACTION THERMODYNAMICS

In order to determine the temperature effects on the reactions of interest, detailed "ground state" thermodynamic calculations are presented in this section, Tables 7-25. Values of the enthalpy and Gibbs free energy of formation, the ΔC_p° of reaction, and $\ln K_a$, from 298 to 1000K are presented; along with the temperature dependent functions. The Tables are divided into three (3) reaction groups: synthesis, oxidation of aromatics, and oxidation of olefins.

1) Tables 7-12 incl. - synthesis reactions; dehydrocyclization, alkylation, and hydroxylation. It will be noted, that overall the dehydrocyclization plus alkylation and hydroxylation of n-paraffins is endothermic; whereas, the alkylation and hydroxylation of aromatics separately or combined are exothermic. In other words, the chemical conversion of n-paraffins to aromatics being endothermic is a key to the utilization of solar energy during the day time.

2) Tables 13-16 incl. - oxidation reactions of aromatics. Last year (1987) Professor Prengle suggested that to produce a cost effective commercial process a combination of day-time (endothermic) and night-time (exothermic) reactions would be required. Also that certain oxidation reactions leading to single step conversion products of high value, or multistep conversions to heterocyclic products of high value should be used. Amongst the simple oxidation products of aromatics are: aldehydes and acids. An important reaction in this series is the first, Table 13, the partial oxidation of toluene to benzaldehyde; the latter being a high value product.

3) Tables 17-25 incl. - oxidation reactions of olefins. The conversion of n-paraffins to aromatics results also in the production of a significant fraction of olefins (ethylene and propylene), which must be converted to high value products as well. Therefore, partial oxidation reactions for this group of compounds is pertinent. Reaction to aldehydes, epoxides, glycols, alcohol, and ketone are presented along with pertinent thermodynamic values. It will be noted that the formation of aldehydes and epoxides are highly favorable.

The attached tables present numerous reactions, in the above three categories, from which a number of day-time night-time sequences can be selected, for experimental study. The ultimate choice of the specific reactions will depend on the experimental results, and where there are choices, will depend on maximizing the \$-value of the products. In this latter connection Table 26 presents the Price/Cost ratios and annual production for the oxidation products of aromatics and olefins.

TABLE 7 - Dehydrocyclization and Hydroxylation
of n-Heptane to o-Cresol

T(K)	n-C ₇ H ₁₆	CH ₃ OH	ΔH_f° (kcal/gmole)		4H ₂	ΔH_R°
			C ₇ H ₈ O	CH ₄		
298.16	-44.88	-48.08	-30.74	-17.89	0	44.33
600	-51.80	-50.34	-33.94	-19.90	0	48.30
1000	-55.26	-51.93	-35.50	-21.40	0	50.29

T(K)	n-C ₇ H ₁₆	CH ₃ OH	ΔF_f° (kcal/gmole)		4H ₂	ΔF_R°	$\ln K_a$
			C ₇ H ₈ O	CH ₄			
298.16	1.91	-38.84	- 8.86	-12.15	0	15.92	-26.87
600	52.32	-28.52	14.72	- 5.51	0	-14.59	12.27
1000	123.01	-13.46	47.75	4.58	0	-57.22	28.80

T(K)	n-C ₇ H ₁₆	CH ₃ OH	ΔC_p° (cals/gmole ^o K)		4H ₂	ΔC_p°
			C ₇ H ₈ O	CH ₄		
298.16	39.67	10.49	31.15	8.54	27.57	17.10
600	68.33	16.02	52.77	12.55	28.03	9.002
1000	91.20	21.38	68.82	17.40	28.87	2.510

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; \quad < \Delta C_p^\circ > = \frac{17.10 + 9.002}{2} = 13.05$$

$$\Delta H_R^\circ = 13.05T + \Delta H^\circ$$

$$44.33 = \frac{13.05(298)}{1000} + \Delta H^\circ ; \quad \Delta H^\circ = 40.44$$

$$\Delta H_R^\circ = 1.305 \times 10^{-2}T + 40.44$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$15.92 = I(298) + 40.44 - (1.305 \times 10^{-2})(298)(\ln 298) ; \quad I = -0.0079$$

$$\Delta F_R^\circ = 40.44 - 0.0079T - (1.305 \times 10^{-2})T \ln T$$

TABLE 8 - Dehydrocyclization and Simultaneous Hydroxylation and Alkylation of n-Hexane to o-Cresol

ΔH_f° (kcal/gmole)

T(K)	n-C ₆ H ₁₄	CH ₃ OH	C ₇ H ₈ O	5H ₂	ΔH_R°
298.16	-39.96	-48.08	-30.74	0	57.30
600	-46.10	-50.34	-33.94	0	62.50
1000	-49.25	-51.93	-35.50	0	65.68

ΔF_f° (kcal/gmole)

T(K)	n-C ₆ H ₁₄	CH ₃ OH	C ₇ H ₈ O	5H ₂	ΔF_R°	$\ln K_a$
298.16	- 0.06	-38.84	- 8.86	0	30.04	-50.71
600	43.02	-28.52	14.72	0	0.220	- 0.1845
1000	103.57	-13.46	47.75	0	-42.36	21.32

ΔC_p° (cal/gmole^oK)

T(K)	n-C ₆ H ₁₄	CH ₃ OH	C ₇ H ₈ O	5H ₂	ΔC_p°
298.16	34.20	10.49	31.15	34.46	20.92
600	58.99	16.02	52.77	35.04	12.80
1000	78.89	21.38	68.82	36.09	4.640

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{20.92 + 12.80}{2} = 16.86$$

$$\Delta H_R^\circ = 16.86T + \Delta H^\circ$$

$$57.30 = \frac{16.86(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = 52.28$$

$$\Delta H_R^\circ = 1.686 \times 10^{-2}T + 52.28$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$30.04 = I(298) + 52.28 - (16.86 \times 10^{-3})(298)(\ln 298) ; I = 0.0214$$

$$\Delta F_R^\circ = 52.28 + 0.0214T - (1.686 \times 10^{-2})T \ln T$$

TABLE 9 - Alkylation of Benzene to Toluene

$$\Delta H_f^\circ \text{ (kcal/gmole)}$$

T(K)	C ₆ H ₆	CH ₃ OH	H ₅ C ₆ -CH ₃	H ₂ O	ΔH_R°
298.16	19.82	-48.08	11.95	-57.80	-17.59
600	16.71	-50.34	8.02	-58.50	-16.85
1000	14.82	-51.93	6.01	-59.24	-16.12

$$\Delta F_f^\circ \text{ (kcal/gmole)}$$

T(K)	C ₆ H ₆	CH ₃ OH	H ₅ C ₆ -CH ₃	H ₂ O	ΔF_R°	ln K _a
298.16	30.99	-38.84	29.16	-54.64	-17.63	29.76
600	43.66	-28.52	48.32	-51.16	-17.98	15.08
1000	62.27	-13.46	75.91	-46.04	-18.94	9.53

$$\Delta C_p^\circ \text{ (cal/gmole}^\circ\text{K)}$$

T(K)	C ₆ H ₆	CH ₃ OH	H ₅ C ₆ -CH ₃	H ₂ O	ΔC_p°
298.16	19.52	10.49	24.77	8.03	2.79
600	37.74	16.02	47.20	8.68	2.12
1000	50.16	21.38	63.32	9.85	1.63

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + k ; < \Delta C_p^\circ > = \frac{2.79 + 2.12}{2} = 2.455$$

$$\Delta H_R^\circ = 2.455T + \Delta H_0^\circ$$

$$-17.59 = \frac{2.455(298)}{1000} + \Delta H_0^\circ ; \Delta H_0^\circ = -18.322$$

$$\Delta H_R^\circ = 2.455 \times 10^{-3}T - 18.322$$

$$\Delta F_R^\circ = IT + \Delta H_0^\circ - < \Delta C_p^\circ > T \ln T$$

$$-17.63 = I(298) - 18.322 - (2.455 \times 10^{-3})(298)(\ln 298) ; I = 0.0163$$

$$\Delta F_R^\circ = -18.322 + 0.0163T - (2.455 \times 10^{-3})T \ln T$$

TABLE 10 - Hydroxylation of Benzene to Phenol

ΔH_f° (kcal/gmole)					
T(K)	C ₆ H ₆	CH ₃ OH	H ₅ C ₆ -OH	CH ₄	ΔH_R°
298.16	19.82	-48.08	-23.03	-17.89	-12.66
600	16.71	-50.34	-25.53	-19.90	-11.80
1000	14.82	-51.93	-26.80	-21.40	-11.09

ΔF_f° (kcal/gmole)						$\ln K_a = \frac{-\Delta F_R^\circ}{RT}$
T(K)	C ₆ H ₆	CH ₃ OH	H ₅ C ₆ -OH	CH ₄	ΔF_R°	
298.16	30.99	-38.84	-7.86	-12.15	-12.16	20.53
600	43.66	-28.52	8.62	-5.51	-12.03	10.09
1000	62.27	-13.46	31.84	4.58	-12.39	6.24

ΔC_p° (cal/gmole°K)					
T(K)	C ₆ H ₆	CH ₃ OH	H ₅ C ₆ -OH	CH ₄	ΔC_p°
298.16	19.52	10.49	24.75	8.54	3.28
600	37.74	16.02	43.54	12.55	2.33
1000	50.16	21.38	55.49	17.40	1.35

$$\Delta H_R^\circ = \int \Delta C_p^\circ dT + k = \langle \Delta C_p^\circ \rangle T + k$$

$$\langle \Delta C_p^\circ \rangle = \frac{3.28 + 2.33}{2} = 2.805$$

$$-12.66 = \frac{2.805(298.16)}{1000} + \Delta H_R^\circ ; \Delta H_R^\circ = -13.496$$

$$\Delta H_R^\circ = 2.805 \times 10^{-3} T - 13.496$$

$$\Delta F_R^\circ = \Delta H_R^\circ - \langle \Delta C_p^\circ \rangle T \ln T$$

$$-12.16 = I(298) - 13.496 - (2.805 \times 10^{-3})(298)(\ln 298) ; I = 0.0205$$

$$\Delta F_R^\circ = -13.496 + 0.0205T - 2.805 \times 10^{-3} T \ln T$$

TABLE 11 - Hydroxylation of Toluene to o-Cresol

ΔH_f° (kcal/gmole)					
T(K)	C ₇ H ₈	CH ₃ OH	C ₇ H ₈ O	CH ₄	ΔH_R°
298.16	11.95	-48.08	-30.74	-17.89	-12.50
600	8.02	-50.34	-33.94	-19.90	-11.52
1000	6.01	-51.93	-35.50	-21.40	-10.98

ΔF_f° (kcal/gmole)						
T(K)	C ₇ H ₈	CH ₃ OH	C ₇ H ₈ O	CH ₄	ΔF_R°	$\ln K_a$
298.16	29.16	-38.84	- 8.86	-12.15	-11.33	19.12
600	48.32	-28.52	14.72	- 5.51	-10.59	8.88
1000	75.91	-13.46	47.75	4.58	-10.12	5.09

ΔC_p° (cal/gmole°K)					
T(K)	C ₇ H ₈	CH ₃ OH	C ₇ H ₈ O	CH ₄	ΔC_p°
298.16	24.77	10.49	31.15	8.54	4.43
600	47.20	16.02	52.77	12.55	2.10
1000	63.32	21.38	68.82	17.40	1.52

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + H^\circ ; < \Delta C_p^\circ > = \frac{4.43 + 2.10}{2} = 3.265$$

$$\Delta H_R^\circ = 3.265T + \Delta H^\circ$$

$$-12.50 = \frac{3.265(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -13.473$$

$$\Delta H_R^\circ = 3.265 \times 10^{-3}T - 13.473$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-11.33 = I(298) - 13.473 - (3.265 \times 10^{-3})(298)(\ln 298) ; I = 0.0258$$

$$\Delta F_R^\circ = -13.473 + 0.0258T - (3.265 \times 10^{-3})T \ln T$$

TABLE 12 - Simultaneous Alkylation and Hydroxylation
of Benzene to o-Cresol

ΔH_f° (kcal/gmole)

T(K)	C ₆ H ₆	CH ₃ OH	C ₇ H ₈ O	H ₂	ΔH_R°
298.16	19.82	-48.08	-30.74	0	-2.48
600	16.71	-50.34	-33.94	0	-0.31
1000	14.82	-51.93	-35.50	0	1.61

ΔF_f° (kcal/gmole)

T(K)	C ₆ H ₆	CH ₃ OH	C ₇ H ₈ O	H ₂	ΔF_R°	$\ln K_a$
298.16	30.99	-38.84	- 8.86	0	-1.01	1.70
600	43.66	-28.52	14.72	0	-0.42	0.352
1000	62.27	-13.46	47.75	0	-1.06	0.534

ΔC_p° (cals/gmole^oK)

T(K)	C ₆ H ₆	CH ₃ OH	C ₇ H ₈ O	H ₂	ΔC_p°
298.16	19.52	10.49	31.15	6.892	8.032
600	37.74	16.02	52.77	7.008	6.018
1000	50.16	21.38	68.82	7.217	4.497

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{8.032 + 6.018}{2} = 7.025$$

$$\Delta H_R^\circ = 7.025T + \Delta H^\circ$$

$$-2.48 = \frac{7.025(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -4.5735$$

$$\Delta H_R^\circ = 7.025 \times 10^{-3}T - 4.5735$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-1.01 = I(298) - 4.5735 - (7.025 \times 10^{-3})(298)(\ln 298) ; I = 0.0520$$

$$\Delta F_R^\circ = -4.5735 + 0.0520T - (7.025 \times 10^{-3})T \ln T$$

TABLE 13 - Partial Oxidation of Toluene to Benzaldehyde

T(K)	ΔH_f° (kcal/gmole)				
	C_7H_8	O_2	$\text{C}_7\text{H}_6\text{O}$	H_2O	ΔH_R°
298	11.95	0	8.80	-57.80	-60.95
600	8.02	0	11.38	-58.50	-55.14
1000	6.01	0	12.60	-59.24	-52.65

T(K)	ΔF_f° (kcal/gmole)					$\ln K_a$
	C_7H_8	O_2	$\text{C}_7\text{H}_6\text{O}$	H_2O	ΔF_R°	
298	29.16	0	5.35	-54.64	-78.45	132.4
600	48.32	0	20.84	-51.16	-78.64	65.96
1000	75.91	0	42.73	-46.04	-79.22	39.87

T(K)	ΔC_p° (cals/gmole $^\circ\text{K}$)				
	C_7H_8	O_2	$\text{C}_7\text{H}_6\text{O}$	H_2O	ΔC_p°
298	24.77	7.017	26.70	8.03	2.94
600	47.20	7.670	47.47	8.68	1.28
1000	63.32	8.335	60.95	9.85	-0.855

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; \quad < \Delta C_p^\circ > = \frac{2.94 + 1.28}{2} = 2.11$$

$$\Delta H_R^\circ = 2.11T + \Delta H^\circ$$

$$-60.95 = \frac{2.11(298)}{1000} + \Delta H^\circ ; \quad \Delta H^\circ = -61.58$$

$$\Delta H_R^\circ = 2.11 \times 10^{-3}T - 61.58$$

$$\Delta F_R^\circ = 1T + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-78.45 = 1(298) - 61.58 - (2.11 \times 10^{-3})(298)(\ln 298) ; \quad 1 = -0.0446$$

$$\Delta F_R^\circ = -61.58 - 0.0446T - (2.11 \times 10^{-3})T \ln T$$

TABLE 14 - Oxidation of p-Xylene to Terephthalic Acid

ΔH_f° (kcal/gmole)					
T(K)	C_8H_{10}	2O_2	$\text{C}_8\text{H}_6\text{O}_4$	2H_2	ΔH_R°
298	4.29	0	-195.1	0	-199.4
500	0.68	0	-195.7	0	-186.7
1000	-3.61	0	-204.9	0	-159.4

ΔF_f° (kcal/gmole)						
T(K)	C_8H_{10}	2O_2	$\text{C}_8\text{H}_6\text{O}_4$	2H_2	ΔF_R°	$\ln K_a$
298	28.95	0	-149.6	0	-178.5	301.5
500	46.73	0	-116.1	0	-162.8	163.8
1000	94.90	0	- 26.8	0	-121.7	61.2

ΔC_p° (cal/gmole $^\circ\text{K}$)					
T(K)	C_8H_{10}	2O_2	$\text{C}_8\text{H}_6\text{O}_4$	2H_2	ΔC_p°
298	30.32	14.03	26.91	13.78	-3.66
500	48.06	14.86	44.74	13.99	-4.19
1000	74.02	16.67	69.36	14.43	-6.90

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{-3.66 - 4.19}{2} = -3.93$$

$$\Delta H_R^\circ = -3.93T + \Delta H^\circ$$

$$-199.4 = \frac{-3.93(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -198.2$$

$$\Delta H_R^\circ = -3.93 \times 10^{-3}T - 198.2$$

$$\Delta F_R^\circ = RT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-178.5 = R(298) - 198.2 - (-3.93 \times 10^{-3})(298)(\ln 298) ; R = 0.0437$$

$$\Delta F_R^\circ = -198.2 + 0.0437T + (3.93 \times 10^{-3})T \ln T$$

TABLE 15 - Carboxylation of Phenol to Salicylic Acid

T(K)	ΔH_f° (kcal/gmole)			
	$\text{C}_6\text{H}_6\text{O}$	CO_2	$\text{C}_7\text{H}_6\text{O}_3$	ΔH_R°
298	-23.03	-94.05	-139.9	-22.37
500	-24.90	-94.09	-142.8	-23.81
1000	-26.80	-94.32	-145.1	-23.98

T(K)	ΔF_f° (kcal/gmole)				$\ln K_a$
	$\text{C}_6\text{H}_6\text{O}$	CO_2	$\text{C}_7\text{H}_6\text{O}_3$	ΔF_R°	
298	-7.86	-94.26	-99.93	2.19	-3.70
500	2.99	-94.39	-72.21	19.19	-19.32
1000	31.84	-94.61	-4.69	121.76	-61.28

T(K)	ΔC_p° (cal/gmole $^\circ\text{K}$)			
	$\text{C}_6\text{H}_6\text{O}$	CO_2	$\text{C}_7\text{H}_6\text{O}_3$	ΔC_p°
298	24.75	8.87	24.22	-9.40
500	38.64	10.66	40.77	-8.53
1000	55.49	12.97	62.20	-6.26

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{-9.40 - 8.53}{2} = -8.97$$

$$\Delta H_R^\circ = -8.97T + \Delta H^\circ$$

$$-22.37 = \frac{-8.97(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -19.70$$

$$\Delta H_R^\circ = -8.97 \times 10^{-3}T - 19.70$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$2.19 = I(298) - 19.70 + (8.97 \times 10^{-3})(298)(\ln 298) ; I = 0.0224$$

$$\Delta F_R^\circ = -19.70 + 0.0224T + (8.97 \times 10^{-3})T \ln T$$

TABLE 16 - Oxidation of Benzene to Maleic Anhydride

$$\Delta H_f^\circ \text{ (kcal/gmole)}$$

T(K)	C ₆ H ₆	$\frac{9}{2} \text{O}_2$	C ₄ H ₂ O ₃	2H ₂ O	2CO ₂	ΔH_R°
298	19.82	0	-112.4	2(-57.80)	2(-94.05)	-435.9
600	16.71	0	-121.0	2(-58.50)	2(-94.12)	-443.0
1000	14.82	0	-132.4	2(-59.24)	2(-94.32)	-454.3

$$\Delta F_f^\circ \text{ (kcal/gmole)}$$

T(K)	C ₆ H ₆	$\frac{9}{2} \text{O}_2$	C ₄ H ₂ O ₃	2H ₂ O	2CO ₂	ΔF_R°	$\ln K_a$
298	30.99	0	-115.3	2(-54.64)	2(-94.26)	-444.1	750.0
600	43.66	0	-135.3	2(-51.16)	2(-94.45)	-470.2	394.4
1000	62.27	0	-161.9	2(-46.04)	2(-94.61)	-505.5	254.4

$$\Delta C_p^\circ \text{ (cal/gmole}^\circ\text{K)}$$

T(K)	C ₆ H ₆	$\frac{9}{2} \text{O}_2$	C ₄ H ₂ O ₃	2H ₂ O	2CO ₂	ΔC_p°
298	19.52	$\frac{9}{2}(7.017)$	17.35	2(8.03)	2(8.87)	0.0535
600	37.74	"(7.670)	30.53	2(8.68)	2(11.31)	-1.745
1000	50.16	"(8.335)	39.49	2(9.85)	2(12.97)	-2.538

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{0.0535 - 1.745}{2} = -0.8458$$

$$\Delta H_R^\circ = -0.8458T + \Delta H^\circ$$

$$-435.9 = \frac{-0.8458(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -435.6$$

$$\Delta H_R^\circ = -8.458 \times 10^{-4}T - 435.6$$

$$\Delta F_R^\circ = RT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-444.1 = I(298) - 435.6 - (-8.458 \times 10^{-4})(298)(\ln 298) ; I = -0.0333$$

$$\Delta F_R^\circ = -435.6 - 0.0333T + (8.458 \times 10^{-4})T \ln T$$

TABLE 17 - Direct Oxidation of Ethylene to Form Acetaldehyde

T(K)	ΔH_f° (kcal/gmole)			
	C_2H_4	$\frac{1}{2} \text{O}_2$	CH_3CHO	ΔH_R°
298	12.50	0	-39.76	-52.26
600	10.60	0	-41.86	-52.46
1000	9.21	0	-43.27	-52.48

T(K)	ΔF_f° (kcal/gmole)				$\ln K_a$
	C_2H_4	$\frac{1}{2} \text{O}_2$	CH_3CHO	ΔF_R°	
298	16.28	0	-31.86	-48.14	81.30
600	20.92	0	-22.97	-43.89	36.81
1000	28.25	0	-9.91	-38.16	19.20

T(K)	ΔC_p° (cal/gmole $^\circ\text{K}$)			
	C_2H_4	$\frac{1}{2} \text{O}_2$	CH_3CHO	ΔC_p°
298	10.41	3.508	13.06	-0.8580
600	17.10	3.835	20.52	-0.4150
1000	22.57	4.168	26.96	0.2220

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{-0.8580 - 0.4150}{2} = -0.6365$$

$$-52.26 = \frac{-0.6365(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -52.07$$

$$\Delta H_R^\circ = -6.365 \times 10^{-4}T - 52.07$$

$$\Delta F_R^\circ = RT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-48.14 = I(298) - 52.07 + (6.365 \times 10^{-4})(298)(\ln 298) ; I = 0.009562$$

$$\Delta F_R^\circ = -52.07 + 0.009562T + (6.365 \times 10^{-4})T \ln T$$

TABLE 18 - Oxidation of Ethylene to Form Ethylene Oxide

ΔH_f° (kcal/gmole)

T(K)	C ₂ H ₄	$\frac{1}{2} \text{O}_2$	C ₂ H ₄ O	ΔH_R°
298	12.50	0	-12.58	-25.08
600	10.60	0	-14.85	-25.45
1000	9.21	0	-16.12	-25.33

ΔF_f° (kcal/gmole)

T(K)	C ₂ H ₄	$\frac{1}{2} \text{O}_2$	C ₂ H ₄ O	ΔF_R°	$\ln K_a$
298	16.28	0	- 3.13	-19.41	32.78
600	20.92	0	7.44	-13.48	11.31
1000	28.25	0	22.74	- 5.51	2.773

ΔC_p° (cals/gmole°K)

T(K)	C ₂ H ₄	$\frac{1}{2} \text{O}_2$	C ₂ H ₄ O	ΔC_p°
298	10.41	3.508	11.54	-2.378
600	17.10	3.835	20.62	-0.3150
1000	22.57	4.168	27.47	0.7320

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H_R^\circ ; < \Delta C_p^\circ > = \frac{-2.378 - 0.3150}{2} = -1.3465$$

$$-25.08 = \frac{-1.3465(298)}{1000} + \Delta H_R^\circ ; \Delta H_R^\circ = -24.68$$

$$\Delta H_R^\circ = -1.3465 \times 10^{-3} T - 24.68$$

$$\Delta F_R^\circ = IT + \Delta H_R^\circ - < \Delta C_p^\circ > T \ln T$$

$$-19.41 = I(298) - 24.68 + (1.3465 \times 10^{-3})(298)(\ln 298) ; I = 0.0100$$

$$\Delta F_R^\circ = -24.68 + 0.0100T + (1.3465 \times 10^{-3})T \ln T$$

TABLE 19 - Direct Oxidation of Propylene to Propylene Oxide

T(K)	C ₃ H ₆	ΔH_f° (kcal/gmole)			ΔH_R°
		$\frac{1}{2} \text{O}_2$	C ₃ H ₆ O		
298	4.88	0	-22.17		-27.05
600	1.98	0	-25.15		-27.13
1000	0.04	0	-26.72		-26.76

T(K)	C ₃ H ₆	ΔF_f° (kcal/gmole)			ΔF_R°	$\ln K_a$
		$\frac{1}{2} \text{O}_2$	C ₃ H ₆ O			
298	14.99	0	- 6.16	-21.15		35.72
600	26.46	0	11.36	-15.10		12.67
1000	43.43	0	36.26	- 7.17		3.608

T(K)	C ₃ H ₆	ΔC_p° (cal/gmole°K)			ΔC_p°
		$\frac{1}{2} \text{O}_2$	C ₃ H ₆ O		
298	15.27	3.508	17.29	-1.488	
600	25.70	3.835	30.07	0.535	
1000	34.46	4.168	39.79	1.162	

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{-1.488 + 0.535}{2} = -0.4765$$

$$-27.05 = \frac{-0.4765(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -26.91$$

$$\Delta H_R^\circ = -4.765 \times 10^{-4} T - 26.91$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-21.15 = I(298) - 26.91 + (4.765 \times 10^{-4})(298)(\ln 298) ; I = 0.0166$$

$$\Delta F_R^\circ = -26.91 + 0.0166T + (4.765 \times 10^{-4})T \ln T$$

TABLE 20 - Catalytic Vapor Phase Oxidation of
Propylene to Acrolein

$$\Delta H_f^\circ \text{ (kcal/gmole)}$$

T(K)	C ₃ H ₆	O ₂	C ₃ H ₄ O	H ₂ O	ΔH_R°
298	4.88	0	-17.79	-57.80	-80.47
600	1.98	0	-19.57	-58.50	-80.05
1000	0.04	0	-20.54	-59.24	-79.82

$$\Delta F_f^\circ \text{ (kcal/gmole)}$$

T(K)	C ₃ H ₆	O ₂	C ₃ H ₄ O	H ₂ O	ΔF_R°	$\ln K_a$
298	14.99	0	-12.86	-54.64	-82.49	139.3
600	26.46	0	-0.500	-51.16	-78.12	65.53
1000	43.43	0	0.660	-46.04	-88.81	44.70

$$\Delta C_p^\circ \text{ (cal/gmole}^\circ\text{K)}$$

T(K)	C ₃ H ₆	O ₂	C ₃ H ₄ O	H ₂ O	ΔC_p°
298	15.27	7.017	16.07	8.03	1.813
600	25.70	7.670	25.59	8.68	0.9000
1000	34.46	8.335	33.61	9.85	0.6650

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H_R^\circ ; < \Delta C_p^\circ > = \frac{1.813 + 0.9000}{2} = 1.356$$

$$-80.47 = \frac{1.356(298)}{1000} + \Delta H_R^\circ ; \Delta H_R^\circ = -80.87$$

$$\Delta H_R^\circ = 1.356 \times 10^{-3} T - 80.87$$

$$\Delta F_R^\circ = IT + \Delta H_R^\circ - < \Delta C_p^\circ > T \ln T$$

$$-82.49 = I(298) - 80.87 - (1.356 \times 10^{-3})(298)(\ln 298) ; I = 0.0023$$

$$\Delta F_R^\circ = -80.87 + 0.0023T - (1.356 \times 10^{-3}) T \ln T$$

TABLE 21 - Hydration of Ethylene Oxide to Form Ethylene Glycol

ΔH_f° (kcal/gmole)				
T(K)	C ₂ H ₄ O	H ₂ O	C ₂ H ₆ O ₂	ΔH_R°
298	-12.58	-57.80	-93.05	-22.67
600	-14.85	-58.50	-94.90	-21.55
1000	-16.12	-59.24	-95.76	-20.40

ΔF_f° (kcal/gmole)					
T(K)	C ₂ H ₄ O	H ₂ O	C ₂ H ₆ O ₂	ΔF_R°	$\ln K_a$
298	- 3.13	-54.64	-72.77	-15.00	25.33
600	7.44	-51.16	-51.40	- 7.68	6.442
1000	22.74	-46.04	-22.09	1.21	- 0.6090

ΔC_p° (cal/gmole°K)				
T(K)	C ₂ H ₄ O	H ₂ O	C ₂ H ₆ O ₂	ΔC_p°
298	11.54	8.03	23.20	3.630
600	20.62	8.68	32.72	3.420
1000	27.47	9.85	39.88	2.560

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H_R^\circ ; < \Delta C_p^\circ > = \frac{3.630 + 3.420}{2} = 3.525$$

$$-22.67 = \frac{3.525(298)}{1000} + \Delta H_R^\circ ; \Delta H_R^\circ = -23.72$$

$$\Delta H_R^\circ = 3.525 \times 10^{-3}T - 23.72$$

$$\Delta F_R^\circ = IT + \Delta H_R^\circ - < \Delta C_p^\circ > T \ln T$$

$$-15.00 = I(298) - 23.72 - (3.525 \times 10^{-3})(298)(\ln 298) ; I = 0.0493$$

$$\Delta F_R^\circ = -23.72 + 0.0493T - (3.525 \times 10^{-3})T \ln T$$

TABLE 22 - Reaction of Propylene Oxide with Water to Form Propylene Glycol

ΔH_f° (kcal/gmole)				
T(K)	H ₂ O	C ₃ H ₆ O	C ₃ H ₈ O ₂	ΔH_R°
298	-57.80	-22.17	-119.3	-39.33
600	-58.50	-25.15	-135.5	-51.85
1000	-59.24	-26.72	-142.2	-56.24

ΔF_f° (kcal/gmole)					
T(K)	H ₂ O	C ₃ H ₆ O	C ₃ H ₈ O ₂	ΔF_R°	$\ln K_a$
298	-54.64	-6.16	-145.3	-84.50	142.7
600	-51.16	11.36	-170.1	-130.3	109.3
1000	-46.04	36.26	-200.7	-190.9	96.06

ΔC_p° (cal/gmole°K)				
T(K)	H ₂ O	C ₃ H ₆ O	C ₃ H ₈ O ₂	ΔC_p°
298	8.03	17.29	24.37	-0.9500
600	8.68	30.07	39.49	0.7400
1000	9.85	39.79	50.92	1.280

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{-0.9500 + 0.7400}{2} = -0.1050$$

$$-39.33 = \frac{-0.1050(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -39.30$$

$$\Delta H_R^\circ = -1.050 \times 10^{-4} T - 39.30$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-84.50 = I(298) - 39.30 + (1.050 \times 10^{-4})(298)(\ln 298) ; I = -0.1523$$

$$\Delta F_R^\circ = -39.30 - 0.1523T + (1.050 \times 10^{-4})T \ln T$$

TABLE 23 - Reaction of Acrolein with Methanol
to Yield Methoxypropionaldehyde

T(K)	ΔH_f° (kcal/gmole)			
	$\text{C}_3\text{H}_4\text{O}$	CH_3OH	$\text{C}_4\text{H}_8\text{O}_2$	ΔH_R°
298	-17.79	-48.08	-114.1	-48.23
600	-19.57	-50.34	-116.3	-46.36
1000	-20.54	-51.93	-116.4	-43.88

T(K)	ΔF_f° (kcal/gmole)				$\ln K_a$
	$\text{C}_3\text{H}_4\text{O}$	CH_3OH	$\text{C}_4\text{H}_8\text{O}_2$	ΔF_R°	
298	-12.86	-38.84	-150.6	-98.90	167.0
600	-0.5000	-28.52	-199.0	-170.0	142.6
1000	0.6600	-13.46	-234.2	-221.4	111.4

T(K)	ΔC_p° (cal/gmole $^\circ\text{K}$)			
	$\text{C}_3\text{H}_4\text{O}$	CH_3OH	$\text{C}_4\text{H}_8\text{O}_2$	ΔC_p°
298	16.07	10.49	32.14	5.580
600	25.59	16.02	48.43	6.820
1000	33.61	21.38	62.71	7.720

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{6.820 + 5.580}{2} = 6.200$$

$$-48.23 = \frac{6.200(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -50.08$$

$$\Delta H_R^\circ = 6.200 \times 10^{-3}T - 50.08$$

$$\Delta F_R^\circ = RT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-98.90 = I(298) - 50.08 - (6.200 \times 10^{-3})(298)(\ln 298) ; I = -0.128$$

$$\Delta F_R^\circ = -50.08 - 0.1285T - (6.200 \times 10^{-3})T \ln T$$

TABLE 24 - Reaction of Acrolein with Isopropanol to Yield Allyl Alcohol and Acetone

T(K)	ΔH_f° (kcal/gmole)				ΔH_R°
	$\text{C}_3\text{H}_4\text{O}$	$\text{C}_3\text{H}_8\text{O}$	$\text{C}_3\text{H}_6\text{O}$	$\text{C}_3\text{H}_6\text{O}$	
298	-17.79	-65.15	-31.55	-52.00	-0.610
600	-19.57	-68.74	-34.39	-55.07	-1.150
1000	-22.54	-70.58	-36.11	-56.93	-1.920

T(K)	ΔF_f° (kcal/gmole)				ΔF_R°	$\ln K_a$
	$\text{C}_3\text{H}_4\text{O}$	$\text{C}_3\text{H}_8\text{O}$	$\text{C}_3\text{H}_6\text{O}$	$\text{C}_3\text{H}_6\text{O}$		
298	-12.86	-41.49	-17.03	-36.58	0.7400	-1.250
600	-0.5000	-15.94	-1.09	-19.65	-4.300	3.607
1000	0.6600	19.93	21.70	4.61	5.720	-2.879

T(K)	ΔC_p° (cal/gmole $^\circ\text{K}$)				ΔC_p°
	$\text{C}_3\text{H}_4\text{O}$	$\text{C}_3\text{H}_8\text{O}$	$\text{C}_3\text{H}_6\text{O}$	$\text{C}_3\text{H}_6\text{O}$	
298	16.07	21.21	18.17	17.90	-1.210
600	25.59	35.76	30.11	29.34	-1.900
1000	33.61	46.82	39.06	39.15	-2.220

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; \Delta C_p^\circ = \frac{-1.210 - 1.900}{2} = -1.555$$

$$-0.610 = \frac{-1.555(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -0.1466$$

$$\Delta H_R^\circ = -1.555 \times 10^{-3}T - 0.1466$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$0.7400 = I(298) - 0.1466 + (1.555 \times 10^{-3})(298)(\ln 298) ; I = -0.0059$$

$$\Delta F_R^\circ = -0.1466 - 0.0059T + (1.555 \times 10^{-3})T \ln T$$

TABLE 25 - Oxidation of Allyl Alcohol to Glycerin with Hydrogen Peroxide

T(K)	ΔH_f° (kcal/gmole)			ΔH_R°
	$\text{C}_3\text{H}_6\text{O}$	H_2O_2	$\text{C}_3\text{H}_8\text{O}_3$	
298	-31.55	-32.53	-159.8	-95.72
600	-34.39	-33.22	-182.6	-115.0
1000	-36.11	-33.57	-184.2	-114.5

T(K)	ΔF_f° (kcal/gmole)			ΔF_R°	$\ln K_a$
	$\text{C}_3\text{H}_6\text{O}$	H_2O_2	$\text{C}_3\text{H}_8\text{O}_3$		
298	-17.03	-25.21	-114.02	-71.78	121.2
600	- 1.09	-17.47	-66.84	-48.28	40.50
1000	21.70	- 6.85	-3.730	-18.58	9.351

T(K)	ΔC_p° (cals/gmole $^\circ\text{K}$)			ΔC_p°
	$\text{C}_3\text{H}_6\text{O}$	H_2O_2	$\text{C}_3\text{H}_8\text{O}_3$	
298	18.17	10.31	40.92	12.44
600	30.11	13.31	43.35	- 0.07
1000	39.06	15.02	55.06	0.98

$$\Delta H_R^\circ = < \Delta C_p^\circ > T + \Delta H^\circ ; < \Delta C_p^\circ > = \frac{12.44 - 0.07}{2} = 6.185$$

$$-95.72 = \frac{6.185(298)}{1000} + \Delta H^\circ ; \Delta H^\circ = -97.56$$

$$\Delta H_R^\circ = 6.185 \times 10^{-3}T - 97.56$$

$$\Delta F_R^\circ = IT + \Delta H^\circ - < \Delta C_p^\circ > T \ln T$$

$$-71.78 = I(298) - 97.56 - (6.185 \times 10^{-3})(298)(\ln 298) ; I = 0.1217$$

$$\Delta F_R^\circ = -97.56 + 0.1217T - (6.185 \times 10^{-3})T \ln T$$

TABLE 26 - Price/Cost Ratios, Oxidation Products

REACTION	REACTANT COST(\$/lb)		PRODUCT(S)	PRODUCT VALUE(\$/lb)	RATIO ⁺	ANNUAL ⁺⁺ PRODUCTION
1) $\text{H}_5\text{C}_6\text{-CH}_3 + \text{O}_2$	0.190		benzaldehyde	1.25	6.58	14.5
2) $\text{H}_4\text{C}_6\text{-(CH}_3)_2 + 2\text{O}_2$	0.195		terephthalic acid	0.38	1.95	5185
3) $\text{H}_5\text{C}_6\text{-OH} + \text{CO}_2$	0.340		salicylic acid	1.68	4.94	55
4) $\text{C}_6\text{H}_6 + \frac{9}{2} \text{O}_2$	0.123		maleic anhydride	0.55	4.47	371.8
1) $\text{C}_2\text{H}_4 + \frac{1}{2} \text{O}_2$	0.180	CH_3CHO	acetaldehyde	0.37	2.06	625
2) $\text{C}_2\text{H}_4 + \frac{1}{2} \text{O}_2$	0.180		ethylene oxide	0.35	1.94	5833
3) $\text{CH}_2=\text{CHCH}_3 + \frac{1}{2} \text{O}_2$	0.154		propylene oxide	0.48	3.12	2210
4) $\text{CH}_2=\text{CHCH}_3 + \text{O}_2$	0.154	$\text{CH}_2=\text{CHCHO}$	acrolein	0.62	4.03	16
5) $\text{C}_2\text{H}_4\text{O} + \text{H}_2\text{O}$	0.355	$\text{HOCH}_2\text{CH}_2\text{OH}$	ethylene glycol	0.31	0.87	4823
6) $\text{C}_3\text{H}_6\text{O} + \text{H}_2\text{O}$	0.485		propylene glycol	0.43	0.89	615
7) $\text{CH}_2=\text{CHCHO} + \text{CH}_3\text{OH}$	0.699	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{CHO}$	methoxypropion-aldehyde	~0.90	1.29	231.1
8) $\text{CH}_2=\text{CH-CHO} +$ $\text{CH}_3\text{CH(OH)CH}_3$	0.827	$\text{CH}_2=\text{CHCH}_2\text{OH}$ + CH_3COCH_3	allyl alcohol + acetone	1.17	1.41	1873
9) $\text{CH}_2=\text{CHCH}_2\text{OH} + \text{H}_2\text{O}_2$	1.13	$\text{CH}_2\text{OH-CHOH-CH}_2\text{OH}$	glycerine	0.91	0.81	315

+ Product Price/Reactant(s) Cost

++ 10^6 lbs/yr

COMMERCIAL PROCESSES

A number of commercial processes are employed to synthesize the desired chemical products. Conventional methods for producing several high \$-value chemicals are discussed as well as processes used by some major companies that directly relate to the paths proposed in this report. The commercial processes discussed follow, in general, the sequence of products listed in Table 5.

Prior to 1955 benzene was produced primarily from coal gas as a by-product of steel manufacturing. Typically one ton of coal yields 1.85-2.40 gallons of benzene [14]; the process flow is shown in Figure 4. By the 1970's however, about 94% of aromatics were produced from petroleum. The petroleum derived methods include [14]: 1) catalytic reforming (conversion of naphthenes to aromatics), shown in Figure 5; 2) hydrodealkylation (HDA) of toluene; 3) extraction from pyrolysis gasoline; and 4) disproportionation of toluene. Very recently a new process has appeared, called the dehydrocyclization of n-paraffins to aromatics [19], which eliminates the use of noble metal catalyst and hydrogen with the feedstock.

Phenol can be obtained from several sources. These methods include [14]: 1) a benzene sulfonate process requiring cheap sulfuric acid and caustic soda (the disposal of sodium sulfate and sodium sulfite must be considered); 2) a chlorobenzene process requiring low cost chlorine; 3) a regenerative (Raschig) process requiring large scale continuous units; and more recently, 4) a cumene peroxidation process requiring cheap acetone; and 5) a toluene oxidation process.

Aniline is most typically produced by catalytic reduction of nitrobenzene in the presence of Raney nickel [15]. Acetanilide is manufactured from refluxing aniline and acetic acid for 6-14 hours [15]. Aniline can be alkylated in the presence of sodamide to give a 75% yield of ethylaniline [16]. Mixtures of 2-ethylaniline and 2,6-diethylaniline are obtained in high yield when aniline

Figure 4. Aromatics from Coal Gas and Tar Light Oil [14]

Reaction (Typical)

Figure 5. Aromatics from Petroleum by Reforming [14]

is heated with ethylene at 200-300°C under pressure in the presence of aluminum anilide.

Resorcinol, hydroquinone, and catechol are isomers of dihydroxybenzene [16]. Sodium dichromate or manganese dioxide can be used to oxidize aniline, with a 90% conversion to hydroquinone. Catechol can be manufactured by hydrolysis of o-chlorophenol in aqueous sodium hydroxide with copper sulfate as catalyst. In the presence of ferrous catalyst, hydroxylation of phenol with hydrogen peroxide directly gives catechol and hydroquinone. Resorcinol and hydroquinone can be prepared from the appropriate isomer of diisopropyl benzene (DIPB) using silica-alumina catalyst, producing 78% yield.

Several methods exist for the production of cresol. Cresol is obtained by refining tar acids, the phenolic constituents present in coal tar [14]. Small amounts of cresol are recovered from cracked naphtha. It can also be produced synthetically by the sulfonation of toluene followed by hydrolysis. Cumene oxidation is another synthesis method. Spence and Sons [17] synthesizes cresols by the alkylation of phenol with methanol; this catalytic (Al_2O_3) process can be operated in fixed-bed or fluid bed, at temperatures of 340-355°C.

Salicylic acid can be made by treating phenol with sodium hydroxide, followed by treatment with dry carbon dioxide gas and precipitation with acid [14].

Benzaldehyde is an industrially important organic used in the production of dyes, pharmaceuticals, and perfumeries. Any of three different methods is employed, depending upon the desired use of the product [16]: 1) Rhone-Poulenc S.A. [18] uses a catalytic ($\text{V}_2\text{O}_5/\text{SiO}_2\text{-Al}_2\text{O}_3$) vapor phase air oxidation process operated in fixed- or fluid bed, at a temperature of ~400°C, which simultaneously produces benzaldehyde and benzoic acid; 2) Liquid phase oxidation of toluene is recommended for small scale production; 3) If chlorine free benzaldehyde is not required, liquid phase chlorination of toluene is the cheapest route.

Benzyl alcohol is manufactured on a commercial scale from benzyl chloride and sodium carbonate with a yield of 70-72% [16]. The still residues from this process can be distilled separately to yield benzyl ether as a by-product.

2,4-Xylidine and 2,5-xylidine are widely employed in the dye industry [16]. They are usually obtained by iron reduction of nitrobenzene.

Guaiacol can be prepared by the mercuric oxide oxidation of lignin, the zinc chloride reduction of acetovanillone, and diazotization and hydrolysis of o-anisidine [16].

As discussed previously, the conversion of n-paraffins to aromatics also results in the production of olefins, such as ethylene and propylene. The Shell Development Company [20] utilizes a direct oxidation process to form epoxides. This catalytic vapor phase oxidation process at 327°C in fixed- or fluid bed has conversions of 50-70%, also producing some complete oxidation products, CO₂ and H₂O.

CONCLUSIONS AND RECOMMENDATIONS

The original objective of the report was to present a list of endothermic chemical reactions, which could utilize broad band IR-VIS-UV radiation, to synthesize high \$-value products from low cost feedstocks. In the intervening year and a half since the original report, laboratory experiments have been conducted, more literature search and calculations have been carried out, and a broader view of the entire project has evolved.

Two additional considerations have been recognized: 1)-the dehydrocyclization of C_6-C_8 n-paraffins to aromatics also produces a significant fraction of C_2-C_3 olefins; and 2)-a cost effective solar chemical plant must utilize its equipment during the night-time as well as the day-light hours. The combination of these two facts leads to the possibility of carrying out exothermic olefin oxidation reactions during the night-time hours.

Conclusions are:

- 1) -pricewise chemicals fall into four (4) ascending value classes: primary feedstocks, intermediates, products, and specialty products ($> \$1.00/lb$);
- 2) -conversion of paraffins to aromatics appears to be a very important primary step towards high \$-value products;
- 3) -the endothermic dehydrocyclization of C_6-C_8 n-paraffins to aromatics, followed by alkylation and hydroxylation of the aromatic ring can lead to a class of high \$-value products;
- 4) -the product price to reactant(s) cost ratio values are in the range of 5 to 8 for the best reaction possibilities;
- 5) -to be cost effective catalyst choice for paraffin to aromatic conversion should optimize aromatic w% yield $> 60\%$;
- 6) -the conversion of olefins to oxygenated compounds is an important supplement to producing an overall cost effective process.

Recommendations are:

- 1) -that the initial experimental reaction sequence should be, n-paraffins to aromatics to alkylated-hydroxylated rings; and,
- 2) -then followed by experiments on the oxidation of C_2-C_3 olefins to judicious choices amongst epoxides, aldehydes, glycols, alcohols, and ketones.

SELECTED REFERENCES

- [1] Wentworth, W.E., C.F. Batten, Gong Wei, "The Photo-Assisted Thermal Decomposition of Methanol and Isopropanol in a Fluidized Bed."
- [2] Lawler, Gloria M., ed. "Chemical Origins And Markets," 5th edition, Stanford Research Institute, Calif., 1977.
- [3] "Top 50 Chemicals," C&E News, p. 35, June 9, 1986.
- [4] "Chemical Prices," Chemical Marketing Reporter, published by Schnell Publishing Company, Inc., New York, pp. 32-39, June 9, 1986.
- [5] Prengle, H.W., Private Communications, 8/86 to 12/86.
- [6] Stull, Daniel R., Edgar F. Westrum, Jr., and Gerald C. Sinke, "The Chemical Thermodynamics of Organic Compounds," John Wiley & Sons, New York, 1969.
- [7] Reid, Robert C., John M. Prausnitz, and Thomas K. Sherwood, "The Properties of Gases and Liquids," 3rd ed., McGraw-Hill Book Company, New York, 1977.
- [8] Weast, Robert C., ed., CRC Handbook of Chemistry and Physics," 61st edition, CRC Press, Florida 1980.
- [9] Streitwieser, Andrew, Jr., and Clayton H. Heathcock, "Introduction to Organic Chemistry," Macmillan Publishing Co., New York, 1981.
- [10] Hatch, Lewis F., "The Chemistry of Petrochemical Reactions," Gulf Publishing Company, Houston, Texas, 1955.
- [11] Schuster, Herbert F. and Gary M. Coppola, "Allenenes in Organic Synthesis," John Wiley & Sons, New York, 1984.
- [12] Timmons, C.J., ed., "Modern Reactions in Organic Synthesis," Van Nostrand Reinhold Company, 1970.
- [13] Saghafi, Majid, Senior Honors Thesis on: "Fundamental Mechanisms of Photo-Thermal Catalysts."
- [14] Faith, W.L., Donald B. Keyes, Ronald L. Clark, "Industrial Chemicals," 3rd ed., John Wiley & Sons, New York, 1965.
- [15] McKetta, J.J., "Chemical Processing and Design," Vol 1-14, Marcel Decker, Inc., New York, 1976.
- [16] Kirk - Othmer, "Encyclopedia of Chemical Technology," 3rd ed., Vol 1-24, John Wiley & Sons, New York, 1978.

- [17] N.M. Cullinane and W.C. Davies, Brit. Pat. #600,837, 20 Apr 1948; *ibid*, Brit. Pat., #600,838, 20 Apr 1948; *ibid*, Brit. Pat. #602,257. 24 May 1948.
- [18] J. Bonnard and G. Poilane, U.S. Pat. #3,387,036, 4 June 1969; J.C. Brunie, U.S. Pat. #3,658,875, 25 Apr 1972; M. Jouffret, U.S. Pat. #3,948,995, 6 Apr 1976.
- [19] F-M Lee, "Catalytic Dehydrocyclization-Dehydrogenation of Hydrocarbons" U.S. Patent #4,607,129 (1986).
- [20] Shell Development Company, U.S. Pat., #3,119,837, (1964).

APPENDIX

A. Estimation of Thermodynamic Properties

For the estimation of heats of formation, ΔH_f° , and free energies of formation, ΔF_f° , an accurate method has been developed by Benson and his colleagues [6-8]. It is a group-contribution method applicable for C_p° , ΔH_{f298}° , and ΔS_{298}° . Once these fundamental properties are obtained, the free energy of formation and the properties at elevated temperatures can be calculated using the following relationships.

$$\Delta F_T^\circ = \Delta H_T^\circ - T\Delta S_T^\circ \quad (a-1)$$

$$\Delta H_{fT}^\circ = \Delta H_{f298}^\circ + \int_{298}^T \Delta C_p^\circ dT \quad (a-2)$$

$$\Delta S_T^\circ = \Delta S_{298}^\circ + \int_{298}^T \Delta C_p^\circ d \ln T \quad (a-3)$$

Contributions are given for atoms with valences greater than one. It is necessary to become familiar with the notation used in Table A-1 to employ this method. For example, contributions for a methyl group attached to a carbon atom, $-\text{CH}_3$, are represented as $\text{C}-(\text{C})(\text{H})_3$. For a description of other shorthand used, refer to the footnotes following Table A-1.

To demonstrate the method for future use, the following example is provided.

example: Estimation of ΔH_{f298}° and ΔS_{298}° for m-cresol,

#	atom	notation	ΔH_{f298}°	ΔS_{298}°	actual values
4	2,3,4,6	$\text{C}_B-(\text{H})$	3.30	11.53	$H_{f298}^\circ = -31.63$
1	5	$\text{C}_B-(\text{O})$	-0.90	-10.20	
1	1	$\text{C}_B-(\text{C})$	5.51	-7.69	$S_{298}^\circ = 85.27$
1	7	$\text{O}-(\text{C}_B)(\text{H})$	-37.9	29.1	
1	8	$\text{C}-(\text{C}_B)(\text{H})_3$	-10.8	30.41	
Σ			-30.17	87.74	

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties*

Group ^a	$\Delta H_{f, \text{gas}}^\circ$ kcal/g-mol	$S_{f, \text{gas}}^\circ$ cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Hydrocarbon Groups								
C—(C)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C) ₂ (H) ₂	-4.95	9.42	5.50	6.95	8.25	9.35	11.07	12.34
C—(C) ₃ (H)	-1.90	-12.07	4.54	6.00	7.17	8.05	9.31	10.05
C—(C) ₄	0.50	-35.10	4.37	6.13	7.36	8.12	8.77	8.76
C _d —(H) ₃	6.26	27.61	5.10	6.36	7.51	8.50	10.07	11.27
C _d —(C)(H)	8.59	7.97	4.16	5.03	5.81	6.50	7.65	8.45
C _d —(C) ₂	10.34	-12.7	4.10	4.61	4.99	5.26	5.80	6.08
C _d —(C _d)(H)	6.78	6.38	4.46	5.79	6.75	7.42	8.35	8.99
C _d —(C _d)(C)	8.88	-14.6	(4.40)	(5.37)	(5.93)	(6.18)	(6.50)	(6.62)
C _d —(C _d) ₂	4.6							
C _d —(C _u)(H)	6.78	6.4	4.46	5.79	6.75	7.42	8.35	8.99
C _d —(C _u)(C)	8.64	(-14.6)	(4.40)	(5.37)	(5.93)	(6.18)	(6.50)	(6.62)
C _d —(C _u) ₂	8.0							
C _d —(C _i)(H)	6.78	6.4	4.46	5.79	6.75	7.42	8.35	8.99
C _d —(C _i)(C)	8.53	...	4.40	5.37	5.93	6.18	6.50	6.62
C—(C _d)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C _d) ₂ (H) ₂	-4.29	(10.2)	(4.7)	(6.8)	(8.4)	(9.6)	(11.3)	(12.6)
C—(C _d) ₃ (C) ₂	1.16	...	3.57	5.98	7.51	8.37	9.00	9.02
C—(C _d)(C) ₄	1.68	(-34.72)	(3.99)	(6.04)	(7.43)	(8.26)	(8.92)	(8.96)
C—(C _d)(C)(H) ₂	-4.76	9.8	5.12	6.86	8.32	9.49	11.22	12.48
C—(C _d)(C) ₂ (H)	-1.48	(-11.7)	(4.16)	(5.91)	(7.34)	(8.19)	(9.46)	(10.19)
C—(C _d) ₂ (C)(H)	-1.24	...	3.74	5.85	7.32	8.30	9.54	10.31
C—(C _i)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C _i)(C)(H) ₂	-4.73	10.3	4.95	6.56	7.93	9.08	10.86	12.19
C—(C _i)(C) ₂ (H)	-1.72	(-11.2)	(3.99)	(5.61)	(6.85)	(7.78)	(9.10)	(9.90)
C—(C _u)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C _u)(C)(H) ₂	-4.86	9.3	5.84	7.61	8.98	10.01	11.49	12.54
C—(C _u)(C) ₂ (H)	-0.98	(-12.2)	(4.88)	(6.66)	(7.90)	(8.75)	(9.73)	(10.25)
C—(C _u)(C) ₃	2.81	(-35.18)	(4.37)	(6.79)	(8.09)	(8.78)	(9.19)	(8.96)
C—(C _u) ₂ (C)(H)	-1.24	...	3.74	5.85	7.32	8.30	9.54	10.31
C—(C _u) ₂ (C) ₂	1.16	...	3.57	5.98	7.51	8.37	9.00	9.02
C—(C _u)(C _d)(H) ₂	-4.29	(10.2)	(4.7)	(6.8)	(8.4)	(9.6)	(11.3)	(12.6)
C _i —(H)	26.93	24.7	5.27	5.99	6.49	6.87	7.47	7.96
C _i —(C)	27.55	6.35	3.13	3.48	3.81	4.09	4.60	4.92
C _i —(C _d)	29.20	(6.43)	(2.57)	(3.54)	(3.50)	(4.92)	(5.34)	(5.50)
C _i —(C _u)	(29.20)	6.43	2.57	3.54	3.50	4.92	5.34	5.50
C _u —(H)	3.30	11.53	3.24	4.44	5.46	6.30	7.54	8.41
C _u —(C)	5.51	-7.69	2.67	3.14	3.68	4.15	4.96	5.44
C _u —(C _d)	5.68	-7.80	3.59	3.97	4.38	4.72	5.28	5.61
C _u —(C _i)	5.7	-7.80	3.59	3.97	4.38	4.72	5.28	5.61
C _u —(C _u)	4.96	-8.64	3.33	4.22	4.89	5.27	5.76	5.95
C _n	34.20	6.0	3.9	4.4	4.7	5.0	5.3	5.5
Next-Nearest-Neighbor Correction								
Alkane gauche	0.80							
Alkene gauche	0.50							
Cis	1.00		-1.34	-1.09	-0.81	-0.61	-0.39	-0.26
Ortho	0.57	-1.61	1.12	1.35	1.30	1.17	0.88	0.66

Ring (σ) ^{a/}	$\Delta H^\circ_{\text{ring}}$ kcal/g-mol	S°_{ring} cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Corrections to be Applied for Ring Compounds								
Cyclopropane (6)	27.6	32.1	-3.05	-2.53	-2.10	-1.90	-1.77	-1.62
Cyclopropene (2)	53.7	33.6						
Cyclobutane (8)	26.2	29.8	-4.61	-3.89	-3.14	-2.64	-1.88	-1.38
Cyclobutene (2)	29.8	29.0	-2.53	-2.19	-1.89	-1.68	-1.48	-1.33
Cyclopentane (10)	6.3	27.3	-6.5	-5.5	-4.5	-3.8	-2.8	-1.9
Cyclopentene (2)	5.9	25.8	-5.98	-5.35	-4.89	-4.14	-2.93	-2.26
Cyclopentadiene	6.0	28.0	-4.5					
Cyclohexane (6)	0	18.8	-5.8	-4.1	-2.9	-1.5	1.1	2.2
Cyclohexene (2)	1.4	21.5	-4.28	-3.04	-1.98	-1.43	-0.29	0.08
Cycloheptane (1)	6.4	15.9						
Cyclooctane (8)	9.9	16.5						
Naphthalene	...	8.1						

Table A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Group ^a	ΔH_f° , kcal/g-mol	S° , cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Oxygen-containing Compounds								
CO—(CO)(H)	-26.0	...	6.72	7.83	8.90	9.89	11.43	12.12
CO—(CO)(C)	-29.2	...	5.46	6.32	7.16	7.87	9.00	9.76
CO—(O)(C ₂)	-32.5	...	5.97	6.7	7.41	8.02	8.87	9.36
CO—(O)(C ₃)	-32.5	...	2.18	2.75	3.98	5.03	6.29	7.06
CO—(O)(C)	-33.1	4.78	5.97	6.70	7.40	8.02	8.87	9.36
CO—(O)(H)	-32.1	34.93	7.03	7.87	8.82	9.68	11.16	12.20
CO—(C ₂)(H)	-31.7	...	7.03	7.87	8.82	9.68	11.16	12.20
CO—(C ₃) ₂	-38.1	...	5.26	6.77	7.67	8.48	9.62	9.85
CO—(C ₃)(C)	-30.9	...	5.68	6.92	7.70	8.36	9.39	9.76
CO—(C ₃)(H)	-34.6	...	6.40	7.72	8.91	9.85	11.49	12.09
CO—(C) ₂	-31.4	15.01	5.59	6.32	7.09	7.76	8.89	9.61
CO—(C)(H)	-29.1	34.93	7.03	7.87	8.82	9.68	11.16	12.20
CO—(H) ₂	-26.0	53.67	8.47	9.38	10.46	11.52	13.37	14.81
O—(C ₂)(CO)	-32.5	...	2.06	2.70	3.11	3.42	3.88	4.18
O—(CO) ₂	-50.9	...	-0.41	1.78	3.20	4.00	5.13	5.85
O—(CO)(O)	-19.0	...	3.7	3.7	3.7	3.7	4.2	4.2
O—(CO)(C ₂)	-46.9	...	1.44	2.98	3.98	4.49	4.97	5.20
O—(CO)(C)	-44.3	8.39	3.90	3.61	4.19	4.62	4.99	4.82
O—(CO)(H)	-58.1	24.52	3.81	4.98	5.80	6.34	7.19	7.75
O—(O)(C)	(-4.5)	(9.4)	(3.7)	(3.7)	(3.7)	(3.7)	(4.2)	(4.2)
O—(O) ₂	(-19.0)	(9.4)	(3.7)	(3.7)	(3.7)	(3.7)	(4.2)	(4.2)
O—(O)(H)	-16.27	27.85	5.17	5.79	6.28	6.66	7.15	7.51
O—(C ₂) ₂	-32.8	10.1	3.4	3.7	3.7	3.8	4.4	4.6
O—(C ₂)(C)	-31.9	9.7	3.4	3.7	3.7	3.8	4.4	4.6
O—(C ₃) ₂	-21.1	...	1.09	1.22	1.50	1.99	2.85	3.51
O—(C ₃)(C)	-22.6	...	3.4	3.7	3.7	3.8	4.4	4.6
O—(C ₃)(H)	-37.9	29.1	4.3	4.5	4.8	5.2	6.0	6.6
O—(C) ₂	-23.7	8.68	3.4	3.7	3.7	3.8	4.4	4.6
O—(C)(H)	-37.9	29.07	4.33	4.45	4.82	5.23	6.02	6.61
C ₂ —(CO)(O)	9.0	...	5.59	7.00	7.48	7.75	8.02	8.13
C ₂ —(CO)(C)	9.4	...	3.73	4.48	5.02	5.40	5.95	6.37
C ₂ —(CO)(H)	8.5	...	3.79	4.90	5.84	6.64	7.80	8.74
C ₂ —(O)(C ₂)	8.9	...	(4.40)	(5.37)	(5.93)	(6.18)	(6.50)	(6.62)
C ₂ —(O)(C)	10.3	...	4.10	4.61	4.99	5.26	5.80	6.08
C ₂ —(O)(H)	8.6	...	4.16	5.03	5.81	6.50	7.65	8.45
C ₂ —(CO)	9.7	...	2.67	3.14	3.68	4.15	4.96	5.44
C ₂ —(O)	-0.9	-10.2	3.9	5.3	6.2	6.6	6.9	6.9
C—(CO) ₂ (H) ₂	-7.6	...	5.60	7.05	8.39	9.68	11.58	12.87
C—(CO)(C) ₂ (H)	-1.8	-12.0	6.21	7.56	8.00	8.21	9.18	9.63
C—(CO)(C)(H) ₂	-5.2	9.6	6.2	7.7	8.7	9.5	11.1	12.2
C—(CO)(C) ₂	1.6	...	5.07	6.88	7.81	8.27	8.80	8.62
C—(CO)(H) ₃	-10.1	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(O) ₂ (C) ₂	-18.6	...	1.59	3.95	6.20	7.39	7.62	8.48
C—(O) ₂ (C)(H)	-16.3	...	5.06	7.28	9.03	9.41	10.31	10.75
C—(O) ₂ (H) ₂	-15.1	...	2.83	5.06	7.52	9.12	10.32	11.29
C—(O)(C ₃)(H) ₂	-8.1	9.7	3.71	6.27	8.28	9.79	11.79	13.20
C—(O)(C ₃)(C)(H)	-6.08	...	5.14	7.30	8.83	9.43	10.23	10.6
C—(O)(C ₂)(H) ₂	-6.9	...	4.66	6.97	8.65	9.88	11.54	12.73
C—(O)(C) ₃	-6.60	-33.56	4.33	6.19	7.25	7.70	8.20	8.24
C—(O)(C) ₂ (H)	-7.2	-11.00	4.80	6.64	8.10	8.73	9.81	10.40
C—(O)(C)(H) ₃	-8.1	9.8	4.99	6.85	8.30	9.43	11.11	12.33
C—(O)(H) ₃	-10.1	30.41	6.19	7.84	9.40	10.79	13.03	14.77

Table A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Strain or rings	$\Delta H_{f,298}^\circ$ kcal/g-mol	S_{298}° cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Strain and Ring Corrections for Oxygen-containing Compounds (Continued)								
Ether oxygen, gauche	0.3	...	-0.10	-0.89	-1.10	-0.73	-0.60	-0.23
Ditertiary ethers	7.8	...	-3.94	-5.64	-7.15	-8.83	-12.04	-14.90
Ethylene oxide,	27.6	31.4	-2.0	-2.8	-3.0	-2.6	-2.3	-2.3
Trimethylene oxide,	26.4	27.7	-4.6	-5.0	-4.2	-3.5	-2.6	0.2
Tetrahydrofuran,	6.7	...	-4.25	-4.54	-4.07	-3.55	-3.09	-2.61
Tetrahydropyran,	2.2	...	-4.28	-3.04	-1.98	-1.43	-0.29	0.08
1,3-Dioxane,	0.9	...	-2.51	-2.88	-2.28	-1.49	-0.26	0.56
1,4-Dioxane,	5.4	...	-4.16	-4.57	-3.11	-1.88	-1.09	-0.47
1,3,5-Trioxane,	5.1	...	1.79	0.56	-0.61	-0.65	-1.20	-2.43
Furan,	-5.8	...	-4.19	-3.63	-2.92	-2.39	-1.99	-1.72
Dihydropyran,	1.2	...	-4.44	-3.20	-1.56	-0.45	0.42	0.66

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Strain or rings	$\Delta H_{f, \text{gas}}^\circ$, kcal/g-mol	$S_{f, \text{gas}}^\circ$, cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Strain and Ring Corrections for Oxygen-containing Compounds (Continued)								
Cyclopentanone,	5.2	...	-8.53	-7.19	-5.31	-3.72	-2.26	-1.22
Cyclohexanone,	2.2	...	-8.10	-6.57	-4.24	-1.91	0.70	1.97
Succinic anhydride,	4.5	...	-7.90	-6.02	-4.49	-3.58	-3.35	-3.06
Glutaric anhydride,	0.8	...	-7.93	-6.04	-4.50	-3.59	3.35	-3.07
Maleic anhydride,	3.6	...	-5.12	-3.38	-2.02	-2.19	-0.37	-0.01

Group*	$\Delta H_{f, \text{gas}}^\circ$, kcal/g-mol	$S_{f, \text{gas}}^\circ$, cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Nitrogen-containing Compounds								
C-(N)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C-(N)(C)(H) ₂	-6.6	9.8	5.25	6.90	8.28	9.39	11.09	12.34
C-(N)(C) ₂ (H)	-5.2	-11.7	4.67	6.32	7.64	8.39	9.56	10.23
C-(N)(C) ₃	-3.2	-34.1	4.35	6.16	7.31	7.91	8.49	8.50
N-(C)(H) ₂	4.8	29.71	5.72	6.51	7.32	8.07	9.41	10.47
N-(C) ₂ (H)	15.4	8.94	4.20	5.21	6.13	6.83	7.90	8.65
N-(C) ₃	24.4	-13.46	3.48	4.56	5.43	5.97	6.56	6.67
N-(N)(H) ₂	11.4	29.13	6.10	7.38	8.43	9.27	10.54	11.52
N-(N)(C)(H)	20.9	9.61	4.82	5.8	6.5	7.0	7.8	8.3
N-(N)(C) ₂	29.2	-13.80	1.56	2.50	3.31	3.87	4.62	4.99
N-(N)(C ₂)(H)	22.1	...	3.28	4.05	4.75	5.31	6.28	6.91
N ₁ -(H)	(16.3)	(12.3)	2.95	4.58	6.45	7.71	9.13	9.92
N ₁ -(C)	21.3	...	2.48	3.34	3.95	4.29	4.59	4.60
N ₁ -(C ₂)	16.7	...	2.60	3.22	3.81	4.22	4.79	5.12
N ₂ -(H)	25.1	26.8	4.38	4.89	5.44	5.94	6.77	7.42
N ₂ -(C)	32.5	8.0	2.70	4.10	4.92	5.34	5.60	5.71
N-(C ₂)(H) ₂	4.8	29.71	5.72	6.51	7.32	8.07	9.41	10.47
N-(C ₂)(C)(H)	14.9	...	3.82	4.89	5.71	6.28	7.19	7.73
N-(C ₂)(C) ₂	26.2	...	0.62	2.02	3.27	4.13	5.23	5.59
N-(C ₂)(H)	16.3	...	2.16	3.12	4.13	5.10	6.76	7.88
C ₂ -(N)	-0.5	-9.69	3.95	5.21	5.94	6.32	6.53	6.56
N ₂ -(N)	23.0	...	2.12	4.18	5.51	6.77	6.86	7.05
CO-(N)(H)	-29.6	34.93	7.03	7.87	8.82	9.68	11.16	12.20
CO-(N)(C)	-32.8	16.2	5.37	6.17	7.07	7.66	9.62	11.19
N-(CO)(H) ₂	-14.9	24.69	4.07	5.74	7.13	8.29	9.96	11.22
N-(CO)(C)(H)	-4.4	3.9	3.87	5.08	5.95	6.76	6.87	6.54
N-(CO)(C) ₂	4.7	...	1.83	3.79	5.24	6.19	7.11	7.42
N-(CO)(C ₂)(H)	0.4	...	3.03	3.91	4.60	5.58	6.23	6.32

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Group ^a	$\Delta H_{f,298}^\circ$ kcal/g-mol	S_{298}° cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Nitrogen-containing Compounds (Continued)								
N-(CO) ₂ (H)	-18.5	...	3.59	5.54	6.70	7.39	7.95	8.19
N-(CO) ₂ (C)	-5.9	...	1.07	3.10	4.31	5.00	5.48	6.47
N-(CO) ₂ (C ₂)	-0.5	...	0.98	3.06	4.23	4.85	5.28	5.29
C-(CN)(C)(H) ₂	22.5	40.20	11.10	13.40	15.50	17.20	19.7	21.30
C-(CN)(C) ₂ (H)	25.8	19.80	11.00	12.70	14.10	15.40	17.30	18.60
C-(CN)(C) ₃	29.0	-2.80	8.65	11.16	12.89	14.05	15.51	16.19
C-(CN) ₂ (C) ₂	...	28.40	14.72	17.79	20.00	21.61	23.78	24.96
C ₂ -(CN)(H)	37.4	36.58	9.80	11.70	13.30	14.50	16.30	17.30
C ₂ -(CN)(C)	39.15	15.91	9.74	11.28	12.48	13.26	14.45	14.93
C ₂ -(CN) ₂	84.1	...	13.60	16.55	18.68	20.25	22.34	23.59
C ₂ -(NO) ₂ (H)	...	44.4	12.5	15.1	17.4	19.2	21.6	23.2
C ₂ -(CN)	35.8	20.50	9.8	11.2	12.5	13.1	14.2	14.9
C ₂ -(CN)	63.8	35.40	10.30	11.30	12.10	12.70	13.60	14.30
C-(NO ₂)(C)(H) ₂	-15.1	48.4	12.59	15.82	18.52	20.66	23.79	25.90
C-(NO ₂)(C) ₂ (H)	-15.8	26.9	11.99	15.21	17.72	19.61	22.18	23.70
C-(NO ₂)(C) ₃	...	3.9	9.89	13.34	15.86	17.62	19.41	20.86
C-(NO ₂) ₂ (C)(H)	-14.9	...	17.32	22.82	27.07	30.21	34.35	36.83
O-(NO)(C)	-5.9	41.9	9.10	10.30	11.2	12.0	13.3	13.9
O-(NO ₂)(C)	-19.4	48.50	9.54	11.54	13.26	15.60	16.39	17.38

Ring	$\Delta H_{f,298}^\circ$ kcal/g-mol	S_{298}° cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Ring Corrections for Nitrogen-containing Compounds								
Ethyleneimine,	27.7	31.6	-2.07	-2.18	-2.17	-2.05	-1.94	-1.88
Azetidine,	26.2	29.3	-4.73	-4.52	-4.08	-3.61	-2.66	0.01
Pyrrolidine,	6.8	26.7	-6.17	-5.58	-4.80	-4.00	-2.87	-2.17
Piperidine,	1.0	...	-0.56	0.37	1.08	1.56	1.71	-0.46
C ₆ H ₁₂ N ₂ ,	3.4	...	-9.44	-8.70	-7.00	-5.36	-3.21	-1.84
Succinimide,	8.5	...	2.16	4.08	6.14	8.01	9.11	9.77

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Group	$\Delta H_{f,298}^\circ$, kcal/g-mol	S_{298}° , cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Halogen Groups								
C—(F) ₃ (C)	-158.4	42.5	12.7	15.0	16.4	17.9	19.3	20.0
C—(F) ₂ (H)(C)	-109.3	39.1	9.9	12.0	13.7	15.1	16.7	17.8
C—(F)(H) ₂ (C)	-51.5	35.4	8.1	10.0	12.0	13.0	15.2	16.6
C—(F) ₂ (C) ₂	-97.0	17.8	9.9	11.8	13.5	14.4	16.1	16.6
C—(F)(H)(C) ₂	-49.0	14.0	7.30	9.04	10.47	11.56	13.10	14.01
C—(F)(C) ₃	-48.5	...	6.80	8.86	10.20	11.16	12.43	12.72
C—(F) ₂ (Cl)(C)	-106.3	40.5	13.7	16.1	17.5	18.6	19.8	20.4
C—(Cl) ₃ (C)	-20.7	50.4	16.3	18.0	19.1	19.8	20.6	21.0
C—(Cl) ₂ (H)(C)	(-18.9)	43.7	12.1	14.0	15.4	16.5	17.9	18.7
C—(Cl)(H) ₂ (C)	-16.5	37.8	8.9	10.7	12.3	13.4	15.3	16.7
C—(Cl) ₂ (C) ₂	-22.0	22.4	12.2	14.88	15.95	16.48	16.96	17.02
C—(Cl)(H)(C) ₂	-14.8	17.6	9.0	9.9	10.5	11.2	13.9	14.6
C—(Cl)(C) ₃	-12.8	-5.4	9.3	10.5	11.0	11.3	12.4	12.7
C—(Br) ₃ (C)	...	55.7	16.7	18.0	18.8	19.4	19.9	20.3
C—(Br)(H) ₂ (C)	-5.4	40.8	9.1	11.0	12.6	13.7	15.5	16.8
C—(Br)(H)(C) ₂	-3.4	...	8.93	10.66	11.96	12.84	14.05	14.72
C—(Br)(C) ₃	-0.4	-2.0	9.3	11.0	11.5	12.3	13.3	13.3
C—(I)(H) ₂ (C)	8.0	43.0	9.2	11.0	12.9	13.9	15.8	17.2
C—(I)(H)(C) ₂	10.5	21.3	9.2	10.9	12.2	13.0	14.2	14.8
C—(I)(C)(C) ₂ (H)	13.32	...	8.13	10.02	11.82	12.6	14.0	14.9
C—(I)(C) ₂ (H) ₂	8.19	...	8.82	10.91	12.97	14.04	15.95	17.34
C—(I)(C) ₃	13.0	0	9.83	11.75	12.92	13.45	13.79	13.60
C—(C)(Br)(H)(C)	...	45.7	12.4	14.0	15.6	16.3	17.9	19.0
N—(F) ₃ (C)	-7.8	...	8.25	10.13	11.52	12.80	14.37	14.98
C—(Cl)(C)(O)(H)	-21.6	15	9.85	10.41	11.05	11.57	12.45	13.14
C—(I) ₂ (C)(H)	(26.0)	(54.6)	12.69	14.78	16.21	17.12	18.31	19.03
C—(I)(O)(H) ₂	3.8	40.7	8.22	10.49	12.23	13.55	15.35	16.57
C ₂ —(F) ₂	-77.5	37.3	9.7	11.0	12.0	12.7	13.8	14.5
C ₂ —(Cl) ₂	-1.8	42.1	11.4	12.5	13.3	13.9	14.6	15.0
C ₂ —(Br) ₂	...	47.6	12.3	13.2	13.9	14.3	14.9	15.2
C ₂ —(F)(Cl)	...	39.8	10.3	11.7	12.6	13.3	14.2	14.7
C ₂ —(F)(Br)	...	42.5	10.8	12.0	12.8	13.5	14.3	14.7
C ₂ —(Cl)(Br)	...	45.1	12.1	12.7	13.5	14.1	14.7	14.7
C ₂ —(F)(H)	-37.6	32.8	6.8	8.4	9.5	10.5	11.8	12.7
C ₂ —(Cl)(H)	-1.2	35.4	7.9	9.2	10.3	11.2	12.3	13.1
C ₂ —(Br)(H)	11.0	38.3	8.1	9.5	10.6	11.4	12.4	13.2
C ₂ —(I)(H)	24.5	40.5	8.8	10.0	10.9	11.6	12.6	13.3
C ₂ —(C)(Cl)	-2.1	15.0	8.0	8.4	8.5	9.0	9.2	9.4
C ₂ —(C)(I)	23.6	...	8.9	9.2	9.1	9.4	9.5	9.6
C ₂ —(C ₂)(Cl)	-3.56	...	8.3	9.2	9.4	9.9	9.9	9.9
C ₂ —(C ₂)(I)	22.14	...	9.2	9.9	10.0	10.3	10.3	10.1
C ₂ —(Cl)	...	33.4	7.9	8.4	8.7	9.0	9.4	9.6
C ₂ —(Br)	...	36.1	8.3	8.7	9.0	9.2	9.5	9.7
C ₂ —(I)	...	37.9	8.4	8.8	9.1	9.3	9.6	9.8
C ₂ —(F)	-42.8	16.1	6.3	7.6	8.5	9.1	9.8	10.2
C ₂ —(Cl)	-3.8	18.9	7.4	8.4	9.2	9.7	10.2	10.4
C ₂ —(Br)	10.7	21.6	7.8	8.7	9.4	9.9	10.3	10.5
C ₂ —(I)	24.0	23.7	8.0	8.9	9.6	9.9	10.3	10.5
C—(C ₂)(F) ₂	-162.7	42.8	12.5	15.3	17.2	18.5	20.1	21.0
C—(C ₂)(Br)(H) ₂	-6.9	...	9.29	11.10	12.47	13.69	15.59	16.71
C—(C ₂)(I)(H) ₂	8.4	...	9.78	11.56	12.90	14.08	15.88	16.91
C—(Cl) ₂ (CO)(H)	-17.8	...	12.8	14.75	15.85	16.65	17.93	18.56
C—(Cl) ₂ (CO)	-19.6	...	17.0	18.75	19.55	19.95	20.63	20.86
CO—(Cl)(C)	-30.2	...	8.87	9.44	10.24	11.08	12.53	13.59

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Next nearest neighbors	$\Delta H_{f,298}^\circ$ kcal/g-mol	S_{298}° cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Corrections for Next-Nearest-Neighbor Halogen Compounds								
Ortho (F)(F)	5.0	...	0	0	0	0	0	0
Ortho (Cl)(Cl)	2.2	...	-0.50	-0.44	-0.55	-0.53	-0.28	-0.02
Ortho (alkane) (halogen)	0.6	...	0.42	0.44	0.28	0.19	0.12	0.14
Cis (halogen) (halogen)	0.3	...	-0.19	-0.01	-0.03	-0.17	0	-0.03
Cis (halogen) (alkane)	-0.8	...	-0.97	-0.70	-0.53	-0.47	-0.24	-0.13

Group	$\Delta H_{f,298}^\circ$ kcal/g-mol	S_{298}° cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Organosulfur Groups								
C—(H) ₃ (S)	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C)(H) ₂ (S)	-5.65	9.88	5.38	7.08	8.60	9.97	12.26	14.15
C—(C) ₂ (H)(S)	-2.64	-11.32	4.85	6.51	7.78	8.69	9.90	10.57
C—(C) ₃ (S)	-0.55	-34.41	4.57	6.27	7.45	8.15	8.72	8.10
C—(C ₂)(H) ₂ (S)	-4.73	...	4.11	6.75	8.70	10.15	11.93	13.10
C—(C ₃)(H) ₂ (S)	-6.45	...	5.00	6.99	8.67	10.07	12.41	14.29
C ₂ —(S)	-1.8	10.20	3.90	5.30	6.20	6.60	6.90	6.90
C ₂ —(H)(S)	8.56	8.0	4.16	5.03	5.81	6.50	7.65	8.45
C ₂ —(C)(S)	10.93	-12.41	3.50	3.57	3.83	4.09	4.41	5.00
S—(C)(H)	4.62	32.73	5.86	6.20	6.51	6.78	7.30	7.71
S—(C ₂)(H)	11.96	12.66	5.12	5.26	5.57	6.03	6.99	7.84
S—(C) ₂	11.51	13.15	4.99	4.96	5.02	5.07	5.41	5.73
S—(C)(C ₂)	9.97	...	4.22	5.08	5.56	5.77	5.87	5.87
S—(C ₂) ₂	-4.54	16.48	4.79	5.58	5.53	6.29	7.94	9.73
S—(C ₂)(C)	19.16	...	3.02	3.39	3.71	4.04	4.62	5.00
S—(C ₂) ₂	25.90	...	2.00	2.01	2.24	2.74	3.80	4.71
S—(S)(C)	7.05	12.37	5.23	5.42	5.51	5.51	5.38	5.12
S—(S)(C ₂)	14.5	...	2.89	3.39	3.72	4.15	4.78	5.10
S—(S) ₂	3.04	13.36	4.7	5.0	5.1	5.2	5.3	5.4
C—(SO)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C)(SO)(H) ₂	-7.72	...	4.55	6.42	7.95	9.16	10.95	12.22
C—(C) ₂ (SO)	-3.05	...	3.06	4.58	4.84	6.60	7.54	7.96
C—(C ₂)(SO)(H) ₂	-7.35	...	4.40	6.36	6.94	9.25	10.97	12.25
C ₂ —(SO)	2.3	...	2.67	3.14	3.68	4.15	4.96	5.44
SO—(C) ₂	-14.41	18.10	8.88	10.03	10.50	10.79	10.98	11.17
SO—(C ₂) ₂	-12.0	...	5.72	9.09	9.70	11.45	11.46	11.25
C—(SO ₂)(H) ₃	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C—(C)(SO ₂)(H) ₂	-7.68	...	5.38	7.08	8.60	9.97	12.26	14.15
C—(C) ₂ (SO ₂)(H)	-2.62	...	4.42	6.25	7.56	8.48	9.64	10.30
C—(C) ₃ (SO ₂)	-0.61	...	2.32	4.38	5.70	6.49	7.27	7.46
C—(C ₂)(SO ₂)(H) ₂	-7.14	...	5.00	6.99	8.67	10.07	12.41	14.29
C—(C ₃)(SO ₂)(H) ₂	-5.54	...	3.71	6.57	8.28	9.79	11.89	13.20
C ₂ —(SO ₂)	2.3	...	2.67	3.14	3.68	4.15	4.96	5.44
C ₂ —(H)(SO ₂)	12.53	...	3.04	4.67	5.93	6.84	7.87	8.67
C ₂ —(C)(SO ₂)	14.47	...	1.85	3.11	3.98	4.60	5.33	5.67
SO ₂ —(C ₂)(C ₂)	-68.58	...	9.89	11.50	13.35	14.61	15.72	15.92
SO ₂ —(C ₂) ₂	-73.58	...	11.52	11.97	13.35	14.28	15.38	15.88
SO ₂ —(C) ₂	-69.74	20.90	10.18	11.74	12.92	13.77	15.13	16.0
SO ₂ —(C)(C ₂)	-72.29	...	9.94	11.50	13.45	14.51	15.62	15.92
SO ₂ —(C ₂) ₂	-68.58	...	8.36	11.03	13.55	14.94	15.86	15.96
SO ₂ —(SO ₂)(C ₂)	-76.25	...	9.81	11.50	13.52	14.73	15.71	16.03
CO—(S)(C)	-31.56	15.43	5.59	6.32	7.09	7.76	8.89	9.61
S—(H)(CO)	-1.41	31.20	7.63	8.09	8.12	8.17	8.50	8.24
C—(S)(F) ₃	...	38.9	9.88	13.01	14.83	16.37	18.17	19.11

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Group ^a	$\Delta H_{f,298}^\circ$, kcal/g-mol	S_{298}° , cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
CS—(N) ₂	-31.56	15.43	5.59	6.32	7.09	7.76	8.89	9.61
N—(CS)(H) ₂	12.78	29.19	6.07	7.28	8.18	8.91	10.09	10.98
S—(S)(N)	-4.90	...	3.7	3.7	3.7	3.7	4.2	4.2
N—(S)(C) ₂	29.9	...	3.97	5.17	6.21	6.94	7.39	9.24
SO—(N) ₂	-31.56	...	5.59	6.32	7.09	7.76	8.89	9.61
N—(SO)(C) ₂	16.0	...	4.20	5.88	6.12	6.53	6.83	8.34
SO ₂ —(N) ₂	-31.56	...	5.59	6.32	7.09	7.76	8.89	9.61
N—(SO ₂)(C) ₂	-20.4	...	6.02	6.35	7.54	8.23	9.03	9.19

Ring (σ) ^{a,1}	$\Delta H_{f,298}^\circ$, kcal/g-mol	S_{298}° , cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Ring Corrections for Sulfur-containing Compounds								
Thiirane (2),
17.7	29.47	-2.85	-2.59	-2.66	-3.02	-4.32	-5.82
Trimethylene sulfide (2),
19.37	27.18	-4.59	-4.18	-3.91	-3.91	-4.60	-5.70
Tetrahydrothiophene (2),
1.73	23.56	-4.90	-4.67	-3.68	-3.66	-4.41	-5.57
Thiacyclohexane (2),
0	17.46	-6.22	-4.26	-2.24	-0.69	0.86	1.29
Thiacycloheptane (2),
3.89	...	-7.75	-4.92	-1.22	2.59	4.79	4.61
3-Thiocylopentene (2),
5.07	...	-6.44	-4.24	-4.23	-4.18	-4.80	-5.96
2-Thiocylopentene (1),
5.07	...	-6.44	-4.24	-4.23	-4.18	-4.80	-5.96
C ₄ H ₆ SO ₂ (2),
5.74	...	-4.90	-4.67	-3.68	-3.66	-4.41	-5.57

TABLE A-1 Benson Group Contributions to Ideal-Gas Properties* (Continued)

Ring (σ) ^{a,1}	$\Delta H_{f,298}^\circ$, kcal/g-mol	S_{298}° , cal/g-mol K	C_p° , cal/g-mol K at:					
			300 K	400 K	500 K	600 K	800 K	1000 K
Ring Corrections for Sulfur-containing Compounds (Continued)								
Thiophene (2),
1.73	23.56	-4.90	-4.67	-3.68	-3.66	-4.41	-5.57

*Data were obtained largely from Refs. 4 and 6. Some $H_{f,298}^\circ$ values were obtained from Ref. 11. Many C_p° values are from Ref. 20. Grateful acknowledgment is extended to Shell Development Co., Bellaire Research Center, and particularly to Dr. P. Chueh, for supplying missing values as well as additional contributions. Finally, Dr. S. W. Benson provided an up-to-date errata list that showed recent modifications of the original tables.

¹ C_d represents a carbon atom that is joined to another carbon atom by a double bond. It is considered divalent. For example, 2-pentene would have the groups $C-(C_d)(H)_2$, $C_d-(C)(H)$ twice, $C-(C_d)(H)_2$, and $C-(C)(H)_2$. C_t represents a carbon atom that is joined to another carbon atom by a triple bond. It is considered monovalent. For example, propyne would have the groups $C_t-(H)$, $C_t-(C)$, and $C-(C_t)(H)_2$. C_a represents a carbon atom in an aromatic ring. It is considered monovalent. For example, *p*-ethyl toluene would have the groups $C-(C)(H)_2$, $C-(C_a)(C)(H)_2$, $C-(C_a)(H)_2$, $C_a-(C)$ twice, and $C_a-(H)$ four times. C_c represents the allene group, $>C=C=C<$; the end carbons are treated as normal C_d atoms. For example, 1,2-butadiene would have the groups C_d , $C_d-(H)_2$, $C_d-(C)(H)$, and $C-(C_d)(H)_2$.

²When one of the groups is *t*-butyl, the cis correction = 4.0; when both are *t*-butyl, the cis correction = 10.0; and when there are two cis corrections around one double bond, the total correction = 3.0.

³Value is 1.2 for but-2-ene but zero for other dienes and 0.6 for trienes.

⁴The number in parentheses beside each ring is the symmetry number.

⁵For $\Delta H_{f,298}^\circ$ contributions for other ring structures, see *Chem. Rev.*, 69: 279 (1969).

⁶ N_d represents a double-bonded nitrogen in imines; $N_t-(C_a)$ represents a pyridine nitrogen. N_a represents a double-bonded nitrogen in azo compounds. For ortho or para substitution in pyridine add -1.5 kcal/g-mol per group to $\Delta H_{f,298}^\circ$.

Example: Estimation of ΔH_{f298}° and ΔS_{298}° for vanillin

#	atom (see below)	notation	ΔH_{f298}°	ΔS_{f298}°
1	1,2	(CO)-(C _B)(H)	-34.6	-
1	3	C _B -(CO)	9.7	-
3	4,5,11	C _B -(H)	3(3.30)	3(11.53)
2	6,8	C _B -(O)	2(-0.90)	2(-10.20)
1	7	O-(C _B)(H)	-37.9	29.1
1	9	O-(C _B)(C)	-22.6	-
1	10	C-(O)(H) ₃	-10.1	30.41
Σ =			-87.40	73.70

Vanillin

**TABLE B-1 - RECOMMENDED CATALYSTS FOR THE
ENDOTHERMIC CYCLIZATION-ADDITION REACTIONS**

CHEM REACTION:

Reaction type:

<i>Recomm. Catalyst</i>	<i>comments</i>	<i>ref</i>
1) Pt on Alumina	Proven to be the best aromatization catalyst. The hydroxylation state of alumina has a determining influence on the particle size of platinum crystallites, and it can also effect selectivity and the rate of production of arenes. Possible side products: iC ₆ , and 1,3,5 trans Hexatriene.	(1),(3),(4)
2) Pt on alumina & silica	A possible candidate; more frequently used for dehydrocyclization of alkylaromatics.	(3)
3) HSZM Zeolite	characterized as SiO ₂ /Al ₂ O ₃ =21, Surface area of 300 sq-m /gr, and restriction coefficient of 12	(7)

CHEM REACTION:

Reaction type:

<i>Recomm. Catalyst</i>	<i>comments</i>	<i>ref</i>
1) Mg-Cu	Good yields have been obtained when used for alkylation of mono substituted arenes with methanol.	(6)
2) X,Y Type Zeolites , HY Zeolite	An alkylation catalyst tested in the laboratories for converting phenol to O,P cresols, which may also be capable of promoting dehydrocyclization	(5), (12)
3) Cr ₂ O ₃ -Al ₂ O ₃	A toluene selective catalyst when used for cyclization of higher hydrocarbons.	(2)

CHEM REACTION:

Reaction type:

Dehydrocyclization-addition

Recomm. Catalyst***comments******ref***

- 1) Pt on Al₂O₃, Cobalt(III) alkylperoxy complexes are proven to be followed by Co(III) selective catalysts for hydroxylation of arenes

(9)

- 2) Pt on Al₂O₃, followed by molybdo- Second step does not involve a powdered catalyst, however vanadophosphoric acid it involves a transition metal complex which may be activated when introduced to UV photons.

CHEM REACTION:

Reaction type:

Dehydrocyclization-addition

Recomm. Catalyst***comments******ref***

- 1) Pt on Al₂O₃ We just might get lucky !!!

none

CHEM REACTION:

Reaction type:

Dehydrocyclization-addition

Recomm. Catalyst***comments******ref***

- 1) Pt on Al₂O₃, In the presence of hydroxyl ions, addition of the the second hydroxyl ion to the phenol ring is promoted followed by Fe(II)

(10)

- 2) Pt on Al₂O₃, Method of preparation of the catalyst is further discussed followed by Vanadium in the article referenced.

(13)

CHEM REACTION:

Reaction type:

Recomm. Catalyst**comments****ref**

1) Fe(II)

This is the reaction I am excited about!!

(10)

This type of reaction is carried out in the presence of hydrogen peroxide, H₂O₂, catalyzed by Fe(II). However, the catalyst will constantly be oxidized to Fe(III) while the reaction is in progress.

Good news:

The UV radiation reduces the catalytically inactive Fe(III) produced, to Fe(II) during the reaction, therefore continuously producing the desired catalyst. It will also break up the water molecules, present in the feed to H⁺ and OH⁻. Hydrogen ions will promote the reduction of Fe(III) ions and the hydroxyl ions will serve as a substitute for the H₂O₂ of the Fenton reagent.

REFERENCES

- [1] Dautzenberg, F.M., and Platteuw, J.C. *J. catal* 19,41 (1970).
- [2] Paal, Z. *Advances in catalysis* 29, 273-329 (1980).
- [3] Csicsery, S.M. *Advances in catalysis* 28, 293-319 (1979).
- [4] Gault, F.G. *Advances in catalysis* 30, 1-90 (1981).
- [5] Balsama, S., and Forni, L. *Appl. catalysis* 13(1), 161-70 (1984).
- [6] Bennet, J.G. Jr. EP. 129065, 27-Dec 1984.
- [7] Lopez, A.F., and Suarez, A.R. *Appl catal* 15(2) 235-45 (1985).
- [8] Mitnik, U.V., and Komorov, A.G. *Deposited doc* 1984, (Moscow chemical technology institute) Moscow, USSR.
- [9] Brazi, E., and Robin, A. *Jour. Amer. chem. soc.* 107(12) 3534-40. (1985).
- [10] Luank, S., and Sedlac, P. *React. kinet. catal. lett.* 26(1-2) 39-44, (1984).
- [11] Essam, Mohammad. *Indian patent document* 151,656. June 18, 1983.
- [12] Fraenkle, Dan. "Eur. pat. appl. 126,245. Nov. 28, 1984.
- [13] U.S. patent 544, 748. Jul. 18, 1983.

C-1

WEEK ENDING JUNE 8, 1988

An index of weekly chemical market reports is on the back cover.

Abies balsamea of Can.	lb.	15.00	-
Acetabularia 99% tanks, Int. ad. B.	lb.	.37	-
Acetic acid, higher W of West			
Acetaminophen (see N-Acetyl-p-aminophenol)			
Acetaminophen tech. packed, bgs., l.f.b.	lb.	1.29	-
Acetic acid, tech. tanks, dhd. E.	lb.	.25	-
Acetic anhydride tanks, dhd. E.	lb.	.45 1/4	-
Acetic anhydride proc. 1c higher in West			
Acetoacetic amide, dms., l.f. dhd.	lb.	1.29	-
Acetoacetyl-o-aniside, dms., l.f., dhd.	lb.	2.70	-
Acetoacetyl-o-chloranide, dms., l.f., dhd.	lb.	2.85	-
Acetoacetyl-o-toluidide, dms., l.f., dhd.	lb.	1.58	-
Acetolactone, dms., l.f.	lb.	3.33	-
Acetone tanks, dhd. E.	lb.	.25	-
Acetone Zone 2 (Calif.)	lb.	.27	-
Acetone Zone 3 (W of Rockies excluding Calif.)	lb.	.27	-
Acetone tanks, Int. ad.	lb.	.53	.54 1/4
Acetophenone (see Phenacetin)			
Acetophenone, tech., tanks, l.f.b.	lb.	.78	.85
perflume grade, extra, dms., l.f.	lb.	2.15	-
N-Acetyl-p-aminophenol, c.f., l.f.	kg		
Acetylene black, imp. 50% com- pressed, 12 1/2-lb. bgs., c.f., l.f. extra	lb.	.98	-
100% 25-lb. bgs., same base	lb.	.95 1/4	-
Acetylene tetrabromide tanks, l.f.b.	lb.	.97	-
Acetylsalicylic acid USP (see Aspirin)			
Acetylacryloyl citrate, bulk, l.f.b.	lb.	1.29	-
Acetylphenyl citrate, bulk, l.f.b.	lb.	2.08	-
Acriden tech tanks works	lb.	.52	-
Acrylonitrile sold Int. works	lb.	1.00	-
Acrylonitrile 100% base tanks, works	lb.	.74	.77
Acrylic acid, special, reg. tanks, dhd.	lb.	.67	-
tech tanks, Int. ad.	lb.	.60	-
Acrylonitrile tanks works	lb.	.39 1/4	.45 1/4
Acrylonitrile-butadiene-styrene resin, high-impact, nat., l.f., dms.	lb.	1.05	1.09
medium-impact, nat., same base	lb.	.94	.98
low-impact, nat., same base	lb.	.75	-
Adipic acid resin grade, bulk, hopper cars, Int. equald.	lb.	.57	-
bgs. 11 c.f. Int. equald.	lb.	.58	-
Agar USP, powd., 80 to 100 mesh, dms.	lb.	9.50	-
Alcohol synth C-8 to C-10, tanks, l.f.b.	lb.	.38	-
C-12 to C-13 tanks, dhd.	lb.	.57	.59
C-14 to C-15 tanks, dhd.	lb.	.57	-
C-16 to C-18 tanks, dhd.	lb.	.60	-
Aldehyde, C-6 dms.	lb.	4.10	5.70
C-7 dms.	lb.	1.95	-
C-8 dms.	lb.	4.30	6.30
C-10 dms.	lb.	4.30	5.35
Algin (see Sodium alginate)			
Alkal blue, dry, flushed, 110-lb dms, dhd.	lb.	3.72	3.83
Alkal blue proc. 1c higher W of Rockies			
Allspice Guatemalan / Honduran, bgs.	lb.	1.00	-
Jamaican bgs.	lb.	1.05	-
Allyl alcohol, tanks, l.f.b., Bayport, Tex.	lb.	.90	-
Allyl bromide 500-450 dms. 2,000 lbs. or more works	lb.	5.50	-
Allyl caproate 25-lb c.f.	lb.	3.90	4.50
Allyl chloride tanks l.f.b. works	lb.	.65	-
Allyl isocyanurate bgs.	lb.	5.40	6.90
Almond oil, nat. bitter (see Benzaldehyde)			
Almond oil, nat. bitter, NF 11 p.a.	lb.	3.50	3.60
sweet	lb.	1.24	1.50
Aloe, Cape c.f.	lb.	2.00	-
powd. c.f.	lb.	2.25	2.75
Curacao kgs.	lb.	2.60	-
powd. kgs.	lb.	3.00	-
Alum NF dms.	lb.	6.00	6.70
Alum ammonium tech gran, bgs.	lb.	35.00	-
c.f., l.f. works	lb.	55.00	-
FCC, 100 lbs.	lb.	55.00	-
Alum, potassium, tech gran bgs, c.f., l.f. works	lb.	35.00	-
FCC powd. 100 lbs.	lb.	55.00	-

Alumina, activated, gran., 100-lb. works, 40,000-lb. min. c.i., bulk, lots	821.00	-
calcined, bulk, same basis	354.00	-
100-lb. bags, same basis	360.00	-
hydrated, white, bulk, same basis	190.00	-
100-lb. bags, same basis	224.00	-
Aluminum acetates, basic, dms., l.c.l., works	3.26	-
Aluminum chloride, anhyd., each, 500-600 lb. dms., c.i., l.l., works, frt. equaled	.30	.56
bulk, same basis	.45H	.56
semi-bulk bin, same basis	.48	.53
Aluminum chloride, cornl., each, 100 lbs. dms., c.i., works	15.00	-
net dms., c.i., works	12.00	-
non-net dms., same basis	20.00	-
Aluminum formate, dibasic, liq. 8% Al_2O_3 l.l., works	.56	-
Aluminum hydrate (see Alumina, hydrated)		
Aluminum hydroxide, dried, gel. NF, 75-lb. dms., c.i., l.l., works	2.75	3.90
Aluminum metal, 99.94% or more, 50-lb. pigs., 30,000-lb. lots, frt. add.	.78	-
Aluminum oxide amorphous (see Alumina, calcined)		
Aluminum oxide, leafing grade, std., lining, 2,400 lb. lots, frt. equaled	1.40	-
linings, same basis	1.99	2.14
Aluminum phos-phosphates, purif., 100-kilo dms., l.l.	6.48	-
Aluminum powder, leafing grade, std. lining, 2,400 lb. lots, divd. lb.	3.17	-
extra fine, same basis	4.04	-
Aluminum sulfate, bags, c.i.	1.25	1.37
Aluminum sulfate, cornl., grd., 100 lb. bags, c.i., works, frt. equaled, basis 17% Al_2O_3 East and Gulf Coast	185.00	-
West Coast	217.80	-
liq., tanks, N.E. same basis	148.00	-
iron-free dry, bags, c.i. same	300.00	-
liq., tanks same basis	225.00	265.00
Aluminum sulfate USP grain, dms. lb. f.o.b. works	2.12	337
rect. l.l., same basis	1.88	-
p-Aminobenzoic acid, 1,000 kilos or more dms., f.o.b. works	9.60	10.10
2-Amino-4-chlorophenol dry and grd., 14,000 lbs. or more, frt. add. lb.	5.79	-
Aminoethyl ethanamine, tanks, frt. collect.	1.37H	-
N-Aminophthalazine, tanks, f.o.b., frt. collect.	1.05	-
2-Amino-2-ethyl-1,3-propanediol dms. 100 f.o.b. works	1.82	-

2-Amino-2-methyl-1-propanol, 96%, dms., c.l., l.l., f.o.b. works, lb.	96	-
tanks, f.o.b. works, lb.	96	-
o-Aminophenol, dms., f.o.b. Charleston, S.C., lb.	3.96	-
p-Aminophenol, dms., f.o.b. Raleigh, N.C., lb.	7.16	-
p-Aminostyrylic acid, USP, 50-40 lb. dms., l.l., lb.	18.50	-
Ammonia, anhyd., fertilizer, wholesale, tanks, divd. Midwest termi- nals, ton	165.00	170.00
tankcars, f.o.b. Gulf Coast, ton	90.00	95.00
aqueous, 29.4% NH ₃ , anhyd. base, tanks, Int. equivd. E. of Rock- land, ton	260.00	315.00
Ammoniacal liquor (see Ammonia, aqueous).		
Ammoniacal sat. galvanizing grade, bag, c.l., f.o.b. works, 100 lbs.	28.80	-
Ammoniacal sat. white (see Ammonium chloride com- pounds).		
Ammonium borates, gran., dms., c.l., works, lb.	90	-
Ammonium bicarbonate, 50 lb. per higher, Ammonium bicarbonate, 300-lb. bgs., dms., c.l., works, 100 lbs.	26.00	-
bgs., c.l., 100 lbs.	25.00	-
Ammonium bichromate, photo-etch grade, gran. 100-lb. dms., l.l., works, lb.	2.00	-
Ammonium bifluoride, bgs., l.l., works, lb.	.70	-
Ammonium bromide, dom. HF, gran., dms., c.l., l.l., f.o.b. works, lb.	1.31	-
Ammonium chloride, white, tech., fine gran., bgs., c.l., USP, gran., dms., 100 lbs.	18.00	-
Ammonium chloride, white, tech., fine gran., dms., 250-lb. bgs., f.o.b. works, lb.	7.40	53.00
Ammonium dimolybdate, approx. 95%, 24,000 lbs. or more, lb.	2.79	-
Ammonium fluoroborate, tech., dms., c.l., l.l., works, Int. equivd. E. of Rockland, lb.	5.48	-
Ammonium heptamolybdate, crys- t., dms., 24,000 lbs. f.o.b. works, lb.	1.79	-
Ammonium lauryl sulfate, tanks, f.o.b. works, lb.	8.57	-
Ammonium lignin, sulfonated, bulk, f.o.b. Houston, Ore., ton	29	35
Ammonium nitrate, dom., fertilizer grade, 33.5% N, bulk, S.E. divd., ton	72.00	-
Ammonium sulfate, tech., fine gran., 300-lb. dms., l.l., f.o.b. works, lb.	1.42	1.80
Ammonium pentaborate, gran. bgs., c.l., works, lb.	.75	-
Ammonium pentaborate powder, 50 lb. per higher, lb.	1.02	-
Ammonium persulfate, 225-lb. dms., 24,000 lbs. or more, f.o.b. works, lb.	.58	-
55-lb. bgs., same basis, lb.	.5614	-
Ammonium phosphate (see Di- and monoammonium phos- phates).		
Ammonium silicofluoride, dms. c.l., l.l., works, lb.	.304	-
Ammonium sulfate, lg. gran., bulk, c.l., works, ton	74.00	79.00
acid, c.l., bulk, f.o.b. works, ton	73.00	-
tech., bgs., c.l., l.l., f.o.b. works, ton	108.00	120.00
Ammonium sulfate, tech., 40-44% tech. 100% basis, Int. equivd. ton	46.00	-
Ammonium thiocyanide, tech (see Ammonium thiocya- nates).		
Ammonium thiocyanate, tech., crys- t., bgs., c.l., works, lb.	1.02	-
tech. soln., 50%, tanks, Int. equivd., lb.	.83	-
Ammonium thiosulfate, photographic, 80%, tanks, f.o.b. works, lb.	.13	-
Ammonium zincanyl carbonate, soln., lb.	.72	-
Amyl acetate, primary mixed isomers, tanks, divd., ton	.57	-
Amyl alcohol, primary mixed isomers, tanks, Int. equivd., ton	.464	-
Amyl crotonic acid, dms., lb.	2.35	2.50
p-Amyl, Amphenol, bulk, works, lb.	.81	1.0
Amyne oil, dms., lb.	11.50	12.00
USP, dms., lb.	10.20	4.00
Angelica root oil, bulk, lb.	700.00	-
Aniline, tanks, f.o.b., lb.	.40	-
Anise oil, dms., lb.	11.75	-

Anise seed, Egypt, bgs.	lb.	83
Santh, bgs.	lb.	94
Turkey, bgs.	lb.	82
Anise, Sicily, bgs.	lb.	80
o-Anisidine, imp. dms. dvid.	lb.	2.27
p-Anisidine, imp., cast. solid, dms., works.	lb.	1.90
Resins, same basis	lb.	2.25
Anthranic acid, pure, 98% min., dms., U. I. I. add.	lb.	1.70
Antimony metal, bulk, c.i., min. 175- dms., U. I. I. work.	lb.	3.02
Antimony metal, bulk, c.i., min. 175- dms., U. I. I. work.	lb.	1.29
Antimony oxide, high- alt. E. of P. dms.	lb.	1.50
Antimony trichloride, anhyd., solid, dms., U. I. I. work.	lb.	3.80
Apomorphine hydrochloride, NF, bot.	gms.	15.00
Apricot seed oil, 100% pure, 100% min.	lb.	2.00
Arabic gum, powdered, 100% min.	lb.	1.85
sprayed dms.	lb.	2.00
USP grade	lb.	6.75
Aromatic petroleum solvents (see Solvent, petroleum, aromatic)		
Arsenic, crude (see Arsenic trioxide)		
Arsenic, red (see Arsenic trioxide)		
Asphalt, refined, crude, 95% bulk, c.i., 100% vacuum, Wash. Ion	100.00	
Asbestine (see Talc, fibrous)		
Ascorbic acid, USP, 100 kilos, dms.	8.00	
Ash, black (see Barium sulfide)		
Asphalt, glaucous, (see Oleoresin)		
Asphalt petroleum cutback, tanks, E. Coast.	gal.	.88
emulsion, tanks, tankwagons, E. Coast.	gal.	.88
steam-refined, 40-300 penetration, tanks, tankwagons	ton	170.00
steep roofing grade, bulk tankwagons	ton	173.00
Aspirin, USP, crystal, powdered, to dms., c.i., 100%	lb.	1.95
10% starch granulation, white, 250- lb. dms., c.i., 100%	lb.	1.97
16% starch granulation, white, same basis	lb.	2.80
Freight equalized, unit, identical quantity over state from N.Y., Phila., Midland, Mich., Chicago, Louis.		
Atropine sulfate, USP, bot.	oz.	10.00
Avocado oil, dms.	lb.	4.00
Azelaic acid, tech., 50-lb. bgs., U. I. I., dvid.	lb.	1.23
Azo orange, bds., dms.	lb.	4.80
Azo yellow, 100 g. bgs., dvid. E. of P. dms.	lb.	4.40
Azo Glycol pigment, bgs., same basis	lb.	2.45

Barbitone, USP, non-stearic, one billion units of more	million tons	8.30
Barbital, NF, 50-kilo drums, dval	kilo	22.50
Barbital-sodium, NF, 50-kilo drums, dval	kilo	23.00
Barrie, dry-grd. Southern, oil-coated, orange, bgs. cl. f.o.b. minn. ls. water-grd., white, bgs. cl. f.o.b. works	lb.	06
unbleached earth-fine pigment grade c.i., f.o.b. works	ton	13
Barium carbonate, anapa, bulk, c.i., works, lt. equid.	lb.	180.00
bgs. same basis	lb.	25
photo grade, bgs. same basis	lb.	25 1/2
Barium chloride, 100-lb. drums, 1:10 dm lots, works	ton	510.00
Barium chloride, tech., crystal, bgs. c.i., works	ton	1.04
anhyd. chrome c.i. same basis	ton	470.00
Barium chloride, purif., cryst. 400-lb. drums, works	lb.	590.00
Barium monohydrate, 55-lb. bgs. c.i., lt. f.o.b. works	100 lbs.	3.78
octal, same basis	100 lbs.	48.00
Barium nitrate, 100-lb. bgs. c.i., works	100 lbs.	33.00
.....	ton	32.50

THE TERMINOLOGY OF THE CHEMICAL MARKETPLACE

a/alpha acid/acidified amorph./amorphous AMP/American melting point anhyd./anhydrous AQAC/Association of Official Agricultural Chemists a.p.s./available phos- phoric acid approx./approximately arbit./arbitrary ASTM/American Soci- ety for Testing & Materials	C./Centigrade carb./carboxy c.c./cubic centimeters CD/completely de- aerated c.f./case insurance freight c./carload cns./cans com./commercial con./concentrated cp./chemically pure cps./canneposes crys./crystalline cs./cases cns./carbons cyls./cylinders	E./East e.p./end point equid./equalized exp./expressed ext./extracted F./Fahrenheit f.s.s./free alongside ferment./fermentation f.f.a./free fatty acid f.c./free from chlorine f.p.a./free from phos- phorus fls./flavor f.o.b./free on board f.p./freezing point fl./flights	Incl./included indus./industrial legs./legs l./leaves lb./pound l.c./leaves carbonized L.I./leaves included lbs./liquids m./miles m.s.p./mixed aniline point mcg./micrograms mfrs./manufacturers mfr./minimum moit./moisten m.p./melting point N./nitrogen n./normal nat./natural neut./neutral NF/National Formulary No./number Nom./nominal	e./orthie ord./ordinary os./ounce P/phosphorus p./parts pac./package p./proof phos./phosphate photo./photography pkgs./packs ages powd./powdered precip./precipitated prod./producer pt./point pub./polymerized puri./purified redist./redistilled retu./retined res./resin resub./resublimed ret./returnable SD/specially denatured s.d./single distilled SE/Southeast sec./secondary	secs./seconds sp.g./specific gravity ship./shipment soln./solution std./standard syn./synthetic tanks./railroad tanks tech./technical tent./tentary LI./liquids tens./tensile tens./tensile of 2,000 pounds TV./television tens./tensile t.w./tennis wagons USP/United States Pharmacopeia vis./viscosity VMA/Vermont Manufacturers & Partners W/W/weight w./w/other white
--	--	--	---	--	--

NOTE: A unit ton is 1 percent of 2,000 pounds of the basic constituent or other standard of the material. Percentage figure of the basic constituent multiplied by the unit ton price shown in Chemical M-Bercentage figure of the price of 2,000 pounds of the material.

NOTE: A unit-ton is 1 percent of 2,000 pounds of the basic constituent or other standard of the material. The percentage figure of the basic constituent multiplied by the unit-ton price shown in Chemical Market Reporter gives the price of 2,000 pounds of the material.

WEEK ENDING JUNE 6, 1988

Diethyl carbonate, tank wagon, f.o.b. works	lb.	1.60
Diethyl etheraniline, CP dms., c.i., dhd.	lb.	7.18
tanks, dhd.	lb.	1.10
Diethyl etheraniline tech. 98 per b. lower.		
Diethyl oxalate, dms., c.i., f.o.b. works	lb.	1.80
Diethylene glycol, tanks, f.o.b.	lb.	.62
odorless cosmetic grades, l.i., works	lb.	
Diethyl sulfate, tanks, f.o.b.	lb.	.87%
Diethyl tetrachloride, dms., c.i., l.i., works	lb.	.59
Di-2-ethylhexyl adipate (see Diocetyl adipate)		
Diethyl toluene, 95-97% min. meta isomer, dms., l.i., f.o.b. works	lb.	2.48
N,N-Diethyl-m-toluidine, tech., liq., dms., c.i., f.o.b.	lb.	2.75
tanks, same basis	lb.	3.10
Diethylenamine, dms., c.i., dhd.	lb.	1.13
tanks, same basis	lb.	1.02
N,N-Diethylaniline, dms., c.i., l.i., f.o.b. works	lb.	1.83
tanks, same basis	lb.	1.75
Diethylenamine, tanks, f.o.b.	lb.	.96
Di-2-ethylhexyl azelate (see Diocetyl azelate)		
Di-2-ethylhexyl phthalate (see Diocetyl phthalate)		
Diethylene glycol, tanks, dhd. E.	lb.	25%
Diethylene glycol monobutyl ether, dms., c.i., Int. std. E.	lb.	.86
tanks, Int. std. E.	lb.	.57
Diethylene glycol monoethyl ether, dms., c.i., Int. std. E.	lb.	.84
tanks, Int. std. E.	lb.	.68
Diethylene glycol monomethyl ether, dms., c.i., Int. std.	lb.	.82
tanks, Int. std.	lb.	.54
Diethylene glycol monobutyl ether acetate, dms., c.i., dhd. E.	lb.	.80
tanks, dhd. E.	lb.	.72
Diethylene glycol monoethyl ether acetate, dms., c.i., Int. std. E.	lb.	.80
tanks, Int. std.	lb.	.72
Diethylenetriamine, tanks, f.o.b. works	lb.	1.80
Diethyleneterephthalic acid, pentasodium salt solution, tank - car/tanktruck, Int-equalized	lb.	.46
Diglycidyl USP, imp., bats	gram	2.80
Diglycol laurate, dms., ion lots	lb.	32%
Diglycol stearate, dms., l.i.	lb.	.82
Dihydroxy sulfide, dms.	lb.	1.10
Dihydroxypropionyl sulfone, bulk lot	kg.	48.00
Dihydroxyacetone, 50- kilo lots, works	kg.	40.00
Di-isobutylene, tanks, f.o.b.	lb.	.80
Di-isobutyl phthalate, tanks, dhd. E.	lb.	.56
Di-n-butylene, tanks, f.o.b. Houston	lb.	
Di-octadecyl phthalate, tanks, dhd. E.	lb.	.37
Di-octadecyl phthalate, tanks, dhd. E.	lb.	36%
Di-octadecyl azelate, tanks, dhd. E.	lb.	.90
Di-octadecyl phthalate, tanks, dhd. E.	lb.	.41
Di-propandiolamine, dms., c.i., Int. std.	lb.	.88%
tanks, same basis	lb.	.54%
Di-propandiolamine, dms., c.i., dhd. E.	lb.	1.17
tanks, same basis	lb.	1.07
Dimethyl 3,3'-thiodipropionate, dms., l.i.	lb.	1.88
Dil oil USP, dms.	lb.	7.00
Dimethyl anthranilate, dms.	lb.	15.80
Dimethyl ethyl carboxyl acetate, 25-lb. cans	lb.	
Dimethyl carbonates, dms., l.i., f.o.b. works	lb.	.80
Dimethyl dichlorovinyl phosphate, 55-gal dms., dhd.	lb.	3.50
Dimethyl etheraniline, anhyd., dms., c.i., dhd. E.	lb.	1.15
tanks, dhd. E.	lb.	1.07
Dimethyl ether, aerosol grade, tanks, dhd.	lb.	.38
Dimethyl phthalate, tanks, f.o.b. works	lb.	.65
Dimethyl sebacate, tanks, f.o.b. works	lb.	2.28
Dimethyl sulfate, ret. dms., c.i., f.o.b. works	lb.	.57
tanks, f.o.b. works	lb.	.46
Dimethyl sulfoxide, tanks, works	lb.	1.08
Dimethyl sulfoxide, tanks, works	lb.	.78
Dimethylsuccinimide, 50 lb. can, tanks, Int. equald., 100% basic	lb.	.87%
40% soln., tanks, Int. equald., 100% basic	lb.	.51%
anhyd., tanks, Int. equald.	lb.	.63%
N,N-Dimethylaniline, l.i., f.o.b.	lb.	54%
l.i. dms.	lb.	1.03
N,N-Dimethylformamide, dms., c.i., l.i., f.o.b. works	lb.	1.11
tanks, same basis	lb.	.57
2,4-Dinitrobenzene, iono-lots, f.o.b.	lb.	.49
Dinitrobenzine, orange loner, CP, bps., dhd. E. of Rockies	lb.	1.22
2,4-Dinitrotoluene, crystallizing at 47° F., l.i., f.o.b. Charleston, N.C.	lb.	5.20
2,4-Dinitrophenol, 250-lb. dms., f.o.b. Charleston, N.C.	lb.	.86
Dinitrotoluene, ntk, tech. f.o.b. works	lb.	1.05
2,4-Dinitrotoluene, dms., c.i., l.i., tanks, works	lb.	.30
tanks, works	lb.	1.25
Diocetyl adipate, tanks, Int. std. E.	lb.	1.20
Diocetyl azelate, tanks, dhd. E.	lb.	.87
Diocetyl phthalates, tanks, dhd. E.	lb.	.90
Diocetyl phthalates, tanks, dhd. E.	lb.	.46
Diocetyl sebacate, 99%, tanks, f.o.b. works	lb.	
1,4-Dioxane, tanks, Int. std. E.	lb.	1.28
l.i., same basis	lb.	1.13
Dipenterythritol, bps., c.i., l.i., dhd. E.	lb.	1.21
Pentapenta steam-dist., tanks, f.o.b. Fla. works	lb.	1.42
sulfate turpentene derived, tanks	lb.	.25
Dip oil (see Tar acid oil)		
Diphenyldichloro hydrochloride USP, dms., 1,000-kilo lots, dms.	kg.	.25
Diphanyl, 99.9%, bps., c.i., l.i., works	lb.	20.00
works	lb.	34

WEEK ENDING JUNE 6, 1988

in tech 95-99% dms. 1.	lb	7.00	-
sulfone, nym amyl USP	80-oz		
lots	oz	1.25	-
sulfone hydrazide NF	cryst.		
less than 100% w/g	kg	38.25	40.25
sulfone sulfate USP cry.	dms.		
less than 1000 w/g	kg	43.00	46.25
thiohydrazide, dms. dms.	lb	.88	

Fennel oil, sweet, USP, anal.	lb.	9.00	-
Fennel seed, Egypt	lb.	.33	-
Indian	lb.	.57	.80
Ferric chloride anhyd., tech., 350-lb. drum, C.I. works	100 lbs.	\$6.00	-
Ferric chloride, 42 lb. photo grade drum, C.I. works	100 lbs.	8.10	-

G			
G test, dms. fr. acid 170% base.....	lb	2.30	-
Gastic acid, 400-450 lots.....	kilo	23.06	-
Gastic oil, dms. Egyptian.....	kilo	85.00	106.00
Gastric, extract, 100 AQAC test, dms., L.L. divd.....	lb.	1.50	1.75
125 AQAC test, dms., L.L.	lb.	1.75	1.85
150 AQAC test, dms., L.L.	lb.	1.85	1.95
175 AQAC test, dms., L.L.	lb.	1.95	2.05
200 AQAC test, dms., L.L.	lb.	2.05	2.15
225 AQAC test, dms., L.L.	lb.	2.10	2.25
250 AQAC test, dms., L.L.	lb.	2.20	2.35
275 AQAC test, dms., L.L.	lb.	2.30	2.45
300 AQAC test, dms., L.L.	lb.	2.50	2.65
German rose (see Methyl rosenating churche)			
Geranio, sy. P-92% dms.....	lb	5.50	-
net 90-92% dms.....	lb	3.25	-
syn 90-98% dms.....	lb	5.75	-
Geranium oil, Moroccan.....	lb	24.00	27.50
Bourbon.....	lb	32.00	38.00
Chinese.....	kilo	70.00	85.00
Egypt.....	lb	53.00	-
Turkish (see Pimpernee oil)			
Geranyl acetate dms.....	lb	5.44	6.00
net dms.....	lb	10.95	-
Geranyl formate, syn. dms.....	lb	8.60	-
net dms.....	lb	15.95	-
Gibsonite, g. p., bulk, c. l., f. o. b. Bo- narize, Utah.....	ton	180.00	-
selects, some basic.....	ton	180.00	-
Ginger, Cochon bags.....	lb	.65	-
Chinese sliced.....	lb	.55	.70
Ginger oil, African dms.....	lb	75.00	-
Indian.....	kilo	70.00	85.00
Ginger oleoresin, HF, botts.....	lb	30.00	-
Glabrous leaf (see Goudon sulfate)			
Glaucous acid test, 50% dms., c. l., L.L., f. o. b. works.....	lb	.50	-
tanks, some basic.....	lb	.44	-

Helictroph, dms.	lb.	8.00	8.25
Hemlock of (see Spruce oil).			
Hemlock leaves, bla.	lb.	.56	-
Hemphactor dms., c.l., l.l., frt. aild., 100% base.	lb.	4.56	-
Heptane, indust., tanks, f.o.b. Beaumont, Tex.	gal.	1.07	-
95%, tanks, f.o.b. Houston, Tex.	gal.	1.18	-
Heptanoic acid, syn., tanks, f.o.b.	lb.	.86	-
n-Heptadecanol, syn., tanks, f.o.b.	lb.	.4375	-
Hexahydrophthalic anhydride, tech. dms., l.l., f.o.b. works.	lb.	1.42	-
Hexamethylenediamine, gran. bgs., c.l., l.l., works.	lb.	.56	-
gran. dms., c.l., l.l., works.	lb.	.58	-
pdf. bgs., c.l., l.l., works.	lb.	.60	-
powd. dms., c.l., l.l., works.	lb.	.83	-
Hexane, indust., tanks, works.	gal.	1.01	1.15
95%, tanks, f.o.b. Houston, Tex.	gal.	1.12	-
n-Hexanol, syn., tanks, f.o.b.	lb.	.50	-
Hexyl alcohol, mixed isomers, tanks.	lb.	.32	-
p-Hexyl methacrylate, dms., c.l., works.	lb.	.75 ¹ / ₂	-
Hexylene glycol, tanks, dhd.	lb.	.52	-
Hexynylcaranol, USP dms., 25-lb lots or more frt. aild.	lb.	30.00	-
Homotriene hydrobromide, USP, 10- 100-oz of lbs. lots.	oz.	10.25	11.25
Homotriene methyl bromide, USP, 10- 250 oz of lbs. lots.	oz.	9.70	10.75
Horshard herb, bla.	lb.	.25	26
Hydrazine hydrate, 85%, l.l., frt. aild.	lb.	1.54	-
55-gal dms., l.l., frt. aild.	lb.	1.61	-
Hydroic acid, purif., 47%-57%, 2- cyls., f.o.b. works.	lb.	7.50	-
Hydrobromic alcohol, tech., acid. dms., c.l., f.o.b. zone 1.	lb.	.85	-
tanks, f.o.b. zone 1.	lb.	.80	-
Hydrobromic acid, 48% dms., c.l., f.o.b.	lb.	38 ¹ / ₂	-
Hydrochloric acid, anhyd. (see Hydrogen chloride)			

WEEK ENDING JUNE 6, 1986

lactamform. NF, 200-kilo dms.,	lb.	4.25	4.50
linoleoleic acid, 98% min., dms.,			
c. L. L. works,	lb.	3.00	-
iodine dms.,	lb.	25.50	-
insolubl. 50-lb. dms., 1000 lbs. or	kilo	17.50	22.00
more f.o.b. works,	kilo	13.50	18.00
iodine crude, dms.,	lb.	14.21	14.50
iodine USP,	lb.		
iodoacetylhydrazine, USP, IVI 50-			
kilo dms., 100-400 lbs., fr.	kilo	36.00	45.00
acid,	kilo		
iodoform. NF, dms., 300-lbs., f.o.b.			
works,	lb.	24.00	-
i-iodone, dms.,	lb.	18.20	-
i-iodone, dms.,	lb.	13.10	-
iodic acid whole bgs.,	lb.	25.00	-
Irish moss, bleached, prime,	lb.		
whole,	lb.	.55	.80
iron blue, alkyl-rosinates, bgs., L.C.L.	lb.	2.70	-
iron blue, dr. E.,	lb.		
iron blue, reg. bgs., L.C.L., iron blue,	lb.	2.00	2.10
same bases,	lb.		

K			
Kacin, water washed, fully cleaned, bags c.i. f.o.b. Georgia.....	ton	256.00	-
HF purd., colloidal, betaine controlled, 50 lb. bags, 5,000 lb. lots.....	lb.	34	-
Kacin, (vacuumed) No. 1 coating, bulk, c.i. f.o.b. Georgia.....	ton	\$4.00	-
No. 2 coating.....	ton	73.00	-
No. 3 coating.....	ton	73.00	-
No. 4 coating.....	ton	70.75	-
fltar, gen'l purpose, same basis.....	ton	58.08	-
determined water washed, unclassified, paint grade 1 micron avg., same basis.....	ton	182.00	-
dry-grd. airfoiled soft, same basis.....	ton	80.00	-
Karyne gum, No. 1, powdered, lots.....	lb.	2.25	-
No. 2, powdered, lots.....	lb.	1.96	-
Note rules, bags.....	lb.	46%	\$1

Lease C, red toner, (rod 53) bbls., fr. add.	6.70	-
Leucolin, anthr., 400-lb. drums	1.00	1.06
leucos, works		
pharmaceutical, 400-lb. drms. works	1.06	1.06
tech., (under 2% I.A.), 400-lb. drms., works	.84	1.04
Lead (See Oils, Fats & Waxes market report.)		
Lead oil, No. 1, drms., c.i., f.o.b.	.34	
tanks, same basis	.38	
Lead oil, extra, winter-strained, drms., c.i.	.41	-
tanks, same basis	.33	-
prime, burning, drms., c.i., same basis, Chicago	.43	-
prime, burning, tanks, same basis	.36	-
NOTE: 300 lb. and higher, except Texas, 20, and West Coast, 30, higher.		
Lead leaves, Turkish	.60	.66
Leaduric's acid, drums, f.o.b.	3.85	-
Laureic acid, com., pure bgs., c.i.	.66	.71
Laureic aldehyde (aldehyde C-12) drms.	7.75	-
n-Lauryl methacrylate, drms., c.i., U. works	1.72	-
Levander oil, Alberta, 30-32% drms.	4.00	-
Levander flowers, ord.	.66	.75
medium, bbs.	.80	.90
select, bbs.	1.10	1.19
Levander flower oil, NF, French, 40-42%, ester, ord.	9.25	13.50
spike, Spanish, drms.	15.00	22.00
Lead acetate, purif., Reka, 400-lb. drms., works	.48	-
tech., Reka, U., 400-lb. drms., works	.37	-
Lead blue, basic, sulfates, bbs., c.i., shipt, pt., f.o.b.	.87	-
Lead carbonate (see Lead white basic carbonate)		
Lead chloride, 400-lb. drms., works	3.25	-
Lead dioxide, tech., powd., 200-lb. drms., U. works	.66	.70
Lead fluoroborate, lg. conc., drms., U. works, int. aqual.	.84	.87
Lead metal, dhd.	.16	.16%
Lead monosulfate, milled, bgs., c.i., f.o.b. works	.55%	-
Lead oxide, bgs., c.i., same basis	.57%	-
Lead resinate, lg., 24% Pb, drms., int. dhd.	.83	-
Lead stearate, tech., 24% Pb, drms., U. works	.32%	-
Lead peroxide (see Lead dioxide)		
Lead red, 95% PbO ₂ , or less, bgs., c.i., works	.37	-
Lead red, 97% PbO ₂ , bgs., c.i., works	.37%	-
Lead red, 98% PbO ₂ , bgs., c.i., same basis	.37%	.40%
Lead silicate (see Lead white basic silicate)		
Lead silicochromate, bgs., c.i., works	.36	-
Lead sulfate (see Lead, blue, basic sulfate and Lead, white basic sulfate)		
Lead, white, basic carbonate, bgs., c.i., int. dhd.	.88	-
Lead, white, basic, silicate, bgs., c.i., same basis	.87	-
Lead, white, basic sulfate, bgs., c.i., same basis	.86	-
Leadline, edible, tech., bleached, non-ret. drms., L.C.I. works	.36	-
unbleached non-ret. drms., L.C.I., same basis	.34	-
edible, tech., bleached, non-ret., drms., U. works	.28	-
unbleached, non-ret., drms., U. same basis	.26	-
Lemon oil, Argentine	14.00	-
Brazil	9.00	7.00
Calif., U.S.P. drms.	6.50	9.36
Italian	11.50	-
Limonene oil, Indian, drms.	11.25	-
Guatemala, drms.	2.25	-
d-Laurene, drms., 1 kilo works	60.00	60.00
Lubricant, red, whole, bbs.	.40	.50
gran. bbs.	.70	.90
powd. bbs.	.96	-
Lignosulfonate (see under Ammonium or Sodium lignin toners)		
Lime, chemical, caustic (quicklime), bds. 50,000 lbs., works, f.o.b. chemical	39.00	45.00
Lime, chemical, hydrated, bulk, same basis	44.00	50.00
bbs., same basis	54.00	57.00
Lime, NF, purif., 100-lb. drms.	.89	-
Lime oil, dist., Mexican, drms.	6.00	-
Hessian, dist., drms.	8.00	-
expressed, drms.	17.50	-
Lime salts (see Calcium)		
d-Limonene, drms.	.70	.86
Unisol oil base de rose oil, drms., syn., 90-100% drms., works	6.35	-
Unisol oxide, syn., 55-gal. drms.	2.50	3.06
Unisol acetate oil base de rose oil, 90-92%, syn., 90-100% drms., works	7.75	-
Unisol benzoin, syn., 55-gal. drms.	18.00	21.00
Unisol cinnamate, syn., 55-gal. drms.	8.00	3.15
Unisol formate, syn., 55-gal. drms.	2.00	-
Unisol isobutyrate, syn., 55-gal. drms.	56.85	-
Unisol propionate, syn., 55-gal. drms.	7.75	6.50
Unders, 23% formulation, to distributors, drms., int. dhd.	6.50	6.56
99.9% tech., f.o.b. Niagara Falls, N.Y., 250-lb. drms., int.	3.50	-
Unalyl propionate, syn., 55-gal. drms.	9.00	-
Uندن Flowers, with leaves, bbs.	7.80	.86
without leaves, bbs.	.90	1.15
Unseed meal (see Oils, Fats & Waxes market report)		
Unseed oil, (see Oils, Fats & Waxes market report)		
Unseed oil fatty acid, dist., drms.	.60	.67
tanks	.53	.62
Uncharge, com., l., powd., bgs., c.i., works	.36%	.80
Urethane bromide, anhyd., drms., ton lots, dhd.	6.27	-
edite, same basis	4.00	-
Urethane carbonate, powd., bgs., c.i., U. dhd.	1.50	-
Urethane chloride, anhyd., ed., U. dhd.	3.52	-
edite, drms., c.i., U. dhd.	2.94	-
Urethane fluoride, drms., c.i., U. dhd.	4.80	-

Mace, East Indian, string,	lb.	4.80
Sisal #2	lb.	4.80
Magnesia, tech., light, neoprene- grade, bgs., c.i., U. works	lb.	.75
Magnesia, syn., tech., chemical- grade, bulk, c.i., U. works	ton	330.00
bagged, c.i., U. same basis	ton	365.00
de-sulfurized, bulk, same	ton	362.00
bgs., same basis	ton	408.00
Magnesia, nat., tech., heavy, 80%, 150 mesh, bulk, c.i., U., f.o.b. Mex.	ton	232.00
80%, 325 mesh, same basis	ton	265.00
Magnesium bromide, 80-b, drms., hes- sychrome	lb.	2.95
Magnesium carbonate, light, tech., bgs., c.i., U. works, fnt. equid.	lb.	.73
USP, 80-bgs., c.i., same basis	lb.	.74
USP, heavy, bgs., c.i., same basis	lb.	.83
Magnesium chloride, anhyd., 92%, flake or pebble drms., c.i., works	lb.	.12
Magnesium chloride, hydrous, 99%, flake, bgs., c.i., works	lb.	.14
Magnesium gluconate, 100-b, drms., f.o.b.	lb.	4.28
Magnesium hydroxide, NF, powd., drms., c.i., U., works fnt. equid.	lb.	.78
Magnesium lauryl sulfate, tank, f.o.b. works	lb.	.22
Magnesium metal, 98.8%, ingots, 10,000-lb. or more, f.o.b. Prepact, Tex.	lb.	1.53
die casting alloy	lb.	1.28
Magnesium nitrate, 100-b, drms., U. works, f.o.b.	lb.	.32
Magnesium oxide, USP light, bgs., c.i., works, fnt. equid.	lb.	1.85
heavy, drms., c.i., same basis	lb.	1.54
Magnesium oxide, tech. (see Magnesia)		
Magnesium phosphide, intabsc, tech. 80-b bgs., f.o.b.	lb.	1.05
Magnesium silicate (see Talc)		
Magnesium silicofluoride, bgs., c.i., U. works	lb.	.15
Magnesium stearate, bulk, U.	lb.	.96
Magnesium sulfate 10% Mg, tech., bgs., mixed c.i., 10,000-lb. min. works	100 lbs.	13.80
bulk, same basis	100 lbs.	11.50
USP, cryst., bgs., mixed c.f., 10,000-lb. min.	100 lbs.	14.35
Magnesium sulfate, 17% Mg, tech., bgs. (synthetic monohydrate) c.i., 30,000 lb. min.	100 lbs.	18.25
bulk, c.i., 100,000 lb. min.	100 lbs.	17.50
Magnesium sulfate, 17% Mg, CP U., 30,000 lb. min. works 100lbs. bulk, c.i., 100,000 lb. min.	100 lbs.	19.25
works	100 lbs.	18.50
Magnesium trioxide, USP, powd., 80- b, drms., 5,000-lb. lots	lb.	.30
USP, micronized powd., drms., 375-lb. lots	lb.	.15
Metachlor, tech., drms., U. works	lb.	1.60
Methac acid, cryst., powd., drums, 100 kilo, f.o.b.	kilo	3.25
drums, tons, f.o.b.	kilo	2.85
Methac anhydride, bgs., U. works, fnt. equid.	lb.	.55
tanica, works, fnt. equid.	lb.	.55
Methac acid, purif. and food grade, 50- b, bgs., U. c.i., dtd.	lb.	.85
Monochlor of (see Terginate oil, talerol)		
Maleic acid, drms., 1,000 kilo lots	kilo	8.05
Manganese acetate, dihydrate, drms., dtd.	lb.	.45
teranhydride, drms., U. dtd.	lb.	.45
Manganese carbonate printing ink dtd.	lb.	1.85
Manganese carbonate, tech., drms., 80-b, drms., 5,000-lb. lots	lb.	.85
Manganese carbonate, chemical grade, 48% Mn bgs., 20,000 lb. lots or more, works	lb.	1.05
Manganese chloride, anhyd., drms., 20,000-lb. lots or more	lb.	.85
Manganese dioxide, nat., African, grd., 74-76% MnO ₂ , 100-b, bgs., 1, works	ton	200.00
84% MnO ₂ , same basis	ton	250.00
Manganese dioxide, syn, cryst., bat- tery grade, 90%-92% MnO ₂ , 100-b, bgs., c.i., works	lb.	.75
chemical, ferrite grade, same ba- sis	lb.	.45
Manganese gluconate, FCC grade, 100-b, drms., f.o.b. works	lb.	3.15
Manganese hydrosulfide, dtd.	lb.	.85
Manganese hypophosphite, NF, drms., 80-b, drms., 5,000-lb. lots	lb.	.85
Manganese metal, electrolytic, No. 1 chip, bulk, c.i., works	lb.	.15
drms., c.i., works	lb.	.15
Manganese naphthalene, liq., 8% Mn, dtd.	lb.	.15

Oleum (see Sulfuric acid, fuming)	lb.	2.10	-
Oleum gum, Haiti, bgs.	gal.	8.00	-
Olive oil, edible, Spanish, drms.	gal.	9.10	-
Osain b-type	ton	12.00	-
Oxline, crs. works	ton	15.00	-
20 mesh, works	ton	15.00	-
100 mesh, works	ton	20.00	-
Optum, USP, gran. powd. 25-40 lots	lb.	125.00	-
Orange oil, expressed, USP, Calif. dms., f.o.b. plant	lb.	1.20	-
expressed Valencia, drms.	lb.	1.00	1.20
Calif. dist., crs. f.o.b. plant	lb.	.40	-
Florida, drms.	lb.	.80	.85
Brazilian	lb.	1.20	-
West Indian, bitter, NF X, crs. dms.	lb.	6.50	-
Orange peel, bitter, Haitian drs.	lb.	.38	-
Oregano, Greece, 30M	lb.	1.01	1.03
Turkey	lb.	.82	-
Mexico	lb.	.80	.85
Origonum of, Spanish, crs.	lb.	35.00	-
Orie rot, Florence, drs.	lb.	4.00	-
Vernon, drs., bulk	lb.	4.80	5.00
Vernon, drs., bulk	lb.	3.00	-
powd., bulk	lb.	4.80	5.00
Curcuy wds. red, pure, bgs.	lb.	3.25	3.35
Oxalic acid, bgs., c.I., world	lb.	.44	-
o-Oxynaphthol acid drms. works, tech.	lb.	2.55	-
Oxyquinoline base, pure, 1,000 lbs., int. sold.	lb.	8.00	-
Oxyquinoline sulfate, 100 lbs. int. sold.	lb.	4.00	-
P			
Palladium metal, works	Troy-oz.	109.00	-
Palm oil, (see Oleo, Fats & Waxes Market Report)			
Palm oil, acid, dib-dist. drms.	lb.	.31% .30	-
tenks.	lb.	.30	-
s.d., drms.	lb.	.42	.46
tenks.	lb.	.35	-
Palm kernel oil, bulk, c.I., U.S. ports	lb.	.11	.14
Palmarosa of, Indian drms.	lb.	36.00	-
Palmitic acid, 90%, tech., bags	lb.	.53	-
tenks	lb.	.51	-
Papaverine hydrochloride, NF powd., imp. bulk	lb.	56.00	-
Paprika, Hungarian, 100 AU bgs.	lb.	.84	-
Spanish, 110 AU bgs.	lb.	.87	-
Paraffin, fully-refd., 127-130 F., ASTM, tenks, nety	lb.	.29	.35
130-135 F., ASTM, tanks, nety	lb.	.33% .35	.39
140-145 F., ASTM, tanks, nety	lb.	.35	.41% .44
150-155 F., ASTM, tanks, nety	lb.	.41% .48	
slack wtd. 5% of tanks nety	lb.	.19	-
12% of tanks nety	lb.	.21	-
20% of tanks nety	lb.	.16	-
AAMP temperatures are an arbitrary 3F higher than ASTM.			
Paraffinohexahydride, 91%, Assu, bgs. c.I., U. divd.	lb.	.29% .39%	-
95% powd., bgs., c.I., U. divd. lb.	lb.	.39%	-
Paraffinohexahydride, 98%, 50-gal. drms. c.I., divd. E.	lb.	.58% .76%	-
tenks, divd. E.	lb.	.76%	-
Paraffin, ethyl, drms., int. sold.	lb.	1.75	-
Paraffin methyl (see Methyl paraffin)			
Pars lower red, bulk	lb.	3.75	-
chlorinated (red 4) bgs.	lb.	3.75	-
Patchouli of, Indonesian, drms.	lb.	20.00	22.00
Peach kernel oil, USP (see Apricot kernel oil)			
Peanut meal (see Oleo, Fats & Waxes market report)			
Peanut oil (see Oleo, Fats & Waxes market report)			
Peanut don., NF, citrus, powd., 100- 400 lots drms.	lb.	3.30	3.70
Pelargonol acid, natl. tenks, int. it. sold.	lb.	.70	-
syn. tenks, f.o.b. int. sold.	lb.	.70	-
Pencilin, potassium, non-aqueous, 200- billion-unit lots.	lb.	20.00	25.00
Pencilin, procaine, sterile 500 billion- unit lots, bulk.	lb.	36.00	-
Pennyroyal oil, drms.	lb.	5.50	-
Pentachlorophenol, 50-lb. bgs., U. f.o.b. Wichita, Kan.	lb.	.86	-
Pentazocine, tech., bgs., c.I., f.o.b. int. sold.	lb.	.71	.72
Pentazocine, di- and tri-acetates (see Dipentazocine and Tri-pentazocine)			
Pentazocine, triacetate, U. drms., f.o.b. works.	lb.	1.80	-
Peracetic acid, drms., 100 lbs. or more, int. sold.	lb.	7.00	-
Peracetic acid-sodium, drms., 100 lbs. or more, divd.	lb.	14.00	-
Perfume, tetrazol, NF, drms., 200-400 lots	lb.	32.00	-
Pepper, black, Brazilian, bgs.	lb.	1.98	-
Lampung, bgs.	lb.	2.03	-
Malabar, bgs.	lb.	2.05	-
Telcherry, bgs.	lb.	2.40	-
Perrier red Chinese Fukien rice bgs.	lb.	.85	-
Harnen, bgs.	lb.	1.00	-
Ling, bgs.	lb.	.78	-
Indian, S-4, bgs.	lb.	.68	-
Pharutan dundulids, bgs.	lb.	.43	-
Pepper, white, Muntok, bgs.	lb.	2.85	-
Peppermint essence, imp. drms.	lb.	2.85	-
Peppermint oil, Madras	lb.	13.00	14.50
Mchweri	lb.	13.00	14.00
Wilemerite	lb.	11.00	13.50
Yale	lb.	8.50	10.00
int. dist. f.o.b. works	lb.	7.00	9.00

CHEMICAL PRICES

WEEK ENDING JUNE 6, 1986

Perchloroethylene, dry cleaning grade, distr. tanks, dhd.	28 1/2	-
Indust. grade, consumers, tanks, dhd.	31	-
Pen acid, dms., 25-lb.	2.56	-
Permanent red 2B, (red 48), calcium salt, dms., 100-lb.	5.25	-
Peru balsam, same base, 100-lb.	5.25	-
Perrigan of Paraguay, 100-lb.	3.00	-
Petrolatum, USP, snow white, dms., c.i., refly.	5.75	6.25
tanks, refly.	3.75	-
USP, soft white, dms., c.i., refly.	3.10	-
tanks, refly.	3.75	-
USP, white, dms., c.i., refly.	3.70	-
Petrolatum, USP, light white, tanks, refly.	3.05	-
USP, cream, dms., c.i., refly.	3.65	-
tanks, refly.	3.30	-
USP, soft yellow, dms., c.i., refly.	3.50	-
tanks, refly.	2.95	-
USP, amber, dms., c.i., refly.	3.45	-
tanks, refly.	2.90	-
Petroleum pitch (see Asphalt, petroleum).		
Petroleum sulfonates, 60-80% sulfonic acid, cont. HW, bulk, works.	48 1/4	49
MWV, same base.	49	-
LMV, same base.	49	49 1/4
Prices for 51% sulfonic content 2¢ per lb. lower on corresponding molecular wt.		
Phenacetin USP, powd., 200-lb. dms., 100-lb. lots dhd.	2.20	-
100-lb. dms., 1000-lb. lots dhd.	2.22	2.45
p-Phenendiol, dms., c.i., f.o.b.	2.00	-
Phenobarbital, USP, dms., 500-lb. lots, f.o.b. works.	19.50	-
Phenobarbital-sodium, USP, 500-lb. lots, f.o.b. works.	27.00	-
Phendyl, sint. tanks, 100-lb. lots.	25	29
p-Phenylsulfonic acid, 65% sol'n, dms., c.i., f.o.b. works.	64	-
tanks, same base.	58	-
Phenothiazine, indust. grade, 50-lb. bags, c.i., f.o.b. works.	2.33	-
purif. grade, same base.	2.89	-
Phenyl acetate, dms., 100-lb. lots, works.	1.04	-
Phenylacetic acid, pure, 25-lb. dms., c.i., f.o.b. works.	4.50	-
di-Phenylalanine, dms., 25-lb. lots.	84.00	-
1-Phenyl-3-carboxy-pyrazolone-5, dms., 200-lb. lots dhd. E.	3.45	-
m-Phenylenediamine, crist. dms., c.i., f.o.b. works.	2.07	-
o-Phenylenediamine, faked, dms., c.i., f.o.b. works.	3.25	-
p-Phenylenediamine, faked, dms., c.i., f.o.b. works.	4.00	-
Phenylethylene hydrochloride, USP, 100-lb. lots or more.	175.00	185.00
Phenylethyl acetate, dms.	3.35	-
2-Phenylethyl alcohol, NF, dms., c.i., f.o.b. works.	2.65	3.40
o-Phenylthiourea, dms., 30,000 lb. or more, 100-lb. lots.	1.50	-
Phenylthiophenyl acetate, 25-lb. dms.	5.50	6.90
Phenylglyoxal, 95% min. dms., 100-lb. lots.	3.50	-
1-Phenyl-3-methyl-5-pyrazolone, dms., 250-lb. lots dhd. E.	1.80	-
p-Phenylphenol, dms., 100-lb. lots.	1.35	2.00
p-Phenylphenol, bgs., 100-lb. lots, or more, 100-lb. lots.	1.85	-
Phenylpropylamine hydrochloride, 100-lb. dms.	24.00	28.00
Phenylsulfate, purif. crist., dms., E.	2.75	-
tech. crist., E.	2.25	-
base, E.	2.35	-
Phosgene (toner red 90), dms., 100-lb. lots.	1.95	2.05
Phosphoric acid, 100-lb. lots, 5 to 8% phosphoric acid, 100-lb. lots.	58	67
Phosphate rock, Fla. and pebble, run of mine washed, 66-68% b.p., bulk, c.i. mines.	23.15	-
vessel, Tampa, same base, 100-lb. lots.	28.00	-
Phosphoric acid, cont. and tech. grades, 75% tanks, 100-lb. lots.	29.00	-
80% tanks, works, 100-lb. lots.	31.00	-
85% N.F. tanks, f.o.b. freight equid., 100-lb. lots.	33.50	-
Food grade prices \$2.00 above tech. grades.		
Phosphoric acid, agricultural grade, 52-54% a.p.a., tanks, 100-lb. lots.	3.10	-
unit-ton super, min. 70% a.p.a., same base, 100-lb. lots.	3.45	-
Phosphorus, white (yellow) solid dms., c.i., works, 100-lb. lots.	1.00	-
tanks, works, f.o.b. works.	31	-
Phosphorus trichloride, tanks, 100-lb. lots.	40	-
Phosphorus pentasulfide, powd., dms., c.i., works, 100-lb. lots.	50.00	-
100-lb. lots, same base.	45.00	-
Phosphorus pentoxide, dms., c.i., works, 100-lb. lots.	82	-
Phosphorus sesquioxide, dms., c.i., works, 100-lb. lots.	38	-
Phosphorus trichloride, dms., c.i., works, 100-lb. lots.	40	-
tanks, works, 100-lb. lots.	35	-
Phthalic anhydride, faked, c.i., U.S. dms., 100-lb. lots.	30	335
moist, tanks, same base.	27	305
Prices 1-1 1/2¢ per lb. higher on the West Coast.		
Phthalic anhydride, faked, works.	85	-
Phthalocyanine blue toner, red shade, 100-lb. lots.	8.10	9.50
green shade, same base.	6.40	8.50
resinated, dms., same base.	6.20	8.75

Phthalocyanine blue toner, water dispersible, 100-lb. lots.	7.05	7.75
Phthalocyanine green toner, 100-lb. lots.	8.10	10.10
Phthalocyanine green toner, resinated, 100-lb. lots.	7.45	9.20
Phthalocyanine blue toner, 100-lb. lots.	6.81	-
Phthalic acid, pure, 25-lb. dms., c.i., dry base, f.o.b. Charlotte, N.C.	2.81	-
tech. base, 25-lb. dms., c.i., dry base, f.o.b. Charlotte, N.C.	6.00	-
tech. base, 25-lb. dms., c.i., dry base, f.o.b. Charlotte, N.C.	5.00	-
Pigment green B, 100-lb. lots.	2.20	-
Pilocarpine hydrochloride, USP, dms., 100-lb. lots.	1,500.00	2,000.00
Pimento leaf oil, dms.	14.50	-
Pine oil, 80% min. alcohol content, bulk, f.o.b. works, 100-lb. lots.	47.00	53.00
dms., c.i., U.S.	91.00	54.00
p-Phenyl, perfume grade, 100-lb. lots.	1.18	-
tech. grade, 100-lb. lots.	1.18	22
p-Phenyl, perfume grade, tanks, 100-lb. lots.	1.90	-
tech. grade, tanks, 100-lb. lots.	1.90	-
Piperazine, anhyd., dms., U.S. refly.	1.80	-
E.	1.80	-
Piperazine, 36%, dms., 100-lb. lots, 100-lb. lots.	2.25	2.35
Piperazine dihydrochloride, 53%, dms., U.S. refly.	2.00	-
Piperazine hexahydrate, 44%, dms., 100-lb. lots, 100-lb. lots.	1.80	-
Piperazine phosphate, 42%, dms., U.S. refly.	1.80	-
Pipendine, 98% min. dms., c.i., U.S. refly.	6.82	-
Picramnyl butoxide dms., dhd. E.	5.00	-
Picramnyl, metal, works, Troy oz.	421.00	-
Polycarbonate resin, pellets, net, U.S. refly.	1.84	1.88
Polyester resin, unsaturated, g.p., orthophthalic, bulk, tanks, 100-lb. lots.	51	53
isophthalic, same base.	58	62
Polyethylene resin, high-density, blow molding, g.p., hopper cars, 100-lb. lots.	43	48
Injection molding, g.p., hopper cars, 100-lb. lots.	43	48
extrusion, g.p., hopper cars, same base.	47	48
wire and cable, net, hopper cars, same base.	45	48
wire and cable, black, same base.	50 1/2	57
Polyethylene resin, low-density, film, 100-lb. lots, 100-lb. lots.	38	-
clarity, film, hopper cars, 100-lb. lots.	37	-
pellet, 100-lb. lots, hopper cars, same base.	36	-
extrusion coating, hopper cars, same base.	39	42
g.p., hopper cars, same base.	38	42
Polyethylene resin, low-density, g.p., 100-lb. lots.	38	40
blown film resin, 40	43 1/2	-
cast film resin, 40	45	-
Polyethylene resin, low-density, injection molding, g.p., hopper cars, same base.	45	48
line wire, CATV, power cable, 100-lb. lots.	647	-
wire and cable, thermoplastic, high-voltage, natural color, same base.	70	74 1/2
wire and cable, XLPE, low voltage, 14% carbon black, same base.	67 1/2	72 1/2
wire and cable, jacking, black, 100-lb. lots.	587	667
Polyimide resin, USP, bulk, 500-lb. lots.	52	-
Polyisobutylene, sorbitan monooleate, dms., 20,000-lb. lots, works.	73	-
Polyacetylene, sorbitan monooleate, dms., 20,000-lb. lots, works.	73	-
Polypropylene resin, homopolymer, g.p., net, U.S. refly.	45	48
copolymer, med. impact, net, 100-lb. lots.	50	56
high impact, same base.	53	60
Colored material, 100-lb. lots, higher for each grade.		
Polyethylene resin, crist., net, hopper cars, 100-lb. lots.	48	-
impact, net, hopper cars, same base.	51	-
high heat, high impact, net, hopper cars, same base.	52	-
expandable beads (EPS), pigging grade, 1,000-lb. lots.	68	-
modified, same base.	73	-
Polyvinyl alcohol, fully hydrolyzed, medium viscosity, bgs., U.S. dhd.	1.00	1.05
partially hydrolyzed, medium viscosity, bgs., U.S. dhd.	1.05	-
Polyvinyl chloride resin, g.p., homopolymer, dispersion, bgs., U.S. dhd.	50	-
g.p. suspension, bulk, same base.	38	-
pipe grade, bulk, same base.	47	-
film grade, bulk, same base.	37	47
Polyvinyl chloride, g.p. copolymer, dispersion, same base.	58	81
g.p. copolymer suspension, same base.	45	49
Poppseed, Dutch, bgs.	38	37
Potash, caustic, 45% base, tanks, 100-lb. lots.	30	-
West Coast, 50% base, tanks, 100-lb. lots.	13.00	-
reg. grade, 50% base, tanks, 100-lb. lots.	18.08	-
reg. grade, 50% base, tanks, 100-lb. lots.	22	-
Potassium acetate, NF, gran., dms., U.S. refly.	80	1.31
Potassium bicarbonate, tech., gran., bgs., c.i., works.	31 1/2	-
Potassium bicarbonate, USP, gran., dms., U.S. refly.	72	-
Potassium bichromate, gran., 400-lb. dms., c.i., U.S. refly.	48	-

Potassium bromide, tech., dms., U.S. refly.	45	48
Potassium bromide, NF, gran., powd., bgs.	80	1.20
Potassium borohydride, powd., dms., 100-1,000 lb. works.	18.00	20.00
Potassium bromate, gran., powd., 200-lb. dms., c.i., f.o.b. works.	1.08	-
Potassium bromide, NF, gran., dms., c.i., f.o.b. works.	1.12	-
Potassium carbonate, 47% K ₂ CO ₃ , tanks, U.S. refly.	15.40	-
dms., c.i., U.S. refly.	20.65	-
calored, 99-100% K ₂ CO ₃ , hopper cars or trucks, 100-lb. lots.	32.50	-
bgs., c.i., U.S. refly.	38.40	-
Potassium carbonate, hydrated, 83-85% K ₂ CO ₃ , dms., c.i., U.S. refly.	34.80	-
works, 100-lb. lots.	33.70	-
bgs., c.i., U.S. refly.	40	46
Potassium carbonate, gran., purif., 400-lb. dms., 5-lb. lots.	40	46
Potassium chlorate, crist., dms., c.i., works.	14 1/2	-
powd., dms., c.i., works.	30	-
purif., gran., 325-lb. dms., f.o.b. shipping point.	40	-
Potassium chloride, chemical grade, 99.95% KCl, bulk, c.i., f.o.b. works.	105.00	-
USP, crist. dms.	1.12	-
USP, gran., dms.	67	-
USP, powd., dms.	67	-
Potassium chloride, agricultural (see Potassium muriate).		
Potassium chromate, purif., crist., dms., works.	57	-
Potassium citrate, NF, gran., 200-lb. dms., 100-lb. lots.	93 1/2	-
Potassium cyanide, dms., 20,000-lb. lots or more, f.o.b. works.	1.32	-
Potassium dichromate (see Potassium bichromate).		
Potassium fluoroborate, tech., dms., c.i., U.S. refly.	1.40	1.42
Potassium fluoride, anhyd., dms., U.S. refly.	1.68	-
Potassium gluconate, dms., U.S. refly.	1.45	-
Price W. of Denver, 100-lb. lots, higher.		
Potassium guano-sulfonate, 300-lb. dms., 600 lb. or more, 100-lb. lots.	2.10	-
Potassium hydroxide, tech. (see Potassium caustic).		
Potassium hydroxide, USP, pellets, 100-lb. dms., c.i., U.S. refly.	10.72	12.38
100-lb. dms., c.i., U.S. refly.	11.32	13.55
Potassium iodide, USP, gran., crist., dms., 100-lb. lots dhd.	10.72	12.38
ACS grade truckload.	11.32	13.55
Potassium-magnesium sulfate, std., bgs., works.	59.00	-
base 40% K ₂ SO ₄ and 55% MgSO ₄ , bulk, works.	67.00	-
Potassium metasilicate, gran., dms., U.S. refly.	44	-
Potassium murexide, 60-62.4% min. K ₂ O, std., bulk, c.i., U.S. refly.	44.00	45.00
Canada, 100-lb. lots.	46.00	47.00
soluble, fine std., f.o.b. 46.00	47.00	50.00
Sask., 100-lb. lots.	50.50	51.50
Potassium nitrate, 100-lb. lots.	267.00	274.00
prilled, 100-lb. lots.	277.00	284.00
tech. gran., bgs., c.i., min. 50 tons, dhd.	470.00	-
Potassium oxalate, neutral, tech., fine gran., powd., 300-lb. dms., 100-lb. lots.	2.54	-
Potassium pentaborate, gran., bgs., c.i., works.	1.01	-
dms., same base.	1.08	-
Potassium persulfate, powder 15c per lb. higher.	78	-
Potassium permanganate, free flowing, bulk, hopper trucks, 100-lb. lots.	1.09	-
50-lb. dms., same base.	1.20	-
150-lb. dms., same base.	1.17	-
Potassium permanganate, USP, 50-lb. kgs., works, c.i., U.S. refly.	1.38	-
Potassium persulfate, 225-lb. dms., 24,000 lb. or more, f.o.b. plant.	78.80	-
cvt. same base.	72.50	-
Potassium pyrophosphate, tribasic, bgs., c.i., U.S. refly.	43.75	47.25
100-lb. lots.	46.00	49.50
Potassium silicate, USP, gran., 200-lb. dms., 2,000 lb. or more, 100-lb. lots.	1.52	-
USP, powd., 300-lb. dms., 2,000 lb. or more, same base.	1.42	-
Potassium silicate, 29.8-30.2 Be., 2.5 ratio, c.i., U.S. refly.	12.40	14.25
dms., c.i., U.S. refly.	18.40	-
40-40.5 Be., 2.1 ratio, c.i., U.S. refly.	18.40	18.85
Potassium silicate, 40-40.5 Be., 2.1 ratio, dms., c.i., U.S. refly.	20.40	-
electronics grade, 30-30.4 Be., 2.1-2.2 ratio, c.i., U.S. refly.	17.10	18.65
Potassium silicate, dms., c.i., U.S. refly.	21.10	-
solid or glass, 2.15 ratio, dms., c.i., U.S. refly.	36.00	-
100-lb. lots.	36.00	-
"Ratio" indicates percentage by weight of SiO ₂ divided by percentage by weight of K ₂ O.		
Potassium silicofluoride, bgs., c.i., U.S. refly.	11 1/2	15
Potassium sulfate, NF, gran., powd., dms.	80	1.20
Potassium sulfate, U.S. refly.	2.20	3.10
Potassium sulfate, agricultural grade, min. 50% K ₂ O, std., bulk, c.i., f.o.b. works.	150.00	160.00
Potassium sulfate, gran., purif., 400-lb. dms.	88	-

Potassium tetraborate, gran., bgs., c.i., works.	1.10	-
dms., same base.	1.15	-
Potassium tetraborate powder 15c per lb. higher.		
Potassium thiocyanate, USP, crist., 225-lb. dms., 5-lb. lots.	4.01	-
tech. crist., dms., U.S. refly.	82	-
Potassium titanate, crist., c.i., works.	71 1/2	-
Potassium-titanium fluoride, tech., dms., U.S. refly.	1.24	-
Potassium permanganate, tech., dms., U.S. refly.	78	-
Prednolone USP, dms., 5 kilos or more.	1.03	-
Prednolone acetate, USP, dms., 5 kilos or more.	1.12	-
Prednolone anhyd., USP, dms., 5 kilos or more.	1.12	-
Procaine hydrochloride, USP, antibiotic grade, dms., 2,000-lb. lots, 100-lb. lots.	4.95	-
Procaine hydrochloride, USP, ampule grade, dms., 1,000-lb. lots, 100-lb. lots.	4.95	-
Propionaldehyde, tanks, f.o.b. works.	35 1/2	-
Propionic acid, syn., pure, tanks, dhd.	33	-
n-Propyl acetate, tanks, dhd.	33 1/2	-
n-Propyl alcohol, tanks, dhd.	42	-
n-Propyl glycol, 100 to 2,000-lb. lots, dhd.	11.50	-
n-Propyl-p-hydroxybenzoate, USP, 500 kilos.	10.80	-
tech. 500 kilos, f.o.b.	10.38	-
Propyl paraben (see n-Propyl-p-hydroxybenzoate).		
Propyl thiolactate, dms., 50-lb. lots or more.	55.00	-
n-Propylthiourea, dms., c.i., dhd.	75	-
Propylene, polymer grade, f.o.b. Tex. and La. Gulf Coast points.	17 1/2	-
chemical grade same base.	15 1/2	-
Propylene glycol, indust., tanks, f.o.b. U.S. refly.	43	-
USP, tanks, f.o.b. U.S. refly.	43	-
Propylene glycol monomethyl ether, tanks, dhd. E.	49	-
Propylene oxide, tanks, f.o.b. works.	47 1/2	-
Pyruvic acid, USP, powd. bgs.	1.50	-
Pumice, dom., fine, 4F-0, bgs., 100-lb. lots.	270.00	-
medium, 0 1/2-1 1/2, bgs., 100-lb. lots.	300.00	-
coarse, 2-extra coarse, bgs., 100-lb. lots.	300.00	-
Pumice, imp., Italian, fines, bgs., 100-lb. lots.	280.00	-
100-lb. lots, East Coast.	350.00	-
coarse, bgs., 100-lb. lots, East Coast.	350.00	-
Pyrazolone red (red 38), dms., works.	5.25	-
Pyrethrum flowers, fine, 0.3% pyrethrins, 100-lb. lots.	1.91	-
Pyrethrum, purif., 20% pyrethrins, dms., works.	37.50	-
Pyridine, refd., 2-deg., c.i., works.	5.90	-
dms., tanks.	5.70	-
Pyridoxine hydrochloride, USP, 100-lb. lots.	29.00	-
Pyrites, Canadian 48-50% S, mines.	4.50	-
Pyrogallic acid (see Pyrogallol).		
Pyrogallol, 100-lb. dms., 1,000-lb. lots, dhd.	13.70	-

Q

Quassia chips	lb.	57
Quinacridone maroon, dms., 100-lb. lots.	lbt.	20 75
red, dms., 100-lb. lots.	lb.	17 75
scarlet, dms., 100-lb. lots.	lb.	21 75
violet, dms., 100-lb. lots.	lb.	17 75
Quince seed, bgs.	lb.	2 00
Quinidine sulfate, USP, 1,000-lb. dms., 2,000 lb. or more.	oz.	4 20
Quinine hydrochloride, NF, 1,000-lb. dms., 2,000 lb. or more.	oz.	2 45
Quinine sulfate, USP XVIII, 1,000-lb. dms., 2,000 lb. or more.	oz.	2 30
Quinine dms., 1-lb. or equal.	lb.	1 49
Tanks, same base.	lb.	1 43

Sorbitan monobenzoate, dms., c.l., l.l.	30.000 lb. min., f.o.b. works	.76	-
Sorbitol	100 lb. min., f.o.b. works	.80	-
Sorbitol, USP, reg. 70% aqueous	dms., c.l., f.o.b. shipping point	.35	-
tanks, f.o.b. shipping point	lb.	.30	-
gran. dms., c.l., l.l. works	lb.	.70	.74
powd. dms., c.l., l.l. works	lb.	.88	.72
Soybean meal (See Oil, Fat & Wax market report)			
Soybean oil acidulated, isopropyl	95% acid, tank, New York lb.	.14	.15
Soybean oil acid, dist., clear, dms.	lb.	.43	.59
tanks	lb.	.43	.44
s.d. dms.	lb.	.47	.58
tanks	lb.	.38	.43
Spermetin leaves, Imp., bal.	lb.	2.50	2.70
Spermetin oil, Far West, native	lb.	10.75	11.50
McWest, native	lb.	12.00	-
Far West, Scotch	lb.	14.50	15.25
McWest, Scotch	lb.	14.50	15.25
Spruce oil, dms.	lb.	8.00	-
St. John's bread, edible, bal.	lb.	.29	.30
Stannic chloride, anhyd., dms.	works	2.125	-
Stannic oxide dms., works	lb.	12.00	-
Stannic chloride, anhyd., dm.	lb.	N.A.	-
Stannic oxide dms., works	lb.	N.A.	-
Stannous chloride, anhyd. dms. wks.	lb.	3.508	-
Stannous fluoroborate, liq. conc. dms.	l.l. works, frt. squared	3.49	3.51
Stannous oxide dms., works	lb.	N.A.	-
Stannous sulfate dms., works	lb.	N.A.	-
Stearic acid double pressed, bulk	lb.	.26	.39
single-pressed, bulk	lb.	.28	.37
triple-pressed, bulk	lb.	.32	.40
l.l. works	lb.	.374	-
Strontium nitrate, 50-15 bags, c.l.	100 lbs.	61.50	-
Styrene monomer, 99.9% min., l.l.	l.l., f.o.b. works	.21	-
Styrene-acrylonitrile resin, natl. bulk	f.o.b. plant	.77	-
cryst. bulk, same base	lb.	.77	.81
clear, same base	lb.	.77	.81
Succinic anhyd., dms.	lb.	2.35	-
Succinic acid, purif., cryst., dms., l.l.	lb.	2.00	2.10
frt. eqd.	lb.	-	-
Succinic anhydride, dms., c.l., l.l., f.o.b.	work	1.71	-
Sucrose, red, white, bags, c.l., f.o.b.	ref'd.	33.10	-
Sucrose acetate, 90% crystals	lb.	1.18	-
dms., l.l., divd.	lb.	1.10	-
tanks, divd.	lb.	1.18	-
100% dms., l.l., divd.	lb.	12.50	13.50
Sulfabenzamide, dms. 500 kilos	kg.	36.50	-
Sulfabenzamide-sodium, dms., 500	kg.	25.00	-
Sulfacetamide, USP, dms., 500	kg.	20.00	23.50
Sulfadiazine, USP, powd. dms., 500	kg.	53.00	-
Sulfadiazine-sodium, USP, dms., 500	kg.	40.70	-
Sulfamerazine, USP, microcrystals,	dms., 500 kilos	33.50	-
USP, powd., dms. 500 kilos.	kg.	32.00	-
Sulfamethazine-sodium, USP, powd.	dms., 50 kilos	13.00	-
Sulfamethazine, powder, dms., 500	kg.	9.50	10.00
Sulfamic acid, cryst., bags, c.l., l.l.	works	36.00	41.00
Sulfamic acid, gran., dms., c.l., l.l.	works	.36	-
Sulfanilamide, NF, reg. 1,000-lb. dms.	frt. eqd.	2.00	-
Sulfuric acid, tech., bags, l.l., f.o.b.	works	.874	-
Sulfisquinoxaline, veterinary, grade,	dms.	8.00	-
Sulfur, crude, bright, molten, cont. f.o.b.	vessels, Gulfports	150.00	-
f.o.b. L.A. refry	long-ton	125.00	-
recovered, divd. Houston	long-ton	125.00	-
as tarsol, Rotterdam	long-ton	135.00	-
f.o.b. tanks, Alberta, Canada, 15	delivery	102.00	-
dark, ex-Tampa, Fla.	long-ton	157.50	-
Sulfur, crude, 50-50 lbs. min. purify. cont.	flour, 50-50 lbs. bags, c.l. mines	13.60	-
base	100 lbs.	13.60	-
lump, same base	100 lbs.	17.50	-
Sulfur, ref'd., 99.5% min. purify. base	50-50 bags, c.l. mines	20.00	-
50-50 bags, c.l. mines	100 lbs.	26.00	-
flour, light, 50-50 bags, same base	100 lbs.	14.80	-
Sulfur, ref'd., sulfamid, NF, 100.65%	min. purify. 50-50 bags, c.l. mines	18.10	-
base	100 lbs.	22.75	-
Sulfur, rubbers, 99.5% min. purify. cont.	flour, 50-50 bags, c.l. mines	210.00	220.00
base	100 lbs.	22.75	-
fine, 98% min. passing through 325	mesh, same base	18.10	-
Sulfur dichloride, dms., c.l., works, frt.	eqd.	.24	-
tanks, same base	lb.	.174	-
Sulfur dioxide, liq. conc. 98% min.	cont. f.o.b. works	275.00	-
tanks, works	lb.	210.00	220.00
Sulfur monochloride, dms., c.l., works,	frt. eqd.	.224	-
tanks, same base	lb.	.164	-

WEEK ENDING JUNE 6, 1986

NOTE: For prices on 60 and 68 Ba., multiply by .7767 and .9319, respectively. For price of 20% turning drum, add \$3.54 to above prices and multiply by 1.046.

The first of these is the *Journal of the American Medical Association* (JAMA), which has been the most influential of the journals in the field of medicine. It was founded in 1883 and has since then published a wide range of research papers, clinical studies, and reviews. The journal is published weekly and is one of the most widely read and cited journals in the field of medicine. It is published by the American Medical Association (AMA) and is available in both print and online formats. The journal is known for its high quality and rigorous peer review process, and it is a key source of information for medical professionals and researchers alike.

Thieryl, chemical, high-purity, 99.8%, 24 000-lb. min. lot, dms. frt. squared	56	-
Thorium nitrate, purif., dms., 100-lb. lots or more, works	2.75	-
d-Threonine, dms. 10 kilos wks.	120.00	-
Thyme leaves, French, bgs.	90	98
Spanish, bgs.	88	1.10
Thyme oil, N.F. red, dms.	12.00	-
N.F. white, dms.	22.00	-
Thymol, N.F.	3.75	6.18
Thymol iodide, dms., 100-lb. f.o.b. works	52.30	64.20
Tin metal (NY composition)	N.A.	-
tin chloride, anhydrous, bgs., 20- ton lots, frt. add.	.73	.74
tin chloride, anhydrous, bgs., 20- ton lots, frt. add.	.72	.74
tin chloride, anhydrous, 30 ton lots, dry bases, frt. add.	.80	-
Non-chalking nuclei material costs 1c. per pound more.	-	-
Titanium hydride powder, electronics grade, dms.	28.50	-
Titanium isocyanide, tech., bulk, c.l.	.30	.36
Titanium sponge 99.3% free of iron, less than 5,000 lbs. f.o.b. wks.	4.85	-
200-gal cylinders c.l., same basis	.50	-
Tablets add, 2,000 lbs. or more	2.45	-
d-Tacophosphate, 67%, dms.	50.08	-
d-Tacophosphoryl acetate, 81% conc., dms.	57.48	-
d-Tacophosphoryl acid succinate, crystall. dms.	78.44	-
d-Tacophosphoryl, dms.	27.40	-
d-Tacophosphoryl acetate, USP 50-kilo dms. 1000 kilo min.	18.00	18.50
Talc, 50 lbs. order	17.00	-
Talc, bulk, same basis	7.80	6.08
Toluene, petroleum, tankers, tanks	-	-
Atlanta, Ga., divd.	1.36	-
Bayonne, N.J., divd.	1.32	-
Baytown, Tex., f.o.b.	1.37	-
Chicago, Ill., divd.	1.2823	-
Clinton, Pa., f.o.b.	1.18	-
Deer Park, Tex., f.o.b.	1.37	-
Elk, Wayne, Ind., divd.	1.34	-
Gulf Coast, spot, barges	64	68
Houston, Tex., divd.	1.22	-
Indianapolis, Ind., divd.	1.32	-
Philadelphia, Pa., divd.	1.26	-
Providence, R.I., divd.	1.36	-
Toluene di-isocyanate (mixed isomers, 80%, 2,4- and 20%, 2,6- isomers, jumbo tankers, divd.)	1.01	-
p-Toluenesulfonamide, powder, dms., Ll. works	3.55	-
m-Toluidine, tech., bulk	3.10	-
o-Toluidine, tech., liq., dms. c.l.	.72	.75
bulk, same basis	80	84
p-Toluidine, tech., cast solid, dms., d. works	1.80	1.86
Ll. works	1.70	-
basic, same basis	1.96	-
Toluidines, mixed, o-m-p, tech., liquid, c.l. f.o.b. works	1.03	-
bulk same basis	.86	-
Tolylacetate, dms., 1,000-lb. lots, f.o.b. Channah, Ohio	2.80	-
Tonka beans, Angostura, prime, 1,000-lb. lots	6.50	-
Tocopherols, dms., c.l., Ll. works	.38	-
Triglyceride gum, No. 1, ribbons, dms.	38.00	40.00
Trisect paper	12.50	15.00
Tricresyl acetate, Ll. drums, f.o.b.	.78	-
Tributyl phosphate, tanks, works	1.70	-
tributyl phosphite, tanks, works	1.65	1.77
Tributylamine, dms., c.l., divd.	1.38	-
tanks, same basis	1.33	-
Trichloroacetic acid, tech., 300-lb. dms. c.l. f.o.b. works	.94	-
USP 100-lb dms. frt. add.	.99%	-
1,2,4-Trichlorobenzene, pure, tanks, divd.	.81%	-
1,1,1-Trichloroethane, tanks, com- sumers, divd.	.40%	-
1,1,2-Trichloroethane, tanks, f.o.b. works	.42	-
Trichloroethylene, tanks, divd.	.38%	-
Trichloroethylene, acid, dms.	1.23	-
Trichlorophenylacetic acid, (see 2,4,5-T)	-	-
Trichlorine crystals, 65%, solid, non-rel. dms., 1,500-lb. lots, divd.	1.38	-
Tricresyl phosphate, tanks, f.o.b. works	1.58	1.88
Tridecyl alcohol, mixed isomers, tanks, divd.	.87	-
Trifluorobenzene, 65% tanks, divd.	.48	.48
99% tanks, same basis	.48	.48
Trifluoromethane lauryl sulfates, tanks, f.o.b. works	.27%	-
Trifluoromethane, dms., c.l., divd.	1.33	27%
tanks, same basis	1.03	-
Trifluoromethane, Ll. drums, f.o.b. works	1.88	-
Trifluorophosphate, tanks, divd.	1.18	-
Trifluorophosphoryl, tanks, f.o.b. Gulf	.47	-
Trifluorophosphoryl diphenylphosphite, tanks f.o.b. works	.29%	-
40-60% tanks, 100% bases, frt. squared	.36	-
Trifluorostyrene tanks, frt. squared	1.43	1.46
Tri-n-butyl trimethylate, f.o.b. works	.51	.58
Tri-n-butylamine, tanks, divd.	.48	-
Tri-n-butylamine, dms., c.l., frt. add. E	.57%	-
Trimethylamine, anhyd., tanks, frt. squared, 100%.	.54%	-
25% tank, tanks, frt. squared, 100% bases	.63%	-
40% tank, tanks, frt. squared, 100% bases	.58%	.57
Trimethylolpropane bgs. c.l., Ll. divd.	.73	-
Trimethylolpropane (triacrylate, Ll. dms. f.o.b. works	1.50	-
Triphenylmethanol, tanks, frt. add. E	1.00	-
Triphenyl phosphite, dms., Ll., frt. squared	1.64	.78
Tripropylene glycol, tanks, frt. add. E	.84	-
Tri-(n-butylphenyl)nitromethane, solid, Ll. works	.805	-
Trisodium phosphate (see Sodium phosphate, tribasic)	-	-
Tri-n-propylamine, dms. 25-kilo lots	62.00	65.00
Tri-n-propylamine, dms. 25-kilo lots	42	43
Tungstic acid 92.14% dms. 1,750- 8,000 lbs. works	12.85	-
Yumurtic, Alcepley	.95	1.10

	1955	1956
Valearian root, Belgian, bgs.	1.05	2.25
Indian, bgs.	.46	—
Verendrum oxytrichloride, 3,000 lb. cyls., works.	5.00	5.10
Verendrum peroxide, tech., gram, per lb. of V_2O_5 , 550-lb. drms., works.	4.10	4.94
fused or flake, per lb. V_2O_5 , 550- lb. drms., works.	3.35	3.85
Vershyde brown, biops., U. H. aquad.	27%	—
Vershyde brown, Mesitogasecar	37.00	—

Vitamin USP, dms., Lab. works	lb.	6.25	-
Imp. dms.	lb.	4.75	5.00
Vitaminol A	lb.	84	-
Vinylalyl acetate, dms.	lb.	60.50	-
Vinylalyl acetate, dms.	lb.	53.00	-
Vinylalyl acetate, dms.	lb.	14.00	17.00
Vitamin	lb.	23.00	24.50
Vitamin	lb.	31.00	-
Vitamin blue toners, molybdenum, PMA	lb.	8.20	6.30
dms.	lb.	10.40	-
Vitaminized, PTA, dms.	lb.	28	-
Vinyl acetate monomer, tonics, dms.	lb.	39	-
Vinyl chloride monomer, polymer	lb.	20	-
grade, tonics, f.a.b.	lb.	20	-
Vinyl ether, USP, anesthetic, 75-cc.	lb.	1.50	-
bot., hospital.	lb.	7.81	-
2-Vinylpyridine U, dms. works	lb.	7.81	-
tonics, works	lb.	27	7.30
Vinyltoluene, bulk, f.a.b.	lb.	27	7.30
Vitamin A, synthetic, dry, phar., 500,000	lb.	31.00	-
A units per gm., 50-100,000	lb.	31.00	-
A, 10, in oil, phar., 100,000	lb.	31.00	-
units per gm., 10-100,000	lb.	31.00	-
Vitamin A, acid grade, 650,000 units	lb.	18.70	23.50
per gm.	lb.	18.70	23.50
Vitamin B ₁ (see Thiamine hydrochloride)	lb.	-	-
Vitamin B ₁ (see Riboflavin and Yeast)	lb.	-	-
Vitamin B ₂ , crystal, non-sterile, USP	lb.	-	-
(cyanocobalamin), yeast, 50-	lb.	-	-
grams, lots, 50-	lb.	-	-
Vitamin B ₂ , 1% solution of	lb.	6.00	8.75
(cyanocobalamin) USP with	lb.	-	-
disodium phosphate, 25-100 dms. lots.	lb.	10.75	12.75
Vitamin B ₂ , 0.1% solution of	lb.	-	-
B ₂ (cyanocobalamin) USP with	lb.	-	-
manned, 25-100 dms. lots.	lb.	15.80	-
Vitamin B ₂ , cobalamin concentrate	lb.	-	-
with manded, 1,000 mcg. per	lb.	-	-
gram dms.	lb.	18.45	-
Vitamin B ₂ , 1% Vitamin B ₂ USP,	lb.	-	-
absorbed on resin, 5-100 dms.	lb.	-	-
500-grams lots, 100 dms. per gram activity	lb.	15.80	-
Vitamin B ₂ , 1% cobalamin concentrate,	lb.	-	-
KF, absorbed on resin, 5-100	lb.	-	-
dms. lot, 100 dms. per gram activity	lb.	15.40	-
Vitamin B ₂ , 1% cyanocobalamin in	lb.	-	-
gelatin, 2.5-100 dms., 100	lb.	-	-
dms. lot, 100 dms. per gram activity	lb.	15.40	-
Vitamin C (see Ascorbic acid)	lb.	-	-
Vitamin D (see Cholecalciferol)	lb.	-	-
Vitamin D ₂ (see Cod liver and Fish liver oil)	lb.	-	-
Vitamin E (see α -Tocopherol and Wheat germ oil)	lb.	-	-
Vitamin H (see Biotin)	lb.	-	-
Violet methyl toner (see Methyl violet toner)	lb.	-	-

Wheatn 0.5%, dms., ton less, Int. std.		
New York or Chicago	lb.	75
Wheat germ oil, cold-pressed	gal.	18.50
Wheat germ oil, cold-pressed	gal.	14.00
White precipitate, USP, power, 100 lb.		
dms., f.o.b. works	lb.	7.882
Whiting (see Calcium carbonate)		
Whitepine oil, syn. (see Methyl eucalyptol)		
Which hazel bark, dms.	lb.	1.25
leaves, dms.	lb.	1.50
401 mesh, bgs., c.i. works	ton	134.00
325 mesh, bgs., c.i. works	ton	117.00
high speed ratio, bgs., works ton		184.00
Wollastonite, U.I., f.o.b., producing plant, general grade	ton	200.00
325 mesh	ton	140.00
400 mesh	ton	180.00
1250 mesh	ton	500.00
Wood grease, USP (see Linseed)		
Wormseed oil (see Chenopodium oil, NF)		
Wormwood oil, cine	lb.	31 00
		38.00
Xanthan gum, food 300-lb. dms., f.o.b. works	lb.	5 85
ind. grade, same basis	lb.	4.54
		6.25

Yarn, yarn, 25-lb. cone.	lb.	2.81
Yeast, pure brewer's, disarmed, NF, Sac-	lb.	
charomyces, U. f. o. b. works	lb.	1.10
Yerba, santa leaves, bla.	lb.	2.40
Yerba, santa, tops	lb.	26.50
Ylang-ylang oil, extra grade	lb.	23.93
grade 1	lb.	19.08
grade 2	lb.	15.90
grade 3	lb.	13.04

Zinc, bgs., 2,000-lb. lots.....	lb.	7.50
Zinc acetate, NF, drms.....	lb.	1.00
tech., dihydrate, bgs., 1-l. works.....	lb.	1.60
Zinc borate, tech., 43% ZnO, 37% B ₂ O ₃ , 50-lb. bgs., 20,000-lb. U. works.....	lb.	
appt., 37% ZnO, 49% B ₂ O ₃ , 250-lb. drms., 20,000 lb. 1 to 6 wks. 1b.	lb.	.58
Zinc chloride, USP, gran. drms.....	100 lb.	.89
Zinc chloride, tech., each, 50% tartrate, 1-l.b. Cleveland, Ohio.....	100 lb.	9.79
Concord, N.C.....	100 lbs.	10.23
Freemont, Tex.....	100 lbs.	20.20
Old Bridge, N.J.....	100 lbs.	20.20
65 degree, same base Cleveland, Ohio.....	100 lbs.	27.90
Concord, N.C.....	100 lbs.	27.90
Old Bridge, N.J.....	100 lbs.	27.90
70 degree, same base Cleveland, Ohio.....	100 lbs.	29.70
Concord, N.C.....	100 lbs.	29.70
Old Bridge, N.J.....	100 lbs.	29.70
72 degree, same base Cleveland, Ohio.....	100 lbs.	31.20
Concord, N.C.....	100 lbs.	31.20
Old Bridge, N.J.....	100 lbs.	31.20
Zinc chromate, bgs., (dvd.).....	lb.	1.12
Zinc cyanide, drms., c.l.....	lb.	1.65
Zinc dust pigment type 1 & 2, drms., c.l. 1-l.b. plant.....	lb.	.59
Zinc ethylenediamine tartrate acid, 8.4% Zn, ammonium salt soln., L.C., 1-l. 1-l.b. works.....	lb.	.54
8% Zn, ammonium salt soln., L.C., 1-l. 1-l.b. works.....	lb.	.48
Zinc fluoroborate, c.l. conc., drms., 1-l. works, 1-l. squalid.....	lb.	.27
Zinc metal, high grade, drms.....	lb.	.93
Zinc naphthalene, c.l. 8% Zn, drms., (dvd.).....	lb.	.35
Zinc nitrate, tech., 1-l. 300-lb. drms., 1-l. work.....	lb.	.34
Zinc oxide photo conductive, bgs., c.l. 1-l. std.....	lb.	.47%
Zinc oxide, USP 50-lb. box, c.l. 1-l. std.....	lb.	.46%
Zinc oxide pigment, American process, lead-free bgs., c.l. 1-l. std.....	lb.	.40
Zinc oxide pigment, French process, regular, bgs., c.l. 1-l. std.....	lb.	.41
Zinc phenolphthalein, purif. gran., 250-lb. drms., 1-l. 1-l. std.....	lb.	1.82
Zinc pyridithione, 48% dispersion, drms., 1-l.b. works.....	lb.	8.50
Industrial grade.....	100 lbs.	14.50
Zinc resinate precip. 7.2-7.8% Zn, drms., 1-l. std.....	lb.	.46
Zinc silicofluoride, drms., c.l., 1-l., works.....	lb.	.17
Zinc stearate, USP, bulk, 1-l. std.....	lb.	.32
Zinc sulfide, gran., monohydrate, 1-l. dust grade 36% Zn, bgs., c.l. works.....	100 lbs.	28.50
agricultural grade powder, bulk, same base.....	100 lbs.	22.50
Zinc yellow (see Zinc chromate).		
Zinc-zincum chloride, bgs., c.l. works.....	lb.	.42
Zinc undecylenate, drms., works.....	lb.	6.87
Zinc-formaldehyde sulfoximate, heat 200-lb. drms., 1-l. std.....	lb.	7.08
Zincum gran. bgs., bulk, c.l. works, 1-l. Zincum milled bgs., 200 and 325 mesh, c.l. works.....	100 lb.	186.00
Zincum acetate soln. 75% ZnO, drms., 30,000 lb. m. works.....	lb.	97
22% ZnO, same base.....	lb.	.78
Zincum hydride, powd., electronic grade, drms., works.....	lb.	.31
Zincum oxide, powd., coml., drms., 2,000 lb. m. works.....	lb.	4.25
electronic, same base.....	lb.	7.25
insulating, stabilized, 325°F same base.....	lb.	3.31
insulating, unstabilized, 325°F same base.....	lb.	3.56
drms., stabilized, 30F same base.....	lb.	2.82
Zincum cyanide, c.l. 5-ton lots, works.....	lb.	.91

.APPENDIX D.

Top 50 Chemicals Production Turned Back Up in 1987

Output increased 5.5% after slipping slightly in 1986 as organic chemicals moved up 5.5% and inorganic chemicals rebounded to rise 4.9%

Marc S. Relsch, C&EN New York

A surge in exports of chemicals last year helped boost U.S. chemical output dramatically. According to C&EN's annual ranking of the 50 largest-volume chemical products, total production in that group in 1987 increased 5.5% from 1986 to 567 billion lb. This marked a dra-

matic improvement over 1986 when total production of the Top 50 chemicals dropped 0.9% from the previous year.

The recovery was all the more dramatic considering that last year imports continued to grow, but at a much slower rate than they did in previous years. A decline in the value of the dollar helped make U.S. chemicals more attractive to overseas purchasers, and thus U.S. export volume increased markedly.

Demand in the U.S. markets for chemicals did little to produce the dramatic improvement in overall chemical production. The housing market was virtually flat. The auto market was down moderately. How-

ever, the U.S. farm economy began to stir and showed some small signs of improvement.

Among the largest-volume chemicals produced in the U.S., sulfuric acid continued to reign as king, with production last year totaling 77.5 billion lb, a 5.2% increase from 1986. Nitrogen continued to rank a distant second, with production at 47.3 billion lb, down 2.7% from 1986.

Organic chemicals, comprising 36% of the Top 50, made a good showing, but that segment of the Top 50 did not do so well as it did in 1986. Production of those substances increased 5.5% from 1986, compared with a 9.3% year-to-year increase in 1986. Between 1982 and

About the Top 50 list of chemical products

Government data, trade association figures, and industry estimates all go into preparing C&EN's annual list of the Top 50 chemical products, ranked by production volume. The federal government is relied upon most heavily, but when government figures are not available, other sources, primarily trade associations, are used. Industry sources and C&EN estimates are used only when other data are lacking.

Government data are not always accurate—they are only as good as the information that individual companies report. But they are an objective measure of production extending back many years. Therefore, relatively accurate indications of growth can be made on a consistent basis.

At this time of year, C&EN has access only to preliminary reports of production for 1987. When the government and trade associations issue their final reports, the outcome can be changed, sometimes dramatically. As a result, the production figures for ear-

lier years that appear in the table on page 31 are different in some cases from those published in last year's Top 50 report. The final reports also can affect the rankings of chemicals. The 1986 rankings of 16 chemicals listed in the table, for example, are different from those in the similar table published by C&EN a year ago (C&EN, April 13, 1987, page 20).

The list itself covers production figures for the U.S. and includes chemicals produced for export. Candidates for the list include all basic, intermediate, and chemically homogeneous finished products. These range from chemical building blocks like ethylene and propylene to downstream products like vinyl acetate and ethylene glycol.

The roster also includes basic inorganic chemicals, but does not include what C&EN considers to be minerals, such as salt, gypsum, and sulfur. Lime is included because it is processed and has many chemical and industrial

applications. Refractory (dead-burned) dolomite is excluded in the calculations of lime production.

In the organics list, such petrochemical feedstocks as ethane, butane, and propane are excluded arbitrarily because they are considered to be products of oil companies and because they have many nonchemical uses.

There are other gray areas besides lime and petrochemical feedstocks. For example, the basic aromatics—benzene, toluene, and xylene—are included. Admittedly, this makes the list less pure chemically.

Production figures are published by the government and trade associations in a variety of units—millions of pounds, thousands of tons, billions of cubic feet, millions of gallons, metric tons. To provide an accurate ranking and to make comparison of production volumes easier, C&EN not only gives production in common units but converts production to billions of pounds as well.

Top 50 chemicals production totaled 567 billion lb in 1987

Rank			Billions of lb		Common units ^b		Average annual growth			
1987	1986 ^a		1987	1986	1987	1986	1986-87	1985-86	1982-87	1977-87
1	1	Sulfuric acid	77.49	73.64	38,748 tt	36,822 tt	5.2%	-7.1%	3.1%	1.1%
2	2	Nitrogen	47.32	48.62	653 bcf	671 bcf	-2.7	2.4	6.2	8.5
3	4	Ethylene	34.99	32.81	34,992 mp	32,811 mp	6.6	9.9	7.4	4.5
4	6	Ammonia	32.32	28.01	16,161 tt	14,005 tt	15.4	-19.1	0.5	-0.3
5	3	Oxygen	32.28	33.03	390 bcf	399 bcf	-2.3	1.5	2.2	0.2
6	5	Lime ^c	30.36	29.67	15,182 tt	14,834 tt	2.3	-6.1	1.5	-2.2
7	7	Sodium hydroxide	23.03	21.38	11,517 tt	10,691 tt	7.7	-1.9	4.2	0.9
8	8	Chlorine	21.93	20.87	10,963 tt	10,436 tt	5.0	0.4	3.6	0.5
9	9	Phosphoric acid	20.95	18.41	10,473 tt	9,206 tt	13.8	-12.5	4.9	2.8
10	10	Propylene	18.53	17.34	18,527 mp	17,343 mp	6.8	16.5	8.1	6.3
11	11	Sodium carbonate ^d	17.78	16.88	8,891 tt	8,438 tt	5.4	-1.8	2.6	5.5
12	14	Urea ^e	14.86	12.06	7,431 tt	6,029 tt	23.3	-9.7	2.8	6.6
13	13	Nitric acid	14.20	13.12	7,102 tt	6,562 tt	8.2	-10.9	-0.8	-0.9
14	12	Ethylene dichloride	13.81	14.53	13,805 mp	14,529 mp	-5.0	20.1	12.6	5.6
15	15	Ammonium nitrate ^f	12.83	11.11	6,416 tt	5,556 tt	15.5	-18.0	-2.0	-1.1
16	16	Benzene	11.67	10.23	1,585 mg	1,389 mg	14.1	8.9	8.5	1.1
17	17	Carbon dioxide ^g	9.84	9.77	4,922 tt	4,883 tt	0.8	-2.3	5.2	9.6
18	18	Ethylbenzene	9.41	8.92	9,409 mp	8,915 mp	5.5	20.7	7.2	5.0
19	19	Vinyl chloride	8.23	8.42	8,230 mp	8,415 mp	-2.2	-11.1	10.9	3.8
20	21	Terephthalic acid ^h	8.11	7.68	8,110 mp	7,684 mp	5.5	18.4	10.9	1.2
21	20	Styrene	8.09	7.84	8,090 mp	7,838 mp	3.2	2.8	6.4	2.5
22	22	Methanol	7.29	7.33	7,292 mp	7,327 mp	-0.5	46.5	-0.7	1.6
23	25	Toluene ⁱ	6.73	5.82	928 mg	802 mg	15.7	14.7	5.4	-0.7
24	24	Formaldehyde ^j	6.08	5.89	6,077 mp	5,885 mp	3.3	5.0	4.8	1.1
25	26	Xylene	5.72	5.55	795 mg	771 mg	3.1	4.5	3.9	1.0
26	23	Ethylene oxide	5.62	5.94	5,623 mp	5,943 mp	-5.4	9.4	2.4	3.0
27	27	p-Xylene	5.15	5.00	5,152 mp	4,995 mp	3.1	4.5	9.9	5.9
28	28	Hydrochloric acid	4.99	4.83	2,495 tt	2,413 tt	3.4	-14.0	0.4	-0.2
29	29	Ethylene glycol	4.52	4.76	4,515 mp	4,759 mp	-5.1	13.9	0.9	3.1
30	30	Ammonium sulfate	4.37	4.17	2,183 tt	2,086 tt	4.7	-0.3	4.3	0.8
31	31	Cumene	4.15	3.70	4,151 mp	3,695 mp	12.3	10.5	8.6	4.3
32	41	Methyl tert-butyl ether ^k	3.37	2.24	3,372 mp	2,237 mp	50.7	18.3	32.6	nm
33	33	Phenol ^l	3.24	2.92	3,241 mp	2,921 mp	11.0	5.2	9.9	4.3
34	32	Acetic acid	3.23	2.93	3,227 mp	2,931 mp	10.1	1.2	3.3	2.7
35	35	Butadiene ^m	3.00	2.59	3,002 mp	2,593 mp	15.8	10.8	9.4	-1.5
35	34	Potash ⁿ	3.00	2.65	1,360 tmt	1,202 tmt	13.1	-7.3	-5.3	-4.6
37	35	Carbon black	2.72	2.59	2,724 mp	2,585 mp	5.4	0.5	3.4	-1.0
38	37	Propylene oxide	2.61	2.48	2,608 mp	2,480 mp	5.2	3.3	9.2	3.6
39	39	Acrylonitrile	2.55	2.31	2,551 mp	2,314 mp	10.2	-1.4	4.6	5.3
40	40	Vinyl acetate	2.51	2.25	2,505 mp	2,249 mp	11.4	6.5	6.0	5.4
41	38	Aluminum sulfate	2.45	2.44	1,223 tt	1,222 tt	0.1	-3.6	1.2	0.2
42	42	Cyclohexane	2.25	2.07	2,250 mp	2,071 mp	8.6	25.0	12.0	0.3
43	43	Acetone	2.08	1.94	2,082 mp	1,936 mp	7.5	8.3	4.2	1.1
44	44	Titanium dioxide	1.90	1.86	952 tt	931 tt	2.3	8.3	7.7	2.9
44	45	Sodium silicate	1.90	1.57	948 tt	786 tt	20.6	9.0	7.4	2.4
46	47	Sodium sulfate ^o	1.61	1.55	805 tt	775 tt	3.9	-6.3	-1.4	-4.2
47	48	Adipic acid	1.57	1.52	1,572 mp	1,522 mp	3.3	4.7	6.3	0.4
48	46	Calcium chloride ^p	1.55	1.56	773 tt	780 tt	-0.9	-2.6	-2.0	-4.3
49	49	Isopropyl alcohol	1.31	1.28	1,313 mp	1,275 mp	3.0	3.2	-1.0	-3.8
50	50	Caprolactam	1.16	0.85	1,161 mp	850 mp	36.6	-21.5	7.9	4.1
TOTAL ORGANICS			201.85	189.17			5.5%	9.3%	7.1%	2.1%
TOTAL INORGANICS			364.83	347.74			4.9%	-5.7%	2.3%	-0.1%
GRAND TOTAL			566.68	536.91			5.5%	-0.9%	4.8%	1.0%

^a Revised. ^b tt = thousands of tons, bcf = billions of cubic feet, mp = millions of pounds, mg = millions of gallons, tmt = thousands of metric tons. ^c Except refractory dolomite. ^d Natural synthetic. ^e 100% basis. ^f Original solution. ^g Liquid and solid only. ^h Includes both acid and ester without double counting. ⁱ All grades. ^j 37% by weight. ^k Production data for earlier years available. ^l Synthetic only. ^m Rubber grade. ⁿ K₂O basis. ^o High and low purity. ^p Solid and liquid. nm = not meaningful.

Business

1987, production of organic chemicals among the Top 50 increased 7.1% a year. Over the long term, however, their output has grown at a much slower pace. Between 1977 and 1987, output of organic chemicals among the Top 50 has grown only at a rate of 2.7% a year.

However, the inorganic segment of the Top 50 made a dramatic recovery in 1987. Production increased 4.9%, following a 5.7% decline in 1986. Within that segment, output of the major fertilizer chemicals was up—including phosphoric acid, up 14%; urea, up 23%; and potash, up 13%. Other big-volume inorganics also showed improvements. Nitric acid production increased 8.2%, and ammonium nitrate production increased 16%.

Of the Top 50 chemicals, production of 42 of them increased in 1987. By contrast, production of only 19 increased in 1986. Although total production of the Top 50 chemicals, at 567 billion lb, was up 5.5% from 1986, it was down 1.4% from the Top 50 historical high of 575 billion lb in 1979. The 1987 production level was 22% higher than the decade-low production level of 464 billion lb registered in 1982.

For the Top 50 group, composite output increased at a compound annual rate of 4.8% between 1982 and 1987. However, indicative of long-term growth, Top 50 production grew at an average annual rate of 1.3% between 1977 and 1987.

As a group, organic chemicals outperformed the inorganics, however. Showing the largest production increase on the organics side was methyl *tert*-butyl ether. Production of that chemical was up 51% to 3.37 billion lb. The gain pushed that chemical to 32nd place in 1987 from 41st place in 1986.

Next in terms of percentage gains among the organics was caprolactam, up 37% to 1.16 billion lb. The increase gained the chemical's entry into the Top 50 at 50th place in 1987. It did not appear on the list for 1986 (C&EN, April 13, 1987, page 20).

Also up appreciably in terms of percentage gains were urea, with output up 23% to 14.9 billion lb following a 9.7% decline in 1986; butadiene, up 16% to 3.0 billion lb; toluene, up 16% to 6.73 billion lb; and benzene, up 14% to 11.7 billion lb.

Not all the organics gained, however. The biggest decreases came from ethylene oxide, whose production was down 5.4% from the previous year, and ethylene glycol, down 5.1%.

For inorganic chemicals, 1987 production was led by sodium silicate, up nearly 21% from 1986 to 1.90 billion lb. Sodium silicate also led the pack in production growth in 1986, when production grew 9.0%. Also showing very good production increases in 1987 were ammonium nitrate, up 16% to 12.8 billion

lb following an 18% decline the previous year; and ammonia, up 15% to 32.3 billion lb following a 19% decline in 1986.

Other large-volume inorganic chemicals that had large swings in production last year were phosphoric acid, up 14% to nearly 21 billion lb, following a 13% decline a year earlier; and potash, up 13% to 3.0 billion lb in 1987 following a 7.3% decline in 1986.

Over the past five years, 43 of the Top 50 chemicals have registered increases in production. Leading in individual increases over the five-year span was methyl *tert*-butyl ether, whose production grew at an annual compound rate of almost 33%. Next were ethylene dichloride, up about 13% annually for the five-year period; cyclohexane, up 12% a year; vinyl chloride, up 11% a year; and terephthalic acid, also up 11% a year.

On the downside of production, leading the products with the largest individual decreases over the five-year period was potash, whose output fell at a compound annual rate of 5.3%. Other chemicals registering decreases over the five-year period were ammonium nitrate, down 2.0% a year; calcium chloride, also down 2.0% a year; sodium sulfate, down 1.4% a year; and isopropyl alcohol, down 1% a year.

Looking at growth over the past 10 years, 39 of the Top 50 chemicals have recorded overall production

Production of Top 50 chemicals is on the upswing

increases. The best long-term performer has been carbon dioxide, whose output grew at an annual compound rate of 9.6% during this period. Production of nitrogen also grew at a remarkable rate of 8.5%. Other chemicals with sizable growth rates were urea, up 6.6% a year, and propylene, up 6.3% per year.

Potash registered the greatest average annual rate of decline over the past 10 years. Its output has fallen at a compound annual rate of 4.6%. Other products with shrinking output were calcium chloride, down 4.3% a year; sodium sulfate, down 4.2% a year; isopropyl alco-

hol, down 3.8% a year; and lime, down 2.2% a year.

The group comprising the 10 largest chemical commodities in terms of production was the same in 1987 as it was in 1986. Some commodities within that group, however, switched places in ranking. Ethylene, for example, which ranked third in volume in 1987, was fourth in terms of production in 1986. Ammonia moved up to fourth place from sixth place a year earlier. Oxygen moved down to fifth place from third place in 1986, and lime slipped a notch to sixth place after ranking fifth place a year earlier.

There was a good deal of reshuffling throughout the rest of the list. In all, 21 of the chemicals either moved up or down in the rankings.

Besides the rise for ethylene and ammonia and the fall of oxygen already noted, there were other significant changes. Some upward movements included methyl tert-butyl ether, with its 51% increase in production in 1987, which jumped to 32nd place from 41st in 1986. Other large changes included urea, which jumped to 12th place from 14th, and toluene, which moved up to 23rd place from 25th.

Polymer output records fifth consecutive annual rise

Overall production of plastics, synthetic fibers, and synthetic rubber polymers reached a record high of 57 billion lb in 1987. The growth marked the fifth consecutive annual increase for the product group, lifting the output total 9.1% above the 1986 record.

Plastics, which account for more than two thirds of the polymer group production totals, continued their buoyant growth in 1987. Production was up 9.9% from 1986. The strongest growth among plastics was registered by the thermoplastic resins, up 10%. Of such resins, polypropylene continued to show the highest growth, up 14%; followed

by high-density polyethylene, up 12%; polyvinyl chloride and copolymers, up 9.9%; low-density polyethylene, up 8.3%; and polystyrene, up 7.7%.

By comparison, output of thermosetting resins grew 7.4% in 1987. Melamine resins made a spectacular recovery in 1987, up 23%, following a dismal performance a year earlier when production dropped 9.9%. Production last year at 212 million lb marked a decade high for melamine resins production. The previous high was in 1984, when the industry produced 210 million lb of the resins.

Showing good though less re-

markable gains were epoxy resin, up 8.8% in 1987; urea resins, up 8.7%; polyesters, up 7.8%; and phenol and other tar acid resins, up 5.4%.

For the five-year period between 1982 and 1987, the plastics segment grew at an annual compound rate of 8.5%. For the 10-year period between 1977 and 1987, plastics grew at an annual compound rate of 5.4%.

Synthetic rubber output increased 8.4% in 1987. Production of 4.8 billion lb last year matched the 4.8 billion lb high of 1984, but the category's output is still running 17% below the decade high of 5.6 billion lb in 1979.

Output of plastics, synthetic fibers, and synthetic rubbers continued to climb in 1987

Business

Total polymer production exceeded 57 billion lb last year

	Billions of lb		Common units					Average annual growth			
	1987	1986	1987	1986	1985	1982	1977	1986-87	1985-86	1982-87	1977-87
PLASTICS (millions of lb)											
Thermosetting resins	6.26	5.83	6,266	5,834	5,631	4,367	3,943	7.4%	3.6%	7.5%	4.7%
Phenol, other tar acid resins	2.87	2.72	2,869	2,721	2,621	2,060	1,458	5.4	3.8	6.8	7.0
Urea resins	1.38	1.27	1,382	1,271	1,210	998	963	8.7	5.0	6.7	3.7
Polyesters (unsaturated)	1.37	1.27	1,370	1,271	1,223	865	1,061	7.8	3.9	9.6	2.6
Epoxies (unmodified)	0.43	0.40	433	398	385	286	261	8.8	3.4	8.6	5.2
Melamine resins	0.21	0.17	212	173	192	158	200	22.5	-9.9	6.1	0.6
Thermoplastic resins	37.03	33.57	37,028	33,569	31,525	24,425	21,569	10.3%	6.5%	8.7%	5.6%
Low-density polyethylene	9.63	8.89	9,629	8,888	8,889	7,503	6,471	8.3	-1.1	5.1	4.1
PVC and copolymers	7.97	7.26	7,971	7,256	6,772	5,326	5,253	9.9	7.1	8.4	4.3
High-density polyethylene	8.00	7.17	7,999	7,171	6,671	4,928	3,652	11.5	7.5	10.2	8.2
Polystyrene ^a	4.78	4.44	4,782	4,442	4,054	3,191	3,446	7.7	9.6	8.4	3.3
Polypropylene	6.65	5.81	6,647	5,812	5,139	3,477	2,747	14.4	13.1	13.8	9.2
TOTAL^b	43.29	39.40	43,294	39,403	37,156	28,792	25,512	9.9%	6.0%	8.5%	5.4%
SYNTHETIC FIBERS (millions of lb)											
Cellulosics	0.61	0.62	605	619	558	584	888	-2.3%	10.9%	0.7%	-3.8%
Rayon	0.41	0.40	414	404	353	388	598	2.5	14.4	1.3	3.6
Acetate ^c	0.19	0.22	191	215	205	196	290	-11.2	4.9	-0.5	-4.1
Noncellulosics^b	8.30	7.82	8,298	7,815	7,564	6,442	6,911	6.2%	3.3%	5.2%	1.8%
Polyester ^d	3.54	3.31	3,541	3,305	3,341	3,168	3,642	7.1	-1.1	2.3	-0.3
Nylon	2.69	2.51	2,688	2,514	2,343	1,927	2,326	6.9	7.3	6.9	1.5
Olefins ^e	1.48	1.38	1,478	1,380	1,249	723	234	7.1	10.5	15.4	20.2
Acrylic ^f	0.59	0.62	591	616	631	624	709	-4.1	-2.4	-1.1	-1.8
TOTAL^b	8.90	8.43	8,903	8,434	8,122	7,026	7,799	5.6%	3.8%	4.8%	0.4%
SYNTHETIC RUBBER (thousands of metric tons)											
Styrene-butadiene ^g	1.87	1.75	850	792	735	876	1,395	7.3	7.8	-0.6	-4.8
Polybutadiene	0.82	0.74	371	336	330	288	361	10.4	1.8	5.2	0.3
Ethylene-propylene	0.56	0.51	254	230	215	123	158	10.4	7.0	15.6	4.9
Nitrile	0.15	0.13	69	59	53	45	70	16.9	11.3	8.9	-0.1
Other ^h	1.41	1.31	638	596	505	497	433	7.0	18.0	5.1	4.0
TOTAL^b	4.81	4.44	2,182	2,013	1,838	1,829	2,417	8.4%	9.5%	3.6%	-1.0%

^a No longer includes acrylonitrile-butadiene-styrene or styrene-acrylonitrile resins; historical data are restated. ^b Totals are for those products listed. Totals may not add because of rounding. ^c Includes diacetate and triacetate yarn, but does not include cigarette tow. Beginning with 1985, includes rayon yarn. ^d Excludes aramid after 1982. ^e Includes olefin film, olefin fiber, spud-bonded polypropylene, and vinyon. ^f Includes modacrylic. ^g Excludes high-styrene latex. ^h Beginning with 1985, includes neoprene; historical data are restated. Also includes butyl; polyisoprene; chlorosulfonated polyethylene; polyisobutylene; and acrylo, fluoro, and silicone elastomers. Sources: Society of the Plastics Industry, Textile Economics Bureau, Rubber Manufacturers Association

All segments of the synthetic rubber category registered increases in production in 1987. Nitrile rubber showed the largest rise at 17%, following a 11% increase in 1986. Both polybutadiene and ethylene-propylene rubber production grew 10% in 1987. The increase in polybutadiene production was the most remarkable, following a 1.8% increase in 1986. Ethylene-propylene rubber's 1987 growth follows a 1986 growth of 7.0%.

Production growth of styrene-butadiene rubber slowed in 1987 to 7.3% from a 7.8% growth rate in 1986. Also putting in a showing that was not quite as good as the year earlier were the synthetic rubber products reported as a group by the Rubber Manufacturers Association: neoprene; butyl; polyisoprene; chlorosulfonated polyethylene; polyisobutylene; and acrylo, fluoro, and silicone elastomers. Production of that group grew 7.0% in 1987, compared with 18% in 1986.

Over the past five years, the synthetic rubber category has not grown so rapidly as it has in the past two years. Synthetic rubber production has grown at an average annual rate of 3.6% over the past five years. For the past 10 years, production has actually declined at an average annual rate of 1.0%.

Though still the slowest-growing of the entire polymers product group, synthetic fibers nevertheless recorded a respectable production increase of 5.6% in 1987 over the year-earlier level. Despite continued pressure from imported textiles, noncellulosics showed the most

impressive gains. However, cellulosics declined in part because of discontinuation of U.S. triacetate production.

Output of noncellulosics recorded the best gain, rising 6.2% in 1987. The group accounted for more than 90% of total fiber production last year. Olefins continued to lead in growth; they were up 7.1% in 1987. Matching that rate of increase was polyester; staple and filament production both grew in 1987. The recovery for polyester followed a 1.1% decline in 1986. Nylon was up 6.9% in 1987 as demand for new generation broadloom continued. The only poor showing was from acrylic fiber, production of which continued to flounder. Output dropped 4.1% in 1987 following a 2.4% decline in 1986. □

.APPENDIX E.

World Chemical Outlook

United States	26
Foreign Trade	29
Western Europe	32
Japan	40
Canada	43
U.S.S.R./Eastern Europe	45
Mexico	47

What a difference four or five years can make.

Reporting on the outlook for the world's chemical industry at the end of 1982, "widespread malaise" were the words that C&EN used. "Production of chemicals has been cut back drastically," we reported at that time, and plant capacity is "far in excess of any foreseeable demand. Producers increasingly find themselves strapped financially."

Since then, however, chemical companies throughout the industrialized Western World have inflicted a massive revamping upon themselves. It has not been without pain. Tens of thousands of jobs have been wiped out. Hundreds of millions of dollars worth of operating assets have been jettisoned. Research and investment programs have been radically reshaped.

Demand for chemicals has increased in the past five years. As a result, output of industrial chemicals and synthetic materials in the U.S. this year is more than 40% higher than in 1982. And in the major chemical-producing countries of Western Europe and in Japan, chemical output is up 15 to 30%, as well.

That's hardly exceptional growth, by any means. But with production capacity slashed, in many key product areas plants now are being pushed to their design limits and producers are reaping the rewards—in higher prices, higher productivity, and increased profitability—that stem from a seller's market.

In a relatively slow-growing global economy, next year's harvest may be less bountiful. But the world's chemical industry now seems to have its house in good enough order that it will be able to withstand an economic slump, should it occur, better than it could in the early 1980s.

United States

A much slimmed down chemical industry is running many of its plants full out to meet current demand

David M. Kiefer, C&EN Washington

This autumn has witnessed chaotic turmoil in U.S. investment and currency exchange markets. Deep and recalcitrant deficits continue to sorely beset both the federal budget and the U.S. balance of foreign trade. Auto sales are slumping and new housing starts are on a decline. Consumer confidence may be ebbing, and, in the wake of the recent debacle on Wall Street, predictions of impending recession, although still not the consensus among economists, are becoming more common.

So who is crying all the way to the bank? U.S. chemical producers, that's who.

This year, producers of basic industrial chemicals, plastics, and fibers will boost their sales at least 10% to about \$90 billion (exclusive of sales of foreign subsidiaries), thanks to increased demand both at home and abroad and to higher selling prices. For the broader chemicals and allied products industry, sales will show a slightly stronger gain of 11 to 12%.

But it is the industry's profitability that really is on a roll. Net income for makers of basic chemicals and synthetic materials will surge 40% or even a bit more in 1987, to about \$7 billion. That's a new high by a long shot. Profit margins are

Profits up sharply as producers reap benefits of restructuring

higher than at any time in the past decade as cutbacks in overhead and improved operating efficiency permit many companies to carry more of each dollar of revenue down to the bottom line. Earnings for the overall chemicals and allied products industry, meanwhile, will increase almost as much to about \$18 billion.

Several factors are responsible for this year's favorable performance by the chemical industry. Strong consumer demand throughout the year has benefited many chemical end markets, including packaging, apparel, appliances, furniture, and electronics. As a result, output of plastics and fibers and of the chemical intermediates that are needed to make them has risen steadily throughout the year. For the year as a whole, production of industrial chemicals and synthetic materials will be about 9% higher than in 1986. And for the overall chemicals and allied products industry the increase in output will be about 6%. Both figures are considerably higher than the 3% increase in production likely this year for U.S. industry as a whole.

Meanwhile, for the second year in a row, the chemical industry is harvesting the fruits of the drastic reorganization that it imposed upon its operations in recent years, especially in 1985. The chronic excess capacity, slow market growth, depressed prices, cutthroat competition, and consequent lackluster earnings that plagued chemical companies throughout the early 1980s led to widespread programs for repositioning, rationalization, and restructuring. Marginally profitable and money-losing businesses were shut down or sold off. Staffs were cut to the bone. And many major companies shifted their emphasis away

from their traditional commodity chemical lines to focus more on the development of specialty products. These, presumably, will be less prone to competition and more likely to garner a favorable return on investment. Monsanto, for one, has slashed its investment in commodity petrochemicals from about a quarter of its total assets in 1980 to only 3% now.

The combination of increased output and scaled back plant capacity has greatly enhanced the industry's productivity. For the first time in several years, too, producers of many chemical products are enjoying a sellers' market and operating their facilities full out. In this year's third quarter, according to Federal Reserve Board statistics, plant utilization in the chemical industry rose to 84% of capacity, up from 80% a year earlier. The industry's operating rate has not been higher since 1974, and just five years ago it was as low as 68%.

In some notable old-line product areas, moreover, plants now are operating even closer to their upper limit. Chlorine and caustic soda producers, for example, were running their plants at about 90% of capacity—or over 100% of "effective" capacity, allowing for plants temporarily shut down—this fall, according to the Chlorine Institute. Output of the two workhorse coproducts will be up 4 to 6% this year to the highest level since 1980, largely because of strong demand from producers of vinyl chloride and other organic chemicals and from the pulp and paper industry. At the same time, however, with some 20 plants shuttered during the 1980s, chlor-alkali producers are making do with about 13% less capacity than they had in 1982. As a result, supplies are tight and suppliers, for a change,

now are thinking of expanding rather than shrinking capacity.

Ethylene producers, too, are hard pressed to keep customers supplied. Output this year will rise only a modest 3%, more because of capacity restraints than slow growth in demand. Current U.S. capacity of about 35 million lb per year is operating at well over 90%, so that further growth will be dependent on debottlenecking present units, upgrading their technology, or restarting plants now mothballed.

Makers of a number of other basic petrochemicals, especially monomers for making plastics, are similarly hard pressed to keep customers satisfied. And in the plastics industry itself, with total production for the first nine months of 1987 up 10% to 36 billion lb, producers are having little problem in finding a buyer for every pound they can turn out. Even man-made fiber plants are humming close to capacity.

All of which leave chemical companies, for a change, sitting in the catbird seat. Selling prices reflect this. Producer prices for industrial chemicals, after having sagged steeply last year in response to the sharp decline in the cost of petroleum-based feedstocks, have rebounded sharply throughout most of 1987. Although for the year as a whole they will be up, on average, just about 4%, right now they stand 7 to 8% higher than at the end of 1986. Plastics prices, meanwhile, are more than 15% higher. Feedstocks also cost more this year, but chemical makers are now in a good position to pass such increases, and more, along to their customers.

The continuing slump in the value of the U.S. dollar relative to major foreign currencies has been yet another fundamental plus for the chemical industry. For one thing, earnings from the industry's far-flung operations abroad are magnified when translated into dollars at increasingly favorable exchange rates. Even more important now, however, U.S. chemicals are gaining a strong price advantage in overseas markets when competing with products made in Europe or Japan. As a result, \$26 billion worth of U.S.-made chemicals will be sold to foreign customers this year, 15%

Growth in production is likely to taper off ...

Industrial production index, ^a 1977 = 100

^a Seasonally adjusted. ^b C&EN estimates. Source: Federal Reserve Board

... and a steady rise in sales volume ...

Net sales, ^a \$ billions

^a Excludes sales of foreign subsidiaries. ^b C&EN estimates. Source: Bureau of the Census

... but prices of many chemicals are climbing ...

Producer price index, 1967 = 100

^a C&EN estimates. Source: Bureau of Labor Statistics

... brings sharp growth for chemical producers' profits

Net income, \$ billions

^a C&EN estimates. Source: Bureau of the Census

more than in 1986, a hefty \$3.4 billion increase over 1986 to U.S. companies' sales volume. At the same time, of course, the weak dollar has made U.S. chemical companies look like cheap and tempting targets to foreign companies interested in expanding in the big U.S. market.

All in all, increased exports, higher sales prices, strong demand, and improved efficiency have provided the U.S. chemical industry with the leverage, this year and last, to put its finances in better shape than at

any time in more than a decade. Can the string be extended to this year?

That depends, of course, on the continued health of the overall U.S. economy. Economists and business analysts generally have been moderately optimistic for 1988. Most are looking for gross national product after adjustment for inflation, to increase about 1.5 to 2.5%—a rate of growth that probably will be slightly below this year's. And the consensus is that, although the rate of

United States

pansion may slow during the latter half of 1988, the U.S. will enjoy an unprecedented sixth year of steady economic progress.

Probably the biggest uncertainty at present is the potential fallout from the stock market collapse. It is still too early to assess just how the \$1 trillion or more in losses suffered by investors this past October and November will impact consumers' spending plans or businessmen's investment decisions. But consumers have been the linchpin in the economic expansion of the past few years, and if they indeed have been badly shaken by events on Wall Street, further growth in gross national product would likely be slowed—or even stopped.

If the economy does manage to eke out another year of slow growth, 1988 will be one more period of high production, rising sales, and strong profits for U.S. chemical firms. It is true, certainly, that two of their most important markets—housing and the automobile industry, both of which have been slumping this year—are likely to be less buoyant in 1988 as well. Housing starts this past October were at the lowest level since early 1983, and,

for the year as a whole, starts probably will be down 10% from 1986. A decline of nearly the same size would not be surprising for next year, also. That, in turn, will undercut a big market for plastics, coatings, and fibers.

The automotive market, another vast outlet for these chemical products, also is facing cloudy prospects. For all of 1987, its sales probably will total about 10 million cars, down sharply from the 11.5 million sold last year. Consequently, car makers have been trimming back their output since midyear. Auto production is likely to total about 7.1 million units for the entire year, down 9 to 10% from 1986. And already, the three major U.S. car companies have trimmed their production schedules for 1988's first quarter to a level 13% below what they rolled off their assembly lines in this year's first three months. For all of 1988, output is likely to drop at least 5%.

Growth is likely to slow, also, for demand from other outlets, such as appliance and furniture makers and the textile industry. But even if chemical production does not rise appreciably above its present level,

it will average about 4% higher next year than in 1987.

It's more likely, though, that production of chemicals, plastics, and synthetic fibers will rise 5 to 6%. Prices, too, probably will be 4 to 5% higher next year. That works out to at least a 10% increase in sales, to about \$100 billion. And with costs under good control and plants operating mostly at optimum rates, profits are expected to continue to move up smartly. Although growth is not likely to be as extensive as this year's, a further rise of 15 to 20% seems a good bet.

On the other hand, if the economy should fall into recession before the end of 1988—an event that every business analyst is thinking about at present but most are still dismissing—chemical industry executives argue that most companies are now well positioned to ward off its effects better than at any time in the recent past.

The industry's leaner, fitter stature and robust balance sheets have made it more resistant to a slump in demand for its products. Earnings would be hurt, certainly, but they probably would hold up reasonably close to this year's levels. □

Continued moderate growth is likely for production of major chemicals and polymers

	1985	1986	1987	1988
INORGANIC CHEMICALS				
Thousands of tons unless otherwise indicated				
Ammonia	17,319	14,474	15,650	16,000
Chlorine gas	10,402	10,436	10,900	11,200
Hydrochloric acid	2,803	2,413	2,450	2,550
Nitric acid	7,645	6,734	6,800	7,100
Oxygen, billions of cu ft	357	347	365	380
Phosphoric acid	10,607	9,578	10,300	10,600
Sodium hydroxide	10,790	10,691	11,400	11,700
Sulfuric acid	39,890	35,759	36,500	40,000

ORGANIC CHEMICALS				
Millions of lb unless otherwise indicated				
Acetic acid	2,897	2,723	3,000	3,150
Acetone	1,788	1,910	2,050	2,100
Acrylonitrile	2,349	2,182	2,400	2,550
Benzene, millions of gal ^a	1,283	1,362	1,520	1,600
Butadiene	2,340	2,546	2,700	2,800
Cumene	2,627	3,745	2,200	4,500
Cyclohexane	1,657	2,070	2,250	2,400
Ethylene	29,847	32,859	33,800	35,500
Ethylene glycol	4,178	4,771	4,400	4,700

	1985	1986	1987	1988
Formaldehyde	5,606	5,549	5,800	6,000
Methanol	5,003	7,205	7,450	7,800
Phenol	2,841	3,115	3,300	3,450
Phthalic anhydride	820	863	1,050	1,100
Propylene	14,887	16,522	18,200	19,600
Styrene	7,622	7,888	8,200	8,600
o-Xylene	675	788	950	1,000
p-Xylene	4,779	5,035	5,300	5,600

PLASTICS				
Millions of lb				
Polyethylene, high density	6,671	7,187	8,000	8,500
Polyethylene, low density	8,889	8,903	9,700	10,200
Polypropylene	5,139	5,812	6,650	7,100
Polystyrene	4,054	4,470	4,800	5,000
Polyvinyl chloride	6,772	7,256	7,950	8,250

SYNTHETIC FIBERS				
Millions of lb				
Acrylic and modacrylic	631	616	630	650
Nylon	2,343	2,514	2,750	2,950
Olefin and vinyon	1,249	1,393	1,480	1,550
Polyester	3,341	3,305	3,550	3,650

^a Petroleum refiners only. Sources: Bureau of the Census; International Trade Commission; Society of the Plastics Industry's Committee on Resins Statistics; Textile Economics Bureau. C&EN estimates (for 1987 and 1988)

Foreign Trade

Thanks to the sagging dollar, U.S. chemicals are making bigger inroads in European and Asian markets

Earl V. Anderson, C&EN New York

The U.S. chemical industry is continuing to rebuild its once-dwindling chemical trade surplus, which had gradually shrunk from an impressive \$12.1 billion surplus in 1980 to \$7.2 billion by 1985. The industry pushed its surplus slightly higher to \$7.8 billion last year, on the strength of a 5% increase in chemical exports.

This year, however, foreign trade of the U.S. chemical industry will increase substantially more. Thanks to a dollar that continues to grow weaker against major foreign currencies, chemical exports will jump a healthy 15% to about \$26.2 billion. Meanwhile, the weaker dollar will help to limit chemical import growth to only 5%. If chemical imports do, in fact, weigh in at \$15.7 billion this year, as expected, the U.S. chemical trade surplus will expand to \$10.5 billion. And the chemical industry will have recouped two thirds of the decline in its trade surplus that it suffered in the early 1980s.

Given this year's strong export performance, the U.S. share of world chemical exports might be expected to increase. It won't. It will slip slightly to 14.4% from last year's 14.5% and the even higher 15 to 17% range in which it hovered dur-

U.S. surplus in chemical trade broadens markedly as exports of chemical products surge

ing the early 1980s, when U.S. chemical export performance was much less vigorous than it is this year.

Why this statistical aberration? It certainly isn't because other countries are outperforming the U.S. when it comes to exporting chemicals. The answer, again, lies with the depreciating dollar.

Measured in dollars, world chemical exports this year will fall just short of \$200 billion. That's about a 15% advance, roughly the same percentage that world chemical exports (measured again in dollars) grew last year.

But realistically, the economies of the world's major industrialized nations just aren't strong enough to support that kind of growth in chemical exports. And the debt problems of many of the developing countries, particularly some of those in Latin America, make it impossible for them to boost their imports by such a large amount.

A more reasonable "guesstimate" of world chemical export growth this year is about 5% in real terms. The inflated 15% dollar-based figure results from the swiftly changing exchange rates of the dollar against other currencies.

Take, for example, the chemical exports of the European Community (EC). Most recent complete EC statistics (through June) show that EC chemical exports are actually running about 1% lower than they were last year. But EC reports its export figures in European currency units, or ECUs, which are weighted "basket" units based on the gross national product and trade of member countries.

In 1986, one U.S. dollar equaled 1.063 ECUs. This year, however, a dollar equals only 0.880 ECU, 17% lower than last year's conversion rate. Not surprisingly, EC chemical exports, measured in dollars, are up 19.5% over last year, even though they are down 1.2% in ECUs.

Cheaper dollar inflates value of world chemical trade ...

Total world chemical exports, \$ billions

... but deflates U.S. share of the total value

U.S. chemical exports, % of world total^b

^a C&EN estimates. ^b Defined as U.S. chemical exports as percent of world chemical exports less U.S. chemical imports. Sources: United Nations, Organization for Economic Cooperation & Development, European Community, Japan Ministry of International Trade & Industry

Foreign Trade

Deficit mounts higher for total U.S. foreign trade ...

... but surplus in chemical trade shows improvement

This is why U.S. chemical companies seem to be denied a higher share of world chemical exports, despite their strong export effort this year. The same story is repeated in Japan, where 1987 chemical exports will be 20% higher in terms of dollars, but less than 5% higher when valued in yen.

Because the weaker dollar makes imports from the U.S. less expensive in those countries whose currencies have appreciated most against the dollar, it is not surprising that most of the gains in U.S. chemical exports have come in the major industrialized nations. U.S. chemical exports to Western Europe, for in-

stance, will be about 18% higher than they were last year. And the countries of Western Europe, as a bloc, make up the largest single U.S. chemical export market.

Although the Canadian dollar has not appreciated as much against the U.S. dollar as some European currencies have (especially the West German deutsche mark), U.S. chemical exports to Canada will score an impressive 30% advance this year. Chalk that up primarily to Canada's strong economy. U.S. chemical exports to Japan, however, are only 10% higher than last year, even though the yen has appreciated dramatically against the U.S. dollar.

U.S. is stepping up its petroleum imports ...

... as higher prices escalate their total cost

Surge in U.S. chemical exports is sparked by organic chemicals and plastics

\$ Millions	U.S. chemical exports				U.S. chemical imports			
	1987 ^a	1986	1985	1984	1987 ^a	1986	1985	1984
Organic chemicals	\$ 7,650	\$ 6,293	\$ 6,012	\$ 6,093	\$ 5,385	\$ 4,704	\$ 4,576	\$ 4,265
Plastics and resins	5,400	4,301	3,777	4,050	1,930	1,837	1,631	1,484
Medicinals and pharmaceuticals	3,275	3,090	2,708	2,628	1,380	1,240	1,084	931
Fertilizers, manufactured	2,095	1,935	2,160	1,815	770	865	967	1,142
Inorganic chemicals	2,045	1,771	1,967	2,184	1,995	1,894	1,984	2,064
Radioactive materials	915	1,302	1,311	1,421	800	1,382	1,417	1,394
Pesticides and disinfectants	785	664	614	707	410	580	399	308
Flavors, perfumes, cosmetics, and toiletries	575	488	478	498	740	679	577	521
Pigments, paints, and varnishes	420	333	322	335	425	348	266	232
Cleaning and polishing materials	365	302	292	308	205	214	172	118
Dyeing, tanning, and coloring materials	210	174	139	158	580	487	352	358
Explosives and pyrotechnics	80	81	100	146	125	111	103	84
Miscellaneous	2,385	2,032	1,878	1,992	955	660	1,005	797
TOTAL	\$26,200	\$22,766	\$21,759	\$22,336	\$15,700	\$15,001	\$14,533	\$13,697

^a C&EN estimates. Source: Bureau of the Census

U.S. trade deficit rises despite higher exports of chemicals and other manufactured goods

\$ Billions	U.S. exports				U.S. imports			
	1987 ^a	1986	1985	1984	1987 ^a	1986	1985	1984
Chemicals	\$ 26.2	\$ 22.8	\$ 21.8	\$ 22.3	\$ 15.7	\$ 15.0	\$ 14.5	\$ 13.7
Machinery and transportation equipment	104.7	95.3	94.3	90.0	173.7	161.6	137.3	119.2
Other manufactured goods	35.7	30.6	29.3	30.8	118.6	105.5	95.0	88.6
Food and live animals	18.7	17.3	19.3	24.5	20.8	20.8	18.6	18.0
Beverages and tobacco	3.4	2.9	3.0	2.9	4.0	3.9	3.7	3.7
Animal and vegetable oils and fats	1.0	1.0	1.4	1.9	0.4	0.5	0.7	0.7
Mineral fuels, lubricants, and related materials	7.6	8.1	10.0	9.3	43.6	37.3	53.9	61.0
Crude materials (inedible) other than fuels	19.5	17.3	16.9	20.2	11.2	10.4	10.4	11.1
All other products	19.9	11.0	11.0	10.1	12.0	14.9	11.1	9.8
TOTAL	\$245.0^b	\$217.3^b	\$213.1^b	\$217.9^b	\$400.0	\$370.0	\$345.3	\$325.7

^a C&EN estimates. ^b Includes re-exported imports (not included in individual product categories). Source: Bureau of the Census

Where U.S. chemical exports, along with the exports of most other products, continue to suffer is in the debt-ridden developing nations. The most indebted nations, such as Brazil, are in Latin America, which is a very large and important outlet for U.S. chemicals.

In 1980, Latin America accounted for almost 24% of all U.S. chemical exports. But the debt problems of several of the region's largest and most economically advanced nations have put the brake on their ability to import. This year, for instance, U.S. chemical exports to Latin America will increase a miserly 1% over 1986. And the share of U.S. chemical exports that Latin America takes in will drop to only 15%.

On the other hand, U.S. chemical exports to developing nations in Asia are increasing rapidly. These nations, in fact, now rival Latin America as a big market for U.S. chemical exports.

The developing countries of Asia don't have the debt burden that Latin American countries do. And, led by the so-called newly industrialized countries, such as Taiwan, South Korea, Hong Kong, and Singapore, industrial development in this area has been swift.

As a result, this year U.S. chemical exports to the area likely will be up almost 20%. And its importance as a chemical market, compared with Latin America, has grown markedly. This year, the developing countries of Asia will import 15% of all U.S. chemical exports, about the same as Latin America's share. In 1980, those shares were 13% and 24%, respectively.

The betting is that developing Asia will continue to grow as a market for U.S. chemicals. But whether Latin America can regain its former stature is a big question. Most observers don't think it will happen any time soon.

It's a catch-22 situation in Latin America. The most heavily indebted countries there can't increase their borrowing to pay for additional imports. And to pay for their existing debt, they must run large trade surpluses. This, in turn, means throttling back on imports and increasing their own exports.

For them to increase their exports, however, implies that the industrialized countries, especially the U.S., must increase their imports from the developing country debtors. But the U.S., of course, has a troublesome trade deficit of its own.

Total U.S. exports this year probably will increase 13% to \$245 billion. Imports, although held to only an 8% advance, nevertheless will reach a record \$400 billion. This combination will push the U.S. trade deficit up slightly to \$155 billion from last year's \$152.7 billion, despite the effects of a much weaker dollar.

The U.S. will pay \$45 billion for imports of oil and related products, almost \$5 billion more than last year. Without this increase, the trade deficit would have been slightly lower than it was last year.

One expert sees a danger in lowering the trade deficit too rapidly. Albert M. Wojnilower, senior adviser and managing director of First Boston Corp., says that the deficit can be lowered by increasing ex-

ports and substituting domestic products for imports, by reducing domestic demand, or by protectionism. The first proposal, he says, could lead to inflation. The second is a euphemism for recession. And the third, protectionism, is hardly desirable. "I see no choice," says Wojnilower, "but to be content with glacial progress until the [newly industrialized countries] become vigorous buyers as well as sellers." □

Sales of U.S. chemicals to Europe, Asia, Canada rise sharply

\$ Millions	1987 ^a	1986
U.S. CHEMICAL EXPORTS		
Canada	\$ 3,460	\$ 2,460
Latin America	4,060	4,060
Western Europe	8,330	7,330
Eastern Europe	365	365
Japan	3,380	3,380
Australasia	840	840
Communist Asia	605	605
Middle East	365	365
Rest of Asia	4,035	3,035
Africa	500	500
Other	260	260
TOTAL	\$26,200	\$22,200
U.S. CHEMICAL IMPORTS		
Canada	\$ 2,985	\$ 2,985
Latin America	960	960
Western Europe	7,865	7,865
Eastern Europe	155	155
Japan	2,070	2,070
Australasia	235	235
Communist Asia	205	205
Middle East	265	265
Rest of Asia	645	645
Africa	80	80
Other	235	235
TOTAL	\$15,700	\$15,700

^a C&EN estimates. Source: Bureau of the Census

HPI's role in chemicals' future

Many hydrocarbon processes must have catalyst and chemicals to operate, but some of these processes are also major suppliers of important chemicals

H. L. Hoffman and L. Riddle,
HYDROCARBON PROCESSING Staff

WHAT DO FUELS and chemicals have in common? A foolish question. Fuels ARE chemicals. Indeed, excluding atomic and subatomic particles, all matter has a chemical composition. Therefore, an attempt to assess the role of fuels manufacture on chemical manufacture must start with definitions that somehow isolate these two manufacturing entities.

Arbitrary definition. One common distinguishing characteristic is chemical purity of products. Chemical manufacturing makes products that are usually composed of single, specific chemical compounds. Fuels manufacturing makes products having an assortment of chemical compounds—normally grouped within a given boiling range.

There are many exceptions to this sort of characterization. Processed natural gas, a common fuel, is almost pure methane. On the other hand, modern plastics—recognized as chemical products—are combinations of many chemical compounds. Even a purer form of plastics such as polyethylene has varying molecular weights symbolized by $(-CH_2-CH_2-)_n$. Isn't it ironic that many compounds in refined petroleum fuels have this same molecular symbol?

Defining HPI. The hydrocarbon processing industry (HPI) concerns manufacturing of products from natural gas and crude petroleum oils. Also included are those other natural raw materials such as coal, kerogen and shale oil that are sources of fuels called "synfuels"—denoting these products are made from raw materials other than natural gas or crude petroleum oil.

So the HPI is a major producer and consumer of hundreds (more like thousands) of different chemicals. Gathering supporting statistics remains largely a problem of definition and convention. Whether one chemical or another is included in a specific list often depends on the way its manufacturer is classified.

For the fuels portion of the HPI, U.S. refiners, for example, consumed more than 1,500 billion pounds of crude petroleum oil in 1986. After a few processing steps, about 85% of the prime feed was converted to fuels products. Most of these products were quickly burned for heat or motive power. Thus, statistics for the fuels portion are less complicated than for the chemicals portion.

Of the remaining 15% of the raw material, about half was used for energy needed by the refining processes. The other half was divided among a host of processes to become intermediates and final chemical products. Depending on the nature of the processing, and the classification of the manufacturer, these chemicals might be part of the USITC's reported 114 billion pounds of "primary products from pe-

TABLE 1—U.S. production of selected petrochemicals, 1987

Products	Billions of pounds	
	Jan-Sept. ¹	Yrly est. ²
Acetic acid	2.403	3.21
Acrylic acid	0.622	0.83
Acrylonitrile	1.881	2.51
Alpha olefins, C ₆ +	1.134	1.52
Aniline	0.709	0.95
Benzene, all grades**	8.361	11.18
Butadiene	2.096	2.80
Butanol, n-	0.696	0.93
Chlorobenzene, mono	0.175	0.23
Cumene	3.106	4.15
Cyclohexane	1.657	2.22
Diethylene glycol	0.343	0.46
Ethyl hexanol, 2-	0.484	0.65
Ethanol, synthetic	0.391	0.52
Ethylbenzene	7.243	9.68
Ethyl chloride	0.130	0.17
Ethylene	25.182	33.67
Ethylene glycol	3.213	4.30
Ethylene oxide	4.326	5.78
Isopropanol	1.019	1.36
Maleic anhydride	0.288	0.39
Methanol	5.445	7.28
MTBE	2.549	3.41
Phenol, synthetic	2.427	3.24
Phthalic anhydride	0.782	1.05
Polyethers and polyesters	1.476	1.97
Polyethylene and copolymers	12.845	17.17
Polypropylene	4.879	6.52
Propylene	13.943	18.64
Propylene glycol	0.593	0.79
Styrene monomer	6.015	8.04
Toluene, all uses**	5.056	6.76
Toluene diisocyanate	0.525	0.70
Vinyl acetate monomer	1.847	2.47
Vinyl chloride monomer	6.469	8.65
Xylene, o- and p-	4.644	6.13

* Yearly estimate, 1987 = (Jan. thru Sept.)(365 days/273 days)

** Originally reported in gallons. Converted to pounds using: 7.365 lb/gal, benzene; 7.26 lb/gal, toluene.

roleum and natural gas" and of the 235 billion pounds of "synthetic organic chemicals" for 1986.¹

ETHYLENE AND ITS DERIVATIVES—Relating estimated 1987 production data

Product	Prod.	Billions of pounds		Equiv. ethylene	(%)
		(Lb feed per lb)	Inter-mediate		
LDPE/LLDPE	9.35	-	-	9.2	
HDPE	7.82	-	-	8.2	
Polyethylene	17.17	-	-	17.4	52
Ethylene glycol	4.30	0.80	3.44		
Diethylene glycol	0.46	0.94	0.43		
Other	-	-	1.91		
Ethylene oxide	-	-	5.78	0.89	15
PVC	7.47	1.00	7.47		
Other	-	-	1.18		
Vinyl chloride	-	-	8.65	0.49	12
Styrene	8.04	1.13	9.09		
Other	-	-	0.59		
Ethylbenzene	-	-	9.68	0.27	8
Olefins, alpha	1.52	-	-	1.30	6
Vinyl acetate	2.47	-	-	0.37	3
Ethanol	0.52	-	-	0.63	1
Ethyl chlorides	0.17	-	-	0.49	
Other	-	-	-	1.1	3
Ethylene				33.7	100

Note: Bold numbers are from Table 1. Other numbers are calculated or inferred. Excludes inventory changes. NPRA report² includes inventory changes and other data sources.

CATALYST PRODUCTION and USE

TABLE 1—1987 estimated refining catalysts associated with fuels processing

Refining processes	Capacity, millions of bpcd	Typical U.S. averages			1987 Estimated catalyst market, millions of U.S. dollars			
		Catalyst consump., lb/bbl	Usage, millions of lb	Unit price, \$/lb	U.S.	W. Europe	Others	World
Cat. cracking	4.9	0.2	380	0.70	250	76	131	457
Hydrotreating	8.4	0.009	28	3.00	84	42	63	189
Alkylation, H ₂ SO ₄	0.50	18	3,300	0.03	99	1	6	106
Hydrocracking	1.0	0.013	4.7	10.00	47	9	42	98
Cat. reforming	3.5	0.0033	4.5	6.50	27	15	21	63
Alkylation, HF	0.40	0.15	22	0.70	15	3	5	23
Isomerization	0.25	0.015	1.4	6.00	8	4	2	14
Oligomerization	0.05	—	—	—	—	—	—	—
Total					530	150	270	950

TABLE 2—Estimated capacities used to ratio catalyst market

Refining processes	Millions of bpcd			
	U.S.	W. Europe	Others	World*
Hydrotreating	8.4	4.3	6.3	19.0
Cat. cracking	4.9	1.5	2.6	9.0
Cat. reforming	3.5	2.0	2.7	8.2
Hydrocracking	1.0	0.2	0.9	2.1
Alkylation, HF	0.40	0.09	0.13	0.62
Alkylation, H ₂ SO ₄	0.50	0.007	0.03	0.54
Isomerization	0.25	0.11	0.06	0.42
Oligomerization	0.05	—	—	—

* World excludes countries in centrally planned economies (C.P.E.).

TABLE 3—Major catalysts for refining, 1988 estimate

Catalyst type	Millions of dollars			
	U.S.	W. Europe	Other	World
Catalytic cracking	\$250	\$ 75	\$130	\$455
Hydrotreating	85	40	65	190
Alkylation, H ₂ SO ₄	100	—	5	105
Hydrocracking	45	10	40	95
Catalytic reforming	25	15	20	60
Alkylation, HF	15	5	5	25
Isomerization	10	5	—	15
Total	\$530	\$150	\$265	\$945

* Less than \$5 million. Others rounded to nearest \$5 million.

TABLE 4—Catalysts for petrochemicals, 1988 estimate

Process	Millions of dollars		
	U.S.	O.U.S.	Worldwide
Polymerization	\$265	\$ 480	\$ 745
Organic synthesis	95	170	265
Oxidation	85	155	240
Syngas	60	110	170
Hydrogenation	40	70	110
Dehydrogenation	10	15	25
Total	\$555	\$1,000	\$1,555

Source: JACA Corp., 1986, adjusted.

TABLE 3—1987 estimated petrochemical catalysts

Petrochemical processes	Millions of U.S. dollars		
	U.S.	Others	World
Polymerization	256	458	714
Organic synthesis	91	164	255
Oxidation	84	151	235
Syngas	58	105	163
Hydrogenation	37	67	104
Dehydrogenation	9	15	24
Total	535	960	1,495

[1] H.L.Hoffman, "Catalyst Usage Tied to Processes", *Hydrocarbon Processing*, p41, Feb 1987.

[2] H.L.Hoffman and L. Riddle, "HPI's Role in Chemical's Future", *Hydrocarbon Processing*, p41, Feb 1988.

END

**DATE
FILMED**

7 / 10 / 92