APR 13 1992

**Whole-Body Autoradiographic Microimaging:
Applications in Radiopharmaceutical and Drug Research***

**Prantika Som and Donald F. Sacker
Brookhaven National Laboratory
Upton, New York 11973**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

*Research supported under DOE Contract #DE-AC02-76CH00016

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

The whole-body autoradiographic (WBARG) microimaging technique is used for evaluation of the temporo-spatial distribution of radiolabeled molecules in intact animals as well as to determine the sites of accumulation of parent compounds and their metabolites. This technique is also very useful to determine the metabolism of a compound, toxicity, and effects of therapeutic interventions on the distribution of a compound in the whole body, by studying animals at different time intervals after injection of the radiolabeled compound.

The routine evaluation of a compound or radiopharmaceutical involves dissecting tissue distribution studies (DTDS), and gamma or positron imaging. DTDS has the following disadvantages:

- a) Since not all tissues can always be sampled, sites of radiopharmaceutical uptake may be missed.
- b) Because the procedure involved weighing of dissected tissue samples, the "spatial resolution" of this method is low and determined by the smallest amount that can be weighed accurately.
- c) Very small amounts of radioactivity present in a small region of a tissue which are barely detected by a scintillation counter can be efficiently detected by autoradiography because the nuclear emulsion is able to integrate the incident radiation over a much longer time than practicable with tissue-counting method.
- d) When using β^- -labeled compounds, it is necessary to completely dissolve the tissue to enable scintillation counting. It involves a time-consuming procedure. On the contrary, WBARG technique allows the pictorial depiction of the distribution of a compound globally, as well as regionally, in minute detail within discrete areas of an organ.

Although the gamma camera imaging (planar, SPECT) or positron emission tomography (PET) have a larger field of view enabling evaluation of global distribution of a compound, they have relatively low spatial resolution,

compared to WBARG.

The development of a simple but ingenious technique to study the macroscopic distribution of soluble compounds developed by Ullberg in 1954 revolutionized the field of WBARG and led to the study of the distribution and metabolism of endogenous as well as foreign substances in the living organisms (1). Since then, the technique has been used extensively in biomedical and toxicological research (2-14).

Another giant step in the field of whole-body autoradiographic microimaging was accomplished when Berlin and Ullberg developed a method for quantitating the distribution of radioactivity in whole-body autoradiographs (3). Subsequently, Appelgren developed a semi-quantitative method for analysis of autoradiographs (15), and Cross and associates developed a quantitative method to relate the optical density to actual radioactivity (16). More recently, a computer-assisted image analysis system (based on videodensitometry), a quantitative method of WBARG applied to positron, beta and gamma emitters (17) has been described which produces quantitative values comparable to those obtained by tissue counting with the advantage of depicting the global distribution at high spatial resolution of very minute areas within a region.

Table 1. Whole-Body Autoradiographic Microimaging: Advantages

-
- Pictorial mapping of the temporo-spatial distribution pattern of compounds
 - Determination of binding sites of potentially toxic substances in an organ or a region.
 - Comprehensive information on global and regional distribution of a compound or drug.
 - No preselection of tissue samples needed; therefore, unforeseen (unsuspected) localization can be detected.
 - Higher resolution than other imaging techniques.
 - Quantitative information derived by videodensitometric digitization can be correlated visually with regional distribution patterns.
-

Over the last four decades, the technique of whole-body autoradiography has been simplified and modified by various researchers (8,9). A brief description of the general method of WBARG is shown in Table 2.

Table 2. WBARG: General Method

-
1. Administration of radiolabeled substance to animals.
 2. Animals sacrificed at specific time intervals.
 3. Animals frozen in mixture of hexane/solid CO₂ or in liquid nitrogen.
 4. Embedding in carboxymethyl cellulose.
 5. Sectioning of wholebody or organs by cryomicrotome at -20°C (section thickness 20-50 μ m, average 30 μ m).
 6. Sections lifted with scotch-tape (3M, 810), air dried in freezer.
 7. Exposure to x-ray film along with step-wedge standards of known amount of radioactivity for specified time periods.
 8. Development of x-ray film.
 9. Quantitation: Digital videodensitometry.
-

Choice of radioisotopes: Negatron (β^-), γ and positron (β^+) emitters.

Most commonly, WBARG has been obtained using negatron emitters; however, more recently a variety of radioisotopes have been used for labeling compounds which include γ and positron emitters. Consideration should be given when choosing a radionuclide based on the following criteria:

1. Half-life of radionuclide:

An isotope with a longer half-life permits more time to perform the study so that small amounts of radioisotope can be used and exposed on film for longer time periods, reducing the cost. However, the problem of radioisotope disposal is an environmental problem (viz. C-14, I-125, Fe-59), contrary to very short-lived isotopes (viz. C-11, F-18) which can also be easily introduced into an organic molecule. The major disadvantages with very short-lived compounds are the high cost of labeling molecules, need for in-house cyclotron for production of

radionuclide, the need of expert synthetic organic chemists, limited time available for study (which may require special handling, viz. force-drying before putting on the film, use of screen with the film).

2. Other Physical Characteristic:

I-125 is a useful isotope for labeling proteins and peptides, and its emission of low-energy photons makes it a suitable radionuclide for WBARG. A relatively shorter half-life I-131 with β^- emission, however, produces a better quality image. Very high energy isotopes such as positron emitters may produce "fuzzy" images depending on the film type and some difficulties are encountered with the high-energy photons. Tritiated compounds are usually easy to prepare; however, low-energy particles of H-3 require use of large amounts of isotope for satisfactory autoradiograms. The special film needed and manual film-processing technique for H-3 labeled compounds are also expensive and time consuming. A list of isotopes which have been used for WBARG is given in Table 3.

Table 3. Radioisotopes Used for WBARG

Isotope	t _{1/2}	Decay Mode and Emission	Reference
H-3	12.3 y	β^-	13
C-14	5760 y	β^-	10
F-18	11.83 h	β^- (97%), EC (3%)	8,18
Na-22	2.60 y	β^+ (90.5%), EC (9.5%), γ	19
Mg-28	21.10 h	β^- , γ	20
P-32	14.26 d	β^-	21
S-35	87.20 d	β^-	1
Cl-38	37.24 m	β^- , γ	22
K-43	22.30 h	β^- , γ	23
Ca-45	163 d	β^-	18
V-48	15.97 d	β^- , EC (51%), γ	24
Cr-51	27.70 d	EC (100%), γ	25
Co-58	70.78 d	β^+ , EC (86%), γ	26
Fe-59	45.1 d	β^- , γ	27
Co-60	5.26 y	β^- , γ	28
Ni-63	100.10 y	β^-	29
Zn-65	243.90 d	β^+ , EC (98%), γ	30
Ga-67	3.24 d	EC (100%), γ	31
As-73	80.30 d	EC (100%), γ	32
Se-75	120.40 d	EC (100%), γ	33
Br-80m	4.42 h	IT (100%), γ	34
Br-82	35.30 h	β^- , γ	34
Sr-85	65.19 d	EC (100%), γ	35
Rb-86	18.66 d	β^- , γ	23
Sr-90	28.60 y	β^-	36
Y-91	58.51 d	β^- , γ	37
Ru-97	2.9 d	EC (100%), γ	38
Mo-99	2.76 d	β^- , 100%, γ	39
Tc-99m	6.02 h	IT (100%), γ	11,40

Table 3: (continued)

Isotope	t _{1/2}	Decay Mode and Emission	Reference
Ru-103	39.35 d	β^- , γ	41
Cd-109	1.24 y	EC (100%), γ	2
In-111	2.83 d	EC (100%), γ	42
Sn-117m	13.61 d	IT, γ	11
I-123	13.20 h	EC (100%), γ	43
I-125	60.14 d	EC (56%), γ	12
Te-127	9.35 h	β^- , γ	44
I-131	8.04 d	β^- , γ	14,45
Ba-133	10.50 y	EC (10%), γ	46
Cs-137	30.17 y	β^-	47
Ba-140	12.79 d	β^- , γ	46
La-140	1.68 d	β^- , γ	46
Ce-144	284.31 d	β^- , γ	48
Au-198	2.69 d	β^- , γ	49
Tl-201	3.08 d	EC (10%), γ	10
Hg-203	46.60 d	β^- , γ	50
Po-210	138.38 d	α , γ	51
Pu-238	89 y	α , γ	52
Am-241	432.20 y	α , γ	53

Animal Models

Animal models chosen for a particular study depend on various factors (54). It is of utmost importance to select an animal model carefully since the distribution, kinetics, metabolism, excretion, and toxicity of a compound may vary between species, and may be affected by sex, genetic variations between species, age, nutritional status, disease, and pregnancy. Some species (mice, rats) are used because they are relatively inexpensive and therefore large numbers can be used for obtaining statistically meaningful data. These animals can be bred easily in a laboratory environment and they reach sexual maturity

rapidly. They can, therefore, be used within a short period of time. Moreover, because of their body size, they require a lesser amount of injected dose of the compound and many animals can be blocked and cut simultaneously, making this model economical.

Larger animals (up to 5 kg body weight) such as minipigs, rabbits, dogs, and small monkeys have also been studied using more modern microtomes with large cutting blades and stage capabilities.

Choice of Adhesive Tapes

Several types of adhesive tapes suitable for adhering to the surface of the frozen animal section at -20°C without losing their flexibility have been described. Synthetic adhesive appears to adhere better than resin-type adhesive. Silicon- treated adhesive is more suitable for lower temperatures.

Table 4: Adhesive Tape for WBARG

Manufacturer and Brand	Type	Use
3M 800	Clear cellulose backing	Adheres very well to frozen section at -20°C Good for staining
3M 810	Matt backing	Adhere very well to frozen section at -20°C , most commonly used Good for staining
3M 352	Polyester (Rubber and resin adhesive)	Suitable for WBARG inexpensive
3M 850	Polyester (Synthetic adhesive)	Suitable for WBARG inexpensive
Hisamitsu Seiyaku Salotape RI-50	Cotton/nylon back	Easily transferable to glass slides for staining

Choice of Films

A wide variety of films have been used for WBARG depending on the radionuclides used and film sensitivity. High-sensitivity films often tend to

have high levels of background fog. Non-screen x-ray cassettes are generally used for WBARG. However, for short-lived positron emitters, intensifying screens are necessary (8). Single (SB5) and double (X-RP1) coated films have been shown to behave differently when using positron emitting radionuclides. At high count rates, X-RP1 (double coated) is more sensitive whereas SB5 (single coated) is more sensitive at lower count rates (8). A list of films which have been used for WBARG is given in Table 5. Other films which have been recommended are Industrex AA (Kodak), Betamax (Amersham).

Table 5: Film types used for WBARG

Film type	Characteristics
DEF-5 (Kodak)	Fastest film, expensive
Ilflex 90 (Ilford)	High sensitivity
Industrex C (Kodak)	Medium sensitivity, high contrast
Lo-Dose Mammography (Du Pont)	Single coated, medium sensitivity, good for wide-range of energy spectrum and emission, high contrast
Osray M3 (Agfa Gevaert)	High sensitivity
SB 5 (X-OMAT) (Kodak)	Single coated, good resolution, but slower than DEF 5
Singul X/RP (CEA Verken)	High sensitivity
Structurix D7 (Agfa Gevaert)	Medium sensitivity high contrast
Ultrofilm (LKB)	Single coated More sensitive to H-3
XAR (Kodak)	Slower than SB-5
XRPI (Kodak)	Double coated medium sensitivity high contrast

Quantitations

The pictorial mapping of the distribution patterns of various substances is complemented when quantitative information is also extracted from ARG films. Berlin and Ullberg first developed a semiquantitative technique using a "radioactive scale" which was visually matched with the corresponding ARGs (3). Significant progress was made when these film images were quantified by using various types of manually operated densitometers (16,55). Although the spatial resolution of an ARG is far better than can be obtained in vivo by external detection (gamma cameras, positron-emission tomography), the procedures for quantitating ARGs are less well established. Computer-aided in vivo imaging systems in routine use provide useful numerical estimates of activity in different organs as well as quantitative changes of activity with time. In order to obtain similar information from autoradiography, it is essential to be able to quantify the distribution of activity in an ARG. By quantifying data from animals killed at different times after tracer injection, one can estimate the temporal kinetic of tracer distribution.

Semi-quantitative analysis of ARGs has been performed by many investigators (55-57). Gooche et al. (58) developed a system to quantitate ARGs using a computer-coupled scanning densitometer. Yonekura et al. (17) described a simple low cost system for digital film analysis based on videosensitometry (59) which uses a TV camera coupled to a microcomputer. Quantitation with special emphasis on positron-emitting radiopharmaceuticals was accomplished with this system which was applied for short-lived positron-emitters as well as for gamma emitters and more conventional beta emitters for simultaneous multiple-tracer studies (8,17):

At present several commercial systems are available which are simple to operate and provide various types of software and capabilities for image enhancement (contrast stretching, spatial filtering), normalization, restoration,

pseudo coloring, superimposition and subtraction/addition/division of images. These capabilities are of importance when using double or multiple tracer studies.

Radioactive Standards for Quantitations:

For long-lived β^- -emitting radioisotopes (v.z. H-3, C-14) serial step-wedge standards of known amounts of radioactive dilution as tissue equivalent standards are available commercially (Amersham Corporation, American Radiolabeled Chemicals). However, the use of these standards is limited by the fact that they do not provide a wide enough range of radioactive dilutions sometimes needed to quantitate temporal wholebody distribution of radioactivity where radioactivity concentration in different regions/organs show wide variations. "Home-made" step-wedge dilution standards are prepared for short-lived gamma and positron emitters. Filter papers soaked in serial dilutions of known amounts of radioactivity had been used as standards (8), however, these standards only provide a relative number for quantitation, since the thickness of the filter paper is often different than that of the tissue sections. Tissue-equivalent standards using several types of media (carboxymethyl cellulose, chicken liver paste, gelatin) has been used to quantitative ARG (60) which are cut alongside with the wholebody/organs. These types of standards provide more accurate quantitative data (cpm/g, $\mu\text{Ci/g}$) since these are tissue equivalents and of the same thickness as the tissue sections. These standards can also be calibrated by punching out tissue sections, and direct counting of the tissue samples from the animal that has been frozen and sectioned (17). This procedure provides the absolute values of activity ($\mu\text{Ci/g}$ or cpm/g) as well as uptake ratios (% injected dose/g) in small regions. The standards are exposed to the film under same conditions as the tissue sections.

Applications:

Single tracer:

The possible applications with single tracer ARG are shown in table 6. Single tracer ARG techniques have shown different affinity of labeled compounds for 1) the same organ in different species, 2) different organs in the same species, 3) specific cellular affinity in the same organ. The distribution of L-tyrosine-0- ^{35}S -sulphate was found to be very similar in mice and rats except that in the rat it accumulates in the inner cortex of the kidney and in the mouse it accumulates in the renal medulla and pelvis. Metabolic data showed that L-tyrosine sulphate is metabolized in the rats and rabbits but not in mice, and that the renal cortex is the site of metabolism (61,62).

WBARG technique can be used to simultaneously visualize the distribution of a compound and subsequent identification of the nature of radioactivity (metabolite) in the same tissue by punching out small pieces of tissue and body fluids from the wholebody section and by chromatographic separations (63). Utilizing a more sophisticated technique such as the three-step WBARG, total radioactivity and non-volatile metabolites as well as non-extractable metabolites were determined by Bergman and colleagues (64). In this study, the ARGs were exposed at different temperature to separate different types of metabolites. Drug binding sites and receptor density mapping in discrete regions is possible using this technique (65,66). The active placental transfer of drugs was also demonstrated by using WBARG (67-69).

WARBG is a useful technique to determine a treatment regimen. It was shown by Shindo et al. (70) that dosages of less than 50 mg/kg of ^{14}C -L-DOPA showed little uptake in the brain while at higher doses the brain uptake was significantly increased. See examples of single tracer studies shown in Figures 2-6.

Table. 6: Applications of Single Tracer Autoradiography

-
- Wholebody and regional temporo-spatial distribution of drugs and newly developed radiopharmaceuticals
 - Determination of drug binding sites
 - Drug receptor density mapping
 - Determination of transfer function (placental, CNS, mammary)
 - Toxicology of drugs
 - Determination of treatment regimens
 - Determination of metabolites in tissues
-

Double tracers:

The method also lends itself to use of double or multiple tracers. The applications for double tracer technique is described in Table 7. The advantages of double tracer ARG are: a) same animal can be used as its own control, therefore any biological variations are eliminated. b) It is economical since fewer animals are needed. In double-tracer studies, the discrimination between radionuclides are made on the basis of: 1) physical half-lives; 2) energy (8). An example of quantitative data using double-tracer technique is shown in table 8. In this study, the animals were injected with 2 glucose analogs (C-14-DG and F-18-FDG). The brain samples were first exposed for shorter half-life F-18 ($t_{1/2}$ = 109 min). The sections were then allowed to decay for 10 half-lives of F-18 and re-exposed on film for longer half-life C-14 ($t_{1/2}$ = 5730 y). Examples of double tracer ARG are shown in Figures 7-11.

Table 7. Applications of Double Tracer Autoradiography

-
1.	Evaluation of new radiopharmaceuticals (same animal, same compound, different radionuclide).
2.	Metabolism and bioavailability of compound (same animal, one compound labeled at different positions).
3.	Uptake and metabolism of different compound (same animal, different compound, different radionuclide).
4.	Effects of therapeutic interventions (same animal, same compound, different radionuclide).
5.	Measure two physiological function (flow/metabolism) (same animal, different compound, different radionuclide).
-

The WBARG microimaging technique provides extremely valuable information on the distribution and kinetics of compounds. Sites of accumulation of radioactivity, may reflect the metabolic fate or therapeutic and toxic effect of a compound. Although this technique does not provide biochemical information at the cellular or molecular levels, however, the information can be used in conjunction with other techniques for complete understanding of drug metabolism and toxicity. WBARG seems to be a very useful survey technique for drug distribution and receptor mapping. Receptor mapping is currently done by using small pieces of tissue and microautoradiography. Wholebody receptor mapping may enable to select organs or regions of organs which need further investigation.

Wholebody autoradiographic determinations of placental transport of drugs at different gestation periods is of utmost importance for assessing the effects of these drugs for development of fetal abnormalities, congenital malformations and other toxic effects on the fetus.

The potential applications of this technique are summarized in table 9. Quantitative WBARG may become a very useful tool in the development of new radiopharmaceuticals. It complements the existing methods and provides a more comprehensive and detailed data on biodistribution, metabolic pathways radiation

dosimetry, at lower cost and by using less animals.

Table 8. Concentrations of C-14 2DG and F-18 FDG in Mouse Brain*

Compound	Frontal Cortex	Caudate nucleus	White matter	Thalamus	Cerebellum
C-14 2DG†	182.4±15.4	145.7±23.5	53.8±13.8	183.4±14.5	166.7±25.7
F-18 FDG‡	143.9±35.5	129.9±19.1	53.2±11.3	160.8±10.4	148.8±22.5
F-18/C-14	0.89	0.89	0.99	0.88	0.89

* Quantitation by digitized videodensitometry from double-isotope study.

† Calibrated by methylmethacrylate standards (nCi/g).

‡ Calibrated by filter-paper standards (cps).

Table 9. Wholebody ARG: Possible Future Applications

1. Evaluation of new PET and SPECT radiopharmaceuticals.
2. Measure simultaneously different physiological parameters (blood flow, metabolism).
3. Pictorial visualization of wholebody distributions of compounds will allow determination of metabolic forms of compounds globally as well regionally in conjunction with other techniques (tissue section punching and metabolite determinations).
4. WBARG will be increasingly used not only to identify the sites of accumulation of labeled compounds in-vivo but also will point to the organs/tissues which require further investigations by other physiological/pharmacological and biochemical techniques.
5. Be of help in directing molecules of specific designs to specific organs in the body.
6. Determine unexpected areas of receptor binding sites in wholebody in animals.
7. WBARG technique will be very useful in evaluating the distribution of labeled compounds in relation to aging, infection, nutritional status, drug therapy, and various disease states.
8. Determination of the rate and amount of transfer of pharmaceuticals and toxic agents through biological membranes (placenta, CNS, retina) is of importance to assess the effects of therapeutic interventions or potential toxicity. WBARG technique will be very useful to obtain such information easily with low cost.

REFERENCES

1. Ullberg S: Studies on the distribution and fate of 35 S-labelled benzyl penicillin in the body. *Acta Radiol.*, (Suppl. 118): 1-110, 1954.
2. Berlin M. and Ullberg: The fate of Cd^{109} in the mouse. An autoradiographic study after a single intravenous injection of $Cd^{109}Cl_2$. *Arch. envir. Hlth.* 7: 686-693, 1963.
3. Berlin M. and Ullberg: Accumulation and retention of mercury in the mouse. *Arch. envir. Hlth.* 6: 589-616, 1963.
4. Cohen EM and Hood M: Application of low temperature autoradiography to studies of the uptake and metabolism of volatile anaesthetics in mouse. *Anaesthesiology* 30: 1969.
5. Busch J: Wholebody autoradiography (WBARG): use for pilot studies of pharmacokinetics in rats. *Acta Pharmacol. Toxicol.* 41: 28-29, 1977.
6. Cross SAM: The localization of metiamide and cimetidine using autoradiographical techniques. *Proc. Eur. Soc. Toxicol.* 18: 288-290, 1977.
7. Cross SAM: Localisation of histamine H_2 -receptor antagonists in the gastric mucosa. *Histochem. J.* 9: 619-644, 1977.
8. Som P, Yonekura Y, Oster ZH, et al.: Quantitative autoradiography with radiopharmaceuticals. Part II: Applications in radiopharmaceutical research. *J. Nucl. Med.* 24: 238-244, 1983.
9. d'Argy R, Ullberg S, Stalnacke C-G, and Langstrom B: Whole-body autoradiography using ^{11}C with double-tracer applications. *Ind. J. Appl. Radiat. Isot.* 35: 129-134, 1984.
10. Yonekura Y, Brill AB, Som P, et al.: Quantitative autoradiographic measurement of regional myocardial substrate utilization in hypertensive rats. *Science* 227: 1494-1496, 1985.

11. Oster ZH, Som P, Srivastava SC, et al.: The development and in vivo behavior of tin-containing radiopharmaceutical. II: Autoradiographic and scintigraphic studies in normal animals and in animal models of bone disease. Int. J. Nucl. Med. Biol. 12: 175-184, 1985.
12. Som P, Oster ZH, Yamamoto K, et al.: Some factors affecting the cerebral and extracerebral accumulation of N-isopropyl-p-iodo-amphetamine (IAMP). Int. J. Nucl. Med. Biol. 12: 185-196, 1985.
13. Fand I, McNally WP, Koul O, et al.: Whole-body autoradiographic localization of [³H]Phenylcycline and its metabolites in mice. Toxicol. Appl. Pharmacol. 93:421-432, 1988.
14. Som P, Oster ZH, Goodman MM and Knapp FF, Jr.: Microimaging studies of myocardial substrate utilization and perfusion in two models of non-coronary heart disease. NucCompact 21: 259-262, 1990.
15. Appelgren LE: Sites of steroid hormone for median. Autoradiographical studies using labelled precursors. Acta Physiol. Scand. (Suppl. 301), 1-108, 1967.
16. Cross SAM, Groves AD and Hesselbo T: A quantitative method for measuring radioactivity in tissues sectioned for whole-body autoradiography. Int. J. Appl. Radiat. Isotopes 25, 381-386, 1974.
17. Yonekura Y, Brill AB, Som P, et al.: Quantitative autoradiography with radiopharmaceuticals. Part I: Digital film analysis system by videodensitometry. J. Nucl. Med. 24: 231-237, 1983.
18. Ericsson Y, and Hammarström L: Mouse placental transfer of F¹⁸ in comparison with Ca⁴⁵. Acta odont. scand. 22: 523-538, 1964.
19. Huggert A, Odelblad E, Söremark R, and Ullberg S: Distribution and kinetics of sodium (Na²²) in mice and rabbits. Acta Isot. (Padova) 2: 151-175, 1961.
20. Arnaud MJ: Autoradiographic study in the rat of the transit of magnesium

- from mineral water. In: "Conference on Autoradiography in Pharmacology and Toxicology", Uppsala. Acta pharmac. tox. 41 (Suppl. 1): 154-155, 1977.
21. Bengtsson G: Autoradiographic distribution studies of injected ^{32}P -labelled polydiethylstilboestrol phosphate. Acta endocr. 43: 581-586, 1963.
 22. Ullberg S, Appelgren L-E, Clemedson C-J, et al: A comparison of the distribution of some halide ions in the body. Biochem. Pharmac. 13:407-412, 1964.
 23. Söremark R, Olsson K, and Wing K: Uptake and distribution of rubidium-86 and potassium-43 in mice and rats. An autoradiographic study. Acta physiol. scand. 77: 322-332, 1969.
 24. Söremark R, Ullberg A, and Appelgren LE: Autoradiographic localisation of vanadium pentoxide ($\text{V}_2^{48}\text{O}_5$) in developing teeth and bones of rats. Acta odont. scand. 20: 225-232, 1962.
 25. Christian RA, Baker JRJ, and Bullock GR: The use of ^{51}Cr for autoradiography at all levels. Acta pharmac. tox. 41 (Suppl. 1), 30-31, 1977.
 26. Ullberg S, Kristofferson H, Flodh H, and Hanngren Å: Placental passage and fetal accumulation of labelled vitamin B_{12} in the mouse. Archs. int. Pharmacodyn. Théor. 167:431-449, 1967.
 27. Ullberg S, Sörbo B, and Clemedson C-J: Distribution of radioactive iron in pregnant mice studied in whole-body autoradiography. Acta radiol. 55: 145-155, 1961.
 28. Blomquist L, Flodh H, and Ullberg S: Uptake of labelled vitamin B_{12} and 4-iodophenylalanine in some tumors of mice. Experientia 25: 294-296, 1969.
 29. Oskarsson A, and Tjälve H: Autoradiography of nickel chloride and nickel carboxyl in mice. In "Conference on Autoradiography in Pharmacology and Toxicology", Uppsala. Acta pharmac. tox. 41 (Suppl. 1): 158-159, 1977.

30. Denker L: Tissue localization of some teratogens at early and late gestation related to fetal effects. *Acta pharmac. tox.* 39 (Suppl. 1): 1-31, 1976.
31. Kamada M: Macroautoradiographic analysis on the concentration of ^{67}Ga -citrate in the tumor tissues and nontarget tissues of tumor-bearing mice. *J. Jpn. Soc. Cancer. Ther.* 11(3): 528-534, 1976.
32. Deak ST, Csaky KG, and Wadell WJ: Localisation and histochemical correlation of arsenic-73 by whole-body autoradiography in mice. *J. Toxicol. Environ. Health* (6): 981-984, 1976.
33. Bérnard P, Rico AG, Braun JP, and Burgat-Sacaze V: New approach to the visualization of receptors by autoradiography. In "Conference on Autoradiography in Pharmacology and Toxicology" Uppsala, *Acta pharmac. tox.* 41 (Suppl. 1): 20-21, 1977.
34. Söremark R: Distribution and kinetics of bromide ions in the mammalian body, some experimental investigations using $\text{Br}^{80\text{m}}$ and Br^{82} . *Acta radiol.* (Suppl. 190), 1960.
35. Appelgren LE, Nilsson A, and Ullberg S: Autoradiographic localisation of strontium-85 in osteosarcomas. *Acta radio (Ther)* 1: 459-464, 1963.
36. Nilsson A, and Ullberg S: Uptake and retention of strontium-90 in mouse tissues studied by whole animal autoradiography and impulse counting. *Acta radiol.* 58:81-98, 1962.
37. Appelgren L-E, Nelson A, and Ullberg S: Distribution of yttrium-91 in mice studied by whole-body autoradiography. *Acta radiol. (Ther)* 4: 41-48, 1966.
38. Som P, and Oster ZH: Unpublished data, 1985.
39. Bawden JW, and Hammerström LE: Autoradiography of ^{99}Mo in developing rat teeth and bone. *Scand. J. Dent. Res.* 84(3): 168-174, 1976.

40. Rohlin M, and Hammarström L: Whole-body autoradiography of ^{99}Tcm -labelled pyrophosphate and related compounds in young rats. *Acta radiol. (Ther) (Stockh.)* 15(1): 71-80, 1976.
41. Som P, Oster ZH, Matsui K, et al: Transferrin uptake in tumor and abscess. *Eur. J. Nucl. Med.* 8: 491-494, 1983.
42. Som P, and Oster ZH: In-111 HIG for tumor and abscess. Unpublished data, 1991.
43. Som P, Oster ZH, Knapp FF, Jr: Effects of cocaine on I-123 BMIPP distribution in the heart. Unpublished data, 1991.
44. Agnew WF: Trans-placental uptake of Tellurium-127^m studied by whole-body autoradiography. *Teratology* 6(3): 331-338, 1972.
45. Nair V, and Roth LJ: Autoradiography of whole animals as an experimental tool in pharmacological research. *Atomlight* 18: 7-10, 1961.
46. Dencker L, Nilsson A, Rönnbäck C, and Walinder G: Uptake and retention of ^{133}Ba and ^{140}Ba - ^{140}La in mouse tissues. *Acta radiol. (Ther) (Stockh.)* 15: 273-287, 1976.
47. Nelson A, Ullberg S, Kristoffersson H, and Rönnbäck C: Distribution of radiocesium in mice. *Acta radiol.* 55: 374-384, 1961.
48. Ewaldsson B, and Magnusson G: Distribution of radiocerium and radiopromethium in mice. An autoradiographic study. *Acta Radiol.* 2: 65-72, 1964.
49. Fries B, and Walinder G: Absorption of colloidal ^{198}Au from peritoneum and pleura and inhibition by polyphlorentin phosphate. *Acta radiol.* 2: 113-122, 1957.
50. Takahashi T, Kimura T, Sato Y, Shiraki H, and Ulita T: Time dependent distribution of mercury-203 compounds in rats and monkey as studies by whole-body autoradiography. *J. hyg. chem. Soc., Japan* 17(2): 93-107, 1971.

51. Söremark R, and Hunt VR: Autoradiographic studies of the distribution of polonium-210 in mice after a single intravenous injection. *Int. J. Radiat. Biol.* 11:43-50, 1966.
52. Matsuoka O, Kashima M, Joshima H, and Noda Y: Whole-body autoradiographic studies on plutonium metabolism as affected by its physiochemical state and route of administration. *Hlth. Phys.* 22(6): 713-722, 1972.
53. Hammarström L, and Nilsson A: Radiopathology of americium 241.
1. Distribution of americium in adult mice. *Acta radiol.* 9: 433-442, 1970.
54. Som P, and Oster ZH: Animal models of human disease. In *Clinical Aspects of Nuclear Medicine*. WB Hladik, GB Saha, KT study, Eds., Williams and Wilkine, Baltimore, pp 333-356, 1986.
55. Sokoloff L, Revich M, Kennedy C, et al.: The (^{14}C) deoxyglucose method for measurement of local cerebral glucose utilization: Theory, procedure and normal values in the conscious and anaesthetized albino rats. *J. Neurochem* 28: 897-916, 1977.
56. Irons RD and Gross EA: Standardization and calibration of wholebody autoradiography for routine semiquantitative analysis of the distribution of ^{14}C -labeled compounds in animal tissues. *Toxicol. Appl. Pharmacol.* 59: 250-256, 1981.
57. Masuoka DT, Pfeiffer E: An in-vivo brain autoradiographic technique and a digital microdosimeter for the quantitation and autoradiogram. *Acta Pharmacol. Toxicol. (Suppl. 1)* 41: 54-55, 1977.
58. Gooche C, Rasband W and Sokoloff L: Computerized densitometry and color coding of (^{14}C) deoxyglucose autoradiographs. *Ann. Neurol.* 7: 359-370, 1980.
59. Kedem D, Kedem D, Smith CW et al: Velocity distribution and blood flow measurements using videodensitometric methods. *Invest. Radiol.* 13: 46-56,

1978.

60. Som P, Bhat M, Rice C, Sacker DF: Autoradiographic standard catalog. Unpublished data, 1991.
61. Curtis CG, Powell GM, Dodgson KS: The site of metabolism of potassium L-tyrosine O- ^{35}S sulphate in the rat. *Biochem. Pharmac.* 21: 2551-2558, 1960.
62. Curtis CG, Bartholomew TC, Rose FA, Dodgson KS: Detoxification of sodium ^{35}S sulphide in the rat. *Biochem. Pharmacol.* 21: 2313-2321, 1972.
63. d'Argy R: Tissue pieces from wholebody sections used for chromatographic separation of metabolites. *Acta Pharmacol. Toxicol.* 41 (Suppl. 1): 16-17, 1977.
64. Bergman K, Tjalve H: Three-step autoradiography of organic and plastic monomers to register total radioactivity, non-volatile metabolites, and non-extractable metabolites. *Acta Pharmacol. Toxicol.* 41 (Suppl. 1): 22-23, 1977.
65. Benard P, Rico AG, Braun JP, Burgat-Sacaze V: New approach by autoradiography. *Acta Pharmacol. Toxicol.* 41 (Suppl. 1): 20-21, 1977.
66. Som P, Oster ZH, Volkow ND, Sacker DF, Weber DA: Organ variations in cocaine receptors. *J. Nucl. Med.* 31: 886, 1990.
67. Kleijn VD, Rijntjes NVM, Vree TB et al: Similarities and differences in the kinetics of distribution and of placental transfer of antic-epileptic drugs of extremely varying physico-chemical properties. *Acta Pharmacol. Toxicol.* 41 (Suppl. 1): 168-169, 1977.
68. Ullberg S, Dencker L: Localization in fetal tissues. *Acta Pharmacol. Toxicol.* 41 (Suppl. 1) 170-171, 1977.
69. Som P: Localization of cocaine in foetal tissue. Unpublished data, 1991.

70. Shindo H, Miyakoshi N, Nakajama E. Studies on the metabolism of D-and L-isomers of 3,4-dihydroxyphenil ala.. (DOPA). Autoradiographic study of the distribution of ^{14}C -labeled D-and L-DOPA and dopamine after intravenous administration in rats. Chem. Pharm. Bull. Tokyo 20(5): 966-973, 1972.

AUTORADIOGRAM

STANDARD

DIGITAL IMAGE

Fig. 1 Method of quantitation digital autoradiography. Film of ARG and standards are digitized at same light intensity. Digital images (middle upper image) was converted to map of radioactivity using the standard curve (middle lower image). (Yonekura, Brill, Som, et. al., JNM, 1983).

^{125}I -I AMP

60 min

Fig. 2

ARG of a rat head 60 min. after IAMP injection. Notice fine details of brain structures which can easily be identified.

Fig. 3 Single Tracer: Temporal distribution of IAMP. Digitized ARG of mice after I-123 IAMP (radioiodinated amphetamine) injection. Left-hand side panel 2 min. and right hand panel 45 min. after injection. Notice high uptake of IAMP in grey matter of cerebrum (A), cerebellum (B), basal ganglia (C) and Harder's gland. Cerebellar (B) and lung activity (E) decrease in time while liver (F) and kidneys (G) increase. Notice the intense uptake in the juxta-medullary region in the kidneys and non-uniform distribution in the lungs.

Fig. 4 Single tracer: WBARG of a rat 3 hr after i.v. injection of Sn-117m
+4-DTPA (a bone-seeking agent). Notice intense uptake in the
cortical bone and lack of uptake in bone marrow.

Fig 5 Single Tracer Wholebody distribution of I 131-SP-21 (a monoclonal anticolon/ovarian cancer antibody SP-21 in a nude mouse transplanted with human colon cancer xenograft on the back. Notice high uptake in the tumor. Also note blood pool activity in the heart and thyroid uptake of I-131 due to deiodination (metabolism of the antibody). Thyroid was not blocked prior to injection of I-131-SP-21.

Fig. 5 Single Tracer. Wholebody distribution of I-131-SP-21 (a monoclonal anticolon/ovarian cancer antibody SP-21 in a nude mouse transplanted with human colon cancer xenograft on the back. Notice high uptake in the tumor. Also note blood pool activity in the heart and thyroid uptake of I-131 due to deiodination (metabolism of the antibody). Thyroid was not blocked prior to injection of I-131-SP-21.

6A

Fig. 6 Time-course WBARG of mice injected with fatty acids. A: palmitic acid (a natural fatty acid) B: C-14-BMHA (a synthetic fatty acid analog). Notice the natural fatty acid clears too fast from the heart which would not allow sufficient time for imaging. On the contrary, the retention of the synthetic fatty acid (B) in the heart is longer and the clearance from the liver and lungs is optimal at 2 hr which will provide better heart/background ratios for imaging.

14C-2DG

18F-FDG

Fig. 7 Double tracer (same animal, different radionuclide) comparison of two glucose analogs C-14-2DG and F-18-FDG distribution in the brain of a mouse injected simultaneously and sacrificed 30 min. later. Upper panels: analog images with step-wedge standards. Lower panel: digitized images. High uptake areas are caudate, basal ganglia and grey matter.

A

B

Figs

Fig. 8 Double tracer (same animal, different compound).

Double-tracer WBARG in a mouse injected with (A) I-131 HIPDM (a diamine) and (B) C-14-tryptophan (amino acid). Notice high uptake of both tracers in the pancreas, however, the distribution in other organs are quite different (I-131-HIPDM: high uptake in brain, kidneys, lungs, and salivary gland; C-14 Try: high uptake in the dividing cells of GI tract). B - Brain, G - Stomach, H - Liver, I - Intestine, K - Kidney, L - Lung, P - Pancreas, S - Salivary gland.

Fig. 9 Single tracer WBARG of neurotransmitter receptor blocking. WBARG of a mouse (4 min) following C-14-Cocaine administration (A). Notice rapid and high uptake in the brain (cortex, basal ganglia), spinal cord and the heart. C-14-Cocaine distribution in a mouse pretreated with desipramine (a norepinephrine/serotonine transporter blocker) (B). Notice significantly reduced heart uptake, while the brain and spinal cord uptake was not affected.

C-14-Cocaine distribution in a mouse following pretreatment with GBR (a dopamine re-uptake blocker) (C). Notice significantly reduced uptake in the brain, spinal cord as well as the heart.

^{14}C -2DG

↓
SALINE

^{18}F -FDG

↓
CHLOROPROMAZINE

Fig. 10 Double tracer (same animal, different radionuclide): Effect of intervention. Double tracer study in two mice given C-14-DG (upper panel). Forty five min. later one mouse received 12 mg/kg of chlorpromazine (CPZ) (right) and the other received same volume of saline as control. Subsequently, both mice received F-18-FDG and sacrificed 45 min. later. Notice decrease in muscle uptake and increase in brown fat uptake after CPZ sedation.

DMIPP

Verapamil

Normal hamsters

Saline

Verapamil

Cardiomyopathy

Saline

11A

FDG (+DMIPP)

Verapamil

Saline

Normal hamsters

Verapamil

Saline

Cardiomyopathy

Fig. 11 Double-tracer study: Therapeutic intervention with glucose analog (C-14-FDG) and fatty acid analog (I-131-DMIPP) (same animals different compounds, different isotopes):

- A. Myocardial uptake of C-14 FDG (a part of dual tracer study with I-131-DMIPP) in verapamil treated normal (upper left) and cardiomyopathic (lower left) hamster compared to saline treated normal (upper right) and cardiomyopathic (lower left) hamsters. Verapamil treatment prevented the appearance of changes seen in the saline injected CM animals.
- B. Myocardial of uptake of I-131-DMIPP (a part of the above study with C-14-FDG). Notice verapamil treatment also prevented the changes seen in saline treated CM animals and there is homogeneous distribution of I-131-DMIPP.

18F-FDG ARGs - SHEEP

A

B

C

A'

B'

C'

1

2

3

Fig. 12 Autoradiography microimaging vs. PET imaging. Example of finer resolution. F-18-FDG PET images of sheep brain (bottom row, 1-3) as compared to F-18-FDG ARG (upper row A-C) and corresponding histological sections (middle row, A-C). No detail of brain structure is seen in PET images. The resolution of ARG sections approximate that of corresponding histological levels, showing intense glucose metabolism.

END

**DATE
FILMED
5/18/92**