STUDIES OF FLUCTUATION PROCESSES IN NUCLEAR COLLISIONS

PROGRESS REPORT

MARCH 1, 1992 — FEBRUARY 28, 1993

S. AYIK

DEPARTMENT OF PHYSICS

TENNESSEE TECHNOLOGICAL UNIVERSITY

COOKEVILLE, TENNESSEE 38505

FEBRUARY 1993

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE U. S. DEPARTMENT OF ENERGY
UNDER GRANT NUMBER DE-FG05-89ER40530

MASTER

REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

eb

TABLE OF CONTENTS

PREFACE	1
A. STOCHASTIC TWO-FLUID MODEL FOR RELATIVISTIC HEAVY-ION COLLISIONS	2
B. NUCLEAR FISSION WITH A LANGEVIN EQUATION	9
APPENDIX I Publications, etc.	18
APPENDIX II Personnel	19

PREFACE

This report summarizes the progress on grant No. DE-FG05-89ER40530 during the period March 1, 1992 to February 28, 1993.

We continued our investigations of various aspects of heavy-ion collisions in the framework of the Boltzmann-Langevin Model (BLM). In a previous work, by projecting the BLM onto a collective space, we deduced a memory-dependent collective transport model. We applied this model to thermal fission and investigated the influence of the memory effects on the fission dynamics. Some results of the calculations are contained in this report. The principle investigator spent a fruitful summer (1992) at GSI, where together with Y. B. Ivanov, V. N. Russkikh and W. Nörenberg, we carried out a reduction of the relativistic BLM to a two-fluid model and extracted transport coefficients associated with fluid dynamical variables. Then we applied this model to investigate equilibration and fluctuation properties in a counter-streaming nuclear fluid. Two papers have been published (Phys. Lett. B and Nucl. Phys. A), two manuscripts have been accepted for publication (Phys. Rev. C and Nucl. Phys. A) and one manuscript is in preparation for publication. Several seminars and invited/contributed talks were presented at various meetings.

Most of the work was done in collaboration with E. Suraud, M. Belkacem, D. Boilley, Y. Abe, Y. B. Ivanov, V. N. Russkikh and W. Nörenberg. Their collaboration has been very stimulating and very much appreciated.

The secretarial and accounting duties for this grant have been performed by Gloria Julian of the TTU Department of Physics, and I wish to express my appreciation. I also wish to thank the scientists at GANIL and at GSI for their hospitality during my visits.

A. STOCHASTIC TWO-FLUID MODEL FOR RELATIVISTIC HEAVY-ION COLLISIONS

(with Y. B. Ivanov, V. N. Russkikh and W. Nörenberg)

ABSTRACT

A reduction of the relativistic BLE to a stochastic two-fluid model is presented and transport coefficients associated with fluid dynamical variables are extracted. The approach is applied to investigate equilibration in a counter-streaming nuclear system.

1. Introduction

In transport models, the system is characterized by its one-body density (the phase-space distribution function), which is determined by a Lorentz covariant transport equation containing mesonic mean-fields and a collision term of the Boltzmann-Uehling-Uhlenbeck (BUU) form [1–7]. As an alternative to the full-scale kinetic simulations, mean-field multi-fluid models have also been developed for describing heavy-ion collisions at high energies [2,8,9]. In these models, the full distribution function is decomposed into components representing the group of particles well separated in momentum space. And the collision process is described in terms of fluid dynamical variables rather than the full phase-space distribution function.

In order to provide a description of the fluctuation processes in many-body systems, an extension of the one-body transport models has been recently formulated [10,11]. This gives rise to a stochastic one-body transport model, which is referred to as the Boltzmann-Langevin Model (BLM), and it provides a consistent basis for treating dissipation and fluctuation properties of the collision process. For relativistic heavy-ion collisions, this approach was extended further on the basis of the Walecka-type of field theory [12].

In this work, we carry out a reduction of the relativistic BLE to a two-fluid model and extract transport coefficients, i.e., interflow friction and diffusion coefficients. These quantities enter as an input into the multi-fluid description of nuclear collisions. As an

application of the two-fluid model, we investigate equilibration and fluctuation properties in a counter-streaming nuclear fluid.

2. Relativistic BLE for a Mixture

A derivation of the relativistic BLE for a mixture of different hadronic species in the framework of the Walecka-type field theory was presented in ref [12]. Here, we restrict our treatment to the semi-classical evolution of the spin-isospin averaged phase-space distribution functions,

$$F^{(\alpha)}(x,p) = \frac{2d_\alpha}{(2\pi)^3} \theta(p_0) \delta(p^2 - M_\alpha^{*2}) f^{(\alpha)}(x,p) \quad (2.1)$$

where $f^{(\alpha)}(x,p)$ is the occupation number, M_α^{*2} and d_α stand for the effective mass and the spin-isospin degeneracy factor for the hadron type- α , respectively. In the semi-classical limit, we obtain a set of covariant BLE's for the fluctuation distribution functions,

$$\left[p_\mu \partial_x^\mu - Q_\mu^{(\alpha)}(x,p) \partial_p^\mu \right] F^{(\alpha)}(x,p) = K^{(\alpha)}(x,p) + \delta K^{(\alpha)}(x,p) \quad (2.2)$$

Here, the left-hand-side describes the Vlasov propagation in terms of the quantity $Q_\mu^{(\alpha)}(x,p)$, which is related to the real part of the self energy of the particle type- α . In the Walecka model, this quantity is determined by means of the effective scalar and vector meson fields, $\sigma(x)$ and $\omega_\mu(x)$, (for details, see [13]). The right-hand-side of Eq. (2.2) describes the collisional coupling due to elastic and inelastic scattering between hadron species, which leads to equilibration and particle production. The additional term $\delta K^{(\alpha)}(x,p)$ in Eq. (2.2) represents the fluctuating part of the collision term. Such an additional term always arises in a transport theory whenever one deals with a reduced treatment. It describes the coupling to the remaining degrees of freedom which are not explicitly considered. In analogy to the treatment of the Brownian motion, it is assumed that Eq. (2.2) describes a stochastic process, in which the entire distribution function $F^{(\alpha)}(x,p)$ is a stochastic function and $\delta K^{(\alpha)}(x,p)$ acts as a random force. In such a description the fluctuating collision term is characterized by a correlation matrix,

$$\langle \delta K^{(\alpha)}(x,p) \delta K^{(\beta)}(x',p') \rangle = C^{(\alpha\beta)}(x,p;x',p') \delta^{(4)}(x-x') \quad (2.3)$$

which is assumed to be local in space-time. The correlation matrix is closely related to the collision term and entirely determined by the average properties of the one-body density. The fluctuation and dissipation properties of the one-body density, like in any relaxation process, are not independent properties, but related to each other through a "fluctuation-dissipation theorem." Therefore, the close relation between the correlation matrix and the collision term can be regarded as a consequence of the fluctuation-dissipation theorem associated with the stochastic evolution of the distribution functions. The relativistic BLE approach can be considered as a basis for deriving simpler stochastic multi-fluid models for describing high-energy heavy-ion collisions.

3. Two-Fluid Model

Here, we consider a reduction of the BLM to a stochastic two-fluid model in the spirit of refs [2,9]. In a two-fluid approach, the full nucleonic distribution function is decomposed into two components which corresponds to the "p = projectile-like" and "t = target-like" nucleons, assumed to be well separated in momentum space,

$$F(x,p) = \sum_{\alpha=p,t} F^{(\alpha)}(x,p). \quad (3.1)$$

This approach is similar to the kinetic treatment of a mixture (non-identical) different particle species, where each species is characterized by its own distribution function. As a result, the partial distribution functions, $F^{(p)}(x,p)$ and $F^{(t)}(x,p)$, evolve according to the set of relativistic Eqs. (2.2–2.3). Furthermore, it is assumed that each α -fluid is in thermal equilibrium in its co-moving frame and represented by a local equilibrium Fermi-Dirac function,

$$f^{(\alpha)}(x,p) = \left\{ \exp \left[(\mathbf{u}^{(\alpha)} \cdot \mathbf{p} - \mu^{(\alpha)}) / T^{(\alpha)} \right] + 1 \right\}^{-1} \quad (3.2)$$

where the quantities $\mathbf{u}^{(\alpha)}(x)$, $T^{(\alpha)}(x)$ and $\mu^{(\alpha)}(x)$ denote the 4-velocity, temperature and the effective chemical potential of the α -fluid. We carry out the reduction into a two-fluid model by evaluating the zeroth and the first momentum moments of the BLE (2.2). This yields a set of fluid dynamical equations for the parameters of the Fermi-Dirac functions,

$$\partial^\mu \mathbf{J}_\mu^{(\alpha)} = 0 \quad (3.3)$$

$$\partial^\mu \mathbf{T}_{\mu\nu}^{(\alpha)} - \mathbf{Q}_\nu^{(\alpha)} = \mathbf{R}_\nu^{(\alpha)} + \delta \mathbf{R}_\nu^{(\alpha)} \quad (3.4)$$

where $\mathbf{T}_{\mu\nu}^{(\alpha)} = \left[\mathbf{E}^{(\alpha)} + \mathbf{P}^{(\alpha)} \right] \mathbf{u}_\mu^{(\alpha)} \mathbf{u}_\nu^{(\alpha)} - g_{\mu\nu} \mathbf{P}^{(\alpha)}$ and $\mathbf{J}_\mu^{(\alpha)} = n^{(\alpha)} \mathbf{u}_\mu^{(\alpha)}$ are the energy-momentum tensor and the current density of the α -fluid, and $\mathbf{Q}_\mu^{(\alpha)}(\mathbf{x}) = \int d^4p \mathbf{F}^{(\alpha)}(\mathbf{x}, p) \partial_p^\mu \mathbf{Q}_\mu^{(\alpha)}$ denotes the mean-field contribution to the equation of motion. The pressure $\mathbf{P}^{(\alpha)}$, the internal energy density $\mathbf{E}^{(\alpha)}$ and the local density $n^{(\alpha)}$ are expressed by the standard statistical expressions in terms of the temperature and the chemical potential of the α -fluid. The quantities $\mathbf{R}_\mu^{(\alpha)}$ and $\delta \mathbf{R}_\mu^{(\alpha)}$ on the right-hand-side of Eq. (3.4) arise from the interflow collisional coupling, and they represent the friction- and stochastic-forces on the α -fluid. Using explicit forms of the collision term in Eq.(2.2) and the correlation matrix in Eq. (2.3), we find the friction force and the correlator of the stochastic force are given by

$$\mathbf{R}_\mu^{(\alpha)}(\mathbf{x}) = -D(\mathbf{J}_\mu^{(\alpha)} \rho^{(\bar{\alpha})} - \mathbf{J}_\mu^{(\bar{\alpha})} \rho^{(\alpha)}) \quad (3.5)$$

$$\langle \delta \mathbf{R}_\mu^{(\alpha)}(\mathbf{x}) \delta \mathbf{R}_\nu^{(\alpha)}(\mathbf{x}') \rangle = \delta^{(4)}(\mathbf{x}-\mathbf{x}') \sum_\beta C_{\mu\nu}^{(\beta)}(\mathbf{x}) \quad (3.6)$$

$$\begin{aligned} C_{\mu\nu}^{(\beta)} &= (A_L - A_T) \left[\mathbf{T}_{\mu\nu}^{(\beta)} \rho^{(\beta)} - M^* \mathbf{J}_\mu^{(\beta)} \mathbf{J}_\nu^{(\beta)} \right] \\ &+ A_T g_{\mu\nu} M^* \left[\rho^{(\beta)} \rho^{(\beta)} - \mathbf{J}^{(\beta)} \cdot \mathbf{J}^{(\beta)} \right] \end{aligned} \quad (3.7)$$

where $\rho^{(\alpha)}$ is the scalar density and $\bar{\alpha}$ denotes the fluid component complementary to the α -fluid. Here, the interflow friction coefficient D , and the diffusion coefficients A_L and A_T are determined in terms of the scattering cross-section. In the case of only elastic scatterings they are given by

$$\begin{bmatrix} D \\ A_L \\ A_T \end{bmatrix} = \frac{\sqrt{\lambda}}{4} \int_{\theta < \pi/2} d\sigma \begin{bmatrix} 1 - \cos \theta \\ \frac{1}{2}(1 - \cos \theta)^2 \\ \frac{1}{6} \sin^2 \theta \end{bmatrix} \left[(1 - f(1')) [1 - f(2')] \right] \quad (3.8)$$

with $\lambda = \bar{s}(\bar{s} - 4M^{*2})$, where $\bar{s}$ is an average value of the square of the invariant total

energy of the colliding nucleons, $s = (p_1 + p_2)^2$. These transport coefficients D , A_L and A_T represent the mean value of longitudinal momentum transfer, the variances of the longitudinal and transverse momentum transfers in a binary collision, respectively. Eqs. (3.3 and 3.4) constitute a stochastic description for the space-time evolution of the collision process in terms of the fluid dynamical variables. These equations involve the relativistic mean-fields, which are self-consistently determined via the nuclear matter characteristics.

4. Equilibration of Counter-Streaming Fluids

As an application of the two-fluid approach, we investigate a simple situation of spatially homogeneous, counter-streaming nuclear fluids at normal density, n_0 , without the mean-fields [9]. In order to estimate the magnitude of local fluctuations, we introduce a coarse-graining over cells of correlation volume V , and consider the integrated variables over the correlation volume,

$$\int_V d^3x J_{\delta}^{(\alpha)} = N^{(\alpha)} \quad (4.1)$$

$$\int_V d^3x T_{\delta\mu}^{(\alpha)} = N^{(\alpha)} \epsilon^{(\alpha)} u_{\mu}^{(\alpha)} = N^{(\alpha)} P_{\mu}^{(\alpha)} \quad (4.2)$$

where $N^{(\alpha)}$ is the number of nucleons in a cell, $\epsilon^{(\alpha)}$ and $P_{\mu}^{(\alpha)}$ denote the average internal energy and the energy-momentum flux per nucleon in the α -fluid. The correlation volume can be estimated to be $V = 2\sigma_{\text{tot}} \ell_{\text{tr}}$, where σ_{tot} is the total cross-section and the second factor is determined in terms of transport cross-section, $\sigma_{\text{tr}} = 4D/\sqrt{\lambda}$, and the total nucleon density, $n = J_{\delta}^{(p)} + J_{\delta}^{(t)}$, as $\ell_{\text{tr}} = 1/(n \sigma_{\text{tr}})$. Since the coarse-grained cells are assumed to be uncorrelated, the number of nucleons, $N^{(\alpha)}$, and the sum of energy-momentum flux, $P^{(p)} + P^{(t)}$, in each cell are conserved quantities, and the relative energy-momentum flux $P_{\mu} = P_{\mu}^{(p)} - P_{\mu}^{(t)}$ is a fluctuating quantity and determined by a Langevin equation. In equal-velocity frame the equation for P_{μ} is given by

$$\frac{d}{dt} P_{\mu} = -\frac{2\bar{D}}{N} P_{\mu} + \frac{2}{N} \delta R_{\mu} \quad (4.3)$$

where $N = N^{(p)} = N^{(t)}$ and the correlator of the random force is

$$\langle \delta R_{\mu}(t) \delta R_{\nu}(t') \rangle = \delta(t-t') \left[(\bar{A}_L - \bar{A}_T) P_{\mu} P_{\nu} + \bar{A}_T g_{\mu\nu} P^2 \right]. \quad (4.4)$$

Here, the new transport coefficients are related to those given by Eq. (3.8) through

$$(\bar{D}, \bar{A}_L, \bar{A}_T) = (Nn_0/ME_{cm})(D, A_L, A_T) \quad (4.5)$$

where $E_{cm}^2 = p_{cm}^2 + M^{*2}$ is the initial energy per nucleon in the equal-velocity frame. We calculate the transport coefficients employing Cugnon parametrization, which incorporates both the elastic and inelastic nucleon-nucleon cross-sections. Fig. 1 shows the evolution of $\bar{D}$ and $\bar{A}_L$ as a function of an effective energy $E = (s - 4M^{*2})/2M^*$, at an incident energy of 1 GeV/nuc., where the values of E indicated by s_{in} and s_{fin} correspond to the incident energy of 1 GeV/nuc. and the complete stopping, respectively. Regarding the transport coefficients are constant, we can solve eq. (4.3) and find analytical expressions for the mean value and dispersion of P^2 . Fig. 2 shows the time evolution of the mean value and the dispersion of P^2 at an incident energy of 1 GeV/nuc. The calculations indicate that the magnitude of fluctuations associated with P^2 is large and an appreciable fraction of its mean value during non-equilibrium stages of the collision process. The thermal components of the diffusion coefficients, $\bar{A}_L$ and $\bar{A}_T$, are neglected in our calculations. As a result, the dispersion of P^2 goes to zero for large times, rather than saturating at its thermal value.

5. Conclusions

In this work we carry out a reduction of the relativistic BLE to a stochastic two-fluid model and extract transport coefficients, i.e., interflow friction and diffusion coefficients, associated with the fluid dynamical variables. This approach is particularly suitable for studying the role of fluctuations on dynamics during non-equilibrium, two-flux regime of heavy-ion collisions. In order to illustrate the formalism and estimate magnitude of fluctuations, we carry out calculations for a simple situation of two homogeneous, counter-streaming nuclear fluids at normal density. Our calculations indicate that large local fluctuations occur in the non-equilibrium stages of the collision process. It will be

interesting to investigate the influence of local fluctuations in energy-momentum space on the particle production mechanism and on the development of instabilities that may occur during the two-flux regime in heavy-ion collisions.

References

1. H. T. Elze, et al., *Mod. Phys. Lett. A* **2** (1987) 451.
2. Y. B. Ivanov, *Nucl. Phys. A* **474** (1987) 693, in "Relativistic Heavy-Ion Physics," eds. L. P. Csernai and D. Strottman, Singapore, World Scientific (1991).
3. B. Blättel, et al., *Phys. Rev. C* **38** (1988) 1767.
4. W. Bottermans and R. Malfliet, *Phys. Lett. B* **215** (1988) 617.
5. Q. Li, J. Q. Wu and C. M. Ko, *Phys. Rev. C* **39** (1989) 849.
6. M. Cubero, et al., *Nucl. Phys. A* **519** (1990) 345c.
7. S. J. Wang, et al., *Ann. of Phys.* **209** (1991) 251.
8. A. A. Amsden, et al., *Phys. Rev. C* **17** (1978) 2080.
9. Y. B. Ivanov, et al., *Z. Phys. A* **340** (1991) 385.
10. S. Ayik and C. Gregoire, *Phys. Lett. B* **212** (1988) 269; *Nucl. Phys. A* **513** (1990) 187.
11. J. Randrup and B. Remaud, *Nucl. Phys. A* **514** (1990) 339.
12. S. Ayik, *Phys. Lett. B* **265** (1991) 47.
13. S. Ayik, et al., Proc. Ninth Workshop on Nuclear Dynamics Key West, Florida (1993), and in preparation for publication.

Fig. 1. Transport coefficients as a function of an effective energy $E = (s - 4M^2)/2M$ at an incident energy of 1 GeV/nuc.

Fig. 2. Time evolution of the mean value and dispersion of p^2 at an incident energy of 1 GeV/nuc.

B. **NUCLEAR FISSION WITH A LANGEVIN EQUATION**
(with D. Boilley, E. Suraud and Y. Abe)

ABSTRACT

A Langevin equation with memory effects is applied to investigate thermal fission. A strong two-body friction coefficient, which is estimated in terms of microscopic quantities, tends to decrease the stationary limit of the fission rate and increase the transient time. The fission rates calculated at different temperatures are compared with other available results.

1. **Introduction**

There are mainly two competing ways of de-excitation of a hot compound nucleus, neutron evaporation and fission. Here, we consider fission. In 1940, Kramers, in a cornerstone paper [1], proposed to describe fission as a diffusion process over the fission barrier and to use a Langevin Equation (LE) to study this phenomenon. In such a description, the slow collective motion with its large inertia behaves like a Brownian particle and the fast nucleonic degrees of freedom provide the heat bath. He performed a study of the stationary flow over the saddle point and derived a formula for the stationary fission rate. More recently, a complete study of the fission process was performed numerically with both the LE and the Fokker-Planck Equation [2,3]. A long transient time that could allow more precision neutrons to evaporate, was pointed out by both groups. These calculations were completely phenomenological in which all the parameters of the model were fitted to reproduce the experimental data. Recently, a derivation of LE has been presented from the microscopic Boltzmann-Langevin Model (BLM), in which transport coefficients are expressed in terms of nuclear matter properties [4]. Here, we use this model to investigate the influence of the memory effect and the large two-body friction on the fission dynamics.

2. **A Langevin Equation for Collective Motion**

The microscopic BLM provides a basis for deriving transport models for large amplitude collective motion. In ref. [4], derivation of a collective transport model was

presented in a diabatic approximation assuming the collective motion is characterized by an irrotational-incompressible velocity field

$$\mathbf{v}(\mathbf{r}, t) = \dot{q}(t) \nabla \phi(\mathbf{r}) \quad (2.1)$$

and carrying out a fluid dynamical reduction. This gives rise to a memory dependent LE for the collective variable $q(t)$, which is usually called as a generalized Langevin Equation (GLE),

$$M\ddot{q} + \frac{1}{2} \frac{dM}{dq} \dot{q}^2 + \frac{dV}{dq} = - \int_{-\infty}^t dt' \gamma(t-t') M \dot{q}(t') + \delta F(t) \quad (2.2)$$

where the friction kernel and the correlation function of the random force are given by

$$\gamma(t-t') = \frac{\Gamma}{M} \exp\left(-\frac{t-t'}{\tau}\right) \quad (2.3)$$

$$\langle \delta F(t) \delta F(t') \rangle = TM \gamma(t-t'). \quad (2.4)$$

Here the relaxation time τ , the inertia M and the quantity Γ are determined in terms of nuclear matter properties according to (using the notation of ref. [4])

$$M(q) = m \int d\mathbf{r} \rho_0(\mathbf{r}, q) \nabla \phi \cdot \nabla \phi \quad (2.5)$$

$$\Gamma = \frac{24}{5} A \epsilon_F T \xi \quad (2.6)$$

$$\frac{1}{\tau} = 8\sigma v_F \rho \left(\frac{T}{\epsilon_F}\right)^2 \quad (2.7)$$

with ξ as the average of the quantity $\Sigma[\nabla_i \nabla_j \phi(\mathbf{r})][\nabla_i \nabla_j \phi(\mathbf{r})]/6$ over density, and ϵ_F and v_F denote the Fermi energy and velocity, respectively. Here, the relaxation time τ is given for the quadrupolar mode [5].

One of the main differences between Eq. (2.2) and the phenomenological LE is the presence of memory dependent damping in Eq. (2.2). In the limit of sufficiently high temperature T , the relaxation time becomes shorter and one recovers a Markovian LE with $\gamma(t-t') \simeq \beta \delta(t-t')$ and $\beta = \tau\Gamma/M$. By following the individual trajectories of the GLE, we can investigate the influence of the memory effect on the fission dynamics. Without the random force, the solutions of Eq. (2.2) can be given analytically as,

$$q(t) = A \exp\left[-\frac{\pm \omega_0^2 t}{\beta \pm \omega_0^2 \tau}\right] + B \exp\left[-\frac{t}{2\tau}\right] \cos(\omega t + \varphi) \quad (2.8)$$

with

$$\frac{1}{M} \frac{dV}{dq} = \pm \omega_0^2 q \text{ and } \omega = \sqrt{\left[\frac{\beta}{\tau} \pm \omega_0^2\right] - \frac{1}{4\tau^2}} \quad (2.9)$$

where A , B and φ are determined by the initial conditions. In addition to the drift term, some damped oscillations appear in the solutions. As seen from fig. 1, the memory dependent random force reactivates these oscillations with much larger amplitudes than those obtained with a Markovian random force. As a result, the memory effect modifies the time evolution of the individual events which can have a large influence on the fission rates as we discuss below.

3. Application to Fission

We investigate time evolution of a symmetric fission in terms of quadrupole moment $q(t)$ of the density distribution with a quadrupolar velocity field $\phi(r) = (2z^2 - x^2 - y^2)/2$. For a numerical solution, it is more convenient to write the equation of motion (2.2) in the following form,

$$\begin{aligned} \dot{q} &= \frac{p}{M} \\ \dot{p} &= -\frac{dV}{dq} + \frac{1}{2} \left[\frac{p}{M} \right]^2 \frac{dM}{dq} + F \end{aligned} \quad (3.1)$$

$$\dot{F} = \frac{1}{\tau} [-\beta p - F + \sqrt{M\beta T} w(t)]$$

where $\beta = \tau\Omega^2$ with Ω as the frequency of giant quadrupolar vibrations, and $\langle w(t) w(t') \rangle = 2\delta(t-t')$. We approximate the potential energy along the fission path in terms of two parabola as

$$V(q) = \begin{cases} 135.5 q^2 & q < 0.122 \\ -135.5(q-0.24)^2 + 4 & q > 0.122 \end{cases} \quad (3.2)$$

which gives the appropriate position of the saddle point and the fission barrier of a heavy nuclei ($A \simeq 250$) consistent with the liquid drop model. We generate a large number of events by solving the GLE (3.1) numerically. The initial conditions of p and q are

randomly determined from a cold Gaussian

$$d(p,q) = \exp\left\{-\frac{1}{T_0} \left[\frac{p^2}{M} + 135.5q^2\right]\right\} \quad (3.3)$$

with $T_0 = 0.3$ MeV. The time dependent fission rate is calculate according to

$$r(t) = \frac{1}{P(t)} \frac{dP(t)}{dt} \quad (3.4)$$

where $P(t)$ denotes the fraction of the events that corresponds to the compact configuration, i.e., with a quadrupole deformation behind the saddle point $q < q_s$.

4. Results and Conclusions

We first calculate the fission rate of ^{248}Cf at a temperature of $T = 4.88$ MeV. Due to a different density parameter, this temperature corresponds to 4 MeV in a similar calculation done in ref. [2]. The relevant parameters corresponding to this temperature are $\beta = 0.1$ c/fm $= 30 \times 10^{21}\text{s}^{-1}$ and $\tau = 33$ fm/c $= 1.1 \times 10^{-22}\text{s}$. In this case, the relaxation time τ is sufficiently short and we carry out the calculations in a Markovian approximation. Since the friction coefficient is large, six times larger than those employed in ref. [2], we perform the calculation in the overdamped limit as well. By neglecting $\dot{p}$ and $\dot{F}$ in the overdamped limit in Eq. (3.1), the equation of motion reduces to a simple form,

$$\dot{q} = -\frac{1}{M\beta} \frac{dV}{dq} + \sqrt{\frac{T}{M\beta}} w(t). \quad (4.1)$$

The results of calculations are shown in fig. 2. Due to large friction, the stationary fission rate is smaller and the transient time is longer than those of ref. [2], but the order of magnitude of these quantities are the same in both cases. Also, the results of the calculations in the overdamped limit are close to the other ones.

At a lower temperature of $T = 3\text{MeV}$, the relevant parameters are $\beta = 0.27$ c/fm $= 80 \times 10^{21}\text{s}^{-1}$ and $\tau = 88$ fm/c $= 3 \times 10^{-22}\text{s}$. In this case, the Markovian approximation is not appropriate; we use the full GLE to evaluate the fission rate. The results are shown in fig. 3. They are not accurate due to small statistics, but we can extract the bulk behavior. The stationary fission rate is about ten times smaller and the transient time is ten times longer than those obtained with the phenomenological parameters [5]. These differences

are mainly due to a large two-body friction employed in the present calculations. A more accurate determination of the two-body friction coefficient can be carried out by incorporating the memory effect into the microscopic BLE [6]. Then, the relaxation time associated with quadrupolar vibrations is modified according to

$$\frac{1}{\tau} = 8\sigma v_F \rho \left[\left[\frac{T}{\epsilon_F} \right]^2 + \left[\frac{\hbar\Omega}{2\pi \epsilon_F} \right]^2 \right] \quad (4.2)$$

which reduces the two-body friction by about a factor of 2 in the temperature range that we consider.

In this work, we carry out a stochastic calculation of thermal fission in a diabatic approximation by incorporating only the two-body dissipation mechanism. These approximations may be reasonable for sufficiently high temperatures, but not very realistic at low temperatures. At low temperatures, the fission process is close to adiabatic rather than diabatic, in which one-body dissipation plays a dominant role. Therefore, for an accurate description of thermal fission, both the one-body and the two-body dissipations must be incorporated into the calculations.

References

1. H. A. Kramers, *Physica VII*, 4 (1940) 284.
2. P. Grange, J. Q. Li and H. A. Weidenmuller, *Phys. Rev. C* **27** (1983) 2063.
3. Y. Abe, C. Gregoire and H. Delagrangé, *J. Phys. (Paris)*, C4, **47** (1986) 329.
4. S. Ayik, E. Suraud, J. Stryjewski and M. Belkacem, *Z. Phys.* **A337** (1990) 413.
5. B. W. Bush, G. F. Bertsch and B. A. Brown, *Phys. Rev. C* **45** (1992) 1709.
6. S. Ayik and D. Boilley, *Phys. Lett.* **B276** (1992) 263.

figure 1: Effect of the Langevin force on a single trajectory, in the Markovian case and the non-Markovian case. The initial condition is far beyond the saddle point. The oscillations are a signature of the memory effect.

$T = 4 \text{ MeV}$

Figure 2: Fission rate as a function of time for the ^{248}Cf at $T=4.88$ MeV. The calculation was made with a LE in the Markovian approximation and in the overdamped limit. The initial condition distribution is "cold", $T_0=0.3\text{MeV}$. The number of events calculated for both graphs is $N=250\,000$. The results of reference [2] with $T=4\text{MeV}$ and $\beta=5 \times 10^{21}\text{s}$ are indicated. The stationary limit is calculated analytically [6].

Figure 3: Fission rate as a function of time for the ^{248}Cf at $T=3$ MeV. The calculation was made with the GLE and in the overdamped limit. The result of a calculation made in the phenomenological case of the reference [2] for $\beta=5 \times 10^{21} \text{ s}$ and at $T=2.46 \text{ MeV}$, but at the same excitation energy, is also shown. The initial condition distribution is "cold", $T_0=0.3 \text{ MeV}$. The stationary limit is calculated analytically [6]. N is the number of events calculated.

APPENDIX I

Papers

"Damping of Collective Vibrations in a Memory-Dependent Transport Model," S. Ayik and D. Boilley, *Phys. Lett. B* **276** (1992) 263.

"Applications of Boltzmann-Langevin Equation to Nuclear Collisions," E. Suraud, S. Ayik, M. Belkacem and J. Stryjewski, *Nucl. Phys. A* **542** (1992) 141.

"Boltzmann-Langevin Model for Nuclear Collisions," S. Ayik, E. Suraud, M. Belkacem and D. Boilley, *Nucl. Phys. A* **545** (1992) 35c.

K⁺ Production Far Below Free Nucleon-Nucleon Threshold in Heavy-Ion Collisions, M. Belkacem, E. Suraud and S. Ayik, *Phys. Rev. C* (in press).

"Nuclear Fission with a Langevin Equation," D. Boilley, E. Suraud, Y. Abe and S. Ayik, *Nucl. Phys. A* (in press).

Seminars

"The Boltzmann-Langevin Approach," GSI, Darmstadt, Germany (1992).

"The Boltzmann-Langevin Approach," University of Munich, Munich, Germany (1992).

Conference Contributions

"The Boltzmann-Langevin Model for Nuclear Collisions," S. Ayik, E. Suraud, M. Belkacem and D. Boilley, Int. Workshop on Dynamical Fluctuations and Correlations in Nuclear Collisions, Aussois, France (1992).

"Stochastic Two-Fluid Model for Relativistic Heavy-Ion Collisions," S. Ayik, Y. B. Ivanov, V. N. Russkikh and W. Nörenberg, Winter Workshop on Nuclear Dynamics, Key West, Florida (1993).

APPENDIX II PERSONNEL

Principal Investigator

S. Ayik, Professor

Sources of Support

TTU, DOE, ORAU

Collaborators

E. Suraud*, Research Staff
M. Belkacem, Graduate Student
D. Boilley*, Graduate Student
Y. Abe, Professor
Y. B. Ivanov, Research Staff

GANIL, Caen, France
GANIL, Caen, France
GANIL, Caen, France
YI, Kyoto, Japan
GSI, Darmstadt, Germany
and KI, Moscow, Russia
GSI, Darmstadt, Germany
and KI, Moscow, Russia
GSI, Darmstadt, Germany

V. N. Russkikh, Research Staff

W. Nörenberg, Professor

Secretary/Account Clerk

G. J. Julian, CPS

TTU

Other Nuclear Physics Faculty at TTU

M. M. Hindi, Associate Professor
R. L. Kozub, Professor
K. Kumar, University Professor
D. P. Murdock, Assistant Professor
S. J. Robinson, Assistant Professor
P. B. Semmes, Associate Professor
J. F. Shriner, Associate Professor
J. C. Wells, Professor

TTU, DOE
TTU, DOE
TTU
TTU, IU
TTU
TTU, DOE
TTU, DOE
TTU, ORAU

DOE = Department of Energy
IU = Indiana University
ORAU = Oak Ridge Associate Universities
TTU = Tennessee Technological University
YI = Yukawa Institute for Theoretical Physics
GSI = Gesellschaft für Schwerionenforschung
KI = Kurchatov Institute

*Present address: University of P. Sabatier, Toulouse, France.

END

DATE
FILMED

5 / 27 / 93