MARTIN MARIETTA

Received
MAY 11 1992

ORNL/ER-47

**ENVIRONMENTAL
RESTORATION
PROGRAM**

**Suggestions for Improvement of the
Methodology and Use of MEPAS,
the Multimedia Environmental
Pollutant Assessment System**

L. Shevenell
F. O. Hoffman

ENERGY SYSTEMS

MANAGED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from 615-576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

Environmental Restoration Division
ORNL Environmental Restoration Program

Suggestions for Improvement of the Methodology and Use of MEPAS, the Multimedia Environmental Pollutant Assessment System

L. Shevenell
F. O. Hoffman

Manuscript Completed--October 1991
Date Issued--April 1992

Prepared by
Environmental Sciences Division
Oak Ridge National Laboratory
ESD Publication 3723

Prepared for
U.S. Department of Energy
Office of Environmental Restoration and Waste Management
under budget and reporting code EW 20

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6285
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Author Affiliations

L. Shevenell and F. O. Hoffman are members of the Environmental Sciences Division, Oak Ridge National Laboratory, Martin Marietta Energy Systems, Inc.

CONTENTS

TABLES AND FIGURE	v
EXECUTIVE SUMMARY	vii
ABSTRACT	ix
1. INTRODUCTION	1
2. GENERAL COMMENTS	4
3. MEPAS METHODOLOGY	5
3.1 USE OF MEPAS	7
3.2 MEPAS OUTPUT	8
4. COMPARISONS WITH OTHER RISK ASSESSMENT METHODOLOGIES	9
5. RECOMMENDATIONS	15
6. CONCLUSIONS	17
ACKNOWLEDGMENTS	17
REFERENCES	18

TABLES

1	Hazard/cancer risk factors for various population sizes, all yielding an HPI of 20	5
2	Pathways considered by MEPAS and ORNL/ER-9	11
3	Transfer coefficients of contaminant to milk (d/L)	12
4	Soil-to-plant transfer factors	12
5	Transfer coefficient of contaminant to meat (d/kg)	13
6	Bioaccumulation factors for freshwater fish	13

FIGURE

1	Site location map for the ORNL WAGs	3
---	---	---

EXECUTIVE SUMMARY

The Multimedia Environmental Pollutant Assessment System (MEPAS) was developed at Pacific Northwest Laboratories for the purpose of providing a consistent way to rank the relative risks associated with waste sites at Department of Energy facilities in the United States. These rankings are being used to prioritize sites for remediation activities. The studies documented in this report were conducted to evaluate the reliability of the rankings of waste sites obtained with the use of MEPAS. Although this study is limited in scope, it considered both the internal calculations and default values of MEPAS and the use of MEPAS at a specific waste site [Waste Area Grouping (WAG) 2, Oak Ridge National Laboratory].

The major findings of this study indicate that the use of the final hazard potential index (HPI) produced by MEPAS should be discontinued and that user input must be much more consistent and considerably improved. In general, the methodology and internal calculations of MEPAS are sound; yet, actual risks to human populations become obscured in the final HPI number. Although it may be desirable to obtain one, final numerical value associated with risks, arriving at such a value is not practicable, given the current state of the art in risk assessment. The use of one number such as the HPI is not recommended because those conducting the prioritization cannot have a full appreciation of the meaning of the number and the relevant issues associated with it. The final results of the MEPAS scheme should be altered to eliminate the HPI and to include risks to the maximally exposed individual, average individual, and population (noting the population size) so that users can evaluate and weight these risks. Reliable decisions on prioritization cannot be made without this information.

The other required improvement is user input to the model. Evaluation of one waste site, WAG 2, indicates that the selection of pathways to be considered and of exposed population sizes was often arbitrary. The evaluation also indicates that the MEPAS code was not used to its full potential in some cases. To fully understand the site and associated conditions, users must take more time to develop input parameters. Without realistic input to any model, one can only expect unreliable results.

ABSTRACT

The Multimedia Environmental Pollutant Assessment System (MEPAS) has been evaluated for the purpose of determining if the current ranking of waste sites is realistic and reliable. There are two main reasons for the uncertainty of the rankings identified in this study: the use of the hazard potential index (HPI) and the user input. The HPI contributes to unreliable rankings because risks to human populations are summarized in a single numerical value. A final result from MEPAS should include risks to the maximally exposed individual, average individual, and the population (noting the population size) so that users can evaluate and weight these risks.

Examination of user input indicates that exposure pathways and exposed populations were sometimes arbitrarily selected. Users must be better informed about the characteristics of the waste site and its potential interaction with human populations for realistic input to the MEPAS model to be developed. Without realistic and consistent input to the MEPAS, or any model, reliable results cannot be expected and any prioritizations based on the results will be questionable.

1. INTRODUCTION

The Multimedia Environmental Pollutant Assessment System (MEPAS) (Droppo et al. 1989; Whelan et al. 1987) has been used to rank the various waste areas at Department of Energy (DOE) facilities in the United States. The ranking effort was directed by DOE's Office of Environmental Audit, and technical support was provided by NUS Corporation, Washington, D.C. An overview of the preliminary environmental surveys of waste sites at Oak Ridge National Laboratory (ORNL) is found in USDOE (1991). No single, comprehensive document describing the results of the MEPAS rankings has been published, yet some results are discussed in this report. The rankings were conducted to identify waste sites that pose immediate risk to public health. The results produced by MEPAS were used to prioritize sites, the ultimate objective being to allocate characterization and remediation funds among sites (or groups of sites) that have previously been identified as potentially hazardous. The screening of sites and the setting of action levels and cleanup goals are not MEPAS objectives. MEPAS is used solely to prioritize sites. A model used to prioritize sites should do the following:

- Consider both off-site and on-site (intruder) risks.
- Consider both measured and modeled sources.
- Contend with chemical carcinogens, radionuclides, and noncarcinogens consistently. Results must be expressed in comparable terms and degrees of conservatism must be similar.
- Deal with multiple pathways consistently. Results must be expressed in comparable terms and degrees of conservatism must be similar.
- Include differences in uncertainty between risk estimates for different contaminant types or pathways.
- Allow combination of pathways and stressor types so that sites can be compared on a common scale.
- Use site-specific information where it exists and is relevant and incorporate the most up-to-date generic values (i.e., transfer coefficients) for parameters that are not site-specific.

A specific risk assessment using MEPAS was conducted by NUS Corporation on the Waste Area Groupings (WAGs) at Oak Ridge National Laboratory (USDOE 1991). In this report, we focus on the WAG 2 assessment to identify potential shortcomings in the methodology and use of the MEPAS code. WAG 2 is a man-made lake (White Oak Lake) which was dammed in order to provide a settling basin for any sediment-bound contaminants that might discharge into surface waters from the adjacent WAGs. Figure 1 illustrates the

position of WAG 2 relative to the nearby WAGs and also shows the location of WAG 1, which is the main ORNL plant area.

The purpose of this study is to determine if MEPAS suitably meets the requirements in the preceding list for a reliable prioritization model. Although this study is rather limited in scope, it is used to identify various deficiencies associated with the MEPAS assessment of WAG 2 that can be attributed to either the improper use of MEPAS or to the specific methodology and default values used. The results of this study are intended to help improve the MEPAS risk assessment methodology as well as to identify issues related to risk assessment in general. Following identification of concerns to be addressed in the use and methodology of MEPAS, several recommendations intended to improve the overall risk assessment and ranking activities being conducted at ORNL are provided.

The evaluation of MEPAS as it relates to WAG 2 revealed that several improvements can be made to this risk assessment scheme. Each of the following practices leads to misleading rankings of hazardous sites or confusing results:

1. User selection of contaminants and pathways to be considered was often unjustified.
2. A single numeric value is used to represent the diverse risks associated with radionuclides, carcinogens, and noncarcinogens.
3. The hazard ranking is distorted by using a logarithmic scale in the final ranking scheme.
4. Selected default values for exposure times, bioaccumulation factors, and transfer coefficients are not those typically used for screening calculations and are not applicable for site-specific assessments.
5. Risk factors were normalized for radionuclides, noncarcinogens, and carcinogens for a risk level of 10^{-6} .

Fig. 1. Site location map for the ORNL WAGs.

2. GENERAL COMMENTS

Although this report is intended to outline the suitability of MEPAS in determining priorities and to evaluate risk assessment schemes in general, it is noted that MEPAS generally utilizes widely accepted equations for transport and exposure estimation that are commonly used in other risk assessment schemes (i.e., IAEA 1982; NCRP 1989; Hoffman et al. 1990; CAPCOA 1990; NCRP 1991).

Because the objective is to identify realistic risks to the general population, MEPAS methodology estimates off-site health risks. Calculation of off-site contaminant concentrations based on on-site concentrations can result in unrealistic contaminant concentrations at usage locations. Yet, in using measured off-site contaminant concentrations, it is often impossible to distinguish between particular sources of contaminants and to identify their ultimate source. In MEPAS, off-site concentrations are generally calculated based on the flux from a particular ranking unit (RU), that is, a waste area grouping, contaminated site, etc., and risks associated with a specific RU are based on these estimated concentrations. When the objectives of a study are to prioritize specific waste sites (waste sources) for cleanup, the MEPAS method is preferable to methods that don't distinguish among the risks associated with particular RUs. However, if measured off-site contaminants can be directly attributed to a specific RU, these values should be used, and MEPAS is equipped to handle this situation. In addition, on-site risks could be estimated by MEPAS, if desired, using measured concentrations at the site.

The following terms specific to MEPAS are used throughout this report:

- RU = ranking unit; a waste site or source of contaminants.
- HPI = hazard potential index; $10 \log_{10} \text{PI}$.
- PI = preliminary hazard potential index; equals the sum over all contaminants and pathways of the product of exposed population and RF.
- RF = risk factor: for noncarcinogens, dose/reference dose; for radionuclides, $270(\text{dose})$, or 2.7×10^{-4} per rem (dose in rem)/ 10^{-6} ; and for carcinogens, (cancer potency factor)(dose, in milligrams per kilogram per day)/ 10^{-6} .
- RfD = reference dose.

3. MEPAS METHODOLOGY

The equations used in the MEPAS ranking of WAG 2 in relation to the other Oak Ridge Reservation sites are generally standard, well-accepted equations used in transport modeling, exposure pathway modeling and risk assessment. Nevertheless, improvements could be made in the MEPAS formulation. By far, the largest and most easily remedied problems with MEPAS are the limited output from the model and the use of the hazard potential index (HPI), which distorts the rankings of waste sites. For instance, the HPI uses a logarithmic scale [$HPI = 10 \log_{10} PI$, where PI, the preliminary hazard potential index, is the sum over all pathways of the product of risk factor (RF) and exposed population]. The logarithmic scale compresses the final ranking scale and, hence, tends to distort it. As an example, an HPI of 10 corresponds to a PI of 10, while an HPI of 100 corresponds to a PI of 10^{10} . Thus, it becomes more difficult to distinguish between risks associated with particular sites that have similar HPI values. Similar values of HPI (i.e., 35 and 40) could be construed to represent similar levels of risk, yet the PIs associated with these values differ by approximately a factor of 3 (i.e., 3.2×10^3 and 1×10^4).

Further complicating the matter is the incorporation of population size associated with each exposure pathway without representing to users of the output the risk and the population size used to obtain the final HPI. Naturally, the population associated with one exposure pathway will not necessarily be the same as that used for other exposure pathways at the same usage location. Hence, any particular HPI value can have several interpretations. The RFs used in MEPAS are a function of the calculated dose as a result of exposure over several pathways and either a reference dose (RfD), a cancer potency factor, or lifetime dose equivalent (for radionuclides). The RFs used in MEPAS for carcinogens are normalized to a 10^{-6} risk and are a relatively consistent measure of risk; yet, these are combined with population in MEPAS to obtain an HPI. As an example of the distortion that this practice causes, note in Table 1 that several RFs and population combinations produce the identical HPI of 20.

Table 1. Hazard/cancer risk factors for various population sizes, all yielding an HPI of 20

Noncarcinogen		Carcinogen		Population size	PI	HPI
Hazard index	RF	Cancer risk	RF ^a			
0.1	0.1	10^{-7}	0.1	1000	100	20
1.0	1.0	10^{-6}	1.0	100	100	20
10	10	10^{-5}	10	10	100	20
100	100	10^{-4}	100	1	100	20

^aThis is the MEPAS normalized risk factor = $10^6(\text{cancer risk})$.

Hence, a relatively high (10^{-4}) risk to one person is assigned the same HPI as a 10^{-7} risk to 1000 people, which is below the level of concern. Because the HPI results can be easily

misinterpreted, a printout containing columns for risk, population size, and PI for each contaminant and pathway would greatly assist anyone in the interpretation of relative risk between sites.

The calculated RFs of noncarcinogens are somewhat misleading in MEPAS because the hazard index of 1.0 (dose in milligrams per kilogram per day) divided by an RfD is assumed to be weighted the same as a 10^{-6} cancer risk. The risk factor for radiation is also calculated as a ratio of the individual lifetime dose and the dose corresponding to a health risk of 10^{-6} (i.e., 0.0037 rem = 10^{-6} lifetime risk), whereas risk from exposure to carcinogens is estimated using cancer potency factors divided by 10^{-6} . Although these practices tend to normalize the RFs calculated for the three types of exposures to a 10^{-6} risk, the comparison of risk factors between radionuclides, carcinogens, and nonradionuclide noncarcinogens can be difficult. A noncarcinogen RF (or hazard index) of 0.1 indicates that the particular dose of a given contaminant would be of no concern to an individual's health. However, an RF for noncarcinogens of 100 would indicate substantial risk to individual health for a given dose; yet, this is equated with a 10^{-4} cancer risk, which is not distinguishable from background (10^{-3}) cancer risk and is of little concern. Hence, the two types of risk are not directly comparable. In MEPAS, all three types of risk factors are calculated separately and their respective HPI values can be compared. Ultimately, all three types of risk are combined with population and summed over all pathways such that the risks from radionuclides, carcinogens, and noncarcinogens are combined into a final HPI value. Hence, interpretation of the final HPI is difficult because the exact values of each type of risk and their weights (population size) are unknown.

RFs associated with exposure to radionuclides, carcinogens, and noncarcinogens are all risks which are normalized to a risk level of 10^{-6} to have a common basis for comparison. This practice necessitates the assumption that exposure at the RfD corresponds to a health risk of 10^{-6} for noncarcinogens and that a linear dose-response relationship exists for exposures that exceed the reference dose. Thus, an exposure to a noncarcinogen that exceeds the RfD by a factor of 1000 will be equivalent to a carcinogen risk estimate of 10^{-3} . A cancer risk of 10^{-3} is not detectable in human populations, for which the spontaneous lifetime cancer incidence rate is approximately 2×10^{-1} . However, for many noncarcinogens, exceeding the reference dose by a factor of 1000 may lead to serious, observable, deleterious effects in humans. Although, in reality, exceeding the RfD by 1000 is not likely because of decreases in source concentrations by processes such as dilution and adsorption, the assumption that the RfD corresponds to a 10^{-6} risk makes comparisons between carcinogens and noncarcinogens unrealistic. From the previous discussion, it is clear that, by this method of comparison, a 10^{-3} risk associated with a carcinogen is relatively innocuous compared with a 10^{-3} risk from noncarcinogens.

The remaining criticisms are more specific and are generally associated with particular exposure pathways. The maximum individual exposure resulting from consumption of fish in MEPAS is underestimated because most of the people who eat substantial quantities of fish from this usage location are fisherman, who make up a small percentage of the population. The default consumption rate of 0.006 kg/d is probably too low for the maximum individual exposure expected for fishermen who eat the fish but may be a realistic average for the population as a whole. This value is especially low for the evaluation of exposure periods that are only on the order of a few years. Similarly, MEPAS uses an average exposure rate of 12 h per year as a default value (as suggested by USNRC 1977) for each of the following:

fishing, boating, and swimming. In reality, a smaller percentage of the population tends to spend a much greater portion of their time in these activities (i.e., local residents). Hence, MEPAS is not actually calculating the maximum individual risk in these cases but an average risk.

MEPAS lists values identified as maximum individual risk values; however, the calculations are based on average exposure and consumption rates. Hence, the reported values are not truly maximum individual risks but risks associated with the average individual exposed. The values listed in the outputs should be labeled as average individual risks to avoid confusion and misinterpretation. If values for maximum individual risks are desired, more conservative exposures and consumption rates should be incorporated into MEPAS. In addition, because of different exposure averaging times and exposure durations, different consumption rates should be used for carcinogens and noncarcinogens.

MEPAS does not consider dredging or soil ingestion by animals. Inclusion of these two would increase the contaminant concentration in the soils, as well as in the meat and milk products of the animals that inadvertently consume soil. Neglect of these two processes by MEPAS results in low values of calculated RF.

Many of the water treatment purification factors used by MEPAS range from 0.5 to 1.0. These values tend to slightly overestimate the actual posttreatment contaminant concentration because it is likely that many radionuclides are removed from the water by adsorption onto particulate matter, which is later filtered out of solution in the purification process. When values for maximum individual exposures are desired, it may not be necessary to alter these values.

Contaminant concentrations in surface waters tend to be overestimated by MEPAS because complexing, precipitation and adsorption are not considered in the model. Because estimations of lifetime risks associated with the contaminants are often sought in risk assessments, this is a somewhat desirable deficiency. Often, contaminants have been released from a RU for periods of time prior to risk modeling; consequently, concentrations at receiving locations can be much greater than calculated values because of long-term exposure of water to contaminants at the site and accumulation of contaminants in sediment. However, MEPAS attempts to estimate current and future risks associated with particular waste sites, and these values are not directly attributable to a particular waste site in its current condition. A separate risk analysis or screening process would be required to estimate actual risks associated with exposures at particular contaminated usage locations.

3.1 USE OF MEPAS

The previous section outlined the major deficiencies in the MEPAS formulation. In many cases, these deficiencies will be overwhelmed by those caused by user error. Without realistic, reliable input to MEPAS for each waste site being ranked, results of the ranking process will be unreliable. The following observations were made in the evaluation of the WAG 2 ranking, yet it is quite likely that other input and user errors would be identified if the rankings and results from other waste areas were scrutinized.

The users of MEPAS did not consider arsenic (As) or polychlorinated biphenyls (PCBs) in fish at WAG 2, both of which could contribute significantly to the risk associated with discharges from this RU. The PCBs were associated entirely with WAG 1 (the main plant area of ORNL). Both the types of contaminants and their concentrations at specific RUs must be determined, and input that is as realistic as possible must be provided to MEPAS for more meaningful rankings to be obtained. The neglect of PCBs in fish and As at WAG 2 may have resulted in an unrealistic ranking of this site.

Users of MEPAS at WAG 2 (and all other ORNL sites studied) only considered selected exposure pathways at each usage location rather than all possible pathways. The selection of the pathways considered in the study of WAG 2 does not appear to have any foundation. More pathways were considered for the Clinch River receptor (eight pathways) than were considered for the K-25 Site, Kingston, or Rockwood receptors (two pathways each); hence, the Clinch River receptor ranked highest (No. 1) in MEPAS for WAG 2 because the RFs were summed over a greater number of pathways than at the other locations. Users must select the pathways to be modeled in a consistent manner, such that the selection itself is not the cause of a higher estimated risk at one site than at another. Provisions in the formatting of user input should be adjusted in MEPAS so that comments that explain the rationales for selection of particular pathways can be included.

3.2 MEPAS OUTPUT

Several types of output are generated by MEPAS, some of which appear to be of limited use. For instance, one listing includes the "maximum individual risk values divided by the RfD ratios" for radionuclides, in which RfD is undefined for radionuclides because it is a value used in noncarcinogen evaluation. The values listed in MEPAS output are actually the RF times 10^{-6} (i.e., risks that are not normalized). We recommend that the unaltered risk estimates be reported for maximally exposed individuals. Although "maximum" individual exposures are listed, they are lumped by usage location, making it impossible to determine which usage locations are associated with the higher risks for particular contaminants and pathways considered. In addition, as stated previously, these values do not take into account maximally exposed individuals.

4. COMPARISONS WITH OTHER RISK ASSESSMENT METHODOLOGIES

Comparisons with other risk assessment methodologies (NCRP 1991; Hoffman et al. 1990) were made to evaluate differences in computed risks at specific usage locations and to help determine how MEPAS might be improved. The results in ORNL/ER-9 (Hoffman et al. 1990) are based on the formulations presented in NCRP (1991). Several differences between MEPAS and these models were identified; generally, the calculated MEPAS risks are less conservative than those of ORNL/ER-9.

Although MEPAS and ORNL/ER-9 use similar models and equations describing exposure pathways, they have different applications. The ORNL/ER-9 method provides a biased estimate of exposure associated with specific usage locations. The total, measured concentrations of contaminants at these sites, regardless of origin, are used to estimate screening indices based on health risks. In contrast, MEPAS calculates concentrations at usage locations based on estimated source concentrations. Hence, the MEPAS concentrations attributed to particular usage locations may not equal the total concentration of contaminants at the usage locations because sources of contamination other than the considered waste area may contribute to the total observed concentration at the considered location. MEPAS is designed to rank the relative risk of waste areas, whereas ORNL/ER-9 ranks the relative risks to human health between different usage locations by means of screening indices. Although the objectives of the two methods are somewhat different, it is useful to make comparisons for the purpose of identifying additional areas in which MEPAS could be improved.

The following is a list of some of the differences between the two risk assessment methods.

1. The final product in the MEPAS calculation is the HPI, whereas ORNL/ER-9 uses screening indices (SIs), which are equivalent to unaltered risk estimates in MEPAS (i.e., $RF \text{ values} \times 10^{-6}$ for carcinogens and $RF = SI$ for noncarcinogens).
2. The calculated concentrations at given receptor locations in MEPAS for nonradionuclides are three to four orders of magnitude less than the measured concentrations reported in ORNL/ER-9. The corresponding calculated RF values in MEPAS are three to seven orders of magnitude less than the conservative SI estimates of ORNL/ER-9. The discrepancies in the final results are a combination of the large differences in contaminant concentrations used in the calculations and the fact that the MEPAS users considered fewer exposure pathways and did not consider the possibility of dredged sediment used as agricultural soil.
3. MEPAS defines a health risk for exposure to radionuclides as

$$R_{ij} = 2.7 \times 10^{-4} D_{ij} .$$

The factor 2.7×10^{-4} per rem in MEPAS is less than one-half of the weighted cancer incidence risk conversion factor of 7.2×10^{-4} per rem ($7.2 \times 10^{-2}/Sv$) currently recommended by the ICRP (1990). In ORNL/ER-9, $5.7 \times 10^{-2}/Sv$ was assumed.

4. The calculated concentrations of radionuclides in MEPAS are three to four orders of magnitude less than the measured concentrations reported in ORNL/ER-9 at corresponding sites; yet, the unnormalized RF values are two to five orders of magnitude less than SI values of ORNL/ER-9. The discrepancies between the unnormalized RF and SI can be attributed in large part to the higher concentrations used in the conservative ORNL/ER-9 model, as well as to differences in the combination of transfer coefficients and bioaccumulation factors for specific constituents considered.
5. MEPAS equates a hazard index of 1.0 for noncarcinogens to a cancer risk of 10^{-6} , which is not appropriate. Noncarcinogens are treated as contaminants associated with threshold concentrations or dose limits below which there are no observable health risks. As indicated in the discussions in "MEPAS Methodology" (Sect. 3), evaluation of noncarcinogens should preferably remain separate from that of carcinogens or a different nonlinear scale should be adopted for noncarcinogens as the dose/RfD ratio approaches toxic levels (~ 100).
6. MEPAS uses calculated concentrations at the usage location, whereas ORNL/ER-9 uses measured concentrations at the usage location. Water and fish concentrations in the Clinch River, which are measured in ORNL/ER-9, are generally one to two orders of magnitude higher than those calculated by MEPAS. Although all the ^{137}Cs , which is the main contaminant at WAG 2, can be attributed solely to WAG 2, measured concentrations at the usage location cannot always be attributed to one particular source (a particular WAG). This tends to increase the calculated risk in the ORNL/ER-9 method because the measured concentrations at the usage location is the combined effect of contaminants originating from several sources (RUs). For the purposes of ranking particular waste sources, the MEPAS methodology is more appropriate because it calculates the portion of a total risk at a usage location that can be attributed to a particular waste site.
7. The particular MEPAS calculation for WAG 2 included the nonradionuclides Cr, Pb, Mn, and Hg, only, whereas ORNL/ER-9 investigators considered other contaminants such as As (which may not be an ORNL contaminant) and PCBs in fish for usage locations corresponding to those of WAG 2. The selection of contaminants tends to increase the calculated risk in the ORNL/ER-9 method.
8. MEPAS calculated HPI for selected pathways at each receptor site, whereas ORNL/ER-9 considered a larger range of pathways for each site (see Table 2). Again, this is a user error, not directly attributable to MEPAS, which results in somewhat higher risks calculated by ORNL/ER-9 than those estimated by MEPAS.
9. For screening purposes, ORNL/ER-9 calculates the maximum individual dose based on 70 years of exposure to the maximum measured concentration. MEPAS estimates an "average" concentration based on current discharge from the site and the expected useful lifetime of the site (i.e., 40 years for WAG 2). To ensure correct interpretation, RF results should be listed in the printouts as average individual exposure if the MEPAS methodology continues to be used.

Table 2. Pathways considered by MEPAS and ORNL/ER-9

Usage location	MEPAS exposure pathways	ORNL/ER exposure pathways
Clinch River	Fish ingestion	Fish ingestion
	Vegetable ingestion (leafy)	
	Vegetable ingestion (other)	Vegetable ingestion
	Meat ingestion	Meat ingestion
	Milk ingestion	Milk ingestion
	External exposure	External exposure
	Ingestion of water while swimming	
Kingston, Oak Ridge K-25 Site, and Rockwood	Water ingestion	Water ingestion
	Dermal contact during bathing	
		Fish ingestion
		Vegetable ingestion
		Meat ingestion
		Milk ingestion
		Soil ingestion
		Dust inhalation
		External Exposure

10. Exposure times used in the ORNL/ER-9 methodology are much greater than those used in MEPAS (i.e., shoreline use in MEPAS is only 12 h per year, whereas ORNL/ER-9 exposure time is 2000 h per year). This difference will also cause the risks calculated with the ORNL/ER-9 method to be greater than those of MEPAS. To obtain more appropriate estimates of risk with MEPAS for maximum exposures to individuals, these higher usage factors could be incorporated into MEPAS.
11. Many transfer coefficients used by MEPAS are lower, and sometimes higher, than those used by ORNL/ER-9. Tables 3 through 5 list selected values for comparison. Transfer coefficients and bioaccumulation factors (Table 6) were compared between the two methods but only for selected contaminants. Hence, the following comments apply only to a small subset of all possible contaminants and is merely provided to indicate there may be some improvements required in the selection of these values in MEPAS. All values used by MEPAS and listed in Tables 3 through 6 are compiled in Napier et al. (1980), which, unlike the National Council on Radiation Protection and Measurement (NCRP), did not develop the constants for screening purposes, calling into question their use in site-specific assessments. In addition, the Napier et al. (1980) values do not include soil uptake for human consumption and ingestion of soil by animals as do

ORNL/ER-9-calculated SI values, which are based on data bases developed by NCRP explicitly for screening. Although the two methods have different objectives, comparisons of these values indicate that some updates may be required in MEPAS to make selected values more appropriate for site-specific assessments.

Table 3. Transfer coefficients of contaminant to milk (d/L)

	MEPAS ^a	NCRP ^b
Cr	0.0011	0.002
Hg	0.019	0.0005
¹³⁷ Cs	0.005	0.01
⁶⁰ Co	0.0005	0.002
⁹⁰ Sr	0.0015	0.002
²³⁴ U	2.5×10^{-6}	0.0004
²³⁵ U	2.5×10^{-6}	0.0004

^aNapier et al. 1980.

^bNg 1982; Ng et al. 1977; Ng et al. 1982b.

Table 4. Soil-to-plant transfer factors

	MEPAS ^a	NCRP ^b
Cr	2.5×10^{-4}	0.01
Hg	0.38	0.03
¹³⁷ Cs	0.002	0.04
⁶⁰ Co	0.0094	0.08
⁹⁰ Sr	0.2	0.3
²³⁴ U	0.0025	0.002
²³⁵ U	0.0025	0.002

^aNapier et al. 1980.

^bBaes et al. 1984; IAEA 1982; Ng et al. 1982. Baes et al. (1984) contains provisions for inadvertent soil ingestion from unwashed vegetables.

Table 5. Transfer coefficients of contaminant to meat (d/kg)

	MEPAS ^a	NCRP ^b
Cr	0.00024	0.03
Hg	0.1	0.01
¹³⁷ Cs	0.03	0.05
⁶⁰ Co	0.001	0.03
⁹⁰ Sr	0.0004	0.01
²³⁴ U	0.00034	0.0008
²³⁵ U	0.00034	0.0008

^aNapier et al. 1980; Streng et al. 1986 (for Cr, ²³⁴U, and ²³⁵U).

^bNg 1982; Ng et al. 1977; Ng et al. 1982b.

Table 6. Bioaccumulation factors for freshwater fish

	MEPAS ^a (L/kg)	NCRP ^b
Cr	20	200
Hg	2×10^5	1×10^3
¹³⁷ Cs	2×10^3	2×10^3
⁶⁰ Co	50	300
³ H	1.0	1.0
⁹⁰ Sr	30	60
²³⁴ U	2.0	10
²³⁵ U	2.0	10

^aNapier et al. 1980.

^bValues compiled from Thompson et al. 1972; IAEA 1982; IAEA 1989; Vanderploeg et al. 1975; Hoffman and Baes 1979; Killough and McKay 1976; Newman 1985; Poston and Klopfer 1986; Copeland et al 1973.

Most of the MEPAS transfer coefficients of contaminants from daily diet to milk are 10^{-1} to 10^{-2} of those reported in NCRP, except for Hg. The milk transfer coefficient in MEPAS for Hg is $\sim 10^2$ times that used by NCRP. Regarding the soil to plant transfer factors, most of those used in MEPAS are similar to those used in NCRP; yet, the values for Cr, ¹³⁷Cs, and ⁶⁰Co are 10^{-2} to 10^{-3} times those used by NCRP.

Variable differences in the transfer coefficients of contaminants from daily diet to meat are observed between the two methods. Some MEPAS values are 10^{-1} to 10^{-2} times those in

NCRP (i.e., Cr, ^{60}Co , and ^{90}Sr), while others (i.e., Hg) are $\sim 10^1$ times those in NCRP, and others are approximately the same.

Additional differences between the two methods were found in the bioaccumulation factors from water to fish. For instance, MEPAS uses values for Cr and ^{60}Co that are $\sim 10^{-1}$ times those of NCRP, whereas the Hg factor in MEPAS is 10^2 times that in NCRP.

These are only a few of the differences in selection of transfer coefficients and bioaccumulation factors between the two methods. Only a small percentage of total contaminants that could be modeled by each method were selected by the authors for comparison. The differences mentioned simply indicate that it may be necessary to determine some of these values for the indicated, and other, contaminants in the MEPAS code for site-specific evaluations. When site-specific information is not available, the more conservative values should be selected, be they derived from Napier et al. (1980) or NCRP (1991), so that consistently conservative assessments can be made. This conservatism will at least partially offset the often low calculated concentrations associated with the specific waste sites.

5. RECOMMENDATIONS

Based on the findings of this study of the MEPAS methodology, we make several recommendations, many of which are relevant to any risk assessment. Incorporation of the following recommendations into future risk assessment studies will lead to a more conservative assessment of risk as well as results that are more easily and reliably interpreted.

1. **Abandon HPI:** MEPAS should be used without the HPI index, which artificially distorts the results of any risk assessment and makes their interpretation difficult.
2. **On-site vs off-site risks:** Considerations need to be made to distinguish between on- and off-site risks. MEPAS has not yet been used to evaluate on-site risks. However, MEPAS users should estimate on-site risks using measured, on-site concentrations. In considering on-site risks, the risk associated with the maximally exposed individual may be far greater than that attributed to an individual at an off-site usage location. In contrast, an on-site hazard area may be restricted from the population, in which case the risks associated with it are substantially diminished. A scheme that weights these opposing effects in terms of overall calculated risk should be developed by which on- and off-site risks can be, at least, qualitatively weighted.
3. **Uncertainty analysis:** To obtain more meaningful results, a hazard assessment should include population risks, both conservative and nonconservative estimates of individual risk, and incorporate uncertainty analysis. Because some sites have been studied to a far greater extent than others, those studied less may have a substantial amount of uncertainty associated with parameters needed as input (e.g., source concentration) to a risk assessment. This uncertainty should be recognized when sites are ranked for their potential hazards to human health. A nonconservative estimate in which risk values above a certain predetermined limit can be identified as a definite health concern could be incorporated into the methodology. This approach will assist in evaluating the hazards associated with sites because more meaningful interpretations can be made when an investigator is aware of the potential uncertainty associated with a particular value of hazard to the population, to the maximally exposed individual, and to the average individual. One, final, all-encompassing number corresponding to a risk cannot provide an investigator with any insight into the reliability of that number unless the ancillary information is also included for examination. Although a final PI summed over all pathways and contaminants may be needed in the final assessment of risk, risks, population sizes, and uncertainties must also be provided so that investigators may make informed decisions.
4. **Update default values:** Exposure data, including bioaccumulation factors, usage factors, and transfer factors, should be updated in the MEPAS code for use in calculating risks to the maximally exposed individual. Values listed in NCRP (1991) for screening purposes are compiled from a variety of sources, and these transfer factors and bioaccumulation factors are often more conservative than those used by MEPAS. The data in NCRP (1991) will, in general, contribute to a more conservative estimate of risk at the studied sites. Perhaps two separate data bases should be incorporated into the MEPAS code: one data set to calculate screening estimates and one to calculate realistic

exposures. Also, the MEPAS weighted cancer incidence risk conversion factor of 2.7×10^{-4} per rem should be updated to a value of 7.2×10^{-4} per rem.

5. Improved user input: Better user input into the model is needed. It should become the policy of any risk assessment study to consult local professionals most knowledgeable about a particular waste area. These professionals should provide the MEPAS users with a listing of relevant and realistic data to be used for the site-specific studies. Advice from experts in bioaccumulation, sources of the contaminants, and usage and transfer factors appropriate for the studied area should be obtained so that realistic input can be obtained and used. Discussions with the local experts should also occur so that the MEPAS user is familiar with the geologic and hydrologic background of the site as well as with the history of contaminant disposal (including potential contaminants present and their probable volumes and concentrations, if known). From the evaluation of the WAG 2 MEPAS risk assessment, it is evident that users of the model must be better informed about the particular sites under consideration.
6. Improve printout results: The results/printouts from MEPAS should include the risk for each contaminant considered. The output should be tabulated by usage location such that a risk can be associated with each exposure pathway in the particular usage location for each contaminant for three different risks: maximally exposed individual, average individual, and the population, with the numerical value of the population included in the printout. This information should be included so that those prioritizing these sites will be able to easily determine the origin of the numbers and not be forced to use the results without pertinent information. A summation of the RF for each contaminant should follow each listing, as well as a final summation over all pathways and contaminants for the three different risks so that a total risk can be assigned to the location. These three numbers can then be used to prioritize each RU. Without the listing of risk for each individual exposure pathway and contaminant type, it is impossible to assign a priority to a waste site and plan for cleanup because it is not known which contaminants may be the most important at a particular site.

6. CONCLUSIONS

MEPAS methodology and use were evaluated by scrutinizing the equations, default values, and input used to rank WAG 2 at Oak Ridge National Laboratory. Although it may be desirable to regulators to obtain one, final numerical value of risks associated with each waste site, this is not practicable, given the current state of the art in risk assessment. From the evaluation of MEPAS, it becomes clear that the use of the all-encompassing HPI values should be discontinued, and risks to the average individual, maximally exposed individual, and population should replace the HPI so that the final numbers can be qualitatively evaluated. For instance, a 10^{-3} risk to one individual who may be an intruder is less important than an immediate 10^{-4} or 10^{-5} risk to a larger portion of the general population, and this difference may not be reflected in the final HPI values. Although the use of one final number such as the HPI may be easily inserted into expressions from which prioritizations are based, the use of such a number in risk assessment is not recommended because those using the number cannot have a full appreciation of the meaning of the number and the complexity of the relevant issues associated with it. A scheme must be developed that can qualitatively evaluate and weight carcinogen versus noncarcinogen risks, hypothetical (intruder) versus real risks, and maximum individual and average individual versus population risks. Uncertainty analyses should accompany each of these values so that an investigator may determine if any real differences exist between various risk estimates and evaluate if the calculations made are consistently conservative.

MEPAS default values for transfer coefficients, bioaccumulation factors, and usage factors should also be updated in some cases so that the risk to a maximally exposed individual can be estimated. If site-specific off-site information is available, it should be used. If site-specific information is not available, values developed for screening purposes should be used to obtain a consistently conservative result.

Of equal importance to the improvement of MEPAS results is better input information. Users must consult local experts concerning the latest information about the site, and users must be consistent in their selection of exposure pathways to be considered. Sufficient time should be spent in developing appropriate input parameters for each site under investigation, including selection of certain bioaccumulation factors and transfer coefficients that may be more applicable to the site than the default values provided in MEPAS. Without realistic input parameters and assumptions, results from MEPAS or any risk assessment scheme will be unreliable. Using input that is not fully developed may save time, but the time savings will be meaningless if unreliable or completely incorrect results are obtained, which will be of little use in the overall risk assessment activities.

ACKNOWLEDGMENTS

The authors wish to thank L. W. Barnthouse and J. R. Trabalka for reviewing this document. The first author would like to thank J. Droppo of Pacific Northwest Laboratory for his assistance and clarifications of the MEPAS development and formulation.

REFERENCES

- Baes, C. F., III, R. D. Sharp, A. L. Sjoreen, and R. W. Shore. 1984. A review and analysis of parameters for assessing transport of environmentally released radionuclides through agriculture. ORNL-5786. Oak Ridge National Laboratory, Oak Ridge, Tenn.
- CAPCOA (California Air Pollution Control Officers Association). 1990. Air toxics "hot spots" program risk assessment guidelines, July 1990. Sacramento, Calif.
- Copeland, R. A., R. H. Beetfe, and W. W. Prater. 1973. Trace element distribution in Lake Michigan fish: A baseline study with calculation of concentration factors and equilibrium radioisotopes distribution. ERG Special Report No. 2. University of Michigan, Ann Arbor. 139 pp.
- Droppo, J. G., G. Whelan, J. W. Buck, D. L. Streng, B. L. Hoopes, and M. B. Walter. 1989. Supplemental mathematical formulations, Pacific Northwest Laboratory: The multimedia environmental pollutant assessment system (MEPAS). PNL-7201. Pacific Northwest Laboratory, Richland, Wash.
- Hoffman, F. F., and C. F. Baes III (eds.). 1979. A statistical analysis of selected parameters for predicting food chain transport and internal dose of radionuclides. NUREG/CR-1004, ORNL/NUREG/TM-282. Oak Ridge National Laboratory, Oak Ridge, Tenn.
- Hoffman, F. O., B. G. Blaylock, M. L. Frank, L. A. Hook, E. L. Etnier, and S. S. Talmage. 1990. Preliminary screening of contaminants in the off-site surface water environment downstream of the U.S. Department of Energy Oak Ridge Reservation. ORNL/ER-9. Oak Ridge National Laboratory, Oak Ridge, Tenn. 151 pp.
- IAEA (International Atomic Energy Agency). 1982. Generic models and parameters for assessing the environmental transfer of radionuclides from routine releases: Exposures of critical groups. Safety Series No. 57. International Atomic Energy Agency, Vienna.
- ICRP (International Commission on Radiological Protection and Measurement). 1991. The 1990 recommendations of the commission. The recommendations of the International Commission on Radiological Protection, National Radiological Protection Board. Publication 60. Annals of the ICRP, Pergamon Press, Oxford, England.
- Killough, G. G., and L. K. McKay. 1976. A methodology for calculating radiation doses from radioactivity released in the environment. ORNL-4992. Oak Ridge National Laboratory, Oak Ridge, Tenn.
- Napier, B. A., R. L. Roswell, W. E. Kennedy, Jr., and D. L. Streng. 1980. Assessment of effectiveness of geologic isolation systems: ARRRG and FOOD: Computer programs for calculating radiation dose to man from radionuclides in the environment. PNL-3180. Pacific Northwest Laboratory, Richland, Wash.

NCRP (National Council on Radiation Protection and Measurement). 1989. Screening techniques for determining compliance with environmental standards. NCRP Commentary No. 3. Bethesda, Md.

NCRP Scientific Committee 64-6. 1991. Screening models for releases of radionuclides to air, surface water, and ground water: Description of models. NCRP report No. XX. vol. 1 (Draft report).

Newman, G. 1985. Concentration factors for stable metals and radionuclides in fish, mussels, and crustaceans—A literary survey. SNUPM 1976E. National Swedish Environmental Protection Board, Solna, Sweden. 36 pp.

Ng, Y. C.. 1982. A review of transfer factors for assessing the dose from radionuclides in agricultural products. Nucl. Saf. 23: 57.

Ng, Y. C., C. S. Colsher, D. J. Quinn, and S. E. Thompson. 1977. Transfer coefficients for the prediction of the dose to man via the forage-cow-milk pathway from radionuclides released to the biosphere. UCRL-51939. Lawrence Livermore National Laboratory, Livermore, Calif.

Ng, Y. C., C. S. Colsher, and S. E. Thompson. 1982a. Soil-to-plant concentration factors for radiological assessments. UCID-19463. Lawrence Livermore National Laboratory, Livermore, Calif.

Ng, Y. C., C. S. Colsher, and S. E. Thompson. 1982b. Transfer coefficients for assessing the doses from radionuclides in meat and eggs. UCID-19464. Lawrence Livermore National Laboratory, Livermore, Calif.

Poston, T. M., and D. C. Klopfer. 1986. A literature review of the concentration ratios of selected radionuclides in freshwater and marine fish. PNL-5484. Pacific Northwest Laboratory, Richland, Wash.

Streng, D. L., R. A. Peloquin, and G. Whelan. 1986. LADTAPII-A technical reference and user guide. PNL-5270. Pacific Northwest Laboratory, Richland, Wash.

Thompson, S. E., C. A. Burton, D. J. Quinn, and Y. C. Ng. 1972. Concentration factors of chemical elements in edible aquatic organisms. UCRL-50564, Rev. 1. Lawrence Livermore National Laboratory, Livermore, Calif.

USDOE (U.S. Department of Energy). 1991. Environmental survey preliminary report, Oak Ridge National Laboratory (X-10), Oak Ridge, Tennessee. DOE/EH/OEV-31P. U.S. Department of Energy, Washington, D.C.

USNRC (U.S. Nuclear Regulatory Commission). 1977. Calculation of annual doses to man from routine releases of reactor effluents for the purpose of evaluating compliance with 10 CFR Part 50, Appendix I. Regulatory Guide 1.109. U.S. Nuclear Regulatory Commission, Washington, D.C.

Whelan, G., D. L. Streng, J. G. Droppo, Jr., B. L. Steelman, and J. W. Buck. 1987. The remedial action priority system (RAPS): Mathematical formulations. DOE/RL/97-09. U.S. Department of Energy, Washington, D.C.

DISTRIBUTION

1. L. V. Asplund/N. W. Durfee 2. L. Barnthouse 3. L. D. Bates 4. R. B. Clapp 5. K. W. Cook 6. J. H. Cushman 7. M. F. P. DeLozier 8. C. Easterly 9. M. P. Farrell 10. D. E. Fowler 11. S. B. Garland 12. C. W. Gehrs 13. C. D. Goins 14. J. A. Greene 15. P. J. Halsey 16. S. G. Hildebrand 17-19. F. O. Hoffman 20. L. D. Hyde 21. D. C. Kocher 22. A. J. Kuhaida 23. V. Legg 24. K. L. Michel	25-26. P. T. Owen 27. D. E. Reichle 28. G. E. Rymer 29. P. A. Schrandt 30-32. L. A. Shevenell 33. D. S. Shriner 34. D. K. Solomon 35. S. H. Stow 36. G. W. Suter 37. D. W. Swindle 38. J. R. Trabalka 39. C. C. Travis 40. R. I. Van Hook 41. D. R. Watkins 42. R. K. White 43-47. ER Document Management Center 48. Central Research Library 49-53. ESD Library 54. ORNL Y-12 Technical Library 55. Laboratory Records Dept. 56. Laboratory Records, ORNL-RC 57. ORNL Patent Section
-
58. Office of Assistant Manager for Energy Research and Development, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8600
 59. J. G. Droppo, Battelle Northwest, P.O. Box 999, Richland, WA 99352
 60. P. H. Edmonds, Radian Corporation, 120 South Jefferson Circle, Oak Ridge, TN 37830
 61. J. F. Franklin, Bloedel Professor of Ecosystem Analysis, College of Forest Resources, University of Washington, Anderson Hall (AR-10), Seattle, WA 98195
 62. G. M. Hornberger, Professor, Department of Environmental Sciences, University of Virginia, Charlottesville, VA 22903
 63. G. Y. Jordy, Director, Office of Program Analysis, Office of Energy Research, ER-30, G-226, U.S. Department of Energy, Washington, DC 20545
 - 64-65. J. R. Kannard, Program Manager, Bechtel National, Inc., P.O. Box 350, Oak Ridge Corporate Center, 151 Lafayette Drive, Oak Ridge, TN 37830
 - 66-69. W. E. Murphie, Department of Energy, Office of Environmental Restoration, Eastern Area D&D Branch, EM-423 (GTN), Washington, DC 20545
 70. R. H. Olsen, Vice President for Research, University of Michigan, 6643 Medical Science Building II, Ann Arbor, MI 48109-0620
 71. A. Patrinos, Acting Director, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
 - 72-73. S. S. Perkins, DOE Oak Ridge Field Office, Information Resource Center, 105 Broadway, Oak Ridge, TN 37830
 - 74-75. S. P. Riddle, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8541
 - 76-77. R. C. Sleeman, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8541

78. F. J. Wobber, Environmental Sciences Division, Office of Health and Environmental Research, Office of Energy Research, ER-74, U.S. Department of Energy, Washington, DC 20584
- 79-80. Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831

END

DATE
FILMED

6 / 23 / 92