DOE/ER/60422--T2

DOE/ER/60422--T2

DE93 009947

Surface Energy Balances of Three General Circulation

Models: Current Climate and Response to

Increasing Atmospheric CO₂

W. J. Gutowski, D. S. Gutzler, D. Portman and W.-C. Wang

Atmospheric and Environmental Research, Inc.

Cambridge, Massachusetts 02139

April 1988

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

	<u>Page</u>
Executive Summary.....	1
1. Introduction.....	7
2. Models and Data.....	9
3. Horizontal Averages.....	11
4. Surface Energy Balance Equations.....	13
5. Global Mean.....	15
6. Zonal Mean.....	18
7. Regions.....	23
8. Summary.....	33
Acknowledgements	35
References.....	37
List of Tables.....	39
List of Figures.....	77

Executive Summary

We examine the surface energy balance simulated by state-of-the-art general circulation models at GFDL, GISS and NCAR for climates with current levels of atmospheric CO₂ concentration (control climate) and with twice the current levels. The work is part of an effort sponsored by the U.S. Department of Energy to assess climate simulations produced by these models. The surface energy balance enables us to diagnose differences between models in surface temperature climatology and sensitivity to doubling CO₂ in terms of the processes that control surface temperature. Our analysis compares the simulated balances by averaging the fields of interest over a hierarchy of spatial domains ranging from the entire globe down to regions a few hundred kilometers across.

Principal Findings

1. For the global average control climate, individual surface fluxes of sensible heat (SH), latent heat (LH), shortwave radiation (SW) and upward (LW↑) and downward (LW↓) longwave radiation agree between the models to within 25 Wm⁻² (Figure E-1). In all three models the surface energy balance is dominated by longwave radiation (the "greenhouse" effect). Other surface fluxes are smaller, but the models also agree on their relative magnitudes. The 25 Wm⁻² difference is only a small fraction of the magnitudes of the largest fluxes, LW↓ and LW↑. However, the difference is a large fraction of the net longwave radiation at the surface (~60 Wm⁻²), and it is as large as any of the global-average seasonal changes in these fluxes. From this perspective, differences between surface fluxes in these state-of-the-art models are substantial even for the global average.

2. Global average changes in surface fluxes when CO₂ doubles agree between models to within 3 Wm⁻² (Figure E-2); intermodel discrepancies are only small fractions of the flux changes from CO₂ doubling. However, the apparent agreement between models in Figure E-2 should be viewed with caution because flux changes associated with doubled CO₂ are about the same size as the differences between models in their control climates: 25 Wm⁻². Furthermore, averaging over the globe obscures much larger differences occurring on smaller scales

(cf. points 4 and 6 below).

3. Spatial variation of net longwave radiation at the surface is small compared to the spatial variability of near-surface air temperature and moisture, both of which exert significant control on the net surface longwave flux. The relatively small variability of LW_{net} appears to be a consequence of compensation among variations of temperature, atmospheric moisture and clouds.

4. Intermodel differences in doubled- CO_2 surface flux changes of up to 100 Wm^{-2} occur near the limits of polar ice caps. Differences in ice modeling approach contribute strongly to the flux differences. Differences in control climate sea ice limits also help produce the strong differences in flux changes.

5. Differences between models for surface fluxes in regions a few hundred km across are often closely linked to differences in zonal mean climatology.

6. Intermodel surface flux discrepancies in the regional averages are as large as 50 Wm^{-2} , which is twice the difference found for the global average. Furthermore, models often disagree over the sign of regional flux changes when CO_2 doubles (Figure E-3). Such differences appear to be closely associated with differences in model hydrology.

7. Fields that are subject to orographic control show relatively small intermodel discrepancy, i.e., differences between models for the same region tend to be smaller than interregional differences. A most notable example is precipitation, examined here because of its indirect relation to the surface energy balance through the hydrological cycle.

Recommendations

1. Effort should be given to developing a better understanding of the hydrological cycle and how to simulate it. Progress in this area appears to be especially important for improving regional climate modeling (point 6 above). Relatively large flux discrepancies on the global scale (points 1

and 2) may also be reduced by such progress because the hydrological cycle, through water vapor and cloud cover, strongly influences the longwave radiation transmitted from the atmosphere to the surface. Examples of the work needed are Manabe and Wetherald (1987) and Meehl and Washington (1987) who have analyzed soil moisture sensitivity in NCAR and GFDL model simulations of current and doubled CO₂ simulations.

2. Better agreement in zonal mean climatology between models is needed, since much of the intermodel difference in regional climatology is attributable to zonal mean differences between models.

3. Effects of increased resolution on regional simulation should be studied. Our results suggest that improvements in orographically dependent fields are possible simply by increasing resolution. Grid nesting needs to be explored as a means of improving regional modeling over land areas where orographic control may be important for regional climatology.

SURFACE FLUX
Annual, global mean
(W/m-m)

Figure E-1. Annual, global mean surface fluxes from the GFDL, NCAR and GISS general circulation models using current atmospheric CO₂ levels. Key: SH - sensible heat flux, LH - latent heat flux, SW - net shortwave radiative flux, LW↓ - downward longwave radiation, LW↑ - upward longwave radiation, S - other contributors to the surface energy balance.

Figure E-2. Same as Fig. E-1 but for the change in the annual, global mean surface fluxes when atmospheric CO₂ doubles.

Figure E-3. Same as Fig E-1 but for the changes by CO₂ doubling for the season June-July-August averaged over selected regions: central United States (USC), western United States (USW), northeast China (CNE) and southwest China (CSW). (See Section 3 of this report for the definitions of the regions.)

1. INTRODUCTION

This report presents an analysis of the surface energy balance in simulations of current and doubled- CO_2 climates made by three general circulation models (GCMs). The surface energy balance determines a quantity of primary interest for climate change, the surface temperature. This perspective enables us to diagnose differences between models in surface temperature and surface temperature sensitivity to increasing CO_2 in terms of the heat fluxes that control surface temperature. Furthermore, although the ultimate source of atmospheric energy is solar heating, most of the energy absorbed by the atmosphere enters as fluxes from the surface (Wallace and Hobbs, 1977), so that the surface energy balance describes an important gateway for atmospheric energy. The analysis thus complements studies that consider the earth-atmosphere system as a whole by analyzing the radiation budget at the top of the atmosphere (e.g., Cess and Potter, 1987). Each of the surface fluxes is influenced by atmospheric processes. The analysis thus provides a basis for understanding how other factors such as clouds and moisture are related to surface temperature changes through their effect on surface fluxes. The analysis also provides a simplified starting point for future analyses of upper air energy budgets.

The climate simulations evaluated here were made by models developed at the National Center for Atmospheric Research (NCAR), the Goddard Institute for Space Studies (GISS) and the Geophysical Fluid Dynamics Laboratory (GFDL). The intercomparison has two parts: (1) an assessment of the model simulations of the current climate, which we refer to as "control" runs and (2) an assessment of the sensitivity of each model's control climate to doubling CO_2 . Like models with fewer spatial dimensions, GCMs are often used to evaluate changes in global mean climate. However, their spatial coverage also allows one to study changes for areas of smaller extent. Here we present climatic data from the GCMs for areas ranging from the entire globe to regions a few hundred kilometers across.

Understanding regional changes is desirable because societal impacts of climate change caused by increasing CO_2 will likely vary on regional scales. However, the reliability of GCM projections of climate change is uncertain on regional scales. Comparison of regional climate changes provides a stringent test of the models' projections, because differences between models may be

smoothed over when averages covering larger domains are compared.

The purpose of the intercomparison is to identify degrees of agreement or disagreement between state-of-the-art climate models in their representation of the surface energy balance. Agreement among the models does not in itself mean that their projections are accurate, but lack of agreement among sophisticated models such as these undermines confidence in their reliability. Comparison is also made with observations when available. The intercomparison is not intended to identify a best model. Although all three models have a common goal, simulating the annual cycle of the earth's climate, the model descriptions in Section 2 show many differences in the approaches taken to achieve this goal. Thus, detailed intercomparison could be misleading, and some differences between the models' surface energy balances should be expected. Also, a model agreeing closely with observations does not necessarily produce the most reliable estimates of climate sensitivity. By evaluating disagreements between the models' surface energy balances we hope to identify the physical processes responsible for discrepancies in model predictions, thereby suggesting promising areas for improvement, particularly on regional scales.

In Section 2, we describe characteristics of the models in brief, emphasizing the aspects of the models directly relevant to this study. Section 3 describes our hierarchy of spatial averages. In Section 4 we discuss the surface energy balance equation and develop a relationship between surface flux changes and surface air temperature changes. We present highlights of results for the global, zonal and regional balances in Sections 5, 6 and 7. Section 8 contains a summary of our findings.

Associated with Sections 5 - 7 is an extensive set of tables and figures that contain data on surface energy fluxes and quantities that influence them such as cloud cover and near-surface atmospheric moisture. Our discussion of them extracts the information we feel is most important for comparing the models, but we do not discuss features of each of these tables and figures in detail, especially those pertaining to individual regions. Rather, the full set has been included in order to provide the reader with an atlas that serves as a reference for the surface energy balance of these models.

2. MODELS AND DATA

The models used here are GCMs developed at NCAR (Washington and Meehl, 1984), GFDL (Manabe and Wetherald, 1987) and GISS (Hansen et al., 1983). Schlesinger and Mitchell (1987) give a comparative summary of their more important features. The GFDL and NCAR models are spectral models of the type first described by Bourke (1974). Both use wavenumbers 0-15 in rhomboidal truncation; the corresponding grid resolution is 4.5° (latitude) x 7.5° (longitude). The GISS model is a grid-point model with 8° (latitude) x 10° (longitude) resolution. All three models use 9 sigma layers for vertical finite differencing. The sigma layers in each are unevenly spaced in order to give highest resolution near the surface and tropopause.

All three models include realistic geography to within the accuracy of their horizontal resolution. In the GFDL and NCAR models, the surface underlying each grid box is either all land or all ocean. The GISS model allows the surface in each box to be partly ocean and partly land, with the portion of each varying according to the actual geographical distribution. The ocean in each model is a simple mixed layer which allows the model to simulate the annual cycle of surface temperature. The GISS model also includes a specified ocean transport that is deduced by requiring a match of the model's simulated control sea surface temperatures with observations. All include a hydrological cycle and interactive cloud cover, snow cover and sea ice. In the NCAR and GFDL models, if snow or ice is present in a grid box, then the entire box is covered with an even depth of snow or ice. The thickness of snow or ice varies according to heat and water budget equations. The GISS model allows a grid box to have fractional snow and ice cover which can vary according to budget equations. Only the GISS model simulates the diurnal cycle.

Surface dynamic fluxes in all three models are computed using bulk aerodynamic formulas:

$$SH = C_p c_p V_a (T_s - T_a) \quad (1a)$$

for sensible heat flux and

$$LH = \mu C_p LV_a \left\{ q_{sat}(T_s) - q_a \right\} \quad (1b)$$

for latent heat flux, where ρ is density, c_p the heat capacity of air, C a transfer coefficient, V wind speed, T temperature, L the latent heat of vaporization, q specific humidity, and q_{sat} the saturation specific humidity. The term μ in (1b) is a moisture availability factor. Over the ocean, $\mu = 1$. Over land, μ is a linear function of soil moisture, varying between 0 for dry ground and 1 for soil moisture greater than a critical value. In the NCAR and GFDL models, the critical value is 75% of field capacity; in the GISS model, it is the full field capacity of the model's upper soil layer. The subscript s in (1) refers to surface (ground or ocean) values. For GFDL and NCAR, the subscript a refers to the value in the lowest model layer, centered at 0.99σ in both cases. For GISS, the subscript a refers to atmospheric values just above the surface that are obtained from a boundary layer model. Table 1 gives a description of coefficient C for each model.

All models include radiation absorption by water vapor, carbon dioxide and ozone, though the bands included vary with model. All three models include scattering by molecules; the GISS model also includes scattering by aerosols. Surface emissivity is unity everywhere in the GFDL model, but has some geographic and spectral dependence in the GISS and NCAR models. In all models, the albedos of snow-free land and open ocean vary geographically but generally do not change when the climate changes. The only exception is the dependence of ocean albedo on surface wind in the GISS model. Each model's surface albedo changes as snow and sea ice cover change. Snow and sea ice albedos are constants in the NCAR model, whereas these albedos depend on factors such as depth and underlying vegetation in the GFDL and GISS models.

Clouds occur in the models as a result of either large-scale condensation or cumulus convection. The GISS model does not allow clouds at pressures less than 100mb. The NCAR model allows clouds to form in all layers except the lowest ($\sigma = 0.991$) and the two highest ($\sigma = 0.074$ and 0.009). The GFDL model allows clouds to form in all layers. Clouds in all three models are produced by diagnostic relations and are not advected, so that clouds can appear and disappear from one time step to the next. Clouds from large-scale condensation occur in each model when a relative humidity criterion is met. For both the GFDL and NCAR models, convective clouds appear when a relative humidity requirement is satisfied for grid box layers undergoing convective

adjustment. The GISS convective cloud parameterization includes a convective mass flux dependence. Cloud optical depths, albedos and emissivities are functions of cloud location and the number of layers a cloud encompasses.

The data available from all models include surface fluxes of sensible and latent heat, net shortwave and net longwave radiation and air temperature near the surface. Available data also include elements of the hydrological cycle that may directly influence the surface energy balance: precipitation, clouds in two layers and near-surface specific humidity. GISS model surface flux and temperature computations are performed separately for the land and ocean fractions in each box. The archived data for each GISS grid box is an area-weighted average of the separately computed quantities.

All data have been supplied as temporal averages obtained from the final years of an experiment's run. For NCAR, the collection period for data is the final 3 years; for GFDL and GISS, the final 10 years. The analysis here uses seasonal means, which were supplied by GISS and NCAR. For GFDL, monthly mean data were averaged to produce analogous seasonal means.

3. HORIZONTAL AVERAGES

We compare the simulations by averaging the model output fields of interest over a hierarchy of spatial domains. First, we examine averages taken over the entire surface of the earth. The global mean provides a context for understanding averages on smaller scales, since part of our analysis considers how smaller scale averages deviate, if at all, from the global mean. Also, changes in global mean quantities, especially surface temperature, are considered key indicators of climate sensitivity. Next, we extend our analysis to zonal averages. Consideration of the zonal mean energy balance is important because the greatest spatial variation of surface temperature is its decrease from equator to pole. Finally, we compare the model output averaged over selected geographical regions to examine explicitly the longitudinal variability of the surface heat balance. Realistic simulation of climate and climate change on the scale of these regions can only be accomplished with three-dimensional models such as the GCMs we examine here.

There are any number of ways that one could choose to examine and compare the longitudinal variability of climate and climate change in these models. Direct comparison of gridpoint values is difficult because the model grids are different; furthermore, the proper physical interpretation of a single model gridbox is unclear. Spectral decomposition of the model output may be quite useful for examination of large-scale dynamics, but the surface energy budget and hydrological cycle are closely tied to the geographical distribution of continents, mountains, and oceans, which do not project cleanly onto spectral harmonics. For this investigation we have chosen to average the model gridpoint output over ten regions (Fig. 1). The regions cover the United States, China, and parts of the Atlantic and Pacific Oceans. The U.S. and China are singled out for examination in order to mesh this study with ongoing research on observed climate changes in those countries. The oceanic regions are included here in order to compare the model surface heat balances over non-continental areas. The regional boundaries were selected a priori from consideration of geography and the climatology of temperature and precipitation. The three U.S. regions correspond to areas where most precipitation occurs in winter (USW), summer (USC), and an area without a pronounced annual cycle in precipitation (USE) (Hsu and Wallace, 1976). China is divided into four regions: CSE, where summer monsoonal rains dominate the annual cycle but precipitation occurs year-round; CSW, a mountainous region far from coastal influences on precipitation; CNW, a desert; and CNE, a grassland region with a pronounced winter dry season.

The GFDL and NCAR regions are identical except for Southeast China, where GFDL includes an extra land box not present in the NCAR model. GISS boxes for China and US regions may include lakes or oceans. For the GISS averages, we weight each box's contribution to the average by the fraction that is land. This lessens but does not remove the contribution of the ocean fraction to the GISS "land" region averages. Similarly, GISS boxes falling into two different regions are included in each regional average with half weight.

4. SURFACE ENERGY BALANCE EQUATIONS

The thermodynamic equation, vertically integrated over all subsurface layers, is

$$\epsilon \sigma T_s^4 + SH + LH + SW + LW\downarrow + S = 0 , \quad (2)$$

where T_s is surface temperature, σ is the Stefan-Boltzmann constant, ϵ is emissivity, SH and LH are surface sensible and latent heat fluxes, SW is the net solar flux at the surface, LW $\downarrow$ is the downward longwave flux from the atmosphere into the surface and S is a residual that includes heat storage. All fluxes are defined as positive upward. Emissivity is unity for the GFDL model but for the NCAR and GISS models it is a function of underlying surface type and wavelength that is ≈ 1 when averaged over the spectrum. We use $\epsilon = 1$ in all calculations.

SH, LH and SW are given explicitly by the model postprocessing routines. The term LW $\downarrow$ is inferred from LW $_{net}$, the net longwave flux at the surface, and T_a , the surface air temperature, by assuming $T_a = T_s$ and using

$$LW_{net} = \sigma T_s^4 + LW\downarrow . \quad (3)$$

Any difference between T_a and T_s is subsumed into LW $\downarrow$. Data available from the GFDL model shows that differences between T_a and T_s can be as large as 4K, implying a bias error in our calculated LW $\downarrow$ of about 20 Wm^{-2} . However, this error applies only to the partitioning of LW $_{net}$ into upward and downward components. Magnitudes of LW $_{net}$ can be compared precisely.

The bias error in estimating LW $\downarrow$ due to differences between T_s and T_a is $\sim 4\sigma T_s^3 (T_s - T_a)$. Sensible heat flux in the models is also proportional to $T_s - T_a$, so comparison of SH between models can indicate the magnitude of the bias. Using (1a) and the typical values $T_s = 290 \text{ K}$, $V_s = 5 \text{ m/s}$, $C_D = 2 \cdot 10^{-3}$ and $\rho = 1 \text{ kg/m}^3$ gives

$$LW\downarrow \text{ bias} \sim SH/2 . \quad (4)$$

Using (4), SH data in Table 2 suggest that bias errors in global mean LW $\downarrow$ are $< 15 \text{ Wm}^{-2}$. Furthermore, Table 2 and (4) suggest that the bias errors are roughly the same size in each model. Thus relative magnitudes of LW $\downarrow$ can be compared with some confidence.

For the GFDL and NCAR models, S includes heating by snowmelt. For GISS, S also includes subsurface heat transport and the flux of heat by precipitation. Snowmelt and precipitation heat flux are generally a few watts per meter squared or less, i.e., relatively small. Subsurface heat transport in the GISS model ocean can be comparable to other fluxes in (2). Miller et al. (1983) and Russell et al. (1985) discuss the derivation of the GISS model's ocean flux in detail and compare it to estimates of the actual ocean flux.

For small climate changes, we can linearize a Taylor expansion of σT_s^4 , yielding

$$\Delta T_a \approx -(\Delta SH + \Delta LH + \Delta SW + \Delta LW\downarrow + \Delta S) / 4\sigma T_a^3, \quad (5)$$

where we have substituted ΔT_a for ΔT_s and T_a for T_s . Differences between T_s and T_a and their changes when CO_2 is doubled enter calculations using (5) primarily in computing $\Delta LW\downarrow$ using (3). The error in this contribution is about the same size as $\Delta T_s - \Delta T_a$. For the GFDL model, $\Delta T_s - \Delta T_a$ is generally much less than 1K except where the edges of the polar ice caps shift. There the two temperature changes differ by about 4K, so that the contribution of $\Delta LW\downarrow$ to ΔT_a will be computed with less accuracy in these regions. However, $\Delta LW\downarrow$'s contribution also tends to be much greater than 4K in the same regions, so that its effect on ΔT_a is not obscured.

We apply equations (2), (3) and (5) to spatially averaged domains. In doing so, we assume that $\{T_s^n\} \approx \{T_s\}^n$, where the brackets refer to a domain average. These assumptions are valid if the variability of T_s about $\{T_s\}$ within the domain is relatively small, which is always the case.

Surface flux changes in succeeding sections are described in terms of associated surface air temperature changes that we calculate using (5). For the range of temperatures encountered in the analysis, flux changes translate to temperature changes in the ratio of approximately 5 Wm^{-2} per degree.

The following sections give results obtained using (2), (3) and (5). We begin in Section 5 with the largest spatial domain considered, the global mean, and then proceed to zonal average and regional domains in Sections 6 and 7.

5. GLOBAL MEAN

Tables 2-4 give global mean values of variables relevant to the surface energy balance for three time averages: the annual mean, the December-January-February (DJF) mean and the June-July-August (JJA) mean. Table 2 contains global mean surface air temperature and each of the terms of the energy balance (2) for the control and doubled- CO_2 simulations from each model. The first columns of Table 2 contain observed climatological fluxes (Budyko, 1974; National Academy of Sciences, 1975) for comparison with the control climates. The observed fluxes are obtained in part through use of empirical formulas. The data sources give LW_{net} , which we separate into upward and downward components using $T_s = 287 \text{ K}$. Table 3 contains elements of the climate that have a direct bearing on the surface energy balance and hydrological cycle: precipitation, specific humidity near the surface, clouds in two layers and lapse rate. For the NCAR and GFDL models low cloud is the cloud cover in the second layer above the surface and high cloud is the cloud cover in the sixth layer. For the GISS model, low cloud is the net cloud cover for the lowest three layers and high cloud is the net cloud cover for the sixth and seventh layers together. Table 4 contains the terms of (5), with each flux change expressed in terms of an effective temperature change.

Differences between the models' global mean T_a are less than 1.5 K in each control climate averaging period. Using (5), this implies a net surface flux difference between the models of about 7.5 Wm^{-2} . As discussed in more detail below, differences between models for a particular flux tend to be greater than this amount, implying some compensation among the different components of the surface energy balance. At least some of the compensation appears to be a result of the models' attempting to achieve a balance with the annual cycle of solar forcing (Meehl and Washington, 1985).

The upward and downward LW fluxes are clearly the dominant terms in the global energy balance throughout the year. This is simply the greenhouse effect: radiative "blanketing" by the atmosphere maintains the surface at a warmer temperature than it would be without an atmosphere. Although the models display a fair qualitative agreement in their longwave fluxes, the range in LW_{net} (the sum of $\text{LW}\downarrow$ and σT_a^4) among them is roughly $20\text{-}30 \text{ Wm}^{-2}$, which is much greater than the 7.5 Wm^{-2} scale derived from temperature differences. This range is also much greater than the change in LW_{net} that

occurs in any model when CO_2 doubles, i.e. the climate change signal is smaller than the discrepancy between models even in the global mean.

Most of the difference between models in LW_{net} is attributable to differences in $\text{LW}\downarrow$, especially for the current climate simulation. The estimated upward longwave fluxes differ by less than 8 Wm^{-2} . Some of the difference in $\text{LW}\downarrow$ appears to be associated with differences in atmospheric moisture: the NCAR model has the least amount of atmospheric moisture of the three models (cf. Table 3), and it has the least $\text{LW}\downarrow$.

Discrepancies between models for the other fluxes are also as large or larger than corresponding flux changes in doubled CO_2 simulations. The GFDL model has about $10\text{-}15 \text{ Wm}^{-2}$ less solar flux reaching the ground than the other two models (despite having a solar constant 7% larger than the other two models), and its latent heat flux is also about $10\text{-}15 \text{ Wm}^{-2}$ less than given by the other models. In all models, sensible heat flux is relatively small compared to the other fluxes, but it also shows a spread of about 10 Wm^{-2} among models.

The observationally based fluxes span about the same range of values the model fluxes. No model appears to be simulating the earth's global mean surface energy balance better than the others. The latent heat flux has the biggest discrepancy between the models and the observations; each model's flux is greater than the observational fluxes by up to 16 W m^{-2} .

The residual term S is generally small but nonzero even in the annual mean. Quantities affecting surface energy balance such as precipitation heat flux in the GISS model and snowmelt in the other two models are included in S and can account for some of its magnitude, but it is still possible that the sum of all processes affecting the surface energy budget may not be in balance. This could occur because of the finite averaging times used to obtain model statistics or because the models may not be in precise equilibrium. For example, Washington and Meehl (1984) note that the NCAR model was probably near but still short of equilibrium at the time they terminated their experiment. Further integration of the NCAR model has shown that a slight downward trend in surface temperature was still occurring at the end of both the current and doubled CO_2 experiments analyzed here (Schlesinger and Mitchell, 1987). However, Washington and Meehl feel that the model is close enough to equilibrium in the two simulations that further integration

would not change the general features of the simulated climates. Changes of S with season are consistent with plausible changes in storage. With a large fraction of ocean-covered surface, the Southern Hemisphere dominates the global mean storage. Oceanic heat storage in the Southern Hemisphere is positive in DJF (summer) and negative in JJA (winter) (cf., Meehl and Washington, 1984). This seasonal behavior dominates the annual cycle of global mean S . Seasonal changes in global mean S tend to be larger than the seasonal change of all other global mean energy balance terms except the LW fluxes. The NCAR model's S , however, has a much smaller seasonal variation in the control climate, and the model's ΔS when CO_2 doubles is much larger than the ΔS in the other two models (cf. Table 4).

As a first approximation, the change in surface air temperature associated with a climate perturbation is $\Delta T_a = -\Delta \text{LW}\downarrow / 4\sigma T_a^3$, i.e., changes in upward and downward thermal radiation fields are highly coordinated (Table 4). However, the instantaneous increase in $\text{LW}\downarrow$ due to increased IR absorption by the additional CO_2 molecules is only $\sim 1 \text{ Wm}^{-2}$ (Wang and Molnar, 1985), implying $\Delta T_a \sim +0.2 \text{ K}$. Comparison of the instantaneous changes with the actual $\Delta \text{LW}\downarrow$ and associated temperature change shows the existence of positive feedbacks in the climate system. The positive feedback results in part because surface warming causes atmospheric temperatures to increase, resulting in further warming. Other positive feedbacks are also involved, most notably that associated with the increase in atmospheric water vapor (e.g., Manabe and Wetherald, 1967; Hansen et al., 1984). The order of water vapor increase among the models for each season matches the order of surface air temperature increase.

After $\Delta \text{LW}\downarrow$, the largest term in (5) for each model is that associated with ΔLH . Global mean LH transports heat into the atmosphere and cools the surface. When CO_2 doubles in these models, global mean LH increases, thereby cooling the surface further. Latent heat flux changes counteract the surface warming caused by the increase in $\text{LW}\downarrow$. The role of moisture in the surface energy balance is thus different than for the balance at the top of the atmosphere. Viewed from the top, an increase in atmospheric moisture warms the surface-atmosphere system (e.g., Manabe and Wetherald, 1967; Hansen et al., 1984), because water vapor is a strong infrared absorber. In the surface energy balance, moisture plays a dual role. Increasing the latent heat flux

can raise the atmosphere's temperature and humidity, causing LW↓ and, hence, surface temperature to increase. However, increasing LH also cools the surface directly.

Changes of other terms in (5) are relatively unimportant. There is a tendency for the intensity of SH to decrease and the intensity of SW to increase, both of which cause warming. Changes in S are even smaller. In both the JJA and DJF seasons, flux changes are almost always the largest in the GISS model, but the flux changes otherwise are similar among all three models.

From Table 3 it can be seen that globally-averaged cloud cover changes are insignificant in all three models, despite precipitation rate increases of 5-10% and increases in LH in all models and all seasons when CO₂ doubles. Hansen et al. (1984) also find that the cloud cover change contribution is of lesser importance compared to other changes occurring as CO₂ doubles.

6. ZONAL MEAN

Figures 2-24 show zonal means of model output corresponding to the global mean output in Tables 2-4. Figures 2, 11 and 18 show T_a and its change when CO₂ doubles for the annual, DJF and JJA means, respectively. Figures 3, 12 and 19 show the terms of the surface energy balance (2) for each model's control climate. Figures 4, 13 and 20 show terms of (5), the surface temperature change associated with each flux change. Figures 5-8, 14-17 and 21-24 show zonal mean plots of the quantities depicted in Table 3. Figures for cloud cover show only the changes when CO₂ doubles; the cloud cover amounts from each model refer to somewhat different vertical layers, so comparison of absolute cloud amounts may be misleading. Figure 9 shows terms of (5) for each model's doubled CO₂ annual mean climate, and Figure 10 shows a comparison of LW↓, LW↑ and LW_{net} from each model for the annual mean.

Surface air temperatures agree fairly well with each other. The biggest differences between the models occur in the winter polar regions and in the vicinity of the Southern Hemisphere ice margin. Differences between the NCAR and GFDL simulations correspond to differences in sea ice limits.

Comparison of Figures 3 and 10 shows that the upward and downward LW fluxes are the dominant terms in the zonal mean surface energy balance, as they were for the global mean. A notable feature of Figure 10 is that LW_{net}

has relatively little variation with latitude despite the variation of the upward and downward fluxes. This feature of zonal mean LW_{net} appears persistently among all models and for all seasons examined (cf. Figures 3, 9, 12, and 19).

We can obtain insight into this behavior by considering an empirical formula for clear sky LW_{net} developed by Brunt (1932) and Berliand and Berliand (1952):

$$LW_{net} = \epsilon \sigma T_a^4 (0.39 - 0.05\sqrt{e}) + 4\epsilon \sigma T_a^3 (T_s - T_a), \quad (6)$$

where e is the vapor pressure (in mb) at anemometer level and ϵ is the emissivity of water vapor ($= 0.97$). Eq. (6) states that LW_{net} increases with T_a and decreases with e . If T_a and T_s are highly coupled, then high T_a implies high T_s and a large flux of longwave radiation emitted from the surface into the atmosphere. Thus LW_{net} increases with T_a . High vapor pressure, e , means that much of the longwave flux emitted by the surface is absorbed by the atmosphere and emitted downward, so that $LW_{net} \approx \sigma T_s^4$ and LW_{net} is small. T and e both decrease with latitude. Their latitudinal changes may have counteracting influences on longwave radiation so that LW_{net} remains fairly independent of latitude. We use (6) to explore this possibility.

Low-level vapor pressure is a function of relative humidity and temperature. The observed relative humidity near the surface is ~80% with little dependence on latitude (London, 1957). Assuming similar behavior in the models' relative humidity fields, then

$$e \approx 0.8 e_{sat}(T_a), \quad (7)$$

where e_{sat} is the saturation vapor pressure. Substituting (7) into the first term on the right side of (6) and using the Clausius-Clapeyron relation,

$$e_{sat}(T) = (6.11 \text{ mb}) \exp[5411(1/273 - 1/T)], \quad (8)$$

we obtain

$$\epsilon \sigma T_a^4 (0.39 - 0.05\sqrt{e}) \approx 86(W/m^2) \times (1 + 0.015T') (1 - 0.014T'), \quad (9)$$

where T' is T_a in $^{\circ}\text{C}$. The first T' factor in (9) comes from T_a^4 ; the second comes from the vapor pressure, e . The two factors have opposing tendencies as T' changes and show that the first term on the right side of (6) has only a weak dependence on temperature. Calculations using representative values of T_a , T_s and e from the GFDL model indicate that this term dominates (6) at all latitudes but the tropics, where the second term is no longer negligible, but still smaller. Thus, the dependence of LW_{net} on temperature and moisture given by (6) indicates that the coupling of atmospheric temperature and humidity by the Clausius-Clapeyron relation constrains the variability of LW_{net} to be much less than $LW_{\uparrow}$ or $LW_{\downarrow}$.

Various modifications of (6) have been used that include effects of cloud cover (Stimpson and Paulson, 1979; Weare et al., 1981; Esbensen and Kushnir, 1981), but the clear sky formula is used here. Model clouds at different levels (not shown) have some latitudinal variability but they generally do not show a clear equator-pole trend. Cloud distribution appears only to modify behavior that is governed by the Clausius-Clapeyron relation.

Noteworthy features of the annual zonal mean energy balance (Figures 3 and 9) are the strong latitudinal dependence of LH and SW in all models, the importance of SH in the NCAR and GFDL polar regions and the departure of S from zero at many latitudes in the GISS model. The last two features are departures from the relative ordering of terms observed in the global mean.

Solar flux decreases with latitude because of the geometry of the planet and because surface albedo tends to increase with latitude. Latent heat flux decreases with latitude because the maximum flux is governed by the Clausius-Clapeyron relation, which constrains the flux to decrease with temperature. The magnitude of SH in the polar regions of the NCAR and GFDL models is not much greater than elsewhere, but the decrease of LH and SW with latitude allows it to assume greater relative importance there. Furthermore, in the polar regions of the NCAR and GFDL models, SH is directed into rather than out of the ground. The GISS model SH, in contrast, is positive at all latitudes and becomes small at high latitudes. During the DJF season part of the discrepancy between the GISS model SH and the SH of the other two models occurs because the GISS model has a region of strong sensible heat flux from the ocean to the atmosphere in the northern Atlantic Ocean (Figure 25a, discussed

in more detail in Section 7). However, the NCAR and GFDL models also have large areas of negative SH over land in the northern hemisphere during DJF, whereas SH is positive over most of the corresponding areas in the GISS model.

The largest contributor to nonzero annual mean S in the GISS model is probably the specified ocean transport, which cools the tropics (producing positive S) and warms the extratropics (producing negative S). Subsurface transport of heat into the polar regions may be a reason why SH does not warm the surface there in the GISS model as it does on the NCAR and GFDL models. Negative SH in these two models may be a compensation for lack of ocean transport. However, polar region heating by GISS ocean transport (negative S) is not as large as the polar heating by SH in the other two models, suggesting that differences in the polar region energy balances of the models are only partially attributable to differences in ocean modeling. In the NCAR and GFDL models, which lack ocean transport, S is nearly zero except at high latitudes, where snowmelt could explain the positive flux. The reason for negative S in the NCAR polar regions is not known.

Average flux values for extreme seasons are shown in Figures 12 for DJF and 19 for JJA. S tends to be negative in the winter hemisphere and positive in the summer hemisphere, indicating the importance of ocean storage in its annual cycle. Seasonal changes in solar flux appear to follow the annual cycle of the solar zenith angle. Seasonal variability of LH is relatively small, but the seasonal change that does occur has a phase opposite the annual cycle of shortwave flux, i.e. LH is strongest in the winter hemisphere. This can be understood by considering the annual cycle of near-surface moisture (Figs. 15 and 22). High summertime atmospheric humidity suppresses the latent heat flux, so the flux from the surface is weakest in summer.

The temperature change in the tropics when CO₂ doubles is greatest in the GISS model and smallest in the NCAR model for all three averaging periods. As with the global mean, changes in surface LW↓ are the greatest contributors to changes in surface temperature over most of the globe (Figures 4, 13 and 20). In the tropics, GISS ΔLW↓ is the largest, followed by GFDL and NCAR. The differences between model-predicted tropical temperature changes correspond closely to the differences in model-predicted low-level moisture (Figures 6, 15 and 22). Differences in surface temperature change in the tropics thus appear to be closely related to differences in moisture change through water

vapor's contribution to greenhouse warming. The same correspondence between water vapor change and surface temperature change also occurs at extratropical latitudes, though with less regularity.

The most prominent features of extratropical surface temperature and flux changes are zones of extreme change in high latitudes. In each model, these zones coincide with the latitudes of retreating sea ice. The extrema in the GISS model flux changes in (Figs. 13 and 20) are not as sharply peaked as for the GFDL and NCAR models, consistent with the difference in sea ice modeling between GISS and the other two. Each grid box in the GISS model can be partially covered with sea ice, whereas sea ice in the GFDL and NCAR models either covers a box completely or not at all. The coarser ice resolution in the GFDL and NCAR models leads to abrupt changes in ice cover and, as a consequence, in surface albedo and surface fluxes. In addition, the GISS sea ice parameterization has lower albedo compared to the other two models, which also results in a less abrupt change in surface heating where sea ice retreats compared to the other two models. Finally, the subsurface heat transport present in the GISS model may also alter its ice line sensitivity relative to the other two models (Meehl and Washington, private communication).

The GFDL model simulates larger LW_{net} and S changes in the Northern Hemisphere during JJA than the NCAR model, despite having a similar sea ice parameterization (Figure 20). The NCAR model's control climate sea ice cover in the Northern Hemisphere extends farther south (to about 60N) than the GFDL model's (70N). The latitude band where NCAR sea ice retreats as the climate warms is covered mostly by land, whereas the band where GFDL ice retreats includes more ocean. As a consequence, the areal extent of sea ice change and its resulting climatic impact in the zonal band where sea ice retreats is greater in the GFDL model. Differences in ice albedo modeling may also contribute to sensitivity differences at the ice margins of the two models. The GFDL parameterization produces greater ice melting than the NCAR model in a warming climate, thus allowing greater sea ice retreat (Washington and Meehl, 1986).

All three models show a peak in the ΔS contribution to ΔT_g at high latitudes in the winter-hemisphere (Figs. 13 and 20). Coincident temperature changes associated with ΔSH and ΔLH are negative because SH and LH increase. As a consequence of the increased energy transport into the atmosphere, the

downward longwave radiation also increases, although over a broader zone.

In the summer hemisphere, ΔS acts to cool the surface near the pole. Downward SW in the same latitude zone increases, especially in the GFDL model, so the S and SW changes tend to balance. During JJA for example, there is only a relatively small temperature change where ΔSW has its Northern Hemisphere minimum, and in each model the Northern Hemisphere ΔT_s maximum occurs further south (Figs. 18 and 20).

The narrow extrema of temperature and flux change in the winter hemispheres of the NCAR and GFDL models are "hot zones." When the climate warms these zones are strongly heated in summer by increased solar flux resulting from a decrease in summer ice cover. Much of the additional absorbed heat is then released in winter, as evidenced by the increased surface cooling by SH and LH (Figs. 13 and 20). The overlying atmosphere is warmed greatly by the additional winter flux, so that $\Delta LW\downarrow$ also peaks in the vicinity. The $\Delta LW\downarrow$ maximum is broader than the extrema of other fluxes, indicating that heat transported into the atmosphere is spread over a wider zone. Lack of an interactive ocean heat transport in the models prevents them from smoothing the surface hot zone within the ocean itself.

7. REGIONS

In this section, we present an evaluation of model data on regional scales by averaging the gridded output over the regions described in Section 3 (Fig. 1). All the regions except one fall in a midlatitude zone extending from 30°N to 50°N. Our chief interest here is seeing how the model behavior on smaller scales departs from the zonal mean averages presented earlier. However, we shall see that many of the regional differences are closely tied to zonal mean differences in model climates. We first show global contour plots of temperature and surface fluxes (Figs. 25 and 26) to illustrate some of the nonzonal characteristics of the model climates and to provide a spatial context for the particular regions we have defined. We then use scatter diagrams (Figs. 27 - 30) to describe features of the regions as a group. Finally, we focus on the behavior of some individual regions to show how regional variability manifests itself in the surface energy balance.

a. Contour plots

1) DJF control climate

For the sake of brevity, we show contour plots derived from model output for the DJF season only. Fig. 25 shows contour plots of sensible heat flux, latent heat flux, short wave radiation, net longwave radiation and surface air temperature for each model's control DJF climate. Fig. 26 shows changes in these fields when CO₂ doubles.

Surface sensible heat flux, SH, for the control climate in each model (Fig. 25a) has substantial longitudinal variability associated with the land/ocean distribution. The most intense fluxes are located off the east coasts of North America and Asia and over land in the southern (summer) hemisphere. Fluxes off the east coasts in the Northern Hemisphere result from cold, continental air advected over warmer ocean waters. Strong fluxes in the Sahara and the Southern Hemisphere continents probably represent responses to intense heating of relatively dry areas by the summer sun.

The most substantial differences in the distribution of SH between the models are broad areas where SH is negative (downward) in the NCAR and GFDL models but positive in the GISS model. Over one such area, north of Scandinavia, the GISS model SH is particularly intense. T_a in this area is warmer in the GISS model than in the other two models (Fig. 25e), suggesting that the flux difference can be attributed to specified sea surface temperatures and implied ocean heat transport in the GISS model. This transport can produce local conditions conducive to large upward SH by maintaining relatively warm ocean temperatures despite the strong cooling of polar air during winter. Why the GISS model has almost no flux into land surfaces is less clear. GISS SH can be negative for $T_a > T_s$. However, the GISS SH parameterization is also stability-dependent, and vertical heat flux is suppressed in highly stable conditions ($T_a > T_s$). The GISS model thus may not allow substantial sensible heat flux into the ground even when the ground is much cooler than the air. The model's coarser resolution and use of fractional land/ocean cover within each grid box may provide an additional reason. Because of the coarse resolution, only a few North American grid boxes and only about half of the Eurasian grid boxes are 100% land covered. Water surface temperatures are specified in the GISS control run, so any amount of water in a grid box is potentially a strong source of sensible heat. Upward sensible heat flux from

the water portion of a grid box may dominate the box-averaged flux. The GISS model's resolution and fractional land/ocean cover within grid boxes may also account for the weaker gradient of sensible heat flux between continents and adjacent oceans in this model. Finally, the GISS model's resolution is probably also responsible to the absence of a local maximum in SH in the vicinity of the Mediterranean that is present in the other two simulations.

Departures of the latent heat flux, LH, from zonal symmetry are also governed primarily by land/ocean distribution (Fig. 25b). The strongest fluxes occur over oceans, where there is abundant moisture supply. In the extratropics, relatively strong fluxes appear along the east coasts of North America and Asia, where cool, dry continental air passes over the warmer ocean. In each of these areas, the sea surface temperatures are about the same in all three models. However, the GISS model fluxes are considerably more intense, suggesting that its atmosphere's circulation transports more dry air into the areas of large LH to absorb heat from the ocean than the other two models do. The GISS model's implied ocean heat transport appears to be especially important in the energy balance of these areas because it helps maintain local sea surface temperatures that are similar to those in the NCAR and GFDL models despite its much stronger LH. The strong fluxes along continental coasts cause the GISS model's zonal mean latent heat flux to peak in the northern subtropics (Fig. 12). The GISS model also shows a region of relatively low flux along the west coast of North America, presumably because it specifies observed ocean temperatures that are cooler than simulated in the NCAR and GFDL models, thus inhibiting latent heat flux.

Net surface solar fluxes, SW, (Fig. 25c) display departures from zonal symmetry in the southern subtropics of all models that appear to be associated with land/ocean distribution. Compared to the NCAR and GISS models, the GFDL model simulates greater departures from zonal symmetry. Furthermore, the zone of greatest departures, where longitudinal variations of SW are as large as latitudinal variations, extends from about 60°S to 30°N in the GFDL model, but from only about 45°S to 15°N in the GISS and NCAR models. The NCAR zone of strongest flux ($> 240 \text{ Wm}^{-2}$ into the surface) is displaced to the south compared to the other two models, a feature that also appears in the zonal mean (Fig. 12).

Strong gradients in SW appear in the NCAR and GFDL models near

Antarctica, but the variation with latitude in the GISS model is less abrupt there. The smoother variation of GISS model SW appears to be the result of fractional ice cover being allowed in each grid box (yielding a less abrupt gradient of the surface albedo) and coarser resolution.

The net longwave flux, LW_{net} , (Fig. 25d) varies more smoothly over the globe than the other fluxes, a feature in accord with its behavior in the zonal mean (e.g., Fig. 12). Spatial variability of LW_{net} is especially small over the ocean; the strongest gradients are associated with land masses, particularly in the NCAR and GFDL models. Recall from Section 6 that if relative humidity remains constant, then radiative contributions to LW_{net} from changes in atmospheric moisture compensate contributions from changes in T_a so that LW_{net} is insensitive to changes in T_a . Local maxima of LW_{net} in the subtropics appear to be associated with relatively dry conditions, since they coincide with local minima in LH. These extrema also coincide with dry areas in the observed climate. The GISS control climate's gradients in LW_{net} are weaker than the gradients in the other models. As with the other surface fluxes, this characteristic again appears to be a result of the GISS model's coarser resolution and fractional land/ocean distribution within a grid box.

Compared to the fluxes shown in Figs. 25a-25d, the surface air temperature, T_a , (Fig. 25e) shows more zonal symmetry. The GFDL model's T_a has some small scale spatial variability in the Indian Ocean and tropical Pacific that is barely resolved by its grid and may be some type of 2Δ noise. Otherwise, departures from zonal symmetry are associated with land/ocean distribution. Gradients associated with the departures tend to be relatively weak compared to gradients in the surface fluxes. There are no discernible differences in gradients that might be attributable to model details such as resolution or grid box definition. Instead, horizontal heat transports in the atmosphere, which tend to reduce temperature gradients, appear to be smoothing the T_a distribution that would be forced by the surface fluxes alone. The smoothing is of interest because in each model T_a is the temperature for a shallow layer near the surface, where one might expect horizontal heat transports to be weak and relatively unimportant in the local thermodynamics. Surface temperature, T_s , is controlled by the surface energy balance (2) and, hence, is not affected directly by horizontal heat transport in the atmosphere. The T_s field is not as smooth as the T_a field, as indicated by the distribution of SH

(Fig. 25a) which is proportional to $T_s - T_a$.

2) DJF ($2CO_2$ - control) changes

The biggest changes in SH when CO_2 doubles (Fig. 26a) occur where the polar sea ice retreats in the NCAR and GFDL models. Maximum values of ΔSH are much weaker in the GISS model, a feature that again may be attributed to its coarser resolution and fractional ice cover within the grid box. The ΔSH maxima in the NCAR and GFDL models are about the same magnitude but the GFDL ice line shift exposes more ocean, so that zonal mean ΔSH is larger in the GFDL model, as noted earlier in Section 6. The NCAR model simulates large ΔSH increases over land in the southern subtropics, where the other two models have at most fairly modest increases. Except for this feature, regional changes in ΔSH are explained well by zonal averages.

The spatial variability of LH changes when CO_2 doubles (Fig. 26b) is smaller than for changes in SH. Also, ΔLH varies more uniformly with latitude than control climate LH, which is concentrated toward the tropics in each model (Fig. 25b). Increases in LH occur in the NCAR and GFDL models where the Arctic ice margin shifts, but the LH changes are smaller than collocated SH increases. Cool temperatures near the ice margins constrain the amount of water vapor the atmosphere can hold, thereby limiting the amount that LH can increase. The sensible heat flux is not so constrained.

Maximum GISS model ΔLH occurs over near-equatorial South America and the tropical western Pacific, where LH increases by more than 20 Wm^{-2} . There are no corresponding regions in the other two models; on the contrary, the NCAR model simulates a relatively strong decrease in LH over South America. The NCAR model has regions of strong LH decrease over each of the southern subtropical land masses. Decreases in LH in these regions in the NCAR model partially compensate for the large increases in SH occurring there.

The shortwave fluxes have their greatest changes (Fig. 26c) where the polar ice margins shift in the Southern Hemisphere. Otherwise there is little spatial variability in the ΔSH field and no evidence of systematic changes that might be caused by other factors such as clouds. Net longwave flux change (Fig. 26d) also shows relatively small spatial dependence and generally does not deviate much from the zonal mean. The largest changes appear to be associated with ice line shifts and, for the NCAR model, with southern

subtropical land masses. Large temperature changes in the NCAR and GFDL models (Fig. 26e) occur over ocean areas where sea ice retreats.

b. Scatter diagrams

For each of the regions in Figure 1 we have constructed tables of model output that correspond to Tables 2 - 4 for global averages. The data thus assembled (Tables 5 - 7) comprise many tables that are difficult to study en masse. In order to summarize some aspects of the regional climate, we have constructed scatter diagrams in which regional means for one variable are plotted against the regional means for another. If two models have very similar regional climatology, then points from the two models representing the same region should lie close to each other. Table 8 gives an identification key for the plotted points.

Figure 27 shows plots of ΔSW vs. ΔT_a . The points tend to cluster by model, especially in the JJA season. This behavior also occurs in diagrams (not shown) using Δ (mixing ratio) or Δ (cloud cover) as a variable. Typically, the clustering of points by model corresponds to differences in the model zonal means. For example, zonal mean JJA ΔT_a in midlatitudes is largest in the GFDL model and smallest in the NCAR model (Fig. 18). Also, NCAR zonal mean solar flux weakens ($\Delta SW > 0$) in midlatitudes during JJA, whereas GFDL and GISS zonal mean solar flux tends to strengthen (Fig. 20). The zonal mean differences between models are larger than the regional differences simulated by a particular model, so locations of the points for each model in Figure 27b follow the same ordering as the zonal mean. The stratification of regional temperature behavior by model in this figure suggests that for some fields better agreement among models on zonal mean scales is necessary before regional climate can be simulated with accuracy.

Plots of Δ (precipitation) vs. precipitation (Fig. 28) exhibit less clustering by model. Instead, points representing the same region in different models tend to lie close to each other relative to the overall spread of points on the diagram. The models thus appear to simulate regional differences in precipitation with some accuracy.

Precipitation does not play a direct role in the surface energy balance. Nonetheless, it is of interest to understand why regional consistency among the models occurs for this field but not for others examined, especially since

precipitation is such a notoriously difficult field to forecast. Although a definitive diagnosis of why the model simulations of regional precipitation change are more consistent than their simulations of regional temperature change is beyond the scope of this report, we can hypothesize one possible reason. Precipitation is partly determined by local orography, a regional feature that is specified for each model and, hence, not dependent on a model's zonal mean climatology. Except for WEP, the regions studied here are located in the Northern Hemisphere midlatitudes, where the three models show their best agreement in zonal mean control climate precipitation and precipitation change (Fig. 14). Much of the Northern Hemisphere midlatitudes are land-covered, so that topography plays an important role in forcing precipitation there. Precipitation over ocean-covered regions of the globe is not controlled by topography, and we observe that the precipitation simulations are less consistent in the tropics and southern hemisphere (Fig. 14).

In contrast to precipitation, fields such as temperature and mixing ratio may be more highly dependent on a model's transports. Accurate modeling of regional features of transport climatology should be far more difficult than specifying the topography. Precipitation depends on temperature and moisture transports too, but our results suggest that the dependence is weaker than for other fields.

Three other sets of diagrams are included here for the insight they provide on regional energy balance in the models. Figure 29 shows $\Delta LW\downarrow$ vs. $\Delta LW\uparrow$ for winter and summer. In the global and zonal means, these two changes tend to balance: $4\sigma T_a^3 \Delta T_a \approx -\Delta LW\downarrow$. The same approximate balance is found in individual regions. Points for the same region in different models do not lie so close together as in Fig. 28, indicative of less regional consistency between the model climates for these fields. Nonetheless, there is some apparent consistency, in that the balance between $\Delta LW\downarrow$ and $\Delta LW\uparrow$ is about the same for each model.

Figure 30 shows ΔSH vs. ΔLH for summer and winter. In both seasons, but especially in summer, there is a tendency for ΔSH and ΔLH to compensate so that their sum is about the same for all regions in all models. The points in Fig. 30 are not clustered by model, which suggests that the behavior depicted is fairly robust. Taking all data together indicates that $\Delta SH + \Delta LH \approx$

+ 5Wm^{-2} for both seasons. This compensation places a constraint on how the surface energy balance can vary, especially if we recall the constraint observed earlier that prevents LW_{net} from changing when relative humidity is constant. Using (2) and the two constraints suggests that for these regions $\Delta\text{SW} + \Delta\text{S} \approx -5\text{Wm}^{-2}$, i.e., that storage changes are primarily a direct response to changes in solar radiation.

The final scatter diagrams show mixing ratio vs. T_a for the control and doubled CO_2 climates (Fig. 31). In winter, there is a clear relationship between T_a and mixing ratio consistent with near-constant relative humidity. In summer there is considerably more scatter, indicating that relative humidity has widely different values between models for the same region. Many simplified climate models assume that relative humidity does not vary horizontally. In the summer, this assumption is not validated by the NCAR model. Furthermore, because atmospheric moisture strongly affects longwave radiative transfer, the disparity between models in atmospheric mixing ratio will contribute to differences in the models' surface energy balances through their influence on $\text{LW}\downarrow$. Differences in atmospheric moisture will also affect LH . Figure 31 illustrates the need to model atmospheric moisture transports more consistently on regional scales before regional climate can be simulated accurately.

c. Individual regions

Tables 5-7 present data for each region that correspond to the global averages that are presented in Tables 2-4. We focus here on the surface energy balance and its changes for the central United States region (USC) during JJA. This region is important for agriculture, and projected changes in the climate of this region caused by increasing CO_2 have been the subject of some controversy (Manabe and Wetherald, 1986a, 1986b, 1987; MacCracken et al., 1986; Meehl and Washington, 1988) because the GFDL model predicts that the summer climate will become significantly hotter and drier. In this section we compare this GFDL prediction with simulations by the NCAR and GISS models in the context of the surface energy balance, illustrating how hydrological cycle differences can dramatically alter the energy balance on a regional scale.

As with larger scale averages, USC temperature changes associated with

LW↓ are largest and they approximate the actual temperature change (Table 7b). The models give nearly the same $\Delta LW\downarrow$, but for other flux changes, there are considerable differences among the models. For the GFDL model, in particular, ΔSH and ΔSW assume an importance generally absent in the global and zonal averages. In contrast with the global mean, GFDL LH decreases, giving a strong positive contribution to ΔT_s . For NCAR and GISS, the other flux changes, including ΔLH , are relatively small.

Some insight into the differences between the GFDL model and the others can be obtained by considering aspects of the hydrological cycle. Table 6b shows changes in precipitation and high cloud cover. GFDL precipitation diminishes considerably: the decrease is one third of the 1 CO_2 climate seasonal mean. NCAR and GISS, on the other hand, show slight increases. The model-predicted changes in high cloud cover are consistent with these precipitation changes, considering that most precipitation over the Midwest in summer is convective. The total cloud cover in the GFDL model (not shown) also decreases.

The GFDL hydrological changes in Table 6b are associated with a soil moisture decrease in the central US that occurs when CO_2 doubles in this model (Manabe and Wetherald, 1987). Drier ground means less water can evaporate into the atmosphere. As a consequence, LH decreases which contributes to surface heating. The decrease in total cloud cover (not shown) allows more surface heating by short wave radiation. The greater temperature increase for GFDL in this region compared to the other models is consistent with the additional heating. Associated with GFDL's surface warming and LH decrease is a strong increase in SH, which opposes the net warming of the surface.

Comparison of the USC climate changes produced by the three models indicates that the GFDL simulation is inconsistent with the other two. However, in other regions one of the other models may differ from the rest in its surface energy balance. For example, in the adjacent western United States region during JJA (Table 7a and Fig. E-3), the GISS model SH and LH change in the opposite direction of the corresponding fluxes in the GFDL and NCAR models. In the northeastern China region during JJA (Table 7g and Fig. E-3), SH increases and LH decreases in the NCAR simulations whereas the GISS and GFDL simulations produce the opposite changes. The discrepancies are not as big as for the USC region, but nonetheless they demonstrate that significant

inconsistency between model simulations on regional scales is typical. The inconsistency between model simulations discussed here is not a manifestation of zonal mean differences between the models for the three regions analyzed are located in the same latitude belt. Rather, the inconsistency is emblematic of differences in model simulations occurring on the regional scale.

Table 5 also contains observationally-based surface fluxes for China (Gao and Lu, 1981) and for the ocean regions (Esbensen and Kushnir, 1981). As with the global fluxes in Table 2, these fluxes are obtained using empirical formulas with observations such as wind speed and air temperature as input. Weare and Strub (1981) and Fung et al. (1984), among others, have discussed the accuracy of the fluxes produced by such a procedure. Sampling biases, random errors and errors in the formulas themselves may produce net errors of 30% or more, an accuracy also suggested by the two sets of observationally-based fluxes in Table 2. For the Chinese regions, the tables contain LW_{net} (denoted by brackets) since appropriate temperatures for separating $LW_{\uparrow}$ and $LW_{\downarrow}$ are not available.

In the Chinese regions, model fluxes often differ from the observationally based fluxes by more than the 30% error estimate. The differences occur in part because the models show greater disparity on the regional scale (cf. Table 2) and in part because of model biases relative to the observational fluxes. Simulated latent heat fluxes are consistently much larger than observed. Modeled sensible heat fluxes often have the opposite sign of observed fluxes. There is also a tendency for modeled solar fluxes to be stronger than observational values.

In the ocean regions, discrepancies between the model and observational fluxes tend to be smaller. This may occur because the oceans present a more uniform lower surface to the atmosphere than land regions do: the ocean surfaces are flat and always 100% saturated. Intermodel discrepancies tend to be larger for these regions than for the global mean, but they are also less than those occurring over China. The most significant model bias relative to the observations appears to be a tendency for modeled solar fluxes to be stronger than the observational values.

8. SUMMARY

We examine the surface energy balance simulated by three state-of-the-art general circulation models for climates with current levels of atmospheric CO₂ concentration (control climate) and with twice the current level and compare the simulated balances by averaging fields of interest over a hierarchy of spatial domains ranging from the entire globe down to regions a few hundred kilometers across.

Differences between models' representations of the surface energy balance become greater as the spatial scale of comparison decreases. For all models, scales and climates, the first-order energy balance is between upward and downward surface longwave fluxes (LW↑ and LW↓, respectively). This is simply the greenhouse warming of the surface. Furthermore, the net longwave flux at the surface has small spatial variability, which we ascribe to the small variability of relative humidity. Because changes in LW↑ are closely tied to surface temperature changes, changes in LW↓ and surface temperature are also tightly coupled. Thus, accurate modeling of the longwave radiation field is essential if GCM climate projections of surface temperature change are to be reliable.

Although qualitative agreement is found among the models' energy balances, there are important quantitative discrepancies at all spatial scales. Most important, differences in LW↓ between models are as large as the changes in LW↓ that occur when CO₂ doubles. Because longwave radiation dominates the surface energy balance, the differences place a limit on the accuracy of temperature change simulated by state-of-the-art models.

Different parameterizations of ocean ice also create quantitative discrepancy between models. The NCAR and GFDL models have abrupt changes in their zonal mean surface energy balances and strong surface heating where the ice line retreats when CO₂ doubles. The abruptness is not present in the GISS zonal mean energy balance. Differences between modeled flux changes near ice margins may be as large as 100 Wm⁻². In the NCAR and GFDL models, ocean ice covers the entire surface of any grid box containing ice. In the GISS model, a grid box can be partially ice-covered. Ice cover in the NCAR and GFDL models thus shifts abruptly when CO₂ doubles in these models, producing large increases in surface heating near ice margins. The location of the control climate ice margins also appears to be important. Differences in surface

energy balance change between the NCAR and GFDL models can be attributed to differences in the location of ocean ice margins in the model control climates.

Regional-scale differences between model-generated surface temperatures are often attributable to zonal-average differences. Thus, better agreement among model simulations of zonal mean climatology must be obtained to improve climate modeling on regional scales.

Additional discrepancies also occur on regional scales that are not only quantitative but also qualitative. Control climate fluxes may be considerably different in different models. Equally important, they may change in different directions when CO₂ doubles. These discrepancies are not attributable to zonal mean differences between models, but rather pertain to intermodel differences in simulating the climate of regions themselves. A prominent example is latent heat flux change over the central United States, a region crucial for agriculture.

A better understanding of the hydrological cycle and how to simulate it appears to be especially important for improving regional climate model capabilities. The ordering among models for surface air temperature change in a region typically matches the ordering of atmospheric humidity change, implying a close association between the hydrological cycle and longwave radiation. Water vapor and clouds interact strongly with LW↓, and differences in both between models may account for the large discrepancies in modeled LW↓. Other hydrological cycle differences, such as soil moisture, also appear to be associated with the qualitative discrepancies between models on regional scales discussed above. The results suggest that accurate modeling of moisture transport through the atmosphere is important for achieving reliable regional climate modeling. Accurate modeling of moisture transport may also be important for reducing differences in zonal and global mean surface energy budget climatology.

Precipitation and its change on regional scales are simulated with more consistency than surface temperatures and fluxes. We speculate that precipitation amounts over land are constrained in these models by topography, which can be accurately specified. Since most of the regions we examine are land-covered, topographic control could account for the intermodel consistency we observe. If this is so, then precipitation modeling may be improved by

simply increasing model resolution. For studies of specific regions, resolution could be increased selectively using a nested grid.

Acknowledgements

We thank W. Washington (NCAR), S. Manabe (GFDL) and J. Hansen (GISS) for providing us with the model data analyzed here, and L. Ver Plank (NCAR), R. Wetherald (GFDL) and G. Russell (GISS) for taking the time to answer our numerous questions about the models and the data. G. Meehl and W. Washington provided extensive reviews of the manuscript as it was prepared. M. B. Phillips, A. Aubin, and T. Foley performed the text editing; M. B. Phillips and M. Frunke patiently prepared the figures. This work was supported by Department of Energy grant DE-FG02-86ER60422.

REFERENCES

- Berliand, M. E., and T. G. Berliand, 1952: Determining the net long-wave radiation of the Earth with consideration of the effect of cloudiness. Isv. Akad. Nauk. SSSR Ser. Geofiz., No. 1.
- Brunt, D., 1932: Notes on radiation in the atmosphere. Quart. J. R. Met. Soc., 58, 389-420.
- Cess, R. D., and G. L. Potter, 1987: Exploratory studies of cloud radiative forcing with a general circulation model. Tellus, 39A, 460-473.
- Esbensen, S. K., and Y. Kushnir, 1981: The Heat Budget of the Global Ocean: An Atlas Based on Estimates from Surface Marine Observations. Report No. 29, Climate Research Institute, Oregon State University, 244 pp.
- Fung, I. Y., D. E. Harrison and A. A. Lacis: On the variability of the net longwave radiation at the ocean surface. Rev. Geoph. Sp. Phys., 22, 177-193.
- Gao, G.-D., and Y.-R. Lu, 1981: Atlas of Physical Climatology of China. Aquaculture Press, Beijing 183 pp.
- Hansen, J., A. Lacis, D. Rind, G. Russell, P. Stone, I. Fung, R. Ruedy and J. Lerner, 1984: Climate sensitivity: Analysis of feedback mechanisms. In Climate Processes and Climate Sensitivity, Geophys. Monogr. Ser., Vol. 29, J. E. Hansen and T. Takahashi (eds.), AGU, Washington, D.C., 130-163.
- Hansen, J., G. Russell, D. Rind, P. Stone, A. Lacis, S. Lebedeff, R. Ruedy, and L. Travis, 1983: Efficient three-dimensional global models for climate studies: Models I and II. Mon. Weather Rev., 111, 609-662.
- Hsu, C.-P. F., and J. M. Wallace, 1976: The global distribution of the annual and semiannual cycles in precipitation. Mon. Wea. Rev., 104, 1093-1101.
- London, J., 1957: A study of the atmospheric heat balance. Final report, AFCRC Cont. AF19 (122)-165, College of Engineering, New York University, 99 pp. [ASTIA 117227].
- MacCracken, M. C., M. E. Schlesinger and M. E. Riches, 1986: Atmospheric carbon dioxide and summer soil wetness. Science, 234, 659.
- Manabe, S., and R.J. Stouffer, 1980: Sensitivity of a global climate model to an increase of CO₂ concentration in the atmosphere. J. Geophys. Res., 85, 5529-5554.
- Manabe, S., and R. T. Wetherald, 1986a: Reduction in summer soil wetness induced by an increase in atmospheric carbon dioxide. Science, 232, 626-628.
- Manabe, S., and R. T. Wetherald, 1986b: Atmospheric carbon dioxide and summer soil wetness. Science, 234, 660.

- Manabe, S., and R. T. Wetherald, 1987: Large-scale changes of soil wetness induced by an increase in atmospheric carbon dioxide. J. Atmos. Sci., 44, 1211-1235.
- Manabe, S., and R. T. Wetherald, 1967: Thermal equilibrium of the atmosphere with a given distribution of relative humidity. J. Atmos. Sci., 24, 241-259.
- Meehl, G. A., and Washington, W. M., 1988: A comparison of soil moisture sensitivity in two global climate models. J. Atmos. Sci. (in press).
- Miller, J. R., G. L. Russell and L.-C. Tsang, 1983: Annual oceanic heat transports computed from an atmospheric model. Dyn. Atmos. Oc., 7, 95-109.
- National Academy of Sciences, 1975: Understanding Climate Change. National Academy of Sciences, Washington, D. C.
- Russell, G. L., J. R. Miller and L.-C. Tsang, 1985: Seasonal oceanic heat transports computed from an atmospheric model. Dyn. Atmos. Oc., 9, 253-272.
- Schlesinger, M. E., and J. F. B. Mitchell, 1987: Climate model simulations of the equilibrium climate response to increased carbon dioxide. Rev. Geophys., 25, 760-798.
- Wallace, J. M., and P. V. Hobbs, 1977: Atmospheric Science: An Introductory Survey, Academic Press, 467 pp.
- Wang, W.-C., and G. Molnar, 1985: A model study of the greenhouse effects due to increasing atmospheric CH₄, N₂O, CF₂Cl₂, and CFCl₃. J. Geophys. Res., 90, 12,971-12,980.
- Washington, W. M., and G. A. Meehl, 1984: Seasonal cycle experiments on the climate sensitivity due to a doubling of CO₂ with an atmospheric general circulation model coupled to a simple mixed layer ocean model. J. Geophys. Res., 89, 9475-9503.
- Washington, W. M., and G. A. Meehl, 1986: General circulation model CO₂ sensitivity experiments: Snow-sea ice albedo parameterizations and globally averaged surface air temperature. Climate Change, 8, 231-241.
- Weare, B. C., and P. T. Strub, 1981: The significance of sampling biases on calculated monthly mean oceanic surface heat fluxes. Tellus, 33, 211-224.

List of Tables

Table 1	Transfer Coefficient C in Each Model
Table 2	Global Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates and for Annual Average and Extreme Seasons
Table 3	Global Mean Components of the Hydrological Cycle
Table 4	Global Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons
Table 5a	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region USW
Table 5b	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region USC
Table 5c	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region USE
Table 5d	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region CSE
Table 5e	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region CSW
Table 5f	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region CNW
Table 5g	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region CNE
Table 5h	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region NP
Table 5i	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region NA
Table 5j	Regional Mean Surface Energy Balance in the Three GCMs for Control and Doubled-CO ₂ Climates, for Average and Extreme Seasons - Region WEP
Table 6a	Regional Mean Components of the Hydrological Cycle - Region USW
Table 6b	Regional Mean Components of the Hydrological Cycle - Region USC
Table 6c	Regional Mean Components of the Hydrological Cycle - Region USE
Table 6d	Regional Mean Components of the Hydrological Cycle - Region CSE
Table 6e	Regional Mean Components of the Hydrological Cycle - Region CSW
Table 6f	Regional Mean Components of the Hydrological Cycle - Region CNW

Table 6g	Regional Mean Components of the Hydrological Cycle - Region CNE
Table 6h	Regional Mean Components of the Hydrological Cycle - Region NP
Table 6i	Regional Mean Components of the Hydrological Cycle - Region NA
Table 6j	Regional Mean Components of the Hydrological Cycle - Region WEP
Table 7a	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region USW
Table 7b	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region USC
Table 7c	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region USE
Table 7d	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region CSE
Table 7e	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region CSW
Table 7f	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region CNW
Table 7g	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region CNE
Table 7h	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region NP
Table 7i	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region NA
Table 7j	Regional Temperature Change Associated with Separate Terms of the Surface Energy Balance in the Three GCMs for Annual Average and Extreme Seasons - Region WEP
Table 8	Point Identification Key for Scatter Plots

Table 1
Transfer Coefficient C in Each Model

	<u>land</u>	<u>ocean</u>
NCAR	$4 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
GFDL	$3 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
GISS	A function of geographically-varying surface roughness and boundary layer stability.	

Table 2
Global Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates and
for Annual Average and Extreme Seasons

		[units: T _s -K, others-W/m ²]							
		a) current (control)				b) 2 CO ₂			
		OBS		NCAR	GFDL	GISS	NCAR	GFDL	GISS
		NAS	Budyko						
ANNUAL	T _a			286.9	287.7	287.1	290.5	291.8	291.3
	SW	-176.0	-143.6	-181.8	-164.0	-173.0	-183.1	-165.5	-175.9
	LW↓	-312.2	-336.8	-311.3	-326.2	-335.3	-335.2	-353.7	-362.5
	LW↑	+384.7	+384.7	+384.2	+388.5	+385.2	+403.9	+411.1	+408.3
	LH	+79.4	+79.8	+96.1	+83.0	+91.5	+103.2	+90.1	+101.9
	SH	+24.2	+16.0	+14.0	+17.1	+25.6	+11.3	+16.5	+22.3
	S	-0.1	-0.1	-1.1	+1.6	+6.0	-0.1	+1.5	+5.9
DJF	T _a			284.4	285.7	285.6	288.1	289.9	290.0
	SW			-182.6	-168.5	-177.1	-185.9	-169.0	-180.2
	LW↓			-299.2	-315.8	-327.9	-322.4	-343.6	-356.1
	LW↑			+370.9	+377.8	+377.2	+390.6	+400.5	+401.0
	LH			+96.3	+81.0	+91.1	+103.4	+88.6	+101.5
	SH			+10.4	+14.6	+23.9	+8.0	+13.9	+20.5
	S			+4.2	+10.9	+12.8	-6.3	+9.6	+13.3
JJA	T _a			289.4	290.1	288.7	292.8	293.9	292.7
	SW			-178.5	-161.1	-168.6	-177.7	-163.4	-171.3
	LW↓			-323.8	-339.1	-344.5	-347.4	-365.0	-370.5
	LW↑			+397.7	+401.6	+393.9	+416.7	+423.0	+416.2
	LH			+95.7	+85.4	+94.8	+102.9	+92.3	+104.9
	SW			+17.6	+21.3	+27.6	+15.4	+20.8	+24.0
	S			-.7	-8.1	-3.2	-9.9	-7.7	-3.3

Table 3
Global Mean Components of the Hydrological Cycle

		a) current (control)		b) 2 CO ₂			c) 2 CO ₂ - control			
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation [mm/day]	3.25	2.87	3.18	3.48	3.11	3.53	+0.23	+0.24	+0.35
	Moisture [g/kg]	9.75	11.91	10.34	11.62	14.48	13.19	+1.87	+2.57	+2.85
	Low Cloud [fraction]	0.19	0.26	0.36	0.19	0.26	0.34	0.00	0.00	-0.02
	High Cloud [fraction]	0.12	0.13	0.29	0.11	0.13	0.29	-0.01	0.00	0.00
	Lapse Rate [K/km]	5.85	5.58	5.48	5.85	5.52	5.23	-0.00	-0.06	-0.25
DJF	Precipitation	3.24	2.79	3.15	3.46	3.05	3.50	+0.22	+0.26	+0.36
	Moisture	9.21	11.36	9.74	10.96	13.81	12.52	+1.75	+2.45	+2.78
	Low Cloud	0.21	0.25	0.37	0.20	0.25	0.35	-0.00	+0.01	-0.02
	High Cloud	0.11	0.13	0.29	0.11	0.13	0.30	-0.01	-0.01	+0.01
	Lapse Rate	5.59	5.30	5.47	5.62	5.30	5.20	+0.03	0.00	-0.27
JJA	Precipitation	3.24	2.94	3.28	3.49	3.18	3.63	+0.25	+0.23	+0.35
	Moisture	10.19	12.48	11.19	12.09	15.11	14.12	+1.90	+2.62	+2.94
	Low Cloud	0.18	0.27	0.35	0.18	0.27	0.33	+0.00	-0.00	-0.02
	High Cloud	0.11	0.13	0.29	0.11	0.12	0.28	-0.00	-0.01	-0.01
	Lapse Rate	6.08	5.85	5.46	6.07	5.74	5.23	-0.10	-0.10	-0.23

Table 4
Global Temperature Change Associated with Separate
Terms of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
[units: K]

		2 CO ₂ - control		
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+3.6	+4.1	+4.2
	ΔSW	+0.2	+0.3	+0.5
	$\Delta LW\downarrow$	+4.5	+5.1	+5.1
	ΔLH	-1.3	-1.3	-1.9
	ΔSH	+0.5	0.1	+0.6
	ΔS	-0.2	+0.0	+0.0
DJF	ΔT_a	+3.7	+4.2	+4.3
	ΔSW	+0.6	+0.1	+0.6
	$\Delta LW\downarrow$	+4.4	+5.2	+5.2
	ΔLH	-1.3	-1.4	-2.0
	ΔSH	+0.5	+0.1	+0.6
	ΔS	-0.4	+0.2	-0.1
JJA	ΔT_a	+3.5	+3.8	+4.0
	ΔSW	-0.2	+0.4	+0.5
	$\Delta LH\downarrow$	+4.4	+4.7	+4.8
	ΔLH	-1.3	-1.2	-1.9
	ΔSH	+0.4	+0.1	+0.7
	ΔS	+0.2	-0.1	0.0

Table 5a
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region USW

[units: T_s-K, others-W/m²]

		a) current (control)			b) 2 CO ₂		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _a	284.7	282.0	281.2	287.2	286.9	286.0
	SW	-188.4	-156.1	-150.7	-184.3	-163.1	-151.7
	LH↓	-283.7	-277.4	-307.9	-302.9	-303.0	-334.5
	LH↑	+372.5	+358.4	+354.6	+385.8	+384.2	+379.6
	LH	+59.0	+44.9	+59.3	+64.6	+48.3	+67.0
	SH	+40.8	+26.2	+42.3	+37.0	+31.6	+38.2
	S	-0.2	+4.0	+2.4	-0.2	+2.0	+1.4
DJF	T _s	273.7	271.0	272.8	276.4	275.4	277.9
	SW	-88.6	-66.0	-67.8	-86.3	-74.2	-68.0
	LH↓	-257.0	-255.5	-282.4	-272.8	-276.6	-308.2
	LH↑	+318.0	+305.9	+314.0	+311.2	+326.3	+338.3
	LH	+62.8	+33.4	+40.3	+66.6	+40.2	+46.2
	SH	-34.8	-25.8	+18.1	-38.2	-20.4	+13.7
	S	-0.3	+7.9	-22.1	-0.4	+4.8	-22.0
JJA	T _s	297.7	295.7	290.0	300.8	301.3	293.9
	SW	-287.7	-267.9	-231.6	-281.0	-270.6	-231.7
	LH↓	-322.6	-306.6	-344.7	-349.2	-342.0	-372.2
	LH↑	+445.3	+433.3	+400.8	+463.8	+467.0	+423.1
	LH	+42.8	+49.2	+87.5	+40.8	+45.6	+101.9
	SH	+122.5	+91.3	+61.3	+125.5	+99.4	+52.3
	S	-0.3	+0.7	+26.7	+0.2	+0.7	+26.6

Table 5b
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region USC

[units: T_s-K, others-W/m²]

		a) current (control)			b) 2 CO ₂		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _a	287.4	285.3	282.8	289.8	290.8	287.7
	SW	-194.1	-170.1	-148.2	-189.4	-178.3	-150.7
	LW↓	-291.4	-297.2	-312.5	-312.4	-325.7	-338.4
	LW↑	+386.8	+375.4	+362.9	+399.9	+405.2	+388.5
	LH	+43.6	+62.4	+53.2	+54.0	+62.2	+55.3
	SH	+55.1	+27.1	+43.7	+48.0	+35.0	+45.1
	S	0.0	+2.4	+0.0	-0.1	+1.6	+0.2
DJF	T _a	271.6	270.1	272.1	274.9	275.3	277.6
	SW	-101.1	-88.9	-71.6	-97.3	-93.8	-72.4
	LW↓	-235.2	-235.2	-274.2	-256.6	-264.1	-299.4
	LW↑	+308.5	+301.8	+311.0	+323.8	+325.5	+336.4
	LH	+40.2	+39.6	+19.0	+50.0	+46.6	+20.9
	SH	-12.3	-23.0	+22.3	-20.0	-18.2	+21.8
	S	0.0	+5.6	-6.6	-0.3	+4.0	-7.5
JJA	T _a	303.0	300.7	292.8	306.4	307.1	296.6
	SW	-282.3	-256.6	-219.8	-276.0	-276.5	-222.4
	LW↓	-359.5	-365.5	-361.2	-387.4	-393.4	-385.5
	LW↑	+478.2	+463.6	+416.9	+499.4	+504.5	+438.8
	LH	+42.6	+80.8	+101.6	+42.9	+64.3	+104.4
	SH	+121.1	+77.0	+54.2	+120.9	+100.4	+57.1
	S	-0.1	+0.8	+8.2	+0.2	+0.7	+7.5

Table 5c
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region USE

[units: T_s-K, others-W/m²]

		a) current (control)			b) 2 CO ₂		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _a	286.7	283.8	283.8	289.6	289.7	287.9
	SW	-184.0	-157.6	-149.6	-177.4	-165.3	-153.6
	LW↓	-297.0	-298.8	-321.5	-322.9	-329.2	-343.8
	LW↑	+382.9	+367.8	+367.9	+399.0	+399.3	+389.7
	LH	+54.2	+66.0	+89.3	+75.2	+67.6	+96.6
	SH	+44.1	+19.8	+47.3	+26.3	+26.0	+45.4
	S	0.0	+2.7	-33.1	-0.2	+1.6	-34.3
DJF	T _s	270.3	268.4	275.1	273.6	274.5	279.6
	SW	-89.0	-72.6	-67.8	-85.0	-79.7	-68.1
	LW↓	-235.6	-238.3	-285.6	-256.9	-269.8	-308.3
	LW↑	+302.6	+294.2	+324.5	+317.7	+321.9	+346.4
	LH	+46.6	+38.8	+69.8	+59.0	+48.2	+78.1
	SH	-24.7	-28.5	+44.3	-34.4	-24.8	+39.1
	S	+0.1	+6.3	-85.2	-0.5	+4.1	-87.3
JJA	T _s	303.1	299.1	292.9	305.9	305.7	296.3
	SW	-273.3	-244.5	-229.3	-268.4	-262.1	-232.8
	LW↓	-374.9	-365.2	-368.6	-398.1	-398.4	-392.3
	LW↑	+478.7	+453.6	+417.5	+496.3	+495.2	+437.0
	LH	+53.7	+86.8	+122.0	+67.2	+77.9	+137.7
	SH	+115.6	+68.4	+40.1	+102.8	+86.7	+34.9
	S	+0.2	+0.9	+18.3	+0.3	+0.7	+15.5

Table 5d
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons
Region CSE

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _a		288.7	286.8	287.3	291.9	291.4	291.9
	SW	-117	-206.5	-157.7	-155.5	-200.7	-169.2	-163.6
	LW↓		-320.5	-334.6	-343.1	-346.3	-358.4	-367.8
	LW↑	[49]	+393.8	+383.6	+386.1	+411.7	+408.7	+411.8
	LH	50	+113.2	+96.3	+104.4	+127.8	+103.3	+111.1
	SH	20	+19.5	+9.3	+28.0	+7.4	+13.6	+28.6
	S		+0.5	+3.0	-19.8	+0.1	+2.1	-20.2
DJF	T _a		273.8	271.8	275.4	277.5	276.8	280.5
	SW	-77	-143.4	-110.1	-93.9	-135.4	-122.1	-98.2
	LW↓	[20]	-242.7	-256.7	-294.7	-272.0	-279.5	-308.9
	LW↑		+318.5	+309.2	+326.3	+336.4	+332.7	+350.8
	LH	20	+80.1	+54.4	+78.4	+100.2	+65.8	+82.8
	SH	18	+18.1	-3.5	+50.2	+0.8	-2.3	+52.4
	S		-30.5	+6.8	-73.1	-29.9	+5.3	-79.0
JJA	T _s		299.5	298.4	297.4	302.4	301.8	302.4
	SW	-160	-255.8	-192.0	-215.2	-252.1	-204.4	-229.8
	LW↓	[79]	-390.6	-406.2	-398.3	-412.9	-425.9	-427.5
	LW↑		+456.0	+449.8	+443.2	+473.9	+470.1	+474.2
	LH	79	+151.9	+133.0	+136.9	+164.1	+139.3	+147.6
	SH	38	+4.4	+14.3	+7.2	-1.0	+20.0	+9.1
	S		+34.2	+1.1	+26.2	+26.5	+0.9	+28.0

Table 5e
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region CSW

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _s		280.8	275.1	276.6	283.4	279.2	281.3
	SW	-150	-206.8	-130.7	-161.1	-205.3	-138.0	-164.5
	LW↓		-263.4	-259.7	-275.6	-280.0	-280.2	-299.2
	LW↑	[78]	+352.7	+324.7	+332.1	+366.0	+344.6	+355.1
	LH	16	+87.0	+55.3	+48.6	+96.0	+62.4	+50.9
	SH	49	+30.6	+3.6	+52.4	+23.5	+5.5	+55.1
	S		-0.2	+6.9	+3.6	-0.2	+5.5	+2.8
DJF	T _s		268.0	264.6	267.8	270.3	267.9	272.8
	SW	-106	-130.0	-68.3	-83.8	-128.2	-71.0	-89.5
	LW↓		-210.9	-221.5	-252.0	-222.3	-236.0	-274.0
	LW↑	[69]	+292.6	+277.9	+291.8	+302.6	+300.0	+314.1
	LH	6	+45.6	+28.2	+31.0	+53.6	+31.5	+35.3
	SH	26	+2.9	-22.1	+13.4	-5.5	-23.9	+14.1
	S		-0.2	+5.9	-0.5	-0.2	+7.5	+0.1
JJA	T _s		293.5	285.9	285.7	295.7	290.1	290.2
	SW	-187	-274.3	-187.8	-237.8	-275.1	-195.5	-235.3
	LW↓		-322.4	-312.6	-302.4	-335.6	-331.1	-330.2
	LW↑	[81]	+420.7	+378.7	+377.5	+433.6	+401.3	+402.4
	LH	30	+120.1	+89.6	+66.1	+118.6	+93.6	+56.3
	SH	45	+55.8	+31.1	+90.2	+58.8	+30.9	+101.4
	S		+0.1	+1.0	+6.3	-0.4	+0.7	+5.5

Table 5f
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region CNW

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _s		278.4	276.0	277.7	281.2	280.7	281.7
	SW	-128	-177.0	-132.4	-119.3	-175.6	-138.4	-132.7
	LW↓		-261.1	-261.0	-288.8	-280.1	-286.6	-311.5
	LW↑	[67]	+340.8	+328.9	+337.2	+354.6	+351.9	+357.0
	LH	15	+77.7	+62.6	+60.1	+87.9	+70.0	+54.1
	SH	50	+19.8	-2.1	+26.5	+13.5	-0.2	+29.6
	S		-0.1	+4.0	+3.9	-0.3	+3.3	+3.5
DJF	T _s		262.9	261.2	265.4	265.0	265.8	269.7
	SW	-69	-102.0	-64.1	-58.7	-97.4	-68.9	-58.8
	LW↓		-183.0	-191.1	-246.5	-198.5	-214.5	-268.0
	LW↑	[47]	+270.9	+264.1	+281.2	+279.6	+283.0	+299.9
	LH	4	+23.0	+23.6	+16.8	+29.9	+29.0	+20.4
	SH	21	-8.8	-36.8	+10.0	-13.4	-33.8	+7.5
	S		-0.1	+4.4	-2.8	-0.3	+5.3	-0.9
JJA	T _s		294.1	290.6	290.1	295.8	295.2	292.8
	SW	-188	-247.5	-197.3	-206.8	-247.3	-203.6	-207.3
	LW↓	[80]	-348.1	-337.4	-341.6	-362.2	-364.2	-365.4
	LW↑		+424.3	+404.4	+401.4	+434.0	+430.8	+422.4
	LH	28	+134.5	+104.3	+87.7	+147.4	+109.0	+92.3
	SH	52	+36.9	+24.9	+48.7	+28.4	+26.9	+48.4
	S		0.0	+1.0	+10.6	-0.4	+1.1	+0.6

Table 5g
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons
Region CNE

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _s		275.8	271.2	275.1	279.9	277.1	278.9
	SW	-112	-167.4	-117.6	-119.2	-163.4	-126.8	-120.5
	LW↓		-247.6	-249.4	-284.7	-273.3	-276.9	-305.1
	LW↑	[56]	+327.9	+306.6	+324.5	+348.0	+334.2	+343.0
	Latent	29	+72.2	+62.7	+60.1	+80.3	+71.2	+67.2
	SH	37	+14.9	-5.7	+26.5	+8.6	-4.9	+22.7
	S		-0.1	+3.4	-7.2	-0.2	+3.1	-7.3
DJF	T _s		252.1	248.8	258.6	256.0	255.1	263.1
	SW	-77	-80.3	-58.3	-52.1	-76.2	-59.0	-49.0
	LW↓	[49]	-144.4	-151.2	-218.6	-164.0	-174.9	-239.5
	LW↑		+228.9	+217.4	+253.5	+243.4	+240.3	+271.6
	Latent	4	+17.9	+24.2	+23.6	+22.2	+25.6	+26.2
	SH	7	-21.8	-35.1	+30.2	-24.1	-35.4	+24.4
	S		-0.3	+3.1	-36.6	-1.2	+3.4	-33.7
JJA	T _s		298.3	293.5	290.2	300.4	298.4	293.2
	SW	-160	-250.3	-167.5	-188.3	-248.4	-182.7	-193.4
	LW↓		-372.6	-379.4	-361.2	-384.0	-407.2	-379.4
	LW↑	[57]	+449.0	+420.7	+402.4	+461.4	+449.2	+418.7
	Latent	66	+134.8	+110.2	+119.6	+121.5	+125.7	+132.9
	SH	35	+38.8	+15.2	+14.5	+49.1	+13.9	+9.2
	S		+0.2	+0.7	+12.9	+0.3	+1.0	+12.0

Table 5h
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region NP

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _a	286.9	286.7	285.4	288.1	290.6	289.8	291.5
	SW	-118.0	-174.2	-144.3	-144.8	-171.0	-141.5	-148.0
	LW↓	-335.0	-316.5	-325.3	-348.8	-342.6	-354.0	-369.4
	LW↑	+384.2	+383.0	+376.2	+390.8	+404.4	+399.7	+409.5
	LH	+76.7	+91.7	+68.3	+98.7	+101.8	+75.5	+106.8
	SH	+8.2	+15.5	+25.2	+20.1	+8.5	+20.0	+16.3
	S		+0.4	0.0	-16.0	-1.1	+0.5	-15.2
DJF	T _a	283.7	283.6	282.0	285.5	287.6	286.4	288.7
	SW	-70.8	-101.8	-70.0	-69.5	-98.1	-70.2	-72.8
	LW↓	-310.2	-283.1	-308.1	-333.7	-309.5	-334.7	-351.1
	LW↑	+367.3	+367.0	+358.4	+376.5	+387.9	+381.5	+393.9
	LH	+92.1	+132.9	+109.2	+116.2	+149.1	+122.6	+135.6
	SH	+18.5	+40.0	+53.9	+30.5	+29.5	+45.0	+26.2
	S		-155.0	-143.4	-120.0	-158.9	-144.1	-131.8
JJA	T _s	290.2	290.0	289.0	291.6	300.8	293.0	295.2
	SW	-152.6	-233.0	-210.8	-220.2	-229.1	-206.6	-221.8
	LW↓	-364.9	-397.2	-350.0	-369.9	-349.2	-378.8	-393.7
	LW↑	+402.1	+445.3	+395.3	+410.0	+463.8	+418.1	+430.5
	LH	+47.9	+54.4	+27.0	+79.1	+66.7	+34.2	+79.6
	SH	-1.3	-8.0	-0.8	+10.0	-7.6	-1.1	+7.3
	S		+138.4	+139.2	+91.0	+124.0	+134.2	+98.0

Table 5i
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons
Region NA

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _s	290.0	288.3	287.4	290.6	291.2	291.4	294.1
	SW	-143.5	-182.6	-154.9	-159.8	-177.8	-150.6	-163.7
	LW↓	-345.3	-319.1	-330.1	-352.6	-340.2	-359.5	-376.1
	LW↑	+401.0	+391.7	+386.5	+404.6	+407.5	+408.8	+424.4
	LH	+99.9	+96.3	+75.2	+126.6	+100.7	+82.2	+136.6
	SH	11.8	+13.3	+23.6	+19.9	+8.6	+19.5	+15.5
	S		+0.4	-0.3	-38.6	+1.2	-0.4	-36.7
DJF	T _s	287.1	286.0	284.8	288.5	288.8	288.8	292.1
	SW	-76.2	-96.7	-74.0	-78.4	-94.3	-72.7	-80.7
	LW↓	-325.8	-302.1	-320.6	-339.9	-321.6	-346.2	-363.7
	LW↑	+385.2	+379.1	+373.1	+392.7	+394.6	+394.4	+412.9
	LH	+121.6	+111.1	+101.9	+151.6	+123.6	+113.8	+158.1
	SH	+24.6	+23.1	+41.1	+28.7	+18.3	+35.8	+21.6
	S		-114.5	-121.6	-154.7	-120.6	-125.0	-148.1
JJA	T _s	292.7	291.1	290.0	293.3	293.9	293.9	296.6
	SW	-201.8	-256.0	-243.2	-238.4	-245.1	-235.0	-241.7
	LW↓	-366.1	-343.3	-340.2	-369.4	-365.2	-373.3	-385.5
	LW↑	+416.2	+407.2	+401.1	+419.5	+423.0	+423.0	+438.8
	LH	+66.7	+73.4	+48.0	+88.4	-75.7	+50.9	+103.7
	SH	+0.4	-0.6	+7.4	+10.1	-2.2	+3.9	+9.2
	S		+119.3	+126.8	+89.8	+113.9	+130.6	+83.8

Table 5j
Regional Mean Surface Energy Balance in the Three GCMs
for Control and Doubled-CO₂ Climates,
for Average and Extreme Seasons

Region WEP

[units: T_s-K, others-W/m²]

		a) current (control)				b) 2 CO ₂		
		OBS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	T _s	301.7	300.8	300.5	300.3	302.6	303.0	303.8
	SW	-170.6	-219.0	-203.0	-263.2	-219.2	-204.4	-266.7
	LW↓	-428.8	-409.0	-410.9	-396.9	-422.5	-432.6	-428.8
	LW↑	+469.8	+464.1	+462.6	+460.9	+475.1	+477.8	+483.1
	LH	+108.7	+155.8	+132.6	+134.0	+161.0	+142.7	+149.2
	SH	+3.5	+7.4	+18.0	+12.1	+5.2	+15.9	+9.3
	S		+0.7	+0.8	+53.1	+0.4	+0.7	+54.0
DJF	T _s	301.4	301.0	300.6	299.8	302.7	303.1	303.5
	SW	-165.6	-216.6	-204.0	-270.0	-219.6	-205.6	-271.3
	LW↓	-428.7	-413.3	-411.1	-387.9	-424.6	-433.6	-424.3
	LW↑	+467.9	+465.4	+462.7	+457.9	+475.9	+478.6	+481.2
	LH	+119.3	+156.7	+141.2	+116.7	+170.3	+152.4	+138.8
	SH	+5.0	+6.6	+19.1	+13.0	+5.0	+16.6	+9.8
	S		+1.2	-7.9	+70.3	-7.0	-8.5	+65.8
JJA	T _s	301.5	300.5	300.4	300.7	302.4	302.8	304.0
	SW	-166.1	-210.1	-188.9	-247.7	-206.4	-192.7	-283.6
	LW↓	-427.6	-403.7	-411.5	-405.0	-420.9	-432.8	-430.6
	LW↑	+468.5	+462.2	+461.9	+463.5	+474.2	+476.9	+484.0
	LH	+105.5	+160.2	+127.7	+152.1	+159.9	+140.8	+159.1
	SH	+5.0	+9.0	+17.6	+10.8	+5.5	+15.9	+9.0
	S		-17.6	-6.7	+26.2	-12.2	-8.1	+32.2

Table 6a
Regional Mean Components of the Hydrological Cycle
Region USW

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	2.71	2.86	2.55	2.97	2.98	2.90	+0.26	+0.12	+0.35
	Moisture	5.98	7.11	7.85	7.33	9.36	9.43	+1.35	+2.25	+1.58
	Low Cloud	0.15	0.44	0.38	0.15	0.45	0.35	0.00	+0.01	-0.03
	High Cloud	0.15	0.17	0.34	0.14	0.16	0.35	-0.01	-0.01	+0.01
	Lapse Rate	6.91	6.33		6.76	6.31		-0.15	-0.02	
DJF	Precipitation	4.47	4.73	3.06	4.60	4.97	3.44	+0.13	+0.24	+0.38
	Moisture	3.72	3.61	4.71	4.49	4.99	6.03	+0.77	+1.38	+1.32
	Low Cloud	0.28	0.40	0.54	0.25	0.41	0.52	-0.03	+0.02	-0.02
	High Cloud	0.22	0.13	0.39	0.19	0.13	0.39	-0.03	-0.00	+0.01
	Lapse Rate	5.65	5.34		5.70	5.27		+0.05	-0.07	
JJA	Precipitation	0.96	0.92	2.22	1.01	0.98	2.67	+0.05	+0.06	+0.45
	Moisture	8.54	11.80	11.21	10.42	15.16	13.75	+1.88	+3.36	+2.54
	Low Cloud	0.06	0.48	0.25	0.06	0.50	0.26	+0.00	+0.02	+0.01
	High Cloud	0.08	0.19	0.29	0.09	0.18	0.32	+0.01	-0.01	+0.03
	Lapse Rate	8.22	7.58		8.08	7.61		-0.14	+0.03	

Table 6b
Regional Mean Components of the Hydrological Cycle
Region USC

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	1.80	2.68	2.39	2.13	2.62	2.41	+0.33	-0.06	+0.02
	Moisture	6.61	8.74	7.82	8.22	10.91	10.29	+1.61	+2.17	+2.47
	Low Cloud	0.11	0.42	0.37	0.12	0.44	0.34	+0.01	+0.02	-0.03
	High Cloud	0.13	0.15	0.21	0.12	0.15	0.21	-0.01	0.00	0.00
	Lapse Rate	6.98	6.10		6.90	6.24		-0.08	+1.14	
DJF	Precipitation	2.04	2.09	1.99	2.48	2.52	2.03	+0.44	+0.43	+0.04
	Moisture	2.97	3.07	4.03	3.82	4.54	5.26	+0.85	+1.47	+1.23
	Low Cloud	0.18	0.35	0.50	0.20	0.37	0.46	+0.02	+0.02	-0.04
	High Cloud	0.13	0.10	0.23	0.13	0.10	0.21	+0.00	+0.00	-0.02
	Lapse Rate	4.99	4.37		5.15	4.48		+0.16	+1.11	
JJA	Precipitation	1.37	2.74	3.05	1.50	1.86	3.10	+0.13	-0.88	+0.05
	Moisture	10.64	15.14	12.53	12.66	17.35	16.56	+2.02	+2.21	+4.03
	Low Cloud	0.06	0.48	0.29	0.06	0.50	0.27	+0.00	+0.02	-0.02
	High Cloud	0.10	0.20	0.18	0.11	0.18	0.21	+0.01	-0.02	+0.03
	Lapse Rate	8.75	7.73		8.69	8.04		-0.06	+0.31	

Table 6c
Regional Mean Components of the Hydrological Cycle
Region USE

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	2.20	3.11	3.13	3.03	3.20	3.21	+0.83	+0.09	+0.08
	Moisture	6.90	8.24	7.76	8.74	10.61	9.64	+1.84	+2.37	+1.88
	Low Cloud	0.12	0.44	0.41	0.15	0.44	0.37	+0.03	0.00	-0.04
	High Cloud	0.14	0.15	0.20	0.14	0.15	0.20	0.00	0.00	0.00
	Lapse Rate	6.55	5.56		6.53	5.77		-0.02	+0.21	
DJF	Precipitation	2.53	3.42	2.85	3.12	3.95	3.01	+0.59	+0.53	+0.16
	Moisture	2.80	2.88	4.51	3.49	4.35	5.60	+0.69	+1.47	+1.09
	Low Cloud	0.21	0.34	0.54	0.22	0.31	0.52	+0.01	-0.03	-0.02
	High Cloud	0.14	0.13	0.29	0.15	0.12	0.33	+0.01	-0.01	+0.04
	Lapse Rate	4.59	3.74		4.59	4.02		0.00	+0.28	
JJA	Precipitation	1.83	2.69	3.69	2.06	2.17	4.21	+0.23	-0.52	+0.52
	Moisture	11.58	14.35	12.58	13.62	17.21	15.04	+2.04	+2.86	+2.46
	Low Cloud	0.03	0.51	0.31	0.04	0.55	0.30	+0.01	+0.04	-0.01
	High Cloud	0.14	0.21	0.19	0.14	0.19	0.23	+0.00	-0.02	+0.04
	Lapse Rate	8.31	7.2		8.29	7.57		-0.02	+0.37	

Table 6d
Regional Mean Components of the Hydrological Cycle
Region CSE

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	5.55	5.83	4.59	6.17	5.98	4.95	+0.62	+0.15	+0.36
	Moisture	10.37	11.75	11.50	12.75	14.36	14.67	+2.38	+2.61	+3.17
	Low Cloud	0.14	0.20	0.43	0.15	0.21	0.39	+0.01	+0.01	-0.04
	High Cloud	0.12	0.10	0.27	0.14	0.09	0.24	+0.02	-0.01	-0.03
	Lapse Rate	5.54	4.54		5.45	4.65		+0.09	-0.11	
DJF	Precipitation	3.08	3.89	2.03	4.66	3.89	1.83	+1.58	+0.00	-0.20
	Moisture	3.61	3.75	4.70	5.00	5.29	6.98	+1.39	+1.54	+2.28
	Low Cloud	0.14	0.11	0.50	0.18	0.10	0.45	+0.04	-0.01	-0.05
	High Cloud	0.10	0.08	0.14	0.14	0.07	0.16	+0.04	-0.01	+0.02
	Lapse Rate	4.01	2.79		3.99	2.97		-0.02	+0.15	
JJA	Precipitation	8.77	8.26	7.20	8.75	8.37	7.46	-0.02	+0.11	+0.26
	Moisture	16.64	19.57	18.99	20.36	22.37	23.95	+3.72	+2.80	+4.96
	Low Cloud	0.14	0.31	0.36	0.15	0.34	0.32	+0.01	+0.03	-0.04
	High Cloud	0.14	0.11	0.46	0.14	0.10	0.27	+0.00	-0.01	-0.19
	Lapse Rate	6.36	5.58		6.23	5.53		-0.13	-0.05	

Table 6e
Regional Mean Components of the Hydrological Cycle
Region CSW

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	4.28	5.44	2.34	4.87	6.46	2.47	+0.59	+1.02	+0.03
	Moisture	7.33	6.72	6.78	8.50	8.58	8.80	+1.17	+1.86	+2.02
	Low Cloud	0.13	0.33	0.36	0.13	0.33	0.32	0.00	0.00	-0.04
	High Cloud	0.14	0.12	0.31	0.16	0.12	0.34	+0.02	0.00	+0.03
	Lapse Rate	6.66	6.44		6.55	6.28		-0.11	-0.16	
DJF	Precipitation	2.98	4.37	2.75	3.94	4.88	3.03	+0.96	+0.51	+0.28
	Moisture	2.84	2.68	4.43	3.43	3.50	5.72	+0.59	+0.82	+1.29
	Low Cloud	0.17	0.31	0.53	0.17	0.30	0.49	+0.00	-0.01	-0.04
	High Cloud	0.14	0.10	0.29	0.16	0.10	0.34	+0.02	+0.00	+0.05
	Lapse Rate	5.83	5.92		5.6	5.67		-0.23	-0.25	
JJA	Precipitation	5.37	7.71	1.88	5.44	10.09	1.76	+0.07	+2.38	-0.12
	Moisture	12.45	12.17	9.74	13.64	14.45	12.50	+1.19	+2.28	+2.76
	Low Cloud	0.11	0.34	0.22	0.09	0.33	0.19	-0.02	-0.01	-0.03
	High Cloud	0.12	0.15	0.26	0.10	0.14	0.29	-0.02	-0.01	+0.03
	Lapse Rate	7.51	6.84		7.24	6.55		-0.27	-0.29	

Table 6f
Regional Mean Components of the Hydrological Cycle
Region CNW

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	3.58	4.01	2.80	3.87	4.48	3.13	+0.29	+0.47	+0.33
	Moisture	6.48	6.75	6.52	7.81	8.80	8.51	+1.33	+2.05	+1.99
	Low Cloud	0.16	0.39	0.43	0.15	0.42	0.42	-0.01	-0.03	-0.01
	High Cloud	0.15	0.16	0.31	0.15	0.15	0.33	0.00	-0.01	+0.02
	Lapse Rate	6.27	5.53		6.11	5.45		-0.16	-0.08	
DJF	Precipitation	1.22	1.40	1.66	1.75	1.66	2.08	+0.53	+0.26	+0.42
	Moisture	1.28	1.43	2.59	1.66	2.23	3.86	+0.38	+0.80	+1.27
	Low Cloud	0.17	0.31	0.57	0.17	0.37	0.56	+0.00	+0.06	-0.01
	High Cloud	0.12	0.13	0.29	0.13	0.12	0.33	+0.01	-0.01	+0.04
	Lapse Rate	5.68	4.68		5.36	4.65		-0.32	-0.03	
JJA	Precipitation	5.53	6.31	4.08	5.90	6.50	3.83	+0.37	+0.19	-0.25
	Moisture	13.09	13.47	11.94	14.99	16.70	14.51	+1.90	+3.23	+2.57
	Low Cloud	0.13	0.48	0.31	0.12	0.49	0.29	-0.01	+0.01	-0.02
	High Cloud	0.3	0.19	0.31	0.12	0.18	0.31	-0.01	-0.01	+0.00
	Lapse Rate	6.84	6.18		6.54	6.07		-0.30	-0.11	

Table 6g
Regional Mean Components of the Hydrological Cycle
Region CNE

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	3.33	3.26	2.76	3.85	3.43	3.06	+0.52	+0.17	+0.30
	Moisture	6.04	6.30	5.92	7.32	8.55	7.35	+1.28	+2.25	+1.43
	Low Cloud	0.15	0.45	0.48	0.17	0.47	0.47	+0.02	+0.02	-0.01
	High Cloud	0.16	0.18	0.24	0.14	0.19	0.24	-0.02	+0.01	0.00
	Lapse Rate	5.29	4.22		5.25	4.40		-0.04	+0.18	
DJF	Precipitation	1.00	1.14	1.26	1.36	1.25	1.32	+0.36	+0.11	+0.06
	Moisture	0.39	0.13	1.52	0.56	0.46	2.16	+0.17	+0.33	+0.64
	Low Cloud	0.13	0.36	0.57	0.15	0.39	0.55	+0.02	+0.03	-0.02
	High Cloud	0.14	0.16	0.17	0.12	0.16	0.17	-0.02	+0.00	+0.00
	Lapse Rate	3.40	1.96		3.44	2.42		+0.04	+0.46	
JJA	Precipitation	5.09	5.64	4.90	5.11	5.50	5.49	+0.02	-0.14	+0.59
	Moisture	14.35	15.95	11.96	15.25	20.38	14.89	+0.90	+4.43	+2.93
	Low Cloud	0.13	0.50	0.40	0.14	0.53	0.39	+0.01	+0.03	-0.01
	High Cloud	0.15	0.20	0.31	0.12	0.21	0.33	-0.03	+0.01	+0.02
	Lapse Rate	6.60	5.90		6.38	5.79		-0.22	-0.11	

Table 6h
Regional Mean Components of the Hydrological Cycle
Region NP

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	3.24	2.72	3.74	3.63	2.92	4.10	+0.39	+0.20	+0.36
	Moisture	7.88	8.68	9.12	10.05	11.46	10.37	+2.17	+2.78	+1.25
	Low Cloud	0.25	0.42	0.52	0.25	0.44	0.49	0.00	+0.02	-0.03
	High Cloud	0.15	0.17	0.27	0.14	0.16	0.26	-0.01	-0.01	-0.01
	Lapse Rate	5.98	5.29		6.04	5.23		+0.06	-0.06	
DJF	Precipitation	4.31	4.09	4.36	4.83	4.35	4.72	+0.52	+0.26	+0.36
	Moisture	6.58	7.12	7.02	8.44	9.43	8.39	+1.86	+2.31	+1.37
	Low Cloud	0.15	0.38	0.57	0.14	0.40	0.53	-0.01	+0.02	-0.04
	High Cloud	0.17	0.15	0.25	0.16	0.15	0.21	-0.01	+0.00	-0.04
	Lapse Rate	6.14	5.65		6.17	5.56		+0.03	-0.09	
JJA	Precipitation	2.54	1.84	3.18	2.84	1.97	3.90	+0.30	+0.13	+0.72
	Moisture	9.18	10.32	11.48	11.54	13.42	13.09	+2.36	+3.10	+1.61
	Low Cloud	0.36	0.45	0.48	0.35	0.50	0.45	-0.01	+0.05	-0.03
	High Cloud	0.14	0.19	0.32	0.12	0.17	0.34	-0.02	-0.02	+0.02
	Lapse Rate	6.05	5.09		6.05	5.03		0.00	-0.06	

Table 6i
Regional Mean Components of the Hydrological Cycle
Region NA

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	2.63	2.32	3.52	2.74	2.45	3.44	+0.11	+0.13	-0.08
	Moisture	8.49	9.63	9.60	10.19	12.53	12.15	+1.70	+2.90	+2.55
	Low Cloud	0.22	0.37	0.43	0.23	0.39	0.40	+0.01	+0.02	-0.03
	High Cloud	0.09	0.17	0.24	0.09	0.16	0.24	0.00	-0.01	0.00
	Lapse Rate	6.40	5.68		6.29	5.61		-0.11	-0.07	
DJF	Precipitation	3.32	3.52	4.31	3.74	3.69	4.14	+0.42	+0.17	-0.17
	Moisture	7.36	8.37	8.53	8.91	10.85	11.14	+1.55	+2.48	+2.61
	Low Cloud	0.20	0.29	0.47	0.18	0.30	0.43	-0.02	+0.01	-0.04
	High Cloud	0.14	0.16	0.25	0.14	0.15	0.26	+0.00	-0.01	+0.01
	Lapse Rate	6.59	5.83		6.43	5.81		-0.16	-0.02	
JJA	Precipitation	1.84	1.00	2.77	1.88	1.09	2.82	+0.04	+0.09	+0.05
	Moisture	9.76	10.84	11.05	11.60	14.24	13.74	+1.84	+3.40	+2.69
	Low Cloud	0.27	0.45	0.39	0.30	0.48	0.37	+0.03	+0.03	-0.02
	High Cloud	0.04	0.19	0.25	0.04	0.17	0.25	+0.00	-0.02	+0.00
	Lapse Rate	6.21	5.33		6.18	5.26		-0.03	-0.07	

Table 6j
Regional Mean Components of the Hydrological Cycle
Region WEP

[units: same as Table 3]

		a) current (control)			b) 2 CO ₂			c) 2 CO ₂ control		
		NCAR	GFDL	GISS	NCAR	GFDL	GISS	NCAR	GFDL	GISS
ANNUAL	Precipitation	7.79	4.91	6.21	8.04	5.39	6.63	+0.25	+0.48	+0.42
	Moisture	18.59	21.29	15.87	20.72	24.55	19.70	+2.13	+3.26	+3.83
	Low Cloud	0.27	0.24	0.10	0.26	0.23	0.08	-0.01	-0.01	-0.02
	High Cloud	0.14	0.13	0.52	0.13	0.11	0.43	-0.01	-0.02	-0.09
	Lapse Rate	5.99	5.93		5.81	5.68		-0.18	-0.25	
DJF	Precipitation	8.40	4.77	4.84	9.04	5.71	5.93	+0.64	+0.94	+1.09
	Moisture	18.74	21.33	15.35	20.83	24.66	19.22	+2.09	+3.33	+3.87
	Low Cloud	0.31	0.24	0.09	0.27	0.22	0.08	-0.04	-0.02	-0.01
	High Cloud	0.14	0.13	0.45	0.15	0.12	0.47	+0.01	-0.01	+0.02
	Lapse Rate	6.04	5.95		5.87	5.69		-0.21	-0.26	
JJA	Precipitation	7.27	5.30	7.47	7.75	5.25	7.18	+0.48	-0.05	-0.29
	Moisture	18.33	21.23	16.44	20.58	24.42	20.14	+2.25	+3.19	+3.70
	Low Cloud	0.25	0.26	0.11	0.26	0.25	0.09	+0.01	-0.01	-0.02
	High Cloud	0.12	0.14	0.58	0.10	0.11	0.39	-0.02	-0.03	-0.19
	Lapse Rate	6.01	5.97		5.86	5.71		-0.15	-0.26	

Table 7a
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region USW

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+2.5	+4.9	+4.8
	ΔSW	-0.8	+1.4	+0.2
	$\Delta LW\downarrow$	+3.7	+5.0	+5.3
	ΔLH	-1.1	-0.7	-1.5
	ΔSH	+0.7	-1.1	+0.8
	ΔS	-0.0	+0.4	+0.2
DJF	ΔT_a	+2.7	+4.4	+5.1
	ΔSW	-0.5	+1.8	+0.0
	$\Delta LW\downarrow$	+3.4	+4.7	+5.6
	ΔLH	-0.8	-1.5	-1.3
	ΔSH	+0.7	-1.2	+1.0
	ΔS	+0.0	+0.7	-0.0
JJA	ΔT_a	3.1	+5.6	+3.9
	ΔSW	-1.1	+0.5	+0.0
	$\Delta LW\downarrow$	+4.4	+6.0	+5.0
	ΔLH	+0.3	+0.6	-2.6
	ΔSH	-0.5	-1.4	+1.6
	ΔS	-0.1	-0.0	+0.0

Table 7b
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region USC

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+2.4	+5.5	+4.9
	ΔSW	-0.9	+1.6	+0.5
	$\Delta LW\downarrow$	+3.9	+5.4	+5.0
	ΔLH	-1.9	+0.0	-0.4
	ΔSH	+1.3	-1.5	-0.3
	ΔS	+0.0	+0.2	+0.0
DJF	ΔT_a	+3.3	+5.2	+5.5
	ΔSW	-0.9	+1.1	+0.2
	$\Delta LW\downarrow$	+4.7	+6.5	+5.5
	ΔLH	-2.2	-1.6	-0.4
	ΔSH	+1.7	-1.1	+0.1
	ΔS	+0.1	+0.4	+0.2
JJA	ΔT_a	+3.3	+6.4	+3.8
	ΔSW	-1.0	+3.2	+0.5
	$\Delta LW\downarrow$	+4.4	+4.5	+4.2
	ΔLH	-0.0	+2.7	-0.5
	ΔSH	+0.0	-3.8	-0.5
	ΔS	-0.0	+0.0	+0.1

Table 7c
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region USE

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+3.0	+5.9	+4.1
	ΔSW	-1.2	+1.5	+0.8
	$\Delta LW\downarrow$	+4.8	+5.9	+4.3
	ΔLH	+3.9	-0.3	-1.4
	ΔSH	+3.3	-1.2	+0.3
	ΔS	+0.0	+0.2	+0.2
DJF	ΔT_a	+3.3	+6.1	+4.5
	ΔSW	-0.9	+1.6	+0.1
	$\Delta LW\downarrow$	+4.8	+7.2	+4.8
	ΔLH	-2.8	-2.1	-1.8
	ΔSH	+2.2	-0.8	+1.1
	ΔS	+0.1	+0.5	+0.5
JJA	ΔT_a	+2.8	+6.6	+3.4
	ΔSW	-0.8	+2.8	+0.6
	$\Delta LW\downarrow$	+3.7	+5.3	+4.2
	ΔLH	-2.1	+1.4	-2.8
	ΔSH	+2.0	-2.9	+0.9
	ΔS	-0.0	+0.0	+0.5

Table 7d
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region CSE

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+3.2	+4.6	+4.6
	ΔSW	-1.1	+2.1	+1.5
	$\Delta LW\downarrow$	+4.7	+4.4	+4.6
	ΔLH	-2.7	-1.3	-1.2
	ΔSH	+2.2	-0.8	-0.1
	ΔS	+0.1	+0.2	+0.1
DJF	ΔT_a	+3.7	+5.0	+5.1
	ΔSW	-1.7	+2.6	+0.9
	$\Delta LW\downarrow$	+6.3	+5.0	+3.0
	ΔLH	-4.3	-2.5	-0.9
	ΔSH	+3.7	-0.3	-0.5
	ΔS	-0.1	+0.3	+1.2
JJA	ΔT_a	+2.9	+3.4	+5.0
	ΔSW	-0.6	+2.1	+2.4
	$\Delta LW\downarrow$	+3.7	+3.3	+4.9
	ΔLH	-2.0	-1.0	-1.8
	ΔSH	+0.9	-0.9	-0.3
	ΔS	+1.0	+0.0	-0.0

Table 7e
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region CSW

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+2.6	+4.1	+4.7
	ΔSW	-0.3	+1.5	+0.7
	$\Delta LW\downarrow$	+3.3	+4.3	+4.9
	ΔLH	-1.8	-1.5	-0.5
	ΔSH	+1.4	-0.4	-0.6
	ΔS	+0.0	+0.3	+0.2
DJF	ΔT_a	+2.3	+3.3	+5.0
	ΔSW	-0.4	+0.6	+1.3
	$\Delta LW\downarrow$	+2.6	+3.5	+5.1
	ΔLH	-1.8	-0.8	-1.0
	ΔSH	+1.9	+0.4	-0.2
	ΔS	+0.0	-0.4	-0.1
JJA	ΔT_a	+2.2	+4.2	+4.5
	ΔSW	+0.1	+1.5	-0.5
	$\Delta LW\downarrow$	+2.3	+3.5	+5.3
	ΔLH	+0.3	-0.8	+1.9
	ΔSH	-0.5	+0.0	-2.1
	ΔS	+0.1	+0.1	+0.2

Table 7f
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region CNW

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+2.8	+4.7	+4.0
	ΔSW	-0.3	+1.3	+2.8
	$\Delta LW\downarrow$	+3.9	+5.4	+4.7
	ΔLH	-2.1	-1.6	+1.2
	ΔSH	+1.3	-0.4	-0.6
	ΔS	+0.0	+0.1	+0.1
DJF	ΔT_a	+2.1	+4.6	+4.3
	ΔSW	-1.1	+1.2	+0.0
	$\Delta LW\downarrow$	+3.8	+5.8	+5.1
	ΔLH	-1.7	-1.3	-0.8
	ΔSH	+1.1	-0.7	+0.6
	ΔS	+0.0	-0.2	-0.5
JJA	ΔT_a	+1.7	+4.6	+3.7
	ΔSW	-0.0	+1.1	+0.1
	$\Delta LW\downarrow$	+2.4	+4.8	+4.3
	ΔLH	-2.2	-0.8	-0.8
	ΔSH	+1.5	-0.4	+0.1
	ΔS	+0.1	-0.0	+0.2

Table 7g
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region CNE

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+4.1	+5.9	+3.8
	ΔSW	-0.8	+2.0	+0.3
	$\Delta LW\downarrow$	+5.4	+6.1	+4.3
	ΔLH	-1.7	-1.9	-1.5
	ΔSH	+1.3	-0.2	+0.8
	ΔS	+0.0	+0.1	+0.0
DJF	ΔT_s	+3.9	+6.3	+4.5
	ΔSW	-1.1	+0.2	-0.8
	$\Delta LW\downarrow$	+5.4	+6.8	+5.3
	ΔLH	-1.2	-0.4	-0.7
	ΔSH	+0.6	+0.1	+1.5
	ΔS	+0.3	-0.1	-0.7
JJA	ΔT_a	+2.1	+4.9	+3.0
	ΔSW	-0.3	+2.7	+0.9
	$\Delta LW\downarrow$	+1.9	+4.8	+3.3
	ΔLH	+2.2	-2.7	-2.4
	ΔSH	-1.7	+0.2	+1.0
	ΔS	-0.0	-0.3	+0.2

Table 7h
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region NP

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+3.9	+4.4	+3.4
	ΔSW	-0.6	-0.5	+0.6
	$\Delta LW\downarrow$	+4.9	+5.4	+6.8
	ΔLH	-1.9	-1.4	-1.5
	ΔSH	+1.3	+1.0	+0.7
	ΔS	+0.3	-0.1	-0.1
DJF	ΔT_a	+4.0	+4.4	+3.2
	ΔSW	-0.7	+0.0	+0.6
	$\Delta LW\downarrow$	+5.1	+5.2	+3.3
	ΔLH	-3.1	-2.6	-3.7
	ΔSH	+2.0	+1.7	+0.8
	ΔS	+0.8	+0.1	+2.2
JJA	ΔT_a	+3.1	+4.0	+3.6
	ΔSW	-0.7	-0.8	+0.3
	$\Delta LW\downarrow$	-8.0	+5.3	+4.2
	ΔLH	-2.1	-1.3	-0.1
	ΔSH	-0.1	+0.1	+0.5
	ΔS	+2.6	+0.9	-1.2

Table 7i
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region NA

		2 CO ₂ - Control [units: K]		
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+2.9	+4.0	+3.5
	ΔSW	-1.0	-0.8	+0.7
	$\Delta LW\downarrow$	+4.4	+5.5	+4.2
	ΔLH	-0.9	-1.3	-1.8
	ΔSH	+1.0	+0.8	+0.8
	ΔS	-0.1	+0.0	-0.3
DJF	ΔT_a	+2.8	+4.0	+3.6
	ΔSW	-0.5	-0.2	+0.4
	$\Delta LW\downarrow$	+3.7	+4.9	+4.4
	ΔLH	-2.4	-2.3	-1.2
	ΔSH	+0.9	+1.0	+1.3
	ΔS	+1.2	+0.7	-1.2
JJA	ΔT_s	+2.8	+3.9	+3.3
	ΔSW	-1.9	-1.5	+0.6
	$\Delta LW\downarrow$	+3.9	+6.0	+4.3
	ΔLH	-0.4	-0.5	-2.7
	ΔSH	+0.3	+0.6	+0.2
	ΔS	+1.0	-0.7	+1.1

Table 7j
Regional Temperature Change Associated with Separate Terms
of the Surface Energy Balance in the Three GCMs
for Annual Average and Extreme Seasons
Region WEP

2 CO ₂ - Control [units: K]				
		NCAR	GFDL	GISS
ANNUAL	ΔT_a	+1.8	+2.5	+3.5
	ΔSW	+0.0	+0.2	+0.6
	$\Delta LW\downarrow$	+2.2	+3.5	+5.2
	ΔLH	-0.8	-1.6	-2.5
	ΔSH	+0.4	+0.3	+0.5
	ΔS	+0.0	+0.0	-0.1
DJF	ΔT_a	+1.7	+2.5	+3.7
	ΔSW	+0.5	+0.3	+0.2
	$\Delta LW\downarrow$	+1.8	+3.7	+6.0
	ΔLH	-2.2	-1.8	-3.6
	ΔSH	+0.3	+0.4	+0.5
	ΔS	+1.3	+0.1	+0.7
JJA	ΔT_a	+1.9	+2.4	+3.3
	ΔSW	-0.6	+0.6	+1.0
	$\Delta LW\downarrow$	+2.8	+3.5	+4.2
	ΔLH	+0.0	-2.1	-1.1
	ΔSH	+0.6	+0.3	+0.3
	ΔS	-0.9	+0.2	-1.0

Table 8
Point Identification Key for Scatter Plots

First letter:

N - NCAR

L - GFDL

S - GISS

Second and third letters:

Western U.S. (U1)

Southeast

List of Figures

- Figure 1(a). Areas selected for regional analysis in the GISS model.
- Figure 1(b). Areas selected for regional analysis in the NCAR and GFDL models.
- Figure 2. Annual mean near surface air temperature (K) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.
- Figure 3. Annual mean surface energy balance as a function of latitude for each model's 1 CO₂ climate. [Units: W/m²]
- Figure 4. Annual mean temperature change associated with each of the terms in (5), as a function of latitude. [Units: K]
- Figure 5. Annual mean precipitation (mm/day) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.
- Figure 6. Annual mean near surface moisture (g/kg) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.
- Figure 7. Changes in (a) high and (b) low annual mean cloud cover as a function of latitude for CO₂ doubling. [units: %]
- Figure 8. Annual mean lapse rate (K/km) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.
- Figure 9. Annual mean surface energy balance as a function of latitude for each model's 2 CO₂ climate. [Units: W/m²]
- Figure 10. Upward, downward and net longwave fluxes (W/m²) as functions of latitude in each model's annual mean 1 CO₂ climate.
- Figure 11. DJF near surface air temperature (K) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.
- Figure 12. DJF surface energy balance as a function of latitude for each model's 1 CO₂ climate. [Units: W/m²]
- Figure 13. DJF temperature change associated with each of the terms in (5), as a function of latitude. [Units: K]
- Figure 14. DJF precipitation (mm/day) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 15. DJF near surface moisture (g/kg) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 16. Changes in (a) high and (b) low DJF cloud cover as a function of latitude for CO₂ doubling. [units: %]

Figure 17. DJF lapse rate (K/km) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 18. JJA near surface air temperature (K) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 19. JJA surface energy balance as a function of latitude for each model's 1 CO₂ climate. [Units: W/m²]

Figure 20. JJA temperature change associated with each of the terms in (5), as a function of latitude. [Units: K]

Figure 21. JJA precipitation (mm/day) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 22. JJA near surface moisture (g/kg) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 23. Changes in (a) high and (b) low JJA cloud cover as a function of latitude for CO₂ doubling. [units: %]

Figure 24. JJA lapse rate (K/km) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 25(a). Contour plots for DJF surface sensible heat flux. The contour interval is 20 W m⁻².

Figure 25(b). Contour plots for DJF surface latent heat flux. The contour interval is 20 W m⁻².

Figure 25(c). Contour plots for DJF surface net shortwave radiation. The contour interval is 20 W m⁻².

Figure 25(d). Contour plots for DJF surface net longwave radiation. The contour interval is 20 W m⁻².

Figure 25(e). Contour plots for DJF surface air temperature. The contour interval for is 10 K.

Figure 26(a). Contour plots for DJF (2CO₂ - 1CO₂) surface sensible heat flux differences. Contour interval is 10 W m⁻².

Figure 26(b). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface latent heat flux differences. Contour interval is 10 W m^{-2} .

Figure 26(c). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface net shortwave radiation differences. Contour interval is 20 W m^{-2} .

Figure 26(d). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface net longwave radiation differences. Contour interval is 20 W m^{-2} .

Figure 26(e). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface air temperature differences. Contour interval is 2 K .

Figure 27. Scatter plots of ΔSW vs. ΔT_a in the regions for (a) DJF and (b) JJA.

Figure 28. Scatter plots $\Delta(\text{precipitation})$ vs. control precipitation in the regions for (a) DJF and (b) JJA. Points representing the same region from different models are connected by a solid line (U.S. regions), a dotted line (China regions) or a dashed line (ocean regions).

Figure 29. Scatter plots of $\Delta\text{LW}\downarrow$ vs. control $\Delta\text{LW}\uparrow$ in the regions for (a) DJF and (b) JJA.

Figure 30. Scatter plots of ΔLH vs. control ΔSH in the regions for (a) DJF and (b) JJA.

Figure 31. Scatter plots of mixing ratio vs. T_a for the (a) DJF

Figure 1(a). Areas selected for regional analysis in the GISS model. Name Key:
 Western U.S. (USW) Southeast China (CSE) Northwest China (CNW) North Atlantic (NA) Western Equatorial Pacific (WEP)
 Central U.S. (USC) Southwest China (CSW) Northeast China (CNE) North Pacific (NP)
 Eastern U.S. (USE)

Figure 1(b). Areas selected for regional analysis in the NCAR and GFDL models. Name Key:

Western U.S. (USW)	Southeast China (CSE)	North Atlantic (NA)	Western Equatorial Pacific (WEP)
Central U.S. (USC)	Southwest China (CSW)	North Pacific (NP)	
Eastern U.S. (USE)

Figure 2. Annual mean near surface air temperature (K) as a function of latitude in each model for the 1CO_2 climate (control), the 2CO_2 climate and the $(2\text{CO}_2 - 1\text{CO}_2)$ change.

Figure 3. Annual mean surface energy balance as a function of latitude for each model's 1 CO_2 climate. [Units: W/m^2]

Figure 4. Annual mean temperature change associated with each of the terms in (5), as a function of latitude. [Units: K]

PRECIPITATION

— NCAR
 GFDL
 ---- GISS

Figure 5. Annual mean precipitation (mm/day) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

NEAR
SURFACE
MOISTURE

— NCAR
..... GFDL
- - - GISS

Figure 6. Annual mean near surface moisture (g/kg) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 7. Changes in (a) high and (b) low annual mean cloud cover as a function of latitude for CO₂ doubling. [units: %]

Figure 8. Annual mean lapse rate (K/km) as a function of latitude in the NCAR and GFDL models for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 9. Annual mean surface energy balance as a function of latitude for each model's 2 CO_2 climate. [Units: W/m^2]

Figure 10. Upward, downward and net longwave fluxes (W/m^2) as functions of latitude in each model's annual mean 1 CO_2 climate.

Figure 11. DJF near surface air temperature (K) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 12. DJF surface energy balance as a function of latitude for each model's 1 CO₂ climate. [Units: W/m²]

Figure 13. DJF temperature change associated with each of the terms in (5), as a function of latitude. [Units: K]

PRECIPITATION

— NCAR
 GFDL
 - - - - GISS

Figure 14. DJF precipitation (mm/day) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

— NCAR
 GFDL
 --- GISS

Figure 15. DJF near surface moisture (g/kg) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 16. Changes in (a) high and (b) low DJF cloud cover as a function of latitude for CO₂ doubling. [units: %]

Figure 17. DJF lapse rate (K/km) as a function of latitude in the NCAR and GFDL models for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 18. JJA near surface air temperature (K) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 19. JJA surface energy balance as a function of latitude for each model's 1 CO_2 climate. [Units: W/m^2]

Figure 20. JJA temperature change associated with each of the terms in (5), as a function of latitude. [Units: K]

Figure 21. JJA precipitation (mm/day) as a function of latitude in each model for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

NEAR
SURFACE
MOISTURE

— NCAR
..... GFDL
--- GISS

Figure 22. JJA near surface moisture (g/kg) as a function of latitude in each model for the 1 CO_2 climate (control), the 2 CO_2 climate and the $(2\text{ CO}_2 - 1\text{ CO}_2)$ change.

Figure 23. Changes in (a) high and (b) low JJA cloud cover as a function of latitude for CO₂ doubling. [units: %]

Figure 24. JJA lapse rate (K/km) as a function of latitude in the NCAR and GFDL models for the 1 CO₂ climate (control), the 2 CO₂ climate and the (2 CO₂ - 1 CO₂) change.

Figure 25(a). Contour plots for DJF surface sensible heat flux. The contour interval is 20 W m⁻².

Figure 25(b). Contour plots for DJF surface latent heat flux. The contour interval is 20 W m^{-2} .

NCAR

GFDL

GISS

Figure 25(c). Contour plots for DJF surface net shortwave radiation. The contour interval is 20 W m⁻²

NCAR

GFDL

GISS

Figure 25(d). Contour plots for DJF surface net longwave radiation. The contour interval is 20 W m^{-2} .

NCAR

GFDL

GISS

Figure 25(e). Contour plots for DJF surface air temperature. The contour interval for is 10 K.

Figure 26(a). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface sensible heat flux differences. Contour interval is 10 W m^{-2} .

NCAR

GFDL

GISS

Figure 26(b). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface latent heat flux differences. Contour interval is 10 W m^{-2} .

Figure 26(c). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface net shortwave radiation differences. Contour interval is 20 W m^{-2} .

NCAR

GFDL

GISS

Figure 26(d). Contour plots for DJF ($2\text{CO}_2 - 1\text{CO}_2$) surface net longwave radiation differences. Contour interval is 20 W m^{-2} .

Figure 26(e). Contour plots for DJF (2CO₂ - 1CO₂) surface air temperature differences. Contour interval is 2 K.

(A) DJF $D(SW)$ VS. $D(T)$

(B) JJA $D(SW)$ VS. $D(T)$

Figure 27. Scatter plots of ΔSW vs. ΔT_a in the regions for (a) DJF and (b) JJA.

(A) DJF P VS. DEL(P)

(B) JJA DEL(P) VS. P

Figure 28. Scatter plots of $\Delta(\text{precipitation})$ vs. control precipitation in the regions for (a) DJF and (b) JJA. Points representing the same region from different models are connected by a solid line (U.S. regions), a dotted line (China regions) or a dashed line (ocean regions).

(A) DJF D(LDN) VS D(LUP)

(B) JJA D(LDN) VS D(LUP)

Figure 29. Scatter plots of $\Delta LW\downarrow$ vs. control $\Delta LW\uparrow$ in the regions for (a) DJF and (b) JJA.

(A) DJF D(SH) VS. D(LH)

(B) JJA D(SH) VS. D(LH)

Figure 30. Scatter plots of ΔLH vs. control ΔSH in the regions for (a) DJF and (b) JJA.

Figure 1 is a scatter plot showing the temperature dependence of the magnetic susceptibility χ (emu/mole) versus temperature T (K) for various samples of the compound. The x-axis ranges from 245.0 to 305.0 K, and the y-axis ranges from 0.0 to 25.0 emu/mole. Data points are labeled with sample identifiers: L03, N03, S03, L01, S01, L02, N01, S02, N02, SC2, SU1, SC3, SU2, LUT, NC1, LUT, NC2, LEE3, NC3, SC4, LC2, NC2, L04, NC4. The plot shows a general increase in susceptibility with increasing temperature, with some samples showing a sharp increase at higher temperatures.

Figure 1 is a scatter plot showing the relationship between Temperature (T in K) on the x-axis and Magnetic Resistance (MR in G/KG) on the y-axis. The x-axis ranges from 245.0 to 305.0 K, and the y-axis ranges from 0.0 to 25.0 G/KG. Data points are labeled with codes: LC1, SC1, NO3, LC4, NC3, LU2, NCW3, LC3, NC3, LU1, SC3, NO2, SC1, LU1, SC2, NO1, NO2, LU1, NU3, NU2, and NU1. The points generally show an increasing trend of MR with increasing T , with some scatter at higher temperatures.

119

END

**DATE
FILMED**

5 / 11 / 93