

STIMULATION OF BREAST GROWTH BY HYPNOSIS

APPROVED:

Donald L. Whaley  
Major Professor

Jack R. Haynor  
Minor Professor

Harold D. Followay  
Chairman of the Psychology Department

Robert B. Toulous  
Dean of the Graduate School

DW

Williams, James E., Stimulation of Breast Growth by Hypnosis. Master of Science (Clinical Psychology), May, 1973, 27 pp., 3 tables, 3 illustrations, bibliography, 30 titles.

The bosom is a major sex symbol in our society. Because the female breast currently holds a place of prominence as an index of female sexuality, a frequent complaint among women is the failure of the breasts to develop to a fashionably desirable size.

Mechanical exercisers and cosmetic preparations have frequently been utilized in an effort to enlarge the breasts of women who felt theirs were less than ideal size, but the effectiveness of these procedures has not been adequately substantiated. Surgical augmentation of the breasts offers a solution, but it is expensive and it requires hospital confinement.

Normal maturation and growth of the female breasts depends on the development and functioning of the pituitary/gonadal/adrenal mechanisms in the production of gonadotropic hormones. Research has shown that psychic factors can influence these mechanisms; emotional states in both humans and animals can alter gonadotropic functions.

The purpose of this study was to determine whether hypnotic suggestions could influence the physiological mechanisms associated with breast growth to produce a significant increase in breast size in an adult female population.

DW

The subjects consisted of nineteen volunteer female students from North Texas State University. The students ranged in age from eighteen to forty years.

The first phase was a pilot study designed to examine whether hypnosis with suggestions for breast enlargement was significantly more effective in stimulating breast growth than was hypnosis without specific suggestions for breast enlargement. Three subjects acted as a control group, and were hypnotized once weekly, but did not receive suggestions for breast growth. The experimental group consisted of three subjects who were hypnotized once weekly and were given suggestions for breast growth.

Based on a comparison of expired breast measurements before treatment was initiated and at the completion of the treatment procedure, difference between treatment groups was examined using the Mann-Whitney U test. The results indicate there was a significant difference ( $U=0$ ,  $p.<.05$ ) between the control group and the experimental group. There was no increase in breast measurements in the control group, while all subjects in the experimental group exhibited an increase in breast measurements.

In the second phase, each of the remaining thirteen subjects acted as her own control in examining the effectiveness of hypnotic suggestions in stimulating breast growth. The treatment procedure consisted of weekly hypnosis during which suggestions were given for breast enlargement.

2

Each subject's record of breast measurements for three weeks prior to initiation of the treatment procedure was averaged, and this mean was used as the baseline measure. The breast measurements taken during the last three weeks of the treatment procedure were averaged, and this mean was used as the treatment measure.

Paired comparisons of baseline and treatment data were examined, using the Wilcoxon signed-rank test. An analysis of the data indicated there was a significant increase in breast dimensions following the treatment procedure ( $U=0$ ,  $p < .005$ ). Increases in individual breast measurements ranged from one to three and one-half inches. All subjects experienced an increase in breast measurements following treatment, with an average increase of two and one-eighth inches.

Further investigation may show this procedure to be a satisfactory alternate method to surgical breast augmentation. The preliminary evidence indicates it could meet the need for a method which would produce a significant increase in breast size in a minimum amount of time, and it merits further study on this basis.

STIMULATION OF BREAST GROWTH BY HYPNOSIS

THESIS

Presented to the Graduate Council of the  
North Texas State University in Partial  
Fulfillment of the Requirements

For the Degree of

MASTER OF SCIENCE

By

James E. Williams, B.S. Ed.

Denton, Texas

May, 1973

LIST OF TABLES

Table	Page
I. Expired Breast Measurements . . . . .	11
II. Expired Breast Measurements . . . . .	13
III. Supplemental Breast Measurement Means . . . . .	23

LIST OF ILLUSTRATIONS

Figure	Page
Excerpts from Suggestions Used . . . . .	20
1. Areas of Breast Measurement . . . . .	21
2. Contrast of Chest Reduction to Breast Enlargement . . . . .	22

## STIMULATION OF BREAST GROWTH BY HYPNOSIS

The bosom is a major sex symbol in our culture. Because of this, many women are concerned with the appearance of their breasts (Jesser, 1971). Some who feel their breasts are too small, sag an unusual amount, or otherwise deviate measurably from the ideal image, often feel deformed, and suffer self-consciousness or anxiety. Since the female breast currently holds a place of prominence as an index of female sexuality in our society, a frequent complaint among women is the failure of the breasts to develop to a fashionably desirable size. In keeping with other American values, "big" is equated with "good" (Winch, 1952). To this end, manufacturers of foundation garments claim styles designed to "lift", "separate", "pad", "cradle", "form", and "shape" in such a way that they enhance the contour of the breasts and give the impression of greater size.

Superficial enhancement of this sort has never been completely satisfactory to all women. The extent to which females are concerned with exhibiting breasts deemed provocative and desirable by popular standards is reflected in the variety of mechanical exercisers, cosmetic preparations, and surgical procedures which have been employed from time to time in an effort to produce lasting enlargement of the breasts.



The value of mechanical exercisers and cosmetic preparations is questionable. The paucity of empirical data which might support the effectiveness of mechanical exercisers precludes an evaluation of their worth. Little more is available regarding the effectiveness of cosmetic preparations. Estrogen creams applied to the breasts usually result in stimulation of the nipples, but there is little, if any, resultant breast growth in the woman who has normal circulatory levels of estrogen. Simultaneous administration of prolactin and estrogen in large amounts may stimulate the breasts of the normal woman, but evidence for this is not good (Lloyd and Leatham, 1964; Lloyd, 1964, 1968).

Early attempts at surgical breast augmentation were effective for a time, but ultimately proved to be either troublesome, uncomfortable, or harmful. Fatty tissue from the buttocks grafted beneath the breasts was successful for a time, but in some instances the fat was absorbed into the system, leaving the breasts reduced to their previous size. Injections of paraffin tended to migrate from one part of the breast to another, and to form lumpy deposits. Injections of liquid silicone appeared successful in increasing the breast size, but the Food and Drug Administration forbade its use because it was felt that it was not safe.

An apparently successful solution was an operation involving the implantation of a flexible silicon rubber bag filled with liquid silicon between the breast tissue and

the chest wall. After healing, breasts treated by this method retained the natural feel and appearance of untreated breasts (Brown, 1968). The disadvantages of this procedure are that the operation necessitates hospital confinement for about one week, and costs from \$750 to \$1700, plus the hospital expenses.

Normal maturation and growth of the female breasts depends on the development and functioning of the pituitary/gonadal/adrenal mechanism in the production of gonadotropic hormones. During the period of development there are temporary imbalances in the amounts of the numerous hormones secreted, and there are individual differences in the responsiveness of the sexual end-organs to the various hormones (Brasel and Blizzard, 1968). The activity of the final effector organ is controlled by the anterior pituitary, which is regulated in turn by the hypothalamus (Rasmussen, 1968).

Reichlin (1968) stated that it is important to recognize that nerve pathways exist which bring the hypothalamus "...under the influence of the 'visceral brain', now recognized to be the anatomic substrate of emotion. Through these pathways, emotional states in both humans and animals can alter gonadotropic function (p. 987)". Hitschmann (1928) reported such changes in his analytic treatment of frigidity. In the women he treated, he found such masculine signs as flat breasts, little mustaches, and large hands.

Hitschmann did not consider it proven that there was a change in the endocrine metabolism in these women when they changed during an analysis, but the breasts grew and the figure became better rounded. Groddeck (1921) and Deutsch (1926) reported cases in which psychic factors were decisive in producing breast growth. During psychoanalysis in these cases, anomalous breast changes occurred in women in whom one breast had been retarded in development.

Mohr (1925) reported the case of a girl in whom strong emotions during puberty resulted in a psychically conditioned inhibition of pubescence for a period of ten years. When psychotherapy was initiated, pubescence was completed in a few months, with menses, development of the breasts, increase of the thyroid, and disappearance of hairiness of the chin, which had been present at the beginning of treatment. Mohr expressed no doubt that endocrine factors had played a considerable role as a connecting link, but was equally certain that the psychic factors were primary, activating the endocrines secondarily. Mohr stressed the fact that here "psychic" is by no means synonymous with "suggestion"; that this psychosomatic interaction occurred without any suggestion.

That suggestion can play an active role has been more than amply demonstrated, however. Hypnotic suggestions can produce a variety of effects which transcend normal voluntary capacities. It is practicable to influence both

mentally and physically the function of any reactive system within the organism. Even in light hypnosis, there is increased control over the autonomic nervous system, and indirectly, of all the glands and organs it supplies. Many of the phenomena which have been reported lend evidence to the thesis that hypnotic suggestion can bring about psychobiological changes in the total organism which are quite impossible to obtain in the waking state (Wolberg, 1948; Gorton, 1949; Weitzenhoffer, 1951, 1953; Van Pelt, 1964; Reiter, 1965).

Among these phenomena, hypnotic regression is one of the most remarkable. There is evidence that when some individuals are regressed, experience and behaviors which existed at the earlier age are reactivated, and the organic conditions of that period may be re-established. Wolberg (1945) described this as an actual organic reproduction of an earlier period of life in which past patterns of ideation and behaviors are revived. Experiences subsequent to the regressed age appear to have no influence on the subject's awareness or behavior (Wolberg, 1948; Weitzenhoffer, 1957; Le Cron, 1965; Norgarb, 1965).

Kupper (1945) reported appearance of a pre-convulsive normal electroencephalogram in an epileptic patient during hypnotic regression to an age that predated the onset of his epilepsy.

Ford and Yeager (1948) reported the demonstration of a right homonymous hemianopsia during the hypnotic regression of a patient who had exhibited a right homonymous hemianopsia prior to removal of a colloid cyst from the floor of the third ventricle and subsequent return of normal vision.

Girido-Frank and Bowersbuch (1948) reported the recovery of the Babinski sign of plantar dorsiflexion in three adult subjects regressed to the age of five or six months. They also found that changes in peripheral chromaxie accompanied the change in plantar reflex. Le Cron (1965) confirmed their findings, using three different subjects. At the regressed age of five months, he also found that the sucking reflex of infancy revived.

Conversely to age regression, Erickson (1954) employed a technique of orientation into the future as an hypnotherapeutic procedure. With this procedure of "time projection", Erickson reported the patient was enabled to achieve "...a detached, dissociated, objective and yet subjective view of what he believed at the moment he had already accomplished, without awareness that those accomplishments were the expression in fantasy of his hopes and desires (p. 261)".

Klemperer (1953, 1954) reported on changes of the body image in directed regressions and visualizations during

hypnoanalysis. These body images are built up and accompanied by perceptions, emotions, conceptions, actions, and social connections. She reported the occurrence of felt and seen changes in tissue, organs, and body systems.

Research suggests that the hypnotic procedures of regression, time projection, and changes in body image can produce psychological phenomena which are capable of stimulating a variety of physiological responses. The purpose of this study was to determine whether these procedures could influence the physiological mechanisms associated with breast growth to produce a significant increase in breast size in an adult female population.

## Method

### Subjects

The subjects consisted of nineteen volunteer female students from North Texas State University. Five were graduate students, and the remaining fourteen were undergraduates. Subjects ranged in age from eighteen to forty years, with a mean age of twenty-four years. Fifty-three per cent of the subjects were married, and forty-seven per cent were single. Of the married subjects, sixty per cent had borne children. Sixty per cent of the married subjects and fifty-five per cent of the single subjects reported they were taking birth control pills. The age at menarche ranged from nine years to fifteen years, with a mean age

of twelve years. All subjects reported their weight had been constant (within five pounds) for a minimum period of six months preceeding the experiment.

### Apparatus

Measuring apparatus consisted of Starrett ten-inch outside calipers, Johnson No. 46 vernier calipers, and a seventy-two inch flexible measuring tape.

### Procedure

The study consisted of two phases. The first phase was a pilot study designed to compare two treatments. The experimental group consisted of three subjects who were hypnotized once weekly and received suggestions for breast growth. The control group consisted of three subjects who were hypnotized once weekly, but received no suggestions for breast growth.

At the initial treatment period of each subject, the expired breast measurements were taken around the bust on the horizontal plane of the nipples. Each subject, under the direction of the experimenter, took the measurements of another subject during the initial and all subsequent treatment periods. In addition, all measurements were verified by a third subject. Then hypnosis was induced and suggestions were given.

The suggestions given to each subject in the control group consisted only in the establishment of a variety of

sensory hallucinations. The suggestions given to each subject in the experimental group consisted of regression to a period when the breasts were developing, and the sensation of breast growth was suggested during this period. Then suggestions of time projection to an unspecified future date were given and the subject was directed to visualize her body image with increased breast size (see appendix).

Treatment periods averaged from one to two hours each, and were continued for a period of twelve weeks. Each subject's expired breast measurements were recorded at each of the weekly hypnosis sessions. The same suggestions each subject received during the initial treatment period, were repeated at each subsequent session.

The second phase of the study involved the remaining thirteen subjects. These subjects acted as their own controls to determine the effectiveness of hypnotic suggestion in breast enlargement.

Each subject's breast measurements were taken weekly for a period of three weeks to establish a baseline prior to the initiation of the treatment procedure. Inspired and expired measurements were taken around the bust on the horizontal plane of the nipples. Expired measurements were taken around the chest on a horizontal plane immediately below the base of the cup of the breasts. Measurements were also taken from the base of the cup to the nipple, from the



sternum to the nipple, from the lateral periphery to the nipple, and the span from nipple to nipple. Figure 1 of the appendix illustrates the areas of measurement. Each measurement throughout the baseline and treatment procedure was made by the experimenter and confirmed by the subject, and verified by a second subject.

During the baseline period no hypnosis was induced, and no suggestions were given for breast growth. Immediately following each subject's final baseline measurements on the third week, hypnosis was induced and the treatment procedure was initiated.

The treatment procedure consisted of a series of suggestions for regression to a period when the breasts were developing, and the sensations of breast growth were suggested during this period. Suggestions were then given for time projection to an unspecified future date, and the subject was directed to visualize her body image with increased breast size (see appendix).

Treatment periods averaged from one to two hours each, and the treatment procedure was followed once weekly for a period of twelve weeks. Immediately following each treatment procedure, the same measures used to establish the subject's baseline were taken and recorded. The same suggestions given during the initial treatment period were repeated at each subsequent treatment period.

## Results

The first phase examined, in the pilot study, whether hypnosis with suggestions for breast enlargement was significantly more effective in stimulating breast growth than was hypnosis without specific suggestions for breast enlargement.

The criterion for effectiveness was the amount of change in the breast measurements of the subjects between the initiation and termination of the treatment procedure.

Table I presents the data on breast measurements for the first phase.

TABLE I  
EXPIRED BREAST MEASUREMENTS

SUBJECT NUMBER	INITIAL	TERMINAL
GROUP 1-Weekly hypnosis without suggestions for breast growth		
1-C	35"	35"
2-C	37"	37"
3-C	32"	32"
GROUP 2-Weekly hypnosis with suggestions for breast growth		
1-E	33"	34 1/2"
2-E	33 1/2"	34"
3-E	30 1/2"	33 3/8"

Differences between the treatment groups were examined using the Mann-Whitney U test described by Siegel (1956).

Results indicate there was a significant difference ( $U=0$ ,  $p.<.05$ ) between the control group (weekly hypnosis without suggestions for breast enlargement) and the experimental group (weekly hypnosis with suggestions for breast enlargement).

The second phase provided a further examination of the effectiveness of hypnotic suggestion in the stimulation of breast growth.

Criterion for the effectiveness of the treatment was a significant increase in breast measurements. Expired breast measurement taken on the horizontal plane of the nipples was selected as the primary index of breast enlargement.

Each subject's record of breast measurements for the three weeks prior to initiation of the treatment procedure was averaged and this mean was used as the baseline measure. The breast measurements taken during the last three weeks of the treatment procedure were averaged and this mean was used as the treatment measure.

All breast measurements were recorded in fractions of an inch. To facilitate machine computation, decimal equivalents were substituted for all fractions of an inch.

Baseline expired breast measurements ranged from 30.21 inches to 39.08 inches, with a mean measure of 33.64 inches. Treatment expired breast measurements ranged from 32.33 inches to 41.33 inches, with a mean measure of 35.75 inches.

The mean increase for the group was 2.11 inches. Individual increases ranged from a minimum of 1.00 inches to a maximum of 3.54 inches.

Table II presents the data on the individual breast measurement means for the second phase.

TABLE II  
EXPIRED BREAST MEASUREMENTS

SUBJECT NUMBER	BASELINE MEAN	TREATMENT MEAN
101	34.04	35.29
102	33.96	35.42
103	33.08	35.67
104	33.58	35.50
105	34.04	36.17
106	39.08	41.33
107	31.92	33.83
108	33.25	35.91
109	37.04	40.58
110	30.21	32.33
111	33.17	35.33
112	32.00	33.00
113	31.96	34.33

Paired comparisons of baseline and treatment data were examined, using the Wilcoxon signed-rank test described by Siegel (1956). An analysis of the data indicated there was a significant increase in breast dimensions following the treatment procedure ( $T=0$ ,  $p.<.005$ ).

As a control measure, expired chest measurements taken at the base of the breasts concurrently with the expired

breast measurements were compared. Baseline expired chest measurements ranged from 28.00 inches to 35.00 inches, with a mean measure of 29.94 inches. Treatment expired chest measurements ranged from 27.00 inches to 34.00 inches, with a mean measure of 29.27 inches. The mean decrease for the group was .67 inches.

The contrast of chest reduction to breast enlargement is graphically illustrated in Figure 2 of the appendix.

The remainder of the breast measurements taken were examined, and a comparison of baseline and treatment means disclosed an average increase in the treatment mean of .16 inches over baseline from the nipple to the cup base, .11 inches from the nipple to the sternum, .39 inches from the nipple to the lateral periphery, and .20 inches in the nipple span.

Data comparing the baseline and treatment means of these measures are presented in Table III of the appendix.

#### Discussion

In phase one, which was primarily a pilot study, no increase in breast measurements was observed in the control group when they were hypnotized weekly without suggestions for breast growth. All subjects in the experimental group exhibited an increase in breast measurements when they were hypnotized weekly with suggestions for breast growth.

Several extraneous variables were held constant for both groups. The control and the experimental group each contained one married and two single subjects. Each of the married subjects had borne children. Each group included one subject over thirty years of age. The average age in each group was twenty-seven years, and the average age at menarche in each group was twelve years, with none occurring earlier than eleven or later than thirteen years.

Since the experimental and control groups were approximately equal in marital status, child bearing, age, and age at menarche, it does not appear that these factors influenced the results to any significant degree.

This preliminary evidence suggests that hypnosis per se had no direct effect on the breast enlargement, but that hypnosis with suggestions for breast growth was effective in stimulating breast enlargement. This is in accord with Weitzenhoffer's (1951) observation that "...hypnosis per se does not produce transcendence (of normal capacities and functions), while the addition of various suggestions can bring about such an effect (p. 277)".

In the second phase, each subject acted as her own control in examining the effectiveness of hypnotic suggestions in stimulating breast growth.

Baseline data taken prior to treatment was compared with the data of the last three weeks of treatment and it was determined that increases in individual breast measurements

ranged from one to three and one-half inches. All subjects experienced some degree of enlargement following treatment, with an average increase of two and one-eighth inches.

Expired chest measurements were taken weekly to determine if the increase in breast measurements might be due to enlargement of the rib cage or upper torso. To the contrary, it was found that the chest measurements decreased an average of five-eighths of an inch in association with the increase in breast measurements (see Figure 2 of the appendix).

Supplementary breast measurements indicate the increase in breast size was symmetrical, with average increases of about one-eighth of an inch from the cup base to the nipple and the sternum to the nipple, three-eighths of an inch from the lateral periphery to the nipple, and one-fourth of an inch in the span of the nipples (see Table III of the appendix).

An examination of the background data failed to disclose any factors which might have significance in selecting the population with which this particular procedure might be most effective. The average increase in breast measurement was about five-eighths of an inch less in the married subjects than in the single subjects, but the sample of each was too small to draw conclusions from. Married subjects who had borne children exhibited slightly larger average

increases than those who had not. There were no significant differences in breast growth between subjects who were taking birth control pills and those who were not. Two subjects discontinued birth control pills midway through the procedure without noticeably altering their growth curve.

The only variable which might appear to influence the degree of breast enlargement was the age of menarche. The subjects who had an age of menarche of eleven years or less showed an average increase of about one and one-fourth inch, compared to the group average of two and one-eighth inches. However, with a sample of only four subjects in this category, little can be inferred from this data without further investigation.

The degree of suggestibility did not appear to be a determining factor in the effectiveness of the treatment procedure. The basic method of hypnotic induction used was a variation of the hand levitation procedure described by Wolberg (1948). No attempt was made to achieve stability in depth of hypnosis between subjects, since one of the extraneous considerations of the study was to determine if depth of hypnosis was a significant factor in the degree to which suggestions could influence breast growth. Since some subjects who were not able to achieve deep hypnosis showed a greater increase in breast size than some who were, it appears that depth of hypnosis was not a significant factor.



It was not within the scope of this paper to determine what, if any, changes in the hormonal sphere were brought about through the treatment procedure investigated. Whatever the psychobiologic changes which are involved in the mechanisms associated with breast growth, it appears a reality that hypnotic suggestion can influence them to a significant degree.

Further investigation may show this procedure to be a satisfactory alternate method to surgical breast augmentation. The preliminary evidence indicates it could meet the need for a method which would produce a significant increase in breast size in a minimum amount of time, and it merits further study on this basis.

APPENDIX


Excerpts from suggestions used

During regression phase:

"...back to the time when your body was in the maturing process and your breasts began to grow. Feel the pleasant tingling sensation in your breasts as the maturing process goes on and your breasts continue to grow..."

During time projection phase:

"...and now you see yourself in the future, standing nude before a mirror. Your breasts have grown to the size you desired. Notice, see and feel the firmness and fullness of your breasts, now that they have grown as large and shapely as you wanted..."



- A - Expired breast measurement
- B - Expired chest measurement
- C - Measurement from cup base to nipple
- D - Measurement from sternum to nipple
- E - Measurement from lateral periphery to nipple
- F - Measurement of span from nipple to nipple

Figure 1-Areas of breast measurement




Figure 2-Contrast of chest reduction to breast enlargement

TABLE III  
 SUPPLEMENTAL BREAST MEASUREMENT MEANS

AREA OF MEASURE	BASELINE MEAN	TREATMENT MEAN
Nipple to cup base		
Left breast	2.40	2.50
Right breast	2.33	2.55
Nipple to sternum		
Left breast	4.15	4.25
Right breast	4.17	4.29
Nipple to lateral periphery		
Left breast	3.96	4.33
Right breast	3.96	4.36
Nipple span	8.03	8.23

## References

- Brasel, J. A. & Blizzard, R. M. The influence of the endocrine glands upon growth and development. In R. H. Williams (Ed.), Textbook of Endocrinology, (4th ed.). Philadelphia: W. B. Saunders, 1968.
- Brown, W. E. Cosmetic Surgery. New York: Stein & Day, 1968.
- Deutsch, F. Der gesunde und der kranke Körper in psychoanalytischer Betrachtung. Cited by H. F. Dunbar, Emotions and Bodily Changes, (4th ed.). New York: Columbia University Press, 1954.
- Erickson, M. H. Pseudo-orientation in time as an hypnotic procedure. Journal of Clinical and Experimental Hypnosis, 1954, 2, 261-283.
- Ford, L. F. & Yeager, C. L. Changes in electroencephalogram in subjects under hypnosis. Diseases of the Nervous System, 1948, 9, 190-192.
- Girido-Frank, L. & Bowersbuch, M. K. A study of the plantar response in hypnotic age regression. Journal of Nervous and Mental Disorders, 1948, 107, 443-458.
- Gorton, B. E. The physiology of hypnosis, I & II. Psychiatric Quarterly, 1949, 23, 317-343, 457-485.
- Groddeck, G. Ueber die psychoanalyse des organischen im menschen. Cited by H. F. Dunbar, Emotions and Bodily Changes, (4th ed.). New York: Columbia University Press, 1954.

- Hitschmann, E. Psychoanalyse trötzt hormonen. Cited by H. F. Dunbar, Emotions and Bodily Changes, (4th ed.). New York: Columbia University Press, 1954.
- Jesser, C. J. Reflections on breast attention. Journal of Sex Research, 1971, 7, 13-25.
- Klemperer, E. Changes of the body image in hypnoanalysis. Journal of Clinical and Experimental Hypnosis, 1954, 2, 157-162.
- Klemperer, E. Hypnosis and hypnoanalysis. Journal of American Medical Women's Association, 1953, 8, 164.
- Kupper, H. L. Psychic concomitants in wartime injuries. Psychosomatic Medicine, 1945, 7, 15-21.
- Le Cron, L. M. A study of age regression under hypnosis. In L. M. Le Cron (Ed.), Experimental Hypnosis. New York: Citadel Press, 1965.
- Lloyd, C. W. Problems associated with sexual maturation and lactation. In C. W. Lloyd (Ed.), Human Reproduction and Sexual Behavior. Philadelphia: Lea & Febiger, 1964.
- Lloyd, C. W. The ovaries. In R. H. Williams (Ed.), Textbook of Endocrinology, (4th ed.). Philadelphia: W. B. Saunders, 1968.
- Lloyd, C. W. & Leathem, J. H. Growth and development of the breast and lactation. In C. W. Lloyd (Ed.), Human Reproduction and Sexual Behavior. Philadelphia: Lea & Febiger, 1964.



- Mohr, F. Psychophysische Behandlungsmethoden. Cited by H. F. Dunbar, Emotions and Bodily Changes, (4th ed.). New York: Columbia University Press, 1954.
- Norgarb, B. A. Rorschach psychodiagnosis in hypnotic regression. In L. M. Le Cron (Ed.), Experimental Hypnosis. New York: Citadel Press, 1965.
- Rasmussen, H. Organization and control of endocrine systems. In R. H. Williams (Ed.), Textbook of Endocrinology, (4th ed). Philadelphia: W. B. Saunders, 1968.
- Reichlin, S. Neuroendocrinology. In R. H. Williams (Ed.), Textbook of Endocrinology, (4th ed.). Philadelphia: W. B. Saunders, 1968.
- Reiter, P. J. The influence of hypnosis on somatic fields of function. In L. M. Le Cron (Ed.), Experimental Hypnosis. New York: Citadel Press, 1965.
- Siegel, S. Nonparametric Statistics. New York: McGraw-Hill, 1956.
- Van Pelt, S. J. Will hypnosis revolutionize medicine? In R. Rhodes (Ed.), Therapy Through Hypnosis. New York: Citadel Press, 1964.
- Winch, R. F. The Modern Family. New York: Henry Holt, 1952.
- Weitzenhoffer, A. M. The transcendence of normal voluntary capacities in hypnosis: An evaluation. Personality, 1951, 272-282.
- Weitzenhoffer, A. M. Hypnotism: An Objective Study in Suggestibility. New York: John Wiley & Sons, 1953.

Weitzenhoffer, A. M. General Techniques of Hypnotism. New York: Grune & Stratton, 1957.

Wolberg, L. R. Hypnoanalysis. New York: Grune & Stratton, 1945.

Wolberg, L. R. Medical Hypnosis. New York: Grune & Stratton, 1948. 2 vols.