DOE/ER/40264--6

DE92 010014

THEORETICAL RESEARCH IN
NUCLEAR STRUCTURE AND NUCLEAR COLLECTIVE MOTION

Progress Report
For the period March 1, 1991 - February 29, 1992

Abraham Klein

University of Pennsylvania
Philadelphia, Pennsylvania 19104

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

January 1992

Prepared for

THE U. S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-86ER-40264

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ABSTRACT

During the past funding period, we have completed, continued or initiated research on the following topics:

- (i) The formulation of a theory of large amplitude collective motion in the domain of small velocities has been largely completed. Representative applications to the approximate quantization of non-separable systems and to chemical reactions have been carried out. A program of application to nuclear collective motion has been initiated and continues.
- (ii) An exhaustive review article on boson mapping of Lie algebras with applications to nuclear physics has been prepared in collaboration with E. R. Marshalek and published in *Reviews of Modern Physics*.
- (iii) As an outgrowth of the work on the review article, we have been led into a new research program associated with the development of the powerful method of vector coherent states. We have pointed to a relationship between the mathematical problem of mapping general tensors under an algebra and an older idea of nuclear physics origin called the quantized Bogoliubov transformation. The exploitation of this idea has a dual importance, providing on the one hand the basis for constructing the matrices of all tensors under the algebra and on the other hand a rigorous foundation for core-particle coupling models.
- (iv) The Kerman-Klein method of quantization and restoration of broken symmetry has been applied to the standard Skyrme model. A study of the restoration of translational invariance has been carried out. A program for the restoration of rotational and isospin invariance that entails replacing the standard mean-field (hedgehog) approach by a symmetry-preserving coupled-channels approach is well-advanced.
- (v) We have proposed an alternative approach to the derivation of relativistic two-body equations from field theories, and have recently completed a calculation in scalar QED, looking for resonances in "electron-positron" scattering. In contrast to claims found in the literature, we believe that we have shown

that they are absent for the models in question. Calculations for real QED are in progress.

(vi) We have recently developed a new method for the study of quantum fluctuations about mean-field solutions. Though our original motivation was to study quantum corrections in the theory of large-amplitude collective motion, we now intend to undertake a systematic investigation of other applications. So far we have completed studies of both the (old) problem of ground-state correlations about equilibrium solutions and the problem of leading quantum corrections for large-amplitude collective motion. We are also well advanced in a study of quantum fluctuations at high angular momentum.

(vii) In the project undertaken most recently, we are applying the Kerman-Klein method to a critical study (with the ultimate aim of carrying out applications) of core-particle (core-hole) coupling methods for deformed nuclei. For the low-spin region, we are now investigating the relative importance of changes in the self-consistent potential with angular momentum, which in principle, should be considered on the same footing as the Coriolis coupling. We later plan to study the rotating shell model.

Of these topics, only the review article is a closed chapter. The remaining topics contain the main ingredients of our research program for the immediate future. For instance, items (iv) and (vii) are thesis investigations; in the case of project (iv) we foresee completion of the work in not less than 1 and not more than 2 years, whereas the investigation described under (vii) is in its infancy, and may require almost the full extent of the renewal period. Item (i) is in a temporary holding pattern because of problems encountered and because items (iii), (v), and (vi) are in a period of rapid development that could continue for some time. Nevertheless, we view the work on large-amplitude collective motion as a major interest and commitment, quite open-ended in the range of interesting applications that one can attempt.

END

**DATE
FILMED**

4 / 23 / 92