2

Secondary Production in Waters Influenced by Upwelling
Off the Coast of Somalia

Sharon Louise Smith

Oceanographic and Atmospheric Sciences Division
Department of Applied Science
Brookhaven National Laboratory
Upton, NY 11973

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This research was performed under the auspices of the United States Department of Energy under Contract No. DE-AC02-76CH00016.

By acceptance of this article, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

482

Abstract

During the southwest monsoon of 1979, plankton samples were collected between the equator and 12°N. Surface samples collected while the research vessel was surveying this area have been analyzed for the abundance of copepod nauplii, primarily the young of the copepod Calanoides carinatus. These samples revealed a sharp rise in the abundance of nauplii in regions where sea-surface temperatures were less than 19°C. The developmental state of the gonads of female C. carinatus also showed that reproduction was likely to be most intense nearshore in cold, recently upwelled water. Lipid content of females was inversely related to gonad development, suggesting that lipids sequestered at some earlier point in the life cycle were being used to synthesize eggs. Secondary production is therefore highest in the nearshore areas having cool sea-surface temperatures. Growth of phytoplankton and growth of C. carinatus seem to be closely coupled; daily ingestion rates estimated in nearshore areas where potential secondary production was high were equal to or greater than measured rates of primary production.

Introduction

The copepod Calanoides carinatus is a common and prominent member of the plankton of upwelling areas around the African continent from northwestern Africa (Boucher et al., 1975; Schnack, 1982) to Ivory Coast (Binet and Suisse de Sainte Claire, 1975) to Ghana (Mensah, 1975) to southwest Africa (Borchers and Hutchings, 1986) to the Somali coast (Smith, 1982). Its abundance and relatively large size for tropical, epipelagic plankton make it an appealing subject for study. Off Somalia, it has been shown to dominate the biomass of zooplankton during the upwelling season in two different years, 1964 and 1979 (Smith, 1984). Its secondary production off Somalia can now be estimated, using old samples and techniques developed since 1979. The very high concentrations of mesopelagic fish in the region (Banse, 1984) cause one to try to estimate the secondary productivity that might support the mesopelagic fish. Widespread suboxic conditions in the Arabian Sea basin below 200 m (Wyrski, 1973; Swallow, 1984) suggest that coupling between primary and secondary production may not be close. Thus, there is an apparent contradiction. Large biomass of mesopelagic fish suggests a coupled and efficient food chain, while low oxygen conditions suggest a lack of coupling between primary and secondary production. Secondary productivity of a dominant herbivore such as Calanoides carinatus is a key variable that should indicate how the epipelagic food chain of this region is structured.

Materials and Methods

Zooplankton were collected from R.V. Columbus Iselin in oblique hauls of Bongo frames fitted with 223 and 333 μm mesh nets and T.S.K. and General Oceanics flowmeters. A mechanical time-depth recorder (Benthos) was mounted one meter above the net mouth. The collection from the 223 μm mesh net was preserved in 5% neutral formalin-seawater solution. Underway samples of zooplankton were collected on 102 μm mesh for five minutes each half-hour during steaming from the output of a through-hull fitting 2 m below the waterline. Samples were preserved in 5% neutral formalin seawater solution. Temperature, salinity, chlorophyll a, and rate of flow were observed at the same time. A detailed account of the methods and data is in Smith and Lane (1981).

Dry weights of various copepodid stages were determined by picking a number of individuals of a given stage from a sample, rinsing the specimens several times in distilled water and transferring them to a preweighed dish for drying at 50°C. Dishes were reweighed using a Cahn model 26 electrobalance. Dry weights were corrected for losses due to preservation in formalin amounting to 40% on average. Lipid content of adult females and copepodid stage V's were determined by repeated extraction in chloroform:methanol (2:1, V:V). The difference in dry weight before and after extraction was the total weight of lipid in the copepod. Lipid loss due to preservation was estimated to be 47%, based upon comparisons between fresh and preserved lipid contents measured in female Calanus hyperboreus captured in the Greenland Sea. Conversions between dry weight and carbon were according to

Cushing et al. (1958). Respiration was estimated using equations of Vidal and Whitley (1982).

The state of gonadal development in female Calanoides carinatus was noted and scored. The scale used in scoring was derived from those of Marshall and Orr (1972), Runge (1985), and Smith (1990). Categories reflect the relative reproductive maturity of the females (Table 1).

Results

During the southwest monsoon of 1979, various stages of Calanoides carinatus were collected at 23 of 31 stations (Figure 1). The predominance of Calanoides carinatus in the two areas of upwelling (Smith 1982, 1984), the steepness of the gradients associated with upwelling features, and the speed of the Somali Current make it difficult to depict the spatial pattern in secondary production. Therefore, the abundances of various stages are presented as a function of sea-surface temperature whose spatial distribution is shown in Figure 1. The earliest copepodids, CI-CII, were collected in areas where sea-surface temperatures were between 18 and 20°C (94% of total numbers) with a maximum between 18 and 19°C (63% of total numbers; Figure 2). Copepodid stage III also had its peak abundance between 18 and 19°C, where 50% of total numbers were collected (Figure 2). Another 43% of total numbers of CIII were collected between 19 and 21°C (Figure 2). Copepodid stage IV showed a wider distribution, being relatively evenly spread between 18 and 23°C (Figure 2). Copepodid stage V was bimodal, with 43% between 18 and 19°C and 29% between

22 and 23°C (Figure 2). Adult females were most abundant in the temperature range in which CI's were dominant (18-19°C), but 11% of the total numbers of females were collected where sea-surface temperatures were lowest (16-18°C; Figure 2).

Using the maximum specific daily production rates of juvenile Calanoides carinatus observed off southwest Africa (23% dry body weight per day; Verheye, 1989) and the daily rates of egg production observed off southwest Africa applied to females captured off Somalia (60% dry body weight per day; Borchers and Hutchings, 1986), maximum secondary production was calculated for all stations where C. carinatus were captured. Highest potential secondary production occurred between 18 and 19°C (Figure 3). This reflects both the elevated concentrations of juvenile stages and reproductive females at these temperatures. Potential egg production accounted for an average of 84% of the calculated secondary production between 18 and 19°C (Figure 3). The distribution of nauplii confirms the fact reproduction was most intense where sea-surface temperatures were lowest (Figure 4).

Analysis of the state of gonadal development of adult females provides additional evidence that spawning by Calanoides carinatus was taking place primarily nearshore. Thirty females from each of two stations nearshore and two stations offshore (Figure 1) were scored. Between 47% and 77% of the nearshore females had gonads in state 4 (see Table 1 for description of characteristics of various states), the state in which egg laying takes place. Only 3% of nearshore females had gonads in state 1, the state in which only the ovary is visible and oviducts contain no oocytes. In contrast

with nearshore, between 57% and 94% of females collected offshore had gonads in state 1 and no offshore female had reached state 4. In summary, female C. carinatus with gonads fully developed and ready for egg laying were collected only nearshore. (The large numbers of copepod nauplii collected where sea surface temperatures were between 28 and 29°C are therefore not likely to be those of C. carinatus.) When potential secondary production is averaged for nearshore stations (less than 9 km from the coast) and for offshore stations (more than 250 km from the coast) where Calanoides carinatus were collected, we find that potential secondary production is much higher nearshore (Figure 5). Most of the nearshore production is accounted for by egg production of females, while offshore egg production and growth of juveniles are of equal magnitude (Figure 5).

Another aspect of secondary production in this species is the large lipid reserve accumulated by copepodid stage V (up to 71% of total dry body weight; Borchers and Hutchings, 1986). Lipid in females is found in two apparently separate lipid sacs, one in the forehead and one ventral to the gut and oviducts (Figure 6). When females are without large numbers of eggs in the diverticula and oviducts, the lipid sacs are spherical or ellipsoid. As the eggs enlarge and fill the diverticula and oviducts, lipid sacs are deformed as shown in Figure 6. The presence or absence of lipid in females was correlated with gonadal development. All females from the two offshore stations, where gonad state was primarily 1, had observable lipid sacs. Of the adult females collected at the nearshore station where 77% were ready to lay eggs (state 4), only

half had any visible lipid. Of the females at the nearshore station where 47% were ready to lay eggs, 90% of females still had observable lipid. These data suggest that lipid reserves are used by females in the synthesis of eggs. Lipid content of females captured nearshore was 12% of dry body weight near 5°N and 4% of dry body weight near 10°N (Table 2). Copepodid stage V captured offshore near 5°N contained an average of 41% lipid (Table 2).

Discussion

The life cycle of Calanoides carinatus has been most fully described in areas of upwelling off the western coast of Africa, particularly off Ghana and the Ivory Coast. The species is in surface layers during the upwelling season and spends the remainder of the year at mesopelagic depths (Mensah, 1974; Binet and Suisse de St. Claire, 1975) where its respiration is fueled by stored lipids. The lipid that can be accumulated by copepodid stage V during feeding upon the upwelling assemblage of the Benguela Current can be considerable, up to 71% of dry body weight (Borchers and Hutchings, 1986). The evidence from the Somali coast is consistent with patterns established for this species on the west coast of Africa. Secondary production, primarily reproduction, is highest in cool, recently upwelled waters nearshore. As juveniles move offshore in the Somali Current and the offshore-moving component of the Great Whirl, their growth becomes a larger fraction of total potential secondary production of this species. The depths at which C. carinatus survives during the seasons lacking upwelling off Somalia are unknown, but it was not collected

in the upper 200m during the northeast monsoon of 1979 (Smith, 1982).

In another study we estimated ingestion from the summation of secondary production and respiration (Smith and Lane, 1988). Applying the same method to the Calanoides carinatus collected at two nearshore stations (Figure 1), we find that daily ingestion could equal or exceed measured daily primary production at those stations. We can infer that during the upwelling season, growth of phytoplankton is closely coupled to the growth of the Calanoides carinatus population. The life cycle of C. carinatus, wherever it is found, has distinctive characteristics that allow it to exploit a predictably intermittent food supply (Smith, 1982). These characteristics are rapid growth rate (Peterson and Painting, 1990), herbivorous feeding particularly on diatoms (Schnack, 1982), massive lipid storage by subadult stages (Borchers and Hutchings, 1986), and diapause at mesopelagic depths (Mensah, 1974; Binet and Suisse de Sainte Claire, 1975). These characteristics are very similar to those of large calanoid copepods inhabiting polar, subpolar and boreal environments.

References

BANSE, K. (1984) overview of the hydrography and associated biological phenomena in the Arabian Sea, off Pakistan. IN: Marine Geology and Oceanography of Arabian Sea and Coastal Pakistan, B.U. Haq and J.D. Milliman, eds., Van Nostrand Reinhold Company, New York, pp. 271-303.

BINET, D. and E. SUISSSE DE SAINTE CLAIRE (1975) Le copepode planktonique Calanoides carinatus repartition et cycle biologique au large de la Cote d'Ivoire, Cahiers O.R.S.T.O.M., series Oceanographique, 13: 15-30.

BORCHERS, P. and L. HUTCHINGS (1986) Starvation tolerance, development time and egg production of Calanoides carinatus in the southern Benguela Current, Journal of Plankton Research, 8: 855-874.

BOUCHER, J., A. LAUREC, J.-F. SAMAIN, and S. L. SMITH (1975) Etude de la nutrition, du regime et du rythme alimentaire du zooplancton dans les conditions naturelles, par la mesure des activites enzymatiques digestives, Tenth European Symposium on Marine Biology, 2: 85-110.

CUSHING, D. H., G. F. HUMPHREY, K. BANSE and T. LAEVASTU (1958) Report of the committee on terms and equivalents, Rapports et proces-verbaux des reunions, Conseil permanent international pour l'exploration de la Mer, 144: 15-16.

MARSHALL, S. M. and A. P. ORR (1972) The Biology of a Marine Copepod, Springer Verlag, New York.

MENSAH, M. A. (1974) The reproduction and feeding of the marine copepod Calanoides carinatus (Kroyer) in Ghanaian waters, Ghana Journal of Science, 14: 167-191.

PETERSON, W. T. and S. J. PAINTING (1990) Developmental rates of the copepods Calanus australis and Calanoides carinatus in the laboratory, with discussion of methods used for calculation of development time, Journal of Plankton Research, 12: 283-293.

RUNGE, J. A. (1985) Relationship of egg production of Calanus pacificus to seasonal changes in phytoplankton availability in Puget Sound, Washington, Limnology and Oceanography, 30: 382-396.

SCHNACK, S. B. (1982) Feeding habits of Calanoides carinatus (Kroyer) in the upwelling area off northwest Africa, Rapports et proces-verbaux des reunions, Conseil permanent international pour l'exploration de la Mer, 180: 303-306.

SMITH, S.L. (1984) Biological indications of active upwelling in the northwestern Indian ocean in 1964 and 1979, and a comparison with Peru and northwest Africa, Deep-Sea Research, 31: 951-967.

SMITH, S.L. (1990) Egg production and feeding by copepods prior to the spring bloom of phytoplankton in Fram Strait, Greenland Sea, Marine Biology, 106: 59-69.

SMITH, S.L. (1982) The northwestern Indian Ocean during the monsoons of 1979: distribution, abundance, and feeding of zooplankton, Deep-Sea Research, 29: 1331-1353.

SMITH, S. L. and P. V. Z. LANE (1988) Grazing of the spring diatom bloom in the New York Bight by the calanoid copepods Calanus finmarchicus, Metridia lucens and Centropages typicus, Continental Shelf Research, 8: 485-509.

SMITH, S. L. and P. V. Z. LANE (1981) Biological oceanography of the Somali Current Data Report, Informal Technical Report BNL 29098, Brookhaven National Laboratory, Upton, New York, 126 pp.

SWALLOW, J. C. (1984) Some aspects of the physical oceanography of the Indian Ocean, Deep-Sea Research, 31: 639-650.

VERHEYE, H. M. (1989) Distribution, dynamics and production of the copepod Calanoides carinatus (Kroyer, 1849) in the southern Benguela upwelling region. Ph.D. thesis, University of Capetown, Capetown, South Africa, 237 pp.

VIDAL, J. and T. E. WHITLEDGE (1982) Rates of metabolism of planktonic crustaceans as related to body weight and temperature of habitat, Journal of Plankton Research, 4: 77-84.

WYRTKI, K. (1973) Physical oceanography of the Indian Ocean. IN: The Biology of the Indian Ocean, B. Zeitzschel and S. Gerlach, eds., Springer-Verlag, New York, pp. 18-36.

Acknowledgments

The research was supported by National Science Foundation Grant OCE78-25454, and DOE Contract DE-AC02-76CH00016 supported preparation of the manuscript. The Office of Naval Research funded my participation in the International Symposium on the Oceanography of the Indian Ocean. Analyses and data reduction were done by P.V.Z. Lane and E. Schwarting.

Table 1. Calanoides carinatus. Description of reproductive state used for scoring female copepods collected off Somalia (from Smith, 1990)

State	Morphological Description
1	Only ovary visible
2	Small oocytes in ovary with single rows extending forward from ovary in diverticula; single row of small oocytes visible in oviducts in lateral view
3	Oocytes larger with several rows extending forward in diverticula; medial walls of diverticula forced together; more than one row of eggs visible in oviducts in lateral view; eggs and ovary somewhat larger than in state 2
4	Oocytes densely packed into diverticula; abutting medial walls of diverticula extend forward from ovary as dark line to anterior cephalothorax; large oocytes occupy much of cephalothorax; pouches of oocytes visible in oviducts in lateral view extending posteriorly to fifth thoracic segment
5	Spent

Table 2. Gonad development and lipid content of female
Calanoides carinatus collected nearshore and offshore

Station	Location	Gonad Development (%)	Lipid Content (%)	Proportion of Females Lacking Lipid (%)
<u>Nearshore</u>				
13	5°01'N 48°40'E	State 1: 3 State 2: 23 State 3: 27 State 4: 47	12	10
29	10°13'N 51°26'E	State 1: 3 State 2: 0 State 3: 20 State 4: 77	4	50
<u>Offshore</u>				
19	4°20'N 51° 0'E	State 1: 94 State 2: 6 State 3: 0 State 4: 0	41	0
20	4°55'N 51°31'E	State 1: 57 State 2: 27 State 3: 16 State 4: 0	N.D.*	0

*N.D. = Not determined.

Figure Captions

Figure 1. The coast of Somalia showing sea-surface temperature and positions of stations where Calanoides carinatus were collected in 1979. Squares nearshore indicate stations in Table 2 and where ingestion has been estimated from secondary production and respiration. Squares offshore indicate stations used in Table 2.

Figure 2. Abundance of juvenile (A) and adult (B) stages of Calanoides carinatus grouped according to sea-surface temperature at the station of collection.

Figure 3. Secondary production of Calanoides carinatus as a function of sea-surface temperature. Hatched portion of bar indicates estimate of egg production by females, remainder of bar indicates estimate of somatic growth by subadult stages.

Figure 4. Abundance of copepod nauplii grouped according the sea-surface temperature at the point of collection.

Figure 5. Secondary production of Calanoides carinatus from stations less than 9 km from the coast (nearshore) and greater than 250 km from the coast (offshore). Hatched portion of bar indicates estimate of egg production by females, remainder of bar indicates estimate of somatic growth by subadult stages.

Figure 6. Diagram of adult female Calanoides carinatus showing positions of lipid sacs (stippled areas) of females bearing eggs. The ovary (o), diverticula (d), oviducts (od) and gut (g) are also indicated. Females without eggs have lipid sacs that are spherical or ellipsoid in shape.

A

END

**DATE
FILMED
51/13/92**