DOE/ER/13512--T2

TECHNICAL PROGRESS REPORT

DOE/ER/13512--T2

FOR

DE93 008396

GRANT NO:

DE-FG02-86ER13512

WITH

U.S. DEPARTMENT OF ENERGY

OFFICE OF BASIC ENERGY SCIENCES

DIVISION OF ENERGY BIOSCIENCES

TITLE:

**The Mechanism of Switching from an Acidogenic to
Butanol-Acetone Fermentation by *Clostridium acetobutylicum***

**PRINCIPAL
INVESTIGATOR:**

**Palmer Rogers
Professor of Microbiology
University of Minnesota
Medical School**

**CURRENT REPORT
PERIOD:**

July 1990 - December 1992

**TYPE OF BUSINESS/
INSTITUTION**

Non-Profit Educational Institution

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**RECEIVED
MAR 04 1993
OSTI**

MASTER

TECHNICAL PROGRESS REPORT

JULY 1990 - DECEMBER 1992

A. Research Objectives

The overall objective of this project is to elucidate the detailed mechanism by which solvent-forming bacteria such as *Clostridium acetobutylicum* regulate the well-known shift in fermentation pathway between alcohol-acetone and organic acid production. We eventually want to isolate and describe: (1) the regulatory genes and protein elements that determine induction of synthesis of the solvent-pathway enzymes; and (2) how this regulation system interacts with the sporulation induction and development program and with related pathways such as granulose and exopolysaccharide formation in clostridia. A working model for how clostridial control systems work can be derived from recent research on stress systems in *E. coli* and sporulation in *Bacillus subtilis* (Losick, R. and P. Stragier, *Nature* 355:601-4, 1992).

The research proposed was designed to initiate a study of the technique of employing transposable elements that create gene fusions *in vivo* in the Gram-positive bacteria, *Bacillus* and *Streptococcus*, in order to identify and clone specific genes and study functional gene regulation in *Clostridium acetobutylicum*. We and others have recently demonstrated transconjugation and insertion of Tn916 into *C. acetobutylicum* ATCC 824. Some of these transconjugates are also presumptive regulatory mutants. Also by electroporative transformation, we attempted to deliver the 5 Kb Tn917 into *C. acetobutylicum* on plasmid PTV1. The success of using transposon Tn917 and its derivatives in analysis of *B. subtilis* sporulation (Youngman, *et al.* 1985) encouraged its use in sorting out and identifying the elements of fermentation regulation in the clostridia.

B. Progress on the Objectives

1. Analysis of Tn916 insertion mutants of *C. acetobutylicum*.

The transposon Tn916 was introduced into *C. acetobutylicum* ATCC 824 by filter mating conjugation using an *Enterococcus faecalis* donor. Tetracycline-resistant (*tet*^r) mutants were selected and screened for granulose production by staining colonies with iodine vapors. Many granulose-negative, *tet*^r mutants were found, by hybridization to a Tn916 probe, to contain a single insertion of the transposon. Granulose mutants that contained a single transposon insertion were differentiated into five classes by their solvents production profiles: Class 1 mutants did not produce acetone or butanol and had reduced ethanol production when compared to the wild-type parent; Class 2 mutants did not make acetone and had lower levels of ethanol and butanol; Class 3 mutants made reduced levels of all solvents; Class 4 mutants were similar to the wild-type parent; and Class 5 mutants made elevated amounts of acetone, ethanol, and butanol. Class 1 and Class 5 appeared to contain potential regulatory mutants for solventogenesis and many Class 1, 2 and 3 mutants also did not sporulate.

Transposon insertion-flanking chromosomal DNA from the mutants was amplified by inverse PCR using divergent primers for Tn916. Briefly, chromosomal DNA from the mutants was cut with *Hind* III, which cuts once in Tn916, and ligated at a dilute DNA concentration to promote circularization of individual fragments. The divergent primers Tn916*tet*, which was

specific for the *tetM* gene, and Tn916L2, which was specific for the left end of the transposon, were then positioned to amplify chromosomal DNA that flanked the left end of the transposon insertion. Single bands of 150 bp to 2.5 kb were amplified from eleven of fourteen Class 1 mutants while the other three Class 1 mutants yielded two bands each.

The PCR products were recovered from agarose gels and digoxigenin-dUTP was incorporated, either by random-primed labeling or by reamplification of the fragments, to produce hybridization probes. Southern blots of restriction enzyme-digested wild-type chromosomal DNA were hybridized using the PCR product probes and the hybridization patterns produced by the different probes were compared. Probes from mutants 224 and 243 hybridized to apparently identical wild-type chromosomal DNA fragments for each of seven different restriction enzyme digests. This suggested that the transposon insertions in those two mutants were in the same region of the chromosome. The results for probes from all of the other mutants suggested that the transposon insertions were in different regions of the chromosome in each case.

The hybridization probes were also used to isolate clones from a library of wild-type chromosomal DNA fragments. *C. acetobutylicum* ATCC 824 chromosomal DNA was partially digested with *Sau3A* and fragments from 4 kb to 6 kb were recovered from an agarose gel. The fragments were ligated into the plasmid pBluescript® KS+ that had been cut with *BamH*I and treated with calf intestinal phosphatase. The ligation products were electroporated into *Escherichia coli* DH5 α and transformants were selected for ampicillin resistance. Fifteen pools of transformants that contained about 400 white colonies each were screened in dot blot assays using the hybridization probes from each Class 1 mutant. Potential clones were identified by colony hybridizations of transformants that were plated from dot blot-positive pools. Two clones, that contained a 3 kb and a 4 kb insert, were isolated using a probe generated from mutant 202 and a probe from mutant 174 identified two clones that contained very small inserts. These clones will be used in genetic complementation experiments with the Class 1 mutants.

2. Delivery of plasmids into *Clostridium acetobutylicum* by electroporative transformation.

Initial studies using the pTV1 plasmid which carries *cat* and also Tn917 transposon with *erm*, showed that transformation was unsuccessful. We decided to develop the electroporation procedure testing a variety of plasmids and a variety of strains of *Clostridium acetobutylicum* in order to ensure transfer of selected cloned DNA into mutant organisms and to transfer Tn917 into the chromosome. We have made excellent progress in achieving the optimal conditions for electroporative transformation of strains of *C. acetobutylicum* with the *B. subtilis* plasmid pIM13. This work is outlined in the enclosed abstract and preprint. We are also in the process of developing shuttle vectors based on pIM13 since recent success with this sort of construction has been reported (Azeddoug, H., Hubert, J., and Reyssset, G. J. Gen. Microbiol. 138:1371-78, 1992).

3. Conjugal transfer of plasmids and chromosomal mobilization in *C. acetobutylicum*.

This is a new approach to genetic analysis in *C. acetobutylicum* based on the observation by O.R. Torres, R.Z. Korman, S.A. Zahler, and G.M. Dunny (Mol. Gen. Genet. 225: 395-400, 1991) that the *E. faecalis* transposon, Tn925 causes general mobilization of chromosomal genes in *B. subtilis*.

In preliminary experiments, we have shown definite mobilization of plasmid pIM13 carried by *E. faecalis* CG110 :: Tn916 during conjugation with *C. acetobutylicum* strain 824. As

shown in table 1, this mobilization is low compared with conjugative transfer of Tc^R and Em^R when the Tn916 is carried on plasmid pAM180. Also this mobilization depends upon a high donor to recipient ratio.

We are planning to develop this conjugative mobilization technique as a means of identifying complementation groups of the various Tn916-mutants of *C. acetobutylicum* described in paragraph 1 above.

Publications

(a) Papers

1. Contag, P.R., Williams, M.G., and Rogers, P. (1990). Cloning of a lactate dehydrogenase gene from *Clostridium acetobutylicum* B643 and expression in *Escherichia coli*. Applied Environ. Microbiol. **56**:3760-3765.
2. Rogers, P. and Gottschalk, G. (1992). Biochemistry and regulation of acid and solvent production in clostridia. In D. Woods, ed., "The Clostridia and Biotechnology", Chapter 2, Butterworth, Inc. (In press).
3. Reding, H.K., Mattsson, D.M., and Rogers, P. (1992). Transformation of intact *Clostridium acetobutylicum* with plasmid pIM13 DNA by electroporation. Applied Environ. Microbiol. (submitted for publication).

(b) Abstracts

4. Contag, P.R. and Rogers, P. (1991). The cloning and expression of an NAD-dependent alcohol dehydrogenase from *Clostridium acetobutylicum* B643. (Abstract) Annual Meeting ASM.
5. Mattsson, D., Rast, T. and Rogers, P. (1992) Transformation of *Clostridium acetobutylicum* NC1B6444 with a chimeric *C. acetobutylicum*-*Escherichia coli* plasmid. (Abstract) Annual Meeting ASM.
6. Reding, H.K., Mattsson, D.M., and Rogers, P. (1993). Electrotransformation of *Clostridium acetobutylicum*. (Abstract) Annual Meeting ASM.
7. Mattsson, D.M. and Rogers, P. (1993). Analysis of transposon-insertion mutants of *Clostridium acetobutylicum* ATCC 824. (Abstract) Annual Meeting ASM.

Appendix

Copies of Publications 1-3 and Abstracts 6, 7.

Reprints + Preprints removed

Table 1. Conjugative transfer of pIM13 from
Enterococcus faecalis to
Clostridium acetobutylicum 824

<i>E. faecalis</i> strain	Donor to Recipient ratio	Efficiency of Conjugation (transconjugants/ 10 ⁶ recipients)		
		Tc ^r	Em ^r	Em ^r /Tet ^r
CG110/pIM13 ¹ (pellet mating ²)	2.1/1	6.3	0	—
	11/1	3.3	0	—
	21/1	4.5	0.28	1/16
	31/1	8.1	1.5	1/5.4
	43/1	5.4	2.6	1/2.1
	66/1	1.8	1.5	1/1.2
CG110 ³ (filter mating ⁴)	5/1	53	0	—
CG110/pIM13	3.5/1	120	1	1/120
CG180 ⁵	7.5/1	120	100	1/1.2

1 CG110/pIM13: chromosomal Tn916 with a Tet^r marker and *B.subtilis* plasmid pIM13 with Em^r marker.

2 Pellet mating technique described in methods.

3 CG110: no pIM13.

4 Filter mating technique described in methods.

5 CG180: both Tn916 and Em^r marker carried on plasmid pAM180.

END

**DATE
FILMED**
5/03/93