**U.S. DEPARTMENT OF ENERGY
OFFICE OF BASIC ENERGY SCIENCES
FINAL PROJECT REPORT**

DOE/ER/13319--T1
DE93 008134

Project Identification Information

INSTITUTION: Stanford University
PRINCIPAL INVESTIGATOR: David D. Pollard
AWARD NUMBER: DE-FG03-85ER13319-M009
AMOUNT: \$298,623
PERIOD OF SUPPORT: 01/01/85 to 06/30/92 (including no-cost extensions)
TITLE: Structure and Emplacement of the Magmatic Intrusions Beneath
the Valley of Ten Thousand Smokes

Summary of Completed Project

The goal of this project was to provide constraints on the subsurface geometry of the vent, feeder conduit(s), and possible intrusive bodies in the Novarupta Basin, Valley of Ten Thousand Smokes (VTTS), Katmai National Park, Alaska. This research was designed to support the Katmai Drilling Project, a consortium involving the DOE, NSF, other government agencies, and several universities. Our work aimed at providing information to help guide site selection and identify targets for the drilling program. However, our research also has contributed to the understanding of silicic volcanic systems in general. Understanding the subsurface geometry of the vent is essential for accurately modeling the present thermal regime in the vent region and the eruption dynamics. Determining the origin of surficial fractures indicated the effect of compaction and consolidation of tephra and orientation of the depositional surface on fracture formation.

Our research plan included:

- (1) completion of a detailed topographic-structural map of surface fractures including spatial distribution and relative displacements;
- (2) numerical modeling studies that related surface fractures to major subsurface structures; and
- (3) interpretation of the origin of surface fractures in light of these model studies.

This project provided full support for the Ph.D. thesis research of Peter Wallmann and partial support for the Ph.D. thesis research of Jake Lowenstern.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Yra

Technical Information

Our research objectives for the Valley of Ten Thousand Smokes (VTTS) and the Novarupta Basin were as follows:

- (1) complete a detailed topographic–structural map of surface fractures in the Novarupta basin including spatial distribution and relative displacements;
- (2) carry out numerical modeling studies that relate surface fractures to major subsurface structures; and
- (3) interpret the origins of the surface fractures in light of these model studies.

These objectives have been completed and the results are summarized in the following paragraphs and in the publications listed in the next section of this report.

New structural data from the Novarupta basin, site of the largest volcanic eruption of this century (1912), has provided limits for the location of magma chambers associated with this eruption. A project of geologic mapping of surficial and bedrock structures in the vent region of the 1912 eruption was undertaken. Landslide scarps, arcuate grabens, a monoclinial fold, and truncated ridges circumscribe the Novarupta basin, marking the inferred outer rim of the vent. A set of radial fissures crosses the southern margin of the basin, striking $\sim 140^\circ$, subparallel to the dominant bedrock joint set. These fissures and joints, along with the local plate-motion vector and the inferred regional stress orientation, are consistent with a feeder dike propagating from a reservoir beneath Trident volcano to the eruptive vent.

Field work discovered two areas in the Novarupta basin which display the effects of long term compaction of the fallback tephra filling the vent. A morphologic monocline which extends for some 500 m along the western edge of the basin is interpreted to have formed by deposition and compaction of tephra over a gently sloping vent rim. The other area displaying continued compaction is located along the northern edge of the basin. Examination of photographs from the Griggs' expeditions in 1916–1919 and topographic profiling have shown that a new drainage is forming there and that the old drainage has been stranded above the present valley bottom. Both of these structures, along with the densely welded fallback tephra that was ejected during the second day of the eruption, can be used to constrain compaction models of the vent region.

Selected photographs from the Griggs' expeditions were re-photographed using the old pictures as guides in relocating the original camera location. These pictures show that the degradation of scarp faces has been significant, although the scarps do not appear to have retreated very far from their original locations. One important result from this comparison is that grabens on the Turtle, a fractured, topographic high in the vent region, may have formed initially as open fractures which then experienced dip–slip movement to become faults. Modeling suggests that

the topographic expression of the Turtle is consistent with the emplacement of an intrusion the size of Novarupta Dome 275 m beneath the top of the Turtle.

Four sets of structures are interpreted to reflect the extent of the Stage I vent at Novarupta basin: 1) truncated topography along the northern and southern basin margins; 2) arcuate grabens on the eastern margin; 3) translational block slides and gulls on the north and south margins; and 4) a monoclinical fold in the tephra blanket along the western margin. These structures define a roughly elliptical vent with axial lengths of 2.5 and 3 km. Other surficial structures include radial slides and fissures, perhaps related to passage of a dike which fed Novarupta Dome. Topographic cross sections, the inferred depth to a possible magma chamber, and the surficial fractures suggest the sub-surface geometry is a flared vent with steeply dipping walls to the north and south, and more gently dipping walls to the east and west.

A large faulted mound of tephra, the Turtle, was produced northeast of the Stage II vent during the last eruption phase. The dominant structures on the Turtle are two graben systems which intersect at nearly right angles. Two conceptual models are proposed for these: 1) doming of the tephra by a shallow igneous intrusion; and 2) compaction of the tephra pile. Numerical modeling of these processes, combined with field observations, suggest the Turtle formed by inflation of a shallow igneous intrusion.

Measurements of extensional strains across fissures and grabens in the Novarupta basin and VTTS have been used to constrain numerical models for compaction of the ash. These models provide estimates for the bedrock topography beneath the compaction structures. Apparently, the western margin of the basin is underlain by a convex-upward, flared vent. In contrast, the geometry of the vent along the eastern margin is inferred to be concave upward and steep-sided. This geometry probably represents truncation of a pre-existing topographic saddle by growth of the flared vent.

We compare the structure and intrusive history of the West Mageik Lake sill complex to possible conduits for magma transport during the 1912 eruption at the nearby Valley of Ten Thousand Smokes (VTTS), on the Alaska Peninsula, to shed light on some of the enigmatic aspects of the 1912 eruption, including its unusual feeder system and eruptive sequence. At West Mageik Lake, Late Tertiary (?) rhyolitic magma propagated along a roughly east-west trend as sills that followed bedding planes and through-going joint sets in the Jurassic Naknek siltstone. This demonstrates the feasibility of arc-parallel transport of rhyolitic magma as sills at shallow crustal levels.

The chemical and isotopic compositions of sills at West Mageik Lake are contrasted with high-silica rhyolite erupted in 1912 at the nearby Valley of Ten Thousand Smokes to gain insight into the petrogenesis of evolved magmas in the Aleutian arc. A high-silica rhyolitic sill has an ϵ_{Nd} value (+8.8) similar to an adjacent dacite porphyry stock, but the elevated Al, Sr, and

$^{87}\text{Sr}/^{86}\text{Sr}$ of the rhyolite suggest that fractionation of a parental dacite magma was accompanied by significant contamination by Sr-rich sediments, probably of Paleozoic age. The chemical and isotopic character of the West Mageik Lake rhyolite suggest that the system was of small volume, making it easily affected by crustal contamination. The fact that the 8 km³ of high-silica rhyolite erupted at the Valley of Ten Thousand Smokes shows only a slight Sr isotopic shift from associated dacite and andesite suggests it was part of a larger system that could assimilate small masses of crust without having its composition significantly affected.

Presentations of Technical Information

Research completed under this grant has been presented at several different scientific meetings including annual meetings of the American Geophysical Union and special meetings organized by the sponsors of the Katmai Drilling Project. References for the abstracts and extended abstracts from these presentations are given below:

- Wallmann, P.C. and Pollard, D.D., 1983, Surficial deformation in the Novarupta Caldera, Valley of Ten Thousand Smokes, Katmai National Park Alaska: *Eos*, Trans AGU, v. 69, p. 1472.
- Lowenstern, J.B. and Wallmann, P.C., 1989, The West Mageik Lakes sill: analogue for the feeder of the 1912 eruption at the Valley of Ten Thousand Smokes?: *New Mexico Bureau of Mines & Mineral Resources*, Bulletin 131, IAVCEI abstracts, p. 169.
- Wallmann, P.C., and Pollard, D.D., 1989, Investigations of the mechanics of surficial deformation in the Novarupta basin, Valley of Ten Thousand Smokes, Katmai National Park, Alaska: *New Mexico Bureau of Mines & Mineral Resources*, Bulletin 131, IAVCEI abstracts, p. 269.
- Wallmann, P.C. and Pollard, D.D., 1989, New insights on the structure of the Novarupta basin, Valley of Ten Thousand Smokes, Katmai National Park, Alaska: *Eos*, Trans AGU, v. 70, p. 1412.

Research results from this grant have been published in four papers to date. Copies of these four papers are attached to this report. References for the four papers are:

- Eichelberger, J.C., Ballard, S. Carrigan, C.R., Goodliffe, A., Hildreth, W., Iwatsubo, E., Kasameyer, P.W., Keith, T.E.C., Kienle, J., Papike, J.J., Pollard, D.D., Stone, D.B., Wallmann, P.C., Ward, P.L., Wilt, M., and Yount, M.E., 1990, Geophysical expedition to Novarupta volcano, Katmai National Park, Alaska: *Eos*, v. 71, p. 733-755.

- Wallmann, P.C., Pollard, D.D., Hildreth, W., and Eichelberger, J.C., 1990, New structural constraints on magma chamber locations at the Valley of Ten Thousand Smokes, Katmai National Park, Alaska: *Geology*, v. 18, p. 1240-1243.
- Lowenstern, J.B., Wallmann, P.C., and Pollard, D.D., 1991, The West Mageik Lake sill complex as an analogue for magma transport during the 1912 eruption at the Valley of Ten Thousand Smokes, Alaska: *Geophysical research letters*, v. 18, p. 1569-1572.
- Lowenstern, J.B. and Mahood, G.A., 1991, Petrogenesis of high-silica rhyolite on the Alaska peninsula: *Geophysical research letters*, v. 18, pp. 1565-1568.

Two additional papers will emerge from Peter Wallmann's thesis. The first paper originally was submitted to the Geological Society of America Bulletin and currently is being revised. The second will be submitted to the Bulletin of Volcanology. The references for these papers in preparation are:

- Wallmann, P.C., and Pollard, D.D., in review, Surficial fractures in the vent region for the 1912 eruption in the Valley of Ten Thousand Smokes and their relationship to the subsurface vent geometry.
- Wallmann, P.C., and Pollard, D.D., in preparation, Origin of a faulted topographic dome in the vent for the 1912 eruption at Novarupta, Alaska, *Bulletin of Volcanology*.

A copy of the manuscript on surficial fractures is attached to this report.

The two Ph.D. theses supported by this project are:

- Peter C. Wallmann, 1991, Structure and subsurface vent geometry of the Novarupta Basin, Valley of Ten Thousand Smokes, Katmai National Park, Alaska: Ph.D. Thesis, Stanford University.
- Jacob B. Lowenstern, 1992, The behavior and distribution of volatile metals and gases in two silicic magmatic systems: Pantelleria, Italy and The Valley of Ten Thousand Smokes, Alaska: Ph.D. Thesis, Stanford University.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

**DATE
FILMED**

3 / 30 / 93