DOE/ER/40694---2

DE93 012934

NUCLEAR STRUCTURE MODELS: APPLICATIONS AND DEVELOPMENT

Progress Report

July 1, 1992 - April 30, 1993

Paul B. Semmes

Tennessee Technological University

Cookeville, TN 38505

April, 1993

PREPARED FOR THE U. S. DEPARTMENT OF ENERGY

UNDER GRANT NUMBER DE-FG05-92ER40694

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

Preface	1
1. Rotational Structure in Odd-Odd Nuclei	2
1.1. Characterizing the Nuclear Shape in ^{82}Y	2
1.2. Signature Inversion in ^{120}Cs - Evidence for a Residual p-n Interaction	6
2. Collective Structure in Odd-A Nuclei	6
2.1 Evolution of Deformation in the $A \approx 130$ Region: ^{133}Nd	6
2.2 Electromagnetic Properties of ^{181}Ir : Evidence of β Stretching?	9
3. Particle-Core Coupling Models	9
References	10
Appendix I: Personnel	11
Appendix II: Publications and Seminars	12
Appendix III: Reprints and Preprints - <i>Removed</i>	13

PREFACE

This report summarizes the progress made during the period July 1, 1992 to April 30, 1993 on Grant No. DE-FG05-92ER40694. The study of rotational structure at moderate spins in odd-odd nuclei (Section 1) has continued into the $A \approx 80$ region. Recent experimental data for the positive parity band in ^{82}Y , including absolute BM1 and BE2 values, are rather well described with a "prolatish" triaxial shape and a $\pi g_{9/2} \otimes \nu g_{9/2}$ configuration. Little effect from a proton-neutron interaction has been found so far. Current studies of odd mass nuclei (Section 2) are mainly focussing on low spin, low energy states in the $A \approx 130$ region as a probe to the evolving collectivity. These projects involve direct collaboration with experimental colleagues. Finally, a view of the microscopic foundation of particle-core coupling models approached from the equation of motion method has been presented (Section 3).

The secretarial and accounting duties for this grant have been performed by Gloria Julian of the TTU Physics Department, and her efforts are greatly appreciated. Discussions with Drs. Ingemar Ragnarsson (LTH, Lund, Sweden), Ragnar Bengtsson (UNISOR, Oak Ridge, TN and Lund, Sweden), and Witold Nazarewicz (Joint Institute for Heavy Ion Research, Oak Ridge) are gratefully acknowledged.

1. Rotational Structure in Odd-Odd Nuclei

Two applications of the 2-quasiparticles + triaxial rotor model are discussed below. The first is the result of a new collaboration with the Florida State University experimental high-spin group, and the second is the conclusion of earlier work.

1.1 Characterizing the Nuclear Shape in ^{82}Y

(with S. Tabor, J. Döring and P. Womble)

The study of odd-odd nuclei has continued into the $A \approx 80$ region. This region is particularly interesting because the fairly small number of particles (compared to $A \approx 150$ for instance) leads to strong variations of the nuclear shape with particle number, configuration and angular momentum [1]. A rotational band built on the $\pi g_{9/2} \otimes \nu g_{9/2}$ configuration has been observed in a number of nuclei, often exhibiting a clear signature inversion at low spins, i.e., a change of the phase of the energy staggering. Recent experimental studies of ^{82}Y have identified both signatures of this band from $I = 6$ to 13. Both $\Delta I = 2$ E2 and $\Delta I = 1$ M1/E2 γ -ray transitions were observed, and $B(\text{M1}; I \rightarrow I-1)/B(\text{E2}; I \rightarrow I-2)$ ratios were deduced. Furthermore, lifetimes were measured for many of these states giving absolute BM1 and BE2 values. Cranking calculations based on the Woods-Saxon potential suggest a rather gamma soft structure for this band, with the minimum energy at a collective triaxial shape, $\beta \approx 0.24$, $\gamma \approx -30^\circ$. Simple cranking estimates of the BE2 values from the calculated deformation agree reasonably well with the data.

In order to explore the observed behavior of the positive parity band in greater detail, calculations were made with the 2-qp + triaxial rotor model. The current version uses the Modified Oscillator potential, and the standard κ , μ parameters of ref. [2] were used, but with the proton parameters for $N=3,4$ adopted for both the protons and neutrons. This choice gives a very similar ordering and structure of the Nilsson orbitals as the Woods-Saxon potential [1]. Calculations were made with the quadrupole deformation $\epsilon = 0.23$ and a variety of γ values ranging from 0° (prolate shape) to 60° (oblate shape). These values correspond to the collective sector in the cranking model; however, note that a single core shape is used for all states in this particle-rotor model. The pairing gaps and Fermi levels were calculated from the BCS method, and the core moments of inertia were calculated from the deformations and pairing gaps. For

the results presented here, no proton-neutron residual interaction was used.

The calculated energy splitting $(E_I - E_{I-1})/2I$ is shown in Fig. 1a with different curves drawn for different values of γ . No attenuation of the Coriolis terms is included. For the prolate shape, a pronounced staggering with odd spins favored is present for $I \geq 10$ or so, and the phase of the staggering changes around $I = 9$, in rough agreement with the data. This basic feature is found for all the shapes considered, even the oblate. The calculated neutron Fermi level is near the middle of the $g_{9/2}$ shell and is closest to the $\Omega = 5/2$ orbital for both prolate and oblate shapes, while the proton Fermi level is below the $g_{9/2}$ shell and is closest to the $\Omega=3/2$ orbital for the prolate shape and the $\Omega=7/2$ orbital for the oblate shape. In all cases, the calculated wave functions are very mixed in the strong coupling basis, with the largest component about 15% (amplitude squared) for $I \approx 10$. Attenuating the Coriolis interaction produces only modest improvement as illustrated for the prolate shape in Fig. 1b.

The $B(E2; I \rightarrow I-2)$ and $B(M1; I \rightarrow I-1)$ transition strengths calculated for the different values of γ are shown in Fig. 2 and correspond to the states of Fig 1a. The general magnitude of the calculated $B(E2)$ values agrees rather well with the data for the prolate shape but not for the oblate, for which the intrinsic quadrupole moment is smaller in magnitude ($Q_0 = +2.33b$ and $Q_0 = -1.84b$, respectively). Note that the observed staggering observed in the $B(E2)$ values is obtained for a "prolatish" triaxial rotor with $\gamma \approx 25^\circ$. Also, the experimental $B(M1)$ strengths are well described in the calculations for a prolate or "prolatish" shape, except for the point at $I = 13$. Again the oblate shape is clearly ruled out by the calculations, although the same phase of the staggering occurs for oblate as well as prolate shape. The calculated $B(M1)$ values are very large (approximately $1.7 \mu_N^2$) for $I = 8$ and $\gamma = 40^\circ$ or 60° and are not shown on the graph. Finally, negative $E2/M1$ mixing ratios are obtained for all shapes considered here, and so agreement with the data for this sign cannot be taken as conclusive evidence for a prolate shape as proposed for ^{84}Y [3]. In particular, the calculated mixing ratio for the $11+ \rightarrow 10+$ transition is negative but small enough in magnitude that it agrees within error bars for prolate, oblate and triaxial shapes.

Thus, qualitative agreement with the experimental positive parity band is obtained for prolate or "prolatish" triaxial shapes. The observed energy splitting including the change of phase at low spins is qualitatively reproduced,

Fig. 1: Calculated energy staggering for the positive parity band in ^{82}Y . The upper graph shows the γ dependence while the lower graph shows the effect of Coriolis attenuation in the case of the prolate shape ($\xi = 1$ is no attenuation). The solid points are the experimental data.

Fig. 2: Calculated E2 and M1 transition strengths as a function of the triaxiality.

while the strong staggering in the $B(M1)$ strengths is very well described. The observed staggering in the $B(E2)$ values suggests a triaxial core shape, with $\gamma \approx 25^\circ$. These results agree qualitatively with the cranking calculations.

Finally, preliminary calculations including a residual interaction do not give any essential changes or improvements. This last observation is interesting because previous studies (including ^{120}Cs , see Appendix III) have found significant effects, although the positions of the proton and neutron Fermi levels relative to the high j -shells is somewhat different here than in the previous studies. These questions are being studied further.

These results are included in a paper submitted for publication (see Appendix II) that also contains the experimental results.

1.2 Signature Inversion in ^{120}Cs - Evidence for a Residual p-n Interaction

Described in a previous report, this work has been published and a copy is given in Appendix III.

2. Collective Structure in Odd-A Nuclei

Applications of the 1-quasiparticle + triaxial rotor model are given below. The first emphasizes low spin, low energy structure of very neutron deficient rare earth isotopes while the second is the conclusion of earlier work on moderate spin states in ^{181}Ir .

2.1 Evolution of Deformation in the $A \approx 130$ Region: ^{133}Nd

(with J. Breitenbach and J. Wood)

Low energy collective modes are a characteristic feature of nuclei, and the observed evolution of collectivity as a function of particle number, angular momentum etc., provide strict tests of nuclear structure models. For mean field models, stably deformed nuclei provide clear tests because the collective properties can be described in terms of a few deformation parameters and the positions of Nilsson orbitals (observed as rotational bandheads) can be related to the underlying single particle parameters of the mean field. The very neutron deficient rare earth nuclei form a new region of deformation that is now coming under scrutiny. Theoretical predictions for the onset and development of collectivity have been in place for a number of years including calculations with different mean field descriptions. For example, Folded Yukawa calculations [4]

indicate a sudden transition to well deformed shapes at $N \approx 73$ in the Pm and Sm isotopes, and Woods-Saxon calculations [5] indicate that the transition from the near spherical ^{142}Sm to well deformed ^{134}Sm proceeds via triaxial shapes.

Recent radioactive decay studies at UNISOR have identified for the first time a number of low lying states in ^{133}Nd , including a $7/2+$ ground state, $1/2+$ and $9/2-$ bandheads and a few low spin negative parity states. According to both refs. [4] and [5], ^{132}Nd is well deformed ($\beta \approx 0.28$) while ^{134}Nd is more transitional in character ($\beta \approx 0.22$). Calculations are underway to determine the shape of ^{133}Nd (and indeed whether a fixed intrinsic shape can provide an adequate description) and to help identify the Nilsson orbitals. A sample calculation is shown in Fig. 3, based on the Woods-Saxon version of the particle + triaxial rotor model. The Woods-Saxon potential is the same as used in ref. [5], and the deformation is the tabulated value for ^{134}Nd in that paper. Qualitative agreement with the data is found, including the $7/2+$ ground state and low lying $1/2+$ and $9/2-$ bandheads. The dominant Nilsson orbital for these states are the $[404]7/2$ for the $7/2+$ and the $[514]9/2$ (part of the $h_{11/2}$ shell) for the $9/2-$ state. At the prolate deformation shown, the $1/2+$ is nearly pure $[400]1/2$, and the low spin negative parity states are mainly due to the $[541]1/2$, which is mainly from the $h_{9/2}$ spherical orbital. However, the positions of these bandheads are not so well reproduced, and the $5/2+$ bandhead expected at very low energy has not yet been identified. Experimentally, the $1/2+$ state is found much closer to the $7/2+$ ground state, and so far this can be achieved in the calculations only with a sizable γ deformation ($\gamma \approx 25^\circ$).

The goal of this work is not just to obtain a good fit to ^{133}Nd , but to develop an understanding of the systematic features of the $N = 73$ and $N = 71$ isotones. The lower Z nuclei are better characterized experimentally, and a systematic description is crucial for testing the mean field models. Preliminary calculations for $^{127,129}\text{Ba}$ also indicate sizable γ deformations are needed to produce the low lying $1/2+$ states, but achieving good descriptions of the $h_{11/2}$ bands and the known moments in these nuclei (including the magnetic moment of the $1/2+$ states) as well remains difficult. A current study of the high spin properties (especially neutron crossing frequencies) of nuclei in the $A \approx 130$ region suggest that the neutron $h_{11/2}$ shell may be somewhat lower than given in the usual Nilsson diagrams [6]. This suggestion is being investigated with the Modified Oscillator version of the particle + triaxial rotor model.

Fig 3. Particle + triaxial rotor calculation for ¹³³Nd using the Woods-Saxon potential. The pairing gap and Fermi level are derived from the BCS equations and the 2+ energy of the core is estimated from Grodzins' rule. The calculated states are organized into rotational bands and the dominant Nilsson orbital is indicated for each band. Some calculated moments and relative M1/E2 strengths are also indicated.

2.2 Electromagnetic Properties of ^{181}Ir : Evidence of β Stretching?

This work has been accepted for publication, and a copy of the preprint is given in Appendix III.

3. Particle-Core Coupling Models

As part of the Workshop on Nuclear Structure Models held in Oak Ridge, TN, March 16-25, 1992, a written contribution was prepared and has now been published. A copy is given in Appendix III.

REFERENCES

- [1] W. Nazarewicz et al., Nucl. Phys. **A435** (1985) 397.
- [2] T. Bengtsson and I. Ragnarsson, Nucl. Phys. **A436** (1985) 14.
- [3] S. Chattopadhyay et al., Phys. Rev **C47** (1993) R1.
- [4] G. A. Leander and P. Möller, Phys. Lett. **110B** (1982) 17.
- [5] B. D. Kern et al., Phys. Rev. **C36** (1987) 1514.
- [6] J-y Zhang, private communication, 1993.

APPENDIX I

Personnel

FACULTY

Paul B. Semmes

COLLABORATORS

J. Breitenbach	Georgia Institute of Technology
J. Döring	Florida State University
S. Tabor	Florida State University
P. Womble	Florida State University
J. Wood	Georgia Institute of Technology

APPENDIX II

PUBLISHED ARTICLES

"Signature Inversion in ^{120}Cs - Evidence for a Residual p-n Interaction", B. Cederwall, F. Liden, A. Johnson, L. Hildingsson, R. Wyss, B. Fant, S. Juutinen, P. Ahonen, S. Mitarai, J. Mukai, J. Nyberg, I. Ragnarsson and P. B. Semmes, Nucl. Phys. **A542** (1992) 454.

"Particle-Core Coupling Models", P. B. Semmes in International Workshop on Nuclear Structure Models, R. Bengtsson et al. (eds.) (World Scientific, 1992), p.287.

ARTICLES IN PRESS

"Electromagnetic Properties of ^{181}Ir : Evidence of β -Stretching?", U. Garg, W. Reviol and P. B. Semmes, Phys. Rev. C.

ARTICLES SUBMITTED FOR PUBLICATION

"Nuclear Structure of odd-odd ^{82}Y ", P. C. Womble, J. Döring, T. Glasmacher, J. W. Holcomb, G. D. Johns, T. J. Petters, M. A. Riley, V. A. Woods, S. L. Tabor and P. Semmes, submitted to Phys. Rev. C.

SEMINARS AND LECTURES

"Invisible Decay of Superdeformed Bands", Sigma Xi Research Award Lecture, Tennessee Technological University, January 14, 1993.

"Signature Splitting in Odd-Odd Nuclei and the Proton-Neutron Residual Interaction", presented to the Nuclear Physics group at Florida State University, February 5, 1993.

END

**DATE
FILMED**

6 / 11 / 93