1 of 1

40/3-30-94 JS②

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2978
UC-420

PPPL-2978

FUSION POWER PRODUCTION FROM TFTR PLASMAS
FUELED WITH DEUTERIUM AND TRITIUM

BY

J.D. STRACHAN, H. ADLER, P. ALLING, ET AL.

MARCH, 1994

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced from the best available copy.
Available in paper copy and microfiche.

Number of pages in this report: 16

DOE and DOE contractors can obtain copies of this report from:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
(615) 576-8401.

This report is publicly available from the:

National Technical Information Service
Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4650

Fusion Power Production from TFTR Plasmas Fueled with Deuterium and Tritium*

J.D. Strachan, H. Adler, P. Alling, C. Ancher, H. Anderson, J.L. Anderson,⁽¹⁾ D. Ashcroft, Cris W. Barnes,⁽¹⁾ G. Barnes, S. Batha,⁽²⁾ M.G. Bell, R. Bell, M. Bitter, W. Blanchard, N.L. Bretz, R. Budny, C.E. Bush,⁽³⁾ R. Camp, M. Caorlin, S. Cauffman, Z. Chang,⁽⁴⁾ C.Z. Cheng, J. Collins, G. Coward, D.S. Darrow, J. DeLooper, H. Duong,⁽⁵⁾ L. Dudek, R. Durst,⁽⁴⁾ P.C. Efthimion, D. Ernst,⁽⁶⁾ R. Fisher,⁽⁵⁾ R.J. Fonck,⁽⁴⁾ E. Fredrickson, N. Fromm, G.Y. Fu, H.P. Furth, C. Gentile, N. Gorelenkov,⁽⁷⁾ B. Grek, L.R. Grisham, G. Hammett, G.R. Hanson,⁽³⁾ R.J. Hawryluk, W. Heidbrink,⁽⁸⁾ H.W. Herrmann, K.W. Hill, J. Hosea, H. Hsuan, A. Janos, D.L. Jassby, F.C. Jobs, D.W. Johnson, L.C. Johnson, J. Kamperschroer, H. Kugel, N.T. Lam,⁽⁴⁾ P.H. LaMarche, M.J. Loughlin,⁽⁹⁾ B. LeBlanc, M. Leonard, F.M. Levinton,⁽²⁾ J. Machuzak,⁽⁶⁾ D.K. Mansfield, A. Martin, E. Mazzucato, R. Majeski, E. Marmar,⁽⁶⁾ J. McChesney⁽⁵⁾, B. McCormack, D.C. McCune, K.M. McGuire, G. McKee,⁽⁴⁾ D.M. Meade, S.S. Medley, D.R. Mikkelsen, D. Mueller, M. Murakami,⁽³⁾ A. Nagy, R. Nazikian, R. Newman, T. Nishitani,⁽¹⁰⁾ M. Norris, T. O'Connor, M. Oldaker, M. Osakabe,⁽¹¹⁾ D.K. Owens, H. Park, W. Park, S.F. Paul, G. Pearson, E. Perry, M. Petrov,⁽¹²⁾ C.K. Phillips, S. Pitcher,⁽¹³⁾ A.T. Ramsey, D.A. Rasmussen,⁽³⁾ M.H. Redi, D. Roberts,⁽⁴⁾ J. Rogers, R. Rossmassler, A.L. Roquemore, E. Ruskov,⁽⁸⁾ S.A. Sabbagh,⁽¹⁴⁾ M. Sasao,⁽¹¹⁾ G. Schilling, J. Schivell, G.L. Schmidt, S.D. Scott, R. Sissingh, C.H. Skinner, J.A. Snipes,⁽⁶⁾ J. Stevens, T. Stevenson, B.C. Stratton, E. Synakowski, W. Tang, G. Taylor, J.L. Terry,⁽⁶⁾ M.E. Thompson, M. Tuszewski,⁽¹⁾ C. Vannoy, A. von Halle, S. von Goeler, D. Voorhees, R.T. Walters, R. Wieland, J.B. Wilgen,⁽³⁾ M. Williams, J.R. Wilson, K.L. Wong, G.A. Wurden,⁽¹⁾ M. Yamada, K.M. Young, M.C. Zarnstorff and S.J. Zweben.

Plasma Physics Laboratory, Princeton University, P.O. Box 451, Princeton, New Jersey 08543

⁽¹⁾*Los Alamos National Laboratory, Los Alamos, New Mexico*

⁽²⁾*Fusion Physics and Technology, Torrance, California*

⁽³⁾*Oak Ridge National Laboratory, Oak Ridge, Tennessee*

⁽⁴⁾*University of Wisconsin, Madison, Wisconsin*

⁽⁵⁾*General Atomics, San Diego, California*

⁽⁶⁾*Massachusetts Institute of Technology, Cambridge, Massachusetts*

⁽⁷⁾*TRINITI, Moscow, Russia*

⁽⁸⁾*University of California, Irvine, California*

⁽⁹⁾*JET Joint Undertaking, Abingdon, United Kingdom*

⁽¹⁰⁾*JAERI Naka Fusion Research Establishment, Naka, Japan*

⁽¹¹⁾*National Institute for Fusion Science, Nagoya, Japan*

⁽¹²⁾*Ioffe Physical-Technical Institute, Russia*

⁽¹³⁾*Canadian Fusion Fuels Technology Project, Toronto, Canada*

⁽¹⁴⁾*Columbia University, New York, New York*

(Received ?? February 1994)

MASTER

*Work supported by U.S. DOE Contract No. DE-AC02-76-CHO-3073.

ABSTRACT

Peak fusion power production of 6.2 ± 0.4 MW has been achieved in TFTR plasmas heated by deuterium and tritium neutral beams at a total power of 29.5 MW. These plasmas have an inferred central fusion alpha particle density of $1.2 \times 10^{17} \text{ m}^{-3}$ without the appearance of either disruptive MHD events or detectable changes in Alfvén wave activity. The measured loss rate of energetic alpha particles agreed with the approximately 5% losses expected from alpha particles which are born on unconfined orbits.

PACS numbers: 28.52.Cx, 52.25.Fi, 52.55.Pi

Most previous experiments in magnetic fusion research have been conducted with hydrogen or deuterium plasmas, even though first generation fusion reactors are expected to operate with equal concentrations of deuterium (D) and tritium (T). One consequence of fueling with D-T is that since the $d(t,n)\alpha$ fusion reactivity is much higher than the D fusion reactivity, more fusion reactions occur and a significant population of the charged fusion products can be created. Potentially, collective phenomena can arise from the 3.5 MeV alpha population influencing their confinement as well as the global plasma stability and energy balance. The Tokamak Fusion Test Reactor (TFTR) has performed initial D-T experiments and has achieved energetic alpha densities which are about 0.2% of the plasma ion density which is about 1/3 - 1/2 of the fraction expected in reactors. TFTR is the second tokamak to use T [1] and the first to use equal concentrations of D and T. A separate paper [2] describes the changes in plasma heating and confinement observed with T and alpha particles present, whereas this paper discusses the energetic ion behavior, the measurements of the fusion reactions, and the search for alpha-induced instabilities.

The primary goal of these experiments was to produce a plasma with greater than 5 MW of peak D-T fusion power. With 29.5 MW of neutral beam heating, a D-T fusion power of 6.2 ± 0.4 MW was produced with a corresponding 14 MeV neutron emission rate of up to $2.2 \pm 0.2 \cdot 10^{18} \text{ sec}^{-1}$. For the high power D-T experiments, TFTR was operated in the supershot regime [3] with 2.0 MA plasma current, 5.0 T toroidal magnetic field, 2.52 m major radius, and 0.87 m minor radius. The D-T was fueled by operating one to eight of the twelve beam sources in pure tritium. There were 7 T discharges at 23-30 MW of total neutral beam power including one with 10% of the beam power in tritium, one with 100% tritium beam power and 5 with 40%-65% of the beam power in tritium. For comparison, 42 similar D and 6 trace T plasmas were produced at the same machine conditions.

The neutron emission rates and yields were measured with fission chambers [4], silicon surface barrier diodes [5], spatially collimated 4 He recoil proportional counters [6] and ZnS scintillators [7], and a variety of elemental activation foils [8]. The activation foils, 4 He counters, and silicon diodes can discriminate between 14 MeV D-T and 2.5 MeV D-D neutrons. The other detectors cannot discriminate between D-D and D-T neutrons, but are more sensitive to the latter. An absolute calibration of the fission chambers, proportional counters and scintillators was performed using an *in situ* 14 MeV neutron generator [9]. The estimated absolute accuracy of each calibration is about $\pm 10\%$ to $\pm 25\%$ while the statistical deviation of all available calibration data, is $\pm 7\%$. The quoted fusion power is the weighted mean of the calibrated signals with the $\pm 7\%$ standard deviation.

The measured D-D neutron emission from D plasmas was 10-15% greater than that calculated from the measured density and temperature profiles and a calculated beam deposition profile (by the steady-state code SNAP [10] and the time dependent code TRANSP [11]). In D-T plasmas, the measured D-T neutron emission is up to 10% less than the calculated value (Fig. 1 and 2a). These discrepancies are comparable to the expected uncertainty in the codes ($\pm 15\%$) and the magnitude of the individual neutron measurement uncertainty, which is based upon separate calibrations for 2.5 MeV and 14 MeV neutrons. The D-D neutron emission from D plasmas [12, 13] and the D-T neutron emission (for D-T plasmas with approximately equal D and T beam heating) increased strongly with the total plasma energy content. The measured neutron source strength normalized to the plasma energy content (Fig. 3) displays a broad maximum near equal injected powers of D and T while the 100% T beam shot had about 65% of the maximum D-T neutron emission. This rate was used to assess the hydrogenic influx and transport models in the SNAP and TRANSP codes and is consistent

with a significant (40%) concentration of thermal D in the plasma core during pure-T injection.

The D-T neutron emission (Fig. 2a) and the TRANSP calculation of it degrades from the peak levels which occur at about 3.5 s. This degradation often occurs in D supershots and can be correlated with several factors including: a) the temporary reduction in the applied neutral beam power (Fig. 2a) due to source faults that occurred at 3.5 and 3.65 s, b) the appearance of $m/n = 4/3$ MHD activity which began at about 3.4 s [14], or c) the increase in the D influx from the limiters throughout the beam heating [15]. The D-T plasmas reported here actually degrade less than comparable D-D plasmas.

The pitch angle and energy of the fusion alpha particles escaping from the plasma was measured with a scintillation detector [16] located near the vacuum vessel wall 90° below the midplane in the ion-gradient-B drift direction. In quiescent low-power plasmas, the relative alpha particle loss decreased by a factor of about 4 between 0.6 MA and 1.8 MA, in rough agreement with the calculated variation in the first-orbit loss to this detector (Fig. 4). The total loss of D-T alphas in the high-power D-T plasmas at 2.0 MA was also roughly consistent with expectations based on the simple first-orbit loss model calibrated by the signal at 0.6 MA (where all the trapped alpha particles are lost). In particular, the alpha loss fraction did not increase significantly between the lowest and highest power D-T shots at 2.0 MA (Fig. 4) while the alpha source rate increased by more than a factor of 10. This indicates that the alpha particles were not being lost as a result of instabilities driven by the alpha particle pressure itself.

This TFTR plasma regime can be unstable to the Toroidal Alfvén Eigenmode (TAE) [17] in the time following the beam heating when the alpha pressure remains high (Fig. 2c) and the average alpha velocity reduces to the Alfvén velocity. However, the plasma fluctuation activity [18, 19] in the TAE range of

frequencies (250 kHz during beam heating, rising to 500 kHz after injection) were the same for D-T and D plasmas (Fig. 5). The level of broad band fluctuations measured by a microwave reflectometer [19] indicates that the upper limit of possible TAE activity is $\tilde{n}/n = 5 \times 10^{-5}$. These levels are one to two orders of magnitude below the trapped-particle-driven TAE modes seen during ion cyclotron heating in TFTR or the beam driven TAE modes detected at low field and high density [20, 21] by the same diagnostics in D plasmas. The behavior of the background turbulence observed during and after the beam heating is very similar in both D-T and D plasmas. There is no indication that the mode amplitude is enhanced by the presence of the alpha population during or following termination of the beam heating (Fig. 2c).

In conclusion, the initial D-T experiments on TFTR produced 6.2 MW of fusion power. The resulting energetic alpha population caused neither detectable anomalous alpha particle losses nor observable instabilities. The TFTR fusion yield can be increased through increases in the T beam voltage and injected neutral beam power as well as through reductions in the hydrogen influx from the limiters, or improvement in the gross energy confinement time by lithium wall conditioning [22].

The authors appreciate the contributions from the technical staff of the Princeton Plasma Physics Laboratory under the leadership of R. Davidson.

Acknowledgment

This work was supported by the United States Department of Energy under contract number DE-AC02-76-CHO-3073.

- [1] JET Team, Nuclear Fusion **32**, 187 (1992).
- [2] R.J. Hawryluk *et al.*, to be published.
- [3] J.D. Strachan *et al.*, Phys. Rev. Lett. **58**, 1004 (1987).
- [4] H. Hendel *et al.*, Rev. Sci. Instrum. **61**, 1900 (1990).
- [5] H.H. Duong and W.W. Heidbrink, Nuclear Fusion **33**, 211 (1993).
- [6] J.S. McCauley and J.D. Strachan, Rev. Sci. Instr. **63**, 4536 (1992).
- [7] L. Johnson, Rev. Sci. Instrum. **63**, 4517 (1992).
- [8] C.W. Barnes *et al.*, Rev. Sci. Instr. **61**, 3190 (1990).
- [9] A.L. Roquemore *et al.*, Proc. 15th IEEE Symposium on Fusion Eng., Hyannis, MA (Oct. 1993).
- [10] H.H. Towner *et al.*, Rev. Sci. Instrum. **63**, 4753 (1992).
- [11] R. Budny *et al.*, Nuclear Fusion **32**, 429 (1992).
- [12] J.D. Strachan *et al.*, Nuclear Fusion **33**, 991 (1993).
- [13] M.G. Bell *et al.*, Plasma Phys. Contr. Fus. Res., Vol. 1, p. 27, (IAEA, Vienna, 1989).
- [14] Z. Chang *et al.*, PPPL-2941 (1993), submitted to Nuclear Fusion.
- [15] J.D. Strachan, PPPL-2933 (1993), submitted to Nuclear Fusion.
- [16] S.J. Zweben *et al.*, to be published in Phys. Fluids B (1993).
- [17] C.Z. Cheng *et al.*, Plasma Phys. and Contr. Nucle. Fus. Res. 1992, Vol II, 51 (IAEA, Vienna, 1993).
- [18] E.D. Fredrickson, K.M. McGuire, A. Cavallo *et al.*, Rev. Sci. Instrum. **59**, 1797 (1988).
- [19] E. Mazzucato and R. Nazikian, Physics Rev. Lett. **71**, 1840 (1993).
- [20] G. Taylor, M.G. Bell, H. Biglari, M. Bitter *et al.*, Phys. Fluids B **5**, 2437 (1993).
- [21] K-L. Wong, R.J. Fonck, S.F. Paul, D.R. Roberts *et al.*, Phys Rev. Lett. **66**, 1874 (1991).
- [22] J. Snipes *et al.*, J. Nuclear Mat. **196-198**, 686 (1992).

FIG. 1. The neutron emission calculated by the equilibrium code SNAP (open squares) and calculated by the time-dependent code TRANSP (solid squares) as a function of the neutron source strength measured by the TFTR fission detectors.

FIG. 2. Time evolution of the plasma with the highest D-T neutron emission. (a) The beam power (in units of 10 MW), the measured D-T neutron emission (in units of 10^{18} sec^{-1}) (solid line) and the TRANSP calculated value (dashed line), including the calculated contributions of beam target, beam beam, and thermonuclear reactions. (b) The measured collection rate of energetic ($> 1 \text{ MeV}$) escaping alphas (solid line), the calculated central alpha particle density (in units of 10^{17} m^{-3}) and the calculated detector signal (by TRANSP) due to classical first orbit loss. (c) The TRANSP-calculated central alpha pressure ($\beta_\alpha = 2\mu_0 p_\alpha/B^2$) (in units of 10^{-3}), the ratio of alpha velocity to Alfvén velocity, and the measured amplitude of the Mirnov signal at the TAE frequency range taken from several D-T and D comparison plasmas.

FIG. 3. D-T neutron emission at peak stored energy divided by the square of the plasma energy content plotted as a function of the fraction of the beam heating coming from T beams. The solid curve is the expected dependence if the fueling of the plasma were entirely from the beams. The dashed curve corresponds to one-half of the fueling from the beams and one-half from the walls (deuterium only).

FIG. 4. The measured alpha particle loss rate to the vessel bottom normalized by the global neutron source strength as a function of plasma current. The shaded region is the calculated alpha first-orbit loss for this location. The x points are low power, quiescent plasmas and the circles are the high power D-T plasmas.

FIG. 5. High frequency fluctuation data taken immediately after the beam heating for D-T and D plasma . The shaded region shows the approximate lower bound on the previously observed TAE mode. (a) Amplitude spectra of an outboard Mirnov coil signal showing a weak peak near the expected TAE frequency. (b) Reflectometer power spectra at a major radius of 2.92 m (the plasma magnetic axis is at 2.63 m).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. I. Kawakami, Hiroshima Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Librarian, Naka Fusion Research Establishment, JAERI, JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	Research Info. Ctr., National Instit. for Fusion Science, JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	Prof. S. Tanaka, Kyoto Univ., JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	Library, Kyoto Univ., JAPAN
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Prof. Dr. I.C. Nascimento, Instituto Fisica, Sao Paulo, BRAZIL	S. Mori, Technical Advisor, JAERI, JAPAN
Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL	Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Documents Office, Atomic Energy of Canada Ltd., CANADA	Dr. G.S. Lee, Korea Basic Sci. Ctr., KOREA
Ms. M. Morin, CCFM/Tokamak de Varennes, CANADA	J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Prof. J. Teichmann, Univ. of Montreal, CANADA	Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Prof. T.W. Johnston, INRS-Energie, CANADA	Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. C.R. James,, Univ. of Alberta, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Dr. P. Lukác, Komenského Univerzita, CZECHO-SLOVAKIA	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
The Librarian, Culham Laboratory, ENGLAND	Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Dr. L. Stenflo, Univ. of UMEA, SWEDEN
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS	Library, Royal Inst. of Technology, SWEDEN
P. Mähönen, Univ. of Helsinki, FINLAND	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
Prof. E. Economou, Univ. of Crete, GREECE	Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. I Apparatus, USSR
Ms. C. Rinni, Univ. of Ioannina, GREECE	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA	Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Librarian, International Center for Theo. Physics, ITALY	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY	Librarian, Max-Planck-Institut, W. GERMANY

DATE

FILMED

4 / 18 / 94

END