Progress Report/ Plan

February 28, 1992

DOE/ER/61134--1

DE92 011033

DoE Grant No. De-FG02-91ER61134

Title: "Construction of a human genome library composed of multimegababase acentric chromosome fragments"

PI/PD: M.J. Lane, Dep't. of Medicine, SUNY-HSC at Syracuse, Syracuse, NY

co-PI/PDs: P.J. Hahn, Dep't of Radiology, SUNY-HSC at Syracuse

J.C. Hozier, Dep't Medical Genetics, FIT, Melbourne, Florida

Overall, given the existence of three established PI/PDs the overall grant management strategy has and will continue to be allowance of the individual PIs to execute those aspects of the project with which they are most comfortable. During the procedural development phase this has served to minimize micromanagement while maximizing the initiative of each investigator. This approach has led to a natural division of labor with surprisingly little experimental overlap. It is the opinion of all involved that the best tactical approach to library construction is to create a unique fragment library of the genome (at 5X density) followed by screening to assign clones to individual human chromosomes. It had been the hope that individual chromosome libraries could be created up-front. This turns out to be neither cost nor labor efficient. The experimental progress of each laboratory is broken out below based on write-ups from the respective PI/PD.

Hozier Laboratory (Florida Institute of Technology)

Progress (3/15/91 - 3/15/92): We have been developing in situ analytical techniques which make rapid cataloguing of human chromosomal DNA content in acentric fragment containing mouse cells via Cot I DNA painting both fairly efficient and effective. The methodology developed is automated and employs a Fisher Code-On slide handling robot and the methodology is requisite for assuring that the human DNA fragments in the fusion cells are maintained unattached to the chromosomes of the mouse recipient cells in an efficient manner. Additional experiments have given us confidence that human repetitive element PCR strategy, followed by painting of the PCR fragments to human metaphase spreads, will allow efficient cytogenetic localization of human fragments retained in the human:mouse hybrids.

Plan (3/15/92 - 3/15/93): We are continuing to develop microcloning techniques which would allow recovery of human DNA in many hybrids on a single chromosome slide. We feel this will dramatically improve analytical efficiency during the library evaluation stage (expectation of number of cell lines in the library of between 2 and 10 thousand). We will continue to improve human repeat PCR efficiency for this purpose as well.

Hahn (Radiology, SUNY-HSC at Syracuse)

Progress (3/15/91 - 3/15/92): We have been concerned with both refining the library construction strategy and physical recovery of acentric human DNA fragments produced from individual cell lines. Our experiments indicate that 1) mouse recipient cells which harbor human DNA fragments produced using human co-transfect (using both a G418 and DHFR plasmid simultaneously) do not lose unselected additional human DNA in an efficient manner; 2) long range pulse field gels allow reasonable recovery of acentric fragment DNA which can be used to construct YAC sublibraries 3) it is not feasible to amplify the introduced DHFR gene without interference from the endogenous mouse DHFR genes and 4) an electroporation strategy can be used to efficiently generate high

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

complexity (with respect to marker insertion position) donors for library construction.

Plan (3/15/92 - 3/15/93): We will continue our efforts to isolate a DHFR- recipient line (EMT-6). Some progress toward this objective has been made already. Refine methodologies for picking, screening and maintaining large numbers of mouse clones. The Hozier group is collaborating in the maintenance function. Focus on strategies to eliminate unselected human DNA carry through in the primary fusion step.

Lane (SUNY-HSC at Syracuse)

Progress (3/15/91 - 3/15/92): We have been focused in several areas: 1) evaluating the feasibility of our proposed mapping strategy for ordering acentric fragments, 2) evaluating the use of retroviral insertion as a means of producing high complexity of insertion position human cell donors for use in primary fusion to the mouse recipient, 3) direct construction of several specific acentric fragments using retrovirally marked human chromosomes retained in monochromosome hybrids and 4) assessing the feasibility of multiple fusion rounds for the purpose of "laundering" unselected human DNA efficiently. Our experiments indicate that 1) an anomalously high number of rearrangements take place in the immediate vicinity (± 10 kb) of the three specific retroviral insertion points we have examined following recovery of DNA in the recipients. This is in contrast to introduction of a marker by transfection strategies (see Hahn above). 2) Multiple fusion rounds significantly reduce the unselected human DNA carry through and are desirable for producing mouse lines with a human acentric DNA fragment as its only material.

Plan (3/15/92 - 3/15/93): We will continue to examine the acentric fragment mapping strategy initially proposed. We have diverted some effort toward understanding the rearrangement frequency encountered with a retroviral marker but should complete this near term. We will continue to refine our multiple fusion round approach for getting rid of unselected human fragments with the aim of minimizing the total radiation dose received in the region surrounding the introduced selectable marker to retain DNA integrity flanking the marker for several million base pairs. We will switch to transfection as a marker introduction strategy. Our focus here is distinct from that of Hahn (above) as we are attempting to create a single copy acentric fragment library (no DHFR introduction).

Overall: In general all three investigators are proceeding along a learning curve which will allow primary library construction and characterization to begin during this fiscal period. The characterization (cloning/in situ /mapping) is and will continue to receive far more attention (read time) than we originally thought necessary based on the object lesson taught with human YAC libraries (eg chimera/recombination issues). It is our objective to define what the library we are building is composed of, rather than learn post-construction that our library is of limited utility in the context of the genome project as a whole. In the coming year these issues will be defined and we feel that taking a cautious construction approach is the best course. -MJL

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

**DATE
FILMED**

5 / 14 / 92