**Progress Report, 1992: "Multiple measurement of the coupling between benthic carbon fluxes and bioturbation activity during the "spring bloom"
DOE No. DE-FG02-92ER61464**

Robert C. Aller, Josephine J. Aller, J. Kirk Cochran, and Cindy Lee
Marine Sciences Research Center
SUNY at Stony Brook

DOE/ER/61464--1

Research 1992-93:

DE93 012327

In accordance with the research plan outlined in our original proposal, we began monitoring indicators of plankton production (Chl-a, cell counts) in surface waters of Long Island Sound from the Port Jefferson / Bridgeport Ferry on Dec. 3, 1992. The project was given the acronym: PULSE. Samples were originally obtained weekly. Sample frequency was increased to 2 - 3 X per week in mid-January 1993 in anticipation of the "spring bloom". We have continued water sampling at this latter frequency to the present. In contrast to past years no dramatic bloom has occurred to date although a significant pattern of increasing chl-a began during the first week of April (Fig. 1). We anticipate that the bloom will be complete by mid to late April. This has been a particularly cold year based on comparison of 1993 bottom water temperatures and previously reported patterns from LIS, possibly reflecting climatic conditions which delayed the bloom. Suspended matter samples, cell samples, and dissolved nutrient samples have been stored in all cases for future analysis.

During December 1992 we also established a permanent bottom sampling site in central Long Island Sound. The site is located in muddy sediments at 41:10.00N 72:57.41W in ~15.5 m water and was originally marked by a surface buoy and acoustic pingers at the bottom. The surface buoy was absent in January, presumably hit by barge or freighter traffic through the area. Dr. Creighton Wirick of BNL made available two shallow water acoustic releases which we refurbished and used to subsequently remark the site. The releases are traded on successive sampling cruises. Exact marking of the site is needed to revisit *in situ* mixing experiments set-up and tended by divers (M. Gerino, M. Green, R. Aller) as outlined below. Bottom sampling has taken place at five times to date with a sixth sampling due 12 April (sampling times indicated in Fig. 1). The frequency of sampling will increase from every 3 weeks to every 2 weeks following direct evidence of the maximum bloom production.

Typical sampling includes the following: (1) surface and bottom water sampling for suspended matter, cell counts, chlorophyll-a, nutrients; (2) triplicate box cores for macro-, meio-, and microfauna analysis; (3) triplicate subcores for O₂ microelectrode profiles; (4) box core for radiochemical analyses (²³⁴Th, ⁷Be); (5) box core subcores for benthic fluxes under aerated and nonaerated conditions (O₂, ΣN, Mn⁺⁺, ΣCO₂); (6) Br⁻ tracer flux core incubation for biogenic irrigation rates; (6) box core for laboratory based luminophore tracer experiments (particle mixing); (7) box core subcores for solid phase and pore water solute profiles (carbonate system, nutrients, Ca, Mg, Sr, F, Mn, Fe) at 0.25

MASTER

- 1 cm intervals; (8) box core subcores for chloropigment profiles and reaction rate incubation experiments.

In addition to remotely obtained samples, we have established diver-tended *in situ* luminophore tracer experiments. Core tubes (15 cm diameter) are placed flush into the bottom, luminophores (fluorescent particles of three different grain sizes) are introduced to the surface sediment by divers, and the exact site marked by small underwater bouys. On the following cruise, previously deployed cores are recovered and luminophore depth penetration profiles measured. A new set of *in situ* cores are set up at each sampling time. We also plan to implace particle tracers without core tubes present to check for mixing artifacts due to the presence of a horizontal barrier within the sediment (core tube wall). These *in situ* mixing experiments are being compared to otherwise identical laboratory core incubations with luminophore tracers (cores are kept in a running sea water table). Non steady state luminophore mixing rates will be compared with mixing rates determined from radionuclides and chloropigment profiles as outlined in the original proposal.

We have also obtained on each sampling cruise: water column samples for plankton biochemical and physiological studies by Dr. J. LaRoche (BNL) and sediment core samples for organic geochemical (protein) analyses by Dr. L Mayer (U. Maine). Samples for sedimentary lipids are being supplied to Dr. S. Wakeham (Skidaway).

Examples of some of the data we are obtaining are given in Figures 1 - 6. We are now in the middle of the most intense field sampling program of the project and most efforts are being spent simply acquiring and processing samples. Analyses of unstable constituents are done immediately, other samples are stored whenever possible.

Research in 1993-94:

We plan to continue a modest field sampling program at increasingly longer intervals (1 month) through the summer in order to follow the remnants of the bloom and to further experimentally compare coherency between mixing tracers in the laboratory and field during warmer periods. The numerous stored samples will also be processed. In particular, biological samples (foraminifera, bivalves, gastropods) for calculation of CaCO_3 precipitation and dissolution processes and bioturbation studies (total benthos) will be worked up (very labor intensive). Solid phase analyses (Mn, CaCO_3 , C_{org} , N, S), including total chloropigments on frozen samples will be completed. Initial chloropigment analyses are made on unfrozen samples, the two treatments allow differentiation of chloropigment pools (bound and free). Pore water solute analyses will be completed. Luminophore profiles will be determined. Data interpretation and analysis will begin in earnest. We anticipate more than enough analytical work within the context of the project to remain busy, and perhaps overwhelmed.

Fig. 1. Surface water chlorophyll-a (HPLC) in central Long Island Sound during December 1992 - April 7, 1993. The 'spring bloom' is building steadily but with substantial fluctuations. The relatively large excursion on March 15, followed a major storm. Storm-related 'mini' blooms are apparently not unusual in LIS.

Chlorophyll-a in surface water

Fig. 2. Chlorophyll-a in surface water of LIS in January - March 1992 compared to December 1992 - April 1993 (Pulse project period). The times of bottom station sampling during 1992-1993 period are indicated by arrows. Note that all bottom sampling periods are pre-bloom for 1993.

Fig. 3. Pore water F^- and Fe^{++} profiles from Pulse 4 and 5. The Fe^{++} peak is relatively deep in the sediment during this time of year because of low decomposition rates and temperatures. O_2 penetrates 5 - 7 mm at this time. The clear separation between F^- and Fe demonstrates that F^- release in this case is not associated with destruction of an Fe -oxide carrier phase, as suggested in the literature, but most likely with $CaCO_3$ dissolution. Quantification of F^- , Mg^{++} , and Sr^{++} along with the carbonate system parameters should allow detailed modeling of dissolution at this time.

Fig. 4. ^{234}Th profiles from Pulse 1 samples as measured by both gamma counting and β counting. Background subtraction results in a large uncertainty for the gamma data, but overall good agreement is found between the two methods for activity gradients and inventories. Both methods are being used in most cases.

Core IS2
22 days in situ conditions
collected on 03/01/93 (PULSE 4)

Fig. 5. Luminophore profiles from retrieved in situ experiment cores, Pulse 4. The upper 0.5 cm initially contained luminophores.

Experimental core M1
 In vitro conditions
 33 days (from 12/16/92 to 01/17/93)

Fig. 6. Luminophore tracer profiles from laboratory incubated cores. In this case temperatures were several degrees above *in situ*.

END

DATE
FILMED

5 / 20 / 93