DOE/ER/40461--5

DE93 009765

**THEORETICAL NUCLEAR REACTION AND STRUCTURE
STUDIES USING HYPERONS AND PHOTONS**

Progress Report

January 1992 - December 1992

Stephen R. Cotanch

Department of Physics

North Carolina State University

Raleigh, North Carolina 27695-8202

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

yes

PREPARED FOR THE U. S. DEPARTMENT OF ENERGY
UNDER GRANT NUMBER DE-FG05-88ER40461

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This report details research progress and results obtained during the 12 month period from January 1992 through 31 December 1992. The research project, entitled "Theoretical Nuclear Reaction and Structure Studies Using Hyperons and Photons", is supported by grant DE-FG05-88ER40461 between North Carolina State University and the United States Department of Energy. In compliance with grant requirements the Principal Investigator, Professor Stephen R. Cotanch, has conducted a research program addressing theoretical investigations of reactions involving hyperons and photons. The Principal Investigator has devoted to this program 50% of his time during the academic year and 100% of his time in the summer. This percent effort will continue for the remaining period of the grant. The new, significant research results are briefly summarized in the following sections. Recent progress has also been discussed in the renewal/continuation grant proposal just submitted to the Department of Energy. Finally, full, detailed descriptions of completed work can be found in the project publications which are listed at the end of this progress report.

ELEMENTARY HADRON STRUCTURE

This project seeks to further the understanding of hadron structure within the framework of Quantum Chromodynamics (QCD) and QCD based models. In the most recent funding period several significant findings have been obtained. As detailed in our publications^{1,9,10} (see end of this report) we (with C. R. Ji and A. Szczepaniak) have performed a comparative study of meson properties using three different relativistic models: our previously extended Dziembowski's model (Ji and Cotanch, Phys. Rev. D**41**,2319(1990)); our generalization^{1,9} of the Hamiltonian light-front approach developed by Coester and Chung; a new completely covariant null plane approach¹⁰. All models have been applied to the open flavor pseudo-scalar (π , K, D), vector (ρ , K*) and pseudo-vector (a_1) mesons and we have obtained a good description of their static properties. Major findings include: 1) all models can describe meson charge radii and decay constants but only the Hamiltonian and covariant null plane approaches can properly account for

the form factor at high Q^2 ; 2) only the extended Dziembowski's model can give a camel shaped (double-peaked) quark distribution amplitude when plotted as a fraction of the meson momentum carried by the quarks; 3) the key ingredient for the success of all models is relativity as a non-relativistic calculation gives poor results; 4) while covariance is desirable it does not appear essential for simple quark models with harmonic oscillator confinement; 5) similar to point 4, the use of light-cone variables, as oppose to conventional instant form variables, does also not appear to be necessary for a successful phenomenological description using harmonic oscillator wavefunctions. Concerning the camel shape quark distribution amplitude for the pion, this is a controversial issue since this signature is not predicted by all QCD lattice and QCD sum-rule calculations. In fact more recent and improved lattice calculations using the unquenched approximation do not obtain a dip in the distribution amplitude. Further, the extended Dziembowski model can only obtain this shape in an inconsistent fashion by using a model parameter which gives an unphysically small decay constant. It is clear that not only does this matter need further study but also that the quark distribution amplitude is very useful for comparing alternative QCD models.

ELECTROMAGNETIC PRODUCTION OF STRANGENESS

This project addresses systems involving the strange quark (hyperons) and investigates hyperon electromagnetic production and radiative capture processes. During the past funding period a number of new, important research results were generated. In collaboration with R. Williams and C. R. Ji, we completed^{2,3,12} our analysis of the world's kaon electroproduction data below 2.5 GeV for $p(e,e' K^+)Y$ with $Y = \Lambda, \Sigma$ and $\Lambda(1405)$ using our unified model based on Quantum Hadrodynamics which incorporates crossing symmetry and duality. We now have a good, consistent and comprehensive description of both cross sections and spin observables for photo and electroproduction of the low-lying strange baryon spectrum as well as the limited crossed channel kaon

radiative capture $p(K^-, \gamma)Y$ branching ratios. We have also completed our theoretical study of the novel $p(K^-, e^+e^-)Y$ reaction and find that this reaction can provide important, unique meaningful information about the kaon form factor in the time-like region. Because of the three-body final state there is no constraint on the 4-momentum transferred by the time-like photon ($Q^2 > 4 m_e^2 \sim 0$) which permits the vector mesons ρ , ω and ϕ to be kinematically accessible and generates a dramatic five order of magnitude enhancement in the cross section. Not only does this large, unambiguous resonance signature facilitate experiments with even low intensity kaon beams it also permits a direct measurement of the ϕ hadronic coupling and therefore an assessment of the strangeness content of the proton. This work has been submitted for publication⁴.

Our results will not only impact upon CEBAF and SLAC but also the kaon facilities (BNL, KEK and, if built, KAON).

MEDIUM ENERGY PHOTONUCLEAR REACTIONS

Significant, interesting results have been obtained in this continuing project during the previous funding period. As detailed in recent invited talks¹¹ and publications^{5,6,7,8} we have performed a large-scale continuum shell model calculation for (γ, N) and (N, γ) , $N = p$ and n , reactions at low and medium energies spanning the Δ isobar region. This is a microscopic calculation in which a realistic many-body Hamiltonian H for the A baryon system is diagonalized in a large Banach space (not Hilbert since the continuum basis states are not square integrable) spanned by products of continuum single-particle states and known bound many-body eigenfunctions of the $A-1$ system. The Hamiltonian contains a realistic effective interaction with finite range, spin, isospin and tensor components. The code is flexible and can accommodate a variety of two-body forces including those based on multi-boson exchange. Diagonalizing H in this space generates a large set of coupled inhomogeneous integro-differential equations in two variables (r

and r' due to non-localities) to be solved, subject to appropriate scattering boundary conditions, for the unknown partial-wave scattering functions. All non-localities are retained in the radial kernels due to wavefunction antisymmetrization, Pauli blocking and exchange. Since the large number (up to 100 currently) of scattering channels rigorously account for loss of flux from the elastic channel there is no direct need for phenomenological absorption and H therefore remains Hermitian which assures proper wavefunction orthogonality that is especially important in photonuclear reactions. Further, the code rigorously respects unitary and gauge-invariance.

During the last funded period, with the assistance of DOE Headquarters, we obtained supercomputer time at Livermore which, in conjunction with the time available on our local North Carolina Cray Y-MP (we have submitted a grant and were funded for 250 hours of time by the North Carolina Supercomputer Center) allowed us to complete several large-scale calculations. In addition to the $^{12}\text{C}(p,\gamma)$ analysis^{5,6,7} at 98 MeV summarized in last years report we have just completed a constrained set of calculations now explicitly including Δ isobar exchange currents for the ground and first excited 2^+ state. Because carbon is not a closed shell nucleus we needed wavefunctions more sophisticated than that provided by the 1 particle - 1 hole basis and this in turn required a significantly larger number of coupled channels. Including the Δ exactly triples the number of channels to 144 which is still too taxing for even our present computing facilities. We therefore performed a constrained calculation by selectively including various Δ channels. Interestingly, we found relatively small Δ contributions. While this should still be confirmed by performing more complete calculations, the result is not too surprising since such effects should only become import around 250 MeV and above. We also did calculations at 176 MeV, without Δ channels, and predicted angular distributions which were impressively confirmed by subsequent measurements at the Svedberg Laboratory in Uppsala, Sweden. Related we completed a full Δ channels analysis for $^{16}\text{O}(\gamma, N)$ for $N = p$ and n at energies between 150 and 250 MeV. For the

first time a simultaneous description has been obtained between theory and experiment (all previous (γ, n) calculations could not reproduce the cross section magnitude). Most significantly, we find a large sensitivity to cut-off parameters in our two-body effective interaction at large momentum transfers. These parameters serve to regulate the unphysical singularities in our Yukawa force. Since there should be no sensitivity, which is the case at lower momentum transfers, there is a clear need to include a more physically correct short-ranged component of the interaction which is currently in process. This work has been submitted for publication⁸.

It is important to stress that our university is implementing a large-scale network of IBM RISC workstations with special hardware/software for providing and optimizing actual parallel computing. With appropriate additional funding we can upgrade our system to participate in this program which will permit production runs comparable and superior to the supercomputers. This is mandatory for not only investigating higher energies but for also properly including all exchange currents.

This project supports experimental investigations at the Bates Laboratory, the Indiana University Cyclotron, the Svedberg Laboratory in Uppsala, Sweden and future CEBAF measurements.

PAPERS PUBLISHED IN PRESS OR SUBMITTED IN THE REPORT PERIOD

1. "Light-Cone Quark Model Axial Vector Meson Wavefunction", C.-R. Ji, P. L. Chung and S. R. Cotanch, Phys. Rev. D**45**,4214 (1992).
2. "Hyperon Electroproduction in a Crossing and Duality constrained Model", R. A. Williams, C.-R. Ji and S. R. Cotanch, Phys. Rev. C**46**,1617(1992).
3. "Kaon Electromagnetic Production and Radiative Capture Near Hyperon Thresholds", S. R. Cotanch, R. A. Williams and C.-R. Ji, Physica Scripta, in press(1992).
4. "Kinematically Accessible Vector Meson Resonance Enhancements in $p(K^-, e^+e^-)\Lambda, \Sigma, \Lambda(1405)$ ", R. A. Williams, C.-R. Ji and S. R. Cotanch, submitted to Phys. Rev. Lett. (1992).
5. "Radiative Proton Capture to Individual States in ^{12}C ", B. Höstad, E. Nilsson, J. Thun, S. Dahlgren, S. Isaksson, G. S. Adams, C. Landberg, T. Bright and S. R. Cotanch, Phys. Lett. B**276**, 294 (1992).
6. "Exclusive Radiative Proton Capture to ^{12}C ", B. Höstad, S. Isaksson, E. Nilsson, J. Thun, G. S. Adams, C. Landberg, T. Bright and S. R. Cotanch, Nucl. Phys. A, in press(1992).
7. "Radiative Proton Capture and Exchange Currents in Light Nuclei" B. Höstad, S. Isaksson, E. Nilsson, J. Thun, G. S. Adams, C. Landberg, T. Bright and S. R. Cotanch, Proceedings of the 1992 International Nuclear Physics Conference, Wiesbaden, Germany, July (1992).
8. "Simultaneous Microscopic Description of $^{16}\text{O}(\gamma, p)$ and $^{16}\text{O}(\gamma, n)$ Including Δ Degrees of Freedom, T. B. Bright and S. R. Cotanch, submitted to Phys. Rev. Lett. (1992).

INVITED PAPERS

9. "Simple Light-Cone Quark Model Description of Meson Properties" 13th International Conference on Few Body Problems in Physics, Adelaide, Australia (Jan., 1992).
10. "Relativistic Quark Model Analysis of Flavored Mesons", Tenth Nordic Meeting on Intermediate and High Energy Nuclear Physics, Gräfvallen, Sweden (Jan., 1993).
11. "Continuum Shell Model Approach to Medium Energy Photonuclear Reactions", Symposium on Nuclear Reactions to Commemorate the Sixty-fifth Birthday of G. R. Satchler, Oak Ridge, Tennessee (April, 1992).
12. "Kaon Electromagnetic Production and Radiative Capture Near Hyperon Thresholds", Second International Conference on Particle Production Near Threshold, Uppsala, Sweden (Aug., 1992).

Also, there were three abstracts detailing contributed papers at several meetings. These references are available upon request.

END

DATE
FILMED

5 / 7 / 93

1

2

3

4