THEORETICAL STUDIES IN NUCLEAR STRUCTURE

Progress Report

for Period December 1, 1991 - November 30, 1992

E. R. Marshalek

University of Notre Dame
Notre Dame, IN 46556

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

December 1992

Prepared for

THE U. S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-91ER40640

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED
yes

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights.

ABSTRACT

Two topics of current interest - strongly anharmonic collective vibrations and collective rotation about a tilted axis - have been related through a theorem proposed by the principal investigator. This theorem, which will be called the Bifurcation Theorem, states that if a system has an axis of rotational symmetry in a static mean-field approximation, then self-consistent cranking about this axis gives rise to symmetry-breaking bifurcations that occur for critical cranking frequencies Ω_c given by $\Omega_c = \omega_\mu / K_\mu$, where ω_μ is the small-oscillation (RPA) frequency of any normal mode carrying $\hbar K_\mu (\neq 0)$ units of angular momentum along the symmetry axis. The bifurcations represent families of anharmonic oscillations built on the axially symmetric mean field. For certain values of K_μ , the solutions appear as rotations about an axis that is tilted with respect to the principal axis of the density distribution. The theorem has been tested on some simple models in preparation for more realistic calculations that are planned for the near future.

I. A CRANKING MODEL FOR COLLECTIVE ANHARMONIC VIBRATIONS

The question of anharmonicity of nuclear collective vibrations has long been one of the most challenging in nuclear structure, both experimentally and theoretically. The most direct experimental evidence for anharmonicity, provided by vibrational states with two or more phonons, has been rather elusive. However, Ani Aprahamian, an experimental nuclear spectroscopist at Notre Dame has recently proposed eighteen candidates for $K = 4$ two-phonon γ vibrations of deformed nuclei.¹ There is also some evidence (though controversial) for a two-phonon octupole level in ^{208}Pb ² and even a suggestion of two-phonon giant resonances in spherical nuclei.³ Since the energy ratios of the proposed two-phonon γ -vibrational states to the known one-phonon states usually deviate considerably from the value of 2, significant anharmonicity seems to be present. While the theorist's toolkit contains a plethora of different microscopic methods for calculating anharmonic corrections,⁴ each method has its pros and cons and all are difficult to implement. Fortunately, I have been able to devise an especially simple method for finding periodic solutions of mean-field equations that is applicable to collective vibrations carrying nonzero angular momentum for small or large amplitudes. This method employs the self-consistent cranking model (SCCM), which has played a central role in elucidating rotational states of high angular momentum. However, the proposal here is to use the SCCM to describe states that are normally regarded as *vibrational* rather than rotational in character. In particular, the SCCM is being applied for the first time to models of vibrating deformed nuclei.

The idea of applying the SCCM to vibrational modes was introduced over twenty years ago by Marshalek and Sabato,^{5,6} who showed that the SCCM could be used not only to derive

the RPA for even-even spherical nuclei, but also the leading-order perturbative anharmonic corrections to the excitation energies as well as the static quadrupole moments and $B(E2)$'s for the aligned n -phonon quadrupole excitations with spins $L = 2n$. Since the essential ingredient of this *spherical cranking model* is the presence of an *axis of symmetry* in the uncranked mean field, the model can be extended to axially symmetric deformed nuclei, thereby providing a new tool with some special advantages for calculating the n -phonon γ -vibrational bandheads with spin projections $K = 2n$ on the axis of symmetry and, possibly, other bandheads as well. Exploratory calculations have been carried out for the interacting boson model (IBM) and the volume-conserving harmonic-oscillator shell model, which can be interpreted as a rough model of giant quadrupole resonances. Unlike the earlier treatment, the present application avoids perturbation theory, which often leads to problems with small resonance denominators. Therefore, it should be applicable to large-amplitude vibrations.

A simple motivation for the new approach is provided by the physical picture of a collective vibration with $K \neq 0$ in an axially deformed nucleus as a surface wave, an instantaneous distortion traveling around the equilibrium axis of symmetry. Since such a mode has a two-fold time-reversal degeneracy, it is always possible to choose the relative phases of the two waves to produce a uniform rotation about this axis. Relative to a uniformly rotating coordinate system, the motion can be viewed as a stationary equilibrium state. The SCCM is nothing more than the condition for equilibrium in a rotating frame. Mathematically, the method is based on the following theorem, which for now will be referred to as the Bifurcation Theorem: *if a system has an axis of rotational symmetry when the angular frequency $\Omega = 0$, then self-consistent cranking about this axis yields families of symmetry-breaking solutions that bifurcate from the trivial solution at the critical rotational frequencies $\Omega_c = \omega_\mu / K_\mu$, where ω_μ is the random-phase approximation (RPA) frequency of each mode with nonzero spin K_μ .* For example, for pure quadrupole excitations, one would have a bifurcation for each mode for which $|K_\mu| = 1, 2$. However, there are no bifurcations for β -vibrational modes, which have $K = 0$, and these cannot be described with the SCCM formalism. At the moment, I have a heuristic proof of this theorem, as well as concrete examples. I believe it is quite general, if one replaces "RPA" by "small-oscillation". It may also apply to generators of continuous symmetries other than spatial rotations. Thus, it may be useful for describing certain types of molecules and may be applicable to some field theories as well.

Prior knowledge of the RPA solutions permits one to locate the critical points Ω_c . However, the solution trajectory emanating from such a point includes anharmonic corrections within a classical approximation. These solutions, which are extrema of the mean-field equations, are not necessarily minima, but can be found by solving nonlinear self-consistency equations. Since the solutions are classical in nature, they can be quantized as usual by

choosing, say, integer values of the spin $\langle \hat{L}_3 \rangle$ about the cranking axis. However, in general, there may be quantal-fluctuation corrections, which are typically of order $1/A$ or, depending on the model, $1/N$ (where N is of the order of the number of valence particles), that are not entirely accounted for by semiclassical quantizations. Such corrections can be systematically taken into account by performing a quantized boson expansion about the SCCM mean field, starting with the corresponding RPA.⁷

The proposed method has been tested on the γ vibrations of the interacting boson model (IBM)⁸ in the deformed SU(3) limit, which is a collective model, and those of the microscopic volume-conserving anisotropic oscillator (VCAO) shell model. The latter work was done partly in collaboration with R. G. Nazmitdinov of Dubna, who was a summer visitor at Notre Dame. For both models, the cranking approximation was used to obtain the energies of the sequence of aligned n -phonon γ -vibrational bandheads with $K = 2n$. Since the exact eigenvalues of the IBM in the SU(3) limit are known analytically,⁸ a direct comparison with the results of the self-consistent cranking approximation can be made. In fact, I find that for this case the error in the approximation is only of order $1/N$, where N is the total number of bosons (half the number of valence nucleons), which is the typical accuracy of the ordinary cranking approximation. In this model the anharmonic contribution is included exactly. As expected from the theorem, the critical cranking frequency is precisely half the small-oscillation frequency.

It has been known for a long time that β and γ vibrations can be described by means of a time-dependent version of the VCAO model called the *vibrating potential model* (VPM), which may include various embellishments such as spin-orbit coupling and the pairing interaction. In its unadorned form, the VPM is interpreted physically as a rough model of giant isoscalar quadrupole resonances.⁹ Since, as recently shown, the VCAO is equivalent to the Hartree approximation applied to a system of nucleons interacting via an exotic many-body effective interaction,¹⁰ it follows that the cranked version of the VCAO model is truly an example of a cranked fermion mean-field approximation. While the cranked VCAO has been amply studied for ordinary rotational bands,¹¹ the present solutions, which represent anharmonic vibrations, are new.

Some calculated results are shown in Figs. 1 - 4. Figs. 1, 2 depict the results for the γ -vibrational bandheads with $K = 2, 4, \dots$, built on axially symmetric configurations with zero angular momentum, such as the ground state. In Fig. 1, the behaviour of the angular momentum $\langle \hat{L}_3 \rangle = K$ along the equilibrium symmetry axis is plotted against the angular frequency Ω in appropriately scaled units, where, in accord with the Bifurcation Theorem, Ω_c is the critical frequency $\Omega_c = \omega_2/2$, ω_2 being the RPA frequency for the $K = 2$ giant resonance mode. It is interesting that this is a backbending trajectory, indicating a compression of the

Fig. 1

Fig. 2

Fig. 3

Fig. 4

level sequence. The corresponding excitation energies are shown in Fig. 2, with an emphasis on the deviation from the harmonic RPA plot. Since $K = 2n$, where n is the number of phonons, and the quantity $\sum_{\perp} \equiv \sum (n_{\perp} + \frac{1}{2})_{\text{occ}}$, a sum over quanta of occupied states perpendicular to the symmetry axis, is large except for the lightest nuclei, the anharmonic corrections to the one- and two-phonon quadrupole giant resonances is actually very small in this model. In spite of the crudeness of the VCAO model, this may be the best estimate thus far of the magnitude of anharmonic corrections to giant resonances.

In addition to γ vibrations built on axially symmetric states with zero spin, one can also use the cranking method to study γ vibrations built on the oblate states that occur at the terminus of normal rotational bands. For the VCAO model, it has been shown that in general there are three distinct RPA modes¹² in this situation. A bifurcation analysis of the cranking theory gives exactly the relationship expected on the basis of the theorem. The three anharmonic solution branches are shown in Fig. 3, 4 for the case of ^{20}Ne , for which the ground-state rotational band terminates at a spin $L_0 = 8\hbar$, when the system becomes axially symmetric about the axis of rotation (1-axis). The frequency ω_0 is just the usual VCAO parameter given by $\hbar\omega_0 = 41/A^{1/3}$. Branch *a*, having a negative bifurcation frequency, corresponds to a $\Delta L = -2$ excitation mode, branch *c* corresponds to a $\Delta L = +2$ excitation mode, while branch *b* is just the ordinary ground-state rotational band, which can be interpreted as a $\Delta L = -2$ de-excitation mode built on the oblate configuration. While branches *b* and *c* have been found previously,¹¹ the interpretation of *c* as a superposition of vibrations rather than a true rotation was, of course, absent in those works. Branch *a* is completely new. Finally, Fig. 4 is a plot of the excitation energy vs. the magnitude of the spin $L \equiv |\langle \hat{L}_1 \rangle|$, which displays an intriguing forking of the γ vibrations from the oblate terminus point. Such a forking may well be a universal signature of band termination. The curious break in segment *a* of Fig. 4 is merely due to the fact that the angular momentum for this segment changes *sign*, as seen in Fig. 3, while the plot in Fig. 4 involves only the *magnitude* of the angular momentum, as is usual for such plots.

In addition to the $K = 2$ vibration in the VCAO model, there also exists a $K = 1$ mode. What is intriguing about this mode is that it can be shown to correspond to a *tilted rotation* in the cranking formalism. While an ordinary rotational motion pivots about a principal axis of the density distribution, a tilted rotation pivots about an axis that is tilted with respect to a principal axis. The $K = 2$ or γ vibration is not tilted, but the $K = 1$ mode is. There has been considerable interest in tilted rotation recently, much of it being generated by the work of Stefan Frauendorf.¹³ However, his tilted solutions are induced by the effects of odd valence nucleons, whereas the tilted solutions I have been studying are purely collective. Further

discussion of tilted rotation is given in my previous progress report.¹⁴ At the time of this writing, the bifurcation point for the $K = 1$ mode built on spin-zero configurations was calculated and again found to agree exactly with the Bifurcation Theorem, i.e., it coincides with the RPA frequency ω_1 for the $K = 1$ giant quadrupole resonance.⁹ I plan to calculate the full anharmonic solution shortly, which corresponds to a sequence of bandhead states with $K = n$, n being the number of phonons, and also to study the behavior of the tilt angle as a function of n . Electromagnetic transitions will also be studied with the objective of finding an experimental signature of the tilting behavior.

An interesting consequence of these studies is that tilted rotation is quite common. In general, for excitations of a given multipole order λ , the mode with spin $|K| = \lambda$ is not tilted, but all the modes with $|K| < \lambda$ have some degree of tilting. This is not only true in the VCAO model, but also in the IBM, which was discussed in the previous progress report. It should be kept in mind, however, that while these solutions correspond classically to uniform rotation, quantally they are bands of anharmonic vibrational levels. An interesting question that I would like to resolve in the next budget period is whether these solutions account for all cases of tilted rotation in the VCAO and IBM models, or whether there are any other physically distinct types.

In the coming year, I hope to apply the cranking approach to collective vibrations in superdeformed nuclei, where it may be the ideal tool for studying octupole vibrations. On the basis of RPA estimates, it appears that octupole vibrations may be very soft in many superdeformed nuclei.¹⁵ Of course, the RPA, which is valid for small-amplitude excitations, would not be reliable in this case, while the SCCM approach is not limited to small amplitudes.

The main task for the coming year, however, will be to develop a realistic SCCM code to calculate the properties of the low-lying γ vibrations in the nuclei with candidate 2-phonon states proposed by Aprahamian. This will be done in several stages, involving addition of more sophisticated components of the mean field. For example, in the first, only the monopole Cooper pairing will be included, while in a later stage, quadrupole pairing will be included. The early work will focus mainly on energies, while later work will include calculation of transition rates.

REFERENCES

1. A. Aprahamian, to be published; see also H. G. Boerner *et al.*, Phys. Rev. Lett. 66 (1991) 691.
2. H. J. Wollersheim *et al.*, Z. Phys. A. 341 (1992) 137.
3. N. Frascaria, H. Emling, in Notre Dame workshop on giant resonances and related phenomena, Oct. 21 - 23, 1991, ed. U. Garg (unpublished).

4. P. Ring and P. Schuck, The nuclear many-body problem (Springer-Verlag, Berlin, 1980) ch. 10.
5. E. R. Marshalek, Phys. Rev. C 3 (1971) 1710; E. R. Marshalek and M. Sabato, Phys. Rev. C 4 (1971) 1006; Phys. Rev. C 5 (1972) 1130; E. R. Marshalek, Nuovo Cim. Lett. 4 (1972) 631.
6. E. R. Marshalek, in Problems of vibrational nuclei, Zagreb, Croatia, Sept. 24 - 27, 1974, eds. G. Alaga, V. Paar and L. Sips (North Holland, Amsterdam, 1975) p. 108.
7. E. R. Marshalek, Nucl. Phys. A 275 (1977) 416.
8. F. Iachello and A. Arima, The interacting boson model (Cambridge UP, Cambridge, 1987).
9. T. Suzuki and D. J. Rowe, Nucl. Phys. A 289 (1977) 461.
10. E. R. Marshalek, Phys. Rev. Lett. 51 (1983) 1534; Phys. Rev. C 29 (1984) 640; Phys. Lett. B 244 (1990) 1; H. Sakamoto and T. Kishimoto, Nucl. Phys. A 486 (1988) 1; Nucl. Phys. A 501 (1989) 205, 242.
11. J. P. Blaizot, N. Kassis and G. Ripka, in Heavy-ion, high-spin states and nuclear structure (STI/PUB/386, IAEA, Vienna, 1975); J. P. Blaizot and G. Ripka, Quantum theory of finite systems (MIT Press, Cambridge, Mass., 1985) ch. 6; V. G. Zelevinsky, Sov. J. Nucl. Phys. 22 (1975) 565; A. P. Stamp, Z. Physik A 284 (1978) 305; D. Glas, U. Mosel and P. G. Zint, Z. Physik A 285 (1978) 83; T. Troudet and R. Arvieu, Ann. Phys. (NY) 134 (1981) 1.
12. H. Kurasawa, Prog. Theor. Phys. 64 (1980) 2055.
13. S. Frauendorf, "Cranking About a Tilted Axis", invited talk at the Workshop Symposium, Strasbourg, France, March 1991; "Tilted Rotation", invited talk at the 21st INS International Symposium on Rapidly Rotating Nuclei, Tokyo, Japan, October 1992.
14. E. R. Marshalek, Progress Report DOE/ER/40640-1 (1991).
15. T. Nakatsukasa, S. Mizutori and K. Matsuyanagi, Prog. Theor. Phys. 87 (1992) 607.

II. PRESENTATIONS & PUBLICATIONS

A. Invited Conference Lectures

1. "Reconstructing Broken Symmetries from Deformed Boson Expansions" - 2.5 hour lecture presented at the International Workshop on Nuclear Structure Models, Oak Ridge, Tennessee, March 16 - 25, 1992.
2. "New Perspectives on the Cranking Model", 21st INS International Symposium on Rapidly Rotating Nuclei, Tokyo, Japan, October 26 - 30, 1992.

B. Departmental Seminars

1. "New Vistas in Cranking", Yukawa Institute, Kyoto, Japan, October 19, 1992.
2. "Cranking Anharmonic Collective Oscillations", Department of Physics, University of Washington, Seattle, WA, November 20, 1992.

Publications

1. E. R. Marshalek, "A New Boson Realization for the Isotropic Three-Dimensional Oscillator", J. Math. Phys. 33 (1992) 2972.
2. E. R. Marshalek, "Tipsy Rotators", in Symposium on Contemporary Physics, in honor of the 65th birthday of Prof. Abraham Klein, Drexel University, Philadelphia, October 31 - Nov. 1, 1991, ed. M. Vallieres (proceedings to be published).
3. E. R. Marshalek, "Reconstructing Broken Symmetries from Deformed Boson Expansions" - 2.5 hour lecture presented at the International Workshop on Nuclear Structure Models, Oak Ridge, Tennessee, March 16 - 25, 1992, eds. J. Draayer and W. Nazarewicz (proceedings to be published, World Scientific).
4. E. R. Marshalek and R. G. Nazmitdinov, "Cranking Anharmonic Gamma Vibrations" (accepted for publication, Physics Letters).
5. E. R. Marshalek, "New Perspectives on the Cranking Model", 21st INS International Symposium on Rapidly Rotating Nuclei, Tokyo, Japan, October 26 - 30, 1992, ed. F. Sakata (proceedings to be published, Nuclear Physics).

**DATE
FILMED**

6 / 7 / 93