STUDIES OF YRAST AND CONTINUUM

STATES IN A - 100-200 NUCLEI

Progress Report for 1991

Patrick J. Daly

Principal Investigator

Purdue University, West Lafayette, IN 47907

February, 1992

Prepared for

THE U.S. DEPARTMENT OF ENERGY

UNDER GRANT NO. DE-FG02-87ER40346

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Abstract

This report summarizes progress in nuclear structure research for the year 1991. The highlights include new spectroscopic results for neutron excessive nuclei (around ^{124}Sn and ^{36}S) formed in deep inelastic heavy ion reactions.

I. Research Activities

This has been a busy and productive year for our research group. We have continued a lively collaboration with the Khoo-Janssens Argonne team in investigating superdeformation and related nuclear structural problems in high-spin physics, as well as the spectroscopy of otherwise hard-to-reach neutron excessive nuclei produced in deep inelastic heavy ion reactions. With the resumption of ATLAS operation in mid-year, several new experiments have been performed and the results have been partially analyzed. The bulk of our data analysis is now performed on our new VAX system at Purdue; this has proven to be an important practical step forward, highly beneficial to the effectiveness and morale of our group.

Ian Bearden and Philippe Benet (until his departure in June 1991) have been based at Argonne NL, where they have concentrated mainly on the superdeformation studies and investigations of other high-spin states of nuclei in the $A \sim 190$ region. Bearden has pushed forward his studies of high-spin structures in $^{188,190,192}\text{Hg}$ nuclei, which will be the main topic of his Ph.D. thesis. At Purdue, Bogdan Fornal, Ralf Mayer and David Nisius have devoted most of their efforts to spectroscopic studies of products of heavy ion reactions just above the Coulomb barrier, namely:

- a) ^{34}S , ^{36}S and ^{37}Cl ions on ^{160}Gd targets, and
- b) ^{76}Ge and ^{80}Se ions on $^{122,124}\text{Sn}$ targets.

Most notably, measurements performed for the system $^{124}\text{Sn} + 344 \text{ MeV } ^{80}\text{Se}$ during a Sept. 1991 ATLAS experiment are turning out to be particularly fruitful. In addition, Fornal has made considerable progress towards elucidating the population patterns of products observed following these deep inelastic heavy ion collisions. Visitors Rafal Broda (Cracow) and Santo Lunardi (Padova) have also contributed to progress, and Jan Blomqvist (Stockholm) has been a constant source of theoretical advice and inspiration.

The Tin Experiments

The original aim of these investigations was to complete a series of $B(E2; 10^+ \rightarrow 8^+)$ determinations for $\pi h_{11/2}^n \nu = 2$ states in the even-A Sn nuclei spanning the mass range $A = 116-130$, where the $\nu h_{11/2}$ subshell is being filled. Previously, $\nu h_{11/2}^n 10^+$ isomers were known in five of these nuclei, but not in ^{122}Sn , ^{124}Sn and ^{126}Sn , which cannot be reached by fusion-evaporation reactions. We sought to identify the missing isomers among the products of heavy ion reactions on ^{124}Sn and ^{122}Sn targets. In the first experiment performed in Sept. 1990 we bombarded 1 mg/cm^2 ^{124}Sn and ^{122}Sn targets with μs pulsed beams of 325 MeV ^{76}Ge ions from ATLAS. As described in our last annual report¹⁾ we identified 10^+ isomers with 62 ± 3 and $45 \pm 5 \mu\text{s}$ half-lives in ^{122}Sn and ^{124}Sn , but could not find a 10^+ isomer in ^{126}Sn . Yield measurements for these reactions indicated that yrast states in In, Sn and Sb nuclei extending over a broad mass range were populated with surprisingly large cross sections. Many of the product nuclei were poorly known, and it became apparent that this method of excitation offered attractive opportunities for wide ranging spectroscopic studies. Much of our recent effort has been focused on follow-up experiments and on analysis and interpretation of the extensive data obtained.

Even-A Tin Isomers and Filling of the $\nu h_{11/2}$ Subshell

The results of the $^{124}\text{Sn} + 325 \text{ } ^{76}\text{Ge}$ experiments virtually completed a series of $B(E2)$ determinations for $(\nu h_{11/2})^n$ states in even-A $^{116-130}\text{Sn}$ isotopes, with ^{126}Sn now the only missing one. In Fig. 1, the transition amplitudes $\sqrt{B(E2; 10 \rightarrow 8)}$ for the Sn isotopes are plotted versus A . In the BCS approximation, the $E2$ transition amplitude is proportional to $u^2 - v^2$, where $v^2 = 1 - u^2$ is the degree of filling of the $h_{11/2}$ subshell. At the point of half-filling, $u^2 = v^2 = 0.5$, particle and hole contributions are equal in magnitude with opposite signs, hence the total amplitude is zero. This happens in the Sn isotopes close to $A = 123$

A

A

Fig. 1. a) Measured E2 transition amplitudes for the $(\nu h_{11/2})^n 10^+ \rightarrow 8^+$ transitions in even-A Sn isotopes. b) A similar presentation of E2 amplitudes for $(\nu i_{13/2})^n 12^+ \rightarrow 10^+$ transitions in even-A Pb isotopes.

or $N = 73$. On the other hand, in $N = 82$ isotones the proton $h_{11/2}$ half-filling occurs²⁾ just below $Z = 71$, two units lower than for neutrons. By means of BCS calculations, this marked difference between protons and neutrons has been traced to the relative $s_{1/2}$, $d_{3/2}$ and $h_{11/2}$ single particle energies. For protons at $N = 82$ the orbitals are actually nearly degenerate, but for neutrons the $s_{1/2}$ state and, to a lesser extent, the $d_{3/2}$ come well below the $h_{11/2}$ state. The differences in single particle energy spacings are a consequence of the Coulomb potential, which for protons raises the $s_{1/2}$ and $d_{3/2}$ energies more than the $h_{11/2}$ energy.

An enlightening comparison between the effective E2 charges observed in tin and lead isotopes has also been developed (Fig. 1). In the two-hole nuclei ^{130}Sn and ^{206}Pb , the neutron effective charges are very similar, close to 0.9 e. However, in both cases the effective charge more than doubles to values of about 2.0 e in the lighter isotopes around the middle of the major shell, where E2 excitations within the shell can build up low-lying E2 collectivity. A paper³⁾ describing these findings has recently been accepted for publication in Phys. Rev. Lett.

New $19/2^+$ Isomers in ^{119}Sn , ^{121}Sn and ^{123}Sn

The off-beam spectra obtained in the $^{122,124}\text{Sn} + 325 \text{ MeV } ^{76}\text{Ge}$ experiment showed unknown γ -rays, decaying with 5-10 μs half-lives, that we suspected might be due to isomerism in odd-A Sn nuclei. In a follow-up experiment at ATLAS, we bombarded ^{122}Sn and ^{124}Sn targets with pulsed 344 MeV ^{80}Se beams in a 10 μs beam-on, 50 μs beam-off configuration, and recorded off-beam γ -ray singles spectra in E_γ versus time mode, as well as $\gamma\gamma$ coincidence events. With the ^{124}Sn target, the off-beam spectra showed six strong new γ -rays that could be grouped in pairs as follows: 818 and 1220 keV, 841 and 1151 keV, and 838 and 1107 keV. Within each pair, the γ -rays had similar intensities, were mutually coincident and

Fig. 2. Energy systematics for even-A and odd-A Sn nuclei in the A = 116-124 mass range, including the new 19/2⁺ isomers.

decayed with similar half-lives:

$$T_{1/2} \text{ (818 and 1220 keV) } = 9.6 \pm 1.2 \text{ } \mu\text{s}$$

$$T_{1/2} \text{ (841 and 1151 keV) } = 5.3 \pm 0.5 \text{ } \mu\text{s}$$

and $T_{1/2} \text{ (838 and 1107 keV) } = 7.4 \pm 2.6 \text{ } \mu\text{s}$

These three new isomers have been interpreted by analogy with a known $1.75 \text{ } \mu\text{s}$ isomer⁴⁾ in ^{117}Sn as $19/2^+$ states of $\nu h_{11/2} \times 5^-$ character in the $^{119,121,123}\text{Sn}$ nuclei, each of them decaying to $15/2^-$ and $11/2^-$ states by a cascade of M2 and E2 transitions (Fig. 2). The 0^+ , 2^+ and 5^- states in the neighboring even-A Sn nuclei are also shown in this figure and it is obvious that the $15/2^-$ and $19/2^+$ energies follow closely the highly regular systematics of the 2^+ and 5^- energies.

The $(\nu h_{11/2}^2 s_{1/2})19/2^+$ states are long-lived because they can decay only by configuration-forbidden M2 transitions to the $15/2^-$ states. The $B(M2)$ values deduced are smaller than 10^{-3} W.u., but there is a possibility that higher-lying μs isomers in these nuclei may have influenced the half-life determinations. In any case, the identification of the $15/2^- \rightarrow 11/2^-$ transitions in $^{119,121,123}\text{Sn}$ is a significant first step toward further studies of $(\nu h_{11/2})^n \nu = 3$ yrast states in these nuclei. A Short Note on this work⁵⁾ has recently been accepted for publication in Z.f. Physik and Ralf Mayer will talk about these results at the Washington DC APS Meeting.

Yrast States in In and Sb Nuclei

From the data obtained in the $^{122,124}\text{Sn} + 344 \text{ MeV } ^{80}\text{Se}$ experiments, it is apparent that this deep inelastic method of excitation delivers a score of nuclei with comparable yields, which are sufficient for yrast spectroscopy in an essentially untouched region of neutron excessive nuclei around ^{124}Sn . Making isotopic assignments for the previously unknown γ -ray cascades that are observed is one of the biggest problems. Thus far we have been able to identify yrast cascades in the $A = 115, 117$ and 119 In isotopes and in the $A = 121, 123$ and 125 Sb isotopes. States of the type $(9/2^+ \times 2^+) 11/2^+, 13/2^+$ in In and $(7/2^+ \times 2^+)$

$9/2^+$, $11/2^+$ in Sb were known from radioactivity work and the ground state transitions from these states are the lowest γ -rays in the six identified yrast cascades. We have also tried to exploit $\gamma\gamma$ coincidences between light and heavy reaction partners for isotopic assignments. As an example, some γ -rays of ^{125}Sb (from lp pickup) could be assigned based on their observed coincidences with known γ -rays of ^{79}As (from lp stripping). Much remains to be done in the analysis.

Most notably, the new yrast cascades identified in ^{123}Sb and ^{125}Sb de-excite μs isomers in the 2-3 MeV excitation range, and we expect to be able to interpret the structure of the yrast states with some confidence. David Nisius will give a talk about these Sb isomeric decays at the Washington DC APS meeting.

Population Patterns in $^{122,124}\text{Sn} + ^{76}\text{Ge}(^{80}\text{Se})$ Collisions

For the reactions of $^{122,124}\text{Sn}$ targets with ^{76}Ge and ^{80}Se ions, 10-15% above the Coulomb barrier, we have performed extensive product yield determinations by in-beam, off-beam and radioactivity measurements. The results include the yields - on the same scale - of:

- 1) 10^+ and $19/2^+$ μs isomers in $^{118-124}\text{Sn}$
- 2) the ^{123}Sn and ^{125}Sn $3/2^+$ "ground states" - from radioactivity
- 3) eight high-spin β -decay isomers in ^{116}In - ^{123}In
- 4) several fusion evaporation products.

One firm and striking result is that the 10^+ and $19/2^+$ isomers in the seven nuclei ^{118}Sn to ^{124}Sn are populated with comparable cross sections from the ^{124}Sn target.

Measurements of $4^+ \rightarrow 2^+$, $2^+ \rightarrow 0^+$ coincidence intensities for even-even Pd, Cd, Sn, Te and Xe nuclei mapped out the distribution of yields with A and Z in these reactions. Analysis of these distributions gave an estimate of the average number of neutrons evaporated from the excited deep inelastic reaction products,

Fig. 3. Relative yields for even-even products of the reaction $^{124}\text{Sn} + 344 \text{ MeV } ^{80}\text{Se}$. Cross-hatching denotes the target and projectile nuclei, and question marks nuclei with unknown level spectra. The larger yields are well correlated with the minimum nuclear potential energy contours calculated for this dinuclear system.

which could then be used to extract the pattern of primary reaction products. It was possible to understand this pattern in terms of N/Z equilibration⁶⁾ in the dinuclear system formed in the deep inelastic collision. Calculations of minimum potential energy surfaces could be used to explain the gross features of the product distribution for the Sn + ^{76}Ge (^{80}Se) systems (Fig. 3), and to predict the outcome of future experiments using other target-projectile combinations.

Bogdan Fornal will talk about these advances at the Washington DC APS Meeting.

Neutron-Rich sd-f Shell Nuclei from Deep Inelastic Reactions

In the Argonne studies of superdeformed (SD) bands in the A ~ 190 region, high quality $\gamma\gamma$ -coincidence data were recorded for reactions of ^{34}S , ^{36}S and ^{37}Cl beams on ^{160}Gd targets. We briefly described in our last report¹⁾ how spectroscopic information about target-like and projectile-like deep inelastic reaction products could also be extracted from these data by emphasizing lower multiplicity subsets of the coincidence events. Gates placed on known γ -rays in the A ~ 160 products selected individual reaction channels and identified coincident γ -rays in light partner products. Transfer of protons from projectile to target, with or without exchange of neutrons, was found to be generally favored, leading to excited neutron-rich light nuclei, some difficult to reach by other means. Generally, yrast states in these neutron-rich products were preferentially populated.

Notable results include the observation in these reaction of yrast cascades up to the highest known yrast states in the four N = 20 isotones ^{34}Si , ^{35}P , ^{36}S and ^{37}Cl . In two N = 19 isotones ^{33}Si and ^{34}P , and two N = 22 isotones ^{38}S and ^{39}Cl , previously unknown yrast states have been identified. An important aspect of these results is that they test the predictions of up to date shell model calculations in a region where such calculations are thought to be rather accurate. A paper based on this work is being prepared.

Superdeformation and Other High-spin Projects

In the projects described up to the point, the Purdue participants led the investigations and performed essentially all the data analysis. Our group participated in several other ATLAS experiments in a supporting role. These experiments are described in some detail in the Argonne Annual Report, and so a listing of the project titles in this report would seem to be sufficient.

- a) Characterization of the Superdeformed Band in ^{189}Hg
- b) Superdeformation Studies in $^{195,196}\text{Pb}$
- c) Population of SD Bands, Competition with Fission and Constraints on the Well Depth
- d) Search for the γ -rays Connecting SD and Normal States
- e) Entrance Channel Dependence in the Population of the SD Bands in ^{192}Hg
- f) Yrast and Near Yrast Spectroscopy in $^{188,189}\text{Hg}$.

References

1. P. J. Daly, Annual Report for 1990, DOE/ER/40346-5 (1991).
2. J. H. McNeill et al., Phys. Rev. Lett. 63 860, (1989).
3. R. Broda et al., Phys. Rev. Lett., in press (1992).
4. O. Hashimoto et al., Nucl. Phys. A318, 145 (1979).
5. R. H. Mayer et al., Z.f. Physik, in press (1992).
6. H. Freisleben and J. V. Kratz, Phys. Reports 106, 1 (1984).

II. Personnel

Senior Staff

Patrick J. Daly, Professor of Chemistry, Principal Investigator
Zbigniew W. Grabowski, Professor of Physics
Rafał Broda, Visiting Assistant Professor
Philippe Benet, Research Associate
Bogdan Fornal, Research Associate

Graduate Students

Ian G. Bearden
Ralf H. Mayer
David Nisius
Laura Richter

III. Publications

"K X-ray Yields Associated with the Superdeformed Band of ^{192}Hg ", E. F. Moore, R. V. F. Janssens, I. Ahmad, M. P. Carpenter, A. M. Baxter, M. E. Bleich, P. B. Fernandez, T. Lauritzen, T. L. Khoo, I. G. Bearden, Ph. Benet, P. J. Daly, U. Garg, W. Reviol, D. Ye, Phys. Lett. **B258**, 284 (1991).

"Superdeformed Bands in $^{189,190}\text{Hg}$ ", M. Drigert, M. P. Carpenter, R. V. F. Janssens, E. F. Moore, I. Ahmad, P. B. Fernandez, T. L. Khoo, F. L. H. Wolfs, I. G. Bearden, Ph. Benet, P. J. Daly, U. Garg, W. Reviol, D. Ye, R. Wyss, Nucl. Phys. **A530**, 452 (1991).

"Proton-rich Shell Model Nuclei with $N \sim 82$ ", Patrick J. Daly in Understanding the Variety of Nuclear Excitations, ed. A. Covello (World Scientific, Singapore, 1991) pp. 53-60.

"Multiple Band Structures in ^{191}Hg ", D. Ye, K. B. Beard, U. Garg, R. V. F. Janssens, M. P. Carpenter, I. Ahmad, T. L. Khoo, E. F. Moore, F. L. H. Wolfs, Ph. Benet, Z. W. Grabowski, M. Drigert, Nucl. Phys. (in press) 1992.

"Yrast Isomers in Tin Nuclei from Heavy Ion Collisions and Filling of the $h_{11/2}$ Subshell", R. Broda, R. H. Mayer, I. G. Bearden, Ph. Benet, P. J. Daly, Z. W. Grabowski, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, T. Lauritsen, E. F. Moore, S. Lunardi, J. Blomqvist, Phys. Rev. Lett. (in press) 1992.

"Characterization of the Superdeformed Band in ^{189}Hg ", I. G. Bearden, R. V. F. Janssens, M. P. Carpenter, E. F. Moore, I. Ahmad, A. M. Baxter, Ph. Benet, P. J. Daly, M. W. Drigert, P. B. Fernandez, U. Garg, Z. W. Grabowski, T. L. Khoo, T. Lauritsen, W. Reviol, D. Ye, Z.f. Physik (in press) 1992.

"New $19/2^+$ Isomers in ^{119}Sn , ^{121}Sn and ^{123}Sn ", R. H. Mayer, B. Fornal, R. Broda, I. G. Bearden, Z. W. Grabowski, S. Lunardi, P. J. Daly, T. Lauritsen, M. P. Carpenter, R. V. F. Janssens, T. L. Khoo, Y. Liang, Z.f. Physik A (in press) 1992.

"Shape Driving Effects in ^{193}Tl from the Spectroscopy of Yrast and Near-Yrast States", W. Reviol, M. P. Carpenter, U. Garg, R. V. F. Janssens, I. Ahmad, I. G. Bearden, Ph. Benet, P. J. Daly, M. W. Drigert, P. B. Fernandez, T. L. Khoo, E. F. Moore, S. Pilotte, D. Ye, submitted to Nucl. Phys. A, Jan. 1992.

"Isomeric Decay Studies Using a Recoil Mass Separator: Exotic $N = 82$ and 83 Nuclei ^{153}Yb , ^{153}Lu , ^{154}Lu and ^{154}Hf " J. H. McNeill, A. A. Chishti, P. J. Daly, W. Gelletly, M. A. C. Hotchkis, M. Piiparinen, B. J. Varley, P. J. Woods, J. Blomqvist, submitted to Z.f. Physik A, Feb. 1992.

END

**DATE
FILMED**

4 / 27 / 92